TITLE:

Tricuspid valve leaflet remodeling in sheep with biventricular heart failure: A comparison

between leaflets

AUTHORS:

Colton J. Kostelnik[a,†], William D. Meador[a,†], Chien-Yu Lin[a], Mrudang Mathur[b], Marcin

Malinowski[c,d], Tomasz Jazwiec[c,e], Zuzanna Malinowska[g], Magda L. Piekarska[c,d], Boguslaw

Gaweda[c,f], Tomasz A. Timek[c], Manuel K. Rausch[a,b,g,*]

AFFILITATIONS:

[a]Department of Biomedical Engineering, The University of Texas at Austin, Austin, TX, USA

[b]Department of Mechanical Engineering, The University of Texas at Austin, Austin, TX, USA

[c]Division of Cardiothoracic Surgery, Corewell Health, Grand Rapids, MI, USA

[d]Department of Cardiac Surgery, Medical University of Silesia, School of Medicine in Katowice,

Katowice, Poland

[e]Department of Cardiac, Vascular, and Endovascular Surgery and Transplantology, Medical

University of Silesia in Katowice, Silesian Centre for Heart Disease, Zabrze, Poland

[f]Department of Cardiac Surgery, Clinical District Hospital No. 2, Faculty of Medicine University

of Rzeszow, Rzeszow, Poland

[g]Department of Aerospace Engineering & Engineering Mechanics, The University of Texas at

Austin, Austin, TX, USA

* Corresponding Author: Manuel K. Rausch, manuel.rausch@utexas.edu, +1 (512) 471-5147,

2501 Speedway, EER Building, Room 7.620

† Colton Kostelnik and William Meador are co-first authors

ABSTRACT: Tricuspid valve leaflets are dynamic tissues that can respond to altered

biomechanical and hemodynamic loads. Each leaflet has unique structural and mechanical

properties, leading to differential *in vivo* strains. We hypothesized that these intrinsic

differences drive heterogeneous, disease-induced remodeling between the leaflets. Although

we previously reported significant remodeling changes in the anterior leaflet, the responses

among the other two leaflets have not been reported. Using a sheep model of biventricular

heart failure, we compared the remodeling responses between all tricuspid leaflets. Our results

show that the anterior leaflet underwent the most significant remodeling, while the septal and

posterior leaflets exhibited similar but less pronounced changes. We found several between-

leaflet differences in key structural and mechanical metrics that have been shown to contribute

to valvular dysfunction. These findings underscore the need to consider leaflet-specific

remodeling to fully understand tricuspid valve dysfunction and to develop targeted therapies

for its treatment and more accurate computational models.

KEYWORDS: Leaflet-specific Remodeling; Tachycardia-induced Cardiomyopathy; Biaxial

Testing; Histology; Two-photon Microscopy

STATEMENT OF SIGNIFICANCE:

Our study is significant as it advances our understanding of tricuspid valve remodeling by

providing a comprehensive analysis of all three leaflets in a sheep model of biventricular heart

failure. Unlike prior works that focused primarily on the anterior leaflet or generalized leaflet

changes, we integrated morphological, histological, immunohistochemistry, biaxial mechanical

testing, and two-photon microscopy to quantify differences between all three tricuspid valve

leaflets (anterior, posterior, and septal) across multiple functional scales. This comprehensive

approach highlights the unique remodeling response of each leaflet. Our findings offer critical

insights for developing targeted therapeutic strategies and improving computational models of

disease progression.

 GRAPHICAL ABSTRACT

56
57
58

1. INTRODUCTION

Tricuspid regurgitation (TR) is a highly prevalent echocardiographic finding detected in 82-86% of people [1]. Approximately 1.6 million Americans suffer from moderate to severe TR, which, if left untreated, can lead to multiple cardiovascular complications [2]. More than 90% of patients with severe TR exhibit a "functional" etiology, wherein the valve dysfunction is secondary to other cardiac pathologies such as left-sided heart disease or pulmonary hypertension [3–5]. The valve itself has historically been considered structurally and mechanically intact, and this assumption has guided a conservative treatment approach that focuses on managing the primary condition to resolve any TR [6,7]. However, early surgical repair of the tricuspid valve during mitral or aortic valve surgeries has shown promise in improving patient outcomes and diminishing cardiac-related mortality [8,9]. Despite this evolution of TR management, it remains undertreated and poorly understood.

Heart valve leaflets were once considered "passive flaps" that merely coordinated unidirectional blood flow between the heart chambers akin to a check valve. However, we now know that heart valve leaflets are highly organized tissues with multiple cell types and intricate extracellular architectures that actively remodel by contracting and reshaping to optimize coaptation [10–12]. In diseased patients, leaflet remodeling induces a fibrotic response (i.e., maladapt), which impedes proper kinematics and coaptation [13]. Animal studies of heart failure and chronic ischemia have shown that maladaptive changes in the mitral valve leaflets manifest by increased size, thickness, and stiffness [13–16]. We have demonstrated in two animal models that similar remodeling patterns exist in the tricuspid valve leaflets (e.g., increases in area, thickness, and stiffness) in response to altered biomechanical and

hemodynamic loads [17,18]. These remodeling-induced leaflet changes affect leaflet stresses

and leaflet contact area (i.e., coaptation area), indicating that valve function is sensitive to

leaflet maladaptation [19].

Moreover, we have previously demonstrated that the tricuspid leaflets in healthy sheep

exhibit distinct, leaflet-specific properties, including unique morphology, mechanical properties,

and *in vivo* mechanics [20,21]. In our prior study of tricuspid valve maladaptation, we found

that the anterior tricuspid leaflet exhibited significant remodeling changes on all functional

scales in sheep with biventricular heart failure after 2-3 weeks of tachycardia-induced

cardiomyopathy (TIC) [17]. Building on this work, here we aim to quantify the remodeling

changes in the septal and posterior tricuspid leaflets from the same sheep and compare the

remodeling patterns across all three leaflets. We hypothesize that the leaflet-specific

mechanics drive heterogeneous remodeling responses between the tricuspid leaflets.

Specifically, we hypothesize that the TIC anterior leaflets exhibit the most significant

remodeling changes relative to its control due to its larger *in vivo* area strains, while the TIC

septal and posterior leaflets remodel to a lesser extent [20]. This distinction of differential

remodeling may be vital in uncovering potential therapeutic targets tailored to specific leaflets

and informing computational models of disease progression.

2. MATERIALS & METHODS

2.1 Animal Model, Medication, and Procedures

This research complied in accordance with the Principles of Laboratory Animal Care,

formulated by the National Society for Medical Research, and adhered to protocols outlined in

the Guide for Care and Use of Laboratory Animals, published by the National Institutes of

Health. Additionally, this protocol was developed, reviewed and performed in accordance with

the approval of a local Institutional Animal Care and Use Committee at Corewell Health. The

specific approval numbers are 2017-357, 2018-402, 2018427, and 2018-439.

The TIC sheep model used in this study has been previously described in detail and

demonstrated as a reliable and repeatable model of biventricular heart failure in sheep [22].

Detailed descriptions of the animal model, medications, and procedures are available in our

prior work [17]. We randomly assigned adult male Dorsett sheep to either the control (CTL) (n =

17, 59.9 ± 4.6 kg) or TIC (n = 33, 60.1 ± 5.3 kg) groups. Before initiating pacing, we monitored

baseline ventricular function and valvular competence in all animals using epicardial

echocardiography. The TIC sheep were then subject to a 180-260 beats per minute pacing

protocol for 19 ± 6 days. Following the pacing period, we performed the terminal procedure on

all TIC sheep and recorded hemodynamic and echocardiographic data over three cardiac cycles.

In CTL sheep, we performed the terminal procedure following baseline epicardial

echocardiography. We reported all hemodynamic and echocardiographic data in Table 1 of our

previous study [17]. After completing the terminal procedures, we isolated the tricuspid valve

from each animal for further analysis.

For disclosure, we previously published data on the anterior leaflet morphology,

thickness, biaxial mechanics, histology, collagen content, and two-photon microscopy from the

sheep included in this study [17]. We include these data here again with expressed permission

from the editorial office.

2.2 Morphology and Storage

We isolated the tricuspid valve leaflet complex from each heart and separated the

leaflets at the posterior-septal commissure. We then floated each valve on 1x PBS and imaged

them on a calibrated grid to compare morphological measures such as leaflet area between CTL

and TIC animals (Figure 1a). The leaflets were then separated and cryopreserved in DMEM

(VWR L0101-0500, Radnor, PA, USA) containing 10% DMSO (VWR BDH1115-1LP) supplemented

with a protease inhibitor (ThermoFisher, A32953, Weltham, MA, USA) and stored at -80°C until

tested. Cryogenic studies have shown that the mechanical properties of soft tissues were

minimally impacted after storage at temperatures below 0°C [23,24].

2.3 Histology and Immunohistochemistry

After thawing, we cut and fixed radial strips (from near-annulus to free edge) of all

leaflets in 10% neutral buffered formalin for 24 hours and then transferred to 70% ethanol for

long-term storage. The fixed leaflet strips were shipped to a histological service (HistoServ Inc.,

Amaranth, MD) for embedding, sectioning at 5 μm slice thickness, and staining with

Hematoxylin & Eosin and immunohistochemistry. Using a light microscope (Ti2-E High-content

Analysis System, Nikon, Tokyo, Japan) with a 10x objective, we acquired full section images for

all histological and immunohistochemistry sections. After imaging, we used ImageJ (National

Institutes of Health, Bethesda, MD, USA) to measure the total length of the tissue section along

the atrial surface and separated each section into three equidistant regions (i.e., near-annulus,

belly, and free edge). We then measured the normal vectors between the atrial and ventricular

surfaces at three random positions within each region. Finally, we averaged all regional

thickness measurements into the total leaflet thickness for each sample.

We assessed four cellular markers associated with remodeling processes: (i) alpha-

smooth muscle actin (α-SMA) (Abcam, ab5694, 1:600, Cambridge, MA, US), (ii) Ki67 (Abcam,

ab15580, 1:1250, Cambridge, MA, US), (iii) matrix-metalloprotease 13 (MMP13) (Abcam,

ab39012, 1:50, Cambridge, MA, US), and (iv) transforming growth factor beta-1 (TGF-β1)

(Abcam, ab9758, 1:100, Cambridge, MA, US). A custom MATLAB program segmented the full

cross-sectional leaflet image into three equidistant length regions (near-annulus, belly, and free

edge). We then interpolated ten thickness regions of the leaflet cross-sectional images between

the atrial and ventricular surfaces, resulting in a total of 30 regions. Each region underwent

analysis through our validated custom MATLAB program to spatially resolve the presence of

positive pixels and normalize their percentage to the total pixels in that region [17].

2.4 Biaxial testing

We tested leaflet samples and analyzed the biaxial mechanics using the methods fully

described in our prior work [17,21]. First, we isolated a 7x7 mm square sample from the belly

region of each septal, anterior, and posterior leaflet. We then applied four fiducial markers to

the atrial surface of each sample to enable strain tracking during testing. Before testing, we

measured sample thickness at four locations using a digital thickness gauge (547-500S,

Mitutoyo Corp., Kawasaki, Japan) and captured an image of the stress-free configuration on a

calibrated grid. Next, we mounted all samples on a biaxial testing device (Biotester, CellScale,

Waterloo, ON, Canada) submerged in 37°C 1x PBS, and performed ten preconditioning cycles of

equibiaxial extension to 300 mN followed by two final equibiaxial cycles to 300 mN. While

testing, we continuously captured circumferential and radial forces and images of the fiducial

markers at 5 Hz.

We used the image analysis software Labjoy (CellScale, Waterloo, ON, Canada) to track

the coordinates of the fiducial markers throughout testing. Our analysis focused on the

downstroke of the last loading cycle, and we reported membrane tension-stretch data relative

to the preloaded stress-free configuration. To characterize tissue deformation, we computed

the deformation gradient tensor $\boldsymbol{F}$ relative to the stress-free reference state and derived the

right Cauchy-Green deformation tensor $\boldsymbol{C} = \boldsymbol{F}^T\boldsymbol{F}$ to determine in-plane stretches throughout

the experiment. Membrane tension was calculated by dividing the measured force by the

perpendicular rake-to-rake distance in the deformed configuration at each time point.

To quantitatively compare the nonlinear membrane tension-stretch curves between CTL

and TIC for all three leaflets, we analyzed four characteristic metrics of the resulting nonlinear J-

shaped curves: (1) toe stiffness, defined as the slope in the lower linear region of the curve; (2)

calf stiffness, measured as the slope near 20 N/m membrane tension; (3) transition stretch,

identified as the stretch value at the curve's inflection point (approximated by the intersection

of toe and calf stiffness lines); and (4) anisotropy index, calculated as the ratio of

circumferential to radial stretches at 20 N/m tension.

2.5 Two-Photon Microscopy

We imaged and analyzed the collagen microstructure and cell nuclei using the same

two-photon microscopy methods described in our previous work [21,25]. The mechanically

tested square sample from the leaflet belly was briefly stained with Hoechst 33342 (Thermo

Fischer Scientific, Hoechst 33342, Waltham, MA, US) for 20 minutes to visualize the cell nuclei.

The tissue sample was then optically cleared under sonication for 30 minutes in a solution

containing glycerol:DMSO: 5x PBS at 50:30:20% (v/v), respectively [26]. Next, we imaged the

leaflet samples with a two-photon microscope (Ultima IV, Bruker, Billerica, MA, USA) to

visualize collagen fiber orientation via second harmonic generation (SHG) and cell nuclei via

fluorescent excitation, respectively. Before imaging, the leaflet samples were placed on a foil-

lined glass slide with a coverslip to flatten the sample during imaging without imposing any

stresses. We captured images from the atrial to the ventricular surface at three centrally

located regions (500x500 μm, for each region) using a 20x water-immersion objective

(Olympus, XLUMPLFLN, Center Valley, PA, US) at an excitation wavelength of 900 nm for SHG

and 800 nm for fluorescence. We collected the backscattered SHG signal through a PMT

channel filter (460 nm ± 25 nm) and acquired a z-stack of images at a step size of 10 μm. To

analyze the SHG images, we used the orientation distribution analysis with the Gaussian

gradient method in the ImageJ plugin OrientationJ (National Institutes of Health, Bethesda, MD,

USA). We then averaged and interpolated across the three z-stacks for each leaflet sample and

then fit a symmetric von Mises distribution to each histogram to estimate the mean fiber angle

μ and the fiber orientation concentration κ [27]. To analyze the cell nuclei morphology and

orientation, we used our custom-validated MATLAB code to quantify individual nuclei contours,

nuclear aspect ratio (NAR), and circularity from each image [17]. Similarly, we fit von Mises

distributions to the nuclear orientation histograms and normal distributions to the NAR and

circularity histograms.

2.6 Statistical Analysis

For all morphological, thickness, biaxial, and two-photon distribution data comparisons,

we used a linear mixed effects model in R via the package *afex* (Version 4.4.2). Our models

included fixed effects for animal group (CTL vs. TIC), leaflet type (anterior, posterior, septal),

direction (X vs. Y) where applicable, and their interactions. Random intercepts were included

for subjects to account for repeated measurements within individuals. We performed a Type III

ANOVA analysis with Satterthwaite's method to assess the significance of main effects and

interactions. We then performed one-tailed pairwise comparisons between CTL and TIC animals

for each leaflet type using the *emmeans* package in R, and performed two-tailed pairwise

comparisons using Tukey's method to compare between leaflets from the CTL and TIC animals.

We defined statistical significance as p-values less than 0.05. We reported data as mean with

the standard deviation of the mean.

3. RESULTS

3.1 Animal model outcomes

We isolated 17 tricuspid valves from CTL sheep and 33 tricuspid valves from TIC sheep.

Our study included all tricuspid valve leaflets to assess their geometric and functional changes.

Echocardiographic and hemodynamic evaluations revealed significant parameter changes in TIC

sheep consistent with biventricular dysfunction and functional TR. Key findings included

reduced biventricular ejection fraction, evidence of biventricular remodeling, tricuspid valve

annular dilation, and increased severity of TR (Table 1) [17].

3.2 TIC-induced changes to leaflet geometry

Leaflet geometry, including area and thickness, is critical for assessing valve function in

TR [16]. While prior studies detailed size and shape differences in healthy tricuspid valve

leaflets [21], geometric changes in TIC leaflets remain underexplored. Building on our previous

findings of geometric changes in TIC anterior leaflets [17], we hypothesized that all TIC leaflets

would exhibit distinct geometric changes. To test this, we quantified leaflet area and thickness

across all leaflets (Figure 1a, Figures 2a–c).

We compared the leaflet area measurements between CTL and TIC animals, and we

found that TIC septal and anterior leaflets significantly increased in area compared to

respective CTL leaflets (p = 0.009 and p = 0.002, respectively) (Figure 1b-c). In contrast, the TIC

posterior leaflets' area remained unchanged (p = 0.153) (Figure 1d). We found a significant

interaction between animal group and leaflet type (p = 0.044), highlighting the heterogeneous

remodeling response among leaflets. For instance, in CTL animals, we found that the septal

leaflet area was significantly smaller than the anterior and posterior leaflets (both p ≤ 0.0001),

while among TIC animals, leaflet area differed significantly across all leaflets (all p ≤ 0.0001). We

compared the average thickness between CTL and TIC animals from our histological cross-

sectional images, and we found that only the TIC anterior and posterior leaflets significantly

increased (p = 0.024 and p = 0.033, respectively) in thickness compared to their respective CTL

leaflets (Figures 2d-f). However, we did not detect any significant differences between the

leaflets of CTL animals or between the leaflets of TIC animals. These findings highlight the

distinct growth of each tricuspid leaflet under disease conditions, suggesting differential

remodeling exists between the tricuspid leaflets.

3.3 TIC-induced changes to leaflet mechanics

Leaflet remodeling in heart failure patients alters the mechanical properties of the valve

leaflets [13]. Building on our previous findings of altered anterior tricuspid leaflet mechanics in

TIC animals [17], we hypothesized that each TIC leaflet would exhibit distinct and leaflet-

specific changes in mechanical behavior. All leaflets exhibited the classic J-shaped loading curve

of biological collagenous tissues (Figures 3a, 3d, 3g) [28]. Our analysis did not detect any

significant change in calf stiffnesses between CTL and TIC animals for any of the leaflets, in

either the circumferential and radial direction (Figures 3b, 3e, 3h). However, we did find that

the TIC anterior leaflet exhibited a significant increase in circumferential toe stiffness when

compared to CTL (p = 0.022). In contrast, we failed to detect a significant change in the

posterior and septal leaflets' toe stiffnesses. Our between-leaflet comparison revealed that

circumferential toe stiffness differed significantly between the TIC septal and anterior leaflets (p

= 0.044) (Figures 3b and 3e).

The most pronounced between-leaflet differences emerged when comparing the

transition stretches and anisotropy indices. We found that the anterior leaflet showed a

significant decrease in the circumferential transition stretch between CTL and TIC animals (p =

0.047), while we failed to detect any significant changes in the posterior and septal leaflets

(Figure 4 a-c). Our between-leaflet comparison within CTL animals revealed that the posterior

leaflet radial transition stretches significantly differed from the other leaflets (anterior-

posterior: p = 0.0009; septal-posterior: p ≤ 0.0001). Among the TIC animals, we found that the

septal leaflet's circumferential transition stretch was significantly different from the anterior

leaflet (p = 0.036), while the posterior leaflet's radial transition stretch was significantly

different from the other leaflets (anterior-posterior: p ≤ 0.0001; septal-posterior: p = 0.0002).

Our analysis of anisotropy indices revealed a near-significant interaction effect between animal

group and leaflet type (p = 0.052). We found that the septal leaflet anisotropy index

significantly decreased (p = 0.004) between CTL and TIC animals to a value below 1, indicating

larger stretches in the radial direction (Figure 4d). In contrast, the anterior and posterior

leaflets retained stable anisotropic properties (Figures 4e and 4f). Our between-leaflet

comparison within CTL animals revealed significant differences between all leaflets (septal-

anterior: p = 0.024; anterior-posterior: p = 0.021; septal-posterior: p ≤ 0.0001). Among TIC

animals, the posterior leaflet's anisotropy differed significantly from both the other leaflets

(anterior-posterior: p = 0. 0009; septal-posterior: p = 0.0005), revealing a disease-induced shift

in the septal leaflet's radial behavior. Overall, TIC induced the most significant remodeling

mechanical changes in the septal leaflet. While the CTL septal leaflet exhibited greater

circumferential compliance, the TIC septal leaflet remodeled in a manner that resulted in

greater radial compliance, aligning its behavior more closely with the anterior and posterior

leaflets. These findings underscore the pronounced remodeling of the septal leaflet under

disease conditions, leading to altered anisotropy and compliance patterns.

3.4 Increased expression of cellular remodeling markers in TIC leaflets

Pathological activation of valvular endothelial cells (VECs) and valvular interstitial cells

(VICs) promotes extracellular matrix (ECM) remodeling through expression, secretion, and

degradation of load-bearing proteins [29]. We hypothesized that each TIC leaflet would exhibit

elevated expression of remodeling markers, with variations in the magnitude and region of

expression between leaflets. Specifically, we anticipated increased expression of all markers

tested (α-SMA, Ki67, MMP13, TGF-β1) [17,18].

In the TIC septal leaflets, we observed region-specific increases in all four remodeling

markers, particularly in the near-annulus and belly regions, while MMP13 expression decreased

at the free edge (Figure 5). These findings align with our previous observations of increased

remodeling marker expression in TIC anterior leaflets (Figure 6). In the TIC posterior leaflets,

expression of α-SMA and Ki67 increased in the near-annulus region (Figures 7a and 7b). In

contrast, the expression levels of MMP13 and TGF-β1 decreased throughout the TIC posterior

leaflets (Figures 7c and 7d). Additionally, all TIC leaflets exhibited regionally increased cell nuclei

density consistent with elevated Ki67 expression, see Supplementary Figure S1.

3.5 Changes in collagen fiber orientation in TIC leaflets

Collagen fiber orientation is a critical determinant of leaflet stiffness, ensuring

mechanical functionality during the cardiac cycle [30]. Building on previous findings of

increased fiber dispersion in the TIC anterior leaflets [17], we hypothesized that all TIC leaflets

would exhibit increased collagen fiber dispersion. To test this, we fit von Mises probability

distribution functions to collagen fiber orientation histograms for leaflets from CTL and TIC

animals (Figures 8a, 8c, 8e). We quantified the concentration parameter κ from the collagen

fiber orientation histograms and measured fiber dispersion as a function of imaging depth. A

large κ value indicates more aligned fibers, while lower values suggest greater dispersion and

more isotropic distribution of collagen fibers.

Our analysis revealed that the TIC septal and anterior leaflet significantly increased in

fiber dispersion near the atrial surface (p = 0.005 and p = 0.011, respectively) (Figures 8b and

8d). In contrast, we failed to detect any significant changes in the fiber dispersion in the other

depth regions and throughout the posterior leaflet. Our between-leaflet comparison revealed

that a significant difference in fiber dispersion was only present in the TIC animals. We found

that the TIC posterior leaflet exhibited significantly different κ values from the other leaflets in

the 0-33% depth (D1), nearest the atrial surface (anterior-posterior: p = 0.017; septal-posterior:

p = 0.033). In the 33-67% depth (D2), we found that the TIC septal leaflet was significantly

different (p = 0.004) from the anterior leaflet. Additionally, we found that the TIC septal leaflet

significantly differed from the other leaflets in the 67-100% depth (D3), nearest the ventricular

surface (septal-anterior: p = 0. 0.005; septal-posterior: p = 0.028). Similar analyses revealed cell

nuclei of TIC septal leaflets concentrating in the circumferential direction within the 0-33%

depth, see Supplementary Fig. 2. In contrast, we observed no notable difference in cell nuclei

orientation, nuclear aspect ratio, or circularity between CTL and TIC anterior and posterior

leaflets, see Supplementary Fig. 3 and Fig. 4. Overall, we observed that TIC-induced changes

resulted in more radially dispersed collagen fibers in TIC septal and anterior leaflets near the

atrial surface.

4. DISCUSSION

Our study provides a comprehensive analysis of the heterogeneous remodeling

responses of tricuspid valve leaflets in a biventricular heart failure model, highlighting leaflet-

specific changes at the tissue and cellular levels. The distinct morphological and mechanical

properties between the tricuspid leaflets are essential for maintaining healthy valve function

[21,31]. Our findings reveal that biventricular heart failure leads to heterogeneous remodeling

between the tricuspid leaflets.

At the cellular level, we observed localized profibrotic and degradative responses driving

heterogeneous ECM remodeling both between and within tricuspid leaflets [32]. Elevated

expression of α-SMA in all TIC leaflets indicates VIC activation as a central mechanism in leaflet

remodeling [33,34]. This activation appears localized, with distinctive changes in α-SMA

expression within each leaflet, suggesting region-specific remodeling processes. Activated VICs

assume a hypersecretory phenotype – known as myofibroblasts – contributing to fibrosis by

collagen secretion [33]. The expression of MMP13 is indicative of ECM degradation [17]. In the

TIC septal and anterior leaflets' near-annulus and belly regions, elevated expression of these

remodeling markers (α-SMA and MMP13) indicate increased matrix turnover. Conversely, the

decreased expression of α-SMA and MMP13 in the free edge region of septal and posterior

leaflets signify limited ECM turnover. The localization of both Ki67 and TGF-β1 to the

endothelial layer TIC leaflets provides evidence of endothelial-to-mesenchymal transition

(EndoMT). These findings highlight the critical role of endothelial cells in responding to

mechanical stress and regulating fibrotic responses [35]. This regional variation in remodeling

markers highlights the leaflet- and region-specific stimuli, possibly related to mechanical

loading or biochemical environment, which drive the localized remodeling response [36].

Spatially heterogeneous pathological strains imposed by annular dilation or leaflet tethering

trigger these responses [37]. Such strains, often localized along the free edge, trigger

mechanobiological pathways promoting remodeling within that region of the TIC leaflets

[29,38].

The tissue and cellular-level changes observed in the tricuspid leaflets after TIC provide

mechanistic insights into the functional alterations observed in previous *in vivo* studies of TIC

sheep models [39,40]. The TIC septal leaflet exhibited a complex remodeling response

characterized by significant geometric, mechanical, and microstructural changes. We observed

an increase in leaflet area and a shift in the anisotropy index towards greater radial compliance.

This was accompanied by increased collagen fiber dispersion near the atrial surface and

elevated expression of α-SMA and MMP13 in the near-annulus and belly regions. These

findings collectively indicate enhanced matrix turnover and suggest a compensatory

mechanism to accommodate the increased septal-lateral annular dilation observed in TIC

animals [32]. The pronounced remodeling observed in the TIC anterior leaflet aligns with

previous findings of increased strain in this leaflet during *in vivo* studies [39]. We observed a

substantial increase in leaflet area and thickness, accompanied by increased circumferential toe

stiffness and decreased circumferential transition stretch. These changes indicate a tissue and

cellular-level response to maintain coaptation and valve function in the face of increased area

strain and annular dilation associated with TIC [17,39]. In contrast, the posterior leaflet

displayed a more stable mechanical profile, with no significant changes in stiffness or transition

stretch. However, localized increases in α-SMA and Ki67 expression suggest that while the

posterior leaflet may not undergo extensive structural remodeling, it remains responsive to

pathological stimuli.

The heterogeneous remodeling response observed in our biventricular heart failure

model reflects the unique properties of each tricuspid leaflet and the differential mechanical

stimuli they experience [20,21]. The TIC model induces complex changes across the tricuspid

valve apparatus, including right ventricular remodeling, asymmetric annular dilation, and

altered chordal tethering forces [22,39,40]. These extrinsic factors create a diverse mechanical

environment, subjecting each leaflet to unique biomechanical stimuli and influencing their

individual propensity to remodel.

Incorporating these findings into future computational models can significantly enhance

our understanding of valvular disease progression, potentially leading to more clinically

relevant results. Computational models, informed by the differential remodeling responses

observed in the tricuspid valve leaflets, may help simulate various disease scenarios and predict

the impact of different therapeutic interventions [19,41]. This approach may allow for a better

understanding of the underlying mechanisms driving leaflet-specific remodeling and aid in

developing more precise and effective treatment strategies.

While our study provides valuable insights into tricuspid valve leaflet remodeling in the

context of biventricular heart failure, several limitations warrant consideration. The TIC sheep

model replicates key features of biventricular heart failure, but induced disease within weeks

rather than years [22]. Therefore, the observed leaflet changes likely reflect early adaptive

responses. Although the TIC sheep model successfully induced differential leaflet growth and

remodeling, we failed to establish clear correlations between these adaptive responses and

potential stimuli like TR severity or right ventricular dilation. Spatial variations in local

mechanical properties and strain distributions within each leaflet may influence the expression

of remodeling-associated genes and lead to heterogeneous maladaptive responses in VICs

across different leaflet regions [31,42]. Beyond our regional immunohistochemistry analysis of

cellular remodeling markers, our study did not fully explore the potential impact of mechanical

microenvironments within individual leaflets. Additionally, interspecies differences in cardiac

structure and response to injury limit the direct translation of these findings to the

pathogenesis of functional TR in humans. To overcome these limitations, future studies may

explore long-term disease progression models and spatially heterogeneous remodeling

responses within tricuspid valve leaflets.

5. CONCLUSION

In conclusion, our study highlights the differential remodeling responses of tricuspid

valve leaflets in biventricular heart failure, emphasizing the unique remodeling changes each

tricuspid leaflet undergoes. These findings underscore the importance of considering leaflet-

specific properties in understanding tricuspid valve dysfunction. By providing insights into how

mechanical and cellular factors contribute to these differences, our work paves the way for

more targeted and effective interventions in treating tricuspid valve disease.

DATA AVAILABILITY

The data for Figures 1-8 and their appendices are publicly available on the Texas Data

Repository https://doi.org/10.18738/T8/5EPBVN. The histology and immunohistochemistry

leaflet images are publicly available on the Texas Data Repository

https://doi.org/10.18738/T8/2TRJGV.

DISCLOSURE

Manuel K. Rausch has a speaking arrangement with Edwards Lifesciences. The other

authors have no conflicts to declare.

AUTHOR CONTRIBUTIONS

Colton J. Kostelnik: Investigation, Data curation, Formal analysis, Visualization, Writing -

original draft, Writing - review & editing. **William D. Meador:** Conceptualization, Data curation,

Software, Formal analysis, Investigation, Methodology, Writing -- original draft, Writing - review

& editing. **Chien-Yu Lin:** Investigation, Writing - review & editing. **Mrudang Mathur:**

Investigation, Writing - review & editing. **Marcin Malinowski:** Conceptualization, Investigation,

Methodology, Writing - review & editing. **Tomasz Jazwiec:** Investigation, Methodology. **Magda**

L Piekarska: Writing - review & editing. **Boguslaw Gaweda:** Writing - review & editing. **Zuzanna**

Malinowska: Formal analysis. **Tomasz A. Timek:** Conceptualization, Resources, Supervision,

Funding acquisition, Writing - review & editing. **Manuel K. Rausch:** Conceptualization, Writing -

original draft, Writing - review & editing, Supervision, Funding acquisition.

ACKNOWLEDGEMENTS

The research reported in this publication was partially supported by the National Heart,

Lung, And Blood Institute of the National Institutes of Health under Award Numbers

1R01HL165251-01 and R21HL161832 (MKR), F31HL145976 (WDM), the American Heart

Association for their support under Award Number 18CDA34120028 (MKR), and internal grants

from Meijer Heart and Vascular Institute at Corewell Health. The opinions, findings, and

conclusions expressed are those of the authors and do not necessarily reflect the official views

of the National Institutes of Health or the American Heart Association.

The anterior leaflet data reported in this publication was previously published in [17].

This publication expands upon our earlier work by incorporating additional leaflet comparisons

to provide new insights into leaflet-specific remodeling as a response to tachycardia-induced

cardiomyopathy.

REFERENCES

[1]   J.P. Singh, J.C. Evans, D. Levy, M.G. Larson, L.A. Freed, D.L. Fuller, B. Lehman, E.J. Benjamin,
Prevalence and clinical determinants of mitral, tricuspid, and aortic regurgitation (the
Framingham Heart Study), The American Journal of Cardiology 83 (1999) 897–902.
https://doi.org/10.1016/S0002-9149(98)01064-9.
[2]   M. Enriquez-Sarano, D. Messika-Zeitoun, Y. Topilsky, C. Tribouilloy, G. Benfari, H.
Michelena, Tricuspid regurgitation is a public health crisis, Progress in Cardiovascular
Diseases 62 (2019) 447–451. https://doi.org/10.1016/j.pcad.2019.10.009.
[3]   F. Condello, M. Gitto, G.G. Stefanini, Etiology, epidemiology, pathophysiology and
management of tricuspid regurgitation: an overview, Reviews in Cardiovascular Medicine
22 (2021) 1115. https://doi.org/10.31083/j.rcm2204122.
[4]   E.A. Fender, C.J. Zack, R.A. Nishimura, Isolated tricuspid regurgitation: outcomes and
therapeutic interventions, Heart 104 (2018) 798–806. https://doi.org/10.1136/heartjnl-
2017-311586.

[5] R.J. Henning, Tricuspid valve regurgitation: current diagnosis and treatment, Am J
Cardiovasc Dis. 12 (2022) 1–18.
[6] C.-H. Lee, D.W. Laurence, C.J. Ross, K.E. Kramer, A.R. Babu, E.L. Johnson, M.-C. Hsu, A.
Aggarwal, A. Mir, H.M. Burkhart, R.A. Towner, R. Baumwart, Y. Wu, Mechanics of the
Tricuspid Valve—From Clinical Diagnosis/Treatment, In-Vivo and In-Vitro Investigations, to
Patient-Specific Biomechanical Modeling, Bioengineering 6 (2019) 47.
https://doi.org/10.3390/bioengineering6020047.
[7] M. Taramasso, H. Vanermen, F. Maisano, A. Guidotti, G. La Canna, O. Alfieri, The Growing
Clinical Importance of Secondary Tricuspid Regurgitation, Journal of the American College
of Cardiology 59 (2012) 703–710. https://doi.org/10.1016/j.jacc.2011.09.069.
[8] J.H. Rogers, S.F. Bolling, The Tricuspid Valve: Current Perspective and Evolving
Management of Tricuspid Regurgitation, Circulation 119 (2009) 2718–2725.
https://doi.org/10.1161/CIRCULATIONAHA.108.842773.
[9] M. Pagnesi, C. Montalto, A. Mangieri, E. Agricola, R. Puri, M. Chiarito, M.B. Ancona, D.
Regazzoli, L. Testa, M. De Bonis, N.E. Moat, J. Rodés-Cabau, A. Colombo, A. Latib, Tricuspid
annuloplasty versus a conservative approach in patients with functional tricuspid
regurgitation undergoing left-sided heart valve surgery: A study-level meta-analysis,
International Journal of Cardiology 240 (2017) 138–144.
https://doi.org/10.1016/j.ijcard.2017.05.014.
[10] K.J. Grande-Allen, A.G. Borowski, R.W. Troughton, P.L. Houghtaling, N.R. DiPaola, C.S.
Moravec, I. Vesely, B.P. Griffin, Apparently normal mitral valves in patients with heart
failure demonstrate biochemical and structural derangements, Journal of the American
College of Cardiology 45 (2005) 54–61. https://doi.org/10.1016/j.jacc.2004.06.079.
[11] M. Chaput, M.D. Handschumacher, F. Tournoux, L. Hua, J.L. Guerrero, G.J. Vlahakes, R.A.
Levine, Mitral Leaflet Adaptation to Ventricular Remodeling: Occurrence and Adequacy in
Patients With Functional Mitral Regurgitation, Circulation 118 (2008) 845–852.
https://doi.org/10.1161/CIRCULATIONAHA.107.749440.
[12] A. Itoh, G. Krishnamurthy, J.C. Swanson, D.B. Ennis, W. Bothe, E. Kuhl, M. Karlsson, L.R.
Davis, D.C. Miller, N.B. Ingels, Active stiffening of mitral valve leaflets in the beating heart,
American Journal of Physiology-Heart and Circulatory Physiology 296 (2009) H1766–
H1773. https://doi.org/10.1152/ajpheart.00120.2009.
[13] K.J. Grande-Allen, J.E. Barber, K.M. Klatka, P.L. Houghtaling, I. Vesely, C.S. Moravec, P.M.
McCarthy, Mitral valve stiffening in end-stage heart failure: Evidence of an organic
contribution to functional mitral regurgitation, The Journal of Thoracic and Cardiovascular
Surgery 130 (2005) 783–790. https://doi.org/10.1016/j.jtcvs.2005.04.019.
[14] M.K. Rausch, F.A. Tibayan, D. Craig Miller, E. Kuhl, Evidence of adaptive mitral leaflet
growth, Journal of the Mechanical Behavior of Biomedical Materials 15 (2012) 208–217.
https://doi.org/10.1016/j.jmbbm.2012.07.001.
[15] T.A. Timek, D.T. Lai, P. Dagum, D. Liang, G.T. Daughters, N.B. Ingels, D.C. Miller, Mitral
Leaflet Remodeling in Dilated Cardiomyopathy, Circulation 114 (2006).
https://doi.org/10.1161/CIRCULATIONAHA.105.000554.
[16] J. Afilalo, J. Grapsa, P. Nihoyannopoulos, J. Beaudoin, J.S.R. Gibbs, R.N. Channick, D.
Langleben, L.G. Rudski, L. Hua, M.D. Handschumacher, M.H. Picard, R.A. Levine, Leaflet
Area as a Determinant of Tricuspid Regurgitation Severity in Patients With Pulmonary

Hypertension, Circ: Cardiovascular Imaging 8 (2015) e002714.
https://doi.org/10.1161/CIRCIMAGING.114.002714.
[17] W.D. Meador, M. Mathur, G.P. Sugerman, M. Malinowski, T. Jazwiec, X. Wang, C.M.
Lacerda, T.A. Timek, M.K. Rausch, The tricuspid valve also maladapts as shown in sheep
with biventricular heart failure, eLife 9 (2020) e63855.
https://doi.org/10.7554/eLife.63855.
[18] A. Iwasieczko, M. Gaddam, B. Gaweda, A. Goodyke, M. Mathur, C.-Y. Lin, J. Zagorski, M.
Solarewicz, S. Cohle, M. Rausch, T.A. Timek, Valvular complex and tissue remodelling in
ovine functional tricuspid regurgitation, European Journal of Cardio-Thoracic Surgery 63
(2023) ezad115. https://doi.org/10.1093/ejcts/ezad115.
[19] M. Mathur, M. Malinowski, T. Jazwiec, T.A. Timek, M.K. Rausch, Leaflet remodeling
reduces tricuspid valve function in a computational model, Journal of the Mechanical
Behavior of Biomedical Materials 152 (2024) 106453.
https://doi.org/10.1016/j.jmbbm.2024.106453.
[20] M. Mathur, T. Jazwiec, W.D. Meador, M. Malinowski, M. Goehler, H. Ferguson, T.A. Timek,
M.K. Rausch, Tricuspid valve leaflet strains in the beating ovine heart, Biomech Model
Mechanobiol 18 (2019) 1351–1361. https://doi.org/10.1007/s10237-019-01148-y.
[21] W.D. Meador, M. Mathur, G.P. Sugerman, T. Jazwiec, M. Malinowski, M.R. Bersi, T.A.
Timek, M.K. Rausch, A detailed mechanical and microstructural analysis of ovine tricuspid
valve leaflets, Acta Biomaterialia 102 (2020) 100–113.
https://doi.org/10.1016/j.actbio.2019.11.039.
[22] M. Malinowski, A.G. Proudfoot, D. Langholz, L. Eberhart, M. Brown, H. Schubert, J.
Wodarek, T.A. Timek, Large animal model of functional tricuspid regurgitation in pacing
induced end-stage heart failure, Interactive CardioVascular and Thoracic Surgery 24 (2017)
905–910. https://doi.org/10.1093/icvts/ivx012.
[23] C.J. Kostelnik, K.J. Crouse, J.D. Goldsmith, J.F. Eberth, Impact of cryopreservation on
elastomuscular artery mechanics, Journal of the Mechanical Behavior of Biomedical
Materials 154 (2024) 106503. https://doi.org/10.1016/j.jmbbm.2024.106503.
[24] G.A. Duginski, C.J. Ross, D.W. Laurence, C.H. Johns, C.-H. Lee, An investigation of the effect
of freezing storage on the biaxial mechanical properties of excised porcine tricuspid valve
anterior leaflets, Journal of the Mechanical Behavior of Biomedical Materials 101 (2020)
103438. https://doi.org/10.1016/j.jmbbm.2019.103438.
[25] K.J. Smith, M. Mathur, W.D. Meador, B. Phillips-Garcia, G.P. Sugerman, A.K. Menta, T.
Jazwiec, M. Malinowski, T.A. Timek, M.K. Rausch, Tricuspid Chordae Tendineae Mechanics:
Insertion Site, Leaflet, and Size-Specific Analysis and Constitutive Modelling, Exp Mech 61
(2021) 19–29. https://doi.org/10.1007/s11340-020-00594-5.
[26] W.D. Meador, J. Zhou, M. Malinowski, T. Jazwiec, S. Calve, T.A. Timek, M.K. Rausch, The
effects of a simple optical clearing protocol on the mechanics of collagenous soft tissue,
Journal of Biomechanics 122 (2021) 110413.
https://doi.org/10.1016/j.jbiomech.2021.110413.
[27] A.J. Schriefl, A.J. Reinisch, S. Sankaran, D.M. Pierce, G.A. Holzapfel, Quantitative
assessment of collagen fibre orientations from two-dimensional images of soft biological
tissues, J. R. Soc. Interface. 9 (2012) 3081–3093. https://doi.org/10.1098/rsif.2012.0339.

[28] M.S. Sacks, Biaxial Mechanical Evaluation of Planar Biological Materials, in: S.C. Cowin, J.D.
Humphrey (Eds.), Cardiovascular Soft Tissue Mechanics, Kluwer Academic Publishers,
Dordrecht, 2004: pp. 199–246. https://doi.org/10.1007/0-306-48389-0_7.
[29] K.M. Kodigepalli, K. Thatcher, T. West, D.P. Howsmon, F.J. Schoen, M.S. Sacks, C.K. Breuer,
J. Lincoln, Biology and Biomechanics of the Heart Valve Extracellular Matrix, JCDD 7 (2020)
57. https://doi.org/10.3390/jcdd7040057.
[30] L.T. Hudson, S.V. Jett, K.E. Kramer, D.W. Laurence, C.J. Ross, R.A. Towner, R. Baumwart,
K.M. Lim, A. Mir, H.M. Burkhart, Y. Wu, C.-H. Lee, A Pilot Study on Linking Tissue
Mechanics with Load-Dependent Collagen Microstructures in Porcine Tricuspid Valve
Leaflets, Bioengineering 7 (2020) 60. https://doi.org/10.3390/bioengineering7020060.
[31] D. Laurence, C. Ross, S. Jett, C. Johns, A. Echols, R. Baumwart, R. Towner, J. Liao, P. Bajona,
Y. Wu, C.-H. Lee, An investigation of regional variations in the biaxial mechanical
properties and stress relaxation behaviors of porcine atrioventricular heart valve leaflets,
Journal of Biomechanics 83 (2019) 16–27.
https://doi.org/10.1016/j.jbiomech.2018.11.015.
[32] T.L. Blevins, S.B. Peterson, E.L. Lee, A.M. Bailey, J.D. Frederick, T.N. Huynh, V. Gupta, K.J.
Grande-Allen, Mitral Valvular Interstitial Cells Demonstrate Regional, Adhesional, and
Synthetic Heterogeneity, Cells Tissues Organs 187 (2008) 113–122.
https://doi.org/10.1159/000108582.
[33] H. Ma, A.R. Killaars, F.W. DelRio, C. Yang, K.S. Anseth, Myofibroblastic activation of
valvular interstitial cells is modulated by spatial variations in matrix elasticity and its
organization, Biomaterials 131 (2017) 131–144.
https://doi.org/10.1016/j.biomaterials.2017.03.040.
[34] G.A. Walker, K.S. Masters, D.N. Shah, K.S. Anseth, L.A. Leinwand, Valvular Myofibroblast
Activation by Transforming Growth Factor-β: Implications for Pathological Extracellular
Matrix Remodeling in Heart Valve Disease, Circulation Research 95 (2004) 253–260.
https://doi.org/10.1161/01.RES.0000136520.07995.aa.
[35] Y. Li, K.O. Lui, B. Zhou, Reassessing endothelial-to-mesenchymal transition in
cardiovascular diseases, Nat Rev Cardiol 15 (2018) 445–456.
https://doi.org/10.1038/s41569-018-0023-y.
[36] K.J. Grande-Allen, J. Liao, The Heterogeneous Biomechanics and Mechanobiology of the
Mitral Valve: Implications for Tissue Engineering, Curr Cardiol Rep 13 (2011) 113–120.
https://doi.org/10.1007/s11886-010-0161-2.
[37] J.P. Dal-Bianco, E. Aikawa, J. Bischoff, J.L. Guerrero, M.D. Handschumacher, S. Sullivan, B.
Johnson, J.S. Titus, Y. Iwamoto, J. Wylie-Sears, R.A. Levine, A. Carpentier, Active
Adaptation of the Tethered Mitral Valve: Insights Into a Compensatory Mechanism for
Functional Mitral Regurgitation, Circulation 120 (2009) 334–342.
https://doi.org/10.1161/CIRCULATIONAHA.108.846782.
[38] S. Korossis, Structure-Function Relationship of Heart Valves in Health and Disease, in: K.
Kirali (Ed.), Structural Insufficiency Anomalies in Cardiac Valves, InTech, 2018.
https://doi.org/10.5772/intechopen.78280.
[39] T. Jazwiec, M.J. Malinowski, H. Ferguson, J. Parker, M. Mathur, M.K. Rausch, T.A. Timek,
Tricuspid Valve Anterior Leaflet Strains in Ovine Functional Tricuspid Regurgitation,

Seminars in Thoracic and Cardiovascular Surgery 33 (2021) 356–364.
https://doi.org/10.1053/j.semtcvs.2020.09.012.
[40] A. Iwasieczko, T. Jazwiec, M. Gaddam, B. Gaweda, M. Piekarska, M. Solarewicz, M.K.
Rausch, T.A. Timek, Septal annular dilation in chronic ovine functional tricuspid
regurgitation, The Journal of Thoracic and Cardiovascular Surgery 166 (2023) e393–e403.
https://doi.org/10.1016/j.jtcvs.2023.04.003.
[41] C.E. Haese, M. Mathur, C.-Y. Lin, M. Malinowski, T.A. Timek, M.K. Rausch, Impact of
tricuspid annuloplasty device shape and size on valve mechanics—a computational study,
JTCVS Open 17 (2024) 111–120. https://doi.org/10.1016/j.xjon.2023.11.002.
[42] K.E. Kramer, C.J. Ross, D.W. Laurence, A.R. Babu, Y. Wu, R.A. Towner, A. Mir, H.M.
Burkhart, G.A. Holzapfel, C.-H. Lee, An investigation of layer-specific tissue biomechanics
of porcine atrioventricular valve anterior leaflets, Acta Biomaterialia 96 (2019) 368–384.
https://doi.org/10.1016/j.actbio.2019.06.049.
600

FIGURES

602   **Figure 1**

Figure 2

Figure 3

611   **Figure 4**

Figure 5

617   **Figure 6**

Figure 7

Figure 8

627 **TABLES**

628 **Table 1**

Parameter	CTL	TIC (baseline)	TIC (terminal)
Echocardiographic data			
TR grade	1.0 (0.0)	0.0 (0.0)	2.0 (1.0) * †
RV EF, %	67.0 ± 8.6	58.8 (20.5)	48.4 ± 12.9 * †
RV FAC, %	52.1 ± 6.0	53.9 ± 8.0	37.4 ± 8.7 * †
RV IDd, cm	2.5 ± 0.4	2.6 (0.6)	2.9 ± 0.7 *
TV annulus dimension, cm	2.5 (0.3)	2.5 ± 0.4	3.2 ± 0.5 * †
MR grade	0.0 (0.0)	0.0 (0.0)	2.0 (1.0) * †
LV EF, %	55.8 ± 4.4	61.3 ± 6.2	30.0 (11.2) * †
LV IDd, cm	4.2 ± 0.3	3.7 ± 0.6	4.6 ± 0.4 * †
Hemodynamic data			
HR, bpm	108 (11)	-	126 ± 23 *
RV P_{Max}, mmHg	28.8 (11.4)	-	44.0 ± 11.5 *
RV P_{ES}, mmHg	22.3 ± 5.6	-	35.7 ± 11.0 *
RA P_{Mean}, mmHg	10.5 (3.2)	-	10.6 (5.6)
LV P_{Max}, mmHg	96.1 ± 9.7	-	92.9 ± 13.8
LV P_{ES}, mmHg	63.1 ± 17.8	-	64.4 ± 13.6

Values are mean ± standard deviation or median (interquartile range).
CTL = control, EF = ejection fraction, ES = end systolic, FAC = fractional area change, HR = heart rate, IDd = inner dimension at diastole, LV = left ventricular, Max = maximal, MR = mitral regurgitation, P = pressure, RA = right atrial, RV = right ventricular, TIC = tachycardia-induced cardiomyopathy, TR = tricuspid regurgitation, TV = tricuspid valve.
* $p < 0.05$ vs. CTL, † $p < 0.05$ vs TIC (baseline)

 CAPTIONS

Figure 1: Tricuspid septal and anterior leaflet area increases in sheep with tachycardia-induced

cardiomyopathy (TIC). (a) Sheep tricuspid valve image on a calibrated grid (1 cm scale).

Measured areas (black line) of septal, anterior, and posterior leaflets are shown. Leaflet areas

were measured between commissures: posterior-septal (PS) to antero-septal (AS) for septal, AS

to antero-posterior (AP) for anterior, and AP to PS for posterior. (b-d) Leaflet area comparisons

between control (CTL, blue, n = 12) and TIC (red, n = 29) for septal, anterior, and posterior

leaflets. The (+) indicates that the anterior leaflet data was previously published and can be

found in [17]. Statistical significance between animal group (CTL vs. TIC) is denoted by (**) for p

< 0.01 using one-tailed pairwise tests. Statistical significance between leaflets is denoted by (‡)

for septal vs. anterior, (#) for anterior vs. posterior, and (§) for septal vs. posterior, all at p ≤

0.0001, using two-tailed pairwise tests.

Figure 2: Tricuspid anterior and posterior leaflet thickness increases in sheep with tachycardia-

induced cardiomyopathy (TIC). Representative images of (a) septal, (b) anterior, and (c)

posterior leaflet strips. Averaged thickness comparisons between control (CTL, blue, n = 8-10)

and TIC (red, n = 11-12) (d) septal, (e) anterior, and (f) posterior leaflets. The (+) indicates that

the anterior leaflet data was previously published and can be found in [17]. Statistical

significance between animal group (CTL vs. TIC) is denoted by (*) for p < 0.05 using one-tailed

pairwise tests.

Figure 3: Tachycardia-induced cardiomyopathy (TIC) anterior leaflets are stiffer at low stretches

in the circumferential (circ) directions. Control (CTL, blue, n = 10 for each leaflet) and TIC (red, n

= 22 for each leaflet) membrane tension (Mem. Tens.) vs stretch average curves (solid) with

standard deviation (shaded) in the circumferential and radial (rad) directions for (a) posterior

and (d) septal leaflets. Comparisons of the (b & e) stiffness at large stretches (calf stiffness) in

the circumferential and radial directions for the posterior and septal leaflets, respectively.

Comparisons of the (c & f) stiffness at small stretches (toe stiffness) in the circumferential and

radial directions for the posterior and septal leaflets, respectively. The (+) indicates that the

anterior leaflet data was previously published and can be found in [17]. Statistical significance

between animal group (CTL vs. TIC) is denoted by (*) using one-tailed pairwise tests at $p < 0.05$,

and statistical significance between septal and anterior leaflets is denoted by (‡) using two-

tailed pairwise tests at $p < 0.05$.

Figure 4: Tachycardia-induced cardiomyopathy (TIC) septal leaflets exhibit increased radial

transition stretch and decreased anisotropy. Comparisons of the transition stretch in

circumferential and radial directions between control (CTL, blue, n = 10) and TIC (red, n = 22) (a)

septal, (b) anterior, and (c) posterior leaflets. Comparisons of the degree of anisotropy between

CTL (blue, n = 10) and TIC (red, n = 22) (d) septal, (e) anterior, and (f) posterior leaflets. The (+)

indicates that the anterior leaflet data was previously published and can be found in [17].

Statistical significance between animal group (CTL vs. TIC) is denoted by (*) for $p < 0.05$ and (**)

for $p < 0.01$ using one-tailed pairwise tests. Statistical significance between leaflets is denoted

by (‡) for septal vs. anterior, (#) for anterior vs. posterior, and (§) for septal vs. posterior, all at p

< 0.05, using two-tailed pairwise tests.

Figure 5: Changes in remodeling cellular markers in tachycardia-induced cardiomyopathy (TIC)

septal leaflets. (a-d) Heat maps (left) showing regional expression of (a) alpha-smooth muscle

actin (α-SMA), (b) Ki67, (c) matrix metalloproteinase 13 (MMP13), (d) transforming growth

factor beta (TGF-β1). The heatmaps are separated into three equidistant regions along the

radial axis (near-annulus (NA), belly (Bel), free edge (FE)) and 10 thickness regions from the

atrial surface (Atr.) to the ventricular surface (Ven.). Fold change (FC) between control (CTL, n =

6) and TIC (n = 6) was determined by the ratio of positively stained pixel percentage between

TIC and CTL. Color map indicates the logarithm base 2 of the fold change, interpreted as

(positive, red): TIC expression is higher than CTL, (0, white): TIC and CTL expression are

approximately equal, and (negative, blue): TIC expression is less than CTL. Representative

images of CTL (middle) and TIC (right) posterior leaflets are shown with the atrial surface

upward. The inset leaflet schematic depicts the longitudinal region from which the

representative image was taken. Scale bars = 500 μm.

Figure 6: Changes in remodeling cellular markers in tachycardia-induced cardiomyopathy (TIC)

anterior leaflets. The (+) indicates that the anterior leaflet data was previously published and

can be found in [17]. (a-d) Heat maps (left) showing regional expression of (a) alpha-smooth

muscle actin (α-SMA), (b) Ki67, (c) matrix metalloproteinase 13 (MMP13), (d) transforming

growth factor beta (TGF-β1). The heatmaps are separated into three equidistant regions along

the radial axis (near-annulus (NA), belly (Bel), free edge (FE)) and 10 thickness regions from the

atrial surface (Atr.) to the ventricular surface (Ven.). Fold change (FC) between control (CTL, n =

6) and TIC (n = 6) was determined by the ratio of positively stained pixel percentage between

TIC and CTL. Color map indicates the logarithm base 2 of the fold change, interpreted as

(positive, red): TIC expression is higher than CTL, (0, white): TIC and CTL expression are

approximately equal, and (negative, blue): TIC expression is less than CTL. Representative

images of CTL (middle) and TIC (right) posterior leaflets are shown with the atrial surface

upward. The inset leaflet schematic depicts the longitudinal region from which the

representative image was taken. Scale bars = 500 μm.

Figure 7: Changes in remodeling cellular markers in tachycardia-induced cardiomyopathy (TIC)

posterior leaflets. (a-d) Heat maps (left) showing regional expression of (a) alpha-smooth

muscle actin (α-SMA), (b) Ki67, (c) matrix metalloproteinase 13 (MMP13), (d) transforming

growth factor beta (TGF-β1). The heatmaps are separated into three equidistant regions along

the radial axis (near-annulus (NA), belly (Bel), free edge (FE)) and 10 thickness regions from the

atrial surface (Atr.) to the ventricular surface (Ven.). Fold change (FC) between control (CTL, n =

6) and TIC (n = 6) was determined by the ratio of positively stained pixel percentage between

TIC and CTL. Color map indicates the logarithm base 2 of the fold change, interpreted as

(positive, red): TIC expression is higher than CTL, (0, white): TIC and CTL expression are

approximately equal, and (negative, blue): TIC expression is less than CTL. Representative

images of CTL (middle) and TIC (right) posterior leaflets are shown with the atrial surface

upward. The inset leaflet schematic depicts the longitudinal region from which the

representative image was taken. Scale bars = 500 μm.

Figure 8: Through-depth collagen microstructure reveals similar mean orientations and regional

concentration changed throughout depth between control (CTL) and tachycardia-induced

cardiomyopathy (TIC) tricuspid leaflets. Heat map visualizations of von Mises probability

distribution functions (VM PDF) fit to collagen fiber orientation histograms of two-photon

acquired images throughout the entire depth (0% - Atrial surface, 100% - Ventricular surface) of

averaged CTL (blue, n = 9 for each leaflet) and TIC (red, n = 9 for each leaflet) (a) septal, (c)

anterior, (e) posterior leaflets. Comparisons of VM concentration parameter, κ, across depth

regions D1, D2, and D3 for the (b) septal, (d) anterior, and (f) posterior leaflets. The (+) indicates

that the anterior leaflet data was previously published and can be found in [17]. Statistical

significance between animal group (CTL vs. TIC) is denoted by (*) for $p < 0.05$ and (**) for $p <$

0.01 using one-tailed pairwise tests. Statistical significance between leaflets is denoted by (‡)

for septal vs. anterior, (#) for anterior vs. posterior, and (§) for septal vs. posterior, all at $p <$

0.05, using two-tailed pairwise tests.

Table 1: Echocardiographic and hemodynamic data of animal model.

a Septal

b Anterior

c Posterior

d
Septal Thickness (μm)

CTL   TIC

e
Anterior⁺ Thickness (μm)

● CTL ● TIC

CTL   TIC

f
Posterior Thickness (μm)

CTL   TIC