TITLE

Proof of concept study deploying CRISPR inhibition and activation opens new avenues for

systematic biological exploration in zebrafish.

AUTHORS

Nelson B. Barrientos[1], Elyse A. Shoppell[1], Rachel J. Boyd[1], Valeria C. Culotta[2], Andrew S.

McCallion[1,3*]

AFFILIATIONS

[1]McKusick-Nathans Department of Genetic Medicine, Johns Hopkins University School of

Medicine, Baltimore, MD 21205, USA

[2]Department of Biochemistry and Molecular Biology, Johns Hopkins University Bloomberg School

of Public Health, Baltimore, MD 21205, USA

[3]Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD 21205,

USA

[*]Corresponding author

Corresponding author's email: andy@jhmi.edu

ABSTRACT

The application of CRISPR interference (CRISPRi) and CRISPR activation (CRISPRa)

technologies in zebrafish has the potential to expand its capacity for the study of gene function

significantly. We report proof-of-principle data evaluating transient expression of a codon

optimized CRISPRi/a system for zebrafish across established pigmentary and growth

phenotypes. A codon-optimized and catalytically inactive *cas9* gene (*dcas9*) was cloned upstream

of codon-optimized Krüppel associated box (KRAB) and methyl-CpG binding protein 2 (MeCP2)

for CRISPRi, and VP64 for CRISPRa. To validate CRISPRi, we targeted key genes in melanocyte

differentiation (*sox10* and *mitfa)*; and melanin production (tyrosinase; *tyr*). Microinjection of

CRISPRi mRNA and single guide RNAs (sgRNAs) targeting the *tyr* promoter or 5'-UTR resulted

in larvae with hypopigmented epidermal melanocytes. Similarly, CRISPRi targeting of the *sox10*

or mitfa promoters results in hypopigmentation of epidermal melanocytes consistent with their

roles upstream of *tyr,* and the role of *sox10* in activation of *mitfa*. Finally, we tested both

CRISPRi/a to modulate a single gene to yield hypomorphic and hypermorphic effects, selecting

mrap2a as our target. This gene regulates energy homeostasis and somatic growth via inhibition

of the melanocortin 4 receptor gene (*mc4r*). We show that inactivating or activating *mrap2a* with

CRISPRi/a significantly decreases or increases larval body length, respectively. We demonstrate

that CRISPRi/a can modulate control of zebrafish gene expression, facilitating efficient assay of

candidate gene function and disease relevance.

KEYWORDS:

Zebrafish, CRISPR-activation, CRISPR-inhibition

INTRODUCTION

Zebrafish (*danio rerio*) are a powerful model system in which to study functional components of vertebrate genomes. Comparative genomics analyses have shown that 61.5% of mouse protein-coding genes and 71% of human protein-coding genes have at least one zebrafish orthologue (1,2). Additional features, including short generation time, high fecundity, external fertilization, and embryonic transparency, have distinguished zebrafish as a desirable model system in which to examine developmental processes. As such, zebrafish have been widely used to study the biological relevance of regulatory elements (3–6), and human disease-associated genes (7–14).

Emerging technologies for transgenesis and genome engineering, such as such as Gal4-UAS (15); Tol2-transposase transgenesis (3); Zinc Finger Nucleases (16); and transcription activator-like effectors (TALENs) (17) have been developed and rapidly implemented in zebrafish. Over the last decade, zebrafish have also been shown to be amenable to CRISPR-Cas-based genomic engineering applications, including targeted mutagenesis, base editing, and generation of knock-in and knockout models (18–22). More recently, the emergence of CRISPR interference (CRISPRi) and activation (CRISPRa) have presented additional opportunities to explore functional components of vertebrate genomes. CRISPRi/a involves the fusion of transcriptional effectors to a catalytically inactive, or "dead" Cas9 (dCas9). CRISPRi employs a transcriptional repressor, such as a Krüppel associated box (KRAB) domain and methyl-CpG binding protein 2 (MeCP2) (23,24), while in the case of CRISPRa, a transcriptional activator, such as VP64, is fused to the dCas9 (25). To date, the CRISPRi/a systems have seen limited application in zebrafish (26), whereas they have been more rapidly employed in mammalian organisms to target human disease-relevant genes (27–29).

Here, we report initial experiments in the development and evaluation of zebrafish codon-optimized zdCas9-KRAB-MeCP2 and zdCas9-VP64 CRISPR systems. We present a systematic

evaluation of these modified proteins in functional proof-of-concept experiments by targeting two

easily scored phenotypes: zebrafish pigmentation and somatic growth. In addition to providing

visual confirmation of the efficacy of both CRISPRi and CRISPRa, these two model phenotypes

and their corresponding pathways are well-characterized and are involved in human disease.

Specifically, we chose to test our CRISPRi system by targeting genes involved in

melanocyte differentiation and pigmentation, *sox10 (SRY*-box transcription factor 10*)*, *mitfa*

(melanocyte inducing transcription factor a), and the gene for the rate-determining enzyme in

pigmentation, *tyr (*tyrosinase*)*. *sox10* directly activates the *mitfa* promoter to stimulate downstream

genes involved in melanocyte differentiation (33,34); this leads to the activation of *tyr*, at

approximately 16 hours post fertilization (16 hpf) to induce melanin production by 24 hpf (30–32).

In the developing zebrafish embryo, due to whole-genome duplication, *mitfa* is co-expressed with

mitfb in the retinal pigment epithelium, but only *mitfa* is expressed in neural crest cells, which

differentiate to give rise to epidermal melanocytes (31,33,34).

We hypothesize that independently targeting each of these genes with functional CRISPRi

will yield a visible reduction in zebrafish pigmentation. In the case of *sox10,* we expect to see

marked reduction in pigmentation globally due to its role in melanocyte cell fate and activation of

mitfa and *tyr* to promote melanin production in zebrafish and other vertebrates (35,36). Due to its

role in multiple stages of melanocyte differentiation and its function as a transcription factor,

silencing *mitfa* is also expected to result in phenotype of epidermal hypopigmentation (37,38).

Lastly, we hypothesize that targeting the rate-determining enzyme in melanin pigment formation

(*tyr)* will result in hypopigmentation of melanocytes throughout the zebrafish body (30,35,39,40).

Zebrafish melanocytes have been extensively studied by our group and others (4,6,41,42), and

targeting genes in this pathway is highly relevant to the study of human disease, including

melanoma (43–45), Waardenburg Syndrome (46,47), microphthalmia, and Hirschsprung's

disease (48,49).

Expanding from this initial study, we sought to apply both CRISPRi and CRISPRa to

mrap2a, a gene with known inhibitory activity to regulate somatic growth in the zebrafish embryo

(50). *mrap2a* encodes the melanocortin receptor accessory protein 2a, which has been found to

inhibit the transcription of the *mc4r* (melanocortin-4 receptor) (50). We hypothesize that targeting

the *mrap2a* promoter with CRISPRa will increase *mrap2a* expression and therefore amplify the

inhibition of *mc4r* and promote visible overgrowth phenotypes in zebrafish larvae. This is

supported by evidence in human patients, wherein autosomal dominant loss of function mutations

in *MCR4* results in severe childhood-onset obesity (51,52). By contrast, we hypothesize that

effective CRISPRi targeting of *mrap2a* will result in decreased *mc4r* inhibition and result in

undergrown zebrafish larvae, consistent with reported morpholino-based knock-down

experiments (50).

Ultimately, we outline the development of zebrafish-optimized zdCas9-KRAB-MeCP2 and

zdCas9-VP64 CRISPR constructs. We establish their utility for efficient screening of candidate

gene function and disease relevance. Ongoing work in the lab will establish stable transgenic

lines expressing these proteins to facilitate rapid and systematic screening of gain and loss of

gene function by leveraging custom guide RNAs (gRNAs) to target promoters, 5'-UTR, or cell-

dependent cis-regulatory elements of selected candidate genes.

RESULTS

CRISPRi modulates tyrosinase expression and its effect is observable between the prim

20-25 stages.

To determine the functional capacity of zebrafish codon-optimized CRISPRi (Fig. 1A),

one-cell stage embryos of wild-type AB (WT AB) zebrafish were injected with CRISPRi mRNA

and target sgRNAs. As part of a series of proof-of-concept experiments, we designed sgRNAs

targeting the promoter region and 5'-UTR of the tyrosinase (*tyr*) gene, predicting that inhibition

would lead to diminished larval pigmentation, similar to established loss of function studies at this

locus (30,32,40).

Initially, sgRNAs were designed to target the *tyr* promoter or 5'-UTR (Methods) and one-

cell embryos (n ≥ 200) were injected with CRISPRi (100-300 ng/µl, final concentration) and a

cocktail of all five sgRNAs (150 ng/µl/each, final concentration). Zebrafish larvae assayed at the

prim 20-25 stages (33-36 hpf) displayed marked reduction in pigmentation of melanocytes (Fig.

1B), consistent with phenotypes arising from loss of function alleles in zebrafish (30) and other

vertebrates (39). By contrast, larvae receiving only the CRISPRi mRNA (n ≥ 200) or uninjected

(WT AB) controls (n ≥ 200), displayed no diminution in melanization at these stages. As expected,

the observed suppression of *tyr* dissipated over time, and by the pec-fin stage (60 hpf), reduction

in melanization was less noticeable in larvae receiving CRISPRi plus the cocktail of all sgRNAs

(Fig. 1B).

We measured the percentage of pigmentated area, imaged laterally, from a sample (n =

10) of larvae from each condition, at the prim 20-25 stage, and found a statistically significant

reduction of pigmentation between uninjected control (WT AB) and larvae injected with CRISPRi

plus all *tyr* sgRNAs (*p-value* = 1.263e-06, Fig. 1C, Supplemental Table 4a). We also observed a

statistically significant reduction of pigmentation between injected control (embryos injected with

CRISPRi mRNA only) and larvae injected with CRISPRi plus all *tyr* sgRNAs (*p-value* = 1.058e-

07, Fig. 1C, Supplemental Table 4a). We did not observe a statistically significant difference

between injected and uninjected controls (*p-value* = 0.5018, Fig. 1C, Supplemental Table 4a). We

subsequently performed the same pigmentation analysis at the pec-fin stage and observed the

same trend of hypopigmentation in CRISPRi plus all *tyr* sgRNAs (Fig. 1B-C, Supplemental Table

4b). However, the results were not statistically significant across conditions (*p-values* = 0.1226,

0.1418, and 0.7671, respectively, Fig. 1C, Supplemental Table 4b), confirming that the effects of

CRISPRi mRNA injection diminish over time.

We tested if delivery of each individual *tyr* sgRNAs could similarly silence transcription of

tyr and reduce melanin production. One of the five sgRNAs that we tested, sgRNA4 targeting the

promoter, efficiently modulated *tyr* expression (Supplemental Fig. 1), resulting in

hypopigmentation within prim 20-25 (33-36 hpf). We used this sgRNA to assess the optimal

concentration of CRISPRi and injected larvae each with 100 ng/µl, 300 ng/µl, and 500 ng/µl of

CRISPRi along with sgRNA4 (150 ng/µl). Our observations suggest no advantage to injecting

more than 300 ng/µl (final concentration) of CRISPRi mRNA (Supplemental Fig. 1). These results

demonstrate the effectiveness in modulating *tyr* by employing our zebrafish codon-optimized

CRISPRi with individual or pooled sgRNAs to target a gene's promoter. To build on these

observations, we sought to confirm the hypopigmentation phenotype by targeting other key

pigment genes that act upstream and are known to regulate *tyr*.

161   CRISPRi targeting of *sox10* impedes epidermal pigmentation zebrafish

Having established that *tyr* can be modulated via CRISPRi, we sought to evaluate a

broader applicability of our strategy, evaluating the effect of modulating genes upstream from *tyr*

in melanocyte differentiation, for which we predict overlapping phenotypic outcomes. We first

investigated whether *sox10*, the gene that encodes for a transcription factor known to be key to

promote melanin production in zebrafish and other vertebrates (35,36) could be silenced via

CRISPRi. We designed sgRNAs to target the promoter of the *sox10* gene, predicting that silencing

it would lead to downregulation of downstream pigment genes, *mitfa* and *tyr*, and subsequent

hypopigmentation in the zebrafish.

Four sgRNAs were designed to target the *sox10* promoter (Methods). Zebrafish larvae (n

≥ 200) injected with CRISPRi (100-300 ng/µl, final concentration) plus a cocktail of all *sox10*

sgRNAs (150 ng/µl/each, final concentration) displayed marked reduction in pigmentation of

epidermal melanocytes, assayed at prim 20-25 (33-36 hpf), consistent with reduction of

pigmentation when targeting downstream gene *tyr* (Fig. 2A, Supplemental Fig. 2). Furthermore,

we observed that sgRNA3 was effective at reducing pigmentation when injected individually with

CRISPRi (Fig. 2A, Supplemental Fig. 2). We measured the percentage of pigmented area,

imaged laterally at the prim 20-25 stage, from a sample (n = 10) of larvae from each condition

and found a statistically significant reduction of pigmentation between uninjected control (WT AB)

and larvae injected with CRISPRi plus all *sox10* sgRNAs and CRISPRi plus *sox10* sgRNA3 (*p-*

value = 7.389e-07, 2.724e-06, respectively, Fig. 2B, Supplemental Table 5a). We also observed

a statistically significant reduction of pigmentation between injected control (embryos injected with

CRISPRi mRNA only) and larvae injected with CRISPRi plus all *sox10* sgRNAs and CRISPRi

plus *sox10* sgRNA3 (*p-value* = 8.704e-09, 3.243e-06, respectively, Fig. 2B, Supplemental Table

5a). Further, we did not observe a statistically significant reduction in pigmentation between larvae

receiving only the CRISPRi mRNA (n ≥ 200) and uninjected (WT AB) controls (n ≥ 200) (*p-value*

= 0.097, Fig. 2B, Supplemental Table 5a). The effects of CRISPRi dissipated by the pec-fin stage

(60 hpf) as the pigmentation patterns between injected and uninjected were indistinguishable (Fig.

2A, Supplemental Table 5b). When measuring the percentage of pigmented area at this stage,

we only observed at statistically significant reduction in pigmentation between WT AB and

CRISPRi plus all *sox10* sgRNAs (*p-value* = 0.019, Fig. 2B, Supplemental Table 5b). We did not

observe a statistically significant reduction in pigmentation across the other conditions (*p-value* =

0.481, 0.090, 0.999, 0.454, and 0.102, respectively, Fig. 2B, Supplemental Table 5b), which

confirms our observation that as larvae grow, the effects of CRISPRi mRNA injections dissipate.

It is noteworthy to mention that larvae injected with CRISPRi plus sgRNAs appear to be slightly

developmentally delayed as compared to controls, which may be the result of injection variability,

nucleic acid toxicity, or off-target effects. This is also consistent with an observed reduction in

retinal pigment epithelium (RPE) pigmentation within Prim 20-25 in larvae injected with CRISPRi

plus sgRNAs as well as in those injected with CRISPRi alone, as compared to WT (*p-value* =

0.048, Supplemental Fig. 2B, Supplemental Table 6). Given that these effects are not consistent

with the established loss of function phenotypes for these genes (33) , we do not score or interpret

RPE phenotypes throughout. However, the overall results are in line with previous work that

demonstrated that *sox10* is important for melanocyte development and melanin production with

its role in activating *tyr*, the rate-determining enzyme in melanin production (30,35,39,40). To

exclude the possibility of off-target effects by *sox10* sgRNA, we employed sgRNA3 and standard

CRISPR/Cas9 to establish correct targeting of the *sox10* promoter. Employing a standard T7

endonuclease I (T7EI) assay (Methods) we confirm that sgRNA3 is effective at guiding Cas9 to

the *sox10* promoter to induce double-strand breaks. We collected various clutches (n = 50) of WT

AB, CRISPR/Cas9 only, and CRISPR/Cas9 plus sgRNA3 and extracted genomic DNA to amplify

the *sox10* promoter. PCR amplification (Oligonucleotide primers: Forward: 5'-

CTCCCAACTCTTATTAACGA-3'; Reverse: 5'- ACACAGAGTAAGCGATAGAG-3') of this locus

and subsequent treatment with T7EI resulted in heteroduplex formation, indicating effective

targeting of *sox10* (Supplemental Fig. 2C).

Effective CRISPRi modulation of *mitfa* limits epidermal melanin production.

Extending from our results modulating *sox10* and *tyr* expression, we next evaluated

whether targeting the gene encoding the transcription factor mitfa would also induce

hypopigmentation. The *mitfa* gene plays critical roles in pigmentation throughout vertebrates

(*MITF/Mitf*), activating *tyr* transcription during early development (35,59). We designed sgRNAs

to target the promoter and 5'-UTR regions of the *mitfa* gene, predicting that, as with known loss-

of-function (LoF) alleles (34,59), titrating activity of this transcription factor would also result in

reduced activation of *tyr* in epidermal melanocytes and thus lower melanization across the body.

Four sgRNAs were designed to target the *mitfa* promoter (Methods). As before, zebrafish

one-cell embryos (n ≥ 200) injected with CRISPRi (100-300 ng/µl, final concentration) and a

cocktail of all *mitfa* sgRNAs (150 ng/µl/each, final concentration) displayed marked reduction in

pigmentation of epidermal melanocytes, assayed at prim 20-25 (33-36 hpf) (Fig. 2C). This is

consistent with prior *mitfa* LoF studies (60) where failure to express *mitfa* prevents the production

of melanocyte precursors, leading to reduced global pigmentation in zebrafish larvae. In contrast,

no reduction in pigmentation was observed in larvae receiving only the CRISPRi mRNA (n ≥ 200)

or uninjected (WT AB) controls (n ≥ 200) (Fig. 2C). Additionally, a single *mitfa* sgRNA targeting

the promoter or 5'-UTR effectively silenced *mitfa,* resulting in marked reduction of melanin

production at prim 20-25 (33-36 hpf) (Supplemental Fig. 3). Silencing of *mitfa* dissipated over time

and by the pec-fin stage (60 hpf), melanization was similar between controls and injected larvae

(Fig. 2C, Supplemental Fig. 3). These results are consistent with the hypopigmentation phenotype

in epidermal melanocytes that is observed when modulating the *tyr* gene (Fig. 1B-C) and

transcription of *sox10* (Fig. 2A-B).

CRISPRi/a affects zebrafish length by modulating transcription of *mrap2a*

To build upon the successful demonstration of CRISPRi-mediated modulation of zebrafish

pigmentation, we set out to test our CRISPRa (Fig. 1A) system focusing on somatic size (length).

The benefit of this phenotype is that it provides the opportunity for evaluation of both reduction

and increase in gene function. We targeted the *mrap2a* gene which is known to regulate the

expression of the melanocortin 4 receptor (*mc4r*) gene by downregulating its transcription (50,62).

Both genes have been shown to be involved in energy homeostasis and somatic growth in mice

and zebrafish; and are thought to be involved in a significant portion of familial early-onset obesity

in humans (63). We predicted that targeting *mrap2a* with CRISPRa and further increasing its

transcription would result in a decrease in *mc4r* transcription, which would lead to an increase in

larval length. By contrast, downregulating *mrap2a* with CRISPRi would lead to an increase in

mc4r expression, resulting in larvae shorter in length (50,63).

While a range of loss-of-function experiments have been performed targeting *mrap2a*

(50,62,63), gain-of-function (GoF) experiments targeting *mrap2a* have not been explored. We

designed four sgRNAs to target the *mrap2a* promoter (Methods) with our CRISPRa system.

Zebrafish larvae (n ≥ 200) injected with CRISPRa (100-300 ng/µl, final concentration) and a

cocktail of all *mrap2a* sgRNAs (150 ng/µl/each, final concentration) displayed marked increase in

body length, assayed at the pec-fin stage (60 hpf) (Fig. 3A). Similarly, we tested if an individual

sgRNA could guide the dCas9-VP64 to modulate body length and found sgRNA3 capable of

inducing an observable phenotype (Fig. 3A-B). By contrast, no increase in larval length was

observed at the pec-fin stage (60 hpf) in larvae receiving only the CRISPRa mRNA (n ≥ 200) or

in uninjected (WT AB) controls (n ≥ 200). Moreover, the effects of CRISPRa were still noticeable

at the day 5 stage (120 hpf) and at the day 7 stage (168 hpf), as the length of these larvae was

still longer than control conditions (Fig. 3A, Supplemental Fig. 4). Furthermore, we measured the

length of larvae receiving CRISPRa plus sgRNA3 (n = 50), sgRNA3-only (n = 50), and uninjected

(WT AB) controls (n = 50) and performed a Welch two-sample t-test to analyze if the average

length between these conditions was statistically different (Fig. 3B, Supplemental Table 7). At the

pec-fin stage (60 hpf), the average length for these conditions was 3673.67 µm, 3468.86 µm, and

3466.28 µm (*p-value* = 8.58e-16; 1.83e-14, and 0.9181), respectively. The longer larval length

phenotype due to increased *mrap2a* transcription and decreased *mc4r* expression is consistent

with increased size and weight in mice that become obese due to *mc4r* haploinsufficiency (29),

and with morbid obesity cases in humans due to frameshift (loss of function) mutations in *mc4r*

(64).

We used the same sgRNAs to target and impede *mrap2a* and increase *mc4r* expression.

Zebrafish larvae (n ≥ 200) injected with CRISPRi (100-300 ng/µl, final concentration) plus all

sgRNAs (150 ng/µl/each, final concentration) displayed a decrease in larval length at the pec-fin

stage (60 hpf) (Fig. 3A, 3C), which is consistent with rescuing obese mice with *mc4r*

haploinsufficiency (29). Additionally, sgRNA3 was able to guide the CRISPRi to modulate

silencing of *mrap2a* individually and led to a decrease in larval length, observed at the pec-fin

stage (60 hpf) (Fig. 3A. 3C). By contrast, larval length observed at the pec-fin stage (60 hpf) for

larvae receiving only the CRISPRi mRNA (n ≥ 200) or in uninjected (WT AB) controls (n ≥ 200)

was higher than our experimental condition (Fig. 3A). Further, larvae injected with CRISPRi still

displayed a shorter length around the day 5 stage (120 hpf); however, their length was similar to

controls at the day 7 stage (168 hpf) (Fig. 3A, Supp Fig. 4). We also measured the larval length

for the conditions CRISPRi plus sgRNA3 (n = 50), sgRNA3-only (n = 50), and uninjected (WT AB)

controls (n = 50) at the pec-fin stage (60 hpf) and found their average length to be 3481.98 μm,

3779.96 μm, and 3748.82 μm, respectively (Fig. 3B, Supplemental Table 8). We analyzed the

statistical significance between these means with a Welch two-sample t-test and found the

difference in means between experimental and control conditions to be statistically significant (*p-*

value = 2.2e-16, 2.2e-16, and 0.0636). Although highly significant, the observed magnitude of

effect may potentially be further enhanced by simultaneous modulation of the agouti-related

protein (agrp), which Sebag et al. demonstrated acted in concert with *mrap2a* in antagonism of

mc4r (50).

DISCUSSION

CRISPRi/a has emerged as a powerful tool in genetic research, offering the ability to

modulate gene expression both for exploration of fundamental mechanisms and for potential

therapeutics. It offers the capacity to modulate gene expression without inducing genetic

alterations which makes CRISPRi/a valuable technology for the study of gene regulation and for

efficient exploration of both gain and loss of function phenotypes. Nevertheless, their development

and application in zebrafish have been slow over the past decade, with only one study describing

an effort to implement it (26), while many others apply CRISPR-based editing technologies

(18,19,21,53,54,65).

In this study, we chose to fuse a catalytically inactive Cas9 with the transcriptional

repressor KRAB-MeCP2. Our choice to use dCas9-KRAB-MeCP2 was guided by previous

studies that demonstrated its ability to induce gene silencing while outperforming dCas9-KRAB

systems (24,66) implemented in previous CRISPRi experiments in zebrafish (26). In contrast, for

our CRISPRa system, we fused the dCas9 with the transcriptional activator VP64. Comparisons

between different Cas9 activators have demonstrated the VP64 to effectively induce gene

activation (25). While other Cas9 activators such as VPR, SAM, and Suntag have been shown to

outperform VP64, these systems have not been tested in zebrafish and/or require special sgRNAs

to induce their effect. By using a catalytically inactive Cas9 fused with the transcriptional

repressors KRAB-MeCP2 or the transcriptional activator VP64, we have achieved transcriptional

control over key genes involved in melanocyte differentiation (*sox10* and *mitfa*), pigmentation (*tyr*),

and somatic growth (*mrap2a*) in zebrafish. Our results underscore the versatility and efficiency of

CRISPRi/a in zebrafish research. For instance, by independently downregulating *sox10, mitfa,*

and *tyr*, we established zebrafish larvae with marked reduction in pigmentation. Similarly, by

activating or downregulating *mrap2a*, we successfully modulated larval length.

These proof-of-principle experiments underscore the potential of CRISPRi/a technology

to control gene expression by targeting gene promoters. CRISPRi/a is relatively straightforward

to implement compared to other genetic editing technologies, which further increases its appeal.

In addition, implementing this technology in transparent one-cell zebrafish embryos makes it

easier to study the effects of gene silencing and activation during development. Further,

CRISPRi/a can be implemented to study the effects of silencing or activating noncoding regulatory

sequences such as enhancers, repressors, and insulators. Taking advantage of the fact that

approximately 70% of human genes have a zebrafish gene homolog facilitates the study of gene

regulatory mechanisms and networks in yet another vertebrate system. The versatility to study

gene expression modulation, by either targeting gene promoters or any potential noncoding

regulatory sequences in a vertebrate system, can offer insights into disease pathogenesis and

therapeutic targets that may translate to clinical applications. Further, our evaluation of *mrap2a*

modulation also demonstrates that modulating the regulation of one gene can elicit indirect

regulation of other target genes, opening the door to a better understanding of gene regulatory

networks and their exploration for potential therapeutic interventions.

CRISPRi/a technologies, just like CRISPR gene editing, can also be limited by off-target

effects and variability in efficiency across genomic loci. Additionally, the ethical implications of

gene expression modulation and the potential for unintended consequences underscore the

importance of careful consideration in the application of this technology. In this current study, the

primary limitation is the delivery of CRISPRi/a via mRNA microinjections, which is titrated as the

embryo develops and the larvae grow older. Furthermore, developmental delays can occur in

larvae injected with foreign mRNA, which further reflects the variability induced by the combination

of mRNA microinjections, potential off-target effects, and nucleic acid toxicity. This results in a

diminution of molecular and observable phenotypes and hinders the potential to study long-term

effects of gene expression modulation, making it difficult to elucidate any potential disease

relevance. To this end, we are actively generating stable transgenic lines expressing CRISPRi/a

to facilitate rapid and systematic screening of gain or loss of gene function. Our ongoing work will

expand the zebrafish toolkit, establishing the utility of CRISPRi/a for evaluation of candidate gene

function and disease relevance.

METHODS

Generation of catalytically inactive Cas9 plasmid vector

The zebrafish optimized *Cas9* gene within the plasmid vector pT3TS-nCas9n (Addgene

#46757) was mutated to produce a catalytically inactive dead *Cas9* (dCas9). The DNA sequence

corresponding to the RuvC-like and HNH nuclease domains of the Cas9 protein were modified

by inducing a change from Aspartic acid to Alanine at position 10 (D10A), and a change from

Histidine to Alanine at position 839 (H839A). This was accomplished by implementing the

NEBuilder HiFi DNA Assembly (Cat. # E5520S) protocol using oligonucleotides primers

(Supplemental Table 1) generated using the NEBuilder Assembly Tool (v.2.10.1). The modified

plasmid vector was then transformed into NEB 5-a competent *E. coli* cells (Cat. # C2987I) and

later mini-prepped using NEB Monarch Plasmid Miniprep Kit (Cat. # T1010L). The plasmid vector

generated was named pT3TS-ndCas9n.

Codon Optimization of KRAB-MeCP2 and VP64 domains for zebrafish usage

The DNA sequence for the KRAB-MeCP2 repressor was obtained from the dCas9-KRAB-

MeCP2 plasmid vector (Addgene #110821); while the DNA sequence for the VP64 activator was

obtained from the dCas9-VP64-GFP plasmid vector (Addgene #61422). The codon optimization

tool from IDT was used to optimize these sequences for zebrafish. The KRAB-MeCP2 repressor

sequence (1,263 bp) containing an SV40 NLS before and after KRAB was optimized and ordered

as a gBlocks HiFi Gene Fragment. The VP64 activator sequence (225 bp) containing an SV40

NLS before VP64 was optimized and ordered as a gBlocks Gene Fragment.

CRISPRi/a plasmid construct generation and *in vitro* transcription

The codon optimized KRAB-MeCP2 repressor and VP64 activator were cloned into the

pT3TS-ndCas9n plasmid vector containing a catalytically inactive dCas9 sequence. Cloning was

carried out by employing NEBuilder HiFi DNA Assembly (Cat. # E5520S) using oligonucleotide

primers (Supplemental Table 2) that overlapped both the target region of the plasmid vector and

the optimized sequences. Once the optimized sequences had been cloned in, the plasmid

constructs were transformed into NEB 5-a competent *E. coli* cells (Cat. # C2987I) and later mini-

prepped using NEB Monarch Plasmid Miniprep Kit (Cat. # T1010L). The plasmid constructs

generated were named zdCas9-KRAB-MeCP2 and zdCas9-VP64 for CRISPRi and CRISPRa,

respectively.

The restriction enzyme XbaI (NEB Cat # R0145) was used to linearize the CRISPRi/a

plasmid constructs. Subsequently, *in vitro* transcription of the d*Cas9* gene plus the repressor or

activator was performed by employing the mMessage mMachine T3 Transcription kit (Cat. #

AM1348). RNA was recovered by using the RNA Clean-up protocol from the Qiagen RNeasy Mini

Kit (Cat. # 74104) and stored at -80°C.

Designing targeted gRNAs for CRISPRi/a

Multiple single guide RNAs (sgRNAs) were designed to target the promoter or 5'-UTR of

the various genes in the zebrafish. All sgRNAs followed a core 5'-N18-3'(53) sequence and were

designed using the online tool CRISPOR (PMID: 29762716). At the 5' end of the forward

oligonucleotide, the nucleotides 5'-TAGG-3' were added. At the 5' end of the reverse

oligonucleotide (reverse complement of N18 sequence), the nucleotides 5'-AAAC-3' were added.

The final form of the oligonucleotides was 5'-TAGG-N18-3' for the forward oligo; and 5'-AAAC-

N18RevCompl-3'. The oligo primers for each sgRNA were ordered from IDT (Supplemental Table

3). Oligonucleotide primers were first annealed and subsequently cloned into the pT7-gRNA

plasmid vector (Addgene #46759) containing a gRNA scaffold, according to restriction enzyme

cloning protocol specifications from previous studies(54). After cloning, each plasmid construct

generated for each sgRNA was transformed into NEB 5-a competent *E. coli* cells (Cat. # C2987I)

and later mini-prepped using NEB Monarch Plasmid Miniprep Kit (Cat. # T1010L). Linearization

of these constructs was performed using the restriction enzyme BamHI-HF (NEB Cat # R3136).

The linear DNA was purified using the Zymo Research DNA Clean & Concentrator-5 protocol

(Cat. # D4013). Subsequently, *in vitro* transcription of linear DNA was performed using the

Thermo Fisher MEGAshortscript T7 Kit (Cat. # AM1354) or the NEB HiScribe T7 High Yield RNA

Synthesis Kit (Cat. #E2040S). After in vitro transcription, RNA was recovered by using the RNA

Clean-up protocol from the Qiagen RNeasy Mini Kit (Cat. # 74104) and stored at -80°C.

Zebrafish husbandry and microinjections of one-cell stage embryos

We used the zebrafish AB strain (ZFIN ID: ZDB-GENO-960809-7) as our wild-type fish

(WT AB). The day before injections, adult AB zebrafish crosses were set up in tanks with dividers

at a ratio of 2:1 or 3:2 females per male. The morning of injections and when the lights came on,

the dividers were removed to allow for mating among zebrafish. Zebrafish embryos were injected

during the first 15-20 mins after being collected to ensure injection occurred at the one-cell stage.

The mixture injected to the embryos contain CRISPRi/a mRNA (100-300 ng/µl, final

concentration), sgRNA mRNA (150 ng/µl/each, final concentration), 0.2 µl of Phenol Red, and

Nuclease Free Water up to 5 µl. After microinjections, zebrafish embryos were incubated at 28°C

in E3 embryo medium provided by Johns Hopkins FinZ Zebrafish Core Center.

Zebrafish length measurement and image collection

All zebrafish larvae used for measurements or image collection were staged according to

anatomical hallmarks described in Kimmel *et al*(55). Larvae were anesthetized using ethyl-3-

aminobenzoate methanesulfonate salt (Tricane/MS-222; Sigma Aldrich #E10521) prepared at a

concentration of 4 mg/mL using 1 M Tris (pH 9, 4% v/v). The Nikon AZ100 Multizoom fluorescent

microscope was used for both measurements and image collection at a ratio of 3.22 µm/px. For

all images, zebrafish larvae were suspended in 3% Methyl Cellulose (Sigma Aldrich #M0387)

solution. Larvae were imaged and measured at pec-fin, day 5, and day 7 stages and p-values

were calculated using the Welch two-sample t-test.

Pigment quantification analysis in ImageJ

To assess variation of pigmentation across conditions, we used ImageJ(56) to measure

the total area of pigmentation in larvae within the prim 20-25 stages and at the pec-fin stages,

across treatment conditions. For this, lateral and dorsal view images were used to assess the

level of pigmentation in epidermal melanocytes and in the RPE, respectively. Briefly, we employed

a previously published pipeline(57) with minor modifications in the selection of the region of

interest (ROI). We first converted all images to 16-bit images before selecting the ROI. We then

created a mask from the ROI and followed by using the threshold function with default

thresholding method. Next, we restored the selection and used the analyze particle function to

measure the area of pigmentation within the ROI. For analyzing lateral view images, the contour

of the larvae, without eyes and the yolk sac, was selected as the ROI. The eyes were excluded

due to the superimposition of one eye over the other from a lateral view increasing the total

pigmentation in one eye. The yolk sac, but not the yolk sac extension, was also excluded from

lateral view analysis due to the level of shade in the periphery of the sac. The dorsal view images

were utilized to analyze the total area of pigmentation in the eyes. Larvae were imaged and

measured at prim 20-25 and pec-fin stages and p-values were calculated using the Welch two-

sample t-test.

T7 Endonuclease I (T7EI) assay for heteroduplex detection

Zebrafish embryos (n > 250) were injected with zebrafish-optimized *Cas9* mRNA (300

ng/µl, final concentration) that was *in-vitro* transcribed from the linearized pT3TS-nCas9n

(Addgene #46757) plasmid vector. The microinjection mixture also included synthesized sgRNA

mRNA (150 ng/µl, final concentration), nuclease-free water, and phenol red. At 48 hours post

injection, clutches of 50 embryos per condition were collected for DNA extraction. DNA extraction

was performed by incubating embryo clutches in 50 µl of zebrafish lysis buffer (1M Tris pH 8.4,

1M KCl, 1M MgCl$_2$, 20% Tween-20, and 10% NP-40) at 94°C for 20 mins, 55°C (Hold to add 10

mg/ml Proteinase K), 55°C for 60 mins, 94°C for 20 mins, and optionally incubating at 4°C.

Following extraction, DNA was amplified following standard PCR amplification protocols and

purified using the Zymo Research DNA Clean & Concentrator-5 protocol (Cat. # D4013). The

standard protocol from the T7EI assay from NEB (Cat. #M0689) was followed and after digestion,

results were visualized in a 2% agarose gel.

Ethics Statement

All zebrafish maintenance, handling, and analyses were performed under approved
protocols from the Johns Hopkins University Animal Care and Use Committee (Protocol #
FI22M203)

AUTHOR CONTRIBUTIONS: A.S.M., V.C.C, and N.B.B. conceptualized the study. N.B.B.
designed codon optimized dCas9-KRAB-MeCP2 and dCas9-VP64 constructs and custom single
guide RNAs. N.B.B., E.A.S., and R.J.B. injected RNA constructs into zebrafish embryos. N.B.B.
and E.A.S. imaged the fish. N.B.B. and R.J.B. wrote the manuscript. All authors edited and
approved the final manuscript for publication.

FUNDING: This research, undertaken at Johns Hopkins University School of Medicine, was
supported in part by awards from the National Institutes of Health to A.S.M. (R21HD120879),
V.C.C. (R35GM136644), as well as N.B.B., E.A.S., and R.J.B. (T32GM148383-01); and by the
Canadian Institutes of Health Research (DFD-181599) to R.J.B.

ACKNOWLEDGEMENTS: The authors would like to acknowledge the FInZ Zebrafish Core
Center for assistance with zebrafish husbandry.

CONFLICTS OF INTEREST: The authors declare no competing interests.

REFERENCES

1.   Giffen KP, Liu H, Kramer KL, He DZ. Expression of Protein-Coding Gene Orthologs in
Zebrafish and Mouse Inner Ear Non-sensory Supporting Cells. Front Neurosci [Internet].
2019 Oct 18 [cited 2024 Aug 28];13. Available from: /pmc/articles/PMC6813431/

2.   Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, et al. The zebrafish
reference genome sequence and its relationship to the human genome. Nature [Internet].
2013 Apr 25 [cited 2024 Aug 28];496(7446):498–503. Available from:
https://pubmed.ncbi.nlm.nih.gov/23594743/

3.   Fisher S, Grice EA, Vinton RM, Bessling SL, Urasaki A, Kawakami K, et al. Evaluating
the biological relevance of putative enhancers using Tol2 transposon-mediated
transgenesis in zebrafish. Nature Protocols 2006 1:3 [Internet]. 2006 Oct 12 [cited 2022
Feb 15];1(3):1297–305. Available from: https://www.nature.com/articles/nprot.2006.230

4.   Gorkin DU, Lee D, Reed X, Fletez-Brant C, Bessling SL, Loftus SK, et al. Integration of
ChIP-seq and machine learning reveals enhancers and a predictive regulatory sequence
vocabulary in melanocytes. Genome Res [Internet]. 2012 Nov [cited 2024 Aug
28];22(11):2290–301. Available from: https://pubmed.ncbi.nlm.nih.gov/23019145/

5.   Fisher S, Grice EA, Vinton RM, Bessling SL, McCallion AS. Conservation of RET
regulatory function from human to zebrafish without sequence similarity. Science
[Internet]. 2006 Apr 14 [cited 2024 Aug 28];312(5771):276–9. Available from:
https://pubmed.ncbi.nlm.nih.gov/16556802/

6.   Antonellis A, Huynh JL, Lee-Lin SQ, Vinton RM, Renaud G, Loftus SK, et al. Identification
of neural crest and glial enhancers at the mouse Sox10 locus through transgenesis in
zebrafish. PLoS Genet [Internet]. 2008 Sep [cited 2024 Aug 28];4(9). Available from:
https://pubmed.ncbi.nlm.nih.gov/18773071/

7.   Doyle AJ, Doyle JJ, Bessling SL, Maragh S, Lindsay ME, Schepers D, et al. Mutations in
the TGF-β repressor SKI cause Shprintzen-Goldberg syndrome with aortic aneurysm.
Nat Genet [Internet]. 2012 Nov [cited 2024 Aug 28];44(11):1249–54. Available from:
https://pubmed.ncbi.nlm.nih.gov/23023332/

8.   Gould RA, Aziz H, Woods CE, Seman-Senderos MA, Sparks E, Preuss C, et al. ROBO4
variants predispose individuals to bicuspid aortic valve and thoracic aortic aneurysm. Nat
Genet [Internet]. 2019 Jan 1 [cited 2024 Aug 28];51(1):42–50. Available from:
https://pubmed.ncbi.nlm.nih.gov/30455415/

9.   Golzio C, Willer J, Talkowski ME, Oh EC, Taniguchi Y, Jacquemont S, et al. KCTD13 is a
major driver of mirrored neuroanatomical phenotypes of the 16p11.2 copy number
variant. Nature [Internet]. 2012 May 17 [cited 2024 Aug 28];485(7398):363–7. Available
from: https://pubmed.ncbi.nlm.nih.gov/22596160/

10.   Panizzi JR, Becker-Heck A, Castleman VH, Al-Mutairi DA, Liu Y, Loges NT, et al.
CCDC103 mutations cause primary ciliary dyskinesia by disrupting assembly of ciliary
dynein arms. Nat Genet [Internet]. 2012 Jun [cited 2024 Aug 28];44(6):714–9. Available
from: https://pubmed.ncbi.nlm.nih.gov/22581229/

11. Roscioli T, Kamsteeg EJ, Buysse K, Maystadt I, Van Reeuwijk J, Van Den Elzen C, et al. Mutations in ISPD cause Walker-Warburg syndrome and defective glycosylation of α-dystroglycan. Nat Genet [Internet]. 2012 May [cited 2024 Aug 28];44(5):581–5. Available from: https://pubmed.ncbi.nlm.nih.gov/22522421/

12. Tobin DM, Roca FJ, Oh SF, McFarland R, Vickery TW, Ray JP, et al. Host genotype-specific therapies can optimize the inflammatory response to mycobacterial infections. Cell [Internet]. 2012 Feb 3 [cited 2024 Aug 28];148(3):434–46. Available from: https://pubmed.ncbi.nlm.nih.gov/22304914/

13. Maragh S, Miller RA, Bessling SL, McGaughey DM, Wessels MW, De Graaf B, et al. Identification of RNA binding motif proteins essential for cardiovascular development. BMC Dev Biol [Internet]. 2011 [cited 2024 Aug 28];11. Available from: https://pubmed.ncbi.nlm.nih.gov/22011202/

14. Wahbeh MH, Boyd RJ, Yovo C, Rike B, McCallion AS, Avramopoulos D. A functional schizophrenia-associated genetic variant near the TSNARE1 and ADGRB1 genes. HGG Adv [Internet]. 2024 Jul 18 [cited 2024 Aug 28];5(3). Available from: https://pubmed.ncbi.nlm.nih.gov/38702885/

15. Asakawa K, Kawakami K. Targeted gene expression by the Gal4-UAS system in zebrafish. Dev Growth Differ [Internet]. 2008 Aug 1 [cited 2024 Sep 8];50(6):391–9. Available from: https://onlinelibrary.wiley.com/doi/full/10.1111/j.1440-169X.2008.01044.x

16. Zhu C, Smith T, Mcnulty J, Rayla AL, Lakshmanan A, Siekmann AF, et al. Evaluation and application of modularly assembled zinc-finger nucleases in zebrafish. Development [Internet]. 2011 Oct 15 [cited 2024 Sep 8];138(20):4555–64. Available from: http://www.r-project.org

17. Huang P, Xiao A, Zhou M, Zhu Z, Lin S, Zhang B. Heritable gene targeting in zebrafish using customized TALENS. Nature Biotechnology 2011 29:8 [Internet]. 2011 Aug 1 [cited 2024 Sep 8];29(8):699–700. Available from: https://www.nature.com/articles/nbt.1939

18. Hwang WY, Fu Y, Reyon D, Maeder ML, Tsai SQ, Sander JD, et al. Efficient genome editing in zebrafish using a CRISPR-Cas system. Nat Biotechnol [Internet]. 2013 Mar [cited 2024 Aug 28];31(3):227–9. Available from: https://pubmed.ncbi.nlm.nih.gov/23360964/

19. Shah AN, Davey CF, Whitebirch AC, Miller AC, Moens CB. Rapid reverse genetic screening using CRISPR in zebrafish. Nat Methods [Internet]. 2015 May 28 [cited 2024 Aug 28];12(6):535–40. Available from: https://pubmed.ncbi.nlm.nih.gov/25867848/

20. Varshney GK, Pei W, Lafave MC, Idol J, Xu L, Gallardo V, et al. High-throughput gene targeting and phenotyping in zebrafish using CRISPR/Cas9. Genome Res [Internet]. 2015 Jul 1 [cited 2024 Aug 28];25(7):1030–42. Available from: https://pubmed.ncbi.nlm.nih.gov/26048245/

21. Zhang Y, Qin W, Lu X, Xu J, Huang H, Bai H, et al. Programmable base editing of zebrafish genome using a modified CRISPR-Cas9 system. Nat Commun [Internet]. 2017 Dec 1 [cited 2024 Aug 28];8(1). Available from: https://pubmed.ncbi.nlm.nih.gov/28740134/

22. Hisano Y, Sakuma T, Nakade S, Ohga R, Ota S, Okamoto H, et al. Precise in-frame
integration of exogenous DNA mediated by CRISPR/Cas9 system in zebrafish. Sci Rep
[Internet]. 2015 [cited 2024 Aug 28];5. Available from:
https://pubmed.ncbi.nlm.nih.gov/25740433/

23. Larson MH, Gilbert LA, Wang X, Lim WA, Weissman JS, Qi LS. CRISPR interference
(CRISPRi) for sequence-specific control of gene expression. Nature Protocols 2013 8:11
[Internet]. 2013 Oct 17 [cited 2023 Jul 25];8(11):2180–96. Available from:
https://www.nature.com/articles/nprot.2013.132

24. Yeo NC, Chavez A, Lance-Byrne A, Chan Y, Menn D, Milanova D, et al. An enhanced
CRISPR repressor for targeted mammalian gene regulation. Nat Methods [Internet]. 2018
Aug 1 [cited 2023 Dec 19];15(8):611–6. Available from:
https://pubmed.ncbi.nlm.nih.gov/30013045/

25. Chavez A, Tuttle M, Pruitt BW, Ewen-Campen B, Chari R, Ter-Ovanesyan D, et al.
Comparison of Cas9 activators in multiple species. Nat Methods [Internet]. 2016 Jun 29
[cited 2024 Aug 28];13(7):563–7. Available from:
https://pubmed.ncbi.nlm.nih.gov/27214048/

26. Long L, Guo H, Yao D, Xiong K, Li Y, Liu P, et al. Regulation of transcriptionally active
genes via the catalytically inactive Cas9 in C. elegans and D. rerio. Cell Research 2015
25:5 [Internet]. 2015 Mar 27 [cited 2023 Aug 22];25(5):638–41. Available from:
https://www.nature.com/articles/cr201535

27. Sastre D, Zafar F, Torres CAM, Piper D, Kirik D, Sanders LH, et al. Inactive S. aureus
Cas9 downregulates alpha-synuclein and reduces mtDNA damage and oxidative stress
levels in human stem cell model of Parkinson's disease. Sci Rep [Internet]. 2023 Dec 1
[cited 2024 Aug 28];13(1). Available from: https://pubmed.ncbi.nlm.nih.gov/37853101/

28. Matharu N, Ahituv N. Modulating gene regulation to treat genetic disorders. Nature
Reviews Drug Discovery 2020 19:11 [Internet]. 2020 Oct 5 [cited 2023 Jul
25];19(11):757–75. Available from: https://www.nature.com/articles/s41573-020-0083-7

29. Matharu N, Rattanasopha S, Tamura S, Maliskova L, Wang Y, Bernard A, et al. CRISPR-
mediated activation of a promoter or enhancer rescues obesity caused by
haploinsufficiency. Science (1979) [Internet]. 2019 Jan 18 [cited 2023 Jul 24];363(6424).
Available from: https://www.science.org/doi/10.1126/science.aau0629

30. Camp E, Lardelli M. Tyrosinase gene expression in zebrafish embryos. Dev Genes Evol
[Internet]. 2001 [cited 2023 Aug 29];211(3):150–3. Available from:
https://link.springer.com/article/10.1007/s004270000125

31. Cooper CD, Raible DW. Mechanisms for reaching the differentiated state: Insights from
neural crest-derived melanocytes. Semin Cell Dev Biol [Internet]. 2009 [cited 2024 Aug
28];20(1):105–10. Available from: https://pubmed.ncbi.nlm.nih.gov/18935965/

32. Haffter P, Odenthal J, Mullins MC, Lin S, Farrell MJ, Vogelsang E, et al. Mutations
affecting pigmentation and shape of the adult zebrafish. Dev Genes Evol [Internet]. 1996
[cited 2023 Aug 30];206(4):260–76. Available from:
https://link.springer.com/article/10.1007/s004270050051

33.   Lane BM, Lister JA. Otx but Not Mitf Transcription Factors Are Required for Zebrafish
601   Retinal Pigment Epithelium Development. PLoS One [Internet]. 2012 Nov 5 [cited 2024
602   Feb 18];7(11):49357. Available from: /pmc/articles/PMC3489725/

603   34.   Lister JA, Close J, Raible DW. Duplicate mitf genes in zebrafish: complementary
604   expression and conservation of melanogenic potential. Dev Biol [Internet]. 2001 Sep 15
605   [cited 2023 Nov 28];237(2):333–44. Available from:
606   https://pubmed.ncbi.nlm.nih.gov/11543618/

607   35.   Murisier F, Guichard S, Beermann F. The tyrosinase enhancer is activated by Sox10 and
608   Mitf in mouse melanocytes. Pigment Cell Res [Internet]. 2007 Jun 1 [cited 2023 Aug
609   29];20(3):173–84. Available from: https://onlinelibrary.wiley.com/doi/full/10.1111/j.1600-
610   0749.2007.00368.x

611   36.   Dutton KA, Pauliny A, Lopes SS, Elworthy S, Carney TJ, Rauch J, et al. Zebrafish
612   colourless encodes sox10 and specifies non-ectomesenchymal neural crest fates.
613   Development [Internet]. 2001 Nov 1 [cited 2024 Aug 11];128(21):4113–25. Available
614   from: https://dx.doi.org/10.1242/dev.128.21.4113

615   37.   Johnson SL, Nguyen ATN, Lister JA. mitfa is required at multiple stages of melanocyte
616   differentiation but not to establish the melanocyte stem cell. Dev Biol [Internet]. 2011 Feb
617   2 [cited 2024 Aug 28];350(2):405. Available from: /pmc/articles/PMC3040983/

618   38.   Wang C, Kocher TD, Wu J, Li P, Liang G, Lu B, et al. Knockout of microphthalmia-
619   associated transcription factor (mitf) confers a red and yellow tilapia with few pigmented
620   melanophores. Aquaculture. 2023 Feb 25;565:739151.

621   39.   Wu YC, Wang IJ. Heat-shock-induced tyrosinase gene ablation with CRISPR in
622   zebrafish. Molecular Genetics and Genomics [Internet]. 2020 Jul 1 [cited 2023 Aug
623   29];295(4):911–22. Available from: https://link.springer.com/article/10.1007/s00438-020-
624   01681-x

625   40.   Camp E, Badhwar P, Mann GJ, Lardelli M. Expression Analysis of a Tyrosinase Promoter
626   Sequence in Zebrafish. Pigment Cell Res [Internet]. 2003 Apr 1 [cited 2023 Sep
627   10];16(2):117–26. Available from: https://onlinelibrary.wiley.com/doi/full/10.1034/j.1600-
628   0749.2003.00002.x

629   41.   Frantz WT, Ceol CJ. From Tank to Treatment: Modeling Melanoma in Zebrafish. Cells
630   [Internet]. 2020 May 22 [cited 2024 Aug 28];9(5). Available from:
631   https://pubmed.ncbi.nlm.nih.gov/32455885/

632   42.   Praetorius C, Grill C, Stacey SN, Metcalf AM, Gorkin DU, Robinson KC, et al. A
633   polymorphism in IRF4 affects human pigmentation through a tyrosinase-dependent
634   MITF/TFAP2A pathway. Cell [Internet]. 2013 Nov 21 [cited 2024 Aug 28];155(5):1022.
635   Available from: https://pubmed.ncbi.nlm.nih.gov/24267888/

636   43.   Rosenbaum SR, Tiago M, Caksa S, Capparelli C, Purwin TJ, Kumar G, et al. SOX10
637   requirement for melanoma tumor growth is due, in part, to immune-mediated effects. Cell
638   Rep [Internet]. 2021 Dec 7 [cited 2024 Aug 28];37(10). Available from:
639   https://pubmed.ncbi.nlm.nih.gov/34879275/

44.   Gudbjartsson DF, Sulem P, Stacey SN, Goldstein AM, Rafnar T, Sigurgeirsson B, et al.
ASIP and TYR pigmentation variants associate with cutaneous melanoma and basal cell
carcinoma. Nat Genet [Internet]. 2008 [cited 2024 Aug 28];40(7):886–91. Available from:
https://pubmed.ncbi.nlm.nih.gov/18488027/

45.   Garraway LA, Widlund HR, Rubin MA, Getz G, Berger AJ, Ramaswamy S, et al.
Integrative genomic analyses identify MITF as a lineage survival oncogene amplified in
malignant melanoma. Nature [Internet]. 2005 Jul 7 [cited 2024 Aug 28];436(7047):117–
22. Available from: https://pubmed.ncbi.nlm.nih.gov/16001072/

46.   Tassabehji M, Newton VE, Read AP. Waardenburg syndrome type 2 caused by
mutations in the human microphthalmia (MITF) gene. Nat Genet [Internet]. 1994 [cited
2024 Aug 28];8(3):251–5. Available from: https://pubmed.ncbi.nlm.nih.gov/7874167/

47.   Bondurand N, Pingault V, Goerich DE, Lemort N, Sock E, Le Caignec C, et al. Interaction
among SOX10, PAX3 and MITF, three genes altered in Waardenburg syndrome. Hum
Mol Genet [Internet]. 2000 Aug 12 [cited 2024 Aug 28];9(13):1907–17. Available from:
https://pubmed.ncbi.nlm.nih.gov/10942418/

48.   Paratore C, Eichenberger C, Suter U, Sommer L. Sox10 haploinsufficiency affects
maintenance of progenitor cells in a mouse model of Hirschsprung disease. Hum Mol
Genet [Internet]. 2002 Nov 15 [cited 2024 Aug 28];11(24):3075–85. Available from:
https://pubmed.ncbi.nlm.nih.gov/12417529/

49.   Mccallion AS, Chakravarti A. EDNRB/EDN3 and Hirschsprung disease type II. Pigment
Cell Res [Internet]. 2001 [cited 2024 Aug 28];14(3):161–9. Available from:
https://pubmed.ncbi.nlm.nih.gov/11434563/

50.   Sebag JA, Zhang C, Hinkle PM, Bradshaw AM, Cone RD. Developmental Control of the
Melanocortin-4 Receptor by MRAP2 Proteins in Zebrafish. Science [Internet]. 2013 Jul 7
[cited 2024 Jun 27];341(6143):278. Available from: /pmc/articles/PMC4255277/

51.   Yeo GSH, Farooqi IS, Aminian S, Halsall DJ, Stanhope RG, O'Rahilly S. A frameshift
mutation in MC4R associated with dominantly inherited human obesity. Nat Genet
[Internet]. 1998 [cited 2024 Aug 28];20(2):111–2. Available from:
https://pubmed.ncbi.nlm.nih.gov/9771698/

52.   Wade KH, Lam BYH, Melvin A, Pan W, Corbin LJ, Hughes DA, et al. Loss-of-function
mutations in the melanocortin 4 receptor in a UK birth cohort. Nat Med [Internet]. 2021
Jun 1 [cited 2024 Aug 28];27(6):1088–96. Available from:
https://pubmed.ncbi.nlm.nih.gov/34045736/

53.   Amorim JP, Bordeira-Carriço R, Gali-Macedo A, Perrod C, Bessa J. CRISPR-Cas9-
Mediated Genomic Deletions Protocol in Zebrafish. STAR Protoc [Internet]. 2020 Dec 18
[cited 2023 Dec 20];1(3). Available from: https://pubmed.ncbi.nlm.nih.gov/33377102/

54.   Jao LE, Wente SR, Chen W. Efficient multiplex biallelic zebrafish genome editing using a
CRISPR nuclease system. Proc Natl Acad Sci U S A [Internet]. 2013 [cited 2023 Dec
19];110(34):13904–9. Available from: https://pubmed.ncbi.nlm.nih.gov/23918387/

55.   Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic
680   development of the zebrafish. Developmental Dynamics [Internet]. 1995 [cited 2025 Feb
681   9];203(3):253–310. Available from: https://pubmed.ncbi.nlm.nih.gov/8589427/

682   56.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
683   analysis. Nat Methods. 2012 Jul 28;9(7):671–5.

684   57.   Siegenthaler A, Mondal D, Benvenuto C. Quantifying pigment cover to assess variation in
685   animal colouration. Biol Methods Protoc. 2017 Jan 1;2(1).

686   58.   Cunningham RL, Kramer ET, DeGeorgia SK, Godoy PM, Zarov AP, Seneviratne S, et al.
687   Functional in vivo characterization of sox10 enhancers in neural crest and melanoma
688   development. Communications Biology 2021 4:1 [Internet]. 2021 Jun 7 [cited 2024 Aug
689   11];4(1):1–15. Available from: https://www.nature.com/articles/s42003-021-02211-0

690   59.   Kawakami A, Fisher DE. The master role of microphthalmia-associated transcription
691   factor in melanocyte and melanoma biology. Laboratory Investigation 2017 97:6
692   [Internet]. 2017 Mar 6 [cited 2023 Aug 22];97(6):649–56. Available from:
693   https://www.nature.com/articles/labinvest20179

694   60.   Cooper CD. Insights from zebrafish on human pigment cell disease and treatment.
695   Developmental Dynamics [Internet]. 2017 Nov 1 [cited 2024 Jul 31];246(11):889–96.
696   Available from: https://onlinelibrary.wiley.com/doi/full/10.1002/dvdy.24550

697   61.   Moreno-Marmol T, Cavodeassi F, Bovolenta P. Setting eyes on the retinal pigment
698   epithelium. Front Cell Dev Biol [Internet]. 2018 Oct 24 [cited 2024 Jul 29];6(OCT):418922.
699   Available from: www.frontiersin.org

700   62.   Baron M, Maillet J, Huyvaert M, Dechaume A, Boutry R, Loiselle H, et al. Loss-of-function
701   mutations in MRAP2 are pathogenic in hyperphagic obesity with hyperglycemia and
702   hypertension. Nature Medicine 2019 25:11 [Internet]. 2019 Nov 7 [cited 2024 Aug
703   13];25(11):1733–8. Available from: https://www.nature.com/articles/s41591-019-0622-0

704   63.   Asai M, Ramachandrappa S, Joachim M, Shen Y, Zhang R, Nuthalapati N, et al. Loss of
705   function of the melanocortin 2 receptor accessory protein 2 is associated with mammalian
706   obesity. Science (1979) [Internet]. 2013 Jul 19 [cited 2024 Aug 13];341(6143):275–8.
707   Available from: https://www.science.org/doi/10.1126/science.1233000

708   64.   Vaisse C, Clement K, Durand E, Hercberg S, Guy-Grand B, Froguel P. Melanocortin-4
709   receptor mutations are a frequent and heterogeneous cause of morbid obesity. J Clin
710   Invest [Internet]. 2000 [cited 2024 Aug 13];106(2):253–62. Available from:
711   https://pubmed.ncbi.nlm.nih.gov/10903341/

712   65.   de Vrieze E, de Bruijn SE, Reurink J, Broekman S, van de Riet V, Aben M, et al. Efficient
713   generation of knock-in zebrafish models for inherited disorders using crispr-cas9
714   ribonucleoprotein complexes. Int J Mol Sci [Internet]. 2021 Sep 1 [cited 2023 Aug
715   22];22(17):9429. Available from: https://www.mdpi.com/1422-0067/22/17/9429/htm

716   66.   Alerasool N, Segal D, Lee H, Taipale M. An efficient KRAB domain for CRISPRi
717   applications in human cells. Nature Methods 2020 17:11 [Internet]. 2020 Oct 5 [cited
718   2023 Nov 28];17(11):1093–6. Available from: https://www.nature.com/articles/s41592-
719   020-0966-x

Figure 1: CRISPRi/a construct design and _tyr_ silencing. **A.** Zebrafish codon-optimized plasmids were
synthesized using NEB HiFi DNA assembly, transformed and amplified in NEB 5-α chemically competent
cells. Plasmid DNA was linearized and _in vitro_ transcribed before microinjection into one-cell embryos. **B.**
Zebrafish embryos injected with CRISPRi (final concentration 100-300 ng/µl) + cocktail of sgRNAs (final
concentration 150 ng/µl) targeting _tyr_ assayed at prim 20-25 (33-36 hpf) and pec-fin (60 hpf) stages for
differences in pigmentation patterns. The larvae injected with CRISPRi and sgRNAs displayed reduced
pigmentation at the prim 20-25 stage when compared to controls. The effect of CRISPRi dissipated by the
pec-fin stage as observable pigmentation patterns were indistinguishable. Scale bars 1000 µm. **C.** Pigment
analysis of larvae at Prim 20-25 stage (33-36 hpf) and Pec-fin stage (60 hpf) to measure the percentage of
pigmented area, as viewed laterally. At Prim 20-25, the percentage of pigmented area is statistically
significant (Welch two-sample t-test); by contrast, at Pec-fin, while not statistically significant,
CRISPRi+sgRNA injected embryos show less percentage of pigmented area than WT AB and injection
controls.

Figure 2: CRISPRi targeting the promoters of the *sox10* and *mitfa* genes. **A.** Zebrafish 1-cell embryos injected with CRISPRi + sgRNAs targeting *sox10* assayed at prim 20-25 (33-36 hpf) and pec-fin (60 hpf) stages for differences in pigmentation patterns. WT AB and injected controls did not show reduced pigmentation around the prim 20-25 stages, whereas embryos injected with CRISPRi + All *sox10* sgRNAs or CRISPRi + *sox10* sgRNA3 were hypopigmented. **B.** At the prim 20-25 stage, differences in pigmentation patterns across conditions are statistically significant (*p-value* < 0.05). At the pec-fin stage, the effect of CRISPRi dissipated as patterns of hypopigmentation are less variable with the only statistically significant difference being between WT AB and CRISPRi + All *sox10* sgRNAs based on Welch two-sample t-test. **C.** Zebrafish embryos injected with CRISPRi + sgRNAs targeting the *mitfa* promoter. Larvae were assayed at prim 20-25 and pec-fin stages for differences in pigmentation patterns. At this stage, embryos injected with CRISPRi + All sgRNAs or CRISPRi + sgRNA3 displayed a hypopigmented pattern as compared to injected and uninjected controls. At the pec-fin stage, the effect of CRISPRi has dissipated as patterns of hypopigmentation displayed no significant differences across conditions. Scale bars: 1000 μm.

747 ***Figure 3:*** Targeting the promoter of *mrap2a* with CRISPRa and CRISPRi. **A.** Length comparison between the pec-fin and day-5 stages between
748 WT AB embryos and embryos injected with CRISPRa and CRISPRi with all *mrap2a* sgRNAs or only sgRNA3. Length measured laterally from tip
749 of the mouth to end of translucent tail fin (dashed lines). Red lines depict length differences between controls and experimental conditions. Scale
750 bars: 1000 μm. **B.** At the pec-fin stage, 50 larvae per condition were measured in length from the tip of the mouth to the end of the translucent tail
751 fin. The Welch two-sample t-test was used to compare the average length between conditions.

    SUPPLEMENTAL FIG.URES

752 ***Supplemental Figure 1:*** Assessment of optimal mRNA concentration for injection

Supplemental Figure 2: Targeting the *sox10* promoter with CRISPRi. **A.** Larvae at prim 20-25 (33-36 hpf) and pec-fin (60 hpf) stages across conditions. Scale bars 1000 µm. **B.** At the prim 20-25 stage, RPE pigmentation is variable across conditions; however, differences are statistically significant between controls and injected conditions. **C.** Gel (2% agarose) electrophoresis of T7 endonuclease I assay to detect heteroduplex formation (red boxes) after targeting the *sox10* promoter with sgRNA3. Expected amplicon length is 761 bp (blue arrow).

759   **Supplemental Figure 3:** Targeting the *mitfa* promoter with CRISPRi. Scale bars 1000 µm.

761   ***Supplemental Figure 4:*** CRISPRa and CRISPRi targeting of the mrap2a promoter. Images taken at pec-fin, day 5, and day 7 stages. Dotted lines
762   represent the translucent tail fin of the larvae. Scale bars 1000 μm.