A transcontinental experiment elucidates (mal)adaptation of a cosmopolitan plant to

climate in space and time

Lucas J. Albano[1,2], Cristina C. Bastias[3,4], Aurélien Estarague[4], Brandon T. Hendrickson[5],

Simon G. Innes[5], Nevada King[5], Courtney M. Patterson[5], Amelia Tudoran[6], François

Vasseur[4], Adriana Puentes[6], Cyrille Violle[4], Nicholas J. Kooyers[5], Marc T. J. Johnson[2]

[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

[2]Department of Biology, University of Toronto Mississauga, Mississauga, ON, Canada

[3]Ecology Area, Department of Botany, Ecology and Plant Physiology, Campus de Rabanales,

Córdoba University, Córdoba, Andalusia, Spain

[4]CEFE, Université Montpellier, CNRS, EPHE, IRD, Montpellier, Occitanie, France

[5]Department of Biology, University of Louisiana at Lafayette, Lafayette, LA, USA

[6]Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Uppland,

Sweden

Correspondence Author:

Lucas J. Albano

Current address: Department of Plant & Microbial Biology, North Carolina State University,

Raleigh, NC, United States, 27695, ljalbano@ncsu.edu

Open Research Statement:

Data are provided for peer review. All data involved in this study is available on the GitHub page

for LJA (https://github.com/ljalbano/transcontinental_common_garden).

Keywords:

Adaptation lag, climatic distance, common garden, cyanogenesis, environmental variation,

invasive species, local adaptation, provenance study, *Trifolium repens*, white clover

Abstract (350 words)

Climate change and the global spread of invasive species are two of the most significant threats to biodiversity. Both of these phenomena subject populations to novel conditions, either in space (species invasion) or in time (climate change), yet the role of adaptation in how populations respond to these rapid environmental shifts is poorly understood. We conducted a large-scale trans-continental common garden experiment using white clover (*Trifolium repens*, Fabaceae) to test if adaptive evolution to spatiotemporal variation in climate could contribute to the ecological success of one of the most widespread invasive plant species in the world. Individuals from 96 populations of *Trifolium repens* (white clover) from both its native (Europe) and introduced (North America) ranges were planted into four experimental common gardens located in northern (Uppsala, Sweden) and southern (Montpellier, France) Europe, and northern (Mississauga, Canada) and southern (Louisiana, USA) North America. We recorded plant sexual and clonal fitness in each common garden and assessed whether the strength of local adaptation differed between the native and introduced ranges and whether populations are rapidly adapting to climate change. Results show that local adaptation was only evident when populations were transplanted into common gardens located in the same range (native or introduced) from which they originated and was driven by stronger selection (due to climatic factors rather than herbivory) at lower latitudes in both ranges. Our results indicate that rapid local adaptation across a large latitudinal gradient has occurred in *T. repens* populations in less than 400 years since its introduction to North America. However, we find evidence of an adaptation lag in the northern common garden in the introduced range, with plants from historically slightly warmer climates exhibiting the greatest fitness. These findings support two major conclusions: 1) white clover can rapidly adapt to spatial variation in climate in its introduced range as well as the

native range, and 2) despite rapid adaptation to novel ranges, introduced white clover

populations are not keeping pace with rapid climate change. Overall, our results provide insight

into the role of adaptation in facilitating the ecological success of invasive species in a rapidly

changing world.

Introduction

Natural selection results in populations evolving traits that are advantageous to the

environmental conditions of a particular location, a phenomenon known as local adaptation

(Turesson, 1922; Kawecki and Ebert, 2004; Hereford, 2009). A fundamental aspect of local

adaptation is that variation in biotic and abiotic factors leads to changes in the relative fitness of

genotypes between environments (Gillespie and Turelli, 1989; Kawecki and Ebert, 2004; Lee

and Mitchell-Olds, 2011). Such genotype-by-environment interactions can result in fitness trade-

offs, whereby genotypes exhibit higher relative fitness in their local environment and lower

fitness in other environments (Gillespie and Turelli, 1989; Hereford, 2009; Leimu and Fischer,

2008; Savolainen et al., 2013). However, while environmental variation can result in local

adaptation, it remains poorly understood how locally adapted populations are affected when

environments rapidly change in space or time (Kawecki and Ebert, 2004; Anderson and Song,

2020). Human activities are exposing populations to novel environments, both across space

through species introductions (Pyšek et al., 2020), and over time through anthropogenically

mediated climate change (Hoffmann and Sgrò, 2011). Understanding how populations

simultaneously adapt to novel biotic and abiotic environments over space and time remains an

important problem in evolutionary ecology (Hendry, 2017; Siepielski et al., 2017), and this

understanding is necessary to determine how increasingly pervasive anthropogenic disturbances

are altering the ecology and evolution of native and introduced species (Parmesan, 2006; Fugère

and Hendry, 2018).

For a species to successfully become invasive in a novel range, an introduced population

must overcome substantial barriers to its establishment and proliferation. Introduced populations

are frequently subject to shifts in selection pressure compared to the native environment to which

they are locally adapted (Erfmeier, 2013; Colautti and Lau, 2015; Lau and terHorst, 2015). This

shift in selection pressure could involve greater stress imposed by the biotic environment (e.g.,

novel species interactions), or the abiotic environment (e.g., climatic factors), which could

inhibit the spread of an introduced population (Erfmeier, 2013; Colautti and Lau, 2015; Lau and

terHorst, 2015). Additionally, classic hypotheses in invasion biology propose that inherent

characteristics of introduced populations will limit their ability to respond to these novel

selection pressures (Allendorf and Lundquist, 2003; Estoup et al., 2016). For example, small,

introduced population sizes can result in Allee effects, such as mate limitation leading to limited

reproductive success for obligate outcrossers (Taylor and Hastings, 2005; Burns et al., 2011;

Pannell et al., 2015). Furthermore, small population size post-introduction can impose genetic

bottlenecks (Tsutsui et al., 2000). Such evolutionary events have the potential to reduce fitness

through inbreeding depression or limit genetic diversity within an introduced population to the

point that stressful, novel selection pressures cannot be overcome (Tsutsui et al., 2000; Kolar and

Lodge, 2001; Lau and terHorst, 2015). In most cases, the ecological and/or genetic constraints

experienced by introduced populations are sufficient to result in the failure of those populations

to establish and their eventual extirpation (Kolar and Lodge, 2001; Jeschke and Pyšek, 2018).

Despite the challenges presented by introduction to novel environments across space,

introduced populations are also presented with substantial ecological opportunity if genetic

constraints can be overcome. For example, selection regimes experienced in novel biotic and

abiotic environments could differ in a manner that is either benign or even advantageous to an

introduced population compared to its native range (Erfmeier, 2013; Colautti and Lau, 2015).

Classic hypotheses in invasion biology, while not universally supported, posit that a release of an

introduced species from its natural enemies in the native range (Keane and Crawley, 2002), the

evolution of increased competitive ability relative to native species (Blossey and Nötzold, 1995),

or the possession of biochemical compounds that are novel to potential enemies in an introduced

range (Callaway and Ridenour, 2004), could each result in the proliferation and success of an

invader. It is also possible that introduced populations do not exhibit strong evidence of

bottlenecks due to reduced genetic diversity (Colautti and Barrett, 2013; Battlay et al., 2024).

This could occur because repeated introductions of novel genotypes can increase effective

population size, or because reductions in genetic variation were not substantial enough to limit

the ability of a population to respond to novel selection pressures to begin with (Maron et al.,

2004; Dlugosch and Parker, 2008a; Dlugosch and Parker, 2008b; van Boheemen et al., 2017). In

fact, when introduced populations experience strong selection in novel environments, it could

facilitate rapid adaptation to those novel environments, which could in turn, contribute to the

invasion success of a species (Maron et al., 2004; Dlugosch and Parker, 2008a; Colautti and

Barrett, 2013; Colautti and Lau, 2015; Simón-Porcar et al., 2021; but see Genton et al., 2005).

However, the extent to which rapid local adaptation to novel biotic and abiotic conditions can

facilitate the success of a widespread invasive species across large environmental gradients

remains poorly understood.

While there are examples in which introduced populations have rapidly adapted to novel

environments across space, it is important to consider that these novel environments may also be

changing through time. Temporal shifts in a local environment can cause selection pressures to

fluctuate, which could result in traits that were previously adapted to that environment becoming

maladaptive (Wilczek et al., 2014; Aguirre-Liguori et al., 2019; Exposito-Alonso et al., 2019;

Anderson and Wadgymar, 2020; Bontrager et al., 2020). If the rate of adaptation is exceeded by

the rate of environmental change, this disruption of local adaptation is likely to result in reduced

fitness (i.e., maladaptation) of populations (St Clair and Howe, 2007; Atkins and Travis, 2010;

Wilczek et al., 2014; Aguirre-Liguori et al., 2019). Recent global warming may result in

populations becoming maladapted to their local climate conditions, whereas populations from

lower latitudes or altitudes become better suited to environmental conditions at higher latitude or

altitudes; a phenomenon known as an adaptation lag (Wilczek et al., 2014; Kooyers et al., 2019;

Anderson and Wadgymar, 2020). Currently, the frequency and magnitude of adaptation lags are

poorly understood and require large scale experiments with sufficient population sampling along

broad environmental gradients to assess local (mal)adaptation as a result of rapid climate change

(Anderson and Song, 2020).

If invasive species rapidly adapt to novel environments across space, similar rapid

adaptation to climate change through time could facilitate their continued range expansion,

persistence, and pervasiveness (Byers, 2002; Maron et al., 2004; Colautti and Barrett, 2013;

Oduor et al., 2016). In fact, invasive species are often better at climate tracking and more

resistant to climate warming than native species, which could contribute to their success as

invaders (Wolkovich et al., 2013). However, the interplay between local adaptation, species

introductions, and anthropogenically mediated climate change remains an understudied, yet

critically important, aspect of understanding the ongoing human-induced biodiversity crisis

(IPBES, 2019). This problem is best addressed through field experiments testing local adaptation

in both the native and introduced ranges of widespread invasive species, alongside elucidating

the environmental conditions that may result in rapid adaptation and invasion success (Colautti

and Lau, 2015; Anderson and Song, 2020).

Plant-herbivore interactions have been highlighted as key selective factors that determine

patterns of local adaptation, species responses to environmental change, and biological invasions.

Local herbivore communities impose selection through preferential feeding on plants with lower

resistance, influencing plant defense evolution through an evolutionary arms race (Ehrlich and

Raven, 1964; Agrawal, 2007; Züst et al., 2012). Spatial variation in herbivore communities is

therefore likely to lead to spatial variation in plant defense traits (Anstett et al., 2018; Baskett and

Schemske, 2018; Moreira et al., 2018). For example, the latitudinal herbivory-defense hypothesis

predicts greater diversity and abundance of plant defenses at lower latitudes due to the presence

of greater herbivore diversity and abundance (Coley and Barone, 1996; Johnson and Rasmann,

2011; Anstett et al., 2016; Moreira et al., 2018). Plant defense evolution is also likely to be

driven by the abiotic environment through growth-defense trade-offs (Coley et al., 1985; Herms

and Mattson, 1992; Züst and Agrawal, 2017). The metabolic costs of defenses can be substantial,

while abiotic environments and the stressors they impose can influence the availability of

resources to plants, exacerbating defense costs (Coley et al., 1985; Fine et al., 2006; Hahn et al.,

2019). Furthermore, it is increasingly recognized that anthropogenic activity can influence the

evolution of plant defenses (Bidart-Bouzat and Imeh-Nathaniel, 2008; Massad and Dyer, 2010;

Miles et al., 2019). Climate warming could result in increased herbivore feeding and/or increased

plant defenses (Dostálek et al., 2020; Lemoine et al., 2013) or could contribute to invasion

success by decoupling locally adapted plant-herbivore interactions (Adams et al., 2009; Costan et

al., 2022; but see Agrawal and Kotanen, 2003). It is therefore important to understand how

populations are adapting to their local biotic and abiotic environments and how anthropogenic

disturbance can affect adaptation.

An ideal study system to test questions surrounding adaptation to novel environments

post-introduction and in response to climate change would be a widespread and well-documented

invasive species that is economically and ecologically important. It would also be advantageous

to have a system with well known agents of biotic and abiotic selection across space and over

time that can drive adaptation in the native and introduced ranges. The cosmopolitan plant white

clover (*Trifolium repens* L.) offers such a system. *Trifolium repens* is an allopolyploid perennial

legume that is native to Eurasia and has been introduced to inhabited continents throughout the

world, including North America (Daday, 1958; Kjærgaard, 2003; Griffiths et al., 2019).

Mechanisms of biotic and abiotic selection in *T. repens* have been studied for over 70 years,

often in the context of the formation of clines – a common manifestation of local adaptation

represented by gradual changes in phenotypes or genotypes across environmental gradients

(Endler, 1977; Stinchcombe et al., 2004; Savolainen et al., 2013).

In *T. repens*, clines in a prominent chemical defense mechanism (cyanogenesis) have

been identified across many types of environmental gradients (Daday, 1954a; Daday, 1954b,

Daday, 1958; Till, 1987; Kooyers and Olsen, 2013; Kooyers et al., 2014; Thompson et al., 2016),

providing opportunities to test hypotheses surrounding the involvement of cyanogenesis in

adaptation to environmental change. Specifically, latitudinal clines in the native European range

of *T. repens* exhibit fixation of cyanogenesis in warmer climates and fixation of acyanogenesis in

cooler climates (Daday, 1954a). Similar, but shallower, clines have also been observed in many

introduced ranges of *T. repens*, possibly indicating weaker local adaptation post-introduction,

which occurred less than 400 years ago in the case of North America (Daday, 1958; Kjærgaard,

2003; Kooyers and Olsen, 2012; Innes et al., 2022). Alternatively, recent studies identify a lack

of involvement of cyanogenesis in local adaptation, and instead provide evidence for other

candidate traits that are likely to vary across environmental gradients in driving local adaptation,

such as flowering time and investment in vegetative growth versus reproduction (Wright et al.,

2018; Wright et al., 2021; Kuo et al., 2024). However, no study has directly compared the

strength of local adaptation in the native and introduced ranges of *T. repens* or directly tested

whether *T. repens* populations exhibit a spatial lag in adaptation to climate change.

To understand adaptation to environmental variation in an invasive species over space

and time, we conducted a trans-continental common garden field experiment across latitudinal

gradients in both the native (Europe) and introduced (North America) ranges of *T. repens*. We

first address four research questions surrounding whether *T. repens* can adapt to environmental

variation in space. (Q1) Are *T. repens* populations locally adapted to their climate of origin and

does the strength of local adaptation differ across space? This question is a prerequisite condition

before any comparisons can be made regarding local adaptation between the native and

introduced ranges. (Q2) Does *T. repens* exhibit stronger local adaptation in its native range than

in its introduced range? We anticipate weaker local adaptation in the introduced range based on

the shorter amount of time that introduced populations have had to locally adapt to novel

environments in North America (less than 400 years). (Q3) What are the roles of cyanogenesis

and herbivory (i.e., biotic factors) in driving local adaptation across space? (Q4) What

bioclimatic variables (i.e., abiotic factors) have the strongest impact on fitness across space

(latitudes and ranges) in *T. repens*? Answering the latter two questions would help to identify the

biotic and abiotic environmental drivers of local adaptation. We then focus on whether *T. repens*

can adapt to climate change through time: (Q5) Are there adaptation lags to climate change in *T.*

repens, such that populations originating from warmer and/or drier climates exhibit greater

fitness than local populations? This study is among the first to investigate the ability of an

introduced species to rapidly adapt to novel environments in space and time simultaneously. Our

results elucidate the potential for rapid adaptation to contribute to initial invasion success and to

continued ecological success of invaders in the face of ongoing environmental change, while

furthering our understanding of local adaptation to climate and the biotic environment.

Methods

Study system

Trifolium repens has been a model for the study of plant defense based on its polymorphism for

the production of the potent chemical defense hydrogen cyanide (HCN; Hughes, 1991). In fact,

the ability to produce HCN as an antiherbivore defense may have contributed to the range

expansion of *T. repens*, allowing it to thrive in human-disturbed habitats such as lawns, fields,

and roadsides in many regions throughout the world (Burdon, 1983; Kjærgaard, 2003; Olsen et

al., 2021). Cyanogenesis occurs through the hydrolyzation of cyanogenic glucosides by the

enzyme linamarase. In *T. repens*, the presence or absence of cyanogenic glucosides, linamarin

and lotaustralin, is controlled by three linked genes in their metabolic pathway (*CYP79D15*,

CYP736A187, and *UGT85K17*; referred to collectively as *Ac*/*ac*), while the presence or absence

of the enzyme linamarase is controlled by the gene *Li*/*li* (Olsen et al., 2007; Olsen and Small,

2018). Thus, four phenotypes exist for HCN and the two metabolic components required for its

production: one in which HCN is constitutively produced (AcLi) and three that are acyanogenic

and therefore lack the defense entirely (acLi, Acli, acli; Corkill, 1940; Atwood and Sullivan,

1943).

Cyanogenesis is widely considered an adaptive polymorphism because *T. repens* exhibits

clines in the presence of HCN across numerous types of environmental gradients (Daday, 1954a;

Daday, 1954b; Daday 1958; Brighton and Horne, 1977; Thompson et al., 2016; Innes et al.,

2022; Santangelo et al., 2022). In general, dominant *Ac* and *Li* alleles (and thus the ability to

produce HCN) are fixed in warmer climates, while recessive *ac* and *li* alleles are fixed in colder

climates (Daday, 1954a; Daday, 1954b; Daday, 1965). One hypothesis to explain cyanogenesis

clines is a gradient in herbivory, with greater diversity and abundance of herbivores in warmer

climates resulting in stronger selection favouring cyanogenic plants (Angseesing, 1974; Dirzo

and Harper, 1982; Kakes, 1989). Alternatively, empirical studies across various environmental

gradients have identified temperature, soil moisture, and soil nutrients as abiotic factors that may

be involved in cyanogenesis cline formation (Daday, 1954a; Daday, 1965; Gleadow and

Woodrow, 2002; Kooyers et al., 2014; Kooyers et al., 2018). Overall, despite 70 years of

research on the drivers of cyanogenesis clines in *T. repens*, there remains little consensus on how

variation in environmental conditions influence selection on cyanogenesis in *T. repens* or if

cyanogenesis is primarily responsible for rapid local adaptation to novel environments (Daday,

1954a; Daday, 1965; Hughes, 1991; Kooyers and Olsen, 2013; Kooyers et al., 2014; Wright et

al., 2018; Wright et al., 2021; Albano and Johnson, 2023; Kuo et al., 2024). Our study sought to

help resolve this debate by directly assessing cyanogenesis and bioclimatic factors as potential

drivers of local adaptation in *T. repens*.

Sampled populations and common garden sites

We addressed our research questions using four common gardens distributed across northern

(Uppsala, Sweden) and southern (Montpellier, France) regions of the native European range of *T.*

repens, and northern (Mississauga, ON, Canada) and southern (Lafayette, LA, USA) regions of

the North American introduced range (**Figure 1**). Seeds were collected from 49 natural *T. repens*

populations scattered across the native European range spanning ~27° of latitude, and 47 natural

T. repens populations along a 2500 km north-south transect spanning ~21° of latitude in the

introduced North American range (**Figure 1**). To minimize maternal and epigenetic effects, we

hand-pollinated plants germinated from field-collected seeds in a growth chamber environment

(25°C day, 15°C night, 14 h:10 h light:dark cycle; Conviron MTPS, Winnipeg, MB, Canada) to

create an outbred F1 cohort of seeds for each population. F1 generation seeds were germinated

and established for four weeks in peat pots (8 cm diameter, 591 mL volume) in uniform growth

chamber or greenhouse conditions near each common garden site. Five replicate individuals

from each population were transplanted in peat pots directly into the natural soil of a typical

cultivated lawn at each common garden site. Inconsistent germination success resulted in some

populations not being included in some common garden locations, and in very limited cases,

populations had four or six individuals planted. The resulting total number of plants per garden

was 465, 465, 458, and 470 in Mississauga, Lafayette, Uppsala, and Montpellier, respectively.

Plants were arranged in rows and columns with 1 m spacing, and black landscaping fabric was

used to cover the entirety of the lawn to prevent the rooting of stolons into the natural soil

between plants. 12 cm diameter holes were cut in the fabric for each individual plant, leaving a 2

cm ring of natural vegetation around each plant to allow for natural competition. This vegetation

was trimmed monthly throughout the experiment. Planting dates matched growing seasons in

each location. The Lafayette and Mississauga common gardens were established in April and

May 2020, respectively, while the Montpellier and Uppsala common gardens were postponed

due to COVID-19 restrictions in Europe, leading to their establishment in March and May 2021,

respectively. Plants were given supplemental water for the first two weeks of the experiment and

intermittently after the first two weeks when necessary to avoid excessive mortality. Plants that

died within the first two weeks of the experiment were deemed to have died due to

transplantation stress and were replaced using extra germinated individuals from the same

population. Plants that died after two weeks were deemed to have died due to the local

environmental conditions at each common garden site and were not replaced. Each common

garden was maintained for two full growing seasons.

Tissue collection, herbivory estimates, and measures of fitness

After 1-2 months of growth, one to three fully-expanded leaves were harvested from each

surviving individual and frozen at -20°C. These leaves were later freeze-dried and used for DNA

extraction, followed by low coverage whole genome sequencing to predict the presence of *Ac/ac*

and *Li/li* alleles (see **Supporting Information**). In total, 586 individuals (Mississauga: 302,

Lafayette: 99, Uppsala: 78, Montpellier: 107) were successfully sequenced and characterized as

possessing either dominant or recessive alleles at *Ac/ac* and *Li/li*. The remaining 1272

individuals were not genotyped because they either died before tissue could be harvested or

genomic library preparations were unsuccessful. Visual herbivory estimates (in % leaf area

consumed) were conducted monthly following the methods of Johnson et al. (2016). Herbivory

was analyzed for the month exhibiting peak herbivory within each garden. Multiple components

of plant fitness were measured throughout the growing seasons at each location. Survival and

onset of flowering were assessed weekly during the first growing season. Mature fruits were

counted and collected weekly, and seeds were separated using a thresher (Precision Machine Co.,

Lincoln, NE, USA) to obtain total seed set mass for each plant across both growing seasons of

the experiment. Photos were taken of each plant monthly during the first growing season at each

common garden location and assessed for plant area using Easy Leaf Area (Version 2.0; Easlon

and Bloom, 2014). The maximum plant area was then determined and the growth rate across the

first growing season was calculated as $(\ln[A_{max}] - \ln[A_1])/(t_{max} - t_1)$, where A_{max} represents the

maximum plant area, A_1 represents the initial plant area (i.e., after one month of growth in the

common garden), and $(t_{max} - t_1)$ represents the number of days between photos taken for

maximum plant area and initial plant area.

Calculating climatic distance

Six bioclimatic variables were extracted from WorldClim (Version 2.1, February 2023) at a

resolution of 30 seconds for each original *T. repens* population and common garden site

(O'Donnell and Ignizio, 2012). The variables included were mean annual temperature ("BIO1";

MAT), maximum temperature of the warmest month ("BIO5"), minimum temperature of the

coldest month ("BIO6"), annual precipitation ("BIO12"), precipitation of the wettest month

("BIO13"), and precipitation of the driest month ("BIO14"). The extracted values of each

bioclimatic variable for each common garden location can be found in **Table S1**. Values for all

six variables were z-score standardized to allow for analysis on an equivalent scale. A principal

component analysis was performed using the *princomp* function in R, with dimensions 1, 2, and

3 explaining 65.8%, 22.1%, and 9.1% of the variation, respectively (**Figure S1**). Coordinates for

each location of origin and each common garden location from all three dimensions were then

extracted using the *get_pca_ind* function in the *factoextra* package (version 1.0.7; Kassambara

and Mundt, 2020) and then plotted on a 3D graph (**Figure S2**). The weighted Euclidean distance

between points in 3D space was calculated using the formula:

343
$$Distance = \sqrt{\sum \left(w_i [O_{ij} - G_i]^2 \right)} \tag{1}$$

Where w_i represents the proportion of variance explained by the ith dimension, O_{ij} represents the

value of the jth population of origin in the ith dimension, and G_i represents the value of a given

common garden in the *i*th dimension. We used this weighted Euclidean distance as a measure of

the climatic distance between the location of each population of origin and each common garden

location based on our six bioclimatic variables of interest (Samis et al., 2019). This measure of

climatic distance was then used as a predictor variable of plant fitness in our statistical analyses.

Statistical analysis

All statistical procedures and analyses were conducted in R Version 4.1.2 (2021) and

RStudio Version 2022.12.0 (R Core Development Team, 2022). Seven response variables were

assessed as components of fitness: i) whether or not plants survived the first growing season, ii)

whether or not plants flowered in the full experiment, iii) whether or not plants produced seeds in

the full experiment, iv) number of flower heads produced in the full experiment, v) seed set mass

produced in the full experiment, vi) maximum plant area in the first growing season, and vii)

growth rate in the first growing season. Herbivory during the first growing season was also

measured as an additional non-fitness response variable. The number of flower heads and seed

set mass were both zero-inflated, which led to the use of a two-step modelling approach as

described previously (Albano and Johnson, 2023). Briefly, both variables were first analyzed in

binomial models to assess whether or not plants produced flowers or seeds. Non-zero flower

head numbers and seed set masses were then subsequently analyzed as separate measures of

fitness. All general linear mixed effects models were conducted using the *glmmTMB* function in

the *glmmTMB* package (version 1.1.5; Brooks et al., 2017). All test statistics and *P* values were

calculated using the *Anova* function in the *car* package (version 3.0-12; Fox and Weisberg, 2019)

with Type II sums-of-squares. For models using quantitative measures of fitness, all

transformations necessary to meet assumptions of normality and homogeneity of variance can be

found in **Table S2**.

Variation in the strength of local adaptation between common garden locations

To determine whether or not *T. repens* populations exhibit local adaptation and to assess the

strength of local adaptation at each garden location (Q1), all response variables were analyzed

using the following structure:

373
$$Response \sim Continent\ of\ Origin\ (O) + Garden\ Location\ (L) + Climatic$$
$$Distance\ (C) + O{:}L + O{:}D + L{:}D + O{:}L{:}D + (1|Population) \qquad (2)$$

Continent of origin (North America or Europe), garden location (Mississauga, Lafayette,

Uppsala, or Montpellier), and climatic distance (the combined measure of six bioclimatic

variables of interest described above) were included as fixed effects, along with all two- and

three-way interactions. Population nested within continent of origin was included as a random

effect to account for non-independence of replicates within each population. An interaction

between continent of origin and climatic distance would indicate that the strength of local

adaptation varies based on whether or not populations originated in the native or introduced

range. An interaction between garden location and climatic distance would indicate that the

strength of local adaptation varies based on the particular common environment in which it is

being tested. The strength of local adaptation refers to the magnitude of the fitness advantage for

local populations compared to populations transplanted an increasingly larger climatic distance

from their origin. We therefore interpret the slope of the relationship between climatic distance

and fitness as being representative of the strength of local adaptation within a given common

garden.

Variation in the strength of local adaptation between ranges

To assess the strength of local adaptation between the native and introduced ranges of *T. repens*

(Q2), the analyses of the previous sub-section were repeated, with the continent of the common

garden (North America or Europe) replacing the location of the common garden in the models:

$$Response \sim Continent\ of\ Origin + Continent\ of\ Garden\ (G) + Climatic$$
$$Distance +\ O{:}G + O{:}D + D{:}G + O{:}D{:}G + (1|Population) \qquad (3)$$

In this model, a 3-way interaction between continent of origin, continent of garden, and climatic

distance may indicate that the strength of local adaptation varies based not only on the range

from which populations originated but also the range in which they are being tested. In other

words, this model aims to test whether locally adapted European populations maintain signatures

of local adaptation in North America and vice versa.

The roles of cyanogenesis and herbivory in local adaptation

The presence of cyanogenesis clines across latitude on both continents was determined

separately for *Ac/ac* and *Li/li* by assessing the proportion of plants in each population containing

at least one dominant *Ac* or *Li* allele using the following model structure:

$$Proportion\ \text{Ac}\ or\ \text{Li}\ Plants \sim Latitude + Continent\ of\ Origin$$
$$+ Latitude{:}Continent\ of\ Origin + (1|Population) \qquad (7)$$

An interaction between latitude and continent of origin would indicate a difference in the slope

of clines in the presence of *Ac* and/or *Li* based on the range in which those clines are occurring

(native European or introduced North American).

The effect of cyanotype (AcLi, Acli, acLi, or acli) on local adaptation (Q3) was assessed

for all eight response variables using the following model structure:

$$Response \sim Cyanotype\ (C) + Garden\ Location\ (L) + Climatic\ Distance\ (D)$$
$$+ C{:}L + C{:}D + C{:}L{:}D + (1|Population) \qquad (8)$$

An interaction between cyanotype and climatic distance would indicate that cyanogenesis or its

metabolic components contribute to the strength of local adaptation in *T. repens*. A 3-way

interaction between cyanotype, garden location, and climatic distance would indicate that

differential selection for or against a given cyanotype in a particular common environment

impacts the strength of local adaptation in that environment. This might be expected due to

consistently observed latitudinal cyanogenesis clines across multiple types of environmental

gradients in *T. repens.*

Bioclimatic variables as individual drivers of local adaptation

We also measured associations between the seven response variables and each of the six

bioclimatic variables of interest individually by determining the difference in z-score

standardized values for each variable between the location of each population of origin and each

common garden location using the following model structure:

$Response \sim Bioclimatic\ Variable + Bioclimatic\ Variable^2 + (1|Population)$ (4)

For this set of analyses, all common gardens were analyzed separately, and all quantitative (non-

binomial) response variables were z-score standardized to allow for simultaneous direct

comparisons of the relative strength of linear and quadratic predictor variables. Within each

garden, we were therefore able to directly compare *P* values across all bioclimatic variables for

both linear and quadratic fits (**Tables S3-S6**). For each response variable, we selected the most

influential bioclimatic variable (i.e., the lowest *P*, but only when at least one *P* value was

significant at a level of *P* < 0.1, highlighted in red in **Tables S3-S6**). Across all response

variables in each garden, we then calculated the percentage of these most influential bioclimatic

variables that were either temperature-based (BIO1, BIO5, or BIO6) or precipitation-based

(BIO12, BIO13, and BIO14) and compared these percentages across the full experiment, across

latitudes, and across ranges (Q4).

Adaptation lags in response to climate warming

We sought to investigate the presence of adaptation lags using a space-for-time substitution (Q5).

We performed an additional analysis using a similar model structure as before, but only using

changes in raw values for MAT (i.e., not z-score standardized, hereafter referred to as ΔMAT) as

the bioclimatic variable of interest.

$$Response \sim \Delta MAT + \Delta MAT^2 + (1|Population) \qquad (5)$$

If the peak of the quadratic relationship in each model is approximately equal to zero, it would be

considered evidence of local adaptation to MAT, because maximum fitness occurs in populations

with the same mean annual temperature as the common garden site. If the peak of the quadratic

relationship is instead shifted towards populations of warmer origins than are occurring at the

common garden site (i.e., these warmer-adapted populations are outperforming the local

populations), this would be considered evidence of an adaptation lag. We tested whether the shift

of peaks away from a ΔMAT of zero and towards populations of warmer origins were significant

by also fitting a similar model to above, but excluding the linear predictor:

$$Response \sim \Delta MAT^2 + (1|Population) \qquad (6)$$

This additional model constrains the peak of the distribution to be centred around a ΔMAT of

zero (approximating the outcome of local adaptation as opposed to an adaptation lag). We then

compared the models with and without the linear predictor using the *anova* and *AIC* functions in

R, with *P* values < 0.05 (along with ΔAIC values < 0) indicating a significant shift of the peak.

We focused our interpretation of this analysis based solely on the North American

populations in the Mississauga common garden because the sampling design of North American

populations along a transect and the location of the Mississauga garden within that sampled

transect provide the most appropriate subset of the full experiment data to detect spatial lags. We

include the results for European populations within the Uppsala common garden in **Table S7** for

completeness, but the Uppsala common garden did not exhibit any trend in fitness based on

climatic distance, precluding the possibility of an adaptation lag being detected. The Lafayette

459 and Montpellier common gardens were excluded from this analysis because they are near the

460 southern extremes of our sampled populations, precluding the ability to detect adaptation lags.

462 **Results**

463 *Q1: Are T. repens populations locally adapted to their climate of origin and does the strength of*

464 *local adaptation differ across space?*

465 We found strong evidence of local adaptation, and the strength of local adaptation in *T. repens*

466 varied among gardens for multiple measures of fitness. Climatic distance between each

467 population of origin and the common garden locations strongly predicted all measures of sexual

468 fitness and survival (**Table 1**). In general, populations tended to have higher fitness with lower

469 climatic distance, indicating that populations of more similar climatic origins to the climate of

470 each common garden were better adapted. The only response variables not showing evidence of

471 local adaptation were the measures of vegetative growth used as proxies for clonal fitness (i.e.,

472 maximum plant area and growth rate). The interaction of common garden location and climatic

473 distance affected the number of flower heads and seed set mass, but not other measures of fitness

474 or herbivory (**Table 1**; **Figure S3**). In general, the strength of local adaptation, measured as the

475 slope of the relationship between sexual fitness and climatic distance, was greater in the more

476 southern common garden location on each continent (**Figure 2**). For the number of flower heads,

477 the slope for Lafayette was 3.4× steeper than the slope for Mississauga and the slope for

478 Montpellier was 349× steeper than the slope for Uppsala. Similarly, for seed set mass, the slope

479 for Lafayette was 2.3× steeper than the slope for Mississauga and the slope for Montpellier was

480 384× steeper than the slope for Uppsala. The main effect of garden location affected all measures

481 of fitness and herbivory, which reflects the large differences in environment between the

gardens. By contrast, the main effect of continent of origin and its interactions with garden

location and/or climatic distance did not affect any measure of fitness or herbivory (**Table 1**).

Q2: Does T. repens exhibit stronger local adaptation in its native range than in its introduced

range?

We observed strong local adaptation in both the native and introduced ranges of *T. repens*.

Continent of origin, continent of the common garden, and climatic distance all interacted to

affect all quantitative measures of sexual fitness, survival, and growth rate of *T. repens* (**Table**

2). Specifically, populations only exhibited strong signatures of local adaptation when they were

transplanted to common gardens on the same continent from which they originated (**Figure 3**).

For number of flower heads, in common gardens located in the native range (Europe), the slope

for European *T. repens* populations was 12× steeper than the slope for North American

populations. In common gardens located in the introduced range (North America), the slope for

North American *T. repens* populations was 20× steeper than the slope for European populations.

Similarly, for seed set mass, in common gardens located in Europe, the slope for European *T.*

repens populations was 24× steeper than the slope for North American populations. In common

gardens located in North America, the slope for North American *T. repens* populations was 7.8×

steeper than the slope for European populations. The main effect of climatic distance affected

whether or not plants flowered and produced seeds, but climatic distance did not interact with the

continent of origin or the continent of the common garden to affect these components of fitness

(**Table 2**; **Figure S4**). However, climatic distance and continent of origin did interact to affect

maximum plant area and herbivory. Together across all gardens, plants from European origins

that were transplanted shorter climatic distances grew larger, despite experiencing greater levels

of herbivory, while size and herbivory for plants from North American origins were not

significantly affected by climatic distance overall (**Table 2**; **Figure S4**).

Q3: What are the roles of cyanogenesis and herbivory in driving local adaptation across space?

Both *Ac/ac* and *Li/li* exhibited strong latitudinal gene frequency clines among *T. repens*

populations from both continents (**Table S8**). However, the lack of an interaction between

latitude and continent of origin for both loci indicates no differences in the steepness of

cyanogenesis clines between the native and introduced ranges of *T. repens* (**Figure S5**). In the

full experiment, the main effect of cyanotype affected survival, whether or not plants produced

flowers and/or seeds, and maximum plant area, with cyanogenic plants generally exhibiting

greater fitness than acyanogenic plants, regardless of the environment in which the common

garden was found (**Table S9**). However, cyanotype alone did not affect the level of herbivory

occurring on *T. repens* plants, and cyanotype did not interact with either common garden

location or climatic distance to affect any measures of fitness or herbivory (**Table S9**).

Q4: What bioclimatic variables have the strongest impact on fitness across space in T. repens?

Climatic variables related to both temperature and precipitation predicted variation in fitness in

T. repens, but the most influential predictors varied between garden locations, with patterns

emerging based on latitude (**Tables S3-S6**, red highlights). For the full experiment, 58% of the

most influential predictors were temperature-based (mean annual temperature, maximum

temperature of the warmest month, or minimum temperature of the coldest month), while 42%

were precipitation-based (annual precipitation, precipitation of the wettest month, or

precipitation of the driest month). In the more southern gardens on each continent (Lafayette and

Montpellier), 67% of the most influential predictors were precipitation-based, while only 33%

were temperature-based. Conversely, in the more northern gardens on each continent

(Mississauga and Uppsala), 80% of the most influential predictors were temperature-based,

while only 20% of were precipitation-based. We did not find evidence that the native and

introduced ranges of *T. repens* were differentially affected by temperature or precipitation

variables. In the native range (Uppsala and Montpellier, combined), 57% of the most influential

predictors were temperature-based and 43% were precipitation-based. In the introduced range

(Mississauga and Lafayette, combined), 58% of the most influential predictors were temperature-

based, while 42% were precipitation-based.

Q5: Are there adaptation lags to climate change in T. repens?

North American populations of *T. repens* demonstrated evidence of an adaptation lag in the

northern garden in the introduced range (**Table 3**). Survival, seed set mass, and maximum plant

area all exhibited a significant quadratic relationship with ΔMAT between locations of origin and

the Mississauga common garden (**Figure 4**). A quadratic relationship centred on a ΔMAT of

zero would represent local adaptation, while a relationship centred on a ΔMAT > 0 would

represent an adaptation lag. For the proportion of plants survived per population, the peak of the

quadratic relationship experienced a significant shift to a climate of origin with a MAT 3.4°C

greater than Mississauga ($\chi^2_{1,4}$ = 4.159, P = 0.041, ΔAIC = −2.159), with the predicted maximum

survival being 5.3% greater than at a ΔMAT of zero. For mean seed set mass per population, the

peak of the quadratic relationship occurred at a MAT of 2.0°C greater than Mississauga, with the

predicted maximum seed set mass being 5.2% greater than at a ΔMAT of zero. However, this

shift was not statistically significant ($\chi^2_{1,5}$ = 0.640, P = 0.423, ΔAIC = 1.360). For mean

maximum plant area per population, the peak of the quadratic relationship experienced a shift to

a climate of origin with a MAT of 2.1°C ($\chi^2_{1,5}$ = 3.721, P = 0.054, ΔAIC = −1.721) greater than

Mississauga, with the predicted maximum plant area being 3.9% greater than at a ΔMAT of zero.

Along our transect of sampled populations, a shift in the peak of the quadratic relationship

towards populations 2.0-3.1°C warmer than the Mississauga common garden represents a 2.1-

3.2° shift in latitude towards the equator. Additionally, whether or not plants flowered or

produced seeds increased linearly with ΔMAT, indicating plants of warmer origins were more

likely to flower and produce seeds (**Figure 4**), while the number of flower heads and growth rate

did not exhibit any significant linear or quadratic trends with ΔMAT (**Table 3**).

Discussion

The aim of our study was to provide insight into the potential for rapid local adaptation in

introduced plant populations in response to novel environments across space and time. The

results of our transcontinental experiment in the native (Europe) and introduced (North America)

ranges of *T. repens* lead to two main conclusions: 1) rapid local adaptation to climatic variation

across space has contributed to the ecological success of *T. repens* in its introduced range; 2)

climate warming has resulted in some evidence of a weakening of local adaptation over time in

the introduced range of *T. repens.* Our conclusions are based on five experimental results. First,

T. repens exhibited local adaptation to climate overall, with greater fitness advantages for local

populations at lower latitudes (Q1). Second, in European common gardens, only native European

populations exhibited local adaptation, while in North American common gardens, only

introduced North American populations exhibited local adaptation (Q2). Third, cyanogenesis did

not differentially affect the fitness of *T. repens* in different environments (Q3). Fourth, both

temperature- and precipitation-based bioclimatic variables predicted fitness, but temperature was

a stronger determinant of fitness at higher latitudes, while precipitation was a stronger

determinant of fitness at lower latitudes (Q4). Fifth, there is some evidence consistent with an

adaptation lag in response to climate warming, but the signature was weak, and this pattern was

only clear in the northern common garden of the introduced range (Q5). Below we discuss our

results in the context of the role of rapid local adaptation in the success of biological invasions

and how introduced populations may (or may not) be able to adapt to future climate change.

Strength of local adaptation is dependent on latitude

Spatial variation in biotic and abiotic factors across a species' range could affect its ability to

adapt to different environments within that range. We found that *T. repens* exhibited significant

local adaptation to the abiotic environment based on a measure of climatic distance that

encompassed mean annual temperature and precipitation along with annual temperature and

precipitation extremes. However, local adaptation differed significantly in strength depending on

the latitude of common gardens. Specifically, local adaptation was stronger at lower latitudes in

both the native and introduced ranges of *T. repens* when fitness was measured as sexual

reproductive success (i.e., number of flower heads and seed set mass). Our results are consistent

with transplant experiments using other plant species, which found sexual fitness advantages for

local versus foreign plants to be more pronounced at lower latitudes (Colautti and Barrett, 2013;

Toräng et al., 2015). Alternatively, life history trade-offs across large latitudinal ranges could

result in differences in the observed strength of fitness advantages, depending on the measure of

fitness involved (Li et al., 2015; Bastias et al., 2024). In fact, life history trade-offs in *T. repens*

specifically have been shown to contribute to local adaptation across divergent environments,

with southern populations investing more into sexual reproduction and northern populations

investing more into vegetative growth (Wright et al., 2021). If warmer climates are generally

more conducive to sexual reproduction than cooler climates, this may explain why sexual fitness

advantages for local populations were more pronounced in Lafayette and Montpellier in our

study, while climatic distance had more limited effects on vegetative growth.

Local adaptation in native and introduced ranges

Invasive species have the potential to become more widespread and persistent by rapidly

adapting to novel environments. For example, latitudinal cyanogenesis clines in introduced

North American *T. repens* populations formed due to rapid local adaptation and sorting of

existing genotypes post-introduction (Kooyers and Olsen, 2012; Kooyers and Olsen, 2013; Olsen

et al., 2021). We found that populations of *T. repens* only exhibited strong signatures of local

adaptation to climate when transplanted into common gardens within the same range as they

originated. The observed strong local fitness advantages for North American populations within

the North American gardens are therefore indicative of rapid local adaptation of *T. repens* within

the past 400 years since its introduction to North America. Selection has been shown to be

particularly strong for introduced populations, setting a precedent for rapid local adaptation

among invasive species on even shorter timescales than exhibited by *T. repens* (Colautti and Lau,

2015; Simón-Porcar et al., 2021). A meta-analysis by Oduor et al. (2016) found that invasive

populations exhibited local adaptation just as frequently as native populations, and that the

strength of local adaptation was not affected by time since introduction (ranging from as little as

20 to as many as 500 years). A reciprocal transplant experiment with *Lythrum salicaria* indicated

local adaptation to all three common garden sites spanning a 10° latitudinal range, which

occurred in ~200 years since its introduction to eastern North America (Colautti and Barrett,

2013). Even faster still is adaptive evolution in the form of trait divergence between coastal and

inland environments in *Eschscholzia californica* in its introduced range, which has occurred in

the past 150 years (Leger and Rice, 2007). Similarly, clines in flowering phenology of invasive

Hypericum canariense have evolved within only 50 years (Dlugosch and Parker, 2008). Taken

together, our results strengthen the mounting evidence that rapid local adaptation to climate plays

an important role in the ecological success of invasive species in their new range.

Interestingly, *T. repens* populations from the introduced range did not exhibit signatures

of local adaptation to similar climates when transplanted back into their native range. Similarly,

native populations did not maintain their strong signatures of local adaptation to climate when

transplanted into the introduced range. These findings demonstrate a clear cost of rapid

adaptation, with local fitness advantages not extending across continents, even when accounting

for similarity among climates. Explanations for this lack of maintenance of local adaptation

across continents could be due to unmeasured biotic and abiotic aspects of the environment.

Abiotic explanations for the lack of local adaptation across continents could involve

shifts in photoperiod or precipitation regimes between Europe and North America due to similar

climates on the two continents being associated with different latitudes. In general, European

environments have longer day lengths in summer and shorter day lengths in winter than those of

similar climates in North America, with differences being more pronounced at progressively

higher latitudes. This photoperiodic shift between the continents could have affected the

initiation of flowering and/or the timing of peak flowering for populations that were transplanted

across continents, leading to differences in the strength of local adaptation. Halbritter et al.

(2015) found that local adaptation was still evident in reciprocal transplants conducted between

high elevation and high latitude *Plantago* populations with similar climates, allowing for

speculation of a role of photoperiod in affecting fitness. Additionally, populations of an obligate

long-day plant, *Mimulus guttatus*, exhibit clinal variation in the critical photoperiod required to

initiate flowering, potentially indicating a role of photoperiod in driving local adaptation

(Kooyers et al., 2015). Saikkonen et al. (2012) speculated that the transition to less extreme

yearly variation in photoperiod at lower latitudes might facilitate greater invasion success of

European species introduced to North America than vice versa, which could be especially true of

a facultative long-day plant like *T. repens* (Beatty, 1956). However, our results do not provide

evidence of increased success of European populations in North American gardens compared to

that of North American populations in European gardens. Additionally, although we accounted

for mean annual precipitation and yearly precipitation extremes in our study, these variables do

not fully capture the exact timing and volume of precipitation, and the resulting hydrological

environment experienced by *T. repens* plants in Europe versus North America. For example, in

the Montpellier garden in southern Europe, plants will experience a Mediterranean climate that is

not only drier but also more seasonally variable, while in the Lafayette garden in southern North

America, plants will experience a sub-tropical environment that is wetter with more limited

seasonal variation. This shift in precipitation availability has caused introduced *T. repens*

populations to evolve greater drought avoidance as opposed to the more desiccation tolerant

strategy observed in Mediterranean populations (Hendrickson et al., 2024).

The biotic environment could also partially explain the lack of local adaptation across

continents based on climatic distance (Hargreaves et al, 2020). *Trifolium repens* is a legume that

forms mutualistic relationships with N-fixing bacteria in root nodules. Murray-Stoker and

Johnson (2024) have demonstrated the existence of locally adapted populations in the *T. repens-*

Rhizobia mutualism, indicating that local adaptation to biotic factors is possible across

environments in this system. Furthermore, in line with classic hypotheses in invasion biology,

herbivory could exhibit substantial differences across environments based on expected shifts in

herbivore communities in different climates or between continents, which could facilitate success

of a plant species post-introduction (Coley et al., 1985; Blossey and Nötzold, 1995; Keane and

Crawley, 2002). If herbivory was likely to explain our results of strong local adaptation within

continents but not across continents, we might have expected to see signatures of local

adaptation based on the presence of chemical defenses in *T. repens* plants.

Cyanogenesis did not affect local adaptation in either the native or introduced range

We tested whether the presence of latitudinal clines in plant defense indicates a role of defense

mechanisms in driving local adaptation in *T. repens*. Among our sampled populations, we

demonstrated rapid evolution of latitudinal clines in the presence of both *Ac* and *Li* alleles in the

introduced range of *T. repens*, based on their similar strength to clines in the native range.

Contrary to our results, all previous work has indicated weaker clines in the introduced range

than the native range (Daday, 1954a; Daday, 1958; Innes et al., 2022). Interestingly, our results

also indicates that neither cyanogenesis nor the presence/absence of cyanogenic glucosides or

linamarase exhibited differential effects on fitness across our common gardens. In fact, local

plants exhibited signatures of local adaptation despite experiencing similar or greater levels of

herbivory than populations transplanted from further climatic distances. Our study therefore does

not demonstrate a role of cyanogenesis or its metabolic components, nor a role of herbivory, in

driving local adaptation of *T. repens*, with no evidence of interactions between cyanotype and

climatic distance in influencing fitness. Previous studies on local adaptation in *T. repens* found

that HCN, cyanogenic glucosides, and linamarase only had marginal, if any, effects on fitness

and local adaptation (Wright et al., 2018; Wright et al., 2021). Further, *T. repens* has not

consistently experienced changes in cyanogenesis clines in response to recent climate change,

which may indicate only marginal involvement of cyanogenesis and/or its metabolic components

in local adaptation (Innes et al., 2022). Instead, divergence in life history strategies in response to

climatic factors have been identified as stronger drivers of local adaptation (Wright et al., 2018;

Wright et al., 2021), which is consistent with our results.

Our findings contrast with previous work that has speculated on the involvement of trade-

offs in the costs and benefits of cyanogenesis in different environments. Namely, previous

studies have hypothesized ecological costs of cyanogenesis in cooler climates due to freezing

autotoxicity (Daday, 1954a; Daday, 1965) or exacerbated allocation costs of producing the

metabolic components of HCN under freezing conditions (Kooyers et al., 2018; but see Fadoul et

al., 2023) as possible explanations for low frequencies of cyanogenesis in cooler climates. In our

study, cyanogenic plants often demonstrated a fitness advantage, even in Mississauga, which

does experience freezing temperatures for much of the year. This finding indicates that even in

cooler climates, the benefits of producing HCN as an anti-herbivore defense could outweigh the

costs of its production. However, despite overall effects of cyanotype on fitness in our common

gardens, we did not find evidence for cyanotype as deterrent to herbivory or as a driver of local

adaptation in *T. repens*. In fact, cyanotype did not affect the level of herbivory occurring on *T.*

repens plants, indicating that greater prevalence of cyanogenesis in warmer climates is likely not

based on an increased ability to defend against greater herbivore pressure.

Alternatively, there is evidence that the ability to produce cyanogenic glucosides and/or

linamarase could be advantageous in periods of drought through the storage and recycling of

nitrogen (Møller, 2010; Kooyers et al., 2014; Machingura et al., 2016). The first growing season

in Mississauga was uncharacteristically dry, resulting in the need for supplemental watering of

the common garden to avoid mortality of the full experiment. It is possible that the dry

conditions may have also contributed to a fitness advantage for *Ac* and/or *Li* plants. However,

any advantages of particular cyanotypes within a given common garden environment did not

confer higher fitness to local plants as opposed to foreign plants based on our measure of

climatic distance. Furthermore, we were unable to harvest enough tissue to determine the

cyanotype of most of the plants in the study due to concerns over effects of tissue harvest on

plant fitness and/or mortality, so the strength of our conclusions with respect to the role of

cyanogenesis in adaptation are limited by the relatively low sample size for all statistical

analyses involving cyanotype. Further field studies are needed to more explicitly test the costs

and benefits of HCN, cyanogenic glucosides, and linamarase, and their effects on local

adaptation to climate in *T. repens* across large environmental gradients.

Spatial variation in environmental drivers of local adaptation

While *T. repens* exhibits local adaptation in its native and introduced ranges based on an overall

measure of climatic distance, populations may spatially vary in the individual bioclimatic

variables responsible for local adaptation. Our results across both ranges demonstrate that at

higher latitudes, where local adaptation is weaker, temperature-based variables are stronger

determinants of fitness, while at lower latitudes, where local adaptation is stronger, precipitation-

based variables are stronger determinants of fitness. Therefore, it is possible that *T. repens* adapts

more readily to local precipitation regimes than local temperature regimes, resulting in stronger

signatures of local adaptation in lower latitude common gardens, consistent between the native

and introduced ranges. This result is supported by a meta-analysis by Siepielski et al. (2017),

which found both precipitation and potential evapotranspiration to be stronger predictors of

selection than temperature across numerous terrestrial taxa. Alternatively, our results could be

affected by the specific environments chosen for the common gardens. In our study, the climates

in Lafayette and Montpellier are among the warmest that occur in the range of *T. repens*, and

therefore could have been especially difficult to tolerate for populations transplanted from cooler

climates. In turn, Uppsala was somewhat isolated, both geographically and climatically, from the

sampled *T. repens* populations, which could have affected our ability to detect local adaptation

(or maladaptation) in that environment. Lastly, it is possible that recent climate warming may

have already disrupted the ability of local populations to outperform foreign populations, thereby

impacting the strength of local adaptation in certain locations.

Evidence of adaptation lags in response to climate warming

The ability of populations to adapt to recent rapid climate warming over time is an important

mechanism by which species can mitigate the risk of local extinction and other downstream

ecological consequences. Previous studies have demonstrated the incidence of adaptation lags,

whereby populations originating from slightly warmer climates would outperform populations

with very similar climatic origins to the common garden location (Wilczek et al., 2014; Aguirre-

Liguori et al., 2019; Anderson and Wadgymar, 2020). These studies indicate the potential for a

disruption in local adaptation, and even the possibility of maladaptation, due to climate warming

that is occurring faster than local populations are able to adapt.

In our northern common garden in North America (Mississauga), populations with

slightly warmer origins (2-3°C in MAT) outperformed local populations in survival and biomass

production (i.e., a proxy for vegetative reproduction) but not measures of sexual fitness. This

finding may signal the presence of an adaptation lag and may also provide a partial explanation

for the observed weaker local adaptation in Mississauga compared to Lafayette. However, the

generalizability of our results should be considered carefully for a few reasons. First, contrary to

our expectations, we provide no evidence of either local adaptation or maladaptation in Uppsala.

In a large-scale common garden experiment with European populations of *Arabidopsis thaliana*,

Wilczek et al. (2014) found that adaptation lags were more pronounced in higher latitude

common garden locations, which was not the case in our study. It would also be pertinent to be

able to test for spatial lags in the native range of *T. repens* to which we could directly compare

spatial lags in the introduced range. Second, we tested for the presence of adaptation lags based

solely on MAT, and Mississauga was the common garden location where MAT most

consistently affected plant fitness compared to other bioclimatic variables. Fitness in lower

latitude common gardens was instead more heavily influenced by precipitation-based variables,

which may not be changing as consistently as are temperature-based variables due to

anthropogenic activity. It is therefore possible that adaptation lags to changing environments

may not occur consistently across different geographic locations, based on the specific biotic and

abiotic factors that most strongly impact local adaptation in those environments. Third, the MAT

at the location of the Mississauga common garden in 2020 and 2021 was 1.6°C and 2.0°C

warmer, respectively, than the average from the historical time period (1970-2000) used to

calculate climatic distances in this study (data extracted from ClimateNA; Wang et al., 2016).

Therefore, it may not be unexpected to see populations from slightly warmer origins outperform

local populations. The adaptation lag for survival does extend to a peak 3.4°C warmer than

Mississauga, and the proportion of plants producing flowers and/or seeds increased linearly with

ΔMAT, which still provides evidence for warm-adapted populations having an advantage that

exceeds expectations based on the MAT during the years of the experiment. However, across our

entire study, we are unable to make broad conclusions about future vulnerability of populations

based on the rate of environmental change exceeding the rate of adaptation. Instead, we conclude

that the fine-tuned nature of local adaptation (in terms of variation in strength, prominent driving

factors, and the potential for maladaptation) to climate in *T. repens* highlights the multifaceted

nature of rapid evolution in response to changing environments.

Limitations

While our study offers strong tests of local adaptation to both biotic and abiotic factors in the

native and introduced ranges of a cosmopolitan plant, there were limitations that should be

considered when interpreting our findings. In particular, we are unable to determine the relative

strength of soil conditions and mutualistic interactions compared to climate as drivers of local

adaptation. Planting *T. repens* individuals from each population along with the natural soil and

microbial communities from their location of origin was infeasible considering our study design

and governmental regulations on importation of soil and bacteria. Instead, the best alternative

was to maintain consistent treatment of all individuals in all gardens, with *T. repens* individuals

planted into the natural soil at each common garden site. It is important to recognize that local

edaphic factors have the potential to provide fitness advantages in *T. repens*, as has been

demonstrated in other systems (Macel et al., 2007). It is also possible that mutualistic biotic

interactions between *T. repens* and nitrogen-fixing bacteria could be involved in providing

fitness advantages to local populations (Murray-Stoker and Johnson, 2024). Despite these

limitations, signatures of local adaptation to climate were still evident across multiple measures

of fitness, strengthening our confidence in the observed patterns.

Conclusions

Our study provides insight into the ability of introduced populations of a widespread invasive

plant to adapt to novel environments across space and over time. Signatures of local adaptation

were only evident when populations were transplanted within the same range as they originated,

indicating both strong and rapid local adaptation in an introduced range of *T. repens* in less than

400 years since introduction. Interestingly, the strength of local adaptation and the most

important environmental predictors of plant fitness differed in the north and south of both the

native and introduced range. Specifically, temperature was the most important driver of local

adaptation in northern gardens, whereas precipitation was most important in southern gardens.

This result indicates that there may be predictable drivers of local adaptation of invasive

populations to novel environments, and that these drivers may vary along broad latitudinal

gradients. Despite evidence of rapid adaptation to changing environments across space, the

northern garden in the introduced range also provided some evidence of local maladaptation

compared to populations with slightly warmer origins, which is characteristic of an adaptation

lag. Therefore, while local adaptation has occurred quickly in introduced populations of *T.*

repens across space, a lack of rapid adaptation to environmental change over time could

eventually result in severe erosion of fitness advantages for local populations in the future.

Continuing to further our understanding of the ability of native and introduced populations to

rapidly adapt to anthropogenic disturbance in space and time is vital to the integrity and

maintenance of biodiversity in the face of future environmental change.

Acknowledgements

We thank James Santangelo for assistance with initial seed collection, and Happy Gill,

Jashandeep Nijjar, and Ruth Thomas for assisting with the generation of F1 plants for use in the

experiment. We are indebted to Aaron Albano, Barry Albano, Paula Albano, Sarah Albano,

Andre Daugeureaux, Hayden Fargo, Axel Hjelm, Antoniette Jansen, Eunice Karinho Betancourt,

Erik Karlsson, Andreas Lundgren, Kavya Manikonda, Lindsay Miles, David Murray-Stoker,

Bernabé Ramirez, Brittany Sutherland, Olivia Toth, Andrea Turcu, Mariona Vinyeta Medina,

and Vicki Zhang for assistance in the field. Keerath Bhachu, Elana Maria, Zain Nasrullah, Nial

Navaratne, Narindra Persaud, and Isabellio Vessio assisted with sample processing. We also

thank the Louisiana Optical Network (LONI) for providing computation resources, along with

Kris Horvath, Joan Lee, Kent Moore, Diane Ross, Vera Velasco, the technical platform

"Plateforme des Terrains d'Expérience" (CEFE-CNRS), and the University of Louisiana

Ecology Center for facilitating and supporting the set-up and execution of the experiment. This

research was supported by an NSERC Discovery Grant, Canada Research Chair, and NSERC

EWR Steacie Fellowship to MTJJ, a CNRS-University of Toronto Ph.D. Student Travel Grant to

MTJJ and CV, NSF grants OIA-1920858 and DBI-2244712 to NJK, and an NSERC CGS

Doctoral Award to LJA.

834   **Conflict of Interest Statement**

835   The authors declare no conflict of interest.

837   **References**

Adams, J. M., Fang, W., Callaway, R. M., Cipollini, D., Newell, E., Transatlantic Acer

platanoides Invasion Network (TRAIN). (2009). A cross-continental test of the Enemy

Release Hypothesis: leaf herbivory on *Acer platanoides* (L.) is three times lower in North

America than in its native Europe. *Biological Invasions*, *11*, 1005-1016.

Agrawal, A. A., Kotanen, P. M. (2003). Herbivores and the success of exotic plants: a

phylogenetically controlled experiment. *Ecology Letters*, *6*, 712-715.

Agrawal, A. A. (2007). Macroevolution of plant defense strategies. *Trends in Ecology and

Evolution*, *22*, 103-109.

Aguirre-Liguori, J. A., Ramírez-Barahona, S., Tiffin, P., Eguiarte, L. E. (2019). Climate change

is predicted to disrupt patterns of local adaptation in wild and cultivated maize.

Proceedings of the Royal Society B, *286*, 20190486.

Albano, L. J., Johnson, M. T. J. (2023). Interactions environmental factors drive selection on

cyanogenesis in *Trifolium repens*. *Oikos*, *2023*, e09629.

Allendorf, F. W., Lundquist, L. L. (2003). Introduction: population biology, evolution, and

control of invasive species. *Conservation Biology*, *17*, 24-30.

Anderson, J. T., Song, B.-H. (2020). Plant adaptation to climate change – Where are we? *Journal*

of Systematics and Evolution, *58*, 533-545.

Anderson, J. T., Wadgymar, S. M. (2020). Climate change disrupts local adaptation and favours

upslope migration. *Ecology Letters*, *23*, 181-192.

Angseesing, J. P. A. (1974). Selective eating of the acyanogenic form of *Trifolium*

repens. *Heredity*, *32*, 73.

Anstett, D. N., Nunes, K. A., Baskett, C., Kotanen, P. M. (2016). Sources of controversy

surrounding latitudinal patterns in herbivory and defense. *Trends in Ecology & Evolution*,

31, 789-802.

Anstett, D. N., Ahern, J. R., Johnson, M. T. J., Salminen, J.-P. (2018). Testing for latitudinal

gradients in defense at the macroevolutionary scale. *Evolution*, *72*, 2129-2143.

Atkins, K. E., Travis, J. M. J. (2010). Local adaptation and the evolution of species' ranges under

climate change. *Journal of Theoretical Biology*, *266*, 449-457.

Atwood, S. S., Sullivan, J. T. (1943). Inheritance of a cyanogenetic glucoside and its hydrolyzing

enzyme in *Trifolium repens*. *Journal of Heredity*, *34*, 311-320.

Baskett, C. A., Schemske, D. W. (2018). Latitudinal patterns of herbivore pressure in a

temperature herb support the biotic interactions hypothesis. *Ecology Letters*, *21*, 578-587.

Bastias, C. C., Estarague, A., Vile, D., Gaignon, E., Lee, C.-R., Exposito-Alonso, M., … ,

Vasseur, F. (2024). Ecological trade-offs drive phenotypic and genetic differentiation of

Arabidopsis thaliana in Europe. *Nature Communications*, *15*, 5185.

Battlay, P., Hendrickson, B. T., Hendez-Reneau, J. I., Santangelo, J. S., Albano, L. J., Wilson, J.,

… , Kooyers, N. J. (2024). Structural variants underlie parallel adaptation following

global invasion. *biorxiv*. doi: 10.1101/2024.07.09.602765

Beatty, D. W. (1956). Effect of photoperiod and temperature on flowering of white clover

(*Trifolium repens*). Iowa State University.

Bidart-Bouzat, M. G., Imeh-Nathaniel, A. (2008). Global change effects on plant chemical

defenses against insect herbivores. *Journal of Integrative Plant Biology*, *50*, 1339-1354.

Blossey, B., Nötzold, R. (1995). Evolution of increased competitive ability in invasive

nonindigenous plants: A hypothesis. *Journal of Ecology*, *83*, 887-889.

Bontrager, M., Muir, C. D., Mahony, C., Gamble, D. E., Germain, R. M., Hargreaves, A. L., … ,

Angert, A. L. (2020). Climate warming weakens local adaptation. *biorxiv*. doi:

10.1101/2020.11.01.364349

Brighton, F., Horne, M. T. (1977). Influence of temperature on cyanogenic polymorphisms.

Nature, *265*, 437-438.

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., … ,

Bolker, B. M. (2014) glmmTMB balances speed and flexibility among packages for zero-

inflated generalized linear mixed modeling. *The R Journal*, *9*, 378-400.

Burdon, J. J. (1983). *Trifolium repens* L. *Journal of Ecology*, *71*, 307-330.

Burns, J. E., Ashman, T.-L., Steets, J. A., Harmon-Threatt, A., Knight, T. M. (2011). A

phylogenetically controlled analysis of the toles of reproductive traits in plant invasions.

Oecologia, *166*, 1009-1017.

Byers, J. E. (2002). Impact of non-indigenous species on natives enhanced by anthropogenic

alteration of selection regimes. *Oikos*, *97*, 449-458.

Callaway, R. M., Ridenour, W. M. (2004). Novel weapons: invasive success and the evolution of

increased competitive ability. *Frontiers in Ecology and the Environment*, *2*, 436-443.

Colautti, R. I., Barrett, S. C. H. (2013). Rapid adaptation to climate facilitates range expansion of

an invasive plant. *Science*, *342*, 364-367.

Colautti, R. I., Lau, J. A. (2015). Contemporary evolution during invasion: evidence for

differentiation, natural selection, and local adaptation. *Molecular Ecology*, *24*, 1999-

2017.

Coley, P. D., Bryant, J. P., Chapin, F. S. (1985). Resource availability and plant antiherbivore

defense. *Science*, *230*, 895-899.

Coley, P. D., Barone, J. A. (1996). Herbivory and plant defenses in tropical forests. *Annual*

Review of Ecology and Systematics, *27*, 305-335.

Corkill, L. D. (1940). Cyanogenesis in white clover (*Trifolium repens* L.). I. Cyanogenesis in

single plants. *New Zealand Journal of Science and Technology, 22*, 65-67.

Costan, C.-A., Godsoe, W. K., Bufford, J. L., Marris, J. W. M., Hulme, P. E. (2022). Can the

enemy release hypothesis explain the success of *Rumex* (Polygonaceae) species in an

introduced range? *Biological Invasions*, *24*, 2665-2677.

Daday, H. (1954a). Gene frequencies in wild populations of *Trifolium repens*. I. Distribution by

latitude. *Heredity*, *8*, 61-78.

Daday, H. (1954b). Gene frequencies in wild populations of *Trifolium repens.* II. Distribution by

altitude. *Heredity*, *8*, 377-384.

Daday, H. (1958). Gene frequencies in wild populations of *Trifolium repens.* III. World

distribution. *Heredity*, *12*, 169-184.

Daday, H. (1965). Gene frequencies in wild populations of *Trifolium repens.* IV. Mechanism of

natural selection. *Heredity*, *20*, 355-365.

Dirzo, R., Harper, J. L. (1982). Experimental studies on slug-plant interactions: III. Differences

in the acceptability of individual plants of *Trifolium repens* to slugs and snails. *Journal of*

Ecology, *70*, 101-117.

Dlugosch, K. M., Parker, I. M. (2008a). Invading populations of an ornamental shrub show rapid

life history evolution despite genetic bottlenecks. *Ecology Letters*, *11*, 701-709.

Dlugosch, K. M., Parker, I. M. (2008b). Founding events in species invasions: genetic variation,

adaptive evolution, and the role of multiple introductions. *Molecular Ecology*, *17*, 431-

449.

Dostálek, T., Rokaya, M. B., Münzbergová, Z. (2020). Plant palatability and trait responses to

experimental warming. *Scientific Reports*, *10*, 10526.

Easlon, H. M., Bloom, A. J. (2014). Easy Leaf Area: Automated digital image analysis for rapid

and accurate measurement of leaf area. *Applications in Plant Sciences*, *2*, 1400033.

Ehrlich, P. R., Raven, P. H. (1964). Butterflies and plants: A study in coevolution. *Evolution*, *18*,

586-608.

Endler, J. A. (1977). Geographic variation, speciation, and clines. Princeton University Press.

Princeton, NJ, USA.

Erfmeier, A. (2013). Constraints and release at difference scales – The role of adaptation in

biological invasions. *Basic and Applied Ecology*, *14*, 281-288.

Estoup, A., Ravigné, V., Hufbauer, R., Vitalis, R., Gautier, M., Facon, B. (2016). Is there a

genetic paradox of biological invasion? *Annual Review of Ecology, Evolution, and*

Systematics, *47*, 51-72.

Exposito-Alonso, M., 500 Genomes Field Experiment Team, Burbano, H. A., Bossdorf, O.,

Nielsen, R., Weigel, D. (2019). Natural selection on the *Arabidopsis thaliana* genome in

present and future climates. *Nature*, *573*, 126-129.

Fadoul, H. E., Albano, L. J., Bergman, M. E., Phillips, M. A., Johnson, M. T. J. (2023).

Assessing the benefits and costs of the hydrogen cyanide antiherbivore defense in

Trifolium repens. *Plants*, *12*, 1213.

Fine, P. V. A., Miller, Z. J., Mesones, I., Irazuzta, S., Appel, H. M., Stevens, M. H. H., … ,

Coley, P. D. (2006). The growth-defense trade-off and habitat specialization by plants in

Amazonian forests. *Ecology*, *87*, S150-S162.

Fox, J., Weisberg, S. (2019). An R companion to applied regression, Third edition. Sage,

Thousand Oaks, CA, USA.

Fugère, V., Hendry, A. P. (2018). Human influences on the strength of phenotypic selection.

Proceedings of the National Academy of Sciences. *115*, 10070-10075.

Genton, B. J., Kotanen, P. M., Cheptou, P.-O., Adolphe, C., Shykoff, J. A. (2005). Enemy

release but no evolutionary loss of defence in a plant invasion: an inter-continental

reciprocal transplant experiment. *Oecologia*, *146*, 404-414.

Gillespie, J. H., Turelli, M. (1989). Genotype-environment interactions and the maintenance of

polygenic variation. *Genetics*, *121*, 129-138.

Gleadow, R. M., Woodrow, I. E. (2002). Constraints on effectiveness of cyanogenic glycosides

in herbivore defense. *Journal of Chemical Ecology*, *28*, 1301-1313.

Griffiths, A. G., Moraga, R., Tausen, M., Gupta, V., Bitton, T. P., Campbell, M. A., … ,

Anderson, S. U. (2019). Breaking free: The genomics of allopolyploidy-facilitated niche

expansion in white clover. *The Plant Cell*, *31*, 1466-1487.

Hahn, P. G., Agrawal, A. A., Sussman, K. I., Maron, J. L. (2019). Population variation,

environmental gradients, and the evolutionary ecology of plant defense against herbivory.

The American Naturalist, *193*, 20-34.

Halbritter, A. H., Billeter, R., Edwards, P. J., Alexander, J. M. (2015). Local adaptation at range

edges: comparing elevation and latitudinal gradients. *Journal of Evolutionary Biology*,

28, 1849-1860.

Hargreaves, A. L., Germain, R. M., Bontrager, M., Persi, J., Angert, A. L. (2020). Local

adaptation to biotic interactions: A meta-analysis across latitudes. *The American*

Naturalist, *195*, 395-411.

Hendrickson, B. T., Stamps, C., Patterson, C. M., Strickland, H., Foster, M., Kim, A., … ,

Kooyers, N. J. (2024). Evolution of drought resistant strategies following the introduction

of white clover (*Trifolium repens* L.). *biorxiv*. doi: 10.1101/2024.09.03.611023

Hendry, A. P. (2017). Eco-evolutionary dynamics. Princeton University Press. Princeton, NJ,

USA.

Hereford, J. (2009). A quantitative survey of local adaptation and fitness trade-offs. *The*

American Naturalist, *173*, 579-588.

Herms, D. A., Mattson, W. J. (1992). The dilemma of plants: to grow or defend. *Quarterly Review of*

Biology, *67*, 478-478.

Hoffmann, A. A., Sgrò, C. M. (2011). Climate change and evolutionary adaptation. *Nature*, *470*,

479-485.

Hughes, M. A. (1991). The cyanogenic polymorphism in *Trifolium repens* L. (white clover).

Heredity, *66*, 105-115.

Hulme, P. E. (2017). Climate change and biological invasions: evidence, expectations, and

response options. *Biological Reviews*, *92*, 1297-1313.

Innes, S., Santangelo, J. S., Kooyers, N. J., Olsen, K. M., Johnson, M. T. J. (2022). Evolution in

response to climate in the native and introduced ranges of a globally distributed plant.

Evolution, *76*, 1495-1511.

IPBES. (2019). Global assessment report on biodiversity and ecosystem services of the

Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.

Brondizio, E. S., Settele, J., Díaz, S., Ngo, H. T. (editors). IPBES secretariat, Bonn,

Germany.

Jeschke, J. M., Pyšek, P. (2018). Tens rule. In *Invasion biology: Hypotheses and evidence* (pp.

124-132). CAB International. Wallingford, UK.

Johnson, M. T. J., Rasmann, S. (2011). The latitudinal herbivory-defence hypothesis takes a

detour on the map. *New Phytologist*, *191*, 589-592.

Johnson, M. T. J., Bertrand, J. A., Turcotte, M. M. (2016). Precision and accuracy in quantifying

herbivory. *Ecological Entomology*, *41*, 112-121.

Kakes, P. (1989). An analysis of the costs and benefits of the cyanogenic system in *Trifolium*

repens L. *Theoretical and Applied Genetics*, *77*, 111-118.

Kassambara, A., Mundt, F. (2020). factoextra: Extract and visualize the results of multivariate

data analyses. R package version 1.0.7. https://CRAN.R-project.org/package=factoextra

Kawecki, T. J., Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, *7*, 1225-
1241.

Keane, R. M., Crawley, M. J. (2002). Exotic plant invasions and the enemy release hypothesis.
Trends in Ecology & Evolution, *17*, 164-170.

Kjærgaard, T. (2003). A plant that changed the world: The rise and fall of clover 1000-2000.
Landscape Research, *28*, 41-49.

Kolar, C. S., Lodge, D. M. (2001). Progress in invasion biology: predicting invaders. *Trends in*
Ecology & Evolution, *16*, 199-204.

Kooyers, N. J., Olsen, K. M. (2012). Rapid evolution of an adaptive cyanogenesis cline in
introduced North American white clover (*Trifolium repens* L.). *Molecular Ecology*, *21*,
2455-2468.

Kooyers, N. J., Olsen, K. M. (2013). Searching for the bull's eye: agents and targets of selection
vary among geographically disparate cyanogenesis clines in white clover (*Trifolium*
repens L.). *Heredity*, *11*, 495-504.

Kooyers, N. J., Gage, L. R., Al-Lozi, A., Olsen, K. M. (2014). Aridity shapes cyanogenesis cline
evolution in white clover (*Trifolium repens* L.). *Molecular Ecology*, *23*, 1053-1070.

Kooyers, N. J., Greenlee, A. B., Colicchio, J. M., Oh, M., Blackman, B. K. (2015). Replicate
altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress
in annual *Mimulus guttatus*. *New Phytologist*, *206*, 152-165.

Kooyers, N. J., Hartman Bakken, B., Ungerer, M. C., Olsen, K. M. (2018). Freeze-induced
cyanide toxicity does not maintain the cyanogenesis polymorphism in white clover
(*Trifolium repens*). *American Journal of Botany*, *105*, 1224-1231.

Kooyers, N. J., Colicchio, J. M., Greenlee, A. B., Patterson, E., Handloser, N. T., Blackman, B.

K. (2019). Lagging adaptation to climate supersedes local adaptation to herbivory in an

annual monkeyflower. *The American Naturalist*, *194*, 541-557.

Kuo, W.-H., Zhong, L., Wright, S. J., Goad, D. M., Olsen, K. M. (2024). Beyond cyanogenesis:

Temperature gradients drive environmental adaptation in North American white clover

(*Trifolium repens* L.). *Molecular Ecology*, *33*, e17484.

Lau, J. A., terHorst, C. P. (2015). Causes and consequences of failed adaptation to biological

invasions: the role of ecological constraints. *Molecular Ecology*, *24*, 1987-1998.

Lee, C.-R., Mitchell-Olds, T. (2011). Quantifying effects of environmental and geographical

factors on patterns of genetic differentiation. *Molecular Ecology*, *20*, 4631-4642.

Leger, E. A., Rice, K. J. (2007). Assessing the speed and predictability of local adaptation in

invasive California poppies (*Eschscholzia californica*). *Journal of Evolutionary Biology*,

20, 1090-1103.

Leimu, R., Fischer, M. (2008). A meta-analysis of local adaptation in plants. *PLoS One*, *3*,

e4010.

Lemoine, N. P., Drews, W. A., Burkepile, D. E., Parker, J. D. (2013). Increased temperature

alters feeding behaviour of a generalist herbivore. *Oikos*, *122*, 1669-1678.

Li, X.-M., She, D.-Y., Zhang, D.-Y., Liao, W.-J. (2015). Life history trait differentiation and

local adaptation in invasive populations of *Ambrosia artemisiifolia* in China. *Oecologia*,

177, 669-677.

Macel, M., Lawson, C. S., Mortimer, S. R., Šmilauerova, M., Bischoff, A., Crémieux, L., … ,

Steinger, T. (2007). Climate vs. soil factors in local adaptation of two common plant

species. *Ecology*, *88*, 424-433.

Machingura, M., Salomon, E., Jez, J. M., Ebbs, S. D. (2016). The β-cyanoalanine synthase

pathway: beyond cyanide detoxification. *Plant, Cell and Environment*, *39*, 2329-2341.

Maron, J. L., Vilà, M., Bommarco, R., Elmendorf, S., Beardsley, P. (2004). Rapid evolution of

an invasive plant. *Ecological Monographs*, *74*, 261-280.

Massad, T. J., Dyer, L. A. (2010). A meta-analysis of the effects of global environmental change

on plant-herbivore interactions. *Arthropod-Plant Interactions*, *4*, 181-188.

Miles, L. S., Breitbart, S. T., Wagner, H. H., Johnson, M. T. J. (2019). Urbanization shapes the

ecology and evolution plant-arthropod herbivore interactions. *Frontiers in Ecology and*

Evolution, *7*, 310.

Møller, B. L. (2010). Functional diversifications of cyanogenic glucosides. *Current Opinion in*

Plant Biology, *13*, 338-347.

Moreira, X., Castagneyrol, B., Abdala-Roberts, L., Berny-Mier y Teran, J. C., Timmermans, B.

G. H., Bruun, H. H., … , Tack, A. J. M. (2018). Latitudinal variation in plant chemical

defences drives latitudinal patterns of leaf herbivory. *Ecography*, *41*, 1124-1134.

Murray-Stoker, D., Johnson, M. T. J. (2024). Mosaic of local adaptation between white clover

and rhizobia along an urbanization gradient. *Journal of Ecology*, *112*, 1150-1163..

O'Donnell, M. S., Ignizio, D. A. (2012). Bioclimatic predictors for supporting ecological

applications in the conterminous United States. *US Geological Survey Data Series*, *10*, 4-

9.

Oduor, A. M. O., Leimu, R., van Kleunen, M. (2016). Invasive plant species are locally adapted

just as frequently and at least as strongly as native plant species. *Journal of Ecology*, *104*,

957-968.

Olsen, K. M., Sutherland, B. L., Small, L. L. (2007). Molecular evolution of the *Li/li* chemical

defence polymorphism in white clover (*Trifolium repens* L.). *Molecular Ecology*, *16*,

4180-4193.

Olsen, K. M., Small, L. L. (2018). Micro- and macroevolutionary adaptation though repeated

loss of a complete metabolic pathway. *New Phytologist*, *219*, 757-766.

Olsen, K. M., Goad, D. M., Wright, S. J., Dutta, M. L., Myers, S. R., Small, L. L., Li, L.-F.

(2021). Dual-species origin of an adaptive chemical defense polymorphism. *New*

Phytologist, *232*, 1477-1487.

Pannell, J. R., Auld, J. R., Brandvain, Y., Burd, M., Busch, J. W., Cheptou, P.-O., … , Winn, A.

A. (2015). The scope of Baker's law. *New Phytologist*, *208*, 656-667.

Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annual*

Review of Ecology, Evolution, and Systematics. *37*, 637-669.

Pyšek, P., Hulme, P. E., Simberloff, D., Bacher, S., Blackburn, T. M., Carlton, J. T. … ,

Richardson, D. M. (2020). Scientists' warning on invasive alien species. *Biological*

Reviews, *95*, 1511-1534.

Saikkonen, K., Taulavuori, K., Hyvönen, T., Gundel, P. E., Hamilton, C. E., Vänninen, I., … ,

Helander, M. (2012). Climate change-driven species' range shifts filtered by

photoperiodism. *Nature Climate Change*, *2*, 239-242.

Samis, K. E., Stinchcombe, J. R., Murren, C. J. (2019). Population climatic history predicts

phenotypic responses in novel environments for *Arabidopsis thaliana* in North America.

American Journal of Botany, *106*, 1068-1080.

Santangelo, J. S., Ness, R. W., Cohan, B., Fitzpatrick, C. R., Innes, S. G., Koch, S., … , Johnson,

M. T. J. (2022). Global urban environmental change drives adaptation in white clover.

Science, *375*, 1275-1281.

Savolainen, O., Lascoux, M., Merilä, J. (2013). Ecological genomics of local adaptation. *Nature*

Reviews Genomics, *14*, 807-820.

Siepielski, A. M., Morrissey, M. B., Buoro, M., Carlson, S. M., Caruso, C. M., Clegg, S. M., … ,

MacColl, A. D. C. (2017). Precipitation drives global variation in natural selection.

Science, *355*, 959-962.

Simón-Porcar, V. I., Silva, J. L., Vallejo-Marín, M. (2021). Rapid local adaptation in both sexual

and asexual invasive populations of monkeyflowers (*Mimulus* spp.). *Annals of Botany*,

127, 655-668.

St Clair, J. B., Howe, G. T. (2007). Genetic maladaptation of coastal Douglas-fir seedlings to

future climates. *Global Change Biology*, *13*, 1441-1454.

Stinchcombe, J. R., Weinig, C., Ungerer, M., Olsen, K. M., Mays, C., Halldorsdottir, S. S., … ,

Schmitt, J. (2004). A latitudinal cline in flowering time in *Arabidopsis thaliana*

modulated by the flowering time gene FRIGIDA. *Proceedings of the National Academy*

of Sciences of the United States of America, *101*, 4712-4717.

Taylor, C. M., Hastings, A. (2005). Allee effects in biological invasions. *Ecology Letters*, *8*, 895-

908.

Thompson, K. A., Renaudin, M., Johnson, M. T. J. (2016). Urbanization drives the evolution of

parallel clines in plant populations. *Proceedings of the Royal Society B: Biological*

Sciences, *283*, 20162180.

Till, I. (1987). Variability of expression of cyanogenesis in white clover (*Trifolium repens*

L.). *Heredity*, *59*, 265.

Toräng, P., Wunder, J., Obeso, J. T., Herzog, M., Coupland, G, Ågren, J. (2015). Large-scale

adaptative differentiation in the alpine perennial *Arabis alpina*. *New Phytologist*, *206*,

459-470.

Tsutsui, N. D., Suarez, A. V., Holway, D. A., Case, T. J. (2000). Reduced genetic variation and

the success of an invasive species. *Proceedings of the National Academy of Sciences*, *97*,

5948-5953.

Turesson, G. (1922). The genotypical response of the plant species to the habitat. *Hereditas*, *3*,

211-350.

van Boheemen, L. A., Lombaert, E., Nurkowski, K. A., Gauffre, B., Rieseber, L. H., Hodgins, K.

A. (2017). Multiple introductions, admixture and bridgehead invasion characterize the

introduction history of *Ambrosia artemisiifolia* in Europe and Australia. *Molecular*

Ecology, *26*, 5421-5434.

Walther, G.-R., Roques, A., Hulme, P. E., Sykes, M. T., Pyšek, P., Kühn, I., … , Settele, J.

(2009). Alien species in a warmer world: risks and opportunities. *Trends in Ecology &*

Evolution, *24*, 686-693.

Wang, T., Hamann, A., Spittlehouse, D., Carroll, C. (2016). Locally downscaled and spatially

customizable climate data for historical and future periods for North America. *PLoS One*,

11, e0156720.

Wilczek, A. M., Cooper, M. D., Korves, T. M., Schmitt, J. (2014). Lagging adaptation to

warming climate in *Arabidopsis thaliana*. *Proceedings of the National Academy of*

Sciences, *111*, 7906-7913.

Wolkovich, E. M., Davies, T. J., Schaefer, H., Cleland, E. E., Cook, B. I., Travers, S. E., … ,

Davis, C. C. (2013). Temperature-dependent shifts in phenology contribute to the success

of exotic species with climate change. *American Journal of Botany*, *100*, 1407-1421.

Wright, S. J., Cui Zhou, D., Kuhle, A., Olsen, K. M. (2018). Continent-wide climatic variation

drives local adaptation in North American white clover. *Journal of Heredity*, *109*, 78-89.

Wright, S. J., Goad, D. M., Gross, B. L., Muñoz, P. R., Olsen, K. M. (2021). Genetic trade-offs

underlie divergent life strategies for local adaptation in white clover. *Molecular Ecology*,

31, 3742-3760.

Züst, T., Heichinger, C., Grossniklaus, U., Harrington, R., Kliebenstein, D. J., Turnbull, L. A.

(2012). Natural enemies drive geographic variation in plant defenses. *Science*, *338*, 116-

119.

Züst, T., Agrawal, A. A. (2017). Trade-offs between plant growth and defense against insect

herbivory: an emerging mechanistic synthesis. *Annual Review of Plant Biology*, *68*, 513-

534.

Tables

Table 1 Results of mixed effects models assessing how continent of origin, common garden location, and climatic distance affected

the fitness of *T. repens*.

1155

Predictor	Plants Producing Flowers			Plants Producing Seeds			# of Flower Heads			Seed Set Mass		
	χ^2	df	P	χ^2	df	P	χ^2	df	P	χ^2	df	P
Continent of Origin (O)	0.330	1	0.566	0.218	1	0.641	2.314	1	0.128	1.214	1	0.271
Garden Location (L)	**31.703**	**3**	**< 0.001**	**8.104**	**3**	**0.044**	**252.144**	**3**	**< 0.001**	**40.068**	**3**	**< 0.001**
Climatic Distance (D)	**6.015**	**1**	**0.014**	**6.882**	**1**	**0.009**	**13.580**	**1**	**< 0.001**	**14.240**	**1**	**< 0.001**
O × L	0.998	3	0.802	0.952	3	0.813	1.442	3	0.696	4.085	3	0.252
O × D	0.421	1	0.516	1.113	1	0.291	0.277	1	0.599	0.978	1	0.323
L × D	4.439	3	0.218	4.185	3	0.242	**12.713**	**3**	**0.005**	**8.808**	**3**	**0.032**
O × L × D	6.860	3	0.077	7.631	3	0.054	7.764	3	0.051	4.882	3	0.181

1156

Predictor	Survival			Maximum Plant Area			Growth Rate			Herbivory		
	χ^2	df	P	χ^2	df	P	χ^2	df	P	χ^2	df	P
Continent of Origin (O)	2.220	1	0.136	0.939	1	0.332	0.094	1	0.759	0.089	1	0.766
Garden Location (L)	**221.162**	**3**	**< 0.001**	**286.435**	**3**	**< 0.001**	**70.328**	**3**	**< 0.001**	**144.426**	**3**	**< 0.001**
Climatic Distance (D)	**6.427**	**1**	**0.011**	0.584	1	0.445	0.089	1	0.766	1.074	1	0.300
O × L	6.155	3	0.104	6.464	3	0.091	1.212	3	0.750	6.635	3	0.084
O × D	0.142	1	0.707	1.747	1	0.186	0.049	1	0.824	0.010	1	0.919
L × D	4.798	3	0.187	5.610	3	0.132	2.887	3	0.409	2.234	3	0.525
O × L × D	7.288	3	0.063	2.693	3	0.441	3.399	3	0.334	3.127	3	0.372

1157

1158

1162 Note: Significant effects ($P < 0.05$) are listed in bold.

1163

1164

Table 2 Results of mixed effects models assessing how continent of origin, continent of garden, and climatic distance affected the fitness of *T. repens*.

Predictor	Plants Producing Flowers		Plants Producing Seeds		# of Flower Heads		Seed Set Mass	
	χ^2	P	χ^2	P	χ^2	P	χ^2	P
Continent of Origin (O)	2.033	0.154	1.754	0.185	0.514	0.474	0.028	0.867
Continent of Garden (G)	**9.184**	**0.002**	1.295	0.255	0.471	0.493	0.037	0.847
Climatic Distance (D)	**28.963**	**< 0.001**	**17.269**	**< 0.001**	**14.310**	**< 0.001**	**16.190**	**< 0.001**
O × G	0.677	0.411	0.942	0.332	0.957	0.328	0.481	0.488
O × D	2.033	0.154	0.036	0.850	1.214	0.271	2.991	0.084
G × D	3.194	0.074	2.490	0.115	1.412	0.235	0.258	0.612
O × G × D	0.501	0.479	0.026	0.871	**19.603**	**< 0.001**	**9.728**	**0.002**

Predictor	Survival		Maximum Plant Area		Growth Rate		Herbivory	
	χ^2	P	χ^2	P	χ^2	P	χ^2	P
Continent of Origin (O)	0.106	0.745	0.797	0.372	0.053	0.817	0.196	0.658
Continent of Garden (G)	**41.220**	**< 0.001**	**170.553**	**< 0.001**	1.468	0.226	**21.180**	**< 0.001**
Climatic Distance (D)	**10.236**	**0.001**	**58.341**	**< 0.001**	**18.137**	**< 0.001**	**15.810**	**< 0.001**
O × G	0.000	0.993	2.375	0.123	0.465	0.495	0.669	0.413
O × D	1.534	0.216	**34.173**	**< 0.001**	**5.145**	**0.023**	**23.686**	**< 0.001**
G × D	**13.870**	**< 0.001**	1.784	0.182	2.263	0.133	0.240	0.624
O × G × D	**6.515**	**0.011**	0.678	0.410	**7.812**	**0.005**	2.178	0.140

Note: Significant effects ($P < 0.05$) are listed in bold. All predictors are assessed using one degree of freedom.

1184 **Table 3** Results of mixed effects models assessing how change in mean annual temperature (ΔMAT) affected the fitness of North

1185 American *T. repens* populations in the Mississauga common garden (linear and quadratic fits).

1186

Predictor	Survival		Plants Producing Flowers		Plants Producing Seeds		# of Flower Heads		Seed Set Mass		Maximum Plant Area		Growth Rate	
	χ^2	P	χ^2	P	χ^2	P	χ^2	P	χ^2	P	χ^2	P	χ^2	P
ΔMAT	3.758	0.053	**7.425**	**0.006**	**5.397**	**0.020**	0.255	0.614	0.644	0.422	3.799	0.051	0.002	0.962
ΔMAT^2	**4.313**	**0.038**	**8.077**	**0.004**	**7.609**	**0.006**	2.908	0.088	**8.542**	**0.003**	**6.741**	**0.009**	1.808	0.179

1187

1188 Note: Significant effects ($P < 0.05$) are listed in bold. All predictors are assessed using one degree of freedom.

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

 Figure Captions

1200

1201 **Figure 1** Maps showing populations of *T. repens* sampled from the (**A**) introduced range (North America, orange circles) and (**B**)

1202 native range (Europe, pink circles), as well as the four common garden locations (green diamonds). Coordinates of common garden

1203 locations are 42°32'40" N, 79°39'38" W (Mississauga, Ontario, Canada), 30°18'23" N, 92°00'33" W (Lafayette, Louisiana, USA),

1204 59°49'08" N, 17°38'49" W (Uppsala, Sweden), and 43°38'16" N, 3°51'43" W (Montpellier, France).

1205

1206 **Figure 2** The effect of climatic distance and common garden location on fitness of *T. repens*. (**A**) Mean number of flower heads and

1207 (**B**) mean seed set mass of *T. repens* populations planted in two common gardens in North America (Lafayette in navy and

1208 Mississauga in orange) and two common gardens in Europe (Montpellier in green and Uppsala in magenta). Shaded areas represent

1209 95% confidence intervals around each line, while asterisks represent *P* values of relevant fixed effects of Climatic Distance (D) and its

1210 interaction with Continent of Origin (O) and Continent of Garden (G) (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$).

1211

1212

1213

1214

Figure 3 The effect of climatic distance on the mean number of flower heads produced by *T. repens* populations originating from its

introduced (North America, orange) and native (Europe, pink) ranges and transplanted into common gardens in the (**A**) introduced and

(**B**) native ranges. The effect of climatic distance on the mean seed set mass produced by *T. repens* populations originating from its

introduced and native ranges and transplanted into common gardens in the (**C**) introduced and (**D**) native ranges. Shaded areas

represent 95% confidence intervals around each line, while asterisks represent *P* values of < 0.05 for relevant fixed effects of Climatic

Distance (D) and its interaction with Continent of Origin (O) and Continent of Garden (G).

1221

1222

Figure 4 The effects of the change in mean annual temperature (ΔMAT) on fitness of North American populations of *T. repens* grown

in the Mississauga common garden. (**A**) Proportion of plants survived, (**B**) proportion of plants that produced flowers, (**C**) proportion

of plants that produced seeds, (**D**) mean seed set mass, and (**E**) mean maximum plant area per population all exhibited evidence of an

adaptation lag in response to climate warming, indicated by a positive linear trend or a shift in the peak of the quadratic curve towards

populations with warmer climates of origin (denoted as x-axis values > 0). Shaded areas represent 95% confidence intervals around

each line.

1229

1230

1231   **Figures**

1232   **Figure 1**

1233

1234   **Figure 2**

1235   **Figure 3**

1236 **Figure 4**