1 **Adaptation to climate in the native and introduced ranges of a cosmopolitan plant**

2

3

4 **Lucas J. Albano[1,2], Cristina C. Bastias[3,4], Aurélien Estarague[4], Brandon T. Hendrickson[5],**

5 **Simon G. Innes[5], Nevada King[5], Courtney M. Patterson[5], Amelia Tudoran[6], François**

6 **Vasseur[4], Adriana Puentes[6], Cyrille Violle[4], Nicholas J. Kooyers[5], Marc T. J. Johnson[2]**

7

8

9 [1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

10 [2]Department of Biology, University of Toronto Mississauga, Mississauga, ON, Canada

11 [3]Ecology Area, Department of Botany, Ecology and Plant Physiology, Campus de Rabanales,

12 Córdoba University, Córdoba, Spain

13 [4]CEFE, Université Montpellier, CNRS, EPHE, IRD, Montpellier, France

14 [5]Department of Biology, University of Louisiana at Lafayette, Lafayette, LA, USA

15 [6]Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

16

17

18 **Correspondence Author:**

19 Lucas J. Albano

20 Current address: Department of Plant & Microbial Biology, North Carolina State University,

21 Raleigh, NC, United States, 27695

22 ljalbano@ncsu.edu

23 (919) 931-5026

Keywords

Local adaptation, invasive species, environmental variation, white clover, adaptation lag,

cyanogenesis, provenance study

Abstract (301 words)

Climate change and the global spread of invasive species are currently two of the most

significant threats to biodiversity worldwide. Yet the role of adaptation in organismal responses

to changing climates or in facilitating species invasions is still poorly understood. We conducted

a large-scale trans-continental common garden experiment to understand adaptation to

spatiotemporal variation in climate in the native and introduced ranges of a cosmopolitan plant

species. Individuals from 96 populations of *Trifolium repens* (white clover) from both its native

(Europe) and introduced (North America) ranges were planted into four experimental common

gardens located in northern (Uppsala, Sweden) and southern (Montpellier, France) Europe, and

northern (Mississauga, Canada) and southern (Louisiana, USA) North America. We recorded

plant sexual and clonal fitness in each common garden and assessed whether plants are best

adapted to local climates, whether the strength of local adaptation differed between the native

and introduced ranges, and whether populations show evidence of rapid adaptation to recent

climate change. Results show that white clover is locally adapted, particularly in lower latitude

locations, but also that the most prominent bioclimatic drivers of local adaptation differed by

latitude. We also found that strong local adaptation was only evident when populations were

transplanted into common gardens located in the same range (native or introduced) from which

they originated, indicating rapid local adaptation across a large latitudinal gradient has occurred

in *T. repens* populations in less than 400 years since its introduction to North America. However,

we do find some evidence of an adaptation lag in the northern common garden in the introduced

range, with plants from historically slightly warmer climates exhibiting the greatest fitness. This

indicates that despite evidence of local adaptation to historic climatic conditions and rapid local

adaptation post-introduction, white clover populations may be evolving more slowly than climate

change is occurring.

Introduction

Natural selection results in populations evolving traits that are advantageous to the

environmental conditions of a particular location, a phenomenon known as local adaptation

(Turesson, 1922; Kawecki and Ebert, 2004; Hereford, 2009). A fundamental aspect of local

adaptation is that spatial variation in biotic and abiotic factors leads to changes in the relative

fitness of genotypes between environments (Gillespie and Turelli, 1989; Kawecki and Ebert,

2004; Lee and Mitchell-Olds, 2011). This can result in fitness trade-offs, whereby genotypes

exhibit higher relative fitness in their local environment and lower fitness in other environments

(Gillespie and Turelli, 1989; Hereford, 2009; Savolainen et al., 2013). Understanding how

populations simultaneously adapt to their local biotic and abiotic environment over space and

time remains an important problem in evolutionary ecology. When changes to the environment

are caused by humans, studies of local adaptation can help to determine how increasingly

pervasive anthropogenic disturbance alters the ecology and evolution of species (Hoffmann and

Sgrò, 2011; Johnson and Munshi-South, 2017).

While spatial variation in the environment can commonly result in local adaptation

(Hereford, 2009; Leimu and Fischer, 2008), it remains poorly understood how locally adapted

populations are affected when environments rapidly change (Kawecki and Ebert, 2004;

Anderson and Song, 2020). Temporal shifts in the local environment can cause selection

pressures to fluctuate, which could result in traits that were previously adapted to that

environment becoming maladaptive (Wilczek et al., 2014; Aguirre-Liguori et al., 2019;

Exposito-Alonso et al., 2019; Anderson and Wadgymar, 2020; Bontrager et al., 2020). If the rate

of adaptation is exceeded by the rate of environmental change, this disruption of local adaptation

is likely to result in reduced fitness (i.e., maladaptation) of local populations (St Clair and Howe,

2007; Atkins and Travis, 2010; Wilczek et al., 2014; Aguirre-Liguori et al., 2019). Recent global

warming due to anthropogenic activity may result in populations from lower latitudes or altitudes

becoming better suited to local environmental conditions than previously locally adapted

populations, which is known as an adaptation lag (Wilczek et al., 2014; Kooyers et al., 2019;

Anderson and Wadgymar, 2020). Currently, the frequency and magnitude at which adaptation

lags are occurring is poorly understood and requires largescale experiments to assess local

(mal)adaptation as a result of rapid environmental change (Anderson and Song, 2020).

While rapid adaptation may facilitate population responses to changing climates, it could

also facilitate establishment and range expansion during species introductions. This introduces

the potential for invasive species to become even more persistent with future environmental

change through rapid local adaptation to novel biotic and abiotic environments (Maron et al.,

2004; Colautti and Barrett, 2013; Oduor et al., 2016). For example, biotic selection regimes

experienced by species introduced to new environments could differ from those experienced in

their native range. Classic hypotheses in invasion biology, while not universally supported, posit

that a release of an introduced species from the natural enemies in its native range (Keane and

Crawley, 2002), or the evolution of increased competitive ability relative to native species

(Blossey and Nötzold, 1995), could each result in the proliferation and success of an invader.

Additionally, populations of introduced species can experience particularly strong abiotic

selection compared to populations of locally adapted native species, facilitating high rates of

adaptation (Maron et al., 2004; Colautti and Barrett, 2013; Colautti and Lau, 2015). Invasive

species are also often better at climate tracking and more resistant to climate warming than

native species, which could contribute to their success as invaders as environments change

(Wolkovich et al., 2013). The interplay between local adaptation, species introductions, and

anthropogenic environmental change is a currently understudied, yet critically important, aspect

of understanding the ongoing human-induced biodiversity crisis (IPBES, 2019). Therefore, direct

tests of local adaptation in both the native and introduced ranges of a widespread invasive

species are needed, alongside elucidating the environmental conditions that may result in rapid

adaptation and invasion success (Colautti and Lau, 2015; Anderson and Song, 2020).

Plant-herbivore interactions can serve as a focal point for simultaneous investigation of

local adaptation, species responses to environmental change, and biological invasions. Local

herbivore communities impose selection through preferential feeding on plants with lower

resistance, influencing plant defense evolution through an evolutionary arms race (Ehrlich and

Raven, 1964; Agrawal, 2007; Züst et al., 2012). Spatial variation in herbivore communities is

therefore likely to lead to spatial variation in plant defense traits (Anstett et al., 2018; Baskett and

Schemske, 2018; Moreira et al., 2018). For example, the latitudinal herbivory-defense hypothesis

predicts greater diversity and abundance of plant defenses at lower latitudes due to the presence

of greater herbivore diversity and abundance (Coley and Barone, 1996; Johnson and Rasmann,

2011; Anstett et al., 2016; Moreira et al., 2018). Plant defense evolution is also likely to be

driven by the abiotic environment through growth-defense trade-offs (Coley et al., 1985; Herms

and Mattson, 1992; Strauss et al., 2002). The metabolic costs of defenses can be substantial,

while abiotic environments and the stressors they impose can influence the availability of

resources to plants, exacerbating defense costs (Coley et al., 1985; Fine et al., 2006; Hahn et al.,

2019). Furthermore, it is increasingly recognized that anthropogenic activity can influence the

evolution of plant defenses (Bidart-Bouzat and Imeh-Nathaniel, 2008; Massad and Dyer, 2010;

Miles et al., 2019). Climate warming could result in increased herbivore feeding and/or increased

plant defenses (Dostálek et al., 2020; Lemoine et al., 2013) or facilitate invasion success by

decoupling locally-adapted plant-herbivore interactions (Adams et al., 2009; Costan et al., 2022;

but see Agrawal and Kotanen, 2003). It is therefore imperative that we understand how

populations are adapting to their local biotic and abiotic environments and how anthropogenic

disturbance can affect adaptation to both ecological conditions.

An often-documented manifestation of local adaptation is a cline – a gradual change in

phenotype or genotype across an environmental gradient (Endler, 1977; Stinchcombe et al.,

2004; Savolainen et al., 2013). Such a cline has been clearly demonstrated for a prominent

chemical defense mechanism (cyanogenesis) in the cosmopolitan plant white clover (*Trifolium*

repens L.) across many different environmental gradients (Daday, 1954a; Daday, 1954b, Daday,

1958; Till, 1987; Kooyers and Olsen, 2013; Kooyers et al., 2014; Thompson et al., 2016). These

clines are consistent with local adaptation, whereby certain *T. repens* genotypes benefit from a

"home field advantage" based on their location along an environmental gradient (Daday, 1965).

One hypothesis to explain cyanogenesis clines is a gradient in herbivory, with greater diversity

and abundance of herbivores in warmer climates resulting in stronger selection favouring

cyanogenic plants (Angseesing, 1974; Dirzo and Harper, 1982; Kakes, 1989). Alternatively,

empirical studies across various environmental gradients have identified temperature, soil

moisture, and soil nutrients as abiotic factors that may be involved in cyanogenesis cline

formation (Daday, 1954a; Daday, 1965; Gleadow and Woodrow, 2002; Kooyers et al., 2014;

Kooyers et al., 2018). Abiotic factors have been shown to be strong drivers of local adaptation in

T. repens, although they exhibit weaker evidence as drivers of cyanogenesis cline formation

(Wright et al. 2018). Furthermore, despite 70 years of research on the drivers of cyanogenesis

clines in *T. repens*, there remains little consensus on how variation in environmental conditions

influence selection on cyanogenesis in *T. repens* (Daday, 1954a; Daday, 1965; Hughes, 1991;

Kooyers and Olsen, 2013; Kooyers et al., 2014; Wright et al., 2018; Albano and Johnson, 2023).

Further investigation of the role that plant defense traits play in local adaptation remains

important to understanding how plant populations adapt to rapidly changing environments.

To understand the roles of biotic and abiotic environments in driving local adaptation, we

conducted a trans-continental common garden field experiment across latitudinal gradients in

both the native (Europe) and introduced (North America) ranges of *T. repens*. This experiment

addressed five unresolved questions: (Q1) Are *T. repens* populations locally adapted to their

climate of origin and does the strength of local adaptation differ across space? (Q2) Does *T.*

repens exhibit stronger local adaptation in its native range than in its introduced range? (Q3)

What bioclimatic variables have the strongest impact on fitness at different latitudes in the native

and introduced ranges of *T. repens*? (Q4) Are there adaptation lags to climate change in *T.*

repens, such that populations originating from warmer and/or drier climates exhibit greater

fitness than local populations? (Q5) What are the roles of cyanogenesis and herbivory in driving

local adaptation? Our results further our understanding of local adaptation to climate and the

biotic environment, the potential for adaptation lags, and the success of invaders in the face of

environmental change.

Methods

Study system

Trifolium repens, or white clover, is an allopolyploid perennial legume that is native to Eurasia

and has been introduced to inhabited continents throughout the world, including North America

(Daday, 1958; Kjærgaard, 2003; Griffiths et al., 2019). *Trifolium repens* has been a model for the

study of plant defense based on its polymorphism for the production of the potent chemical

defense hydrogen cyanide (HCN; Hughes, 1991). Cyanogenesis occurs through the

hydrolyzation of cyanogenic glucosides by the enzyme linamarase. In *T. repens*, the presence or

absence of cyanogenic glucosides, linamarin and lotaustralin, is controlled by three linked genes

in their metabolic pathway (*CYP79D15*, *CYP736A187*, and *UGT85K17*; referred to collectively

as *Ac/ac*), while the presence or absence of the enzyme linamarase is controlled by the gene *Li/li*

(Olsen et al., 2007; Olsen and Small, 2018). Thus, four phenotypes exist for HCN and the two

metabolic components required for its production: one in which HCN is constitutively produced

(AcLi) and three that are acyanogenic and therefore lack the defense entirely (acLi, Acli, acli;

Corkill, 1940; Atwood and Sullivan, 1943).

The ability to produce HCN as an antiherbivore defense may have facilitated the range

expansion of *T. repens*, allowing it to thrive in human-disturbed habitats such as lawns, fields,

and roadsides in many regions throughout the world (Burdon, 1983; Kjærgaard, 2003; Olsen et

al., 2021). However, the presence of HCN is not ubiquitous across environments. In fact, *T.*

repens exhibits clines in the presence of HCN across numerous types of environmental gradients

(Daday, 1954a; Daday, 1954b; Daday 1958; Brighton and Horne, 1977; Thompson et al., 2016;

Innes et al., 2022; Santangelo et al., 2022). Latitudinal cyanogenesis clines in *T. repens* in the

native European range exhibit fixation of dominant *Ac* and *Li* alleles in warmer Mediterranean

climates and fixation of recessive *ac* and *li* alleles in cooler Nordic climates (Daday, 1954a).

Similar clines have also been demonstrated in many introduced ranges of white clover, including

North America (Daday, 1958; Kooyers and Olsen, 2012; Innes et al., 2022). However, these

clines are less steep, possibly indicating weaker local adaptation in introduced ranges due to the

smaller amount of time that local adaptation has had to occur (less than 400 years in North

America; Daday, 1958; Kjærgaard, 2003; Kooyers and Olsen, 2012; Innes et al., 2022). Despite

over seven decades of research on this system, an explicit test of the strength of local adaptation

in the native and introduced ranges of *T. repens* has yet to be conducted.

Sampled populations and common garden sites

We addressed our research questions using four common gardens distributed across northern

(Uppsala, Sweden) and southern (Montpellier, France) regions of the native European range of *T.*

repens, and northern (Mississauga, ON, Canada) and southern (Lafayette, LA, USA) regions of

the North American introduced range (**Figure 1**). Seeds were collected from 49 natural *T. repens*

populations scattered across the native European range spanning ~27° of latitude, and 47 natural

T. repens populations along a 2500 km north-south transect spanning ~21° of latitude in the

introduced North American range (**Figure 1**). To minimize maternal and epigenetic effects, we

hand-pollinated plants germinated from field-collected seeds in a growth chamber environment

(25°C day, 15°C night, 14 h:10 h light:dark cycle; Conviron MTPS, Winnipeg, MB, Canada) to

create a refresher generation of outbred F1 seeds for each population. F1 generation seeds were

germinated and established for four weeks in peat pots (8 cm diameter, 591 mL volume) in

uniform growth chamber or greenhouse conditions near each common garden site. Five replicate

individuals from each population were transplanted in peat pots directly into the natural soil of a

typical cultivated lawn at each common garden site. Inconsistent germination success resulted in

some populations not being included in some common garden locations, and in very limited

cases, populations had four or six individuals planted (the resulting total number of plants per

garden was 465, 465, 458, and 470 in Mississauga, Lafayette, Uppsala, and Montpellier,

respectively). Plants were arranged in rows and columns with 1 m spacing, and black

landscaping fabric was used to cover the entirety of the lawn to prevent the rooting of stolons

into the natural soil between plants. 12 cm diameter holes were cut in the fabric for each

individual plant, leaving a 2 cm ring of natural vegetation around each plant to allow for natural

competition. This vegetation was trimmed monthly throughout the experiment. Planting dates

matched growing seasons in each location. The Lafayette and Mississauga common gardens

were established in April and May 2020, respectively, while the Montpellier and Uppsala

common gardens were postponed due to COVID-19 restrictions in Europe, leading to their

establishment in March and May 2021, respectively. Plants were given supplemental water for

the first two weeks of the experiment and intermittently after the first two weeks when necessary

to avoid excessive mortality. Plants that died within the first two weeks of the experiment were

deemed to have died due to transplantation stress and were replaced using extra germinated

individuals from the same population. Plants that died after two weeks were deemed to have died

due to the local environmental conditions at each common garden site and were not replaced.

Each common garden was maintained for two full growing seasons.

Tissue collection, herbivory estimates, and measures of fitness

After 1-2 months of growth, one to three fully-expanded leaves were harvested from each

surviving individual and frozen at -20°C. These leaves were later freeze-dried and used for DNA

extraction, followed by low coverage whole genome sequencing to predict the presence of *Ac/ac*

and *Li/li* alleles (see **Supplemental Methods**). In total, 586 individuals (Mississauga: 302,

Lafayette: 99, Uppsala: 78, Montpellier: 107) were successfully sequenced and characterized as

possessing either dominant or recessive alleles at *Ac/ac* and *Li/li*. The remaining 1272

individuals were not genotyped because they either died before tissue could be harvested or

genomic library preparations were unsuccessful. Visual herbivory estimates (in % leaf area

consumed) were conducted monthly following the methods of Johnson et al. (2016). Herbivory

was analyzed for the month exhibiting peak herbivory within each garden. Multiple components

of plant fitness were measured throughout the growing seasons at each location. Survival and

whether or not plants flowered were assessed weekly during the first growing season. Mature

fruits were counted and collected weekly, and seeds were separated using a thresher (Precision

Machine Co., Lincoln, NE, USA) to obtain total seed set mass for each plant across both growing

seasons of the experiment. Photos were taken of each plant monthly during the first growing

season at each common garden location and assessed for plant area using Easy Leaf Area

(Version 2.0; Easlon and Bloom, 2014). The maximum plant area was then determined and the

growth rate across the first growing season was calculated as $(\ln[A_{max}] - \ln[A_1])/(t_{max} - t_1)$,

where A_{max} represents the maximum plant area, A_1 represents the initial plant area (i.e., after one

month of growth in the common garden), and $(t_{max} - t_1)$ represents the number of days between

photos taken for maximum plant area and initial plant area.

Calculating climatic distance

Six bioclimatic variables were extracted from WorldClim (Version 2.1, February 2023) at a

resolution of 30 seconds for each original *T. repens* population and common garden site

(O'Donnell and Ignizio, 2012). The variables included were mean annual temperature ("BIO1";

MAT), maximum temperature of the warmest month ("BIO5"), minimum temperature of the

coldest month ("BIO6"), annual precipitation ("BIO12"), precipitation of the wettest month

("BIO13"), and precipitation of the driest month ("BIO14"). The extracted values of each

bioclimatic variable for each common garden location can be found in **Table S1**. Values for all

six variables were z-score standardized to allow for analysis on an equivalent scale. A principal

component analysis was performed using the *princomp* function in R, with dimensions 1, 2, and

3 explaining 65.8%, 22.1%, and 9.1% of the variation, respectively (**Figure S1**). Coordinates for

each location of origin and each common garden location from all three dimensions were then

extracted using the *get_pca_ind* function in the *factoextra* package (version 1.0.7; Kassambara

and Mundt, 2020) and then plotted on a 3D graph (**Figure S2**). The weighted Euclidean distance

between points in 3D space was calculated using the formula:

263
$$Distance = \sqrt{\sum (w_i [O_{ij} - G_i]^2)}$$

Where w_i represents the proportion of variance explained by the ith dimension, O_{ij} represents the

value of the jth population of origin in the ith dimension, and G_i represents the value of a given

common garden in the ith dimension. We used this weighted Euclidean distance as a measure of

the climatic distance between the location of each population of origin and each common garden

location based on our six bioclimatic variables of interest (Samis et al., 2019). This measure of

climatic distance was then used as a predictor variable of plant fitness in our statistical analyses.

Statistical analysis

All statistical procedures and analyses were conducted in R Version 4.1.2 (2021) and

RStudio Version 2022.12.0 (R Core Development Team, 2022). Seven response variables were

assessed as components of fitness: i) whether or not plants survived the first growing season, ii)

whether or not plants flowered in the full experiment, iii) whether or not plants produced seeds in

the full experiment, iv) number of flower heads produced in the full experiment, v) seed set mass

produced in the full experiment, vi) maximum plant area in the first growing season, and vii)

growth rate in the first growing season. Herbivory during the first growing season was also

measured as an additional non-fitness response variable. The number of flower heads and seed

set mass were both zero-inflated, which led to the use of a two-step modelling approach as

described previously (Albano and Johnson, 2023). Briefly, both variables were first analyzed in

binomial models to assess whether or not plants produced flowers or seeds. Non-zero flower

head numbers and seed set masses were then subsequently analyzed as separate measures of

fitness. All general linear mixed effects models were conducted using the *glmmTMB* function in

the *glmmTMB* package (version 1.1.5; Brooks et al., 2017). All test statistics and *P* values were

calculated using the *Anova* function in the *car* package (version 3.0-12; Fox and Weisberg, 2019)

with Type II sums-of-squares. For models using quantitative measures of fitness, all

transformations necessary to meet assumptions of normality and homogeneity of variance can be

found in **Table S2**.

289   *Variation in the strength of local adaptation between common garden locations*

290   To determine whether or not *T. repens* populations exhibit local adaptation and to assess the

291   strength of local adaptation at each garden location (Q1), all response variables were analyzed

292   using the following structure:

293      $Response \sim Continent\ of\ Origin\ (O) + Garden\ Location\ (L) + Climatic\ Distance\ (C)$

294                    $+ O{:}L + O{:}D + L{:}D + O{:}L{:}D + (1|Population)$

Continent of origin (North America or Europe), garden location (Mississauga, Lafayette,

Uppsala, or Montpellier), and climatic distance (the combined measure of six bioclimatic

variables of interest described above) were included as fixed effects, along with all two- and

three-way interactions. Population nested within continent of origin was included as a random

effect to account for non-independence of replicates within each population. An interaction

between continent of origin and climatic distance would indicate that the strength of local

adaptation varies based on whether or not populations originated in the native or introduced

range. An interaction between garden location and climatic distance would indicate that the

strength of local adaptation varies based on the particular common environment in which it is

being tested. The strength of local adaptation refers to the magnitude of the fitness advantage for

local populations compared to populations transplanted an increasingly larger climatic distance

from their origin. We therefore interpret the slope of the relationship between climatic distance

and fitness as being representative of the strength of local adaptation within a given common

garden.

Variation in the strength of local adaptation between ranges

To assess the strength of local adaptation between the native and introduced ranges of *T. repens*

(Q2), the analyses of the previous sub-section were repeated, with the continent of the common

garden (North America or Europe) replacing the location of the common garden in the models:

$Response \sim Continent\ of\ Origin + Continent\ of\ Garden\ (G) + Climatic\ Distance$

$+\ O{:}G + O{:}D + D{:}G + O{:}D{:}G + (1|Population)$

In this model, a 3-way interaction between continent of origin, continent of garden, and climatic

distance may indicate that the strength of local adaptation varies based not only on the range

from which populations originated but also the range in which they are being tested. In other

words, this model aims to test whether locally adapted European populations maintain signatures

of local adaptation in North America and vice versa.

Bioclimatic variables as individual drivers of local adaptation

We also measured associations between the seven response variables and each of the six

bioclimatic variables of interest individually by determining the difference in z-score

standardized values for each variable between the location of each population of origin and each

common garden location using the following model structure:

$Response \sim Bioclimatic\ Variable + Bioclimatic\ Variable^2 + (1|Population)$

For this set of analyses, all common gardens were analyzed separately, and all quantitative (non-

binomial) response variables were z-score standardized to allow for simultaneous direct

comparisons of the relative strength of linear and quadratic predictor variables. Within each

garden, we were therefore able to directly compare P values across all bioclimatic variables for

both linear and quadratic fits (**Tables S3-S6**). For each response variable, we selected the most

influential bioclimatic variable (i.e., the lowest P, but only when at least one P value was

significant at a level of $P < 0.1$, highlighted in red in **Tables S3-S6**). Across all response

variables in each garden, we then calculated the percentage of these most influential bioclimatic

variables that were either temperature-based (BIO1, BIO5, or BIO6) or precipitation-based

(BIO12, BIO13, and BIO14) and compared these percentages across the full experiment, across

latitudes, and across ranges (Q3).

Adaptation lags in response to climate warming

We sought to investigate the presence of adaptation lags using a space-for-time substitution (Q4).

We performed an additional analysis using a similar model structure as before, but only using

changes in raw values for MAT (i.e., not z-score standardized, hereafter referred to as ΔMAT) as

the bioclimatic variable of interest.

$Response \sim \Delta MAT + \Delta MAT^2 + (1|Population)$

If the peak of the quadratic relationship in each model is approximately equal to zero, it would be

considered evidence of local adaptation to MAT, because maximum fitness occurs in populations

with the same mean annual temperature as the common garden site. If the peak of the quadratic

relationship is instead shifted towards populations of warmer origins than are occurring at the

common garden site (i.e., these warmer-adapted populations are outperforming the local

populations), this would be considered evidence of an adaptation lag. We tested whether the shift

of peaks away from a ΔMAT of zero and towards populations of warmer origins were significant

by also fitting a similar model to above, but excluding the linear predictor:

$$Response \sim \Delta MAT^2 + (1|Population)$$

This additional model constrains the peak of the distribution to be centred around a ΔMAT of

zero (approximating the outcome of local adaptation as opposed to an adaptation lag). We then

compared the models with and without the linear predictor using the *anova* and *AIC* functions in

R, with *P* values < 0.05 (along with ΔAIC values < 0) indicating a significant shift of the peak.

We focused our interpretation of this analysis based solely on the North American

populations in the Mississauga common garden because the sampling design of North American

populations along a transect and the location of the Mississauga garden within that sampled

transect provide the most appropriate subset of the full experiment data to detect spatial lags. We

include the results for European populations within the Uppsala common garden in **Table S7** for

completeness, but the Uppsala common garden did not exhibit any trend in fitness based on

climatic distance, precluding the possibility of an adaptation lag being detected. The Lafayette

and Montpellier common gardens were excluded from this analysis because they are near the

southern extremes of our sampled populations, precluding the ability to detect adaptation lags.

The roles of cyanogenesis and herbivory in local adaptation

The presence of cyanogenesis clines across latitude on both continents was determined

separately for *Ac/ac* and *Li/li* by assessing the proportion of plants in each population containing

at least one dominant *Ac* or *Li* allele using the following model structure:

$$Proportion\ \text{Ac}\ or\ \text{Li}\ Plants \sim Latitude + Continent\ of\ Origin$$

$$+ Latitude: Continent\ of\ Origin + (1|Population)$$

An interaction between latitude and continent of origin would indicate a difference in the slope

of clines in the presence of *Ac* and/or *Li* based on the range in which those clines are occurring

(native European or introduced North American).

The effect of cyanotype (AcLi, Acli, acLi, or acli) on local adaptation (Q5) was assessed

for all eight response variables using the following model structure:

$$Response \sim Cyanotype\ (C) + Garden\ Location\ (L) + Climatic\ Distance\ (D)$$

$$+ C:L + C:D + C:L:D + (1|Population)$$

An interaction between cyanotype and climatic distance would indicate that cyanogenesis or its

metabolic components contribute to the strength of local adaptation in *T. repens*. A 3-way

interaction between cyanotype, garden location, and climatic distance would indicate that

differential selection for or against a given cyanotype in a particular common environment

impacts the strength of local adaptation in that environment. This might be expected due to

consistently observed latitudinal cyanogenesis clines across multiple types of environmental

gradients in *T. repens*.

Results

Local adaptation in white clover is dependent on latitude

We found strong evidence of local adaptation, and the strength of local adaptation in *T. repens*

varied among gardens for multiple measures of fitness. Climatic distance between each

population of origin and the common garden locations strongly predicted all measures of sexual

fitness and survival (**Table 1**). In general, populations tended to have higher fitness with lower

climatic distance, indicating that populations of more similar climatic origins to the climate of

each common garden were better adapted. The only response variables not showing evidence of

local adaptation were the measures of vegetative growth used as proxies for clonal fitness (i.e.,

maximum plant area and growth rate). The interaction of common garden location and climatic

distance affected the number of flower heads and seed set mass, but not other measures of fitness

or herbivory (**Table 1**; **Figure S3**). In general, the strength of local adaptation, measured as the

slope of the relationship between sexual fitness and climatic distance, was greater in the more

southern common garden location on each continent (**Figure 2**). For the number of flower heads,

the slope for Lafayette was 3.4× steeper than the slope for Mississauga and the slope for

Montpellier was 349× steeper than the slope for Uppsala. Similarly, for seed set mass, the slope

for Lafayette was 2.3× steeper than the slope for Mississauga and the slope for Montpellier was

384× steeper than the slope for Uppsala. The main effect of garden location affected all measures

of fitness and herbivory, which reflects the large differences in environment between the

gardens. By contrast, the main effect of continent of origin and its interactions with garden

location and/or climatic distance did not affect any measure of fitness or herbivory (**Table 1**).

Local adaptation is more evident within ranges than across ranges

We found that the strength of local adaptation in *T. repens* populations was dependent not only

on latitude, but also a combination of the range from which plants originated and the range to

which they were transplanted. Continent of origin, continent of the common garden, and climatic

distance all interacted to affect all quantitative measures of sexual fitness, survival, and growth

rate of *T. repens* (**Table 2**). Specifically, populations only exhibited strong signatures of local

adaptation when they were transplanted to common gardens on the same continent from which

they originated (**Figure 3**). For number of flower heads, in common gardens located in the native

range (Europe), the slope for European *T. repens* populations was 12× steeper than the slope for

North American populations. In common gardens located in the introduced range (North

America), the slope for North American *T. repens* populations was 20× steeper than the slope for

European populations. Similarly, for seed set mass, in common gardens located in Europe, the

slope for European *T. repens* populations was 24× steeper than the slope for North American

populations. In common gardens located in North America, the slope for North American *T.*

repens populations was 7.8× steeper than the slope for European populations. The main effect of

climatic distance affected whether or not plants flowered and produced seeds, but climatic

distance did not interact with the continent of origin or the continent of the common garden to

affect these components of fitness (**Table 2**; **Figure S4**). However, climatic distance and

continent of origin did interact to affect maximum plant area and herbivory. Together across all

gardens, plants from European origins that were transplanted shorter climatic distances grew

larger, despite experiencing greater levels of herbivory, while size and herbivory for plants from

North American origins were not significantly affected by climatic distance overall (**Table 2**;

Figure S4).

The most prominent driving factors of local adaptation are dependent on latitude

Climatic variables related to both temperature and precipitation predicted variation in fitness in

T. repens, but the most influential predictors varied between garden locations, with patterns

emerging based on latitude (**Tables S3-S6**, red highlights). For the full experiment, 58% of the

most influential predictors were temperature-based (mean annual temperature, maximum

temperature of the warmest month, or minimum temperature of the coldest month), while 42%

were precipitation-based (annual precipitation, precipitation of the wettest month, or

precipitation of the driest month). In the more southern gardens on each continent (Lafayette and

Montpellier), 67% of the most influential predictors were precipitation-based, while only 33%

were temperature-based. Conversely, in the more northern gardens on each continent

(Mississauga and Uppsala), 80% of the most influential predictors were temperature-based,

while only 20% of were precipitation-based. We did not find evidence that the native and

introduced ranges of *T. repens* were differentially affected by temperature or precipitation

variables. In the native range (Uppsala and Montpellier, combined), 57% of the most influential

predictors were temperature-based and 43% were precipitation-based. In the introduced range

(Mississauga and Lafayette, combined), 58% of the most influential predictors were temperature-

based, while 42% were precipitation-based.

Evidence of adaptation lags in northern North America

North American populations of *T. repens* demonstrated evidence of an adaptation lag in the

Mississauga common garden (**Table 3**). Survival, seed set mass, and maximum plant area all

exhibited a significant quadratic relationship with ΔMAT between locations of origin and the

Mississauga common garden (**Figure 4**). A quadratic relationship centred on a ΔMAT of zero

would represent local adaptation, while a relationship centred on a ΔMAT > 0 would represent

an adaptation lag. For the proportion of plants survived per population, the peak of the quadratic

relationship experienced a significant shift to a climate of origin with a MAT 3.4°C greater than

Mississauga ($\chi^2_{1,4} = 4.159$, $P = 0.041$, ΔAIC = −2.159), with the predicted maximum survival

being 5.3% greater than at a ΔMAT of zero. For mean seed set mass per population, the peak of

the quadratic relationship occurred at a MAT of 2.0°C greater than Mississauga, with the

predicted maximum seed set mass being 5.2% greater than at a ΔMAT of zero. However, this

shift was not statistically significant ($\chi^2_{1,5} = 0.640$, $P = 0.423$, ΔAIC = 1.360). For mean

maximum plant area per population, the peak of the quadratic relationship experienced a shift to

a climate of origin with a MAT of 2.1°C ($\chi^2_{1,5}$ = 3.721, P = 0.054, ΔAIC = −1.721) greater than

Mississauga, with the predicted maximum plant area being 3.9% greater than at a ΔMAT of zero.

Along our transect of sampled populations, a shift in the peak of the quadratic relationship

towards populations 2.0-3.1°C warmer than the Mississauga common garden represents a 2.1-

3.2° shift in latitude towards the equator. Additionally, whether or not plants flowered or

produced seeds increased linearly with ΔMAT, indicating plants of warmer origins were more

likely to flower and produce seeds (**Figure 4**), while the number of flower heads and growth rate

did not exhibit any significant linear or quadratic trends with ΔMAT (**Table 3**).

Limited evidence for a role of cyanogenesis in local adaptation

Both *Ac/ac* and *Li/li* exhibited strong latitudinal gene frequency clines among *T. repens*

populations from both continents (**Table S8**). However, the lack of an interaction between

latitude and continent of origin for both loci indicates no differences in the steepness of

cyanogenesis clines between the native and introduced ranges of *T. repens* (**Figure S5**). In the

full experiment, the main effect of cyanotype affected survival, whether or not plants produced

flowers and/or seeds, and maximum plant area, with cyanogenic plants generally exhibiting

greater fitness than acyanogenic plants, regardless of the environment in which the common

garden was found (**Table 4**). However, cyanotype alone did not affect the level of herbivory

occurring on *T. repens* plants, and cyanotype did not interact with either common garden

location or climatic distance to affect any measures of fitness or herbivory (**Table 4**).

Discussion

The aim of our study was to provide insight into adaptation and maladaptation of plants to

climate in native and introduced habitats. The results of our transcontinental experiment in the

native (Europe) and introduced (North America) ranges of *T. repens* lead to four main

conclusions: 1) *T. repens* exhibits stronger local adaptation to the abiotic environment at lower

latitudes compared to higher latitudes; 2) rapid local adaptation has contributed to the ecological

success of *T. repens* in its introduced range; 3) climate warming has resulted in some evidence of

a weakening of local adaptation; 4) cyanogenesis is not a major driver of local adaptation. These

conclusions are based on five experimental results. First, *T. repens* exhibited local adaptation to

climate overall, with greater fitness advantages for local populations at lower latitudes (Q1).

Second, in European common gardens, only native European populations exhibited local

adaptation, while in North American common gardens, only introduced North American

populations exhibited local adaptation (Q2). Third, both temperature- and precipitation-based

bioclimatic variables predicted fitness, but temperature was a stronger determinant of fitness at

higher latitudes, while precipitation was a stronger determinant of fitness at lower latitudes (Q3).

Fourth, there is some marginal evidence of an adaptation lag in response to climate warming, but

the signature was weak, and this pattern was only clear in the northern common garden of the

introduced range (Q4). Fifth, cyanogenesis did not differentially affect the fitness of *T. repens* in

different environments (Q5). Our results have important implications for the study of

environmental drivers of local adaptation, for projecting adaptation to future environmental

change, and for determining the role of rapid local adaptation in the success of biological

invasions.

Strength of local adaptation is dependent on latitude

Variation in biotic and abiotic factors across a species' range could affect its ability to adapt to

different environments within that range. We found that *T. repens* exhibited significant local

adaptation to the abiotic environment based on a measure of climatic distance that encompassed

mean annual temperature and precipitation along with annual temperature and precipitation

extremes. However, local adaptation differed significantly in strength depending on the latitude

of common gardens. Specifically, local adaptation was stronger at lower latitudes in both the

native and introduced ranges of *T. repens* when fitness was measured as sexual reproductive

success (i.e., number of flower heads and seed set mass). Our results are consistent with

transplant experiments using other plant species, which found sexual fitness advantages for local

versus foreign plants to be more pronounced at lower latitudes (Colautti and Barrett, 2013;

Toräng et al., 2015). Alternatively, life history trade-offs across large latitudinal ranges could

result in differences in the observed strength of fitness advantages, depending on the measure of

fitness involved (Li et al., 2015; Bastias et al., 2024). In fact, life history trade-offs, in *T. repens*

specifically, have been shown to contribute to local adaptation across divergent environments,

with southern populations investing more into sexual reproduction and northern populations

investing more into vegetative growth (Wright et al., 2021). If warmer climates are generally

more conducive to sexual reproduction than cooler climates, this may explain why sexual fitness

advantages for local populations were more pronounced in Lafayette and Montpellier in our

study, while climatic distance had more limited effects on vegetative growth.

While *T. repens* exhibits local adaptation based on an overall measure of climatic

distance, populations may vary in the individual bioclimatic variables responsible for local

adaptation. Our results demonstrate that at higher latitudes, where local adaptation is weaker,

temperature-based variables are stronger determinants of fitness, while at lower latitudes, where

local adaptation is stronger, precipitation-based variables are stronger determinants of fitness.

Therefore, it is possible that *T. repens* adapts more readily to local precipitation regimes than

local temperature regimes, resulting in stronger signatures of local adaptation in lower latitude

common gardens. This result is supported by a meta-analysis by Siepielski et al. (2017), which

found both precipitation and potential evapotranspiration to be stronger predictors of selection

than temperature across numerous terrestrial taxa. Alternatively, our results could be affected by

the specific environments chosen for the common gardens. In our study, the climates in Lafayette

and Montpellier are among the warmest that occur in the range of *T. repens*, and therefore could

have been especially difficult to tolerate for populations transplanted from cooler climates. In

turn, Uppsala was somewhat isolated, both geographically and climatically, from the sampled *T.*

repens populations, which could have affected our ability to detect local adaptation (or

maladaptation) in that environment. Lastly, it is possible that recent climate warming may have

already disrupted the ability of local populations to outperform foreign populations, thereby

impacting the strength of detectable local adaptation in certain locations.

Evidence of adaptation lags in response to climate warming

The ability of populations to adapt to recent rapid climate warming is an important mechanism

by which species can mitigate the risk of local extinction and other downstream ecological

consequences. Previous studies have demonstrated the incidence of adaptation lags, whereby

populations originating from slightly warmer climates would outperform populations with very

similar climatic origins to the common garden location (Wilczek et al., 2014; Aguirre-Liguori et

al., 2019; Anderson and Wadgymar, 2020). These studies indicate the potential for a disruption

in local adaptation, and even the possibility of maladaptation, due to climate warming that is

occurring faster than local populations are able to adapt.

In our northern common garden in North America (Mississauga), populations with

slightly warmer origins (2-3°C in MAT) outperformed local populations in survival and biomass

production (i.e., a proxy for vegetative reproduction) but not measures of sexual fitness. This

finding may signal the presence of a weak adaptation lag and may also provide a partial

explanation for the observed weaker local adaptation in Mississauga compared to Lafayette.

However, the generalizability of our results should be considered carefully for a few reasons.

First, contrary to our expectations, we provide no evidence of either local adaptation or

maladaptation in Uppsala. In a large-scale common garden experiment with European

populations of *Arabidopsis thaliana*, Wilczek et al. (2014) found that adaptation lags were more

pronounced in higher latitude common garden locations, which was not the case in our study.

Second, we tested for the presence of adaptation lags based solely on MAT, and Mississauga was

the common garden location where MAT most consistently affected plant fitness compared to

other bioclimatic variables. Fitness in lower latitude common gardens was instead more heavily

influenced by precipitation-based variables, which may not be changing as consistently as are

temperature-based variables due to anthropogenic activity. It is therefore possible that adaptation

lags to changing environments may not occur consistently across different geographic locations,

based on the specific biotic and abiotic factors that most strongly impact local adaptation in those

environments. Third, the MAT at the location of the Mississauga common garden in 2020 and

2021 was 1.6°C and 2.0°C warmer, respectively, than the average from the historical time period

(1970-2000) used to calculate climatic distances in this study (data extracted from ClimateNA;

Wang et al., 2016). Therefore, it may not be unexpected to see populations from slightly warmer

origins outperform local populations. The adaptation lag for survival does extend to a peak 3.4°C

warmer than Mississauga, and the proportion of plants producing flowers and/or seeds increased

linearly with ΔMAT, which still provides evidence for warm-adapted populations having an

advantage that exceeds expectations based on the MAT during the years of the experiment.

However, across our entire study, we are unable to make broad conclusions about future

vulnerability of populations based on the rate of environmental change exceeding the rate of

adaptation. Future studies investigating adaptation in response to environmental change may

benefit from approaches to experimental design that can allow for more accurate characterization

of the context-dependency of adaptation lags across a species' range.

Local adaptation in native and introduced ranges

Invasive species have the potential to become more problematic by rapidly adapting to novel

environments. Latitudinal cyanogenesis clines in North American *T. repens* have been shown to

be most likely a product of rapid local adaptation and sorting of existing genotypes post-

introduction (Kooyers and Olsen, 2012; Kooyers and Olsen, 2013). We found that populations of

T. repens only exhibited strong signatures of local adaptation to climate when transplanted into

common gardens within the same range as they originated. The observed strong local fitness

advantages for North American populations within the North American gardens are therefore

indicative of rapid local adaptation of *T. repens* within the past 400 years since its introduction to

North America. Selection has been shown to be particularly strong for introduced populations,

setting a precedent for rapid local adaptation among invasive species on even shorter timescales

than exhibited by *T. repens* (Colautti and Lau, 2015). A meta-analysis by Oduor et al. (2016)

found that invasive populations exhibited local adaptation just as frequently as native

populations, and that the strength of local adaptation was not affected by time since introduction

(ranging from as little as 20 to as many as 500 years). A reciprocal transplant experiment with

Lythrum salicaria indicated local adaptation to all three common garden sites spanning a 10°

latitudinal range, which occurred in ~200 years since its introduction to eastern North America

(Colautti and Barrett, 2013). Even faster still is adaptive evolution in the form of trait divergence

between coastal and inland environments in *Eschscholzia californica* in its introduced range,

which has occurred in the past 150 years (Leger and Rice, 2007). Similarly, clines in flowering

phenology of invasive *Hypericum canariense* have evolved within only 50 years (Dlugosch and

Parker, 2008). Taken together, our results and those of previous studies highlight the potential

for rapid local adaptation to climate, even across very large geographic ranges, in a small number

of generations post-introduction.

Interestingly, *T. repens* populations from the introduced range did not exhibit signatures

of local adaptation to similar climates when transplanted back into their native range. Similarly,

native populations did not maintain their strong signatures of local adaptation to climate when

transplanted into the introduced range. These findings demonstrate a clear cost of adaptation,

with local fitness advantages not extending across continents, even when accounting for

similarity among climates. Explanations for this lack of maintenance of local adaptation across

continents could be due to both biotic and abiotic aspects of the environment.

Abiotic explanations for the lack of local adaptation across continents could involve

shifts in photoperiod or precipitation regimes between Europe and North America due to similar

climates on the two continents being associated with different latitudes. The photoperiodic shift

between the continents could have affected the initiation of flowering and/or the timing of peak

flowering for populations that were transplanted across continents, leading to differences in the

strength of local adaptation. Halbritter et al. (2015) found that local adaptation was still evident

in reciprocal transplants conducted between high elevation and high latitude *Plantago*

populations with similar climates, allowing for speculation of a role of photoperiod in affecting

fitness. Additionally, populations of *Mimulus guttatus* exhibit clinal variation in the critical

photoperiod required to initiate flowering, potentially indicating a role of photoperiod in driving

local adaptation (Kooyers et al., 2015). Saikkonen et al. (2012) speculated that the transition to

less extreme yearly variation in photoperiod at lower latitudes might facilitate greater invasion

success of European species introduced to North America than vice versa. However, our results

do not provide evidence of increased success of European populations in North American

gardens compared to that of North American populations in European gardens. Additionally,

although we accounted for mean annual precipitation and yearly precipitation extremes in our

study, these variables do not fully capture the exact timing and volume of precipitation, and the

resulting hydrological environment experienced by *T. repens* plants in Europe versus North

America. For example, in the Montpellier garden in southern Europe, plants will experience a

Mediterranean climate that is not only drier but also more seasonally variable, while in the

Lafayette garden in southern North America, plants will experience a sub-tropical environment

that is wetter with more limited seasonal variation. This shift in precipitation availability has

caused introduced *T. repens* populations to evolve greater drought avoidance as opposed to the

more desiccation tolerant strategy observed in Mediterranean populations (Hendrickson et al.,

2024).

The biotic environment could be partially responsible for the lack of signatures of local

adaptation across continents based on climatic distance (Hargreaves et al, 2020). *Trifolium*

repens is a legume that forms mutualistic relationships with N-fixing bacteria in root nodules.

Murray-Stoker and Johnson (2024) have demonstrated the existence of locally adapted

populations in the *T. repens-Rhizobia* mutualism, indicating that local adaptation to biotic factors

is possible across environmental gradients or continents in this system. Furthermore, in line with

classic hypotheses in invasion biology, herbivory could exhibit substantial differences across

environments based on expected shifts in herbivore communities in different climates or across

continents, which could facilitate success of a plant species post-introduction (Coley et al., 1985;

Blossey and Nötzold, 1995; Keane and Crawley, 2002). If herbivory was likely to explain our

results of strong local adaptation within continent but not across continents, we might expect to

also see signatures of local adaptation based on the presence of chemical defenses in *T. repens*

plants.

Cyanogenesis and its metabolic components did not affect local adaptation

We tested whether the presence of latitudinal clines in plant defense indicates a role of defense

mechanisms in driving local adaptation in *T. repens*. Among our sampled populations, we

demonstrated rapid evolution of latitudinal clines in the presence of both *Ac* and *Li* alleles in the

introduced range of *T. repens*, based on their similar strength to clines in the native range.

Contrary to our results, all previous work has indicated weaker clines in the introduced range

than the native range (Daday, 1954a; Daday, 1958; Innes et al., 2022). Interestingly, our study

also did not demonstrate a role of cyanogenesis in local adaptation of *T. repens*, with no evidence

of interactions between cyanotype and climatic distance in influencing fitness. This result

indicates that neither cyanogenesis nor the presence/absence of cyanogenic glucosides or

linamarase exhibited differential effects on fitness across our common gardens. Previous studies

on local adaptation in *T. repens* found that HCN, cyanogenic glucosides, and linamarase only

had marginal effects on fitness and local adaptation, if any (Wright et al., 2018; Wright et al.,

2021). Further, *T. repens* does not consistently experience changes in cyanogenesis gene

frequency clines in response to recent climate change, which may indicate only marginal

involvement of cyanogenesis and/or its metabolic components in local adaptation (Innes et al.,

2022). Instead, divergence in life history strategies in response to climatic factors have been

identified as stronger drivers of local adaptation (Wright et al., 2018; Wright et al., 2021), which

is consistent with our results.

Our findings contrast with previous work that has speculated on the involvement of trade-

offs in the costs and benefits of cyanogenesis in different environments. Namely, previous

studies have hypothesized ecological costs of cyanogenesis in cooler climates due to freezing

autotoxicity (Daday, 1954a; Daday, 1965) or exacerbated allocation costs of producing the

metabolic components of HCN under freezing conditions (Kooyers et al., 2018; but see Fadoul et

al., 2023) as possible explanations for low frequencies of cyanogenesis in cooler climates. In our

study, cyanogenic plants often demonstrated a fitness advantage, even in Mississauga, which

does experience freezing temperatures for much of the year. This finding indicates that even in

cooler climates, the benefits of producing HCN as an anti-herbivore defense could outweigh the

costs of its production. However, despite overall effects of cyanotype on fitness in our common

gardens, we did not find evidence for cyanotype as deterrent of herbivory or as a driver of local

adaptation in *T. repens*. In fact, cyanotype did not affect the level of herbivory occurring on *T.*

repens plants, indicating that greater prevalence of cyanogenesis in warmer climates is likely not

based on an increased ability to defend against greater herbivore pressure.

Alternatively, there is evidence that the ability to produce cyanogenic glucosides and/or

linamarase could be advantageous in periods of drought through the storage and recycling of

nitrogen (Møller, 2010; Kooyers et al., 2014; Machingura et al., 2016). The first growing season

in Mississauga was uncharacteristically dry, resulting in the need for supplemental watering of

the common garden to avoid mortality of the full experiment. It is possible that the dry

conditions may have also contributed to a fitness advantage for *Ac* and/or *Li* plants. However,

any advantages of particular cyanotypes within a given common garden environment did not

confer higher fitness to local plants as opposed to foreign plants based on our measure of

climatic distance. Furthermore, we were unable to harvest enough tissue to determine the

cyanotype of most of the plants in the study due to concerns over effects of tissue harvest on

plant fitness and/or mortality, so the strength of our conclusions with respect to the role of

cyanogenesis in adaptation are limited by the relatively low sample size for all statistical

analyses involving cyanotype. Further field studies are needed to more explicitly test the costs

and benefits of HCN, cyanogenic glucosides, and linamarase, and their effects on local

adaptation to climate in *T. repens* across large latitudinal ranges.

Limitations

While our study offers strong tests of local adaptation to both biotic and abiotic factors in the

native and introduced ranges of a cosmopolitan plant, there were limitations that should be

considered when interpreting our findings. In particular, we are unable to determine the relative

strength of soil conditions and mutualistic interactions compared to climate as drivers of local

adaptation. Planting *T. repens* individuals from each population along with the natural soil and

microbial communities from their location of origin was infeasible considering our study design

and governmental regulations on importation of soil and bacteria. Instead, the best alternative

was to maintain consistent treatment of all individuals in all gardens, with *T. repens* individuals

planted into the natural soil at each common garden site. It is important to recognize that local

edaphic factors have the potential to provide fitness advantages in *T. repens*, as has been

demonstrated in other systems (Macel et al., 2007). It is also possible that mutualistic biotic

interactions between *T. repens* and nitrogen-fixing bacteria could be involved in providing

fitness advantages to local populations (Murray-Stoker and Johnson, 2024). However, we did not

account for local soil conditions or any biotic interactions apart from plant-herbivore interactions

in the analysis for our study. Despite these limitations, signatures of local adaptation to climate

were still evident across multiple measures of fitness, strengthening our confidence in the

observed patterns.

Conclusions

We found evidence of strong local adaptation in the native and introduced ranges of a

cosmopolitan plant species, but also that local adaptation is highly context-dependent. Namely,

local adaptation was stronger in lower latitude common garden locations in both native and

introduced ranges, and signatures of local adaptation were only evident when populations were

transplanted within the same range as they originated. These findings indicate both strong and

rapid local adaptation in an introduced range of *T. repens* in less than 400 years since

introduction. However, the northern garden in the introduced range also provided some evidence

of local maladaptation compared to populations with slightly warmer origins, which is

characteristic of an adaptation lag. Therefore, while local adaptation has occurred quickly in

introduced populations of *T. repens*, it is possible that local adaptation is being weakened by the

effects of recent climate change, which could eventually result in more severe erosion of fitness

advantages for local populations in the future. The ability of plants to rapidly adapt to

anthropogenic environmental change is vital to the integrity of biodiversity in the face of future

change. Understanding ecological factors and the interactions between them that are likely to

impact plant adaptation is therefore essential to informing future management decisions

involving the maintenance of biodiversity.

Acknowledgements

We thank James Santangelo for assistance with initial seed collection, and Happy Gill,

Jashandeep Nijjar, and Ruth Thomas for assisting with the generation of F1 plants for use in the

experiment. We are indebted to Aaron Albano, Barry Albano, Paula Albano, Sarah Albano,

Andre Daugeureaux, Hayden Fargo, Axel Hjelm, Antoniette Jansen, Eunice Karinho Betancourt,

Erik Karlsson, Andreas Lundgren, Kavya Manikonda, Lindsay Miles, David Murray-Stoker,

Bernabé Ramirez, Brittany Sutherland, Olivia Toth, Andrea Turcu, and Mariona Vinyeta Medina

for assistance in the field. Keerath Bhachu, Elana Maria, Zain Nasrullah, Nial Navaratne,

Narindra Persaud, and Isabellio Vessio assisted with sample processing. We also thank the

Louisiana Optical Network (LONI) for providing computation resources, along with Kris

Horvath, Joan Lee, Kent Moore, Diane Ross, Vera Velasco, the technical platform "Plateforme

des Terrains d"Expérience" (CEFE-CNRS), and the University of Louisiana Ecology Center for

facilitating and supporting the set-up and execution of the experiment. This research was

supported by an NSERC Discovery Grant, Canada Research Chair, and NSERC EWR Steacie

Fellowship to MTJJ, a CNRS-University of Toronto Ph.D. Student Travel Grant to MTJJ and

CV, NSF grants OIA-1920858 and DBI-2244712 to NJK, and an NSERC CGS Doctoral Award

to LJA.

Data Availability Statement

All data involved in this study is available on the GitHub page for LJA

(https://github.com/ljalbano/transcontinental_common_garden).

References

Adams, J. M., Fang, W., Callaway, R. M., Cipollini, D., Newell, E., Transatlantic Acer

platanoides Invasion Network (TRAIN). (2009). A cross-continental test of the Enemy

Release Hypothesis: leaf herbivory on *Acer platanoides* (L.) is three times lower in North

America than in its native Europe. *Biological Invasions*, *11*, 1005-1016.

Agrawal, A. A., Kotanen, P. M. (2003). Herbivores and the success of exotic plants: a

phylogenetically controlled experiment. *Ecology Letters*, *6*, 712-715.

Agrawal, A. A. (2007). Macroevolution of plant defense strategies. *Trends in Ecology and*

Evolution, *22*, 103-109.

Aguirre-Liguori, J. A., Ramírez-Barahona, S., Tiffin, P., Eguiarte, L. E. (2019). Climate change

is predicted to disrupt patterns of local adaptation in wild and cultivated maize.

Proceedings of the Royal Society B, *286*, 20190486.

Albano, L. J., Johnson, M. T. J. (2023). Interactions environmental factors drive selection on

cyanogenesis in *Trifolium repens*. *Oikos*, *2023*, e09629.

Anderson, J. T., Song, B.-H. (2020). Plant adaptation to climate change – Where are we? *Journal*

of Systematics and Evolution, *58*, 533-545.

Anderson, J. T., Wadgymar, S. M. (2020). Climate change disrupts local adaptation and favours

upslope migration. *Ecology Letters*, *23*, 181-192.

Angseesing, J. P. A. (1974). Selective eating of the acyanogenic form of *Trifolium*

repens. *Heredity*, *32*, 73.

Anstett, D. N., Nunes, K. A., Baskett, C., Kotanen, P. M. (2016). Sources of controversy

surrounding latitudinal patterns in herbivory and defense. *Trends in Ecology & Evolution*,

31, 789-802.

Anstett, D. N., Ahern, J. R., Johnson, M. T. J., Salminen, J.-P. (2018). Testing for latitudinal

gradients in defense at the macroevolutionary scale. *Evolution*, *72*, 2129-2143.

Atkins, K. E., Travis, J. M. J. (2010). Local adaptation and the evolution of species' ranges under

781   climate change. *Journal of Theoretical Biology*, *266*, 449-457.

782   Atwood, S. S., Sullivan, J. T. (1943). Inheritance of a cyanogenetic glucoside and its hydrolyzing

783   enzyme in *Trifolium repens*. *Journal of Heredity*, *34*, 311-320.

784   Baskett, C. A., Schemske, D. W. (2018). Latitudinal patterns of herbivore pressure in a

785   temperature herb support the biotic interactions hypothesis. *Ecology Letters*, *21*, 578-587.

786   Bastias, C. C., Estarague, A., Vile, D., Gaignon, E., Lee, C.-R., Exposito-Alonso, M., … ,

787   Vasseur, F. (2024). Ecological trade-offs drive phenotypic and genetic differentiation of

788   *Arabidopsis thaliana* in Europe. *Nature Communications*, *15*, 5185.

789   Bidart-Bouzat, M. G., Imeh-Nathaniel, A. (2008). Global change effects on plant chemical

790   defenses against insect herbivores. *Journal of Integrative Plant Biology*, *50*, 1339-1354.

791   Blossey, B., Nötzold, R. (1995). Evolution of increased competitive ability in invasive

792   nonindigenous plants: A hypothesis. *Journal of Ecology*, *83*, 887-889.

793   Bontrager, M., Muir, C. D., Mahony, C., Gamble, D. E., Germain, R. M., Hargreaves, A. L., … ,

794   Angert, A. L. (2020). Climate warming weakens local adaptation. *biorxiv*. doi:

795   10.1101/2020.11.01.364349

796   Brighton, F., Horne, M. T. (1977). Influence of temperature on cyanogenic polymorphisms.

797   *Nature*, *265*, 437-438.

798   Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., … ,

799   Bolker, B. M. (2014) glmmTMB balances speed and flexibility among packages for zero-

800   inflated generalized linear mixed modeling. *The R Journal*, *9*, 378-400.

Burdon, J. J. (1983). *Trifolium repens* L. *Journal of Ecology*, *71*, 307-330.

Colautti, R. I., Barrett, S. C. H. (2013). Rapid adaptation to climate facilitates range expansion of

an invasive plant. *Science*, *342*, 364-367.

Colautti, R. I., Lau, J. A. (2015). Contemporary evolution during invasion: evidence for

differentiation, natural selection, and local adaptation. *Molecular Ecology*, *24*, 1999-

2017.

Coley, P. D., Bryant, J. P., Chapin, F. S. (1985). Resource availability and plant antiherbivore

defense. *Science*, *230*, 895-899.

Coley, P. D., Barone, J. A. (1996). Herbivory and plant defenses in tropical forests. *Annual*

Review of Ecology and Systematics, *27*, 305-335.

Corkill, L. D. (1940). Cyanogenesis in white clover (*Trifolium repens* L.). I. Cyanogenesis in

single plants. *New Zealand Journal of Science and Technology, 22*, 65-67.

Costan, C.-A., Godsoe, W. K., Bufford, J. L., Marris, J. W. M., Hulme, P. E. (2022). Can the

enemy release hypothesis explain the success of *Rumex* (Polygonaceae) species in an

introduced range? *Biological Invasions*, *24*, 2665-2677.

Daday, H. (1954a). Gene frequencies in wild populations of *Trifolium repens*. I. Distribution by

latitude. *Heredity*, *8*, 61-78.

Daday, H. (1954b). Gene frequencies in wild populations of *Trifolium repens*. II. Distribution by

altitude. *Heredity*, *8*, 377-384.

Daday, H. (1958). Gene frequencies in wild populations of *Trifolium repens*. III. World

distribution. *Heredity*, *12*, 169-184.

Daday, H. (1965). Gene frequencies in wild populations of *Trifolium repens*. IV. Mechanism of

natural selection. *Heredity*, *20*, 355-365.

Dirzo, R., Harper, J. L. (1982). Experimental studies on slug-plant interactions: III. Differences

in the acceptability of individual plants of *Trifolium repens* to slugs and snails. *Journal of*

Ecology, *70*, 101-117.

Dlugosch, K. M., Parker, I. M. (2008). Invading populations of an ornamental shrub show rapid

life history evolution despite genetic bottlenecks. *Ecology Letters*, *11*, 701-709.

Dostálek, T., Rokaya, M. B., Münzbergová, Z. (2020). Plant palatability and trait responses to

experimental warming. *Scientific Reports*, *10*, 10526.

Easlon, H. M., Bloom, A. J. (2014). Easy Leaf Area: Automated digital image analysis for rapid

and accurate measurement of leaf area. *Applications in Plant Sciences*, *2*, 1400033.

Ehrlich, P. R., Raven, P. H. (1964). Butterflies and plants: A study in coevolution. *Evolution*, *18*,

586-608.

Endler, J. A. (1977). Geographic variation, speciation, and clines. Princeton University Press.

Exposito-Alonso, M., 500 Genomes Field Experiment Team, Burbano, H. A., Bossdorf, O.,

Nielsen, R., Weigel, D. (2019). Natural selection on the *Arabidopsis thaliana* genome in

present and future climates. *Nature*, *573*, 126-129.

Fadoul, H. E., Albano, L. J., Bergman, M. E., Phillips, M. A., Johnson, M. T. J. (2023).

Assessing the benefits and costs of the hydrogen cyanide antiherbivore defense in

Trifolium repens. *Plants*, *12*, 1213.

Fine, P. V. A., Miller, Z. J., Mesones, I., Irazuzta, S., Appel, H. M., Stevens, M. H. H., … ,

Coley, P. D. (2006). The growth-defense trade-off and habitat specialization by plants in

Amazonian forests. *Ecology*, *87*, S150-S162.

Fox, J., Weisberg, S. (2019). An R companion to applied regression, Third edition. Sage,

Thousand Oaks, CA, USA.

Gillespie, J. H., Turelli, M. (1989). Genotype-environment interactions and the maintenance of

polygenic variation. *Genetics*, *121*, 129-138.

Gleadow, R. M., Woodrow, I. E. (2002). Constraints on effectiveness of cyanogenic glycosides

in herbivore defense. *Journal of Chemical Ecology*, *28*, 1301-1313.

Griffiths, A. G., Moraga, R., Tausen, M., Gupta, V., Bitton, T. P., Campbell, M. A., … ,

Anderson, S. U. (2019). Breaking free: The genomics of allopolyploidy-facilitated niche

expansion in white clover. *The Plant Cell*, *31*, 1466-1487.

Hahn, P. G., Agrawal, A. A., Sussman, K. I., Maron, J. L. (2019). Population variation,

environmental gradients, and the evolutionary ecology of plant defense against herbivory.

The American Naturalist, *193*, 20-34.

Halbritter, A. H., Billeter, R., Edwards, P. J., Alexander, J. M. (2015). Local adaptation at range

edges: comparing elevation and latitudinal gradients. *Journal of Evolutionary Biology*,

28, 1849-1860.

Hargreaves, A. L., Germain, R. M., Bontrager, M., Persi, J., Angert, A. L. (2020). Local

adaptation to biotic interactions: A meta-analysis across latitudes. *The American

Naturalist*, *195*, 395-411.

Hendrickson, B. T., Stamps, C., Patterson, C. M., Strickland, H., Foster, M., Kim, A., … ,

Kooyers, N. J. (2024). Evolution of drought resistant strategies following the introduction

of white clover (*Trifolium repens* L.). *biorxiv*. doi: 10.1101/2024.09.03.611023

Hereford, J. (2009). A quantitative survey of local adaptation and fitness trade-offs. *The

American Naturalist*, *173*, 579-588.

Herms, D. A., Mattson, W. J. (1992). The dilemma of plants: to grow or defend. *Quarterly Review of

Biology*, *67*, 478-478.

Hoffmann, A. A., Sgrò, C. M. (2011). Climate change and evolutionary adaptation. *Nature*, *470*,
479-485.

Hughes, M. A. (1991). The cyanogenic polymorphism in *Trifolium repens* L. (white clover).
Heredity, *66*, 105-115.

Innes, S., Santangelo, J. S., Kooyers, N. J., Olsen, K. M., Johnson, M. T. J. (2022). Evolution in
response to climate in the native and introduced ranges of a globally distributed plant.
Evolution, *76*, 1495-1511.

IPBES. (2019). Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
Brondizio, E. S., Settele, J., Díaz, S., Ngo, H. T. (editors). IPBES secretariat, Bonn,
Germany.

Johnson, M. T. J., Rasmann, S. (2011). The latitudinal herbivory-defence hypothesis takes a
detour on the map. *New Phytologist*, *191*, 589-592.

Johnson, M. T. J., Bertrand, J. A., Turcotte, M. M. (2016). Precision and accuracy in quantifying
herbivory. *Ecological Entomology*, *41*, 112-121.

Johnson, M. T. J., Munshi-South, J. (2017). Evolution of life in urban environments. *Science*,
358, 6363.

Kakes, P. (1989). An analysis of the costs and benefits of the cyanogenic system in *Trifolium*
repens L. *Theoretical and Applied Genetics*, *77*, 111-118.

Kassambara, A., Mundt, F. (2020). factoextra: Extract and visualize the results of multivariate
data analyses. R package version 1.0.7. https://CRAN.R-project.org/package=factoextra

Kawecki, T. J., Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, *7*, 1225-
1241.

Keane, R. M., Crawley, M. J. (2002). Exotic plant invasions and the enemy release hypothesis.

Trends in Ecology & Evolution, *17*, 164-170.

Kjærgaard, T. (2003). A plant that changed the world: The rise and fall of clover 1000-2000.

Landscape Research, *28*, 41-49.

Kooyers, N. J., Olsen, K. M. (2012). Rapid evolution of an adaptive cyanogenesis cline in

introduced North American white clover (*Trifolium repens* L.). *Molecular Ecology*, *21*,

2455-2468.

Kooyers, N. J., Olsen, K. M. (2013). Searching for the bull's eye: agents and targets of selection

vary among geographically disparate cyanogenesis clines in white clover (*Trifolium*

repens L.). *Heredity*, *11*, 495-504.

Kooyers, N. J., Gage, L. R., Al-Lozi, A., Olsen, K. M. (2014). Aridity shapes cyanogenesis cline

evolution in white clover (*Trifolium repens* L.). *Molecular Ecology*, *23*, 1053-1070.

Kooyers, N. J., Greenlee, A. B., Colicchio, J. M., Oh, M., Blackman, B. K. (2015). Replicate

altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress

in annual *Mimulus guttatus*. *New Phytologist*, *206*, 152-165.

Kooyers, N. J., Hartman Bakken, B., Ungerer, M. C., Olsen, K. M. (2018). Freeze-induced

cyanide toxicity does not maintain the cyanogenesis polymorphism in white clover

(*Trifolium repens*). *American Journal of Botany*, *105*, 1224-1231.

Kooyers, N. J., Colicchio, J. M., Greenlee, A. B., Patterson, E., Handloser, N. T., Blackman, B.

K. (2019). Lagging adaptation to climate supersedes local adaptation to herbivory in an

annual monkeyflower. *The American Naturalist*, *194*, 541-557.

Lee, C.-R., Mitchell-Olds, T. (2011). Quantifying effects of environmental and geographical

factors on patterns of genetic differentiation. *Molecular Ecology*, *20*, 4631-4642.

Leger, E. A., Rice, K. J. (2007). Assessing the speed and predictability of local adaptation in

invasive California poppies (*Eschscholzia californica*). *Journal of Evolutionary Biology*,

20, 1090-1103.

Leimu, R., Fischer, M. (2008). A meta-analysis of local adaptation in plants. *PLoS One*, *3*,

e4010.

Lemoine, N. P., Drews, W. A., Burkepile, D. E., Parker, J. D. (2013). Increased temperature

alters feeding behaviour of a generalist herbivore. *Oikos*, *122*, 1669-1678.

Li, X.-M., She, D.-Y., Zhang, D.-Y., Liao, W.-J. (2015). Life history trait differentiation and

local adaptation in invasive populations of *Ambrosia artemisiifolia* in China. *Oecologia*,

177, 669-677.

Macel, M., Lawson, C. S., Mortimer, S. R., Šmilauerova, M., Bischoff, A., Crémieux, L., … ,

Steinger, T. (2007). Climate vs. soil factors in local adaptation of two common plant

species. *Ecology*, *88*, 424-433.

Machingura, M., Salomon, E., Jez, J. M., Ebbs, S. D. (2016). The β-cyanoalanine synthase

pathway: beyond cyanide detoxification. *Plant, Cell and Environment*, *39*, 2329-2341.

Maron, J. L., Vilà, M., Bommarco, R., Elmendorf, S., Beardsley, P. (2004). Rapid evolution of

an invasive plant. *Ecological Monographs*, *74*, 261-280.

Massad, T. J., Dyer, L. A. (2010). A meta-analysis of the effects of global environmental change

on plant-herbivore interactions. *Arthropod-Plant Interactions*, *4*, 181-188.

Miles, L. S., Breitbart, S. T., Wagner, H. H., Johnson, M. T. J. (2019). Urbanization shapes the

ecology and evolution plant-arthropod herbivore interactions. *Frontiers in Ecology and*

Evolution, *7*, 310.

Møller, B. L. (2010). Functional diversifications of cyanogenic glucosides. *Current Opinion in*
Plant Biology, *13*, 338-347.

Moreira, X., Castagneyrol, B., Abdala-Roberts, L., Berny-Mier y Teran, J. C., Timmermans, B.
G. H., Bruun, H. H., … , Tack, A. J. M. (2018). Latitudinal variation in plant chemical
defences drives latitudinal patterns of leaf herbivory. *Ecography*, *41*, 1124-1134.

Murray-Stoker, D., Johnson, M. T. J. (2024). Mosaic of local adaptation between white clover
and rhizobia along an urbanization gradient. *Journal of Ecology*, *112*, 1150-1163..

O'Donnell, M. S., Ignizio, D. A. (2012). Bioclimatic predictors for supporting ecological
applications in the conterminous United States. *US Geological Survey Data Series*, *10*, 4-
9.

Oduor, A. M. O., Leimu, R., van Kleunen, M. (2016). Invasive plant species are locally adapted
just as frequently and at least as strongly as native plant species. *Journal of Ecology*, *104*,
957-968.

Olsen, K. M., Sutherland, B. L., Small, L. L. (2007). Molecular evolution of the *Li/li* chemical
defence polymorphism in white clover (*Trifolium repens* L.). *Molecular Ecology*, *16*,
4180-4193.

Olsen, K. M., Small, L. L. (2018). Micro- and macroevolutionary adaptation though repeated
loss of a complete metabolic pathway. *New Phytologist*, *219*, 757-766.

Olsen, K. M., Goad, D. M., Wright, S. J., Dutta, M. L., Myers, S. R., Small, L. L., Li, L.-F.
(2021). Dual-species origin of an adaptive chemical defense polymorphism. *New*
Phytologist, *232*, 1477-1487.

Saikkonen, K., Taulavuori, K., Hyvönen, T., Gundel, P. E., Hamilton, C. E., Vänninen, I., … ,

Helander, M. (2012). Climate change-driven species' range shifts filtered by

photoperiodism. *Nature Climate Change*, *2*, 239-242.

Samis, K. E., Stinchcombe, J. R., Murren, C. J. (2019). Population climatic history predicts

phenotypic responses in novel environments for *Arabidopsis thaliana* in North America.

American Journal of Botany, *106*, 1068-1080.

Santangelo, J. S., Ness, R. W., Cohan, B., Fitzpatrick, C. R., Innes, S. G., Koch, S., … , Johnson,

M. T. J. (2022). Global urban environmental change drives adaptation in white clover.

Science, *375*, 1275-1281.

Savolainen, O., Lascoux, M., Merilä, J. (2013). Ecological genomics of local adaptation. *Nature*

Reviews Genomics, *14*, 807-820.

Siepielski, A. M., Morrissey, M. B., Buoro, M., Carlson, S. M., Caruso, C. M., Clegg, S. M., … ,

MacColl, A. D. C. (2017). Precipitation drives global variation in natural selection.

Science, *355*, 959-962.

St Clair, J. B., Howe, G. T. (2007). Genetic maladaptation of coastal Douglas-fir seedlings to

future climates. *Global Change Biology*, *13*, 1441-1454.

Stinchcombe, J. R., Weinig, C., Ungerer, M., Olsen, K. M., Mays, C., Halldorsdottir, S. S., … ,

Schmitt, J. (2004). A latitudinal cline in flowering time in *Arabidopsis thaliana*

modulated by the flowering time gene FRIGIDA. *Proceedings of the National Academy*

of Sciences of the United States of America, *101*, 4712-4717.

Strauss, S. Y., Rudgers, J. A., Lau, J. A., Irwin, R. E. (2002). Direct and ecological costs of

resistance to herbivory. *Trends in Ecology & Evolution*, *17*, 278-285.

Thompson, K. A., Renaudin, M., Johnson, M. T. J. (2016). Urbanization drives the evolution of

parallel clines in plant populations. *Proceedings of the Royal Society B: Biological*

Sciences, *283*, 20162180.

Till, I. (1987). Variability of expression of cyanogenesis in white clover (*Trifolium repens*

L.). *Heredity*, *59*, 265.

Toräng, P., Wunder, J., Obeso, J. T., Herzog, M., Coupland, G, Ågren, J. (2015). Large-scale

adaptative differentiation in the alpine perennial *Arabis alpina*. *New Phytologist*, *206*,

459-470.

Turesson, G. (1922). The genotypical response of the plant species to the habitat. *Hereditas*, *3*,

211-350.

Wang, T., Hamann, A., Spittlehouse, D., Carroll, C. (2016). Locally downscaled and spatially

customizable climate data for historical and future periods for North America. *PLoS One*,

11, e0156720.

Wilczek, A. M., Cooper, M. D., Korves, T. M., Schmitt, J. (2014). Lagging adaptation to

warming climate in *Arabidopsis thaliana*. *Proceedings of the National Academy of*

Sciences, *111*, 7906-7913.

Wolkovich, E. M., Davies, T. J., Schaefer, H., Cleland, E. E., Cook, B. I., Travers, S. E., … ,

Davis, C. C. (2013). Temperature-dependent shifts in phenology contribute to the success

of exotic species with climate change. *American Journal of Botany*, *100*, 1407-1421.

Wright, S. J., Cui Zhou, D., Kuhle, A., Olsen, K. M. (2018). Continent-wide climatic variation

drives local adaptation in North American white clover. *Journal of Heredity*, *109*, 78-89.

Wright, S. J., Goad, D. M., Gross, B. L., Muñoz, P. R., Olsen, K. M. (2021). Genetic trade-offs

underlie divergent life strategies for local adaptation in white clover. *Molecular Ecology*,

31, 3742-3760.

Züst, T., Heichinger, C., Grossniklaus, U., Harrington, R., Kliebenstein, D. J., Turnbull, L. A.

(2012). Natural enemies drive geographic variation in plant defenses. *Science*, *338*, 116-

119.

Tables and Figures

Table 1 Results of mixed effects models assessing how continent of origin, common garden location, and climatic distance affected
the fitness of *T. repens*. Measures of fitness include whether or not plants produced flowers and/or seeds, number of flower heads,
seed set mass, survival, maximum plant area, growth rate, and herbivory. Significant effects ($P < 0.05$) are listed in bold.

1012

Predictor	Plants Producing Flowers			Plants Producing Seeds			# of Flower Heads			Seed Set Mass		
	χ^2	df	*P*	χ^2	df	*P*	χ^2	df	*P*	χ^2	df	*P*
Continent of Origin (O)	0.330	1	0.566	0.218	1	0.641	2.314	1	0.128	1.214	1	0.271
Garden Location (L)	**31.703**	**3**	**< 0.001**	**8.104**	**3**	**0.044**	**252.144**	**3**	**< 0.001**	**40.068**	**3**	**< 0.001**
Climatic Distance (D)	**6.015**	**1**	**0.014**	**6.882**	**1**	**0.009**	**13.580**	**1**	**< 0.001**	**14.240**	**1**	**< 0.001**
O × L	0.998	3	0.802	0.952	3	0.813	1.442	3	0.696	4.085	3	0.252
O × D	0.421	1	0.516	1.113	1	0.291	0.277	1	0.599	0.978	1	0.323
L × D	4.439	3	0.218	4.185	3	0.242	**12.713**	**3**	**0.005**	**8.808**	**3**	**0.032**
O × L × D	6.860	3	0.077	7.631	3	0.054	7.764	3	0.051	4.882	3	0.181

1013

1014

1015

Predictor	Survival			Maximum Plant Area			Growth Rate			Herbivory		
	χ^2	df	*P*	χ^2	df	*P*	χ^2	df	*P*	χ^2	df	*P*
Continent of Origin (O)	2.220	1	0.136	0.939	1	0.332	0.094	1	0.759	0.089	1	0.766
Garden Location (L)	**221.162**	**3**	**< 0.001**	**286.435**	**3**	**< 0.001**	**70.328**	**3**	**< 0.001**	**144.426**	**3**	**< 0.001**
Climatic Distance (D)	**6.427**	**1**	**0.011**	0.584	1	0.445	0.089	1	0.766	1.074	1	0.300
O × L	6.155	3	0.104	6.464	3	0.091	1.212	3	0.750	6.635	3	0.084
O × D	0.142	1	0.707	1.747	1	0.186	0.049	1	0.824	0.010	1	0.919
L × D	4.798	3	0.187	5.610	3	0.132	2.887	3	0.409	2.234	3	0.525
O × L × D	7.288	3	0.063	2.693	3	0.441	3.399	3	0.334	3.127	3	0.372

1019

1020

1021

1022 **Table 2** Results of mixed effects models assessing how continent of origin, continent of garden, and climatic distance affected the
1023 fitness of *T. repens*. Measures of fitness include whether or not plants produced flowers and/or seeds, number of flower heads, seed set
1024 mass, survival, maximum plant area, growth rate, and herbivory. Significant effects ($P < 0.05$) are listed in bold. All predictors are
1025 assessed using one degree of freedom.
1026

Predictor	Plants Producing Flowers		Plants Producing Seeds		# of Flower Heads		Seed Set Mass7	
	χ^2	P	χ^2	P	χ^2	P	χ^2	P
Continent of Origin (O)	2.033	0.154	1.754	0.185	0.514	0.474	0.028	0.867
Continent of Garden (G)	**9.184**	**0.002**	1.295	0.255	0.471	0.493	0.037	0.847
Climatic Distance (D)	**28.963**	**< 0.001**	**17.269**	**< 0.001**	**14.310**	**< 0.001**	**16.190**	**< 0.001**
O × G	0.677	0.411	0.942	0.332	0.957	0.328	0.481	0.488
O × D	2.033	0.154	0.036	0.850	1.214	0.271	2.991	0.084
G × D	3.194	0.074	2.490	0.115	1.412	0.235	0.258	0.612
O × G × D	0.501	0.479	0.026	0.871	**19.603**	**< 0.001**	**9.728**	**0.002**

1028

Predictor	Survival		Maximum Plant Area		Growth Rate		Herbivory)29 1030	
	χ^2	P	χ^2	P	χ^2	P	χ^2	P 1031
Continent of Origin (O)	0.106	0.745	0.797	0.372	0.053	0.817	0.196	0.658
Continent of Garden (G)	**41.220**	**< 0.001**	**170.553**	**< 0.001**	1.468	0.226	**21.180**	**< 0.001**
Climatic Distance (D)	**10.236**	**0.001**	**58.341**	**< 0.001**	**18.137**	**< 0.001**	**15.810**	**< 0.001**
O × G	0.000	0.993	2.375	0.123	0.465	0.495	0.669	0.413
O × D	1.534	0.216	**34.173**	**< 0.001**	**5.145**	**0.023**	**23.686**	**< 0.001**
G × D	**13.870**	**< 0.001**	1.784	0.182	2.263	0.133	0.240	0.624
O × G × D	**6.515**	**0.011**	0.678	0.410	**7.812**	**0.005**	2.178	0.140

Table 3 Results of mixed effects models assessing how change in mean annual temperature (ΔMAT) affected the fitness of North American *T. repens* populations in the Mississauga common garden (linear and quadratic fits). Significant effects ($P < 0.05$) are listed in bold. All predictors are assessed using one degree of freedom.

Predictor	Survival		Plants Producing Flowers		Plants Producing Seeds		# of Flower Heads		Seed Set Mass		Maximum Plant Area		Growth Rate	
	χ^2	P	χ^2	P	χ^2	P	χ^2	P	χ^2	P	χ^2	P	χ^2	P
ΔMAT	3.758	0.053	**7.425**	**0.006**	**5.397**	**0.020**	0.255	0.614	0.644	0.422	3.799	0.051	0.002	0.962
ΔMAT ^{2}	**4.313**	**0.038**	**8.077**	**0.004**	**7.609**	**0.006**	2.908	0.088	**8.542**	**0.003**	**6.741**	**0.009**	1.808	0.179

1050

Table 4 Results of mixed effects models assessing how cyanotype (AcLi, Acli, acLi, or acli), common garden location, and climatic distance affected the fitness of *T. repens*. Measures of fitness include survival, whether or not plants produced flowers and/or seeds, number of flower heads, seed set mass, maximum plant area, growth rate, and herbivory. Significant effects ($P < 0.05$) are listed in bold.

1055

Predictor	Plants Producing Flowers			Plants Producing Seeds			# of Flower Heads			Seed Set Mass		
	χ^2	df	P	χ^2	df	P	χ^2	df	P	χ^2	df	P
Cyanotype (C)	**16.862**	**3**	**< 0.001**	**11.731**	**3**	**0.008**	6.635	3	0.084	1.801	3	0.615
Garden Location (L)	**29.547**	**3**	**< 0.001**	**43.214**	**3**	**< 0.001**	**125.477**	**3**	**< 0.001**	**19.881**	**3**	**< 0.001**
Climatic Distance (D)	**9.464**	**1**	**0.002**	**9.816**	**1**	**0.002**	**22.595**	**1**	**< 0.001**	**23.040**	**1**	**< 0.001**
C × L	13.049	9	0.160	11.819	9	0.224	6.558	9	0.683	2.835	9	0.970
C × D	2.089	3	0.554	3.236	3	0.357	3.926	3	0.270	4.463	3	0.216
C × L × D	10.144	12	0.603	13.401	12	0.341	11.509	12	0.486	10.804	12	0.546

1056

Predictor	Survival			Maximum Plant Area			Growth Rate			Herbivory		
	χ^2	df	P	χ^2	df	P	χ^2	df	P	χ^2	df	P
Cyanotype (C)	**19.889**	**3**	**< 0.001**	**9.841**	**3**	**0.020**	0.381	3	0.944	4.956	3	0.175
Garden Location (L)	**56.896**	**3**	**< 0.001**	**116.238**	**3**	**< 0.001**	**79.026**	**3**	**< 0.001**	**85.426**	**3**	**< 0.001**
Climatic Distance (D)	**3.903**	**1**	**0.048**	**6.887**	**1**	**0.009**	3.835	1	0.050	0.237	1	0.626
C × L	8.506	9	0.484	7.049	9	0.632	6.369	9	0.703	3.326	9	0.950
C × D	1.723	3	0.632	2.726	3	0.436	1.251	3	0.741	3.724	3	0.293
C × L × D	6.019	12	0.915	6.575	12	0.884	19.085	12	0.087	8.448	12	0.749

1057

Figure 1 Maps showing populations of *T. repens* sampled from the (**A**) introduced range (North America, orange circles) and (**B**) native range (Europe, pink circles), as well as the four common garden locations (green diamonds). Coordinates of common garden locations are 42°32'40" N, 79°39'38" W (Mississauga, Ontario, Canada), 30°18'23" N, 92°00'33" W (Lafayette, Louisiana, USA), 59°49'08" N, 17°38'49" W (Uppsala, Sweden), and 43°38'16" N, 3°51'43" W (Montpellier, France).

1064
1065   **Figure 2** The effect of climatic distance and common garden location on fitness of *T. repens*. (**A**) Mean number of flower heads and
1066   (**B**) mean seed set mass of *T. repens* populations planted in two common gardens in North America (Lafayette in navy and
1067   Mississauga in orange) and two common gardens in Europe (Montpellier in green and Uppsala in magenta). Shaded areas represent
1068   95% confidence intervals around each line, while asterisks represent *P* values of relevant fixed effects of Garden Location (L) and
1069   Climatic Distance (D) and their interaction (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$).
1070
1071
1072
1073

Figure 3 The effect of climatic distance on the mean number of flower heads produced by *T. repens* populations originating from its
introduced (North America, orange) and native (Europe, pink) ranges and transplanted into common gardens in the (**A**) introduced and
(**B**) native ranges. The effect of climatic distance on the mean seed set mass produced by *T. repens* populations originating from its
introduced and native ranges and transplanted into common gardens in the (**C**) introduced and (**D**) native ranges. Shaded areas
represent 95% confidence intervals around each line, while asterisks represent *P* values of relevant fixed effects of Climatic Distance
(D) and its interaction with Continent of Origin (O) and Continent of Garden (G) (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$).

Figure 4 The effects of the change in mean annual temperature (ΔMAT) on fitness of North American populations of *T. repens* grown in the Mississauga common garden. (**A**) Proportion of plants survived, (**B**) proportion of plants that produced flowers, (**C**) proportion of plants that produced seeds, (**D**) mean seed set mass, and (**E**) mean maximum plant area per population all exhibited evidence of an adaptation lag in response to climate warming, indicated by a positive linear trend or a shift in the peak of the quadratic curve towards populations with warmer climates of origin (denoted as x-axis values > 0). Shaded areas represent 95% confidence intervals around each line.