A transcontinental experiment elucidates (mal)adaptation of a cosmopolitan plant to climate in space and time

Lucas J. Albano[1,2], Cristina C. Bastias[3,4], Aurélien Estarague[4], Brandon T. Hendrickson[5], Simon G. Innes[5], Nevada King[5], Courtney M. Patterson[5], Amelia Tudoran[6], François Vasseur[4], *Adriana Puentes[6], *Cyrille Violle[4], *Nicholas J. Kooyers[5], *Marc T. J. Johnson[2]

[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

[2]Department of Biology, University of Toronto Mississauga, Mississauga, ON, Canada

[3]Ecology Area, Department of Botany, Ecology and Plant Physiology, Campus de Rabanales, Córdoba University, Córdoba, Andalusia, Spain

[4]CEFE, Université Montpellier, CNRS, EPHE, IRD, Montpellier, Occitanie, France

[5]Department of Biology, University of Louisiana at Lafayette, Lafayette, LA, USA

[6]Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Uppland, Sweden

*These authors co-supervised the project

Correspondence Author:

Lucas J. Albano

Current address: Department of Plant & Microbial Biology, North Carolina State University, Raleigh, NC, United States, 27695, lucas.albano@alumni.utoronto.ca

Open Research Statement:

Data are provided for peer review. All data involved in this study is available on the GitHub page

for LJA (https://github.com/ljalbano/transcontinental_common_garden).

Keywords:

Adaptation lag, climatic distance, common garden, cyanogenesis, environmental variation,

introduced range, non-native species, local adaptation, native range, provenance study, *Trifolium*

repens, white clover

Abstract (350 words)

Climate change and the global spread of non-native species are two of the most significant

threats to biodiversity and ecosystem function. Both these phenomena subject populations to

novel conditions, either in space (species introductions) or in time (climate change), yet the role

of adaptation in how populations respond to these rapid environmental shifts is poorly

understood. We conducted a large-scale trans-continental common garden experiment using

white clover (*Trifolium repens*, Fabaceae) to test whether adaptive evolution to spatiotemporal

variation in climate could contribute to the ecological success of one of the most widespread

plant species in the world. Individuals from 96 populations of *Trifolium repens* (white clover)

from both its native (Europe) and introduced (North America) ranges were planted into four

experimental common gardens located in northern (Uppsala, Sweden) and southern (Montpellier,

France) Europe, and northern (Mississauga, Canada) and southern (Lafayette, USA) North

America. We recorded plant sexual and clonal fitness in each common garden and assessed

whether the strength of local adaptation differed between the native and introduced ranges and

whether populations are rapidly adapting to climate change. Results show that local adaptation

was only evident when populations were transplanted into common gardens located in the same

range (native or introduced) from which they originated and was driven by stronger selection

(due to climatic factors rather than herbivory) at lower latitudes in both ranges. Our results

indicate rapid local adaptation across a large latitudinal gradient in introduced *T. repens*

populations, along with an associated adaptation cost when transplanted back into the native

range. We also find evidence of an adaptation lag in the northern common garden in the

introduced range, with plants from historically warmer climates exhibiting the greatest fitness.

These findings support two major conclusions: 1) white clover can rapidly adapt to spatial

variation in climate in its introduced range as well as the native range, and 2) despite rapid

adaptation to novel environments, introduced white clover populations are not keeping pace with

rapid climate change. Overall, our results provide insight into the role of adaptation in facilitating

the ecological success of non-native species in a rapidly changing world.

Introduction

Natural selection results in populations evolving traits that are advantageous to the

environmental conditions of a particular location, a phenomenon known as local adaptation

(Turesson, 1922; Kawecki and Ebert, 2004; Hereford, 2009). A fundamental aspect of local

adaptation is that variation in biotic and abiotic factors leads to changes in the relative fitness of

genotypes between environments (Gillespie and Turelli, 1989; Kawecki and Ebert, 2004; Lee

and Mitchell-Olds, 2011). Such genotype-by-environment interactions can be elucidated by

reciprocal transplant experiments, which are challenging to conduct, but vital in demonstrating

fitness trade-offs, whereby genotypes exhibit higher relative fitness in their local environment

and lower fitness in other environments (Gillespie and Turelli, 1989; Hereford, 2009; Leimu and

Fischer, 2008; Savolainen et al., 2013). Human activities are exposing populations to rapidly

changing novel environments, both across space through species introductions (Pyšek et al.,

2020), and over time through anthropogenically mediated climate change (Hoffmann and Sgrò,

2011). Understanding how populations simultaneously adapt to novel biotic and abiotic

environments over space and time is an important problem that is poorly resolved (Hendry,

2017; Siepielski et al., 2017). Addressing this problem is necessary to determine how

increasingly pervasive anthropogenic disturbances alter the ecology and evolution of native and

introduced species (Parmesan, 2006; Fugère and Hendry, 2018).

Introduced populations are frequently subjected to shifts in selection pressure compared

to the native environment to which they are locally adapted (Erfmeier, 2013; Colautti and Lau,

2015; Lau and terHorst, 2015). This shift in selection pressure in either the biotic environment

(e.g., novel species interactions) or the abiotic environment (e.g., climatic factors) could inhibit

the establishment and spread of an introduced population (Erfmeier, 2013; Colautti and Lau,

2015; Lau and terHorst, 2015). Additionally, inherent characteristics of introduced populations,

such as small population size and low genetic diversity, could limit their ability to respond to

novel selection pressures, with Allee effects and genetic bottlenecks potentially resulting in mate

limitation and inbreeding depression (Tsutsui et al., 2000; Kolar and Lodge, 2001; Allendorf and

Lundquist, 2003; Taylor and Hastings, 2005; Burns et al., 2011; Lau and terHorst, 2015; Pannell

et al., 2015; Estoup et al., 2016). In most cases, the ecological and/or genetic constraints

experienced by introduced populations are sufficient to result in the failure of those populations

to establish and their eventual extirpation (Kolar and Lodge, 2001; Jeschke and Pyšek, 2018).

Despite the challenges presented by introduction to novel environments across space,

introduced populations still often become invasive, experiencing substantial ecological

opportunity if constraints can be overcome. For example, ecological constraints can be overcome

if novel selection regimes in introduced environments are either benign or even favourable to a

non-native population (Erfmeier, 2013; Colautti and Lau, 2015). This is particularly true when an

introduced population has an advantage over its new natural enemies or competitors compared to

in its native range (Blossey and Nötzold, 1995; Callaway and Ridenour, 2004; Müller-Schärer et

al., 2004; Callaway et al., 2022). It is also possible that introduced populations do not exhibit

strong evidence of genetic constraints to begin with (i.e., the genetic paradox of invasions),

because reductions in genetic variation were not substantial enough to limit the ability of a

population to respond to novel selection pressures to begin with (Allendorf and Lundquist, 2003;

Maron et al., 2004; Dlugosch and Parker, 2008a; Dlugosch and Parker, 2008b; Colautti and

Barrett, 2013; Estoup et al., 2016; van Boheemen et al., 2017; Battlay et al., 2024). In fact, when

introduced populations experience strong selection and sufficient genetic variation in novel

environments, it could facilitate rapid adaptation to those novel environments, which could in

turn, contribute to the success of a non-native species (Maron et al., 2004; Dlugosch and Parker,

2008a; Colautti and Barrett, 2013; Colautti and Lau, 2015; Simón-Porcar et al., 2021; but see

Genton et al., 2005). However, the extent to which rapid local adaptation to novel biotic and

abiotic conditions can facilitate the success of a widespread non-native species across large

environmental gradients remains poorly understood.

While there are examples in which introduced populations have rapidly adapted to novel

environments across space, it is important to consider that these novel environments may also be

changing through time. Temporal shifts in a local environment can cause selection pressures to

fluctuate, which could result in traits that were previously adapted to that environment becoming

maladaptive (Wilczek et al., 2014; Aguirre-Liguori et al., 2019; Exposito-Alonso et al., 2019;

Anderson and Wadgymar, 2020; Bontrager et al., 2020; Capblancq et al., 2020). If the rate of

adaptation is exceeded by the rate of environmental change, this disruption of local adaptation is

likely to result in reduced fitness (i.e., maladaptation) of populations (St Clair and Howe, 2007;

Atkins and Travis, 2010; Wilczek et al., 2014; Aguirre-Liguori et al., 2019; Capblancq et al.,

2020). Recent global warming may result in populations becoming maladapted to their local

climate conditions, whereas populations from lower latitudes or elevations (i.e., warmer origins)

become better suited to environmental conditions at higher latitudes or elevations; a phenomenon

known as an adaptation lag (Wilczek et al., 2014; Kooyers et al., 2019; Anderson and

Wadgymar, 2020). Currently, the frequency and magnitude of adaptation lags are poorly

understood (Anderson and Song, 2020).

If non-native species rapidly adapt to novel environments across space, similar rapid

adaptation to climate change through time could facilitate their continued range expansion,

persistence, and pervasiveness (Byers, 2002; Maron et al., 2004; Colautti and Barrett, 2013;

Oduor et al., 2016). In fact, non-native species are often better at climate tracking and more

resistant to climate warming than native species, which could contribute to their success as

invaders (Wolkovich et al., 2013). However, the interplay between local adaptation, species

introductions, and anthropogenically mediated climate change remains an understudied, yet

critically important, aspect of understanding the ongoing human-induced biodiversity crisis

(IPBES, 2019). This problem is best addressed through large field experiments testing local

adaptation in both the native and introduced ranges of widespread non-native species, alongside

elucidating the environmental conditions that may result in rapid adaptation and success in a

novel environment (Colautti and Lau, 2015; Anderson and Song, 2020).

Plant-herbivore interactions have been highlighted as key selective factors that determine

patterns of local adaptation, species responses to environmental change, and biological invasions.

Spatial variation in both biotic and abiotic factors are likely to lead to spatial variation in plant

defense traits by affecting the costs and benefits involved in growth-defense trade-offs (Anstett

et al., 2018; Baskett and Schemske, 2018; Moreira et al., 2018). Local herbivore communities

impose selection through preferential feeding on plants with lower resistance, influencing the

benefits of plant defenses in areas of particularly high herbivore pressure(Ehrlich and Raven,

1964; Coley and Barone, 1996; Agrawal, 2007; Johnson and Rasmann, 2011; Züst et al., 2012).

In turn, the metabolic costs of defenses can be substantial, while abiotic environments and the

stressors they impose can influence the availability of resources to plants, exacerbating defense

costs (Coley et al., 1985; Herms and Mattson, 1996; Fine et al., 2006; Hahn et al., 2019).

Furthermore, it is increasingly recognized that anthropogenic activity can influence the evolution

of plant defenses (Bidart-Bouzat and Imeh-Nathaniel, 2008; Massad and Dyer, 2010; Miles et

al., 2019). Climate warming could result in increased herbivore feeding and/or increased plant

defenses (Dostálek et al., 2020; Lemoine et al., 2013) or could contribute to success of a non-

native species by decoupling locally adapted plant-herbivore interactions (Adams et al., 2009;

Costan et al., 2022; but see Agrawal and Kotanen, 2003). It is therefore important to understand

how populations are adapting to their local biotic and abiotic environments and how

anthropogenic factors can affect adaptation.

An ideal study system to test questions surrounding adaptation to novel environments

post-introduction and in response to climate change would be a widespread species with known

agents of biotic and abiotic selection across space and over time that can drive adaptation in the

native and introduced ranges. *Trifolium repens* is an allopolyploid perennial legume that is native

to Eurasia and now has a cosmopolitan distribution, following introduction to inhabited

continents throughout the world, including North America (Daday, 1958; Kjærgaard, 2003;

Griffiths et al., 2019). Mechanisms of biotic and abiotic selection in *T. repens* have been studied

for over 70 years. In *T. repens*, clines in a prominent chemical defense mechanism, hydrogen

cyanide (HCN), have been identified across many types of environmental gradients (Daday,

1954a; Daday, 1954b, Daday, 1958; Till, 1987; Kooyers and Olsen, 2013; Kooyers et al., 2014;

Thompson et al., 2016), providing opportunities to test hypotheses surrounding the involvement

of biotic factors in adaptation to environmental change.

To understand adaptation to environmental variation in a non-native species over space

and time, we conducted a trans-continental common garden field experiment across latitudinal

gradients in both the native (Europe) and introduced (North America) ranges of *T. repens*. We

first address four research questions focused on adaptation in *T. repens* to biotic and abiotic

environmental variation in space. (Q1) Are *T. repens* populations locally adapted to their climate

of origin and does the strength of local adaptation differ across space? This question is a

prerequisite condition before any comparisons can be made regarding local adaptation between

the native and introduced ranges. (Q2) Does *T. repens* exhibit stronger local adaptation in its

native range than in its introduced range?. (Q3) What are the roles of plant defense and herbivory

(i.e., biotic factors) in driving local adaptation across space? (Q4) What bioclimatic variables

(i.e., abiotic factors) have the strongest impact on fitness across space (latitudes and ranges) in *T.*

repens? We then focus on whether *T. repens* can adapt to climate change through time using a

space-for-time approach to analyzing a subset of the common garden experimental data (Pickett,

1989; Fournier-Level et al., 2011), asking: (Q5) Are there adaptation lags to climate change in *T.*

repens, such that populations originating from warmer and/or drier climates exhibit greater

fitness than local populations? This study is among the first to investigate the ability of an

introduced species to rapidly adapt to novel environments in space and time simultaneously. Our

results elucidate the potential for rapid adaptation to contribute to initial and continued

ecological success of non-native species in the face of ongoing environmental change.

Methods

Study system

Trifolium repens has been a model for the study of plant defense based on its polymorphism for

the production of the potent chemical defense hydrogen cyanide (HCN; Hughes, 1991). In fact,

the ability to produce HCN as an antiherbivore defense may have contributed to the range

expansion of *T. repens*, allowing it to thrive in human-disturbed habitats such as lawns, fields,

and roadsides in many regions throughout the world (Burdon, 1983; Kjærgaard, 2003; Olsen et

al., 2021). Cyanogenesis occurs through the hydrolyzation of cyanogenic glucosides by the

enzyme linamarase. In *T. repens*, the presence or absence of cyanogenic glucosides, linamarin

and lotaustralin, is controlled by three linked genes in their metabolic pathway (*CYP79D15*,

CYP736A187, and *UGT85K17*; referred to collectively as *Ac/ac*), while the presence or absence

of the enzyme linamarase is controlled by the gene *Li/li* (Olsen et al., 2007; Olsen and Small,

2018). Thus, four phenotypes exist for HCN and the two metabolic components required for its

production: one in which HCN is constitutively produced (AcLi) and three that are acyanogenic

and therefore lack the defense entirely (acLi, Acli, acli; Corkill, 1940; Atwood and Sullivan,

1943).

Cyanogenesis is widely considered an adaptive polymorphism because *T. repens* exhibits

clines in the presence of HCN across numerous types of environmental gradients (Daday, 1954a;

Daday, 1954b; Daday 1958; Brighton and Horne, 1977; Thompson et al., 2016; Innes et al.,

2022; Santangelo et al., 2022). In general, dominant *Ac* and *Li* alleles (and thus the ability to

produce HCN) are fixed in warmer climates, while recessive *ac* and *li* alleles are fixed in colder

climates (Daday, 1954a; Daday, 1954b; Daday, 1965). Latitudinal cyanogenesis clines have been

identified in the native European range of *T. repens* (Daday, 1954a), as well as its introduced

ranges, such as North America (Daday, 1965). However, clines within introduced ranges as

generally shallower, possibly indicating weaker local adaptation post-introduction, which

occurred less than 400 years ago in the case of North America (Daday, 1958; Kjærgaard, 2003;

Kooyers and Olsen, 2012; Innes et al., 2022). One hypothesis to explain cyanogenesis clines is a

gradient in herbivory, with greater diversity and abundance of herbivores in warmer climates

resulting in stronger selection favouring cyanogenic plants (Angseesing, 1974; Dirzo and Harper,

1982; Kakes, 1989). Alternatively, empirical studies across various environmental gradients have

identified temperature, soil moisture, and soil nutrients as abiotic factors that may impose

selection on cyanogenesis or its metabolic components to influence cline formation (Daday,

1954a; Daday, 1965; Gleadow and Woodrow, 2002; Kooyers et al., 2014; Kooyers et al., 2018).

Overall, despite 70 years of research on the drivers of cyanogenesis clines in *T. repens*, there

remains little consensus on how variation in environmental conditions influence selection on

cyanogenesis in *T. repens*, or if cyanogenesis is primarily responsible for rapid local adaptation

to novel environments (Daday, 1954a; Daday, 1965; Hughes, 1991; Kooyers and Olsen, 2013;

Kooyers et al., 2014; Wright et al., 2018; Wright et al., 2021; Albano and Johnson, 2023; Kuo et

al., 2024). Our study aims to resolve this uncertainty by directly testing the strength of local

adaptation and its biotic and abiotic drivers in both the native and introduced ranges of *T. repens*.

Sampled populations and common garden sites

We addressed our research questions using four common gardens distributed across northern

(Uppsala, Sweden) and southern (Montpellier, France) regions of the native European range of *T.*

repens, and northern (Mississauga, ON, Canada) and southern (Lafayette, LA, USA) regions of

the North American introduced range (**Figure 1**). Seeds were collected from 49 natural *T. repens*

populations scattered across the native European range spanning ~27° of latitude, and 47 natural

T. repens populations along a 2500 km north-south transect spanning ~21° of latitude in the

introduced North American range (**Figure 1**). Temperature and latitude are highly correlated in

eastern North America but less directly associated in Europe, with temperatures varying across

both north-south and east-west gradients, which led to the more diffuse sampling approach in

Europe and the transect sampling approach in North America. To minimize maternal and

epigenetic effects, we hand-pollinated plants germinated from field-collected seeds in a growth

chamber environment (25°C day, 15°C night, 14 h:10 h light:dark cycle; Conviron MTPS,

Winnipeg, MB, Canada) to create an outbred F1 cohort of seeds for each population. F1

generation seeds were germinated and established for four weeks in peat pots (8 cm diameter,

591 mL volume) in uniform growth chamber or greenhouse conditions near each common

garden site. Five replicate individuals from each population were transplanted in peat pots

directly into the natural soil of a typical cultivated lawn at each common garden site. Inconsistent

germination success resulted in some populations not being included in some common garden

locations, and in very limited cases, populations had four or six individuals planted. The

resulting total number of plants per garden was 465, 465, 458, and 470 in Mississauga, Lafayette,

Uppsala, and Montpellier, respectively. Plants were arranged in rows and columns with 1 m

spacing, and black landscaping fabric was used to cover the entirety of the lawn to prevent the

rooting of stolons into the natural soil between plants. Holes (12 cm diameter) were cut in the

fabric for each individual plant, leaving a 2 cm ring of natural vegetation around each plant to

allow for natural competition. This vegetation was trimmed monthly throughout the experiment.

Planting dates matched growing seasons in each location. The Lafayette and Mississauga

common gardens were established in April and May 2020, respectively, while the Montpellier

and Uppsala common gardens were postponed due to COVID-19 restrictions in Europe, leading

to their establishment in March and May 2021, respectively. Plants were given supplemental

water for the first two weeks of the experiment and intermittently after the first two weeks when

necessary to avoid excessive mortality. Plants that died within the first two weeks of the

experiment were deemed to have died due to transplantation stress and were replaced using extra

germinated individuals from the same population (Mississauga: 3; Lafayette: 32; Uppsala: 72;

Montpellier: 84). Plants that died after two weeks were deemed to have died due to the local

environmental conditions at each common garden site and were not replaced. Each common

garden was maintained for two full growing seasons.

Tissue collection, herbivory estimates, and measures of fitness

After 1-2 months of growth, one to three fully-expanded leaves were harvested from each

surviving individual and frozen at -20°C. These leaves were later freeze-dried and used for DNA

extraction, followed by low coverage whole genome sequencing to predict the presence of *Ac/ac*

and *Li/li* alleles (see the **Supplementary Methods** section of the **Supporting Information**). In

total, 586 individuals (Mississauga: 302, Lafayette: 99, Uppsala: 78, Montpellier: 107) were

successfully sequenced and characterized as possessing either dominant or recessive alleles at

Ac/ac and *Li/li.* The remaining 1272 individuals were not genotyped because they either died

before tissue could be harvested or genomic library preparations were unsuccessful. We exposed

plants to the natural environment of each common garden at the earliest life stage possible to

capture potentially important selection on seedlings (Agrawal et al., 2012). As a result, the

experiment experienced high mortality of young individuals, which is common and a natural part

of the biology of *T. repens* and other plants (Escudero et al., 2000; Moles and Westoby, 2004;

Agrawal et al., 2012). This mortality prevented many individuals from being included in tissue

collection. Visual herbivory estimates (% leaf area consumed) were conducted monthly

following the methods of Johnson et al. (2016). Herbivory was analyzed for the month exhibiting

peak herbivory (highest average % leaf area consumed) within each garden. Multiple

components of plant fitness were measured throughout the growing seasons at each location.

Survival and onset of flowering were assessed weekly during the first growing season. Mature

fruits were counted and collected weekly, and seeds were separated using a thresher (Precision

Machine Co., Lincoln, NE, USA) to obtain total seed set mass for each plant across both growing

seasons of the experiment. Photos were taken of each plant monthly during the first growing

season at each common garden location and assessed for plant area using Easy Leaf Area

(Version 2.0; Easlon and Bloom, 2014). The maximum plant area was then determined and the

growth rate across the first growing season was calculated as $(\ln[A_{max}] - \ln[A_1])/(t_{max} - t_1)$,

where A_{max} represents the maximum plant area, A_1 represents the initial plant area (i.e., after one

month of growth in the common garden), and $(t_{max} - t_1)$ represents the number of days between

photos taken for maximum plant area and initial plant area.

Calculating climatic distance

Six bioclimatic variables were extracted from WorldClim (Version 2.1, February 2023) at a

resolution of 30 seconds for each original *T. repens* population and common garden site

(O'Donnell and Ignizio, 2012). The variables included were mean annual temperature ("BIO1";

MAT), maximum temperature of the warmest month ("BIO5"), minimum temperature of the

coldest month ("BIO6"), annual precipitation ("BIO12"), precipitation of the wettest month

("BIO13"), and precipitation of the driest month ("BIO14"). The extracted values of each

bioclimatic variable for each common garden location can be found in **Table S1**. Values for all

six variables were z-score standardized to allow for analysis on an equivalent scale. A principal

component analysis was performed using the *princomp* function in R, with a scree plot to guide

the decision to retain principal components 1, 2, and 3, which explained 65.8%, 22.1%, and 9.1%

of the variation, respectively (cumulatively explaining 97% of variation; **Figure S1**).

Coordinates for each location of origin and each common garden location from all three

dimensions were then extracted using the *get_pca_ind* function in the *factoextra* package

(version 1.0.7; Kassambara and Mundt, 2020), and then plotted on a 3D graph (**Figure S2**). The

weighted Euclidean distance between points in 3D space was calculated using the formula:

$$Distance = \sqrt{\sum \left(w_i [O_{ij} - G_i]^2 \right)} \#(1)$$

Where w_i represents the proportion of variance explained by the ith dimension, O_{ij} represents the

value of the jth population of origin in the ith dimension, and G_i represents the value of a given

common garden in the ith dimension. We used this weighted Euclidean distance as a measure of

the climatic distance between the location of each population of origin and each common garden

location based on our six bioclimatic variables of interest (Wright et al., 2018; Samis et al.,

2019). This measure of climatic distance was then used as a predictor variable of plant fitness in

our statistical analyses. Bivariate correlations among individual climatic variables and with

climatic distance in each common garden are reported in **Table S2**. We also tested average

climatic distances among sampled populations within each continent transplanted into common

gardens on each continent using the following model (R function *lm*):

$$Climatic\ Distance \sim Continent\ of\ Origin + Continent\ of\ Garden + O:G \qquad (2)$$

Climatic distance varied based on continent of origin ($F_{1,1854} = 93.849$, $P < 0.001$), continent of

garden ($F_{1,1854} = 65.960$, $P < 0.001$), and their interaction ($F_{1,1854} = 142.491$, $P < 0.001$), with

European populations transplanted into European common gardens generally experiencing the

smallest climatic distances between environment of origin and the common garden environment.

Statistical analysis

All statistical procedures and analyses were conducted in R Version 4.1.2 (2021) and

RStudio Version 2022.12.0 (R Core Development Team, 2022). Seven response variables were

assessed as components of fitness: i) plant survival in the first growing season, ii)

presence/absence of flowers during either growing season, iii) presence/absence of seeds during

either growing season, iv) number of flower heads produced across both growing seasons, v)

seed set mass produced across both growing seasons, vi) maximum plant area in the first

growing season, and vii) growth rate in the first growing season. Herbivory during the first

growing season was also measured as an additional non-fitness response variable. The number of

flower heads and seed set mass were both zero-inflated, which led to the use of a two-step

modelling approach as described previously (Albano and Johnson, 2023). Briefly, both variables

were first analyzed in binomial models to assess whether plants produced flowers or seeds. Non-

zero flower head numbers and seed set masses were then subsequently analyzed as separate

fitness components. All linear mixed effects models were conducted using the *glmmTMB*

function in the *glmmTMB* package (version 1.1.5; Brooks et al., 2017). All test statistics and *P*

values were calculated using the *Anova* function in the *car* package (version 3.0-12; Fox and

Weisberg, 2019) with Type II sums-of-squares. We fit three binary fitness measures (i.e., plant

survival, presence/absence of flowers, presence/absence of seeds) to a binomial distribution. All

quantitative measures of fitness (i.e., number of flower heads, seed set mass, maximum plant

area, growth rate) and herbivory were modeled using a normal distribution, with all

transformations necessary to meet assumptions of normality and homogeneity of variance found

in **Table S3** and sample sizes for each response variable and each model found in **Table S4**.

Slope estimates and their standard errors were calculated using the *emtrends* function in the

emmeans package (version 1.10.4; Lenth, 2024).

Variation in the strength of local adaptation between common garden locations

To determine whether *T. repens* populations exhibit local adaptation and to assess the strength of

local adaptation at each garden location (Q1), all response variables were analyzed using the

following structure:

$$Response \sim Continent\ of\ Origin\ + Garden\ Location\ (L) + Climatic$$
$$Distance\ (D) + O:L + O:D + L:D + (1|Population) \#(3)$$

Continent of origin (North America or Europe), garden location (Mississauga, Lafayette,

Uppsala, or Montpellier), and climatic distance (the combined measure of six bioclimatic

variables of interest described above) were included as fixed effects, along with all two- and

three-way interactions. Population was included as a random effect to account for non-

independence of replicates within each population. The strength of local adaptation refers to the

magnitude of the fitness advantage for local populations compared to populations transplanted at

an increasingly greater climatic distance from their origin. We therefore interpret the slope of the

relationship between climatic distance and fitness as being representative of the strength of local

adaptation within a given common garden (a negative slope of greater magnitude represents

greater strength of local adaptation). An interaction between continent of origin and climatic

distance would indicate that the strength of local adaptation varies based on whether populations

originated in the native or introduced range. An interaction between garden location and climatic

distance would indicate that the strength of local adaptation varies based on the common

environment in which it is being tested.

Variation in the strength of local adaptation between ranges

To assess the strength of local adaptation between the native and introduced ranges of *T. repens*

(Q2), the analyses of the previous subsection were repeated, with the continent of the common

garden (North America or Europe) replacing the location of the common garden in the models:

$$Response \sim Continent\ of\ Origin + Continent\ of\ Garden\ + Climatic$$
$$Distance\ +\ O{:}G + O{:}D + D{:}G + O{:}D{:}G + (1|Population)\#(4)$$

385 Interpretation of this model is similar to the model above, with an interaction between continent

386 of origin and climatic distance indicating that the strength of local adaptation varies based on

387 whether populations originated in the native or introduced range. An interaction between

388 continent of garden and climatic distance would indicate that the strength of local adaptation

389 varies based on the continent of the common environment in which it is being tested. In this

390 model, we are particularly interested in a three-way interaction between continent of origin,

391 continent of garden, and climatic distance because it would indicate that the strength of local

392 adaptation varies based not only on the range from which populations originated but also the

393 range in which they are being tested. In other words, this model aims to test whether locally

394 adapted European populations maintain signatures of local adaptation in North America and vice

395 versa.

396   *The roles of cyanogenesis and herbivory in local adaptation*

397 The presence of cyanogenesis clines across latitude on both continents was determined

398 separately for *Ac/ac* and *Li/li* by assessing the proportion of plants in each population containing

399 at least one dominant *Ac* or *Li* allele using the following model structure fit to a binomial

400 distribution:

$$Proportion\ \text{Ac}\ or\ \text{Li}\ Plants \sim Latitude + Continent\ of\ Origin$$
$$+ Latitude{:}Continent\ of\ Origin + (1|Population)\#(5)$$

401 An interaction between latitude and continent of origin would indicate a difference in the slope

402 of clines in the presence of *Ac* and/or *Li* based on the range in which those clines are occurring

403 (native European or introduced North American).

404   The effect of cyanotype (AcLi, Acli, acLi, or acli) on local adaptation (Q3) was assessed

405   for all eight response variables using the following model structure:

$$Response \sim Cyanotype\ (C) + Garden\ Location\ (L) + Climatic\ Distance\ (D) \\ + C:L + C:D + C:L:D + (1|Population)\#(6)$$

An interaction between cyanotype and climatic distance would indicate that cyanogenesis or its

metabolic components contribute to the strength of local adaptation in *T. repens*. A three-way

interaction between cyanotype, garden location, and climatic distance would indicate that

selection for or against a given cyanotype in a particular common environment impacts the

strength of local adaptation in that environment. This might be expected due to consistently

observed latitudinal cyanogenesis clines across multiple types of environmental gradients in *T.*

repens, which indicate selective advantages for particular cyanotypes depending on the location

in which they are found, and the biotic and abiotic factors imposing selection in that location.

Bioclimatic variables as individual drivers of local adaptation

We also measured associations between the seven response variables and each of the six

bioclimatic variables of interest individually by determining the difference in z-score

standardized values for each variable between the location of each population of origin and each

common garden location using the following model structure:

$$Response \sim Bioclimatic\ Variable + \ Bioclimatic\ Variable^2 + (1|Population)\#(7)$$

For this set of analyses, all common gardens were analyzed separately, and all quantitative (non-

binomial) response variables were z-score standardized to allow for simultaneous direct

comparisons of the relative strength of linear and quadratic predictor variables. Within each

garden, we were therefore able to directly compare *P* values across all bioclimatic variables for

both linear and quadratic fits (**Tables S5-S8**). For each response variable, we selected the most

influential bioclimatic variable (i.e., the lowest *P*, but only when at least one *P* value was

significant at a level of $P < 0.1$, highlighted in red in **Tables S5-S8**). Across all response

variables in each garden, we then calculated the percentage of these most influential bioclimatic

variables that were either temperature-based (BIO1, BIO5, or BIO6) or precipitation-based

(BIO12, BIO13, and BIO14) and compared these percentages across the entire experiment,

across latitudes, and across ranges (Q4). This approach was used to identify the most important

abiotic variables driving local adaptation in general, as opposed to focusing on the results of any

one particular test.

Adaptation lags in response to climate warming

We sought to investigate the presence of adaptation lags using a space-for-time substitution (Q5).

We performed an additional analysis using a similar model structure as before, but only using

changes in raw values for MAT (i.e., not z-score standardized, hereafter referred to as ΔMAT) as

the bioclimatic variable of interest.

$$Response \sim \Delta MAT + \Delta MAT^2 + (1|Population) \#(8)$$

If the peak of the quadratic relationship in each model is approximately equal to zero, it would be

considered evidence of local adaptation to MAT, because maximum fitness occurs in populations

with the same mean annual temperature as the common garden site. If the peak of the quadratic

relationship is instead shifted towards populations of warmer origins than are occurring at the

common garden site (i.e., these warmer-adapted populations are outperforming the local

populations), this would be considered evidence of an adaptation lag. We tested whether the shift

of peaks away from a ΔMAT of zero and towards populations of warmer origins were significant

by also fitting a similar model to above, but excluding the linear predictor:

$$Response \sim \Delta MAT^2 + (1|Population) \#(9)$$

This additional model constrains the peak of the distribution to be centred around a ΔMAT of

zero (approximating the outcome of local adaptation as opposed to an adaptation lag). We then

compared the models with and without the linear predictor using the *anova* and *AIC* functions in

R, with *P* values < 0.05 (along with ΔAIC values < 0) indicating a significant shift of the peak.

We focused our interpretation of this analysis based solely on the North American

populations in the Mississauga common garden because the sampling design of North American

populations along a transect and the location of the Mississauga garden within that sampled

transect provide the most appropriate subset of the data to detect spatial lags. We include the

results for European populations within the Uppsala common garden in **Table S9** for

completeness, but the Uppsala common garden did not exhibit any trend in fitness based on

climatic distance, precluding the possibility of an adaptation lag being detected. The Lafayette

and Montpellier common gardens were excluded from this analysis because they are near the

southern extremes of our sampled populations, precluding the ability to detect adaptation lags.

Results

Q1: Are T. repens populations locally adapted to their climate of origin and does the strength of

local adaptation differ across space?

We found strong evidence of local adaptation, and the strength of local adaptation in *T. repens*

varied among gardens for multiple fitness components. Climatic distance between each

population of origin and the common garden locations strongly predicted all sexual fitness

components and survival (**Table 1**). In general, populations tended to have higher fitness with

lower climatic distance, indicating that populations of more similar climatic origins to the

climate of each common garden were better adapted. The only response variables not showing

evidence of local adaptation were the measures of vegetative growth used as proxies for clonal

fitness (i.e., maximum plant area and growth rate). The interaction of common garden location

and climatic distance affected the number of flower heads and seed set mass, but not other fitness

components or herbivory (**Table 1**; **Figure S3**). In general, the strength of local adaptation,

measured as the slope of the relationship between sexual fitness and climatic distance, was

greater in the more southern common garden location on each continent (**Figure 2**). For the

number of flower heads, the slope for Lafayette was 3.4× steeper than the slope for Mississauga

and the slope for Montpellier was 349× steeper than the slope for Uppsala. Similarly, for seed set

mass, the slope for Lafayette was 2.3× steeper than the slope for Mississauga and the slope for

Montpellier was 384× steeper than the slope for Uppsala. The main effect of garden location

affected all fitness components and herbivory, which reflects the large differences in

environment between the gardens. By contrast, the main effect of continent of origin and its

interactions with garden location and/or climatic distance did not affect any fitness component or

herbivory (**Table 1**).

Q2: Does T. repens exhibit stronger local adaptation in its native range than in its introduced

range?

We observed strong local adaptation in both the native and introduced ranges of *T. repens*, but

the presence of local adaptation was dependent on where populations originated. This result is

explained by a three-way interaction between continent of origin, continent of the common

garden, and climatic distance, which affected all quantitative sexual fitness components,

survival, and growth rate of *T. repens* (**Table 2**). Specifically, populations only exhibited strong

signatures of local adaptation when they were transplanted to common gardens on the same

continent from which they originated (**Figure 3**). For number of flower heads, in common

gardens located in the native range (Europe), the slope for European *T. repens* populations was

12× steeper than the slope for North American populations. In common gardens located in the

introduced range (North America), the slope for North American *T. repens* populations was 20×

steeper than the slope for European populations. Similarly, for seed set mass, in common gardens

located in Europe, the slope for European *T. repens* populations was 24× steeper than the slope

for North American populations. In common gardens located in North America, the slope for

North American *T. repens* populations was 7.8× steeper than the slope for European populations.

The main effect of climatic distance affected whether plants flowered and produced seeds, but

climatic distance did not interact with the continent of origin or the continent of the common

garden to affect these fitness components (**Table 2**; **Figure S4**). However, climatic distance and

continent of origin did interact to affect maximum plant area and herbivory. Together across all

gardens, plants from European origins that were transplanted shorter climatic distances grew

larger, despite experiencing greater levels of herbivory, while size and herbivory for plants from

North American origins were not significantly affected by climatic distance overall (**Table 2**;

Figure S4).

Q3: What are the roles of cyanogenesis and herbivory in driving local adaptation across space?

Both *Ac/ac* and *Li/li* exhibited strong latitudinal gene frequency clines among *T. repens*

populations from both continents (**Table S8**). However, the lack of an interaction between

latitude and continent of origin for both loci indicates no differences in the steepness of

cyanogenesis clines between the native and introduced ranges of *T. repens* (**Figure S5**). The

main effect of cyanotype affected survival, whether plants produced flowers and/or seeds, and

maximum plant area, with cyanogenic plants generally exhibiting greater fitness than

acyanogenic plants, regardless of the environment in which the common garden was found

(**Table S9**). However, cyanotype alone did not affect the level of herbivory occurring on *T.*

repens plants, and cyanotype did not interact with either common garden location or climatic

distance to affect any fitness components or herbivory (**Table S9**).

Q4: What bioclimatic variables have the strongest impact on fitness across space in T. repens?

Climatic variables related to both temperature and precipitation predicted variation in fitness in

T. repens, but the most influential predictors varied between garden locations, with patterns

emerging based on latitude (**Tables S3-S6**, red highlights). Overall, 58% of the most influential

predictors were temperature-based (mean annual temperature, maximum temperature of the

warmest month, or minimum temperature of the coldest month), while 42% were precipitation-

based (annual precipitation, precipitation of the wettest month, or precipitation of the driest

month). In the more southern gardens on each continent (Lafayette and Montpellier), 67% of the

most influential predictors were precipitation-based, while only 33% were temperature-based.

Conversely, in the more northern gardens on each continent (Mississauga and Uppsala), 80% of

the most influential predictors were temperature-based, while only 20% were precipitation-

based. We did not find evidence that the native and introduced ranges of *T. repens* were

differentially affected by temperature or precipitation variables. In the native range (Uppsala and

Montpellier, combined), 57% of the most influential predictors were temperature-based and 43%

were precipitation-based. In the introduced range (Mississauga and Lafayette, combined), 58%

of the most influential predictors were temperature-based, while 42% were precipitation-based.

Q5: Are there adaptation lags to climate change in T. repens?

North American populations of *T. repens* demonstrated evidence of an adaptation lag in the

northern garden in the introduced range (**Table 3**). Survival, seed set mass, and maximum plant

area all exhibited a significant quadratic relationship with ΔMAT between locations of origin and

the Mississauga common garden (**Figure 4**). A quadratic relationship centred on a ΔMAT of

zero would indicate local adaptation to the climate surrounding the common garden, while a

relationship centred on a ΔMAT > 0 would represent an adaptation lag. For the proportion of

plants survived per population, the peak of the quadratic relationship experienced a significant

shift to a climate of origin with a MAT 3.4°C greater than Mississauga ($\chi^2_{1,4}$ = 4.159, P = 0.041,

ΔAIC = −2.159), with the predicted maximum survival being 5.3% greater than at a ΔMAT of

zero. For mean seed set mass per population, the peak of the quadratic relationship occurred at a

MAT of 2.0°C greater than Mississauga, with the predicted maximum seed set mass being 5.2%

greater than at a ΔMAT of zero. However, this shift was not statistically significant ($\chi^2_{1,5}$ =

0.640, P = 0.423, ΔAIC = 1.360). For mean maximum plant area per population, the peak of the

quadratic relationship experienced a shift to a climate of origin with a MAT of 2.1°C ($\chi^2_{1,5}$ =

3.721, P = 0.054, ΔAIC = −1.721) greater than Mississauga, with the predicted maximum plant

area being 3.9% greater than at a ΔMAT of zero. Along our transect of sampled populations, a

shift in the peak of the quadratic relationship towards populations 2.0-3.1°C warmer than the

Mississauga common garden represents a 2.1-3.2° shift in latitude towards the equator.

Additionally, whether plants flowered or produced seeds increased linearly with ΔMAT,

indicating plants of warmer origins were more likely to flower and produce seeds (**Figure 4**),

while the number of flower heads and growth rate did not exhibit any significant linear or

quadratic trends with ΔMAT (**Table 3**).

Discussion

The aim of our study was to provide insight into the potential for rapid local adaptation in

introduced plant populations in response to novel environments across space and time. The

results of our transcontinental experiment in the native (Europe) and introduced (North America)

ranges of *T. repens* lead to two main conclusions: 1) rapid local adaptation to climatic variation

across space has contributed to the ecological success of *T. repens* in its introduced range; 2)

climate warming has resulted in some evidence of a weakening of local adaptation over time in

the introduced range of *T. repens.* Our conclusions are based on five experimental results. First,

T. repens exhibited local adaptation to climate overall, with greater fitness advantages for local

populations at lower latitudes (Q1). Second, in European common gardens, only native European

populations exhibited local adaptation, while in North American common gardens, only

introduced North American populations exhibited local adaptation (Q2). Third, cyanogenesis did

not differentially affect the fitness of *T. repens* in different environments (Q3). Fourth, both

temperature- and precipitation-based bioclimatic variables predicted fitness, but temperature was

a stronger determinant of fitness at higher latitudes, while precipitation was a stronger

determinant of fitness at lower latitudes (Q4). Fifth, there is some evidence consistent with an

adaptation lag in response to climate warming, but the signature was weak, and this pattern was

only clear in the northern common garden of the introduced range (Q5). Below we discuss our

results in the context of the role of rapid local adaptation in the success of non-native species and

how introduced populations may (or may not) be able to adapt to future climate change.

Strength of local adaptation is dependent on latitude

Spatial variation in biotic and abiotic factors across a species' range could affect its ability to

adapt to different environments within that range. We found that *T. repens* exhibited significant

local adaptation to the abiotic environment based on a measure of climatic distance that

encompassed mean annual temperature and precipitation along with annual temperature and

precipitation extremes. However, local adaptation differed significantly in strength depending on

the latitude of common gardens. Specifically, local adaptation was stronger at lower latitudes in

both the native and introduced ranges of *T. repens* when fitness was measured as sexual

reproductive success (i.e., number of flower heads and seed set mass). Our results are consistent

with transplant experiments in other plant species, which found sexual fitness advantages for

local versus foreign plants to be more pronounced at lower latitudes (Colautti and Barrett, 2013;

Toräng et al., 2015), likely at least partially due to life history trade-offs across large latitudinal

ranges (Li et al., 2015; Bastias et al., 2024). In fact, life history trade-offs in *T. repens* contribute

to local adaptation across divergent environments, with southern populations investing more into

sexual reproduction and northern populations investing more into vegetative growth (Wright et

al., 2021). If warmer climates are generally more conducive to sexual reproduction than cooler

climates, this may explain why sexual fitness advantages for local populations were more

pronounced in Lafayette and Montpellier in our study, while climatic distance had more limited

effects on vegetative growth.

Local adaptation in native and introduced ranges

Non-native species could become more widespread and persistent by rapidly adapting to novel

environments. For example, latitudinal cyanogenesis clines in introduced North American *T.*

repens populations formed due to rapid local adaptation and sorting of existing genotypes post-

introduction (Kooyers and Olsen, 2012; Kooyers and Olsen, 2013; Olsen et al., 2021). We found

that populations of *T. repens* only exhibited strong signatures of local adaptation to climate when

transplanted into common gardens within the same range as they originated. The observed strong

local fitness advantages for North American populations within the North American gardens are

therefore indicative of rapid local adaptation of *T. repens* within the past 400 years since its

introduction. Strong selection on introduced populations provides opportunity for rapid local

adaptation in non-native species on even shorter timescales than exhibited by *T. repens* (Colautti

and Lau, 2015; Simón-Porcar et al., 2021). A meta-analysis by Oduor et al. (2016) found that

non-native populations exhibited local adaptation just as frequently as native populations, and

that the strength of local adaptation was not affected by time since introduction (ranging from as

little as 20 to as many as 500 years). More specifically, previous studies on invasion in

Hypericum canariense (Dlugosch and Parker, 2008), *Eschcholzia californica* (Leger and Rice,

2007), and *Lythrum salicaria* (Colautti and Barrett, 2013) indicate local adaptation in introduced

ranges within 50, 150, and 200 years, respectively. Taken together, our results strengthen the

mounting evidence that rapid local adaptation to climate plays an important role in the ecological

success of non-native species in their new range.

616   Interestingly, *T. repens* populations from the introduced range did not exhibit signatures

of local adaptation to similar climates when transplanted back into their native range. Similarly,

native populations did not maintain their strong signatures of local adaptation to climate when

transplanted into the introduced range. One important caveat is the overall narrower range of

climatic distances among European populations (particularly within European common gardens),

despite our widespread sampling (Figure S2), which could have limited our ability to compare

the strength of local adaptation between ranges (Lucas et al., 2024). Additionally, while our

common gardens are robust to detecting local adaptation caused by genotype × environment

interactions (Bock et al., 2018), we also acknowledge that phenotypic and/or transgenerational

plasticity (whose determination was beyond the scope of this study) may also be important for

successful establishment or spread of non-native species (Hulme, 2008; Valladares et al., 2014;

de Villemereuil et al., 2018). Despite these considerations, our findings demonstrate a clear cost

of rapid adaptation, with local fitness advantages not extending across continents, even when

accounting for a comprehensive measure of climatic distance between source environments and

common garden environments on both continents. Explanations for this lack of maintenance of

local adaptation across continents could be due to unmeasured biotic or abiotic aspects of the

environment.

Abiotic explanations for the lack of local adaptation across continents could involve

shifts in photoperiod or precipitation regimes between Europe and North America due to similar

climates on the two continents being associated with different latitudes. In general, European

environments have longer day lengths in summer and shorter day lengths in winter than those of

similar climates in North America, with differences being more pronounced at progressively

higher latitudes. This photoperiodic shift between the continents could have affected the

initiation of flowering and/or the timing of peak flowering for populations that were transplanted

across continents, leading to differences in the strength of local adaptation. Halbritter et al.

(2015) found asymmetries in the strength of local adaptation in reciprocally-transplanted high-

elevation and high-latitude *Plantago lanceolata* and *Plantago major* populations with similar

climates of origin, indicating a potential role of photoperiod in local adaptation. Additionally,

populations of an obligate long-day plant, *Mimulus guttatus*, exhibit clinal variation in the

critical photoperiod required to initiate flowering, potentially indicating a role of photoperiod in

driving local adaptation (Kooyers et al., 2015). Saikkonen et al. (2012) speculated that the

transition to less extreme yearly variation in photoperiod at lower latitudes might facilitate

greater ecological success of European species introduced to North America than vice versa,

which could be especially true of a facultative long-day plant like *T. repens* (Beatty, 1956).

However, our results do not provide evidence of increased success of European populations in

North American gardens compared to that of North American populations in European gardens,

indicating that other environmental factors are likely at play. Additionally, although we

accounted for mean annual precipitation and yearly precipitation extremes in our study, these

variables do not fully capture the exact timing and volume of precipitation, and the resulting

hydrological environment experienced by *T. repens* plants in Europe versus North America. For

example, in the Montpellier garden in southern Europe, plants will experience a Mediterranean

climate that is not only drier but also more seasonally variable, while in the Lafayette garden in

southern North America, plants will experience a sub-tropical environment that is wetter with

more limited seasonal variation. This shift in precipitation availability has caused introduced *T.*

repens populations to evolve greater drought avoidance as opposed to the more desiccation

tolerant strategy observed in Mediterranean populations (Hendrickson et al., 2024).

While our experiment offers strong tests of local adaptation to both biotic and abiotic

factors in the native and introduced ranges of a cosmopolitan plant, one limitation of this study is

an inability to determine the relative strength of soil conditions compared to climate as drivers of

local adaptation. Planting *T. repens* individuals from each population along with the natural soil

and microbial communities from their location of origin was infeasible considering our study

design and governmental regulations on importation of soil and bacteria. Instead, the best

alternative was to maintain consistent treatment of all individuals in all gardens, with *T. repens*

individuals planted into the natural soil at each common garden site. It is important to recognize

that local edaphic factors have the potential to provide fitness advantages in *T. repens*, as has

been demonstrated in other systems (Macel et al., 2007; Murray-Stoker and Johnson, 2024).

Despite this, signatures of local adaptation to climate were still evident across multiple fitness

components, strengthening our confidence in the observed patterns of rapid adaptation across

ranges. However, a future study exploring the roles of edaphic and other abiotic factors could

provide more comprehensive insight into how introduced plant populations adapt to new ranges.

Cyanogenesis did not affect local adaptation in either the native or introduced range

The biotic environment could also partially explain the lack of local adaptation across continents

based on climatic distance (Hargreaves et al, 2020). In line with classic hypotheses in invasion

biology, herbivory could exhibit substantial differences across environments based on expected

shifts in herbivore communities between ranges, which could facilitate success of a plant species

post-introduction (Coley et al., 1985; Blossey and Nötzold, 1995; Keane and Crawley, 2002). If

herbivory was likely to explain our results, we might expect to see signatures of local adaptation

based on the presence of chemical defenses in *T. repens* plants. Among our sampled populations,

we demonstrated rapid evolution of latitudinal clines in the presence of both *Ac* and *Li* alleles in

the introduced range of *T. repens*, based on their similar strength to clines in the native range,

demonstrating potential for involvement of cyanogenesis in local adaptation. However, our

experimental results indicate that HCN did not confer higher fitness to local plants compared to

foreign plants in any garden. In fact, local plants exhibited signatures of local adaptation despite

experiencing similar or greater levels of herbivory than populations transplanted from further

climatic distances. Additionally, experimental plants in the introduced range exhibited greater

herbivory than plants in the native range, which is contrary to the prediction that *T. repens* was

released from natural enemies in its native range (Keane and Crawley, 2002). Previous studies on

local adaptation in *T. repens* have similarly found that HCN only had marginal, if any, effects on

fitness (Wright et al., 2018; Wright et al., 2021) or have found weaker clines in the introduced

range than the native range (Daday, 1954a; Daday, 1958; Innes et al., 2022). Taken together, our

study contrasts classic invasion hypotheses, providing little evidence of a role of biotic factors

(cyanogenesis or herbivory) in driving local adaptation of *T. repens* to its introduced range.

Our findings contrast with previous work that has speculated on the involvement of trade-

offs in the costs and benefits of cyanogenesis or its metabolic components in adaptation to

contrasting environments. For example, exacerbated ecological or allocation costs of producing

HCN or its metabolic components in cooler climates could pair with exacerbated benefits of

cyanogenesis in warmer climates with greater herbivore pressure (Daday, 1954a; Daday, 1965;

Kooyers et al., 2018; Fadoul et al., 2023). In our study, cyanogenic plants often demonstrated a

fitness advantage (i.e., benefits outweighing costs), even in cooler environments like

Mississauga. However, despite these fitness advantages, we did not find evidence for HCN as a

deterrent to herbivory in *T. repens*, indicating that greater prevalence of cyanogenesis in warmer

climates is likely not based on an increased ability to defend against greater herbivore pressure.

Previous work instead indicates the potential for cyanogenic glucosides and/or linamarase,

individually, to be advantageous in periods of drought through the storage and recycling of

nitrogen (Møller, 2010; Kooyers et al., 2014; Machingura et al., 2016). However, we also do not

find evidence of the metabolic components of cyanogenesis affecting the fitness of *T. repens* or

its adaptation to local climates. It is important to highlight that we were unable to harvest enough

tissue to determine the cyanotype of most experimental plants due to concerns over effects of

tissue harvest on plant fitness and/or mortality, so the strength of our conclusions with respect to

the role of cyanogenesis in adaptation are limited by low sample sizes. Additionally, we do not

account for other biotic interactions, such as belowground herbivory, soil biota, or competition,

for their potential involvement in selection and local adaptation in *T. repens* (Blossey and

Nötzold, 1995; Mitchell et al., 2006; Pal et al., 2020; Sheng et al., 2022; Murray-Stoker and

Johnson, 2024). Further field studies are needed to more explicitly test a diverse array of biotic

factors as drivers of local adaptation to climate across large environmental gradients, including

direct tests of the costs and benefits of the most prominent defense mechanism in *T. repens*.

Spatial variation in environmental drivers of local adaptation

While *T. repens* exhibits local adaptation in its native and introduced ranges based on an overall

measure of climatic distance, populations may spatially vary in the individual bioclimatic

variables responsible for local adaptation. Our results across both ranges demonstrate that at

higher latitudes, where local adaptation is weaker, temperature-based variables are stronger

determinants of fitness, while at lower latitudes, where local adaptation is stronger, precipitation-

based variables are stronger determinants of fitness. Therefore, it is possible that *T. repens* adapts

more readily to local precipitation regimes than local temperature regimes, resulting in stronger

signatures of local adaptation in lower latitude common gardens, consistent between the native

and introduced ranges. This result is consistent with Siepielski et al. (2017), which found both

precipitation and potential evapotranspiration to be stronger predictors of selection than

temperature across numerous taxa. Alternatively, our results could be affected by the specific

environments chosen for the common gardens. In our study, the climates in Lafayette and

Montpellier are among the warmest that occur in the range of *T. repens*, while representing two

opposing extremes in terms of precipitation regime, and therefore could have been especially

difficult to tolerate for populations transplanted from cooler climates with less extreme

precipitation regimes. In turn, Uppsala was somewhat isolated, both geographically and

climatically, from the sampled *T. repens* populations, which could have affected our ability to

detect local adaptation (or maladaptation) in that environment. Lastly, it is possible that recent

climate warming may have already disrupted the ability of local populations to outperform

foreign populations, thereby impacting the strength of local adaptation in certain locations.

Evidence of adaptation lags in response to climate warming

The ability of populations to adapt to recent rapid climate warming over time is an important

mechanism by which species can mitigate the risk of local extinction and other downstream

ecological consequences. Previous studies have demonstrated the incidence of adaptation lags,

whereby populations originating from warmer climates would outperform populations with very

similar climatic origins to the common garden location (Wilczek et al., 2014; Aguirre-Liguori et

al., 2019; Anderson and Wadgymar, 2020). These studies indicate the potential for a disruption

in local adaptation, and even the possibility of maladaptation, due to climate warming that is

occurring faster than local populations are able to adapt.

In our northern common garden in North America (Mississauga), populations with

warmer origins (2-3°C in MAT) outperformed local populations in survival and biomass

production (i.e., a proxy for vegetative reproduction) but not sexual fitness components. This

finding may signal the presence of an adaptation lag and may also provide a partial explanation

for the observed weaker local adaptation in Mississauga compared to Lafayette. However, the

generalizability of our results should be considered carefully for a few reasons. First, contrary to

our expectations, we provide no evidence of either local adaptation or maladaptation in Uppsala.

In a large-scale common garden experiment with European populations of *Arabidopsis thaliana*,

Wilczek et al. (2014) found that adaptation lags were more pronounced in higher latitude

common garden locations, which was not the case in our study. It would also be pertinent to be

able to test for spatial lags in the native range of *T. repens* to which we could directly compare

spatial lags in the introduced range. Second, we tested for the presence of adaptation lags based

solely on MAT, and Mississauga was the common garden location where MAT most

consistently affected plant fitness compared to other bioclimatic variables. Fitness in lower

latitude common gardens was instead more heavily influenced by precipitation-based variables,

which may not be changing as consistently as temperature-based variables due to anthropogenic

activity but could still play a significant role in maladaptation in response to changing

environments (Nagy et al., 2024). It is therefore possible that adaptation lags to changing

environments may not occur consistently across different geographic locations, based on the

specific biotic and abiotic factors that most strongly impact local adaptation in those

environments. Third, the MAT at the location of the Mississauga common garden in 2020 and

2021 was 1.6°C and 2.0°C warmer, respectively, than the average from the historical time period

(1970-2000) used to calculate climatic distances in this study (data extracted from ClimateNA;

Wang et al., 2016). Therefore, it may not be unexpected to see populations from warmer origins

outperform local populations. The adaptation lag for survival does extend to a peak 3.4°C

warmer than Mississauga, and the proportion of plants producing flowers and/or seeds increased

linearly with ΔMAT, which still provides evidence for warm-adapted populations having an

advantage that exceeds expectations based on the MAT during the years of the experiment.

However, across our entire study, we are unable to make broad conclusions about future

vulnerability of populations based on the rate of environmental change exceeding the rate of

adaptation. Instead, we conclude that the fine-tuned nature of local adaptation (in terms of

variation in strength, prominent driving factors, and the potential for maladaptation) to climate in

T. repens highlights the multifaceted nature of rapid evolution in response to changing

environments.

Conclusions

Our study provides insight into the ability of introduced populations of a widespread plant to

adapt to novel environments across space and over time. Signatures of local adaptation were only

evident when populations were transplanted within the same range as they originated, indicating

both strong and rapid local adaptation to the introduced range of *T. repens* in less than 400 years

since introduction. Interestingly, the strength of local adaptation and the most important

environmental predictors of plant fitness differed in the north and south of both the native and

introduced range. Specifically, temperature was the most important driver of local adaptation in

northern gardens, whereas precipitation was most important in southern gardens. This result

indicates that there may be predictable drivers of local adaptation of non-native populations to

novel environments, and that these drivers may vary along broad latitudinal gradients. Despite

evidence of rapid adaptation to changing environments across space, the northern garden in the

introduced range also provided some evidence of local maladaptation compared to populations

with warmer origins, which is characteristic of an adaptation lag. Therefore, while local

adaptation has occurred quickly in introduced populations of *T. repens* across space, a lack of

rapid adaptation to environmental change over time could eventually result in severe erosion of

fitness advantages for local populations in the future. Continuing to further our understanding of

the ability of native and introduced populations to rapidly adapt to anthropogenic environmental

change in space and time is vital to the integrity and maintenance of biodiversity in the face of

future shifts in environment.

Acknowledgements

We thank James Santangelo for assistance with initial seed collection, and Happy Gill,

Jashandeep Nijjar, and Ruth Thomas for assisting with the generation of F1 plants for use in the

experiment. We are indebted to Aaron Albano, Barry Albano, Paula Albano, Sarah Albano,

Andre Daugeureaux, Hayden Fargo, Axel Hjelm, Antoniette Jansen, Eunice Karinho Betancourt,

Erik Karlsson, Andreas Lundgren, Kavya Manikonda, Lindsay Miles, David Murray-Stoker,

Bernabé Ramirez, Brittany Sutherland, Olivia Toth, Andrea Turcu, Mariona Vinyeta Medina,

and Vicki Zhang for assistance in the field. Keerath Bhachu, Elana Maria, Zain Nasrullah, Nial

Navaratne, Narindra Persaud, and Isabellio Vessio assisted with sample processing. We also

thank the Louisiana Optical Network (LONI) for providing computation resources, along with

Kris Horvath, Joan Lee, Kent Moore, Diane Ross, Vera Velasco, the technical platform

"Plateforme des Terrains d'Expérience" (CEFE-CNRS), and the University of Louisiana

Ecology Center for facilitating and supporting the set-up and execution of the experiment. This

research was supported by an NSERC Discovery Grant, Canada Research Chair, and NSERC

EWR Steacie Fellowship to MTJJ, a CNRS-University of Toronto Ph.D. Student Travel Grant to

MTJJ and CV, NSF grants OIA-1920858 and DBI-2244712 to NJK, and an NSERC CGS

Doctoral Award to LJA.

Conflict of Interest Statement

The authors declare no conflict of interest.

References

Adams, J. M., Fang, W., Callaway, R. M., Cipollini, D., Newell, E., Transatlantic *Acer*

platanoides Invasion Network (TRAIN). (2009). A cross-continental test of the Enemy

Release Hypothesis: leaf herbivory on *Acer platanoides* (L.) is three times lower in North

America than in its native Europe. *Biological Invasions*, *11*, 1005-1016.

Agrawal, A. A., Kotanen, P. M. (2003). Herbivores and the success of exotic plants: a

phylogenetically controlled experiment. *Ecology Letters*, *6*, 712-715.

Agrawal, A. A. (2007). Macroevolution of plant defense strategies. *Trends in Ecology and*

Evolution, *22*, 103-109.

Agrawal, A. A., Hastings, A. P., Johnson, M. T. J., Maron, J. L., Salminen, J.-P. (2012). Insect

herbivores drive real-time ecological and evolutionary change in plant populations.

Science, *338*, 113-116.

Aguirre-Liguori, J. A., Ramírez-Barahona, S., Tiffin, P., Eguiarte, L. E. (2019). Climate change

is predicted to disrupt patterns of local adaptation in wild and cultivated maize.

Proceedings of the Royal Society B, *286*, 20190486.

Albano, L. J., Johnson, M. T. J. (2023). Interactions environmental factors drive selection on

cyanogenesis in *Trifolium repens*. *Oikos*, *2023*, e09629.

Allendorf, F. W., Lundquist, L. L. (2003). Introduction: population biology, evolution, and

control of invasive species. *Conservation Biology*, *17*, 24-30.

Anderson, J. T., Song, B.-H. (2020). Plant adaptation to climate change – Where are we? *Journal*

of Systematics and Evolution, *58*, 533-545.

Anderson, J. T., Wadgymar, S. M. (2020). Climate change disrupts local adaptation and favours

upslope migration. *Ecology Letters*, *23*, 181-192.

Angseesing, J. P. A. (1974). Selective eating of the acyanogenic form of *Trifolium*

repens. *Heredity*, *32*, 73.

Anstett, D. N., Ahern, J. R., Johnson, M. T. J., Salminen, J.-P. (2018). Testing for latitudinal

gradients in defense at the macroevolutionary scale. *Evolution*, *72*, 2129-2143.

Atkins, K. E., Travis, J. M. J. (2010). Local adaptation and the evolution of species' ranges under

climate change. *Journal of Theoretical Biology*, *266*, 449-457.

Atwood, S. S., Sullivan, J. T. (1943). Inheritance of a cyanogenetic glucoside and its hydrolyzing

enzyme in *Trifolium repens*. *Journal of Heredity*, *34*, 311-320.

Baskett, C. A., Schemske, D. W. (2018). Latitudinal patterns of herbivore pressure in a

temperature herb support the biotic interactions hypothesis. *Ecology Letters*, *21*, 578-587.

Bastias, C. C., Estarague, A., Vile, D., Gaignon, E., Lee, C.-R., Exposito-Alonso, M., … ,

Vasseur, F. (2024). Ecological trade-offs drive phenotypic and genetic differentiation of

Arabidopsis thaliana in Europe. *Nature Communications*, *15*, 5185.

Battlay, P., Hendrickson, B. T., Hendez-Reneau, J. I., Santangelo, J. S., Albano, L. J., Wilson, J.,

… , Kooyers, N. J. (2024). Structural variants underlie parallel adaptation following

global invasion. *biorxiv*. doi: 10.1101/2024.07.09.602765

Beatty, D. W. (1956). Effect of photoperiod and temperature on flowering of white clover

(*Trifolium repens*). Iowa State University.

Bidart-Bouzat, M. G., Imeh-Nathaniel, A. (2008). Global change effects on plant chemical

defenses against insect herbivores. *Journal of Integrative Plant Biology*, *50*, 1339-1354.

Blossey, B., Nötzold, R. (1995). Evolution of increased competitive ability in invasive

nonindigenous plants: A hypothesis. *Journal of Ecology*, *83*, 887-889.

Bock, D. G., Kantar, M. B., Caseys, C., Matthey-Dorey, R., Rieseberg, L. H. (2018). Evaluation

of invasiveness by genetic accommodation. *Nature Ecology & Evolution*, *2*, 991-999.

Bontrager, M., Muir, C. D., Mahony, C., Gamble, D. E., Germain, R. M., Hargreaves, A. L., … ,

Angert, A. L. (2020). Climate warming weakens local adaptation. *biorxiv*. doi:

10.1101/2020.11.01.364349

Brighton, F., Horne, M. T. (1977). Influence of temperature on cyanogenic polymorphisms.

Nature, *265*, 437-438.

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., … ,

Bolker, B. M. (2014) glmmTMB balances speed and flexibility among packages for zero-

inflated generalized linear mixed modeling. *The R Journal*, *9*, 378-400.

Burdon, J. J. (1983). *Trifolium repens* L. *Journal of Ecology*, *71*, 307-330.

Burns, J. E., Ashman, T.-L., Steets, J. A., Harmon-Threatt, A., Knight, T. M. (2011). A

phylogenetically controlled analysis of the roles of reproductive traits in plant invasions.

Oecologia, *166*, 1009-1017.

Byers, J. E. (2002). Impact of non-indigenous species on natives enhanced by anthropogenic

alteration of selection regimes. *Oikos*, *97*, 449-458.

Callaway, R. M., Ridenour, W. M. (2004). Novel weapons: invasive success and the evolution of

increased competitive ability. *Frontiers in Ecology and the Environment*, *2*, 436-443.

Callaway, R. M., Lucero, J. E., Hierro, J. L., Lortie, C. J. (2022). The EICA is dead? Long live

the EICA! *Ecology Letters*, *25*, 2289-2302.

Capblancq, T., Fitzpatrick, M. C., Bay, R. A., Exposito-Alonso, M., Keller, S. R. (2020).

Genomic prediction of (mal)adaptation across current and future climatic landscapes.

Annual Review of Ecology, Evolution, and Systematics, *51*, 245-269.

Colautti, R. I., Barrett, S. C. H. (2013). Rapid adaptation to climate facilitates range expansion of

an invasive plant. *Science*, *342*, 364-367.

Colautti, R. I., Lau, J. A. (2015). Contemporary evolution during invasion: evidence for

differentiation, natural selection, and local adaptation. *Molecular Ecology*, *24*, 1999-

2017.

Coley, P. D., Bryant, J. P., Chapin, F. S. (1985). Resource availability and plant antiherbivore
defense. *Science*, *230*, 895-899.

Coley, P. D., Barone, J. A. (1996). Herbivory and plant defenses in tropical forests. *Annual*
Review of Ecology and Systematics, *27*, 305-335.

Corkill, L. D. (1940). Cyanogenesis in white clover (*Trifolium repens* L.). I. Cyanogenesis in
single plants. *New Zealand Journal of Science and Technology, 22*, 65-67.

Costan, C.-A., Godsoe, W. K., Bufford, J. L., Marris, J. W. M., Hulme, P. E. (2022). Can the
enemy release hypothesis explain the success of *Rumex* (Polygonaceae) species in an
introduced range? *Biological Invasions*, *24*, 2665-2677.

Daday, H. (1954a). Gene frequencies in wild populations of *Trifolium repens.* I. Distribution by
latitude. *Heredity*, *8*, 61-78.

Daday, H. (1954b). Gene frequencies in wild populations of *Trifolium repens.* II. Distribution by
altitude. *Heredity*, *8*, 377-384.

Daday, H. (1958). Gene frequencies in wild populations of *Trifolium repens.* III. World
distribution. *Heredity*, *12*, 169-184.

Daday, H. (1965). Gene frequencies in wild populations of *Trifolium repens.* IV. Mechanism of
natural selection. *Heredity*, *20*, 355-365.

de Villemereuil, P., Mouterde, M., Gaggiotti, O. E., Till-Bottraud, I. (2018). Patterns of
phenotypic plasticity and local adaptation in the wide elevation range of the alpine plant
Arabis alpina. Journal of Ecology, *106*, 1952-1971.

Dirzo, R., Harper, J. L. (1982). Experimental studies on slug-plant interactions: III. Differences
in the acceptability of individual plants of *Trifolium repens* to slugs and snails. *Journal of*
Ecology, *70*, 101-117.

Dlugosch, K. M., Parker, I. M. (2008a). Invading populations of an ornamental shrub show rapid

life history evolution despite genetic bottlenecks. *Ecology Letters*, *11*, 701-709.

Dlugosch, K. M., Parker, I. M. (2008b). Founding events in species invasions: genetic variation,

adaptive evolution, and the role of multiple introductions. *Molecular Ecology*, *17*, 431-

449.

Dostálek, T., Rokaya, M. B., Münzbergová, Z. (2020). Plant palatability and trait responses to

experimental warming. *Scientific Reports*, *10*, 10526.

Easlon, H. M., Bloom, A. J. (2014). Easy Leaf Area: Automated digital image analysis for rapid

and accurate measurement of leaf area. *Applications in Plant Sciences*, *2*, 1400033.

Ehrlich, P. R., Raven, P. H. (1964). Butterflies and plants: A study in coevolution. *Evolution*, *18*,

586-608.

Erfmeier, A. (2013). Constraints and release at difference scales – The role of adaptation in

biological invasions. *Basic and Applied Ecology*, *14*, 281-288.

Escudero, A., Iriondo, J. M., Olano, J. M., Rubio, A., Somolinos, R. C. (2000). Factors affecting

the establishment of a gypsophyte: The case of *Lepidum subulatum* (Brassicaceae).

American Journal of Botany, *87*, 861-871.

Estoup, A., Ravigné, V., Hufbauer, R., Vitalis, R., Gautier, M., Facon, B. (2016). Is there a

genetic paradox of biological invasion? *Annual Review of Ecology, Evolution, and*

Systematics, *47*, 51-72.

Exposito-Alonso, M., 500 Genomes Field Experiment Team, Burbano, H. A., Bossdorf, O.,

Nielsen, R., Weigel, D. (2019). Natural selection on the *Arabidopsis thaliana* genome in

present and future climates. *Nature*, *573*, 126-129.

Fadoul, H. E., Albano, L. J., Bergman, M. E., Phillips, M. A., Johnson, M. T. J. (2023).

Assessing the benefits and costs of the hydrogen cyanide antiherbivore defense in

Trifolium repens. *Plants*, *12*, 1213.

Fine, P. V. A., Miller, Z. J., Mesones, I., Irazuzta, S., Appel, H. M., Stevens, M. H. H., … ,

Coley, P. D. (2006). The growth-defense trade-off and habitat specialization by plants in

Amazonian forests. *Ecology*, *87*, S150-S162.

Fournier-Level, A., Korte, A., Cooper, M. D., Nordborg, M., Schmitt, J., Wilczek, A. M. (2011).

A map of local adaptation in *Arabidopsis thaliana*. *Science*, *334*, 86-89.

Fox, J., Weisberg, S. (2019). An R companion to applied regression, Third edition. Sage,

Thousand Oaks, CA, USA.

Fugère, V., Hendry, A. P. (2018). Human influences on the strength of phenotypic selection.

Proceedings of the National Academy of Sciences USA. *115*, 10070-10075.

Genton, B. J., Kotanen, P. M., Cheptou, P.-O., Adolphe, C., Shykoff, J. A. (2005). Enemy

release but no evolutionary loss of defence in a plant invasion: an inter-continental

reciprocal transplant experiment. *Oecologia*, *146*, 404-414.

Gillespie, J. H., Turelli, M. (1989). Genotype-environment interactions and the maintenance of

polygenic variation. *Genetics*, *121*, 129-138.

Gleadow, R. M., Woodrow, I. E. (2002). Constraints on effectiveness of cyanogenic glycosides

in herbivore defense. *Journal of Chemical Ecology*, *28*, 1301-1313.

Griffiths, A. G., Moraga, R., Tausen, M., Gupta, V., Bitton, T. P., Campbell, M. A., … ,

Anderson, S. U. (2019). Breaking free: The genomics of allopolyploidy-facilitated niche

expansion in white clover. *The Plant Cell*, *31*, 1466-1487.

Hahn, P. G., Agrawal, A. A., Sussman, K. I., Maron, J. L. (2019). Population variation,

environmental gradients, and the evolutionary ecology of plant defense against herbivory.

The American Naturalist, *193*, 20-34.

Halbritter, A. H., Billeter, R., Edwards, P. J., Alexander, J. M. (2015). Local adaptation at range

edges: comparing elevation and latitudinal gradients. *Journal of Evolutionary Biology*,

28, 1849-1860.

Hargreaves, A. L., Germain, R. M., Bontrager, M., Persi, J., Angert, A. L. (2020). Local

adaptation to biotic interactions: A meta-analysis across latitudes. *The American*

Naturalist, *195*, 395-411.

Hendrickson, B. T., Stamps, C., Patterson, C. M., Strickland, H., Foster, M., Kim, A., … ,

Kooyers, N. J. (2024). Evolution of drought resistant strategies following the introduction

of white clover (*Trifolium repens* L.). *biorxiv*. doi: 10.1101/2024.09.03.611023

Hendry, A. P. (2017). Eco-evolutionary dynamics. Princeton University Press. Princeton, NJ,

USA.

Hereford, J. (2009). A quantitative survey of local adaptation and fitness trade-offs. *The*

American Naturalist, *173*, 579-588.

Herms, D. A., Mattson, W. J. (1992). The dilemma of plants: to grow or defend. *Quarterly Review of*

Biology, *67*, 478-478.

Hoffmann, A. A., Sgrò, C. M. (2011). Climate change and evolutionary adaptation. *Nature*, *470*,

479-485.

Hughes, M. A. (1991). The cyanogenic polymorphism in *Trifolium repens* L. (white clover).

Heredity, *66*, 105-115.

Hulme, P. E. (2008). Phenotypic plasticity and plant invasions: is it all Jack? *Functional*

Ecology, *22*, 3-7.

Innes, S., Santangelo, J. S., Kooyers, N. J., Olsen, K. M., Johnson, M. T. J. (2022). Evolution in

response to climate in the native and introduced ranges of a globally distributed plant.

Evolution, *76*, 1495-1511.

IPBES. (2019). Global assessment report on biodiversity and ecosystem services of the

Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.

Brondizio, E. S., Settele, J., Díaz, S., Ngo, H. T. (editors). IPBES secretariat, Bonn,

Germany.

Jeschke, J. M., Pyšek, P. (2018). Tens rule. In *Invasion biology: Hypotheses and evidence* (pp.

124-132). CAB International. Wallingford, UK.

Johnson, M. T. J., Rasmann, S. (2011). The latitudinal herbivory-defence hypothesis takes a

detour on the map. *New Phytologist*, *191*, 589-592.

Johnson, M. T. J., Bertrand, J. A., Turcotte, M. M. (2016). Precision and accuracy in quantifying

herbivory. *Ecological Entomology*, *41*, 112-121.

Kakes, P. (1989). An analysis of the costs and benefits of the cyanogenic system in *Trifolium*

repens L. *Theoretical and Applied Genetics*, *77*, 111-118.

Kassambara, A., Mundt, F. (2020). factoextra: Extract and visualize the results of multivariate

data analyses. R package version 1.0.7. https://CRAN.R-project.org/package=factoextra

Kawecki, T. J., Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, *7*, 1225-

1241.

Keane, R. M., Crawley, M. J. (2002). Exotic plant invasions and the enemy release hypothesis.

Trends in Ecology & Evolution, *17*, 164-170.

Kjærgaard, T. (2003). A plant that changed the world: The rise and fall of clover 1000-2000.

Landscape Research, *28*, 41-49.

Kolar, C. S., Lodge, D. M. (2001). Progress in invasion biology: predicting invaders. *Trends in*

Ecology & Evolution, *16*, 199-204.

Kooyers, N. J., Olsen, K. M. (2012). Rapid evolution of an adaptive cyanogenesis cline in

introduced North American white clover (*Trifolium repens* L.). *Molecular Ecology*, *21*,

2455-2468.

Kooyers, N. J., Olsen, K. M. (2013). Searching for the bull's eye: agents and targets of selection

vary among geographically disparate cyanogenesis clines in white clover (*Trifolium*

repens L.). *Heredity*, *11*, 495-504.

Kooyers, N. J., Gage, L. R., Al□Lozi, A., Olsen, K. M. (2014). Aridity shapes cyanogenesis

cline evolution in white clover (*Trifolium repens* L.). *Molecular Ecology*, *23*, 1053-1070.

Kooyers, N. J., Greenlee, A. B., Colicchio, J. M., Oh, M., Blackman, B. K. (2015). Replicate

altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress

in annual *Mimulus guttatus*. *New Phytologist*, *206*, 152-165.

Kooyers, N. J., Hartman Bakken, B., Ungerer, M. C., Olsen, K. M. (2018). Freeze□induced

cyanide toxicity does not maintain the cyanogenesis polymorphism in white clover

(*Trifolium repens*). *American Journal of Botany*, *105*, 1224-1231.

Kooyers, N. J., Colicchio, J. M., Greenlee, A. B., Patterson, E., Handloser, N. T., Blackman, B.

K. (2019). Lagging adaptation to climate supersedes local adaptation to herbivory in an

annual monkeyflower. *The American Naturalist*, *194*, 541-557.

Kuo, W.-H., Zhong, L., Wright, S. J., Goad, D. M., Olsen, K. M. (2024). Beyond cyanogenesis:

Temperature gradients drive environmental adaptation in North American white clover

(*Trifolium repens* L.). *Molecular Ecology*, *33*, e17484.

Lau, J. A., terHorst, C. P. (2015). Causes and consequences of failed adaptation to biological

invasions: the role of ecological constraints. *Molecular Ecology*, *24*, 1987-1998.

Lee, C.-R., Mitchell-Olds, T. (2011). Quantifying effects of environmental and geographical

factors on patterns of genetic differentiation. *Molecular Ecology*, *20*, 4631-4642.

Leger, E. A., Rice, K. J. (2007). Assessing the speed and predictability of local adaptation in

invasive California poppies (*Eschscholzia californica*). *Journal of Evolutionary Biology*,

20, 1090-1103.

Leimu, R., Fischer, M. (2008). A meta-analysis of local adaptation in plants. *PLoS One*, *3*,

e4010.

Lemoine, N. P., Drews, W. A., Burkepile, D. E., Parker, J. D. (2013). Increased temperature

alters feeding behaviour of a generalist herbivore. *Oikos*, *122*, 1669-1678.

Lenth R (2024). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.10.4.

Li, X.-M., She, D.-Y., Zhang, D.-Y., Liao, W.-J. (2015). Life history trait differentiation and

local adaptation in invasive populations of *Ambrosia artemisiifolia* in China. *Oecologia*,

177, 669-677.

Lucas, M. S., Hensen, I., Barratt, C. D., Callaway, R. M., Durka, W., Lekberg, Y., … , Rosche,

C. (2024). Re-focusing sampling, design and experimental methods to assess rapid

evolution by non-native plant species. *Biological Invasions*, *26*, 1327-1343.

Macel, M., Lawson, C. S., Mortimer, S. R., Šmilauerova, M., Bischoff, A., Crémieux, L., … ,

Steinger, T. (2007). Climate vs. soil factors in local adaptation of two common plant

species. *Ecology*, *88*, 424-433.

Machingura, M., Salomon, E., Jez, J. M., Ebbs, S. D. (2016). The β-cyanoalanine synthase

pathway: beyond cyanide detoxification. *Plant, Cell and Environment*, *39*, 2329-2341.

Maron, J. L., Vilà, M., Bommarco, R., Elmendorf, S., Beardsley, P. (2004). Rapid evolution of

an invasive plant. *Ecological Monographs*, *74*, 261-280.

Massad, T. J., Dyer, L. A. (2010). A meta-analysis of the effects of global environmental change

on plant-herbivore interactions. *Arthropod-Plant Interactions*, *4*, 181-188.

Miles, L. S., Breitbart, S. T., Wagner, H. H., Johnson, M. T. J. (2019). Urbanization shapes the

ecology and evolution plant-arthropod herbivore interactions. *Frontiers in Ecology and*

Evolution, *7*, 310.

Mitchell, C. E., Agrawal, A. A., Bever, J. D., Gilbert, G. S., Hufbauer, R. A., Klironomos, J. N.,

… , Vázquez, D. P. (2006). Biotic interactions and plant invasions. *Ecology Letters*, *9*,

726-740.

Moles, A. T., Westoby, M. (2004). Seedling survival and seed size: a synthesis of the literature.

Journal of Ecology, *92*, 372-383.

Møller, B. L. (2010). Functional diversifications of cyanogenic glucosides. *Current Opinion in*

Plant Biology, *13*, 338-347.

Moreira, X., Castagneyrol, B., Abdala-Roberts, L., Berny-Mier y Teran, J. C., Timmermans, B.

G. H., Bruun, H. H., … , Tack, A. J. M. (2018). Latitudinal variation in plant chemical

defences drives latitudinal patterns of leaf herbivory. *Ecography*, *41*, 1124-1134.

Müller-Schärer, H., Schaffner, U., Steinger, T. (2004). Evolution in invasive plants: implications

for biological control. *Trends in Ecology and Evolution*, *19*, 417-422.

Nagy, D. U., Thoma, A. E., Al-Gharaibeh, M., Callaway, R. M., Flory, S. L., Frazee, L. J., … ,

Rosche, C. (2024). Among-population variation in drought responses is consistent across

life stages but not between native and non-native ranges. *New Phytologist*, *243*, 922-935.

Murray-Stoker, D., Johnson, M. T. J. (2024). Mosaic of local adaptation between white clover

and rhizobia along an urbanization gradient. *Journal of Ecology*, *112*, 1150-1163..

O'Donnell, M. S., Ignizio, D. A. (2012). Bioclimatic predictors for supporting ecological

applications in the conterminous United States. *US Geological Survey Data Series*, *10*, 4-

9.

Oduor, A. M. O., Leimu, R., van Kleunen, M. (2016). Invasive plant species are locally adapted

just as frequently and at least as strongly as native plant species. *Journal of Ecology*, *104*,

957-968.

Olsen, K. M., Sutherland, B. L., Small, L. L. (2007). Molecular evolution of the *Li/li* chemical

defence polymorphism in white clover (*Trifolium repens* L.). *Molecular Ecology*, *16*,

4180-4193.

Olsen, K. M., Small, L. L. (2018). Micro- and macroevolutionary adaptation though repeated

loss of a complete metabolic pathway. *New Phytologist*, *219*, 757-766.

Olsen, K. M., Goad, D. M., Wright, S. J., Dutta, M. L., Myers, S. R., Small, L. L., Li, L.-F.

(2021). Dual-species origin of an adaptive chemical defense polymorphism. *New*

Phytologist, *232*, 1477-1487.

Pal, R. W., Maron, J. L., Nagy, D. U., Waller, L. P., Tosto, A., Liao, H., Callaway, R. M. (2020).

What happens in Europe stays in Europe: apparent evolution by an invader does not help

at home. *Ecology*, *101*, e3072.

Pannell, J. R., Auld, J. R., Brandvain, Y., Burd, M., Busch, J. W., Cheptou, P.-O., … , Winn, A.

A. (2015). The scope of Baker's law. *New Phytologist*, *208*, 656-667.

Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annual*

Review of Ecology, Evolution, and Systematics. *37*, 637-669.

Pickett, S. T. A. (1989). Space-for-time substitution as an alternative for long-term studies. In

Long-term studies in ecology: approaches and alternatives (pp. 110-135). Springer. New

York, NY, USA.

Pyšek, P., Hulme, P. E., Simberloff, D., Bacher, S., Blackburn, T. M., Carlton, J. T. … ,

Richardson, D. M. (2020). Scientists' warning on invasive alien species. *Biological*

Reviews, *95*, 1511-1534.

Saikkonen, K., Taulavuori, K., Hyvönen, T., Gundel, P. E., Hamilton, C. E., Vänninen, I., … ,

Helander, M. (2012). Climate change-driven species' range shifts filtered by

photoperiodism. *Nature Climate Change*, *2*, 239-242.

Samis, K. E., Stinchcombe, J. R., Murren, C. J. (2019). Population climatic history predicts

phenotypic responses in novel environments for *Arabidopsis thaliana* in North America.

American Journal of Botany, *106*, 1068-1080.

Santangelo, J. S., Ness, R. W., Cohan, B., Fitzpatrick, C. R., Innes, S. G., Koch, S., … , Johnson,

M. T. J. (2022). Global urban environmental change drives adaptation in white clover.

Science, *375*, 1275-1281.

Savolainen, O., Lascoux, M., Merilä, J. (2013). Ecological genomics of local adaptation. *Nature*

Reviews Genomics, *14*, 807-820.

Sheng, M., Rosche, C., Al-Gharaibeh, M., Bullington, L. S., Callaway, R. M., Clark, T., … ,

Lekberg, Y. (2022). Acquisition and evolution of enhanced mutualism—an

underappreciated mechanism for invasive success? *The ISME Journal*, *16*, 2467-2478.

Siepielski, A. M., Morrissey, M. B., Buoro, M., Carlson, S. M., Caruso, C. M., Clegg, S. M., … ,

MacColl, A. D. C. (2017). Precipitation drives global variation in natural selection.

Science, *355*, 959-962.

Simón-Porcar, V. I., Silva, J. L., Vallejo-Marín, M. (2021). Rapid local adaptation in both sexual

and asexual invasive populations of monkeyflowers (*Mimulus* spp.). *Annals of Botany*,

127, 655-668.

St Clair, J. B., Howe, G. T. (2007). Genetic maladaptation of coastal Douglas-fir seedlings to

future climates. *Global Change Biology*, *13*, 1441-1454.

Taylor, C. M., Hastings, A. (2005). Allee effects in biological invasions. *Ecology Letters*, *8*, 895-

908.

Thompson, K. A., Renaudin, M., Johnson, M. T. J. (2016). Urbanization drives the evolution of

parallel clines in plant populations. *Proceedings of the Royal Society B: Biological*

Sciences, *283*, 20162180.

Till, I. (1987). Variability of expression of cyanogenesis in white clover (*Trifolium repens*

L.). *Heredity*, *59*, 265.

Toräng, P., Wunder, J., Obeso, J. T., Herzog, M., Coupland, G, Ågren, J. (2015). Large-scale

adaptive differentiation in the alpine perennial *Arabis alpina*. *New Phytologist*, *206*,

459-470.

Tsutsui, N. D., Suarez, A. V., Holway, D. A., Case, T. J. (2000). Reduced genetic variation and

the success of an invasive species. *Proceedings of the National Academy of Sciences*, *97*,

5948-5953.

Turesson, G. (1922). The genotypical response of the plant species to the habitat. *Hereditas*, *3*,

211-350.

Valladares, F., Matesanz, S., Guilhaumon, F., Araújo, M. B., Balaguer, L., Benito-Garzón, M.,

… , Zavala, M. A. (2014). The effects of phenotypic plasticity and local adaptation on

forecasts of species range shifts under climate change. *Ecology Letters*, *17*, 1351-1364.

van Boheemen, L. A., Lombaert, E., Nurkowski, K. A., Gauffre, B., Rieseber, L. H., Hodgins, K.

A. (2017). Multiple introductions, admixture and bridgehead invasion characterize the

introduction history of *Ambrosia artemisiifolia* in Europe and Australia. *Molecular*

Ecology, *26*, 5421-5434.

Walther, G.-R., Roques, A., Hulme, P. E., Sykes, M. T., Pyšek, P., Kühn, I., … , Settele, J.

(2009). Alien species in a warmer world: risks and opportunities. *Trends in Ecology &*

Evolution, *24*, 686-693.

Wang, T., Hamann, A., Spittlehouse, D., Carroll, C. (2016). Locally downscaled and spatially

customizable climate data for historical and future periods for North America. *PLoS One*,

11, e0156720.

Wilczek, A. M., Cooper, M. D., Korves, T. M., Schmitt, J. (2014). Lagging adaptation to

warming climate in *Arabidopsis thaliana*. *Proceedings of the National Academy of*

Sciences, *111*, 7906-7913.

Wolkovich, E. M., Davies, T. J., Schaefer, H., Cleland, E. E., Cook, B. I., Travers, S. E., … ,

Davis, C. C. (2013). Temperature-dependent shifts in phenology contribute to the success

of exotic species with climate change. *American Journal of Botany*, *100*, 1407-1421.

Wright, S. J., Cui Zhou, D., Kuhle, A., Olsen, K. M. (2018). Continent-wide climatic variation

drives local adaptation in North American white clover. *Journal of Heredity*, *109*, 78-89.

Wright, S. J., Goad, D. M., Gross, B. L., Muñoz, P. R., Olsen, K. M. (2021). Genetic trade-offs

underlie divergent life strategies for local adaptation in white clover. *Molecular Ecology*,

31, 3742-3760.

Züst, T., Heichinger, C., Grossniklaus, U., Harrington, R., Kliebenstein, D. J., Turnbull, L. A.

(2012). Natural enemies drive geographic variation in plant defenses. *Science*, *338*, 116-

119.

1176

1177 **Table 1** Results of mixed effects models assessing how continent of origin, common garden

1178 location, and climatic distance affected multiple fitness components of *T. repens*.

	Plants Producing Flowers			Plants Producing Seeds		
Predictor	χ^2	df	*P*	χ^2	df	*P*
Continent of Origin (O)	0.377	1	0.539	0.271	1	0.603
Garden Location (L)	**32.302**	**3**	**< 0.001**	**8.023**	**3**	**0.046**
Climatic Distance (D)	**6.308**	**1**	**0.012**	**7.197**	**1**	**0.007**
O × L	0.773	3	0.856	0.724	3	0.868
O × D	0.299	1	0.584	0.876	1	0.349
L × D	4.222	3	0.238	3.944	3	0.268
Garden Slope Estimates	Slope		SE	Slope		SE
Lafayette	− 0.251		0.115	− 0.255		0.124
Mississauga	0.027		0.120	0.021		0.131
Montpellier	0.061		0.172	0.097		0.174
Uppsala	− 0.099		0.113	− 0.091		0.116

	# of Flower Heads			Seed Set Mass		
Predictor	χ^2	df	*P*	χ^2	df	*P*
Continent of Origin (O)	2.270	1	0.132	1.136	1	0.286
Garden Location (L)	**247.413**	**3**	**< 0.001**	**39.550**	**3**	**< 0.001**
Climatic Distance (D)	**13.326**	**1**	**< 0.001**	**14.168**	**1**	**< 0.001**
O × L	1.415	3	0.702	4.059	3	0.255
O × D	0.271	1	0.602	0.873	1	0.350
L × D	**12.474**	**3**	**0.006**	**8.553**	**3**	**0.036**
Garden Slope Estimates	Slope		SE	Slope		SE
Lafayette	− 0.597		0.168	− 0.674		0.223
Mississauga	− 0.169		0.170	− 0.273		0.217
Montpellier	− 0.037		0.252	0.053		0.316
Uppsala	0.036		0.147	0.002		0.185

	Survival			Maximum Plant Area		
Predictor	χ^2	df	*P*	χ^2	df	*P*
Continent of Origin (O)	2.408	1	0.121	0.924	1	0.336
Garden Location (L)	**224.949**	**3**	**< 0.001**	**285.680**	**3**	**< 0.001**
Climatic Distance (D)	**6.633**	**1**	**0.010**	0.582	1	0.446
O × L	6.049	3	0.109	6.467	3	0.091
O × D	0.093	1	0.761	1.752	1	0.186
L × D	4.609	3	0.203	5.620	3	0.132
Garden Slope Estimates	Slope		SE	Slope		SE
Lafayette	− 0.109		0.111	− 0.119		0.124
Mississauga	− 0.010		0.117	− 0.150		0.138
Montpellier	0.032		0.167	0.609		0.339
Uppsala	− 0.260		0.100	− 0.214		0.126

Predictor	Growth Rate			Herbivory		
	χ^2	df	P	χ^2	df	P
Continent of Origin (O)	0.094	1	0.760	0.088	1	0.766
Garden Location (L)	**69.948**	**3**	**< 0.001**	**143.951**	**3**	**< 0.001**
Climatic Distance (D)	0.087	1	0.768	1.070	1	0.301
O × L	1.207	3	0.751	6.613	3	0.085
O × D	0.049	1	0.825	0.010	1	0.920
L × D	2.873	3	0.412	2.226	3	0.527
Garden Slope Estimates	Slope		SE	Slope		SE
Lafayette	0.001		0.003	0.177		0.155
Mississauga	− 0.004		0.003	0.055		0.127
Montpellier	0.002		0.006	− 0.358		0.346
Uppsala	0.002		0.002	0.079		0.136

1179 Note: Significant effects ($P < 0.05$) are listed in bold. Slope estimates and standard errors (SE)

1180 are listed individually for each common garden location.

1181

1182 **Table 2** Results of mixed effects models assessing how continent of origin, continent of garden,

1183 and climatic distance affected the fitness of *T. repens*.

1184

Predictor	Plants Producing Flowers		Plants Producing Seeds	
	χ^2	P	χ^2	P
Continent of Origin (O)	2.033	0.154	1.754	0.185
Continent of Garden (G)	**9.184**	**0.002**	1.295	0.255
Climatic Distance (D)	**28.963**	**< 0.001**	**17.269**	**< 0.001**
O × G	0.677	0.411	0.942	0.332
O × D	2.033	0.154	0.036	0.850
G × D	3.194	0.074	2.490	0.115
O × G × D	0.501	0.479	0.026	0.871

O × G Slope Estimates	Slope	SE	Slope	SE
North America/North America	− 0.207	0.067	− 0.241	0.076
North America/Europe	− 0.072	0.096	− 0.086	0.097
Europe/North America	− 0.363	0.071	− 0.231	0.077
Europe/Europe	− 0.069	0.178	− 0.038	0.181

Predictor	# of Flower Heads		Seed Set Mass	
	χ^2	P	χ^2	P
Continent of Origin (O)	0.514	0.474	0.028	0.867
Continent of Garden (G)	0.471	0.493	0.037	0.847
Climatic Distance (D)	**14.310**	**< 0.001**	**16.190**	**< 0.001**
O × G	0.957	0.328	0.481	0.488
O × D	1.214	0.271	2.991	0.084
G × D	1.412	0.235	0.258	0.612
O × G × D	**19.603**	**< 0.001**	**9.728**	**0.002**

O × G Slope Estimates	Slope	SE	Slope	SE
North America/North America	− 0.482	0.125	− 0.631	0.139
North America/Europe	− 0.237	0.162	− 0.179	0.169
Europe/North America	0.097	0.139	− 0.006	0.151
Europe/Europe	− 1.351	0.301	− 0.826	0.313

Predictor	Survival		Maximum Plant Area	
	χ^2	P	χ^2	P
Continent of Origin (O)	0.106	0.745	0.797	0.372
Continent of Garden (G)	**41.220**	**< 0.001**	**170.553**	**< 0.001**
Climatic Distance (D)	**10.236**	**0.001**	**58.341**	**< 0.001**
O × G	0.000	0.993	2.375	0.123
O × D	1.534	0.216	**34.173**	**< 0.001**
G × D	**13.870**	**< 0.001**	1.784	0.182
O × G × D	**6.515**	**0.011**	0.678	0.410

$O \times G$ Slope Estimates	Slope	SE	Slope	SE
North America/North America	− 0.133	0.067	− 0.109	0.078
North America/Europe	0.062	0.078	− 0.005	0.122
Europe/North America	− 0.357	0.069	− 0.721	0.073
Europe/Europe	0.335	0.152	− 0.385	0.230

	Growth Rate		Herbivory	
Predictor	χ^2	P	χ^2	P
Continent of Origin (O)	0.053	0.817	0.196	0.658
Continent of Garden (G)	1.468	0.226	**21.180**	**< 0.001**
Climatic Distance (D)	**18.137**	**< 0.001**	**15.810**	**< 0.001**
$O \times G$	0.465	0.495	0.669	0.413
$O \times D$	**5.145**	**0.023**	**23.686**	**< 0.001**
$G \times D$	2.263	0.133	0.240	0.624
$O \times G \times D$	**7.812**	**0.005**	2.178	0.140

$O \times G$ Slope Estimates	Slope	SE	Slope	SE
North America/North America	0.002	0.002	0.069	0.091
North America/Europe	0.003	0.002	− 0.122	0.135
Europe/North America	0.012	0.002	− 0.591	0.094
Europe/Europe	− 0.003	0.004	− 0.311	0.257

Note: Significant effects ($P < 0.05$) are listed in bold. All predictors are assessed using one

degree of freedom. Slope estimates and standard errors (SE) are listed individually for each

common garden location.

1191 **Table 3** Results of mixed effects models assessing how change in mean annual temperature

1192 (ΔMAT) affected the fitness of North American *T. repens* populations in the Mississauga

1193 common garden (linear and quadratic fits).

Predictor	Plants Producing Flowers		Plants Producing Seeds		# of Flower Heads		Seed Set Mass	
	χ^2	*P*	χ^2	*P*	χ^2	*P*	χ^2	*P*
ΔMAT	**7.425**	**0.006**	**5.397**	**0.020**	0.255	0.614	0.644	0.422
ΔMAT^2	**8.077**	**0.004**	**7.609**	**0.006**	2.908	0.088	**8.542**	**0.003**

Predictor	Survival		Maximum Plant Area		Growth Rate	
	χ^2	*P*	χ^2	*P*	χ^2	*P*
ΔMAT	3.758	0.053	3.799	0.051	0.002	0.962
ΔMAT^2	**4.313**	**0.038**	**6.741**	**0.009**	1.808	0.179

1194 Note: Significant effects ($P < 0.05$) are listed in bold. All predictors are assessed using one

1195 degree of freedom.

1196

Figure 1

Figure 1 Maps showing populations of *T. repens* sampled from the (**A**, **C**) introduced range

(North America, circles) and (**B**, **D**) native range (Europe, grey circles), as well as the four

common garden locations (black diamonds). Map panels **A** and **B** are set on a backdrop of

colour-graded historical (1970-2000) mean annual temperature (BIO1; with darker red shades

representing warmer climates). Map panels **C** and **D** are set on a backdrop of colour-graded

historical (1970-2000) annual precipitation (BIO12; with darker blue shades representing wetter

1204   climates). Coordinates of common garden locations are 42°32'40" N, 79°39'38" W

1205   (Mississauga, Ontario, Canada), 30°18'23" N, 92°00'33" W (Lafayette, Louisiana, USA),

1206   59°49'08" N, 17°38'49" W (Uppsala, Sweden), and 43°38'16" N, 3°51'43" W (Montpellier,

1207   France). All climate data was extracted from WorldClim (Version 2.1, February 2023).

1208

1209

Figure 2

Figure 2 The effect of climatic distance and common garden location on fitness of *T. repens*. (**A**)

Mean number of flower heads and (**B**) mean seed set mass of *T. repens* populations planted in

two common gardens in North America (Lafayette in navy and Mississauga in orange) and two

common gardens in Europe (Montpellier in green and Uppsala in magenta). Shaded areas

represent 95% confidence intervals around each line, while asterisks represent *P* values of

relevant fixed effects of Garden Location (L), Climatic Distance (D) and their interaction (*$*P <$

0.05, **$P < 0.01$, ***$P < 0.001$).

1218

1219   Figure 3

1220

Figure 3 The effect of climatic distance on the mean number of flower heads produced by *T.*

repens populations originating from its introduced (North America, orange) and native (Europe,

pink) ranges and transplanted into common gardens in the (**A**) introduced and (**B**) native ranges.

The effect of climatic distance on the mean seed set mass produced by *T. repens* populations

originating from its introduced and native ranges and transplanted into common gardens in the

(**C**) introduced and (**D**) native ranges. Shaded areas represent 95% confidence intervals around

each line, while asterisks represent *P* values of < 0.05 for relevant fixed effects of Climatic

Distance (D) and its interaction with Continent of Origin (O) and Continent of Garden (G).

1229

1230

1231   Figure 4

1232

Figure 4 The effects of the change in mean annual temperature (ΔMAT) on fitness of North

American populations of *T. repens* grown in the Mississauga common garden. (**A**) Proportion of

plants survived, (**B**) proportion of plants that produced flowers, (**C**) proportion of plants that

produced seeds, (**D**) mean seed set mass, and (**E**) mean maximum plant area per population all

exhibited evidence of an adaptation lag in response to climate warming, indicated by a positive

linear trend or a shift in the peak of the quadratic curve towards populations with warmer

climates of origin (denoted as x-axis values > 0). Shaded areas represent 95% confidence

intervals around each line.

A transcontinental experiment elucidates (mal)adaptation of a cosmopolitan plant to climate in space and time

Lucas J. Albano[1,2], Cristina C. Bastias[3,4], Aurélien Estarague[4], Brandon T. Hendrickson[5], Simon G. Innes[5], Nevada King[5], Courtney M. Patterson[5], Amelia Tudoran[6], François Vasseur[4], Adriana Puentes[6], Cyrille Violle[4], Nicholas J. Kooyers[5], Marc T. J. Johnson[2]

[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

[2]Department of Biology, University of Toronto Mississauga, Mississauga, ON, Canada

[3]Ecology Area, Department of Botany, Ecology and Plant Physiology, Campus de Rabanales, Córdoba University, Córdoba, Andalusia, Spain

[4]CEFE, Université Montpellier, CNRS, EPHE, IRD, Montpellier, Occitanie, France

[5]Department of Biology, University of Louisiana at Lafayette, Lafayette, LA, USA

[6]Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Uppland, Sweden

Correspondence Author:

Lucas J. Albano

Current address: Department of Plant & Microbial Biology, North Carolina State University, Raleigh, NC, United States, 27695, ljalbano@ncsu.edu

Open Research Statement:

Data are provided for peer review. All data involved in this study is available on the GitHub page

for LJA (https://github.com/ljalbano/transcontinental_common_garden).

Keywords:

Adaptation lag, climatic distance, common garden, cyanogenesis, environmental variation,

introduced range, non-native species, local adaptation, native range, provenance study, *Trifolium*

repens, white clover

Table S1 Historical bioclimatic data (1970-2000) for each common garden site.

Bioclimatic Variable	Mississauga	Lafayette	Uppsala	Montpellier
BIO1 (°C)	8.4	19.7	5.9	14.7
BIO5 (°C)	26.9	33.1	21.9	28.8
BIO6 (°C)	− 9.1	4.4	− 7.0	3.1
BIO12 (mm)	813	1578	568	657
BIO13 (mm)	79	158	73	97
BIO14 (mm)	49	110	26	21

Note: Bioclimatic variables are mean annual temperature (BIO1), maximum temperature of the warmest month (BIO5), minimum temperature of the coldest month (BIO6), annual precipitation (BIO12), precipitation of the wettest month (BIO13), and precipitation of the driest month (BIO14). Data was extracted from WorldClim (Version 2.1, February 2023).

Table S2 Bivariate Pearson's correlation coefficients between all bioclimatic variables of interest, latitude, and climatic distance for each common garden.

	BIO1	BIO5	BIO6	BIO12	BIO13	BIO14	Latitude
BIO1	—	—	—	—	—	—	—
BIO5	0.751	—	—	—	—	—	—
BIO6	0.816	0.264	—	—	—	—	—
BIO12	0.556	0.577	0.181	—	—	—	—
BIO13	0.549	0.564	0.176	0.961	—	—	—
BIO14	0.508	0.506	0.182	0.936	0.841	—	—
Latitude	− 0.759	− 0.937	− 0.273	− 0.733	− 0.722	− 0.647	—
CD (Lafayette)	− 0.828	− 0.754	− 0.478	− 0.921	− 0.891	− 0.874	0.851
CD (Mississauga)	0.661	0.728	0.235	0.789	0.824	0.693	− 0.752
CD (Montpellier)	0.282	0.505	− 0.162	0.741	0.761	0.676	− 0.554
CD (Uppsala)	0.768	0.797	0.342	0.915	0.912	0.834	− 0.882

Note: Bioclimatic variables are mean annual temperature (BIO1), maximum temperature of the warmest month (BIO5), minimum temperature of the coldest month (BIO6), annual precipitation (BIO12), precipitation of the wettest month (BIO13), and precipitation of the driest month (BIO14) for each of the 96 locations of source populations of *T. repens*. Climatic distances (CD) are included separately for the four common garden sites (Lafayette, Mississauga, Montpellier, and Uppsala). Bioclimatic data was extracted from WorldClim (Version 2.1, February 2023).

Table S3 Summary of transformations used for each quantitative response variable.

Model	# of Flower Heads	Seed Set Mass	Maximum Plant Area	Growth Rate	Herbivory
(Q1) Response ~ Continent of Origin * Garden Location * Climatic Distance + (1|Continent of Origin/Population)	Log transformation	Log transformation	Log transformation	Square root transformation	Square root transformation
(Q2) Response ~ Continent of Origin * Continent of Garden * Climatic Distance + (1|Continent of Origin/Population)	Log transformation	Log transformation	Log transformation	Square root transformation	Square root transformation
(Q3; Mississauga Garden) Response ~ Bioclimatic Variable + Bioclimatic Variable^2 + (1|Population)	Log transformation	Log transformation	Log transformation	No transformation	—
(Q3; Lafayette Garden) Response ~ Bioclimatic Variable + Bioclimatic Variable^2 + (1|Population)	Log transformation	Log transformation	Square root transformation	Log transformation	—
(Q3; Uppsala Garden) Response ~ Bioclimatic Variable + Bioclimatic Variable^2 + (1|Population)	Square root transformation	Log transformation	Square root transformation	No transformation	—
(Q3; Montpellier Garden) Response ~ Bioclimatic Variable + Bioclimatic Variable^2 + (1|Population)	Square root transformation	Log transformation	Square root transformation	No transformation	—
(Q4; Mississauga Garden) Response ~ MAT + MAT^2 + (1|Population)	Log transformation	Log transformation	Log transformation	No transformation	—
(Q4; Uppsala Garden) Response ~ MAT + MAT^2 + (1|Population)	Log transformation	Log transformation	Log transformation	No transformation	—
(Q5) Response ~ Cyanotype * Garden Location * Climatic Distance + (1|Population)	Log transformation	Log transformation	Log transformation	Log transformation	Square root transformation

Note: Transformations were performed to meet assumptions of normality and homogeneity of variance. All models described in the table are fit to a normal distribution and all figures display means of raw data to facilitate visual interpretation. All models in the study that were fit to the binomial distribution did not require further transformation and therefore are not included in the table.

Table S4 Sample sizes for each model used in the study, divided by the research question that each model set addresses and each of

the eight response variables analyzed.

	Survival	Plants Producing Flowers	Plants Producing Seeds	# of Flower Heads	Seed Set Mass	Maximum Plant Area	Growth Rate	Herbivory
Q1	1858	1858	1858	406	389	1262	730	949
Q2	1858	1858	1858	406	389	1262	730	949
Q3	554	554	554	218	211	553	343	507
Q4 (Lafayette)	465	465	465	77	66	323	129	—
Q4 (Mississauga)	465	465	465	117	113	451	188	—
Q4 (Montpellier)	470	470	470	102	102	127	102	—
Q4 (Uppsala)	458	458	458	110	108	361	311	—
Q5 (Mississauga)	235	235	235	64	61	229	91	—
Q5 (Uppsala)	233	233	233	58	57	179	153	—

Table S5 Results of mixed effects models assessing how individual bioclimatic variables (both linearly and quadratically) affected the fitness of *T. repens* in the Mississauga common garden.

Predictor	Survival χ^2	P	Plants Producing Flowers χ^2	P	Plants Producing Seeds χ^2	P	# of Flower Heads χ^2	P	Seed Set Mass χ^2	P	Maximum Plant Area χ^2	P	Growth Rate χ^2	P
BIO1	**3.603**	**0.058**	**6.119**	**0.013**	**4.762**	**0.029**	0.116	0.733	1.042	0.307	2.385	0.122	0.054	0.816
BIO1^2	**3.271**	**0.071**	**3.921**	**0.048**	**3.488**	**0.062**	1.807	0.179	**7.080**	**0.008**	**3.734**	**0.053**	2.648	0.104
BIO5	1.354	0.245	**6.704**	**0.010**	**6.684**	**0.010**	0.303	0.582	0.145	0.704	0.652	0.419	1.699	0.193
BIO5^2	0.639	0.424	**2.920**	**0.088**	2.701	0.100	0.024	0.878	0.742	0.389	0.944	0.331	0.154	0.695
BIO6	1.211	0.271	1.588	0.208	1.273	0.259	0.010	0.920	0.037	0.847	0.405	0.525	0.314	0.575
BIO6^2	2.452	0.117	**4.997**	**0.025**	**5.763**	**0.016**	0.720	0.396	**3.318**	**0.069**	2.605	0.107	2.661	0.103
BIO12	1.077	0.299	1.162	0.281	0.066	0.797	0.087	0.768	0.200	0.655	1.554	0.213	2.144	0.143
BIO12^2	0.883	0.347	0.321	0.571	0.004	0.948	0.558	0.455	**3.110**	**0.078**	1.930	0.165	**3.496**	**0.062**
BIO13	0.610	0.435	0.403	0.525	0.099	0.753	0.100	0.752	0.290	0.590	2.254	0.133	**6.168**	**0.013**
BIO13^2	0.377	0.539	0.043	0.837	0.015	0.902	0.055	0.815	1.450	0.229	1.963	0.161	**6.641**	**0.010**
BIO14	0.173	0.677	0.562	0.453	0.064	0.800	2.066	0.151	1.708	0.191	0.031	0.860	0.051	0.822
BIO14^2	1.175	0.279	0.692	0.406	0.002	0.964	0.896	0.344	**3.996**	**0.046**	1.290	0.256	2.330	0.127

Note: Bioclimatic variables are mean annual temperature (BIO1), maximum temperature of the warmest month (BIO5), minimum temperature of the coldest month (BIO6), annual precipitation (BIO12), precipitation of the wettest month (BIO13), and precipitation of the driest month (BIO14). All variables were z-score standardized to allow for simultaneous direct comparisons of the relative strength of linear and quadratic predictor variables. Effects with $P < 0.1$ are bolded and the effect in red exhibited the lowest P-value among all independent variables tested for a given fitness measure (among P values < 0.1). All predictors are assessed using one degree of freedom.

Table S6 Results of mixed effects models assessing how individual bioclimatic variables (both linearly and quadratically) affected the fitness of *T. repens* in the Lafayette common garden.

Predictor	Survival χ^2	P	Plants Producing Flowers χ^2	P	Plants Producing Seeds χ^2	P	# of Flower Heads χ^2	P	Seed Set Mass χ^2	P	Maximum Plant Area χ^2	P	Growth Rate χ^2	P
BIO1	**3.562**	**0.059**	**4.562**	**0.033**	**4.498**	**0.034**	0.514	0.474	2.559	0.110	0.006	0.940	1.062	0.303
BIO1^2	2.237	0.135	0.767	0.381	0.556	0.456	0.291	0.590	0.051	0.821	0.000	0.988	1.467	0.226
BIO5	0.109	0.741	0.866	0.352	0.421	0.516	0.112	0.738	1.950	0.163	**6.484**	**0.011**	0.005	0.946
BIO5^2	0.491	0.484	0.012	0.912	0.108	0.742	0.032	0.859	0.426	0.514	**8.242**	**0.004**	0.047	0.829
BIO6	2.680	0.102	1.279	0.258	2.156	0.142	1.672	0.196	1.100	0.294	0.052	0.820	1.252	0.263
BIO6^2	2.375	0.123	0.264	0.607	0.473	0.492	0.009	0.923	0.000	0.991	0.013	0.909	1.746	0.186
BIO12	0.307	0.580	2.493	0.114	**3.151**	**0.076**	**2.989**	**0.084**	**7.048**	**0.008**	0.426	0.514	0.046	0.830
BIO12^2	0.006	0.937	0.399	0.528	0.659	0.417	0.910	0.340	**2.831**	**0.092**	0.955	0.328	0.024	0.877
BIO13	1.157	0.282	**5.384**	**0.020**	**5.918**	**0.015**	2.139	0.144	**5.549**	**0.018**	0.171	0.679	0.745	0.388
BIO13^2	0.169	0.681	1.328	0.249	1.640	0.200	0.231	0.631	1.577	0.209	0.781	0.377	0.591	0.442
BIO14	0.377	0.539	**4.254**	**0.039**	**5.182**	**0.023**	1.860	0.172	**3.346**	**0.067**	0.879	0.349	0.404	0.525
BIO14^2	0.047	0.829	1.485	0.223	1.823	0.177	0.551	0.458	0.955	0.328	1.407	0.236	0.442	0.506

Note: Bioclimatic variables are mean annual temperature (BIO1), maximum temperature of the warmest month (BIO5), minimum temperature of the coldest month (BIO6), annual precipitation (BIO12), precipitation of the wettest month (BIO13), and precipitation of the driest month (BIO14). All variables were z-score standardized to allow for simultaneous direct comparisons of the relative strength of linear and quadratic predictor variables. Effects with $P < 0.1$ are bolded and the effect in red exhibited the lowest P-value among all independent variables tested for a given fitness measure (among P values < 0.1). All predictors are assessed using one degree of freedom.

Table S7 Results of mixed effects models assessing how individual bioclimatic variables (both linearly and quadratically) affected the fitness of *T. repens* in the Uppsala common garden.

Predictor	Survival χ^2	Survival P	Plants Producing Flowers χ^2	Plants Producing Flowers P	Plants Producing Seeds χ^2	Plants Producing Seeds P	# of Flower Heads χ^2	# of Flower Heads P	Seed Set Mass χ^2	Seed Set Mass P	Maximum Plant Area χ^2	Maximum Plant Area P	Growth Rate χ^2	Growth Rate P
BIO1	1.749	0.186	0.332	0.564	0.226	0.635	0.295	0.587	0.004	0.948	0.512	0.475	0.209	0.648
BIO1^2	0.114	0.736	1.291	0.256	1.137	0.286	0.583	0.445	0.009	0.923	0.242	0.622	1.389	0.239
BIO5	**3.906**	**0.048**	**5.261**	**0.022**	**5.104**	**0.024**	0.377	0.539	1.694	0.193	0.083	0.773	**4.633**	**0.031**
BIO5^2	**6.392**	**0.011**	**6.781**	**0.009**	**6.643**	**0.010**	0.361	0.548	1.665	0.197	0.072	0.789	**5.401**	**0.020**
BIO6	**9.121**	**0.003**	0.457	0.499	0.532	0.466	0.048	0.826	0.325	0.569	0.656	0.418	0.717	0.397
BIO6^2	0.722	0.396	2.530	0.112	2.381	0.123	0.779	0.377	0.027	0.870	0.001	0.979	0.841	0.359
BIO12	**9.972**	**0.002**	0.778	0.378	0.782	0.377	0.973	0.324	0.001	0.972	0.535	0.465	0.454	0.501
BIO12^2	**12.362**	**< 0.001**	1.197	0.274	1.269	0.260	0.862	0.353	0.005	0.944	0.536	0.464	1.929	0.165
BIO13	**8.552**	**0.003**	0.603	0.437	0.713	0.398	0.519	0.471	0.021	0.886	0.379	0.538	0.075	0.785
BIO13^2	**11.256**	**< 0.001**	1.104	0.293	1.343	0.247	0.516	0.473	0.051	0.822	0.240	0.624	1.219	0.270
BIO14	**6.626**	**0.010**	1.180	0.277	1.058	0.304	2.045	0.153	0.177	0.674	0.742	0.389	0.515	0.473
BIO14^2	**8.578**	**0.003**	1.696	0.193	1.673	0.196	1.797	0.180	0.001	0.981	1.155	0.283	1.931	0.165

Note: Bioclimatic variables are mean annual temperature (BIO1), maximum temperature of the warmest month (BIO5), minimum temperature of the coldest month (BIO6), annual precipitation (BIO12), precipitation of the wettest month (BIO13), and precipitation of the driest month (BIO14). All variables were z-score standardized to allow for simultaneous direct comparisons of the relative strength of linear and quadratic predictor variables. Effects with $P < 0.1$ are bolded and the effect in red exhibited the lowest P-value among all independent variables tested for a given fitness measure (among P values < 0.1). All predictors are assessed using one degree of freedom.

Table S8 Results of mixed effects models assessing how individual bioclimatic variables (both linearly and quadratically) affected the fitness of *T. repens* in the Montpellier common garden.

Predictor	Survival χ^2	P	Plants Producing Flowers χ^2	P	Plants Producing Seeds χ^2	P	# of Flower Heads χ^2	P	Seed Set Mass χ^2	P	Maximum Plant Area χ^2	P	Growth Rate χ^2	P
BIO1	0.248	0.618	1.006	0.316	1.024	0.312	1.599	0.206	1.397	0.237	**3.389**	**0.066**	1.002	0.317
BIO1^2	0.317	0.574	0.027	0.869	0.089	0.765	1.771	0.183	1.210	0.271	0.487	0.485	0.001	0.974
BIO5	0.076	0.782	0.360	0.548	0.263	0.608	0.660	0.417	0.043	0.836	1.633	0.201	0.017	0.898
BIO5^2	0.793	0.373	0.780	0.377	1.250	0.264	0.024	0.877	1.274	0.259	0.357	0.550	0.376	0.540
BIO6	0.519	0.471	0.688	0.407	0.685	0.408	1.039	0.308	2.139	0.144	**6.489**	**0.011**	**5.509**	**0.019**
BIO6^2	0.816	0.366	0.225	0.635	0.287	0.592	0.130	0.719	0.070	0.791	**3.978**	**0.046**	**2.862**	**0.091**
BIO12	0.202	0.653	0.244	0.621	0.094	0.760	**2.797**	**0.094**	1.208	0.272	2.606	0.106	0.459	0.498
BIO12^2	0.151	0.697	0.513	0.474	0.430	0.512	2.577	0.108	0.586	0.444	**5.722**	**0.017**	1.085	0.298
BIO13	0.715	0.398	1.007	0.316	0.602	0.438	**3.470**	**0.063**	1.323	0.250	0.701	0.403	0.071	0.790
BIO13^2	0.582	0.446	1.204	0.273	0.960	0.327	**3.588**	**0.058**	1.240	0.266	1.344	0.246	0.000	0.987
BIO14	0.000	0.991	0.021	0.884	0.065	0.799	2.539	0.111	2.177	0.140	**2.758**	**0.097**	1.317	0.251
BIO14^2	0.502	0.579	0.895	0.344	1.017	0.313	1.880	0.170	0.515	0.473	**7.215**	**0.007**	**3.214**	**0.073**

Note: Bioclimatic variables are mean annual temperature (BIO1), maximum temperature of the warmest month (BIO5), minimum temperature of the coldest month (BIO6), annual precipitation (BIO12), precipitation of the wettest month (BIO13), and precipitation of the driest month (BIO14). All variables were z-score standardized to allow for simultaneous direct comparisons of the relative strength of linear and quadratic predictor variables. Effects with $P < 0.1$ are bolded and the effect in red exhibited the lowest P-value among all independent variables tested for a given fitness measure (among P values < 0.1). All predictors are assessed using one degree of freedom.

Table S9 Results of mixed effects models assessing how change in mean annual temperature (not z-score standardized) affected the

fitness of European *T. repens* populations in the Uppsala common garden (linear and quadratic fits).

Predictor	Survival		Plants Producing Flowers		Plants Producing Seeds		# of Flower Heads		Seed Set Mass		Maximum Plant Area		Growth Rate	
	χ^2	*P*	χ^2	*P*	χ^2	*P*	χ^2	*P*	χ^2	*P*	χ^2	*P*	χ^2	*P*
ΔMAT	1.612	0.204	0.429	0.513	0.306	0.580	0.003	0.957	0.006	0.941	0.000	0.999	0.101	0.751
ΔMAT^2	2.267	0.132	0.595	0.440	0.440	0.507	0.028	0.867	0.070	0.792	0.004	0.953	0.184	0.668

Note: All predictors are assessed using one degree of freedom.

Table S10 Results of mixed effects models assessing how latitude and continent of origin affected the proportion of *Ac* and *Li* alleles found in populations of *T. repens*.

Predictor	Proportion *Ac*		Proportion *Li*	
	χ^2	*P*	χ^2	*P*
Latitude	**7.830**	**0.005**	**22.362**	**< 0.001**
Continent of Origin	**14.772**	**< 0.001**	**17.765**	**< 0.001**
Latitude × Continent of Origin	1.044	0.307	0.042	0.837

Note: Significant effects ($P < 0.05$) are listed in bold. All predictors are assessed using one degree of freedom.

Table S11 Results of mixed effects models assessing how cyanotype (AcLi, Acli, acLi, or acli), common garden location, and climatic distance affected the fitness of *T. repens*.

Predictor	Plants Producing Flowers			Plants Producing Seeds			# of Flower Heads			Seed Set Mass		
	χ^2	df	*P*	χ^2	df	*P*	χ^2	df	*P*	χ^2	df	*P*
Cyanotype (C)	**16.862**	**3**	**< 0.001**	**11.731**	**3**	**0.008**	6.635	3	0.084	1.801	3	0.615
Garden Location (L)	**29.547**	**3**	**< 0.001**	**43.214**	**3**	**< 0.001**	**125.477**	**3**	**< 0.001**	**19.881**	**3**	**< 0.001**
Climatic Distance (D)	**9.464**	**1**	**0.002**	**9.816**	**1**	**0.002**	**22.595**	**1**	**< 0.001**	**23.040**	**1**	**< 0.001**
C × L	13.049	9	0.160	11.819	9	0.224	6.558	9	0.683	2.835	9	0.970
C × D	2.089	3	0.554	3.236	3	0.357	3.926	3	0.270	4.463	3	0.216
C × L × D	10.144	12	0.603	13.401	12	0.341	11.509	12	0.486	10.804	12	0.546

Predictor	Survival			Maximum Plant Area			Growth Rate			Herbivory		
	χ^2	df	*P*	χ^2	df	*P*	χ^2	df	*P*	χ^2	df	*P*
Cyanotype (C)	**19.889**	**3**	**< 0.001**	**9.841**	**3**	**0.020**	0.381	3	0.944	4.956	3	0.175
Garden Location (L)	**56.896**	**3**	**< 0.001**	**116.238**	**3**	**< 0.001**	**79.026**	**3**	**< 0.001**	**85.426**	**3**	**< 0.001**
Climatic Distance (D)	**3.903**	**1**	**0.048**	**6.887**	**1**	**0.009**	3.835	1	0.050	0.237	1	0.626
C × L	8.506	9	0.484	7.049	9	0.632	6.369	9	0.703	3.326	9	0.950
C × D	1.723	3	0.632	2.726	3	0.436	1.251	3	0.741	3.724	3	0.293
C × L × D	6.019	12	0.915	6.575	12	0.884	19.085	12	0.087	8.448	12	0.749

Note: Significant effects (*P* < 0.05) are listed in bold.

Figure S1 Eigenvectors of six bioclimatic variables (z-score standardized) on the three most influential dimensions of the principal component analysis used to determine climatic distance between each population of origin and each common garden. (**A**) Plot showing eigenvectors along dimensions 1 and 2. (**B**) Plot showing eigenvectors along dimension 1 and 3. (**C**) Plot showing eigenvectors along dimensions 2 and 3. Bioclimatic variables included in the principal component analysis are mean annual temperature (BIO1), maximum temperature of the warmest month (BIO5), minimum temperature of the coldest month (BIO6), annual precipitation (BIO12), precipitation of the wettest month (BIO13), and precipitation of the driest month (BIO14).

Figure S2 Results of a principal component analysis including six bioclimatic variables of interest, where each point represents the z-score standardized values on the three most influential dimensions (three-dimensional plot viewed from different angles) for each population from the native range (Europe, grey), each population from the introduced range (North America, brown) and each common garden location (black; Mississauga: Mi; Lafayette: La; Uppsala: Up; Montpellier: Mo). Weighted Euclidean distances were calculated between each population and each common garden using their PC1, PC2, and PC3 loading values as weights in order to determine a measure of climatic distance.

Figure S3 The effect of climatic distance and common garden location on fitness of *T. repens*. (**A**) Proportion of plants survived, (**B**) proportion of plants that produced flowers, (**C**) proportion of plants that produced seeds, (**D**) mean maximum plant area, (**E**) mean growth rate and (**F**) mean herbivory (measured as the percentage of leaf area consumed) for *T. repens* populations planted in four common gardens (Lafayette in navy, Montpellier in green, Mississauga in orange, and Uppsala in magenta). Shaded areas represent 95% confidence intervals around each line.

Figure S4 The effect of climatic distance on the proportion of *T. repens* survived when originating from the introduced (North America, orange) and native ranges (Europe, pink) and transplanted into common gardens in the (**A**) introduced and (**B**) native ranges. The effect of climatic distance on the proportion of plants flowered when originating from the introduced and native ranges and transplanted into the (**C**) introduced and (**D**) native ranges. The effect of climatic distance on the proportion of plants producing seeds when originating from the introduced and native ranges and transplanted into the (**E**) introduced and (**F**) native ranges. The effect of climatic distance on the maximum

area of plants originating from the introduced and native ranges and transplanted into the (**G**) introduced and (**H**) native ranges. The effect of climatic distance on growth rate of plants originating from the introduced and native ranges and transplanted into the (**I**) introduced and (**J**) native ranges. The effect of climatic distance on herbivory of plants originating from the introduced and native ranges and transplanted into the (**K**) introduced and (**L**) native ranges. Points represent mean values for each population within each garden. Shaded areas represent 95% confidence intervals around each line.

Figure S5 (**A**) The proportion of *T. repens* plants possessing at least one dominant *Ac* allele per sampled population in either the native range (Europe, pink) or introduced range (North America, orange) as a function of the latitude of origin of each population. (**B**) The proportion of *T. repens* plants possessing at least one dominant *Li* allele per sampled population in either the native or introduced range as a function of the latitude of origin of each population. Shaded areas represent 95% confidence intervals around each line.

Figure S6 Associations between inferred genomic coverage of cyanogenesis genes and Feigl-Anger cyanogenesis phenotypes. Coverage of

cyanogenesis genes is inferred from low coverage whole genome sequencing reads found within loci associated with

the *Ac*/*ac* and *Li*/*li* genes. The x-axis is corrected for the total number of reads because samples with few reads may not have reads within

these regions by chance (i.e., not because the plant possessed the associated gene deletion). Lines represents generalized linear models fit

with a binomial distribution.

Supplementary Methods

DNA Extraction and Library Preparation

Our DNA extraction and library construction have been previously described (Battley et al., 2024). Briefly, we collected 1-3 trifoliate leaves per plant and either stored tissue at -80°C until extraction at University of Louisiana, Lafayette (ULL) or lyophilized tissue prior to shipping to ULL. DNA was extracted with a modified cetyltrimethylammonium bromide (CTAB) procedure using increased concentrations of CTAB and 2-β-mercaptoethanol (Doyle and Doyle, 1987; Saeidi et al., 2018). DNA concentrations were quantified and diluted to 3 ng/uL before undergoing a modified Nextera dual-index library preparation (Therkildsen and Palumbi, 2017). We used Nextera XT Index Kit (Illumina; San Diego, CA, USA) for barcoding libraries. Each library was quantified and concentrations normalized before pooling of 48 samples with unique barcodes into a single tube. Libraries were sequenced on a HiSeq X Ten platform by Novogene (Sacramento, CA, USA) using paired end 150 bp reads. In total, 656 samples (Lafayette, Louisiana, USA = 138; Mississauga, Ontario, Canada = 327; Montpellier, France = 111; Uppsala, Sweden = 79) were sequenced across 14 sequencing lanes.

Bioinformatics Pipeline

For each sample, adapter sequences and poly-g tails were trimmed from reads using *fastp* (Chen et al., 2018). An FM-index was constructed for the *Trifolium repens* reference genome (Santangelo et al., 2023) using the bwa indexing algorithm with default settings (Li and Durbin, 2009). Reads were then aligned to the reference genome using bwa mem (Li and Durbin, 2009). SAM files were converted to BAM file format, coordinate sorted, and indexed using SAMtools (Li et al., 2009). To find the genomic coordinates of the *Ac*/ac and *Li/li* loci in the *T. repens* reference genome, we took 5 sample sequences of the *CYP79D15* locus (the *Ac/ac* gene; Olsen et al. 2008) and *Li* locus (*Li/li* gene; Olsen et al., 2007) from the NCBI database and compared nucleotide similarity of NCBI samples with our reference genome using the basic local alignment search tool (BLAST; Camacho et al., 2009). We

found 99% similarity between 5 NCBI nucleotide sequences of the *Li* gene with Chr04_Pall:28511241 - 28515079. For *CYP79D15*, we found 99% sequence similarity at sites Chr02_Occ:9222822 – 9224484 and Chr02_Occ: 9536228 – 9537890. Constraining our search to these regions, we calculated the number of reads aligning to these regions using the 'view -c' command and calculated read coverage using the 'coverage' command within SAMtools (Li et al., 2009). This resulted in number of reads and read coverage data for 586 samples (Lafayette, Louisiana, USA = 99; Mississauga, Ontario, Canada = 302; Montpellier, France = 107; Uppsala, Sweden = 78) after removing samples that sequenced poorly (< 100 Mb of read data).

Statistics

Our low coverage whole genome sequencing approach for direct determination of genotypes at the *Ac/ac* or *Li/li* genes (referred to below as cyanotypes). We took a statistical genomic approach to determining genotypes and evaluating potential type I and type II error in our cyanotypes. Since variation at *Ac/ac* and *Li/li* is caused by deletions, we used read coverage within the underlying genomic regions as evidence for the presence of the *Ac* or *Li* alleles. We first removed any samples with < 500,000 reads (20 samples). We use read coverage of each locus rather than number of reads as number of reads is likely more influenced by variable gene deletion sizes and misalignment of homeologous sequences. Indeed, our downstream analyses suggest the number of reads metric produces higher error than coverage in genotype estimates. For each locus, we first correct our coverage values for variation in sequencing effort using a general linear model with the *lm* function in R Version 4.2.2 (2022) to associate coverage with total number of reads across the entire genome. The residuals from this model reflect effort-corrected coverage of a locus. We then use a set of individuals (N = 178) whose cyanotypes were previously determined by Feigl-Anger assays to model the relationship between coverage at each locus and the presence of at least one dominant allele (*Ac* or *Li*; Feigl and Anger, 1966). Separate univariate generalized linear models implemented using the *glm*

function with a binomial family and logit link were conducted for each locus with presence or absence of the dominant allele assessed via Feigl-Anger assay as the response variable and the residual of coverage model as the predictor variable. Statistical significance of these associations was assessed via a Wald statistic implemented via the *Anova* function in the *car* package (version 3.1-2; Fox et al., 2013).

These models were highly significant for each locus (*Ac*: $\chi^2 = 77.3$, df = 1, *P* < 2.2e-16; *Li*: $\chi^2 = 110.6$, df = 1, *P* < 2.2e-16; **Figure S6**), indicating that coverage was a valuable predictor of cyanotype. From our predicted models, we calculated the inflection point of the model for each locus and used this value to predict the presence or absence of *Ac* and *Li* alleles for every sample. Plants with higher values than the inflection point that the model estimate for each locus were called as having either the *Ac* or *Li* allele present. Using this method to estimate cyanotypes of samples with known cyanotypes via FA-assay, we find 142/178 genotypes were correctly determined for the *Ac/ac* gene (79.8%) and 154/178 for the *Li/li* gene (86.5%). Errors were skewed toward type II error (false negatives) as would be expected with our low coverage whole genome sequencing approach. We also note that Feigl-Anger assays have some inherent error, especially when conducted on small amount of tissue from field-grown plants. We are unable to attribute the differences between approaches to error in either of the assays and thus just note the substantial concordance.

References

Battlay, P., Hendrickson, B. T., Mendez-Reneau, J. I., Santangelo, J. S., Albano, L. J., Wilson, J., … , Kooyers, N. J. (2024). Structural variants underlie parallel adaptation following global invasion. *biorxiv*.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*, *10*, 1-9.

Chen, S., Zhou, Y., Chen, Y., Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, *34*, i884-i890.

Doyle, J. J., Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical bulletin*, *19*, 11-15.

Feigl, F., Anger, V. (1966). Replacement of benzidine by copper ethylacetoacetate and tetra base as spot-test reagent for hydrogen cyanide and cyanogen. *The Analyst*, *91*, 282-284.

Fox, J., Friendly, M., Weisberg, S. (2013). Hypothesis tests for multivariate linear models using the car package. *R Journal*, *5*, 39-52.

Li, H., Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, *25*, 1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*, 2078-2079.

Olsen, K. M., Hsu, S.-C., Small, L. L. (2008). Evidence on the molecular basis of the *Ac/ac* adaptive cyanogenesis polymorphism in white clover (*Trifolium repens* L.). *Genetics*, *179*, 517-526.

Olsen, K. M., Sutherland, B. L., Small, L. L. (2007). Molecular evolution of the *Li/li* chemical defence polymorphism in white clover (*Trifolium repens* L.). *Molecular Ecology*, *16*, 4180-4193.

Saeidi, S., McKain, M. R., Kellogg, E. A. (2018). Robust DNA isolation and high-throughput sequencing library construction for herbarium specimens. *Journal of Visual Experiments*, *133*, e56837.

Santangelo, J. S., Battlay, P., Hendrickson, B. T., Kuo, W.-H., Olsen, K. M., Kooyers, N. J., … , Ness, R. W. (2023). Haplotype-resolved, chromosome-level assembly of white clover (*Trifolium repens* L., Fabaceae). *Genome Biology and Evolution*, *15*, evad146.

Therkildsen, N. O., Palumbi, S. R. (2017). Practical low-coverage genomewide sequencing of hundreds of individually barcoded samples for population and evolutionary genomics in nonmodel species. *Molecular Ecology Resources*, *17*, 194-208.