The Primate Major Histocompatibility Complex: An Illustrative Example of Gene Family Evolution

Alyssa Lyn Fortier[1,2]* and Jonathan K. Pritchard[1,2]

*For correspondence:
afortier@stanford.edu (ALF);
pritch@stanford.edu (JKP)

[1]Department of Biology, Stanford University, Stanford, CA USA; [2]Department of Genetics, Stanford University, Stanford, CA USA

Abstract   Gene families are groups of evolutionarily-related genes. One large gene family that has experienced rapid evolution is the Major Histocompatibility Complex (MHC), whose proteins serve critical roles in innate and adaptive immunity. Across the ~60 million year history of the primates, some MHC genes have turned over completely, some have changed function, some have converged in function, and others have remained essentially unchanged. Past work has typically focused on identifying MHC alleles within particular species or comparing gene content, but more work is needed to understand the overall evolution of the gene family across species. Thus, despite the immunologic importance of the MHC and its peculiar evolutionary history, we lack a complete picture of MHC evolution in the primates. We readdress this question using sequences from dozens of MHC genes and pseudogenes spanning the entire primate order, building a comprehensive set of gene and allele trees with modern methods. Overall, we find that the Class I gene subfamily is evolving much more quickly than the Class II gene subfamily, with the exception of the Class II MHC-DRB genes. We also pay special attention to the often-ignored pseudogenes, which we use to reconstruct different events in the evolution of the Class I region. We find that despite the shared function of the MHC across species, different species employ different genes, haplotypes, and patterns of variation to achieve a successful immune response. Our trees and extensive literature review represent the most comprehensive look into MHC evolution to date.

Introduction

Gene families are groups of related genes categorized by functional similarity or presumed evolutionary relatedness. Based on clustering of their proteins' sequences, human genes fall into hundreds to thousands of distinct families (*Gu et al., 2002*; *Li et al., 2001*; *Demuth et al., 2006*; *Friedman and Hughes, 2003*). Families originate from successive gene duplications, although particular gene copies or entire families can also be lost (*Nei et al., 1997*; *Demuth et al., 2006*). For example, there are hundreds of genes that are specific to human or chimpanzee and have no orthologs in the other species (*Demuth et al., 2006*). This birth-and-death evolution is distinct from evolution at the nucleotide or protein level (*Thornton and Desalle, 2000*; *Hahn et al., 2005*). However, phylogenetics can still be applied to understand the relationships within families of genes, providing insight into speciation and specialization (*Thornton and Desalle, 2000*).

   One large gene family is united by a common protein structure called the "MHC fold". This

group of genes is involved in diverse functions including lipid metabolism, iron uptake regulation, and immune system function (*Hansen et al., 2007*; *Kupfermann et al., 1999*; *Kaufman, 2022*; *Adams and Luoma, 2013*). This family includes the Class I and Class II MHC genes whose protein products present peptides to T-cells ("classical" genes) and/or interact with other immune cell receptors like killer cell immunoglobulin-like receptors (KIRs) or leukocyte immunoglobulin-like receptors (LILRs) ("classical" and "non-classical" genes). The classical genes are conventionally known to be highly polymorphic, have an excess of missense variants, and even share alleles across species, all indicative of balancing selection at the allele level (*Maccari et al., 2017*, *2020*; *Robinson et al., 2024*; *Hughes and Nei, 1988*, *1989b*; *Arden and Klein, 1982*; *Mayer et al., 1988*). In addition to variation within individual genes, the region is also significantly structurally divergent across the primates (*Mao et al., 2024*). Balancing selection is evident at the haplotype level as well, where haplotypes with drastically different functional gene content are retained in various primate populations (*Hans et al., 2017*; *de Groot et al., 2017b*, *2009*; *Gleimer et al., 2011*; *Heijmans et al., 2020*). This motivates the need to study the MHC holistically as a gene family. Even though species may retain different sets of genes and haplotypes, related genes likely function similarly, facilitating comparisons across species. Thus, by treating the genes as a related set, our understanding improves significantly compared to considering single genes in isolation. Because gene family birth-and-death is important to speciation and the MHC itself is highly relevant to organismal health, this family is an excellent case for studying gene family evolutionary dynamics.

There are two classes of MHC genes within the greater family (Class I and Class II), and each class contains two functionally distinct types of genes: "classical" and "non-classical". "Classical" MHC molecules perform antigen presentation to T cells—a key part of adaptive immunity—while "non-classical" molecules have niche immune roles. The classical Class I molecules are generally highly polymorphic, ubiquitously expressed, and present short, intracellularly-derived peptides to T cells. Many of them also serve as ligands for other types of immune cell receptors and influence innate immunity (see *Appendix 1* for an overview) (*Anderson et al., 2023*; *Parham and Moffett, 2013*; *Guethlein et al., 2015*; *Hans et al., 2017*; *Wroblewski et al., 2019*). The non-classical Class I molecules have limited polymorphism, restricted expression, and perform specific tasks such as mediating maternal-fetal interaction and monitoring levels of MHC synthesis. In humans, the classical Class I genes are HLA-A, -B, and -C, and the non-classical Class I genes are HLA-E, -F, and -G (*Heijmans et al., 2020*). In contrast, the classical Class II molecules are expressed only on professional antigen-presenting cell types and present longer, extracellularly-derived peptides to T cells (*Gfeller and Bassani-Sternberg, 2018*; *Heijmans et al., 2020*; *Neefjes et al., 2011*). The non-classical Class II molecules assist with loading peptides onto the classical Class II molecules before their transport to the cell surface (*Dijkstra and Yamaguchi, 2019*; *Neefjes et al., 2011*). In humans, HLA-DP, -DQ, and -DR are the classical Class II molecules, and HLA-DM and -DO are the non-classical molecules (see *Appendix 3* for more detail on all of these genes).

However, the landscape of MHC genes differs even across closely-related species. Over evolutionary time, the Class I gene subfamily has been extraordinarily plastic, having undergone repeated expansions, neofunctionalizations, and losses (*Hans et al., 2017*; *Wilming et al., 2013*; *Otting et al., 2020*; *Heijmans et al., 2020*). Convergent evolution has also occurred; in different primate lineages, the same gene may be inactivated, acquire a new function, or even evolve similar splice variants (*Hans et al., 2017*; *Wilming et al., 2013*; *Otting et al., 2020*; *Heijmans et al., 2020*; *Walter, 2020*). As a result, it is often difficult to detect orthologous relationships in Class I even within the primates (*Hughes and Nei, 1989a*; *Piontkivska and Nei, 2003*; *Go et al., 2003*; *Flügge et al., 2002*; *de Groot et al., 2020*). Studies that focus only on the highly-polymorphic binding-site-encoding exons are complicated by these phenomena, necessitating a more comprehensive look into MHC evolution across exons and species groups.

In contrast to Class I, the Class II region has been largely stable across the primates, but gene content still varies in other species. For example, the pig has lost the MHC-DP genes while expanding the number of MHC-DR genes, and the cat has lost both the MHC-DQ and -DP genes, relying

entirely on MHC-DR (*Hammer et al., 2020*; *Okano et al., 2020*). The use of the different Class II molecules appears to be fluid, at least over longer timescales, motivating the need to fill in the gaps in knowledge in the primate tree.

Due to the large volume of existing MHC literature, results are scattered across hundreds of papers, each presenting findings from a limited number of species or genes. Thus, we first performed an extensive literature review to identify the genes and haplotypes known to be present in different primate species. We present a detailed summary of these genes and their functions in *Appendix 3*. We also performed a *BLAST* search using a custom IPD-based MHC allele database against several available reference genomes to discover which genes were present on various primate reference haplotypes (*Figure 1—figure Supplement 2* and *Figure 2—figure Supplement 2*). Our *BLAST* search and our search of NCBI RefSeq confirmed the presence of various genes in several species for the first time. *Figure 1* and *Figure 2* show the landscape of MHC genes present in different primate species for Class I and Class II, respectively. The inclusion of sequences from dozens of new species across all genes and the often-ignored pseudogenes helps us paint a more detailed picture of MHC evolution in the primates.

In this work, we present a large set of densely-sampled Bayesian phylogenetic trees using sequences from a comprehensive set of MHC genes across dozens of primate species. These trees permit us to explore the overall evolution of the gene family and relationships between genes, as well as trace particular allelic lineages over time. Across the trees, we see examples of rapid gene turnover over just a few million years, evidence for long-term balancing selection retaining allelic lineages, and slowly-evolving genes where orthology is retained for long time periods. In this paper, we describe broad-scale differences between classes and discuss some specific results about the relationships between genes. In a companion paper (*Fortier and Pritchard, 2024*), we explore the patterns of polymorphism *within* individual genes, finding evidence for deep trans-species polymorphism at multiple genes.

Results

Data

We collected MHC nucleotide sequences for all genes from the IPD-MHC/HLA database, a large repository for MHC alleles from humans, non-human primates, and other vertebrates (*Maccari et al., 2017*, *2020*; *Robinson et al., 2024*). Although extensive, this database includes few or no sequences from several key lineages including the gibbon, tarsier, and lemur. Thus, we supplemented our set of alleles using sequences from NCBI RefSeq (see asterisks in *Figure 1* and *Figure 2*). Because the MHC genes make up an evolutionarily related family, they can all be aligned. Using *MUSCLE* (*Edgar, 2004*), we aligned all Class I sequences together, all Class IIA sequences together, and all Class IIB sequences together. We then constructed trees for various subsets of these sequences using *BEAST2*, a Bayesian MCMC phylogenetic inference method (see Methods for more detail) (*Bouckaert et al., 2014*, *2019*). One major advantage of *BEAST2* over less tuneable methods is that it can allow evolutionary rates to vary across sites, which is important for genes such as these which experience rapid evolution in functional regions (*Wu et al., 2013*). We also considered each exon separately to minimize the impact of recombination as well as to compare and contrast the binding-site-encoding exons with non-binding-site-encoding exons.

Here, we present these densely-sampled Bayesian phylogenetic trees which include sequences from 106 species and dozens of MHC genes. In this paper, we focus on the Class I, Class IIA, and Class IIB multi-gene trees and discuss overall relationships between genes. Our companion paper (*Fortier and Pritchard, 2024*) explores individual clades/gene groups within these multi-gene trees to understand allele relationships and assess support for trans-species polymorphism.

Figure 1. Class I MHC genes present in different species. The primate evolutionary tree (*Kuderna et al., 2023*) is shown on the left hand side (nonprimate icons are shown in beige). The MHC region has been well characterized in only a handful of species; the rows corresponding to these species are highlighted in gray. Species that are not highlighted have partially characterized or completely uncharacterized MHC regions. Asterisks indicate new information provided by the present study, typically discovery of a gene's presence in a species. Each column/color indicates an orthologous group of genes, labeled at the top and ordered as they are in the human genome (note that not all genes appear on every haplotype). A point indicates that a given gene is present in a given species; when a species has 3 or more paralogs of a given gene, only 3 points are shown for visualization purposes. Filled points indicate that the gene is fixed in that species, outlined points indicate that the gene is unfixed, and semi-transparent points indicate that the gene's fixedness is not known. The shape of the point indicates the gene's role, either a pseudogene, classical MHC gene, non-classical MHC gene, a gene that shares both features ("dual characteristics"), or unknown. The horizontal gray brackets indicate a breakdown of 1:1 orthology, where genes below the bracket are orthologous to 2 or more separate loci above the bracket. The set of two adjacent gray brackets in the top center of the figure show a block duplication. Gene labels in the middle of the plot ("W", "A", "G", "B", and "I") clarify genes that are named differently in different species. OWM, Old-World Monkeys; NWM, New World Monkeys.

Figure 1—source data 1. References for *Figure 1*.

Figure 1—figure supplement 1. A version of *Figure 1* without asterisks.

Figure 1—figure supplement 2. *BLAST* hits for MHC Class I genes in various reference genomes.

Figure 2. Class II MHC genes present in different species. The mammal evolutionary tree is shown on the left hand side, with an emphasis on the primates (*Foley et al., 2023*; *Kuderna et al., 2023*). The rest of the figure design follows that of *Figure 1*, except that we did not need to limit the number of points shown per locus/species due to space constraints. OWM, Old-World Monkeys; NWM, New World Monkeys; Strep., *Strepsirrhini*.

Figure 2—source data 1. References for *Figure 2*.

Figure 2—figure supplement 1. A version of *Figure 2* without asterisks.

Figure 2—figure supplement 2. *BLAST* hits for MHC Class II genes in various reference genomes.

The MHC Across the Primates

The MHC is a particularly dynamic example of a gene family due to intense selective pressure driven by host-pathogen co-evolution (*Radwan et al., 2020*; *Ebert and Fields, 2020*). Within the family, genes have duplicated, changed function, and been lost many times in different lineages. As a result, even closely-related species can have different sets of MHC genes. Thus, while the MHC has been extensively studied in humans, there is a limit to how much we can learn from a single species. Leveraging information from other species helps us understand the evolution of the entire family and provides key context as to how it currently operates in humans (*Thornton and Desalle, 2000*; *Adams and Parham, 2001b*). In *Figure 1* and *Figure 2*, we compare the genes present in different species. In both, each column represents an orthologous gene, while the left-hand-side shows the evolutionary tree for primates and our closest non-primate relatives (*Kuderna et al., 2023*; *Foley et al., 2023*). Humans are part of the ape clade (red label), which is most closely related to the old-world monkeys (OWM; blue label). Next, the ape/OWM clade is most closely related to the new-world monkeys (NWM; orange label), and the ape/OWM/NWM clade is collectively known as the *Simiiformes*. Only species with rows highlighted in gray have had their MHC regions extensively studied. Gene presence in each species is indicated by points in each column, and the points also indicate the function of the gene and whether it is fixed in the species. Points with an asterisk indicate contributions from this work.

Figure 1 shows that not all Class I genes are shared by apes and OWM, and much fewer are shared between apes/OWM and NWM. Genes have also been differently expanded in different lineages. While humans and most other apes have a single copy of each gene, the OWM and NWM have multiple copies of nearly all genes. Additionally, many genes exhibit functional plasticity; for example, MHC-G is a non-classical gene in the apes and a pseudogene in the OWM (it is not 1:1 orthologous to NWM MHC-G). The differences between even closely-related primate groups indicate that the Class I region is evolving very rapidly.

In contrast, the Class II genes are more stable, as the same genes can be found in even distantly-related mammals (*Figure 2*). The notable exception to this pattern is the MHC-DRB group of genes, indicated by dark blue points in the middle of *Figure 2*. While some of the individual MHC-DRB genes are orthologous between apes and OWM, indicated by points in the same column, others are limited to the apes alone. Furthermore, no individual MHC-DRB genes (with the possible exception of MHC-DRB9) are shared between apes/OWM and NWM, pointing to their extremely rapid evolution. While the other genes have been relatively stable, there have been expansions in certain lineages, such as separate duplications of the MHC-DQA and -DQB genes in apes/OWM, NWM, and mouse lemur. Thus, both of the MHC Class I and Class I gene subfamilies appear to be subject to birth-and-death evolution, with Class I and MHC-DRB undergoing the process more rapidly than the rest of Class II.

Evolution of a Gene Family

We performed phylogenetic inference using *BEAST2* on our aligned MHC allele sequences collected from NCBI RefSeq and the IPD-MHC database. *BEAST2* is a Bayesian method, meaning the set of trees it produces represents the posterior space of trees (*Bouckaert et al., 2019*). For visualization purposes, we collapsed the space of trees into a single summary tree that maximizes the sum of posterior clade probabilities (*BEA, 2024*). In each tree, the tips represent sequences, either named with their RefSeq identifier or with standard allele nomenclature (see *Appendix 2*). The summary tree for Class I is shown in *Figure 3*, while the summary trees for Class IIA and Class IIB are shown in *Figure 4*.

*Figure 3*A shows the Class I multi-gene tree using sequences from exon 4, a non-peptide-binding-region-encoding (non-PBR) exon equal in size to each of the peptide-binding-region-encoding (PBR) exons 2 and 3. This exon is the least likely to be affected by convergent evolution, making its tree's structure easier to interpret. This tree—which contains hundreds of tips—has been further simplified by collapsing clades of related tips, although two fully-expanded clades are shown in panels

B and C. Sequences do not always assort by locus. For example, ape MHC-J is separated from OWM MHC-J, which is more closely related to ape/OWM MHC-G. Meanwhile, NWM MHC-G does not group with ape/OWM MHC-G, instead falling outside of the clade containing ape/OWM MHC-A, -G, -J and -K. This supports the fact that the NWM MHC-G genes are broadly orthologous to a large group of genes which expanded within the ape/OWM lineage.

However, some clades/genes do behave in the expected fashion; that is, with their trees matching the overall species tree. One such gene is non-classical MHC-F, shown in *Figure 3*B. Although the gene has duplicated in the common marmoset (Caja-F), this subtree closely matches the species tree shown in the upper right. This indicates that MHC-F is orthologous across apes, OWM, and NWM. Orthology between apes and OWM is also observed for pseudogenes MHC-L, -K, -J, and -V and non-classical MHC-E and -G (*Figure 3—figure Supplement 2* and *Figure 5—figure Supplement 1*). For the other NWM genes, orthology with apes/OWM is less clear.

Other genes do not at all follow the species tree, such as NWM MHC-G. This gene group is broadly orthologous to a large set of ape/OWM genes and pseudogenes, as its ancestor expanded independently in both lineages. In NWM, the many functional MHC-G genes are classical, and there are also a large number of MHC-G-related pseudogenes. Shown in *Figure 3*C, NWM MHC-G sequences do not always group by species (colored box with abbreviation), instead forming mixed clades. Thus, while some duplications appear to have occurred prior to speciation events, others are species-specific. Similar expansions are seen among the MHC-A and -B genes of the OWM and the MHC-B genes of the NWM (*Figure 3—figure Supplement 2*, *Figure 3—figure Supplement 3*, and *Figure 3—figure Supplement 4*).

Figure 4 shows summary trees for exon 3 (non-PBR) for the Class IIA and IIB sequence sets. In the Class II genes, exon 3 does not encode the binding site, but is similar in size to binding-site-encoding exon 2. In contrast to Class I (*Figure 3*), Class II sequences group entirely and unambiguously by gene, shown by the collapsed trees in *Figure 4*A. However, the subtrees for each gene exhibit varying patterns. As with Class I, non-classical genes tend to evolve in a "typical" fashion with sequences assorting according to the species tree. This is clearly the case for non-classical MHC-DMA, -DMB, -DOA, and -DOB (*Figure 4—figure Supplement 1* and *Figure 4—figure Supplement 2*). Classical genes MHC-DRA and -DPA also follow this pattern (*Figure 4*B and *Figure 4—figure Supplement 1*). However, the other classical genes' subtrees look very different from the species tree.

There are several reasons why these MHC gene trees can differ from the overall species tree. Incomplete lineage sorting can happen purely by chance, especially if species have recently diverged. However, balancing selection can cause alleles to be longer-lived, resulting in incomplete lineage sorting even among deeply-diverged species; this is called trans-species polymorphism (TSP). *Figure 4*C illustrates this phenomenon for MHC-DPB. Within the OWM clade (shades of green), sequences group by allelic lineage rather than by species. For example, crab-eating macaque allele Mafa-DRB1*09:02:01:01 groups with green monkey allele Chsa-DPB1*09:01 (both members of the DRB1*09 lineage) rather than with the other macaque alleles (Mane-, Mamu-, Math-, and Malo-), despite the fact that these species are 15 million years separated from each other (*Kuderna et al., 2023*). We see this pattern in many Class II genes and some Class I genes (*Figure 3—figure Supplement 3*, *Figure 3—figure Supplement 4*, *Figure 3—figure Supplement 5* , *Figure 4—figure Supplement 8*, *Figure 4—figure Supplement 9*, and *Figure 4—figure Supplement 10*). In our companion paper, we explore each of these genes further and evaluate the strength of support for TSP in each gene.

Another way to obtain discordant trees is in the case of recent expansions of genes. Such expansions make it difficult to assign sequences to loci, resulting in clades where sequences (ostensibly from the same locus) do not group by species. An example of this is shown in *Figure 3*C for the NWM Class I gene MHC-G. The Class II MHC-DRB genes have also expanded, although locus assignments are somewhat clearer. *Figure 4*D shows the Class II subtree for MHC-DRB, where ape sequences (red/orange boxes) are interspersed with OWM sequences (green boxes). The MHC-

A. Sequences do not always group by gene name in the Class I multi-gene tree.

B. All MHC-F sequences group together and generally assort according to the species tree.

C. MHC-G sequences do not group together; the genes have expanded and diversified separately in the NWM.

Figure 3. The Class I exon 4 multi-gene *BEAST2* tree. The Class I multi-gene tree was constructed using exon 4 (non-PBR) sequences from Class I genes spanning the primates. **A)** For the purposes of visualization, each clade in the multi-gene tree is collapsed and labeled according to the main species group and gene content of the clade. The white labels on colored rectangles indicate the species group of origin, while the colored text to the right of each rectangle indicates the gene name. The abbreviations are defined in the species key to the right. **B)** The expanded MHC-F clade (corresponding to the clade in panel A marked by a †). **C)** The expanded NWM MHC-G clade (marked by a ∗ in panel A). In panels B and C, each tip represents a sequence and is labeled with the species of origin (white label on colored rectangle) and the sequence ID or allele name (colored text to the right of each rectangle; see *Appendix 2*). The species key is on the right hand side of panel A. Dashed branches have been shrunk to 10% of their original length (to clarify detail in the rest of the tree at this scale). OWM: old-world monkeys; NWM: new-world monkeys; Cat.: Catarrhini—apes and OWM; Pri.: Primates—apes, OWM and NWM; Mam.: mammals—primates and other outgroup mammals.

Figure 3—figure supplement 1. Color and abbreviation key/tree for all species included in this study.

Figure 3—figure supplement 2. Full, non-collapsed versions of the Class I multi-gene *BEAST2* trees.

Figure 3—figure supplement 3. Full, non-collapsed versions of the MHC-A-related focused *BEAST2* trees.

Figure 3—figure supplement 4. Full, non-collapsed versions of the MHC-B-related focused *BEAST2* trees.

Figure 3—figure supplement 5. Full, non-collapsed versions of the MHC-C-related focused *BEAST2* trees.

Figure 3—figure supplement 6. Full, non-collapsed versions of the MHC-E-related focused *BEAST2* trees.

Figure 3—figure supplement 7. Full, non-collapsed versions of the MHC-F-related focused *BEAST2* trees.

Figure 3—figure supplement 8. Full, non-collapsed versions of the MHC-G-related focused *BEAST2* trees.

Figure 3—source data 1. *GENECONV* results for the Class I focused alignments.

DRB genes are usually assigned to specific named loci, but in this tree only MHC-DRB5 sequences
group by named locus (the collapsed ape/OWM MHC-DRB5 clade is about 1/3 from the bottom of
the tree). The failure of the other named loci to group together indicates a lack of 1:1 orthology
between apes, OWM, and NWM for these genes and thus rapid evolution. This makes the MHC-
DRB genes unique among the Class II genes. We created a "focused" tree with more sequences in
order to explore the evolution of the MHC-DRB genes further, which is presented in a later section
(*Figure 4—figure Supplement 8*).

Gene conversion is a third way that gene trees might differ from the overall species tree. Gene
conversion is the unidirectional copying of a sequence onto a similar sequence (usually another
allele or a related locus), which results in two sequences being unusually similar even if they are
not related by descent. We consider this possibility in the next section.

Detection of Gene Conversion

Because the MHC contains many related genes in close proximity, gene conversion—the unidirec-
tional exchange of sequence between two similar sequences—can occur (*Chen et al., 2007*). We
used the program *GENECONV* (*Sawyer, 1999*) to infer pairs of sequences of which one has likely
been converted by the other (*Figure 3—source data 1* and *Figure 4—source data 1*). We recov-
ered known gene conversion events, such as between human allelic lineages HLA-B*38 and HLA-
B*67:02, as well as novel events, such as between gorilla allelic lineages Gogo-B*01 and Gogo-B*03
and ape/OWM lineages MHC-DQA1*01 and MHC-DQA1*05.

However, most of the tracts we detected involved many different groups of species but im-
plicated the same pair of loci. We interpreted these as gene conversion events that must have
happened a long time ago in the early history of the two genes, and they are likely to blame for
the topological differences from exon to exon among the trees. For example, in exon 2, the Class
I pseudogene MHC-K groups with MHC-G, while in exon 3 it groups with MHC-F, and in exon 4 it
groups outside of MHC-G, -J, and -A (*Figure 3*). The uncertain early branching structure we observe
in our trees may be due to these ancient gene conversion events.

The Importance of the Pseudogenization Process

Gene birth-and-death drives the evolution of a gene family as a whole. The "death" can include the
deletion of all or part of a gene from the genome or pseudogenization by means of inactivating
mutations, which can leave gene remnants behind. In Class I, we find many pseudogenes that
have been produced in this process; while countless more have undoubtedly already been deleted
from primate genomes, many full-length and fragment pseudogenes still remain. Although non-
functional, these sequences provide insight into the granular process of birth-and-death as well as
improve tree inference.

Full Class I haplotypes including the pseudogenes are known only for human, chimpanzee, go-
rilla, and macaque, and even so we do not have sequences for *all* the balanced haplotypes in each
species (*Anzai et al., 2003*; *Wilming et al., 2013*; *Shiina et al., 2017*; *Karl et al., 2023*). From these
studies, we know that few functional Class I genes are shared by apes/OWMs and NWMs, and
so far no shared pseudogenes have been found (*Lugo and Cadavid, 2015*; *Kono et al., 2014*; *Ca-
david et al., 1996*; *Maccari et al., 2017*, *2020*). Therefore, the Class I genes in the two groups have
been generated by a largely separate series of duplications, neofunctionalizations, and losses. This
means that turnover has occurred on a relatively short timescale, and understanding the pseudo-
genes within the apes and OWM can thus shed light on the evolution of the region more granularly.
These ancient remnants could provide clues as to when genes or whole blocks were duplicated,
which regions are more prone to duplication, and how the MHC may have functioned in ancestral
species.

The Class I MHC region is further divided into three polymorphic blocks—α, κ, and β—that each
contain MHC genes but are separated by well-conserved non-MHC genes. The majority of the Class
I genes are located in the α-block, which in humans includes 12 MHC genes and pseudogenes

Figure 4. The Class II exon 3 multi-gene *BEAST2* trees. The trees were constructed using all Class IIA and all Class IIB exon 3 (non-PBR) sequences across all available species. The design of this figure follows ***Figure 3***. **A)** The top tree shows the collapsed Class IIA gene tree, while the bottom tree shows the collapsed Class IIB gene tree. In this case, all collapsed clades are labeled with "Mam." for mammals, because sequences from primates and mammal outgroups assort together by gene. **B)** The expanded MHC-DPA clade (corresponding to the clade in panel A marked by a ∗). **C)** The expanded MHC-DPB clade (marked by a † in panel A). **D)** The expanded MHC-DRB clade (marked by a § in panel A). OWM: old-world monkeys; NWM: new-world monkeys; Cat.: Catarrhini—apes and OWM; Mam.: mammals—primates and other outgroup mammals.

Figure 4—figure supplement 1. Full, non-collapsed versions of the Class IIA multi-gene *BEAST2* trees.

Figure 4—figure supplement 2. Full, non-collapsed versions of the Class IIB multi-gene *BEAST2* trees.

Figure 4—figure supplement 3. Full, non-collapsed versions of the MHC-DRA focused *BEAST2* trees.

Figure 4—figure supplement 4. Full, non-collapsed versions of the MHC-DQA focused *BEAST2* trees.

Figure 4—figure supplement 5. Full, non-collapsed versions of the MHC-DPA focused *BEAST2* trees.

Figure 4—figure supplement 6. Full, non-collapsed versions of the MHC-DMA focused *BEAST2* trees.

Figure 4—figure supplement 7. Full, non-collapsed versions of the MHC-DOA focused *BEAST2* trees.

Figure 4—figure supplement 8. Full, non-collapsed versions of the MHC-DRB focused *BEAST2* trees.

Figure 4—figure supplement 9. Full, non-collapsed versions of the MHC-DQB focused *BEAST2* trees.

Figure 4—figure supplement 10. Full, non-collapsed versions of the MHC-DPB focused *BEAST2* trees.

Figure 4—figure supplement 11. Full, non-collapsed versions of the MHC-DMB focused *BEAST2* trees.

Figure 4—figure supplement 12. Full, non-collapsed versions of the MHC-DOB focused *BEAST2* trees.

Figure 4—source data 1. *GENECONV* results for the Class II focused alignments.

(**Shiina et al., 2017**). The α-block also contains a large number of repetitive elements and gene fragments belonging to other gene families, and their specific repeating pattern in humans led to the conclusion that the region was formed by successive block duplications (**Shiina et al., 1999**). Later, comparison of macaque and chimpanzee α-block haplotypes with the sequenced human haplotype bolstered this hypothesis, although the proposed series of events is not always consistent with phylogenetic data (**Kulski et al., 2005**, **2004**; **Geraghty et al., 1992**; **Hughes, 1995**; **Messer et al., 1992**; **Alexandrov et al., 2023**; **Gleimer et al., 2011**). Improving existing theories about the evolution of this block is useful for disentangling the global pattern of MHC evolution from locus- and gene-specific influences. This could help us understand how selection on specific genes has affected entire linked regions. We therefore created an α-block-focused tree involving sequences from more species than ever before in order to strengthen and update previous hypotheses about the evolution of the block, shown in **Figure 5**.

Figure 5A shows the α-block-focused tree for exon 3, with an expanded MHC-V clade. MHC-V is a fragment pseudogene containing exons 1-3 which is located near MHC-F in the α-block. Previous work disagrees on the age of this fragment, with some suggesting it was fixed relatively early while others claiming it arose from one of the more recent block duplications (**Shiina et al., 1999**; **Kulski et al., 2004**, **2005**). Our tree groups ape and OWM MHC-V together and places them as an outgroup to all of the classical and non-classical genes, including those of the NWM. Thus, the MHC-V fragment may be an ancient remnant of one of the ancestral Class I genes. We also dispute the hypothesis that MHC-V and -P (a 3'-end fragment) are since-separated pieces of the same original gene (**Horton et al., 2008**), as we found that both contain exon 3 and their exon 3 sequences clearly do not group together.

In **Figure 5**B, we zoom in on a different clade within the exon 3 tree, corresponding to the asterisk in panel A. Here we focus on MHC-U, a previously-uncharacterized fragment pseudogene containing only exon 3. Our tree groups it with a clade of human, chimpanzee, and bonobo MHC-A, suggesting it duplicated from MHC-A in the ancestor of these three species. However, it is present on both chimpanzee haplotypes and nearly all human haplotypes. Since these haplotypes likely diverged earlier in the ancestor of human and gorilla, we presume that MHC-U will be found in the gorilla when more haplotypes are sequenced. Ours is the first work to show that MHC-U is actually an MHC-A-related gene fragment.

The exon 4 α-block-focused tree is shown in **Figure 5**C. Here, we expand the clade for MHC-K, a full-length pseudogene present in apes and OWM. In humans, only MHC-K is present, but on some chimpanzee haplotypes both MHC-K and its duplicate MHC-KL are present. In gorillas, haplotypes can contain either MHC-K or -KL, and in OWM there are many copies of MHC-K as they are part of one of the basic block duplication units (**Figure 5—figure Supplement 2**) (**Karl et al., 2023**). **Figure 5**C shows that MHC-K and -KL are closely related and OWM MHC-K groups outside of both, indicating that the duplication (which also copied MHC-W, -A, and -T) occurred after the split of apes and OWM. We did not detect MHC-K or -KL sequences in either the gibbon or orangutan reference genomes during our *BLAST* search, so we cannot date this duplication event more precisely. The pseudogene may have been deleted from both genomes entirely, or it may be present on non-reference haplotypes. Sequencing of more haplotypes may help resolve the timing of this duplication event.

Both the exon 3 and exon 4 trees show a clear separation between the clade of MHC-W, -WL, -P, -T, -TL, and -OLI fragment pseudogenes and the rest of the genes (Figures 5A and C). On the chromosome, members of these two subfamilies are interleaved throughout the α-block, suggesting that both groups are old and a series of block duplications occurred. Previous work on human sequences has shown that HLA-P, -W, -T, and -OLI are related (**Alexandrov et al., 2023**; **Hughes, 1995**; **Kulski et al., 2005**). However, humans do not have orthologs of every single primate gene, so utilizing other primate sequences is critical to understanding this subfamily's evolution.

The MHC-W/WL/P/T/TL/OLI clade, marked with a † in **Figure 5**C, is expanded in panel D. We expected OWM MHC-W sequences to form a monophyletic clade either outside of all of the ape

Figure 5. Class I α-block-focused multi-gene *BEAST2* trees. The α-block-focused trees use the common backbone sequences as well as additional sequences from our custom *BLAST* search of available reference genomes. For the purposes of visualization, some clades are collapsed and labeled with the species group and gene content of the clade (colored text to the right of each rectangle). The white labels on colored rectangles indicate the species group of origin, while the colored text to the right of each rectangle indicates the gene or sequence name (see *Appendix 2*). The species abbreviations are defined in the species key at the bottom. **A)** Exon 3 α-block-focused *BEAST2* tree with expanded MHC-V clade. **B)** The expanded MHC-A/AL/OKO/U/Y clade from the exon 3 tree (corresponding to the clade in panel A marked by a ∗), focusing on MHC-U. **C)** Exon 4 α-block-focused *BEAST2* tree with expanded MHC-K/KL clade. **D)** The expanded MHC-W/WL/P/T/TL/OLI clade from the exon 4 tree (marked by a † in panel C). OWM: old-world monkeys; NWM: new-world monkeys; Cat.: Catarrhini—apes and OWM; Pri.: Primates—apes, OWM and NWM.

Figure 5—figure supplement 1. Full, non-collapsed versions of the Class I α-block pseudogene-focused *BEAST2* trees.

Figure 5—figure supplement 2. Full, non-collapsed versions of the all-pseudogene-focused *BEAST2* trees.

genes or with a single ape MHC gene, demonstrating orthology. Surprisingly, OWM MHC-W sequences instead formed four distinct clades, with one grouping with ape MHC-W/WL, one with ape MHC-P, one with ape MHC-T/TL, and one outside of all. Furthermore, based on the alleles present, each of these OWM MHC-W clades corresponds to a type of basic repeat block (as revealed by the published macaque MHC haplotype) (*Karl et al., 2023*; *Kulski et al., 2004*). The correspondence between the OWM clades and the genes' physical location lends further support to the hypothesis that macaque haplotypes were generated by tandem duplications. Furthermore, the fact that the OWM MHC-W clades each group with a different ape pseudogene suggests that there are three ape/OWM orthologous groups. Thus, there must have been three distinct MHC-W-like genes in the ape/OWM ancestor.

We also learned more about HLA-OLI, a recently-discovered MHC pseudogene found on the same insertion segment that carries HLA-Y in a small fraction of the human population. Its discoverers used only human sequences, finding HLA-OLI was most similar (88%) to HLA-P (*Alexandrov et al., 2023*). Our inclusion of non-human primate genes revealed that HLA-OLI is actually most similar in both structure and sequence to MHC-TL, a gene not found in humans (*Figure 5—figure Supplement 1*). Furthermore, since MHC-Y and -OLI are fully linked and are located in close proximity, it is likely that they duplicated as a unit. MHC-Y's similarity to MHC-AL/OKO and HLA-OLI's similarity to MHC-TL is consistent with them duplicating together from those genes, which are adjacent to each other on non-human haplotypes.

With these findings, in addition to many other observations from our trees and results from past literature (references in *Figure 6—figure Supplement 1*), we propose a new hypothesis for the evolution of the Class I α-block. *Figure 6* shows a possible evolutionary path for α-block haplotypes that could have led to the currently observed haplotypes. Haplotypes found so far in each species are at the bottom of the figure (with additional never-before-reported haplotypes from our *BLAST* search shown in *Figure 1—figure Supplement 2*). Notably, our work has revealed that MHC-V is an old fragment, three MHC-W-like genes were already established at the time of the ape/OWM ancestor, MHC-U is closely related to African ape MHC-A, and MHC-OLI is closely related to MHC-TL. Additionally, the OWM MHC-A fragment pseudogene is actually more similar to the ape MHC-A genes than to the other OWM MHC-A genes, supporting the existence of two MHC-A-like genes in the ape/OWM ancestor.

Evolution of the MHC-DRB Region

The Class I vs. Class II division reflects a major functional distinction within the MHC gene family, but even within these subfamilies evolution is not homogeneous. Among the Class II genes, there are few duplicated genes and generally only one way for the protein products to pair, e.g. MHC-DPA1 with MHC-DPB1. However, the MHC-DR genes are a notable exception to the general pattern; MHC-DRA can pair with any of the multiple functional MHC-DRB genes. In addition, MHC-DRB has many more related pseudogenes compared to the rest of the Class II genes, making the region's pattern of evolution more reminiscent of Class I (see *Figure 2* to see the varied landscape of MHC-DRB genes in different species). We explored the evolution of the MHC-DRB region in greater detail by creating focused trees with a larger set of MHC-DRB sequences.

In exon 2, which codes for the binding site, the MHC-DRB genes group mostly by name (e.g. MHC-DRB3) across apes, OWM, and NWM (*Figure 4—figure Supplement 8*A). The exon 2 tree considered alone thus suggests that the genes are orthologous across apes, OWM, and NWM—which is how the genes were named in the first place. Exon 3 does not encode the binding site and is less likely to be affected by convergent evolution; its tree is shown in *Figure 4—figure Supplement 8*B. In this tree, all NWM sequences group together (clade with blue boxes about halfway up the tree), suggesting that the genes expanded separately in NWM and apes/OWM. Additionally, OWM MHC-DRB1 and -DRB3 form their own clade (green boxes near the top of the tree) and OWM MHC-DRB4 sequences group outside of several ape and NWM clades. We see that only three MHC-DRB genes/pseudogenes (MHC-DRB5, -DRB2/6, and -DRB9) form monophyletic clades of ape and

Figure 6. Evolution of the Class I α-block. The primate evolutionary tree is shown in gray (branches not to scale). The bottom of the tree shows currently known haplotypes in each species or species group. Horizontal gray bars indicate haplotypes shared among the African apes. The history of the genes/haplotypes in the α-block is overlaid on the tree, synthesizing previous work with our own observations (see Methods and *Figure 8*). Genes are represented by colored rectangles, while haplotypes are shown as horizontal lines containing genes. MHC-F—indicating the telomeric end of the α-block—was fixed early on and is located immediately to the left on all haplotypes shown, but is not pictured due to space constraints. Dashed arrows with descriptive labels represent evolutionary events. In the upper right, the "Symbol Key" explains the icons and labels. The "Gene Relationships" panel shows the relationships between the loci shown on the tree, without the layered complexity of haplotypes and speciation events. The "MHC-A Allelic Lineages" panel shows which MHC-A allele groups are present in human, chimpanzee, and gorilla.

Figure 6—figure supplement 1. A version of *Figure 6* with references.

OWM sequences, indicating that these three are the only orthologous MHC-DRB genes. Further, none are orthologous to any particular NWM gene. Thus, the longevity of individual MHC-DRB genes in the primates appears to be less than 38 million years.

The longevity of MHC-DRB haplotypes is even shorter. Only one haplotype is shared between human and chimpanzee, and none are shared with gorilla (*de Groot et al., 2009*; *Heijmans et al., 2020*; *Hans et al., 2015*). This shows that the region is evolving even more rapidly than Class I (where haplotypes are shared among human, chimpanzee, and gorilla; *Figure 6*). These haplotypes, combined with past literature (*Figure 7—figure Supplement 1*) and our trees, allowed us to trace backward and propose a hypothesis for the evolution of the region, shown in *Figure 7*.

Limited data exists for the orangutan, and in our exon 3 tree (*Figure 4—figure Supplement 8*B) orangutan alleles do not group definitively with any other ape lineages. Furthermore, the orangutan MHC-DRB3 gene groups with orangutan MHC-DRB1, suggesting that it may not be orthologous to the African ape MHC-DRB3 gene. We also found an orangutan sequence that groups with the human HLA-DRB2 pseudogene, suggesting that this gene has an ortholog in the orangutan. Several haplotypes have been previously identified in the gibbon, but since they rely on exon 2 sequence alone, it is unclear how these alleles relate to the known ape lineages (*de Groot et al., 2017a*). Analysis of more orangutan and gibbon haplotypes will be essential for understanding how the region has evolved in the apes.

Overall, the MHC-DRB genes are not evolving in the same fashion as the rest of the Class II genes, even though they have a shared structure and function. This peculiar case illustrates that there are multiple ways to achieve a functional immune response from the same basic parts.

Differences between MHC Subfamilies

We explored the evolution of the Class I and Class II genes separately and noticed several differences between the classes. First, sequences group by gene rather than species group in the Class II gene trees (*Figure 4*, *Figure 4—figure Supplement 1*, and *Figure 4—figure Supplement 2*). Our inclusion of RefSeq sequences from distant groups of placental mammals confirms that most of the primate Class II genes have maintained orthology at least since the ancestor of placentals, 105 million years ago (*Foley et al., 2023*). In contrast, our Class I trees (*Figure 3* and *Figure 3—figure Supplement 2*) showed sequences more often grouping by species group than by gene, indicating that the genes turn over quickly and 1:1 orthology is often lost. Only non-classical MHC-F (and possibly MHC-E) are truly orthologous among the apes, OWM, and NWM, consistent with previous findings (*Piontkivska and Nei, 2003*; *Adams and Parham, 2001b*; *Sawai et al., 2004*). Additionally, our tarsier and *Strepsirrhini* sequences group outside of all *Simiiformes* Class I sequences, setting an upper bound on the maintenance of Class I orthology of 58 million years (*Kuderna et al., 2023*; *Flügge et al., 2002*).

This turnover of genes at the MHC—rapid for Class I and slower for Class II—is generally believed to be due to host-pathogen co-evolution (*Radwan et al., 2020*). Although this means the MHC genes are critically important for survival, no single gene is so vital that its role must be preserved. For example, in the apes the MHC-G gene is non-classical, but in the OWM it has been inactivated and its role largely replaced by an MHC-A-related gene called MHC-AG (*Heijmans et al., 2020*). This process of turnover ultimately results in different sets of MHC genes being used in different lineages. For instance, separate expansions generated the classical Class I genes in NWM (all called MHC-G) and the α-block genes in apes/OWM. Similarly, separate expansions generated the MHC-DRB genes of the NWM and of the apes/OWM. Aside from MHC-DRB, the Class II genes have been largely stable across the mammals, although we do see some lineage-specific expansions and contractions (*Figure 2* and *Figure 2—figure Supplement 2*).

Class I and Class II also differ in their degree of gene conversion. Our *GENECONV* analysis revealed two types of gene conversion events: 1) specific, more-recent events involving paralogous genes or particular allelic lineages and 2) broad-scale, very-old events involving two dissimilar loci (*Figure 3—source data 1* and *Figure 4—source data 1*). We discovered far more "specific" events

Figure 7. Evolution of MHC-DRB. The bottom of the tree shows current haplotypes in each species or species group; human, chimpanzee, gorilla, and old-world monkey haplotypes are well characterized, while orangutan, gibbon, and new-world monkey haplotypes are partially known. The history of the genes/haplotypes in the MHC-DRB region is overlaid on the tree, synthesizing previous work with our own observations (see Methods and *Figure 8*). The rest of the figure design follows that of *Figure 6*.

Figure 7—figure supplement 1. A version of *Figure 7* with references.

in Class I, while "broad-scale" events were predominant in Class II. This could reflect the different age of these gene groups: Class I genes turn over more rapidly and allelic lineages are less diverged from each other, making gene conversion more likely. In contrast, Class II genes have much longer-lived allelic lineages, potentially explaining why we mainly picked up older events in the Class II *GENECONV* analysis.

Even within a class, evolutionary patterns are not homogeneous. The non-classical vs. classical distinction is one functionally meaningful way to partition the genes. The classical genes perform peptide presentation to T-cells, making them direct targets of host-pathogen co-evolution. In contrast, the non-classical genes are involved in innate immune surveillance or niche roles, and may be less directly affected by this co-evolution. In our trees, sequences from non-classical genes of both classes often group by gene with topology matching the species tree, while sequences from classical genes do neither (*Figure 3—figure Supplement 2*, *Figure 4—figure Supplement 1*, and *Figure 4—figure Supplement 2*). This shows that classical genes experience more turnover and are more often affected by long-term balancing selection or convergent evolution. Ultimately, selection acts upon functional differences between classical and non-classical genes in a manner that is largely independent of whether they belong to Class I or Class II.

Discussion

The MHC proteins serve diverse roles in innate and adaptive immunity (*Adams and Luoma, 2013*). They are critically important to infection resistance, autoimmune-disease susceptibility, and organ transplantation success, and can provide insight into human evolution, inform disease studies, and improve upon non-human-primate disease models (*Kennedy et al., 2017*). Despite their varied functions, all Class I and Class II MHC genes are derived from a common ancestor, allowing us to compare genes to learn more about the evolution of the gene family as a whole (*Hansen et al., 2007*; *Kupfermann et al., 1999*; *Kaufman, 2022*; *Adams and Luoma, 2013*). A few ~20-year-old studies addressed the overall evolution of the MHC gene family via multi-gene alignment and phylogenetics, but the trees had many polytomies (*Adams and Parham, 2001b*; *Sawai et al., 2004*; *Cardenas et al., 2005*; *Piontkivska and Nei, 2003*; *Takahashi et al., 2000*). Since then, most work has focused on particular genes or small sets of species, meaning our knowledge of primate MHC evolution is scattered across hundreds of papers (*Urvater et al., 2000*; *Van Der Wiel et al., 2013*; *Geller et al., 2002*; *Hans et al., 2017*; *Maibach et al., 2017*; *Wroblewski et al., 2017, 2019*; *Lafont et al., 2004*; *Flügge et al., 2002*; *Go et al., 2003, 2005*; *Shiina et al., 2017*; *Abi-Rached et al., 2010*; *Gleimer et al., 2011*; *de Groot et al., 2015*; *De Groot et al., 2012*; *de Groot et al., 2022*; *Cao et al., 2015*; *Otting et al., 2020*; *Fukami-Kobayashi et al., 2005*; *Lugo and Cadavid, 2015*; *Averdam et al., 2011*; *Figueroa et al., 1994*; *Doxiadis et al., 2012*; *Buckner et al., 2021*; *Doxiadis et al., 2006*; *Diaz et al., 2000*; *Gongora et al., 1997*; *Kasahara et al., 1992*; *de Groot et al., 2009*; *Satta et al., 1996*). In this project, we revisited primate MHC evolution with more data from a wider range of species and a coherent analysis framework. We confirm and unify past findings, as well as contribute many new insights into the evolution of this complex family.

We found that the Class I genes turn over rapidly, with only the non-classical gene MHC-F being clearly orthologous across the *Simiiformes*. In the rest of the Class I α-block, genes expanded entirely separately in the ape/OWM and NWM lineages. This process of expansion generated many full-length and fragment pseudogenes, which we found were equally important as the functional genes to understanding the evolution of the region as a whole. Notably, we found that MHC-U is an MHC-A-related pseudogene, MHC-V is not closely related to MHC-P, and that there were at least three genes of the MHC-W/P/T/OLI family present in the ape/OWM ancestor. Generally, Class II genes do not turn over as rapidly, although there were exceptions. The classical MHC-DRB genes were even shorter-lived than the Class I genes, with most human genes lacking 1:1 orthologs beyond the great apes. We also found that the classical MHC-DQA and -DQB genes likely expanded separately in the ape/OWM and NWM lineages. In contrast, the classical MHC-DPA and -DPB genes were orthologous across the *Simiiformes*, and the non-classical Class II genes were 1:1 orthologous

across most of the mammals. In both Class I and Class II, classical genes turned over more rapidly than non-classical genes and their trees exhibited more deviations from the expected species tree. Overall, our treatment of the genes as related entities instead of distinct cases helped us understand shared patterns of evolution across classes and species groups.

One concern when discussing gene families is the relative importance of birth-and-death and concerted evolution by gene conversion (*Gu and Nei, 1999a*; *Nei and Rooney, 2005*; *Klein et al., 2007*; *Bergström and Gyllensten, 1995*; *Gyllensten et al., 1991*; *Nei et al., 1997*). Gene conversion can cause adjacent small sequence tracts to have wildly different evolutionary histories, making it difficult to interpret a tree constructed from larger regions. Our trees reveal different topologies depending on exon and our *GENECONV* analysis pulled out several different sequence pairs, revealing that gene conversion has played a significant role in the evolution of the MHC genes. With this in mind, comparing trees across exons helps us interpret the overall trees and strengthens our conclusions. Neither birth-and-death nor concerted evolution can be ignored when discussing gene families.

The MHC region is difficult to assemble owing to the large number of related genes, extreme polymorphism, and abundant repetitive regions. Nonetheless, several committed researchers have dedicated effort to sequencing and mapping the region in different species (*Wilming et al., 2013*; *Anzai et al., 2003*; *Karl et al., 2023*; *Okano et al., 2020*; *Hammer et al., 2020*; *Liao et al., 2023*). However, we now know that haplotype variation is just as important as nucleotide variation to maintaining MHC diversity, and haplotypes are surprisingly short-lived (*de Groot et al., 2015*, *2017a*; *Gleimer et al., 2011*; *Hans et al., 2015*; *Heijmans et al., 2020*). Therefore, more research effort should be dedicated to fully characterizing the breadth of MHC haplotypes in different species. This is a difficult problem owing to the plethora of repetitive elements and recently-duplicated genes in the region, and long-read sequencing will be invaluable for parsing these complex haplotypes. Not only will this improve our understanding of health and disease in each species, but it will also help us answer evolutionary questions with better precision.

We hope that our extensive set of trees incorporating classical genes, non-classical genes, pseudogenes, gene fragments, and alleles of medical interest across a wide range of species will provide context to future researchers. This work will provide a jumping-off-point for further exploration of the evolutionary processes affecting different subsets of the gene family as well as necessary context for studies of particular alleles or genes in disease.

Methods and Materials

Data Collection

We downloaded MHC allele nucleotide sequences for all human and non-human genes from the IPD Database (collected January 2023) (*Barker et al., 2023*; *Maccari et al., 2017*, *2020*; *Robinson et al., 2024*). To supplement the alleles available in the database, we also collected nucleotide sequences from NCBI using the Entrez E-utilities with query "histocompatibility AND txidX AND alive[prop]", where X is a taxon of interest. This resulted in a very large collection of sequences from a large number of species. While Class II genes were generally assigned to loci, most Class I sequences had ambiguous or no locus assignments. Therefore, we performed a refined search for additional sequences by running *BLAST* on the available primate reference genomes.

BLAST Search

For the reference genomes, we downloaded human chromosome 6 (GenBank accession `CM000668.2`), chimpanzee chromosome 5 (`CM054439.2`), bonobo chromosome 5 (`CM055477.2`), gorilla chromosome 5 (`CM055451.2`), Sumatran orangutan chromosome 5 (`CM054684.2`), Bornean orangutan chromosome 5 (`CM054635.2`), pileated gibbon linkage group LG22 (`CM038537.1`), siamang chromosome 23 (`CM054531.2`), Northern white-cheeked gibbon chromosome 22a (`CM016966.1`), olive baboon chromosome 6 (`CM018185.2`), Guinea baboon chromosome 6 (`CM053423.1`), gelada chromosome

4 (`CM009953.1`), Tibetan macaque chromosome 4 (`CM045091.1`), crab-eating macaque chromosome
4 (`CP141358.1`), Formosan rock macaque chromosome 4 (`CM049490.1`), mantled guereza chromo-
some 5 (`CM058078.1`), snub-nosed monkey chromosome 4 (`CM017354.1`), cotton-top tamarin chro-
mosome 4 (`CM063172.1`), golden-handed tamarin linkage group LG04 (`CM038394.1`), common mar-
moset chromosome 4 (`CM021918.1`), coppery titi chromosome 3 (`CM080817.1`), gray mouse lemur
chromosome 6 (`CM007666.1`), black-and-white ruffed lemur chromosome 6 (`CM052441.1`), mongoose
lemur chromosome 15 (`CM052867.1`), ring-tailed lemur chromosome 2 (`CM036473.1`), Bengal slow
loris linkage group LG08 (`CM043617.1`), Sunda slow loris chromosome 9 (`CM050145.1`), Philippine
flying lemur chromosome 5 (`CM050031.1`), and mouse chromosome 17 (`CM001010.3`).

To create the *BLAST* databse, we first compiled all nucleotide MHC sequences from the IPD-
MHC and IPD-IMGT/HLA databases into three `fasta` files: one containing the Class I sequences, one
containing the Class II sequences, and one containing MHC-DRB9 sequences. We then constructed
three custom databases from these sets of sequences using the `makeblastdb` command in *BLAST*
version 2.11.0 (*Camacho et al., 2009*).

We then queried each of the three custom databases using the above reference genomes and
screened the hits manually. In most cases, we were able to identify loci unambiguously, resulting
in several newly-reported haplotypes (*Figure 1—figure Supplement 2* and *Figure 2—figure Supple-*
ment 2). The discovery of various genes in various species also allowed us to fill in gaps in *Figure 1*
and *Figure 2*.

Sequence Selection

Because *BEAST2* is computationally limited by the number of sequences, it was necessary to prior-
itize certain sequences. To do this, we (very roughly) aligned as many exon 2 and 3 sequences as
possible (from both NCBI RefSeq and the IPD database) using *MUSCLE* (*Edgar, 2004*) with default set-
tings. We then constructed UPGMA trees in R to visualize the sequences. We preferentially selected
sequences that were 1) in primate species not represented by the IPD database or 2) grouped with
genes not well represented by the IPD database, and which were not similar/identical to other se-
quences. We also included several non-primate species to provide context and explore orthology
beyond the primates. After choosing sequences with this preliminary screening method, we col-
lected the full-length sequences for inclusion in further analyses. We limited sequences to one per
species-gene pair for building the Class I, Class IIA, and Class IIB multi-gene trees (lists of alleles
provided as Supplementary Files).

For Class I, we then re-aligned all genes together for each exon separately using *MUSCLE* (*Edgar,*
2004) with default settings (and manually adjusted). For Class II, alleles for each gene group (MHC-
DMA, -DMB, -DOA, -DOB, -DPA, -DPB, -DQA, -DQB, -DRA, and -DRB) were aligned separately for
each exon using *MUSCLE* (*Edgar, 2004*) with default settings (and manually adjusted). Since some
Class II genes are too far diverged from one another to be reliably aligned automatically, the nu-
cleotide alignments were then combined manually based on published amino acid alignments
(*Radley et al., 1994*; *Dijkstra et al., 2013*; *Dijkstra and Yamaguchi, 2019*; *Cuesta et al., 2006*; *Chen*
et al., 2006; *Chazara et al., 2011*). For Class IIA, exons 4 and 5 were concatenated together be-
fore this manual combination process because some analogous sites between genes are located
across exons. For the same reason, exons 5 and 6 were concatenated together for Class IIB before
combining. This produced three multi-gene alignments: Class I, Class IIA, and Class IIB.

We also aligned a larger set of sequences for each gene group to create our "focused" trees
that each zoomed in on a different subtree of the multi-gene trees. Details for this are located in
the Methods of our companion paper (*Fortier and Pritchard, 2024*).

Bayesian Phylogenetic Analysis

582 We constructed phylogenetic trees using *BEAST2* (*Bouckaert et al., 2014*, *2019*) with package *sub-*
583 *stBMA* (*Wu et al., 2013*). *SubstBMA* implements a spike-and-slab mixture model that simultaneously
584 estimates the phylogenetic tree, the number of site partitions, the assignment of sites to partitions,

the nucleotide substitution model, and a rate multiplier for each partition. Since we were chiefly interested in the partitions and their rate multipliers, we used the RDPM model as described by *Wu et al.* (*2013*). In the RDPM model, the number of nucleotide substitution model categories is fixed to 1, so that all sites, regardless of rate partition, share the same estimated nucleotide substitution model. This reduces the number of parameters to be estimated and ensures that only evolutionary rates vary across site partitions, reducing overall model complexity. We used an uncorrelated lognormal relaxed molecular clock because we wanted evolutionary rates to be able to vary among branches.

Priors

For the Dirichlet process priors, we used the informative priors constructed by *Wu et al.* (*2013*) for their mammal dataset. This is appropriate because they include several of the same species and their mammals span approximately the same evolutionary time that we consider in our study. We also use their same priors on tree height, base rate distribution, and a Yule process coalescent prior. We did not specify a calibration point—a time-based prior on a node—because we did not expect our sequences to group according to the species tree.

Running *BEAST2*

We ran *BEAST2* on various subsets of the three alignments. Considering exons separately helped to minimize the effects of recombination on the tree, while also allowing us to compare and contrast tree topologies for exons encoding the binding site vs. exons encoding the other domains. For Class I, we repeated the analysis for 1) exon 2 only (PBR), 2) exon 3 only (PBR), and 3) exon 4 only (non-PBR). For Class IIA, we used 1) exon 2 only (PBR) and 2) exon 3 only (non-PBR). For Class IIB, we analyzed 1) exon 2 only (PBR) and 2) exon 3 only (non-PBR). In the following, each "analysis" refers to a collection of *BEAST2* runs using a particular subset of either the Class I, Class IIA, or Class IIB alignment. The procedure is exactly the same for the "focused" trees, which each focus on a particular gene group within the Class I, Class IIA, or Class IIB alignment. More detail about the generation of the focused trees is located in the Methods of our companion paper (*Fortier and Pritchard, 2024*).

The xml files we used to run *BEAST2* were based closely on those used for the mammal dataset with the RDPM model and uncorrelated relaxed clock in *Wu et al.* (*2013*) (https://github.com/jessiewu/substBMA/blob/master/examples/mammal/mammal_rdpm_uc.xml). Running a model with per-site evolutionary rate categories and a relaxed clock means there are many parameters to estimate. Along with the large number of parameters, highly-polymorphic and highly-diverged sequences make it difficult for *BEAST2* to explore the state space. Thus, we undertook considerable effort to ensure good mixing and convergence of the chains. First, we employed coupled MCMC for all analyses. Coupled MCMC is essentially the same as the regular MCMC used in *BEAST2*, except that it uses additional "heated" chains with increased acceptance probabilities that can traverse unfavorable intermediate states and allow the main chain to move away from an inferior local optimum (*Müller and Bouckaert, 2020*). Using coupled MCMC both speeds up *BEAST2* runs and improves mixing and convergence. We used four heated chains for each run with a delta temperature of 0.025. Second, we ran each *BEAST2* run for 40,000,000 states, discarding the first 4,000,000 states as burn-in and sampling every 10,000 states. Third, we ran at least eight independent replicates of each analysis. The replicates use the exact same alignment and coupled MCMC settings, but explore state space independently and thus are useful for improving the effective sample size of tricky parameters. As recommended by *BEAST2*, we examined all replicates in *Tracer* version 1.7.2 (*Rambaut et al., 2018*) to ensure that they were sampling from the same parameter distributions and had reached convergence. We excluded replicates for which this was not true, as these chains were probably stuck in suboptimal state space. Additionally, *Tracer* provides estimates of the effective sample size (ESS) for the combined set of states from all chosen replicates, and we required that the combined ESS be larger than 100 for all parameters. If there were fewer than 4 acceptable

₆₃₄ replicates or if the ESS was below 100 for any parameter, we re-ran more independent replicates
₆₃₅ of the analysis until these requirements were satisfied. We obtained between 7 and 14 acceptable
₆₃₆ replicates (median 8) per analysis for the Class I, Class IIA, and Class IIB runs.

₆₃₇   For some analyses, computational limitations prevented *BEAST2* from being able to reach 40,000,000
₆₃₈ states. In these situations, more replicates (of fewer states) were usually required to achieve good
₆₃₉ mixing and convergence. Regardless of how far these *BEAST2* runs got, the first 4,000,000 states
₆₄₀ from each run were still discarded as burn-in even though this represented more than 10% of
₆₄₁ states. The xml files required to run all our analyses are provided as Supplementary Files.

₆₄₂   This extremely stringent procedure ensured that all of the replicates were exploring the same
₆₄₃ parameter space and were converging upon the same global optimum, allowing the ≥ 4 inde-
₆₄₄ pendent runs to be justifiably combined. We combined the acceptable replicates (discarding the
₆₄₅ first 4,000,000 states as burn-in) using *LogCombiner* version 2.6.7 (***Drummond and Rambaut, 2007***),
₆₄₆ which aggregates the results across all states. We then used the combined results for downstream
₆₄₇ analyses.

₆₄₈ **Phylogenetic Trees**
₆₄₉ After combining acceptable replicates, we obtained 17,927 - 28,384 phylogenies per gene/sequence
₆₅₀ subset for the Class I, Class IIA, and Class IIB trees (mean 25,154). We used *TreeAnnotator* version
₆₅₁ 2.6.3 (***Drummond and Rambaut, 2007***) to summarize each set of possible trees as a maximum
₆₅₂ clade credibility tree, which is the tree that maximizes the sum of posterior clade probabilities.
₆₅₃ Since *BEAST2* samples trees from the posterior, one could in principle reduce the large set of trees
₆₅₄ to a smaller 95% credible set of trees representing the "true" tree (***BEA, 2024***). However, given the
₆₅₅ high complexity of the model space, all our posterior trees were unique, meaning this was not
₆₅₆ possible in practice. Throughout this paper, we rely on summary trees for our observations.

₆₅₇ **Integration with Literature**
₆₅₈ Hundreds of authors have contributed to the study of MHC evolution, and their myriad published
₆₅₉ results played a key role in this project. ***Figure 8*** illustrates our approach to this project, including
₆₆₀ how we used existing literature and how we divided results among this paper and its companion
₆₆₁ (***Fortier and Pritchard, 2024***). We first constructed large multi-gene trees encompassing all Class
₆₆₂ I, Class IIA, and Class IIB genes. These provided a backbone for us to investigate subtrees in more
₆₆₃ depth, adding more sequences and more species to construct "focused trees" for each gene group.
₆₆₄ These, in combination with the literature, allowed us to create hypotheses about the evolution of
₆₆₅ the Class I α-block (***Figure 6***) and Class II MHC-DRB region (***Figure 7***).

₆₆₆ **Gene Conversion**
₆₆₇ We inferred gene conversion fragments using *GENECONV* version 1.81a (***Sawyer, 1999***) on each fo-
₆₆₈ cused alignment. It is generally advisable to use only synonymous sites when running the program
₆₆₉ on a protein-coding alignment, since silent sites within the same codon position are likely to be cor-
₆₇₀ related. However, the extreme polymorphism in these MHC genes meant there were too few silent
₆₇₁ sites to use in the analysis. Thus, we considered all sites but caution that this could slightly overes-
₆₇₂ timate the lengths of our inferred conversion tracts. For each alignment, we ran *GENECONV* with
₆₇₃ options `ListPairs`, `Allouter`, `Numsims=10000`, and `Startseed=310`. We collected all inferred "Global
₆₇₄ Inner" (GI) fragments with sim_pval< 0.05 (this is pre-corrected for multiple comparisons by the
₆₇₅ program). GI fragments indicate a stretch of similar sequence shared by two otherwise-dissimilar
₆₇₆ sequences in the alignment. This suggests that a gene conversion event occurred between the
₆₇₇ ancestors of the two sequences.

₆₇₈   Many of the thousands of GI hits were redundant, involving very-closely-related alleles, slightly
₆₇₉ different fragment bounds, or even a wide range of species all implicating the same gene. We
₆₈₀ manually grouped and summarized these hits for ***Figure 3—source data 1*** and ***Figure 4—source***
₆₈₁ ***data 1***. The "start" and "end" columns indicate the smallest start and largest end position (along

Figure 8. *BEAST2* trees provide insight into MHC gene and allele relationships. We first created multi-gene Bayesian phylogenetic trees using sequences from all genes and species, separated into Class I, Class IIA, and Class IIB groups. We then focused on various subtrees of the multi-gene trees by adding more sequences for each subtree and running *BEAST2* using only sequences from that group (in addition to the "backbone" sequences common to all trees). Our trees gave us insight into both overall gene relationships (this paper) and allele relationships within gene groups (see our companion paper, *Fortier and Pritchard* (*2024*)).

682 the alignment) for the group of redundant hits, and the sequences involved are summarized as
683 specifically as possible.

Acknowledgments

685 We acknowledge support from NIH grants R01 HG011432 and R01 HG008140. This material is
686 based upon work supported by the National Science Foundation Graduate Research Fellowship un-
687 der Grant No. DGE-1656518. We appreciate helpful comments from Jeffrey Spence, the Pritchard
688 lab, and the reviewers of the previous version of our companion paper, which jumpstarted this
689 project.

References

SUMMARIZING POSTERIOR TREES. Centre for Computational Evolution; 2024. https://www.beast2.org/
summarizing-posterior-trees/.

Abi-Rached L, Kuhl H, Roos C, ten Hallers B, Zhu B, Carbone L, de Jong PJ, Mootnick AR, Knaust F, Reinhardt R,
Parham P, Walter L. A Small, Variable, and Irregular Killer Cell Ig-Like Receptor Locus Accompanies the
Absence of MHC-C and MHC-G in Gibbons . The Journal of Immunology. 2010; 184(3):1379–1391. doi:
10.4049/jimmunol.0903016.

Adams EJ, Parham P. Genomic analysis of common chimpanzee major histocompatibility complex class I genes.
Immunogenetics. 2001; 53(3):200–208. doi: 10.1007/s002510100318.

Adams EJ, Luoma AM. The adaptable major histocompatibility complex (MHC) fold: Structure and function
of nonclassical and mhc class I-like molecules. Annual Review of Immunology. 2013; 31:529–561. doi:
10.1146/annurev-immunol-032712-095912.

Adams EJ, Parham P. Species-specific evolution of MHC class I genes in the higher primates. Immunological
Reviews. 2001; 183:41–64. doi: 10.1034/j.1600-065X.2001.1830104.x.

Alexandrov N, Wang T, Blair L, Nadon B, Sayer D. HLA-OLI: A new MHC class I pseudogene and HLA-Y are
located on a 60 kb indel in the human MHC between HLA-W and HLA-J. Hla. 2023; 102(5):599–606. doi:
10.1111/tan.15180.

Anderson JL, Sandstrom K, Smith WR, Wetzel M, Klenchin VA, Evans DT. MHC Class I Ligands of Rhesus
Macaque Killer Cell Ig-like Receptors. The Journal of Immunology. 2023; 210(11):1815–1826. doi: 10.4049/jim-
munol.2200954.

Anzai T, Shiina T, Kimura N, Yanagiya K, Kohara S, Shigenari A, Yamagata T, Kulski JK, Naruse TK, Fujimori Y,
Fukuzumi Y, Yamazaki M, Tashiro H, Iwamoto C, Umehara Y, Imanishi T, Meyer A, Ikeo K, Gojobori T, Bahram
S, et al. Comparative sequencing of human and chimpanzee MHC class I regions unveils insertions/deletions
as the major path to genomic divergence. Proceedings of the National Academy of Sciences of the United
States of America. 2003; 100(13):7708–7713. doi: 10.1073/pnas.1230533100.

Arden B, Klein J. Biochemical comparison of major histocompatibility complex molecules from different sub-
species of Mus musculus: Evidence for trans-specific evolution of alleles. Proceedings of the National
Academy of Sciences of the United States of America. 1982; 79(7):2342–2346. doi: 10.1073/pnas.79.7.2342.

Averdam A, Kuschal C, Otto N, Westphal N, Roos C, Reinhardt R, Walter L. Sequence analysis of the grey
mouse lemur (Microcebus murinus) MHC class II DQ and DR region. Immunogenetics. 2011; 63(2):85–93.
doi: 10.1007/s00251-010-0487-3.

Barker DJ, Maccari G, Georgiou X, Cooper MA, Flicek P, Robinson J, Marsh SGE. The IPD-IMGT/HLA Database.
Nucleic Acids Research. 2023; 51(November 2022):1053–1060.

Benton MJ, Donoghue PCJ, Asher RJ, Friedman M, Near TJ, Vinther J. Constraints on the timescale of animal
evolutionary history. Palaeontologia Electronica. 2015; 18(1):1–107. doi: 10.26879/424.

Bergström T, Gyllensten U. Evolution of Mhc Class II Polymorphism: The Rise and Fall of Class II Gene Function
in Primates. Immunological Reviews. 1995; 143(1):13–31. doi: 10.1111/j.1600-065X.1995.tb00668.x.

Biassoni R, Malnati MS. Human natural killer receptors, co-receptors, and their ligands. Current Protocols in
Immunology. 2018; e47:1–52. doi: 10.1002/0471142735.im1410s84.

Bondarenko GI, Burleigh DW, Durning M, Breburda EE, Grendell RL, Golos TG. Passive Immunization against
the MHC Class I Molecule Mamu-AG Disrupts Rhesus Placental Development and Endometrial Responses.
The Journal of Immunology. 2007; 179(12):8042–8050. doi: 10.4049/jimmunol.179.12.8042.

Bondarenko GI, Dambaeva SV, Grendell RL, Hughes AL, Durning M, Garthwaite MA, Golos TG. Characterization
of cynomolgus and vervet monkey placental MHC class i expression: Diversity of the nonhuman primate AG
locus. Immunogenetics. 2009; 61(6):431–442. doi: 10.1007/s00251-009-0376-9.

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu CH, Xie D, Suchard MA, Rambaut A, Drummond AJ. BEAST 2:
A Software Platform for Bayesian Evolutionary Analysis. PLoS Computational Biology. 2014; 10(4):1–6. doi:
10.1371/journal.pcbi.1003537.

Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled J, Jones G, Küh-
nert D, De Maio N, Matschiner M, Mendes FK, Müller NF, Ogilvie HA, Du Plessis L, Popinga A, Rambaut A,
Rasmussen D, Siveroni I, Suchard MA, et al. BEAST 2.5: An advanced software platform for Bayesian evolu-
tionary analysis. PLoS Computational Biology. 2019; 15(4):1–28. doi: 10.1371/journal.pcbi.1006650.

Boyson JE, McAdam SN, Gallimore A, Golos TG, Liu X, Gotch FM, Hughes AL, Watkins DI. The MHC E locus in
macaques is polymorphic and is conserved between macaques and humans. Immunogenetics. 1995; 41(2-
3):59–68. doi: 10.1007/BF00182314.

Buckner JC, Jack KM, Melin AD, Schoof VAM, Gutiérrez-Espeleta GA, Lima MGM, Lynch JW. Major histocompat-
ibility complex class II DR and DQ evolution and variation in wild capuchin monkey species (Cebinae). PLoS
ONE. 2021; 16(8):e0254604. doi: 10.1371/journal.pone.0254604.

Budde ML, Wiseman RW, Karl JA, Hanczaruk B, Simen BB, O'Connor DH. Characterization of Mauritian cynomol-
gus macaque major histocompatibility complex class I haplotypes by high-resolution pyrosequencing. Im-
munogenetics. 2010; 62(11-12):773–780. doi: 10.1007/s00251-010-0481-9.

Budeus B, Álvaro-Benito M, Crivello P. HLA-DM and HLA-DO interplay for the peptide editing of HLA class
II in healthy tissues and leukemia. Best Practice and Research: Clinical Haematology. 2024; 37(2). doi:
10.1016/j.beha.2024.101561.

Buniello A, MacArthur JAL, Cerezo M, Harris LW, Hayhurst J, Malangone C, McMahon A, Morales J, Mountjoy E, Sollis E, Suveges D, Vrousgou O, Whetzel PL, Amode R, Guillen JA, Riat HS, Trevanion SJ, Hall P, Junkins H, Flicek P, et al. The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. Nucleic Acids Research. 2019 1; 47(D1):D1005–D1012. https://academic.oup.com/nar/article/47/D1/D1005/5184712, doi: 10.1093/nar/gky1120.

Cadavid LF, Hughes AL, Watkins DI. MHC class I-processed pseudogenes in New World primates provide evidence for rapid turnover of MHC class I genes. Journal of immunology (Baltimore, Md : 1950). 1996; 157(6):2403–9. http://www.ncbi.nlm.nih.gov/pubmed/8805639.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. BLAST+: Architecture and applications. BMC Bioinformatics. 2009; 10:1–9. doi: 10.1186/1471-2105-10-421.

Cao YH, Fan JW, Li AX, Liu HF, Li LR, Zhang CL, Zeng L, Sun ZZ. Identification of MHC I class genes in two Platyrrhini species. American Journal of Primatology. 2015; 77(5):527–534. doi: 10.1002/ajp.22372.

Cardenas PP, Suarez CF, Martinez P, Patarroyo ME, Patarroyo MA. MHC class I genes in the owl monkey: Mosaic organisation, convergence and loci diversity. Immunogenetics. 2005; 56(11):818–832. doi: 10.1007/s00251-004-0751-5.

Carlini F, Ferreira V, Buhler S, Tous A, Eliaou JF, René C, Chiaroni J, Picard C, Di Cristofaro J. Association of HLA-A and non-classical HLA class i alleles. PLoS ONE. 2016; 11(10):1–17. doi: 10.1371/journal.pone.0163570.

Carter AM. Unique aspects of human placentation. International Journal of Molecular Sciences. 2021; 22(15). doi: 10.3390/ijms22158099.

Chazara O, Tixier-Boichard M, Morin V, Zoorob R, Bed'Hom B. Organisation and diversity of the class II DM region of the chicken MHC. Molecular Immunology. 2011; 48(9-10):1263–1271. http://dx.doi.org/10.1016/j.molimm.2011.03.009, doi: 10.1016/j.molimm.2011.03.009.

Chen JM, Cooper DN, Chuzhanova N, Férec C, Patrinos GP. Gene conversion: Mechanisms, evolution and human disease. Nature Reviews Genetics. 2007; 8(10):762–775. doi: 10.1038/nrg2193.

Chen SL, Zhang YX, Xu MY, Ji XS, Yu GC, Dong CF. Molecular polymorphism and expression analysis of MHC class II B gene from red sea bream (Chrysophrys major). Developmental and Comparative Immunology. 2006; 30(4):407–418. doi: 10.1016/j.dci.2005.06.001.

Cuesta A, Ángeles Esteban M, Meseguer J. Cloning, distribution and up-regulation of the teleost fish MHC class II alpha suggests a role for granulocytes as antigen-presenting cells. Molecular Immunology. 2006; 43(8):1275–1285. doi: 10.1016/j.molimm.2005.07.004.

De Groot NG, Otting N, Robinson J, Blancher A, Lafont BAP, Marsh SGE, O'Connor DH, Shiina T, Walter L, Watkins DI, Bontrop RE. Nomenclature report on the major histocompatibility complex genes and alleles of Great Ape, Old and New World monkey species. Immunogenetics. 2012; 64(8):615–631. doi: 10.1007/s00251-012-0617-1.

Demuth JP, Bie TD, Stajich JE, Christiani N, Hahn MW. The evolution of mammalian gene families. PLoS ONE. 2006; 1(1). doi: 10.1371/journal.pone.0000085.

Dendrou CA, Petersen J, Rossjohn J, Fugger L. HLA variation and disease. Nature Reviews Immunology. 2018; 18(5):325–339. http://dx.doi.org/10.1038/nri.2017.143, doi: 10.1038/nri.2017.143.

Diaz D, Naegeli M, Rodriguez R, Nino-Vasquez JJ, Moreno A, Patarroyo ME, Pluschke G, Daubenberger CA. Sequence and diversity of MHC DQA and DQB genes of the owl monkey Aotus nancymaae. Immunogenetics. 2000; 51(7):528–537. doi: 10.1007/s002510000189.

Dijkstra JM, Grimholt U, Leong J, Koop BF, Hashimoto K. Comprehensive analysis of MHC class II genes in teleost fish genomes reveals dispensability of the peptide-loading DM system in a large part of vertebrates. BMC Evolutionary Biology. 2013; 13(1):1–14. doi: 10.1186/1471-2148-13-260.

Dijkstra JM, Yamaguchi T. Ancient features of the MHC class II presentation pathway, and a model for the possible origin of MHC molecules. Immunogenetics. 2019; 71(3):233–249. doi: 10.1007/s00251-018-1090-2.

Douillard V, Castelli EC, Mack SJ, Hollenbach JA, Gourraud PA, Vince N, Limou S. Approaching Genetics Through the MHC Lens: Tools and Methods for HLA Research. Frontiers in Genetics. 2021; 12(December):1–13. doi: 10.3389/fgene.2021.774916.

Doxiadis GGM, Hoof I, De Groot N, Bontrop RE. Evolution of HLA-DRB genes. Molecular Biology and Evolution. 2012; 29(12):3843–3853. doi: 10.1093/molbev/mss186.

Doxiadis GGM, Rouweler AJM, De Groot NG, Louwerse A, Otting N, Verschoor EJ, Bontrop RE. Extensive sharing of MHC class II alleles between rhesus and cynomolgus macaques. Immunogenetics. 2006; 58(4):259–268. doi: 10.1007/s00251-006-0083-8.

Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling trees. BMC Evolutionary Biology. 2007; 7(1):214. http://bmcevolbiol.biomedcentral.com/articles/10.1186/1471-2148-7-214, doi: 10.1186/1471-2148-7-214.

Ebert D, Fields PD. Host–parasite co-evolution and its genomic signature. Nature Reviews Genetics. 2020 12; 21(12):754–768. https://www.nature.com/articles/s41576-020-0269-1, doi: 10.1038/s41576-020-0269-1.

Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research. 2004 3; 32(5):1792–1797. https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkh340, doi: 10.1093/nar/gkh340.

Figueroa F, O'hUigin C, Tichy H, Klein J. The origin of the primate Mhc-DRB genes and allelic lineages as deduced from the study of prosimians. The Journal of Immunology. 1994; 152(9):4455–4465. doi: 10.4049/jimmunol.152.9.4455.

Flajnik MF, Kasahara M. Comparative genomics of the MHC: Glimpses into the evolution of the adaptive immune system. Immunity. 2001; 15(3):351–362. doi: 10.1016/S1074-7613(01)00198-4.

Flügge P, Zimmermann E, Hughes AL, Günther E, Walter L. Characterization and phylogenetic relationship of prosimian MHC class I genes. Journal of Molecular Evolution. 2002; 55(6):768–775. doi: 10.1007/s00239-002-2372-7.

Foley NM, Mason VC, Harris AJ, Bredemeyer KR, Damas J, Lewin HA, Eizirik E, Gatesy J, Karlsson EK, Lindblad-Toh K, Springer MS, Murphy WJ, Andrews G, Armstrong JC, Bianchi M, Birren BW, Bredemeyer KR, Breit AM, Christmas MJ, Clawson H, et al. A genomic timescale for placental mammal evolution. Science. 2023; 380(6643). doi: 10.1126/science.abl8189.

Fortier AL, Pritchard JK. Ancient Trans-Species Polymorphism at the Major Histocompatibility Complex in Primates. TBD. 2024; .

Friedman R, Hughes AL. The temporal distribution of gene duplication events in a set of highly conserved human gene families. Molecular Biology and Evolution. 2003; 20(1):154–161. doi: 10.1093/molbev/msg017.

Fukami-Kobayashi K, Shiina T, Anzai T, Sano K, Yamazaki M, Inoko H, Tateno Y. Genomic evolution of MHC class I region in primates. Proceedings of the National Academy of Sciences of the United States of America. 2005; 102(26):9230–9234. doi: 10.1073/pnas.0500770102.

Geller R, Adams EJ, Guethlein LA, Little AM, Madrigal AJ, Parham P. Linkage of Patr-AL to Patr-A and-B in the major histocompatibility complex of the common chimpanzee (pan troglodytes). Immunogenetics. 2002; 54(3):212–215. doi: 10.1007/s00251-002-0452-x.

Geraghty DE, Koller BH, Hansen JA, Orr HT. The HLA class I gene family includes at least six genes and twelve pseudogenes and gene fragments. The Journal of Immunology. 1992; 149(6):1934–1946. doi: 10.4049/jimmunol.149.6.1934.

Gfeller D, Bassani-Sternberg M. Predicting antigen presentation-What could we learn from a million peptides? Frontiers in Immunology. 2018; 9(JUL):1–17. doi: 10.3389/fimmu.2018.01716.

Gleimer M, Wahl AR, Hickman HD, Abi-Rached L, Norman PJ, Guethlein LA, Hammond JA, Draghi M, Adams EJ, Juo S, Jalili R, Gharizadeh B, Ronaghi M, Garcia KC, Hildebrand WH, Parham P. Although Divergent in Residues of the Peptide Binding Site, Conserved Chimpanzee Patr-AL and Polymorphic Human HLA-A*02 Have Overlapping Peptide-Binding Repertoires. The Journal of Immunology. 2011; 186(3):1575–1588. doi: 10.4049/jimmunol.1002990.

Go Y, Rakotoarisoa G, Kawamoto Y, Shima T, Koyama N, Randrianjafy A, Mora R, Hirai H. Characterization and evolution of major histocompatibility complex class II genes in the aye-aye, Daubentonia madagascariensis. Primates. 2005; 46(2):135–139. doi: 10.1007/s10329-004-0101-0.

Go Y, Satta Y, Kawamoto Y, Rakotoarisoa G, Randrianjafy A, Koyama N, Hirai H. Frequent segmental sequence exchanges and rapid gene duplication characterize the MHC class I genes in lemurs. Immunogenetics. 2003; 55(7):450–461. doi: 10.1007/s00251-003-0613-6.

Gongora R, Figueroa F, O'Huigin C, Klein J. HLA-DRB9 - Possible remnant of an ancient functional DRB subregion. Scandinavian Journal of Immunology. 1997; 45(5):504–510. doi: 10.1046/j.1365-3083.1997.d01-428.x.

Goyos A, Guethlein LA, Horowitz A, Hilton HG, Gleimer M, Brodsky FM, Parham P. A Distinctive Cytoplasmic Tail Contributes to Low Surface Expression and Intracellular Retention of the Patr-AL MHC Class I Molecule. The Journal of Immunology. 2015; 195(8):3725–3736. doi: 10.4049/jimmunol.1500397.

Grimsley C, Mather KA, Ober C. HLA-H: A pseudogene with increased variation due to balancing selection at neighboring loci. Molecular Biology and Evolution. 1998; 15(12):1581–1588. doi: 10.1093/oxfordjournals.molbev.a025886.

de Groot N, Stanbury K, de Vos-Rouweler AJM, de Groot NG, Poirier N, Blancho G, de Luna C, Doxiadis GGM, Bontrop RE. A quick and robust MHC typing method for free-ranging and captive primate species. Immunogenetics. 2017; 69(4):231–240. doi: 10.1007/s00251-016-0968-0.

de Groot NG, Blokhuis JH, Otting N, Doxiadis GGM, Bontrop RE. Co-evolution of the MHC class I and KIR gene families in rhesus macaques: Ancestry and plasticity. Immunological Reviews. 2015; 267(1):228–245. doi: 10.1111/imr.12313.

de Groot NG, de Groot N, de Vos-Rouweler AJM, Louwerse A, Bruijnesteijn J, Bontrop RE. Dynamic evolution of Mhc haplotypes in cynomolgus macaques of different geographic origins. Immunogenetics. 2022; 74(4):409–429. https://doi.org/10.1007/s00251-021-01249-y, doi: 10.1007/s00251-021-01249-y.

de Groot NG, Heijmans CMC, de Ru AH, Janssen GMC, Drijfhout JW, Otting N, Vangenot C, Doxiadis GGM, Koning F, van Veelen PA, Bontrop RE. A Specialist Macaque MHC Class I Molecule with HLA-B*27–like Peptide-Binding Characteristics. The Journal of Immunology. 2017; 199(10):3679–3690. doi: 10.4049/jimmunol.1700502.

de Groot NG, Heijmans CMC, van der Wiel MKH, Blokhuis JH, Mulder A, Guethlein LA, Doxiadis GGM, Claas FHJ, Parham P, Bontrop RE. Complex MHC Class I Gene Transcription Profiles and Their Functional Impact in Orangutans. The Journal of Immunology. 2016; 196(2):750–758. doi: 10.4049/jimmunol.1500820.

de Groot NG, Heijmans CMC, de Groot N, Doxiadis GGM, Otting N, Bontrop RE. The chimpanzee Mhc-DRB region revisited: Gene content, polymorphism, pseudogenes, and transcripts. Molecular Immunology. 2009; 47(2-3):381–389. doi: 10.1016/j.molimm.2009.09.003.

de Groot NG, Otting N, Maccari G, Robinson J, Hammond JA, Blancher A, Lafont BAP, Guethlein LA, Wroblewski EE, Marsh SGE, Shiina T, Walter L, Vigilant L, Parham P, O'Connor DH, Bontrop RE. Nomenclature report 2019: major histocompatibility complex genes and alleles of Great and Small Ape and Old and New World monkey species. Immunogenetics. 2020; 72(1-2):25–36. doi: 10.1007/s00251-019-01132-x.

Gu X, Nei M. Locus specificity of polymorphic alleles and evolution by a birth-and death process in mammalian MHC genes. Molecular Biology and Evolution. 1999; 16(2):147–156. doi: 10.1093/oxfordjournals.molbev.a026097.

Gu X, Nei M. Locus specificity of polymorphic alleles and evolution by a birth-and death process in mammalian MHC genes. Molecular Biology and Evolution. 1999; 16(2):147–156. doi: 10.1093/oxfordjournals.molbev.a026097.

Gu X, Wang Y, Gu J. Age distribution of human gene families shows significant roles of both large- and small-scale duplications in vertebrate evolution. Nature Genetics. 2002; 31(2):205–209. doi: 10.1038/ng902.

Guethlein LA, Norman PJ, Hilton HHG, Parham P. Co-evolution of MHC class I and variable NK cell receptors in placental mammals. Immunological Reviews. 2015; 267(1):259–282. doi: 10.1111/imr.12326.

Gyllensten UB, Sundvall M, Erlich HA. Allelic diversity is generated by intraexon sequence exchange at the DRB1 locus of primates. Proceedings of the National Academy of Sciences of the United States of America. 1991; 88(9):3686–3690. doi: 10.1073/pnas.88.9.3686.

Hahn MW, De Bie T, Stajich JE, Nguyen C, Cristianini N. Estimating the tempo and mode of gene family evolution from comparative genomic data. Genome Research. 2005; 15(8):1153–1160. doi: 10.1101/gr.3567505.

Hammer SE, Ho CS, Ando A, Rogel-Gaillard C, Charles M, Tector M, Tector AJ, Lunney JK. Importance of the Major Histocompatibility Complex (Swine Leukocyte Antigen) in Swine Health and Biomedical Research. Annual Review of Animal Biosciences. 2020; 8:171–198. doi: 10.1146/annurev-animal-020518-115014.

Hans JB, Bergl RA, Vigilant L. Gorilla MHC class I gene and sequence variation in a comparative context. Immunogenetics. 2017; 69(5):303–323. doi: 10.1007/s00251-017-0974-x.

Hans JB, Haubner A, Arandjelovic M, Bergl RA, Fünfstück T, Gray M, Morgan DB, Robbins MM, Sanz C, Vigilant L. Characterization of MHC class II B polymorphism in multiple populations of wild gorillas using non-invasive samples and next-generation sequencing. American Journal of Primatology. 2015; 77(11):1193–1206. doi: 10.1002/ajp.22458.

Hansen TH, Huang S, Arnold PL, Fremont DH. Patterns of nonclassical MHC antigen presentation. Nature Immunology. 2007; 8(6):563–568. doi: 10.1038/ni1475.

Heijmans CMC, de Groot NG, Bontrop RE. Comparative genetics of the major histocompatibility complex in humans and nonhuman primates. International Journal of Immunogenetics. 2020; 47(3):243–260. doi: 10.1111/iji.12490.

Heimbruch KE, Karl JA, Wiseman RW, Dudley DM, Johnson Z, Kaur A, O'Connor DH. Novel MHC class I full-length allele and haplotype characterization in sooty mangabeys. Immunogenetics. 2015; 67(8):437–445. doi: 10.1007/s00251-015-0847-0.

Hilton HG, Parham P. Missing or altered self: human NK cell receptors that recognize HLA-C. Immunogenetics. 2017; 69(8-9):567–579. doi: 10.1007/s00251-017-1001-y.

Horton R, Gibson R, Coggill P, Miretti M, Allcock RJ, Almeida J, Forbes S, Gilbert JGR, Halls K, Harrow JL, Hart E, Howe K, Jackson DK, Palmer S, Roberts AN, Sims S, Stewart CA, Traherne JA, Trevanion S, Wilming L, et al. Variation analysis and gene annotation of eight MHC haplotypes: The MHC Haplotype Project. Immunogenetics. 2008; 60(1):1–18. doi: 10.1007/s00251-007-0262-2.

Hubert L, Paganini J, Picard C, Chiaroni J, Abi-Rached L, Pontarotti P, Di Cristofaro J. HLA-H*02:07 Is a Membrane-Bound Ligand of Denisovan Origin That Protects against Lysis by Activated Immune Effectors. The Journal of Immunology. 2022; 208(1):49–53. doi: 10.4049/jimmunol.2100358.

Hughes AL, Nei M. Evolution of the major histocompatibility complex: independent origin of nonclassical class I genes in different groups of mammals. Molecular Biology and Evolution. 1989; 6(6):559–579. doi: 10.1093/oxfordjournals.molbev.a040573.

Hughes AL, Nei M. Nucleotide substitution at major histocompatibility complex class II loci: evidence for over-dominant selection. Proceedings of the National Academy of Sciences of the United States of America. 1989; 86(3):958–962. doi: 10.1073/pnas.86.3.958.

Hughes AL. Origin and evolution of HLA class I pseudogenes. Molecular Biology and Evolution. 1995; 12(2):247–258. doi: 10.1093/oxfordjournals.molbev.a040201.

Hughes AL, Hughes MK. Natural selection on the peptide-binding regions of major histocompatibility complex molecules. Immunogenetics. 1995; 42(4):233–243. doi: 10.1007/BF00176440.

Hughes AL, Nei M. Pattern of nucleotide substitution at major histocompatibility complex class I loci reveal overdominant selection. Nature. 1988; 335(6186):167–170. doi: 10.1038/335167a0.

Hurley CK. Naming HLA diversity: A review of HLA nomenclature. Human Immunology. 2021; 82(7):457–465. https://doi.org/10.1016/j.humimm.2020.03.005, doi: 10.1016/j.humimm.2020.03.005.

Karl JA, Prall TM, Bussan HE, Varghese JM, Pal A, Wiseman RW, O'Connor DH. Complete sequencing of a cynomolgus macaque major histocompatibility complex haplotype. Genome Research. 2023; 33(3):448–462. doi: 10.1101/gr.277429.122.

Kasahara M, Klein D, Vincek V, Sarapata DE, Klein J. Comparative anatomy of the primate major histocompatibility complex DR subregion: Evidence for combinations of DRB genes conserved across species. Genomics. 1992; 14(2):340–349. doi: 10.1016/S0888-7543(05)80224-1.

Kaufman J. The new W family reconstructs the evolution of MHC genes. Proceedings of the National Academy of Sciences. 2022; 119(5):119–121.

Kennedy AE, Ozbek U, Dorak MT. What has GWAS done for HLA and disease associations? International Journal of Immunogenetics. 2017; 44(5):195–211. doi: 10.1111/iji.12332.

Klein J, Sato A, Nagl S, Colm O. Molecular trans-species polymorphism. Annual Review of Ecology and Systematics. 1998; 21(1):1–21.

Klein J, Sato A, Nikolaidis N. MHC, TSP, and the origin of species: From immunogenetics to evolutionary genetics. Annual Review of Genetics. 2007; 41:281–304. doi: 10.1146/annurev.genet.41.110306.130137.

Knapp LA, Cadavid LF, Watkins DI. The MHC-E Locus Is the Most Well Conserved of All Known Primate Class I Histocompatibility Genes . The Journal of Immunology. 1998; 160(1):189–196. doi: 10.4049/jimmunol.160.1.189.

Kono A, Brameier M, Roos C, Suzuki S, Shigenari A, Kametani Y, Kitaura K, Matsutani T, Suzuki R, Inoko H, Walter L, Shiina T. Genomic Sequence Analysis of the MHC Class I G/F Segment in Common Marmoset (Callithrix jacchus) . The Journal of Immunology. 2014; 192(7):3239–3246. doi: 10.4049/jimmunol.1302745.

Kuderna LFK, Gao H, Janiak MC, Kuhlwilm M, Orkin JD, Bataillon T, Manu S, Valenzuela A, Bergman J, Rousselle M, Silva FE, Agueda L, Blanc J, Gut M, de Vries D, Goodhead I, Harris RA, Raveendran M, Jensen A, Chuma IS, et al. A global catalog of whole-genome diversity from 233 primate species. Science (New York, NY). 2023; 380(6648):906–913. doi: 10.1126/science.abn7829.

Kulski JK, Anzai T, Inoko H. ERVK9, transposons and the evolution of MHC class I duplicons within the alpha-block of the human and chimpanzee. Cytogenetic and Genome Research. 2005; 110(1-4):181–192. doi: 10.1159/000084951.

Kulski JK, Anzai T, Shiina T, Inoko H. Rhesus macaque class I duplicon structures, organization, and evolution within the alpha block of the major histocompatibility complex. Molecular Biology and Evolution. 2004; 21(11):2079–2091. doi: 10.1093/molbev/msh216.

Kulski JK, Suzuki S, Shiina T. SNP-Density Crossover Maps of Polymorphic Transposable Elements and HLA Genes Within MHC Class I Haplotype Blocks and Junction. Frontiers in Genetics. 2021; 11(January). doi: 10.3389/fgene.2020.594318.

Kupfermann H, Satta Y, Takahata N, Tichy H, Klein J. Evolution of Mhc-DRB introns: Implications for the origin of primates. Journal of Molecular Evolution. 1999; 48(6):663–674. doi: 10.1007/PL00006510.

Lafont BAP, Buckler-White A, Plishka R, Buckler C, Martin MA. Pig-tailed macaques [Macaca nemestrina] possess six MHC-E families that are conserved among macaque species: Implication for their binding to natural killer receptors variants. Immunogenetics. 2004; 56(3):142–154. doi: 10.1007/s00251-004-0663-4.

Lampen MH, Hassan C, Sluijter M, Geluk A, Dijkman K, Tjon JM, de Ru AH, van der Burg SH, van Veelen PA, van Hall T. Alternative peptide repertoire of HLA-E reveals a binding motif that is strikingly similar to HLA-A2. Molecular Immunology. 2013; 53(1-2):126–131. http://dx.doi.org/10.1016/j.molimm.2012.07.009, doi: 10.1016/j.molimm.2012.07.009.

Lenormand C, Bausinger H, Gross F, Signorino-Gelo F, Koch S, Peressin M, Fricker D, Cazenave JP, Bieber T, Hanau D, de la Salle H, Tourne S. HLA-DQA2 and HLA-DQB2 Genes Are Specifically Expressed in Human Langerhans Cells and Encode a New HLA Class II Molecule . The Journal of Immunology. 2012; 188(8):3903–3911. doi: 10.4049/jimmunol.1103048.

Li WH, Gu Z, Wang H, Nekrutenko A. Evolutionary analyses of the human genome. Nature. 2001; 409(6822):847–849. doi: 10.1038/35057039.

Liao WW, Asri M, Ebler J, Doerr D, Haukness M, Hickey G, Lu S, Lucas JK, Monlong J, Abel HJ, Buonaiuto S, Chang XH, Cheng H, Chu J, Colonna V, Eizenga JM, Feng X, Fischer C, Fulton RS, Garg S, et al. A draft human pangenome reference. Nature. 2023; 617(7960):312–324. doi: 10.1038/s41586-023-05896-x.

Lugo JS, Cadavid LF. Patterns of MHC-G-like and MHC-B diversification in new world monkeys. PLoS ONE. 2015; 10(6):1–20. doi: 10.1371/journal.pone.0131343.

Maccari G, Robinson J, Ballingall K, Guethlein LA, Grimholt U, Kaufman J, Ho CS, de Groot NG, Flicek P, Bontrop RE, Hammond JA, Marsh SGE. IPD-MHC 2.0: an improved inter-species database for the study of the major histocompatibility complex. Nucleic Acids Research. 2017 1; 45(D1):D860–D864. https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkw1050, doi: 10.1093/nar/gkw1050.

Maccari G, Robinson J, Hammond JA, Marsh SGE. The IPD Project: a centralised resource for the study of polymorphism in genes of the immune system. Immunogenetics. 2020 2; 72(1-2):49–55. http://link.springer.com/10.1007/s00251-019-01133-w, doi: 10.1007/s00251-019-01133-w.

Maibach V, Hans JB, Hvilsom C, Marques-Bonet T, Vigilant L. MHC class I diversity in chimpanzees and bonobos. Immunogenetics. 2017; 69(10):661–676. doi: 10.1007/s00251-017-0990-x.

Mao Y, Harvey WT, Porubsky D, Munson KM, Hoekzema K, Lewis AP, Audano PA, Rozanski A, Yang X, Zhang S, Yoo DA, Gordon DS, Fair T, Wei X, Logsdon GA, Haukness M, Dishuck PC, Jeong H, del Rosario R, Bauer VL, et al. Structurally divergent and recurrently mutated regions of primate genomes. Cell. 2024; 187(6):1547–1562. https://doi.org/10.1016/j.cell.2024.01.052, doi: 10.1016/j.cell.2024.01.052.

Marsh SGE, Albert ED, Bodmer WF, Bontrop RE, Dupont B, Erlich HA, Fernández-Viña M, Geraghty DE, Holdsworth R, Hurley CK, Lau M, Lee KW, MacH B, Maiers M, Mayr WR, Müller CR, Parham P, Petersdorf EW, Sasazuki T, Strominger JL, et al. Nomenclature for factors of the HLA system, 2010. Tissue Antigens. 2010; 75(4):291–455. doi: 10.1111/j.1399-0039.2010.01466.x.

Mayer WE, Jonker M, Klein D, Ivanyi P, van Seventer G, Klein J. Nucleotide sequences of chimpanzee MHC class I alleles: evidence for trans-species mode of evolution. The EMBO journal. 1988; 7(9):2765–2774. doi: 10.1002/j.1460-2075.1988.tb03131.x.

Messer G, Zemmour J, Orr HT, Parham P, Weiss EH, Girdlestone J. HLA-J, a second inactivated class I HLA gene related to HLA-G and HLA-A. Implications for the evolution of the HLA-A-related genes. The Journal of Immunology. 1992; 148(12):4043–4053. doi: 10.4049/jimmunol.148.12.4043.

Moffett-King A. Natural killer cells and pregnancy. Nature Reviews Immunology. 2002; 2(9):656–663. doi: 10.1038/nri886.

Müller NF, Bouckaert RR. Adaptive metropolis-coupled MCMC for BEAST 2. PeerJ. 2020; 8:1–16. doi: 10.7717/peerj.9473.

Neefjes J, Jongsma MLM, Paul P, Bakke O. Towards a systems understanding of MHC class I and MHC class II antigen presentation. Nature Reviews Immunology. 2011 12; 11(12):823–836. http://www.nature.com/articles/nri3084, doi: 10.1038/nri3084.

Neehus AL, Wistuba J, Ladas N, Eiz-Vesper B, Schlatt S, Müller T. Gene conversion of the major histocompatibility complex class I Caja-G in common marmosets (Callithrix jacchus). Immunology. 2016; 149(3):343–352. doi: 10.1111/imm.12652.

Nei M, Gu X, Sitnikova T. Evolution by the birth-and-death process in multigene families of the vertebrate immune system. Proceedings of the National Academy of Sciences of the United States of America. 1997; 94(15):7799–7806. doi: 10.1073/pnas.94.15.7799.

Nei M, Rooney AP. Concerted and birth-and-death evolution of multigene families. Annual Review of Genetics. 2005; 39:121–152. doi: 10.1146/annurev.genet.39.073003.112240.

Nicholas RE, Sandstrom K, Anderson JL, Smith WR, Wetzel M, Banerjee P, Janaka SK, Evans DT. KIR3DL05 and KIR3DS02 Recognition of a Nonclassical MHC Class I Molecule in the Rhesus Macaque Implicated in Pregnancy Success. Frontiers in Immunology. 2022; 13(March):1–12. doi: 10.3389/fimmu.2022.841136.

Okano M, Miyamae J, Suzuki S, Nishiya K, Katakura F, Kulski JK, Moritomo T, Shiina T. Identification of Novel Alleles and Structural Haplotypes of Major Histocompatibility Complex Class I and DRB Genes in Domestic Cat (Felis catus) by a Newly Developed NGS-Based Genotyping Method. Frontiers in Genetics. 2020; 11(July):1–15. doi: 10.3389/fgene.2020.00750.

Olsson N, Jiang W, Adler LN, Mellins ED, Elias JE. Tuning DO:DM Ratios Modulates MHC Class II Immunopeptidomes. Molecular and Cellular Proteomics. 2022; 21(3):100204. https://doi.org/10.1016/j.mcpro.2022.100204, doi: 10.1016/J.MCPRO.2022.100204.

Otting N, de Groot NG, Bontrop RE. Evolution of HLA-F and its orthologues in primate species: a complex tale of conservation, diversification and inactivation. Immunogenetics. 2020; 72(9-10):475–487. https://doi.org/10.1007/s00251-020-01187-1, doi: 10.1007/s00251-020-01187-1.

Paganini J, Abi-Rached L, Gouret P, Pontarotti P, Chiaroni J, Di Cristofaro J. HLAIb worldwide genetic diversity: New HLA-H alleles and haplotype structure description. Molecular Immunology. 2019; 112(May):40–50. https://doi.org/10.1016/j.molimm.2019.04.017, doi: 10.1016/j.molimm.2019.04.017.

Parham P, Moffett A. Variable NK cell receptors and their MHC class i ligands in immunity, reproduction and
human evolution. Nature Reviews Immunology. 2013; 13(2):133–144. doi: 10.1038/nri3370.

Pierini F, Lenz TL. Divergent Allele Advantage at Human MHC Genes: Signatures of Past and Ongoing Selection.
Molecular Biology and Evolution. 2018 9; 35(9):2145–2158. https://academic.oup.com/mbe/article/35/9/2145/
5034935, doi: 10.1093/molbev/msy116.

Piontkivska H, Nei M. Birth-and-death evolution in primate MHC class I genes: Divergence time estimates.
Molecular Biology and Evolution. 2003; 20(4):601–609. doi: 10.1093/molbev/msg064.

Radley E, Alderton RP, Kelly A, Trowsdale J, Beck S. Genomic organization of HLA-DMA and HLA-DMB. Compari-
son of the gene organization of all six class II families in the human major histocompatibility complex. Journal
of Biological Chemistry. 1994; 269(29):18834–18838. http://dx.doi.org/10.1016/S0021-9258(17)32242-1, doi:
10.1016/s0021-9258(17)32242-1.

Radwan J, Babik W, Kaufman J, Lenz TL, Winternitz J. Advances in the Evolutionary Understanding of MHC
Polymorphism. Trends in Genetics. 2020; 36(4):298–311. doi: 10.1016/j.tig.2020.01.008.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior Summarization in Bayesian Phylogenetics
Using Tracer 1.7. Systematic Biology. 2018 9; 67(5):901–904. https://academic.oup.com/sysbio/article/67/5/
901/4989127, doi: 10.1093/sysbio/syy032.

Riegert P, Wanner V, Bahram S. Genomics, Isoforms, Expression, and Phylogeny of the MHC Class I-Related
MR1 Gene . The Journal of Immunology. 1998; 161(8):4066–4077. doi: 10.4049/jimmunol.161.8.4066.

Robinson J, Barker DJ, Marsh SGE. 25 years of the IPD-IMGT/HLA Database. Hla. 2024; 103(6):e15549. doi:
10.1111/tan.15549.

Salomonsen J, Marston D, Avila D, Bumstead N, Johansson B, Juul-Madsen H, Olesen GD, Riegert P, Skjødt K,
Vainio O, Wiles MV, Kaufman J. The properties of the single chicken MHC classical class II α chain (B-LA) gene
indicate an ancient origin for the DR/E-like isotype of class II molecules. Immunogenetics. 2003; 55(9):605–
614. doi: 10.1007/s00251-003-0620-7.

Satta Y, Mayer WE, Klein J. Evolutionary relationship of HLA-DRB genes inferred from intron sequences.
Journal of Molecular Evolution. 1996 6; 42(6):648–657. http://link.springer.com/10.1007/BF02338798, doi:
10.1007/BF02338798.

Sawai H, Kawamoto Y, Takahata N, Satta Y. Evolutionary Relationships of Major Histocompatibility Complex
Class I Genes in Simian Primates. Genetics. 2004; 166(4):1897–1907. doi: 10.1534/genetics.166.4.1897.

Sawyer SA, GENECONV: a computer package for the statistical detection of gene conversion. Distributed by
the author, Department of Mathematics, Washington University in St. Louis; 1999. http://www.math.wustl.
edu/~sawyer.

Schulze MSED, Wucherpfennig KW. The mechanism of HLA-DM induced peptide exchange in the MHC
class II antigen presentation pathway. Current Opinion in Immunology. 2012; 24(1):105–111. doi:
10.1016/j.coi.2011.11.004.

Shiina T, Blancher A, Inoko H, Kulski JK. Comparative genomics of the human, macaque and mouse major
histocompatibility complex. Immunology. 2017; 150(2):127–138. doi: 10.1111/imm.12624.

Shiina T, Kono A, Westphal N, Suzuki S, Hosomichi K, Kita YF, Roos C, Inoko H, Walter L. Comparative genome
analysis of the major histocompatibility complex (MHC) class i B/C segments in primates elucidated by ge-
nomic sequencing in common marmoset (Callithrix jacchus). Immunogenetics. 2011; 63(8):485–499. doi:
10.1007/s00251-011-0526-8.

Shiina T, Tamiya G, Oka A, Takishima N, Yamagata T, Kikkawa E, Iwata K, Tomizawa M, Okuaki N, Kuwano Y,
Watanabe K, Fukuzumi Y, Itakura S, Sugawara C, Ono A, Yamazaki M, Tashiro H, Ando A, Ikemura T, Soeda E,
et al. Molecular dynamics of MHC genesis unraveled by sequence analysis of the 1,796,938-bp HLA class I
region. Proceedings of the National Academy of Sciences of the United States of America. 1999; 96(23):13282–
13287. doi: 10.1073/pnas.96.23.13282.

Shiina T, Yamada Y, Aarnink A, Suzuki S, Masuya A, Ito S, Ido D, Yamanaka H, Iwatani C, Tsuchiya H, Ishigaki
H, Itoh Y, Ogasawara K, Kulski JK, Blancher A. Discovery of novel MHC-class I alleles and haplotypes in Fil-
ipino cynomolgus macaques (Macaca fascicularis) by pyrosequencing and Sanger sequencing: Mafa-class I
polymorphism. Immunogenetics. 2015; 67(10):563–578. doi: 10.1007/s00251-015-0867-9.

Slierendregt BL, van Noort JT, Bakas RM, Otting N, Jonker M, Bontrop RE. Evolutionary stability of transspecies major histocompatibility complex class II DRB lineages in humans and rhesus monkeys. Human Immunology. 1992 9; 35(1):29–39. https://linkinghub.elsevier.com/retrieve/pii/0198885992900922, doi: 10.1016/0198-8859(92)90092-2.

Takahashi K, Rooney AP, Nei M. Origins and divergence times of mammalian class II MHC gene clusters. Journal of Heredity. 2000; 91(3):198–204. doi: 10.1093/jhered/91.3.198.

Thornton JW, Desalle R. G Ene F Amily E Volution and H Omology. New York. 2000; 1(48):41–73. http://www.ncbi.nlm.nih.gov/pubmed/11701624.

Urvater JA, Otting N, Loehrke JH, Rudersdorf R, Slukvin II, Piekarczyk MS, Golos TG, Hughes AL, Bontrop RE, Watkins DI. Mamu-I: A Novel Primate MHC Class I B-Related Locus with Unusually Low Variability. The Journal of Immunology. 2000; 164(3):1386–1398. https://doi.org/10.4049/jimmunol.164.3.1386.

Van Der Wiel MK, Otting N, De Groot NG, Doxiadis GGM, Bontrop RE. The repertoire of MHC class i genes in the common marmoset: Evidence for functional plasticity. Immunogenetics. 2013; 65(12):841–849. doi: 10.1007/s00251-013-0732-7.

Van Lith M, McEwen-Smith RM, Benham AM. HLA-DP, HLA-DQ, and HLA-DR have different requirements for invariant chain and HLA-DM. Journal of Biological Chemistry. 2010; 285(52):40800–40808. http://dx.doi.org/10.1074/jbc.M110.148155, doi: 10.1074/jbc.M110.148155.

Vollmers S, Lobermeyer A, Körner C. The new kid on the block: Hla-c, a key regulator of natural killer cells in viral immunity. Cells. 2021; 10(11). doi: 10.3390/cells10113108.

Walter L. Nomenclature report on the major histocompatibility complex genes and alleles of the laboratory rat (Rattus norvegicus). Immunogenetics. 2020; 72(1-2):5–8. doi: 10.1007/s00251-019-01131-y.

Watkins DI, Chen ZW, Garber TL, Hughes AL, Letvin NL. Segmental exchange between MHC class I genes in a higher primate: recombination in the gorilla between the ancestor of a human non-functional gene and an A locus gene. Immunogenetics. 1991; 34(3):185–191. doi: 10.1007/BF00205822.

Watkins DI, Chen ZW, Hughes AL, Evans MG, Tedder TF, Letvin NL. Evolution of the MHC class I genes of a New World primate from ancestral homologues of human non-classical genes. Nature. 1990; 346(6279):60–63. doi: 10.1038/346060a0.

Welsh R, Song N, Sadegh-Nasseri S. What to do with HLA-DO/H-2O two decades later? Immunogenetics. 2019; 71(3):189–196. doi: 10.1007/s00251-018-01097-3.

Welsh RA, Sadegh-Nasseri S. The love and hate relationship of HLA-DM/DO in the selection of immunodominant epitopes. Current Opinion in Immunology. 2020; 64(Mhc Ii):117–123. https://doi.org/10.1016/j.coi.2020.05.007, doi: 10.1016/j.coi.2020.05.007.

van der Wiel MKH, Doxiadis GGM, de Groot N, Otting N, de Groot NG, Poirier N, Blancho G, Bontrop RE. MHC class I diversity of olive baboons (Papio anubis) unravelled by next-generation sequencing. Immunogenetics. 2018; 70(7):439–448. doi: 10.1007/s00251-018-1053-7.

Wilming LG, Hart EA, Coggill PC, Horton R, Gilbert JGR, Clee C, Jones M, Lloyd C, Palmer S, Sims S, Whitehead S, Wiley D, Beck S, Harrow JL. Sequencing and comparative analysis of the gorilla MHC genomic sequence. Database. 2013; 2013. doi: 10.1093/database/bat011.

Wroblewski EE, Guethlein LA, Norman PJ, Li Y, Shaw CM, Han AS, Ndjango JBN, Ahuka-Mundeke S, Georgiev AV, Peeters M, Hahn BH, Parham P. Bonobos Maintain Immune System Diversity with Three Functional Types of MHC-B. The Journal of Immunology. 2017; 198(9):3480–3493. doi: 10.4049/jimmunol.1601955.

Wroblewski EE, Parham P, Guethlein LA. Two to tango: Co-evolution of hominid natural killer cell receptors and MHC. Frontiers in Immunology. 2019; 10(FEB):1–22. doi: 10.3389/fimmu.2019.00177.

Wu CH, Suchard MA, Drummond AJ. Bayesian Selection of Nucleotide Substitution Models and Their Site Assignments. Molecular Biology and Evolution. 2013 3; 30(3):669–688. https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/mss258, doi: 10.1093/molbev/mss258.

Yeager M, Hughes AL. Evolution of the mammalian MHC: Natural selection, recombination, and convergent evolution. Immunological Reviews. 1999; 167:45–58. doi: 10.1111/j.1600-065X.1999.tb01381.x.

Zhou Y, Song L, Li H. Full resolution HLA and KIR gene annotations for human genome assemblies. Genome Research. 2024; p. gr.278985.124. doi: 10.1101/gr.278985.124.