More than fruity scents: floral biology, scent and spectral reflectance of

twelve Annonaceae species

Ming-Fai Liu[1,2], Junhao Chen[3,4], Chun-Chiu Pang[5], Tanya Scharaschkin[6], Richard M. K.

Saunders[1]

1: Flora Conservation Department, Kadoorie Farm & Botanic Garden, Lam Kam

Road, Lam Tsuen, Hong Kong.

2: Area of Ecology & Biodiversity, School of Biological Sciences, The University

of Hong Kong, Pokfulam Road, Hong Kong.

3: Singapore Botanic Gardens, National Parks Board, 1 Cluny Road, Singapore

259569, Singapore.

4: Department of Biological Sciences, National University of Singapore, 16

Science Drive 4, Singapore 117558, Singapore

5: Coalition for Research on Ecology and Wildlife Limited, Room F, 25/F, 18

Farm Road, Tokwawan, Kowloon, Hong Kong.

6: Botanical Research, Art and Training, 54 Mill Road, Collinsvale, TAS, 7012,

Australia.

Correspondence

1: Ming-Fai Liu, Flora Conservation Department, Kadoorie Farm & Botanic

Garden, Lam Kam Road, Lam Tsuen, Hong Kong;

Email: nedlmf0@gmail.com

2: Richard M.K. Saunders, Area of Ecology & Biodiversity, School of Biological

Sciences, The University of Hong Kong, Pokfulam Road, Hong Kong.

Email: saunders@hku.hk

ORCIDs

Ming-Fai Liu: https://orcid.org/0000-0002-4732-2512

Junhao Chen: https://orcid.org/0000-0001-8822-169X

Chun-Chiu Pang: https://orcid.org/0000-0002-5286-1238

Tanya Scharaschkin: https://orcid.org/0000-0003-3081-0066

Richard M. K. Saunders: https://orcid.org/0000-0002-8104-7761

Abstract

The pantropical family Annonaceae possesses a broad array of floral phenotypes and

pollination specialisations. Being the most species-rich lineage in the magnoliids,

Annonaceae are a major flowering plant component of tropical rainforests, and are important

in the plant-pollinator network of this biome. Insects such as beetles, flies, bees, cockroaches

and thrips have been reported as their pollinators. Although there has been considerable effort

to assess their interactions with pollinators, attempts to characterise the visual and olfactory

communication channels between them are scarce. Here, we report the floral phenology and

characterise the spectral reflectance and odour of 12 Annonaceae species from five genera,

viz. *Meiogyne*, *Monoon*, *Polyalthia*, *Pseuduvaria*, and *Uvaria*. *Pseuduvaria* species are either

monoecious or dioecious while the four genera produce hermaphroditic flowers. The

Meiogyne species are likely pollinated by small beetles; the *Polyalthia* and *Pseuduvaria*

species are likely pollinated by both beetles and flies; and the *Uvaria* species is likely

pollinated by both beetles and bees. All species produce flowers that are not reflective in the

300–350 nm UV spectrum, and have various spectral reflectance profile across the remaining

UV and visible spectra. We detected diverse floral volatiles across all the species assessed.

The flowers of the small beetle-pollinated species, including *Meiogyne* species and

Polyalthia xanthocarpa, emitted mostly branched-chain esters, including isobutyl acetate and

isoamyl acetate. The *Pseuduvaria* species emitted scent reminiscent of rotten fruit, largely

consisting of 2,3-butanediol, a common by-product in yeast fermentation. The flowers of

Uvaria concava, which were visited by both bees and beetles, released mainly straight-chain

esters, including methyl hexanoate and methyl (Z)-4-octenoate. A corrugated inner petal

outgrowth was observed in *Meiogyne* and *Pseuduvaria*; this functions as a food body reward

in *Meiogyne*, but as a nectary gland for *Pseuduvaria*. This study provides insights into how

Annonaceae flowers attract different guilds of pollinators.

Introduction

The pantropical family Annonaceae is an early-divergent angiosperm lineage of trees and lianas, comprising *ca.* 2500 species from 110 genera (Couvreur et al., 2019). Annonaceae species are one of the major components of tropical lowland to lower montane rainforests (Sosef et al., 2017). Almost all Annonaceae species possess a basic floral architecture of one whorl of three sepals, two whorls of three petals, numerous spirally arranged stamens and often free carpels (van Heusden, 1992). These flowers are largely pollinated by small beetle pollinators from Curculionidae, Nitidulidae, Staphylinidae and Chrysomelidae (Gottsberger, 2012; Saunders, 2012). Because beetle pollinators spend considerable time in the flowers and can often inflict damage to floral tissues, Annonaceae flowers have evolved protective structures like fleshy corolla and shield-like stamen connectives (Gottsberger, 1999). These adaptations are particularly pronounced in Neotropical species that are pollinated by large scarab beetles from Cetoniinae, Dynastinae, Rutelinae and Trichiinae (Gottsberger, 1999). While small beetles pollinate the majority of Annonaceae species, some are reported to be pollinated by thrips (Gottsberger, 1999) or small flies (Silberbauer-Gottsberger et al., 2003), and more rarely, flowers can be pollinated by bees (Teichert et al., 2009) or cockroaches (Nagamitsu et al., 1997). Pollinators are often rewarded with pollen, stigmatic exudates, floral nectar and thickened nutritious floral tissues (Gottsberger and Webber, 2018; Saunders, 2020). The floral chamber created by the corolla often additionally serves as tryst sites, and more rarely as oviposition sites (Saunders, 2020). Highly specialised pollination systems have been previously identified in Annonaceae, such as floral mimicry of fruits (Goodrich and Jürgens, 2018, and references therein), fermentation substrates (Goodrich et al., 2006), mushrooms (Teichert et al., 2012) and aerial litter (Liu et al., 2024). Other derived pollination strategies, including private channel communication with beetles (Maia et al., 2012) and euglossine bees (Teichert et al., 2009) have also been reported. Among these floral specialisations, mimicry of fruits was proposed to be the ancestral condition for the family (Johnson and Schiestl, 2016), and was likely pervasive, as around half of the recorded species emit fruity floral aromas (Goodrich, 2012).

How flowers achieve specialised pollination is a research interest for pollination ecologists. In particular, visual and olfactory cues have been recognised for their roles in mediating these interactions (Schaefer and Ruxton, 2011). Annonaceae flowers display diverse combinations of colour and odour, which might have enabled the evolution of specialisation in pollinators.

Strong floral fragrance is often noticeable in Annonaceae flowers. Based on a review by

Goodrich (2012), there appears to be a myriad of floral odour reported in the family based on

textual description. Although delineation of the chemical components of floral aromas is rare

for Annonaceae, volatiles from over 12 chemical classes have been reported so far (Goodrich,

2012). The type of pollination adaptation seems to be associated with molecules of specific

classes. For instance, proposed fruit mimics appear to be associated with aliphatic esters

(Jürgens, 2000; Johnson and Schiestl, 2016); flowers mimicking fermenting substates

reportedly emit fermentation by-products, including short-chain alcohols and ketones

(Goodrich et al., 2006); and monoterpenoids are associated with aerial litter mimics (Liu et

al., 2024) and flowers pollinated by Euglossines bees (Teichert et al., 2009).

Floral coloration is also diverse within the family, and is more frequently documented in the

literature. Floral colours, including yellow, red, pink, green, orange, white and dark maroon

or indigo flowers have been recorded in Annonaceae (van Heusden, 1992). It has been

reported that colour can shape the choice decision of one of the major pollinating beetle

family Nitidulidae (Döring et al., 2012; Vuts et al., 2022), highlighting the importance of

floral visual cues. Surprisingly, virtually no attempt has been made to characterise

Annonaceae floral colour based on spectral reflectance with only one exception (Liu et al.,

2024). Given the diversity of pollination strategies, floral colour is an overlooked area which

can potentially help elucidate how Annonaceae flowers interact with their pollinators.

Previous studies have revealed that Annonaceae species in Australia display a variety of

floral adaptations to small beetle, fly or bee pollinators (Silberbauer-Gottsberger et al., 2003).

For instance, *Meiogyne* and *Polyalthia* (as *Haplostichanthus*), both generally with yellowish

corolla, are reportedly pollinated by small beetles from the family Nitidulidae and

Curculionidae, respectively. *Pseuduvaria* is reportedly pollinated by flies, and *Uvaria* is

reportedly pollinated by stingless bees. The aim of the current study is to characterise the

phenotypic adaptations for various types of pollination systems in Annonaceae. We selected

12 species from the genera *Meiogyne*, *Monoon*, *Polyalthia*, *Pseudvaria* and *Uvaria* in

Australia. These systems offer arenas for studying specialised bee pollination and potential

floral mimicry of fruit and fermenting substrates. Their floral biology, floral spectral

reflectance and floral odour are characterised here, laying a foundation for the future study of

floral traits and pollination ecology of specialised pollination systems. Furthermore,

magnoliids including Annonaceae preserve many architectural elements of ancestral

angiosperm flowers (Sauquet et al., 2017). Characterisation of the pollination strategies

employed by early-divergent angiosperms like magnoliids may therefore offer insights into

the pollination system of the ancestral flower.

Materials and methods

Study species and study sites

We examined three *Meiogyne* Miq. species (*Meiogyne cylindrocarpa* (Burck) Heusden,

Meiogyne trichocarpa (Jessup) D.C.Thomas & R.M.K. Saunders, *Meiogyne stenopetala*

(F.Muell.) Heusden), one *Monoon* Miq. species (*Monoon patinatum* (Jessup) B.Xue &

R.M.K.Saunders), two *Polyalthia* Blume species, (*Polyalthia hispida* B.Xue &

R.M.K.Saunders, *Polyalthia xanthocarpa* B.Xue & R.M.K.Saunders), four *Pseuduvaria* Miq.

species (*Pseuduvaria hylandii* Jessup, *Pseuduvaria glabrescens* (Jessup) Y.C.F.Su &

R.M.K.Saunders, *Pseuduvaria mulgraveana* Jessup, *Pseuduvaria villosa* Jessup) and two

Uvaria species (*Uvaria concava* Teijsm. & Binn. and *Uvaria rufa* Blume) (See Table 1 for

detailed information on the studied plants. Observations were conducted in the wild for

individuals of *Me. stenopetala* in Mount Tamborine in 2013 (permit number:

WITK08297010). All other observations were made on cultivated individuals at various sites:

Gardens by the Bay, Singapore, in 2018; Garry and Nada Sankowsky Arboretum, Tolga,

Australia, in 2019 and 2023; Cairns Botanic Gardens, Australia, in 2023; and Graham Wood

Arboretum, Wondecla, Australia, in 2023 (see Table 1 for details).

Phenology

To assess floral ontology, 10 flowers for each species were tagged and observed for seven

consecutive days. Observations were made daily to assess the duration of the pistillate and

staminate phase. Stigmatic peroxidase activity has been widely used as a gauge for stigma

receptivity (Dafni and Maués, 1998). However, the stigmas of some species, especially

Meiogyne spp., displayed peroxidase activity even during bud stage based on our preliminary

assessment with Peroxtesmo KO paper (MacheryNagel, Düren, Germany). The onset of

stigmatic receptivity was therefore defined by stigmatic exudate secretion, while the end of

stigmatic receptivity was determined by the abscission of stigmas. Staminate activity was

similarly determined by anther dehiscence and the abscission of petals. A pistillate phase is

defined here as the period that the flower has only the pistillate function, and likewise a

staminate phase is defined as the period that the flower only exhibits the staminate function.

An overlap phase describes the period that the flower has both the pistillate and staminate

functions, while an interim phase refers to the non-receptive period between the pistillate and

staminate phase. Floral thermogenesis was assessed using a handheld digital thermometer

every two hours throughout the day in the floral chamber, with additional readings *ca.* 10 cm

away from the flower to measure ambient temperature.

Floral visitors

Floral visitors were recorded during the entire observation period. A representative number of

floral visitors were collected and either preserved in 50% isopropanol for identification or

frozen at -20°C and silica-dried for inspection of pollen deposition under a dissecting

microscope. Probable pollinators were assessed by these criteria: 1) visitation to flowers of

both sexes or sexual phases; 2) contact with floral sexual organs; and 3) presence of pollen

grains on visitors in pistillate-phase flowers. Since the pistillate phase precedes the staminate

phase, the presence of these pollen-laden insects provide evidence for inter-floral transfer of

pollen grains.

Spectral reflectance of the petals

Pollinators often access Annonaceae floral reproductive organs via apical and basal apertures

created by the inner petals (Saunders, 2012). For genera with similar colour and morphology

between the inner and outer petals, namely *Meiogyne*, *Monoon*, *Polyalthia* and *Uvaria*, petal

spectral reflectance was represented by that of the inner petals. The abaxial surface of the

inner petals was measured for genera with drooping inner petals, viz. *Meiogyne*, *Monoon* and

Polyalthia, whereas the adaxial surface of the inner petals were measured for *Uvaria*, which

has spreading inner petals. *Pseuduvaria* has varying coloration among different parts of the

petals, and therefore three perianth areas were measured: outer petal adaxial surface, inner

petal abaxial surface, and inner petal gland. Differences between sexes and sexual phases

were not assessed due to logistic constraints. Spectral reflectance of the flowers at 300–700

nm was measured using an Ocean Optics FLAME-S-UV-VIS-ES spectrometer (Dunedin, FL,

USA), a DH-mini UV-VIS-NIR light source and an R200-7-UV-VIS probe positioned at 45°.

For each species, individual flowers (n=1–8) selected randomly from one to five biologically

distinct individuals were used as replicates.

Floral odour

To sample the floral volatiles, up to 10 intact flowers were enclosed in a "Toppits" oven bag

for each sample. To control for ambient contaminants, an empty bag was likewise sampled.

Two slits were made in the headspace bag, with one end connecting to a charcoal cartridge

(ORBO 32, Supelco, USA), supplying filtered air, and the other connected to a PAS 500 air

pump (Spectrex, USA). Floral volatiles were collected in a Porapak-Q cartridge (ORBO

1103, 50/80, Supelco, USA) at a flow rate of 100 m/min for 4 h. The porapak-Q cartridge

was then sealed and transported to the laboratory at The University of Hong Kong. The floral

volatiles then were eluted in 1 ml hexane (HPLC-grade, Sigma-Aldrich, USA). The eluents

were then concentrated to 200 µl by nitrogen blowdown. For each species, technical

replicates were made. The internal standard toluene was added to each sample at 34.68 ng/µl.

Gas Chromatography-Mass Spectrometry was performed on an Agilent 6890N/5973 gas

chromatograph-mass selective detector to characterise floral headspace. For each sample, 1

µl eluent was injected into the inlet at 230 °C for 1 min, and then injected into a DB-WAX

column (30 m × 0.25 mm, 0.25 µm thick film; J&W) in splitless mode. Helium was supplied

at a flow rate of 1 ml/min as the carrier gas. Oven temperatures were held at 60 °C for 3 min,

followed by a ramp up to 250 °C at 10 °C/min, and finally held for 7 min. Peaks were

manually integrated and identified tentatively using the Wiley and National Institute of

Standards and Technology mass spectral libraries (NIST, USA; 2020), with an 85% quality

threshold. Compound identities were verified by published standardised retention index

values or by co-injecting with commercially available standards. Molecules were quantified

with the internal standard toluene (see also Kantsa et al., 2018). The relative abundance of

each molecule was calculated. Volatiles recovered in the ambient control sample were

excluded in other samples.

Results

218   The floral biology of the studied species is summarised in Table 1. Most species assessed

219   displayed strong levels of protogyny and none were thermogenic. All *Meiogyne*, *Monoon*,

220   *Polyalthia* and *Uvaria* species assessed here bear hermaphroditic flowers, whereas the

221   *Pseuduvaria* species bear unisexual flowers (*Ps. hylandii* and *Ps. glabrescens*) or a

222   combination of staminate and hermaphroditic flowers (*Ps. mulgraveana*).

Floral biology and pollinator activities—*Meiogyne* species

The three *Meiogyne* species (Fig. 1a–c) produced yellow flowers with longitudinally grooved

basal outgrowths on the adaxial surface of the inner petals (Fig. 1c, Fig. 2a, c) and emitted a

banana-like scent. Their pistillate phase lasted two days, while the duration of staminate

phase differs among species (1–3 days) (Table 1). They exhibited a short overlap between the

pistillate and staminate phases that lasts *ca.* 0–5 hours. Small beetles were frequently

observed visiting the flowers of *Me. cylindrocarpa* (Curculionidae and Nitidulidae), *Me.*

trichocarpa (Curculionidae and Nitidulidae), and *Me. stenopetala* (Nitidulidae and

Staphylinidae) in both staminate and pistillate phase, during which they contacted the

stamens and stigmas. Pollen-laden curculionid beetles from the tribe Ochyromerini and

pollen-laden nitidulid beetles from the subfamily nitidulinae observed in pistillate-phase

flowers were likely the pollinators of both *Me. cylindrocarpa* and *Me. trichocarpa*. The

staphylinid beetles from the subfamily Omaliinae and the nitidulid beetles from the subfamily

Epuraeinae were likely the pollinators of *Me. stenopetala.* These floral pollinators consumed

pollen grains on the flowers where the beetles also sought shelter and copulated. Stigmas of

all three *Meiogyne* species produced a thin film of stigmatic exudate, which became more

copious towards the end of the pistillate phase. Consumption of stigmatic exudate was

invariably observed in curculionid, nitidulid and staphylinid beetle visitors. Consumption of

floral tissues, specifically petals and stigmas, were observed, primarily by curculionid

beetles, and less frequently by nitidulid and staphylinid beetles, leaving gnaw marks on the

floral tissues of the three *Meiogyne* species. However, consumption of the inner petal

corrugation was particularly pronounced compared to the surrounding floral tissues (Fig. 2).

Nectar secretion was not observed on the inner petal corrugation of *Me. cylindrocarpa*, *Me.*

trichocarpa and *Me. stenopetala*, thus they likely functioned as a food body reward for the

floral visitors.

Floral biology and pollinator activities—*Polyalthia* species

Polyalthia xanthocarpa (Fig. 1f) exhibited a 2–3-day pistillate phase followed by a 1-day

staminate phase (Table 1). The transition between pistillate and staminate phase was not

observed directly and could have occurred at night: it is therefore unclear whether an interim

or overlap phase was present. As with *Meiogyne*, *Po. xanthocarpa* bore yellow flowers with a

banana-like odour (Fig. 1f) and were visited by curculionid beetles that accessed the floral

sexual organs in both the staminate and pistillate phases (Table 1). Pollen grains were not

observed on the limited number of pistillate-phase visiting beetles, however. The curculionid

beetles were observed to consume pollen and utilise the flower for shelter. Secretion of

stigmatic exudate in *Po. xanthocarpa* was weak and the consumption of stigmatic exudate

was not observed. In contrast, *Polyalthia hispida* bore yellow flowers without detectable

odour (Fig. 1e). Its anthesis comprised a three-day pistillate phase followed by a two-day

staminate phase. No floral visitors were observed.

Floral biology and pollinator activities—*Uvaria* species

Uvaria concava bore flowers with red petals and emitted a fruity odour (Fig. 1k). It had a 2–

3-day pistillate phase, followed by a prolonged overlap phase (1–2 days), in which stigmas

remained attached to the ovaries as the anthers dehisce. The drying and abscission of stigmas

usually occurred one day after the overlap phase and were sometimes synchronised with petal

abscission. In most flower, petal abscission occurred two days after the staminate function

initiated. Because of the variation in the timing of stigma drying, the duration of staminate

phase varied between 0–1 day. Nitidulid beetles (subfamily Nitidulinae) were the primary

floral visitors observed: they accessed the floral sexual organs and were present throughout

anthesis. They were probable pollinators since pollen-laden individuals were found in

pistillate-phase flowers. We observed the visitation of a single honeybee (*Apis* sp.) during the

overlap phase; although the bee contacted the floral sexual organs while collecting pollen, we

failed to capture it and were unable to assess pollen deposition.

Uvaria rufa flowers bore orange petals with tints of red and green (Fig. 1l). The flower

emitted a fruity odour and had a one-day pistillate phase, followed by a one-day interim

phase and a four-day staminate phase. Only one unidentified fly was observed, which visited

during the interim phase. It is unclear whether pollen grains were deposited onto this fly

visitor.

Floral biology and pollinator activities—*Pseuduvaria*

Pseuduvaria species (Fig. 1g–j) displayed an array of sexual systems (Table 1). *Pseuduvaria*

hylandii was assessed to be dioecious, while *Ps. mulgraveana* and *Ps. glabrescens* were

assessed to be structurally andromonoecious, in which the same individuals bear staminate

flowers that lack carpels, as well as bisexual flowers with a few stamens. *Pseuduvaria*

hylandii had prolonged anthesis for both pistillate and staminate flowers (> 6 days), but we

were unable to assess the anthesis duration (and floral phenology) for *Ps. glabrescens* due to

logistic limitations. Anthesis in staminate flowers of *Ps. mulgraveana* lasts two days, whereas

in hermaphroditic flowers the pistillate phase lasts for two days, followed immediately by

petal abscission, with anthers dehiscing two days after petal abscission. We were only able to

assess one staminate individual of *Ps. villosa* and were therefore unable to determine its

sexual system.

The flowers of all four *Pseuduvaria* species possessed white outer petals (Fig. 1g–j). They

also possessed white to reddish inner petals with dark maroon inner petal glands on their

adaxial surface (fig. 2d). These flowers emitted a strong fermentation odour that resembled

rotten fruits. They had a mitriform corolla, with large basal apertures that allowed larger

insects to access the floral chamber. The inner petals were apically connivent and the three

pairs of glands on the inner petals form a platform in the floral chamber (Fig. 1h, Fig. 2d–f).

Nectar secretion was observed on the inner petal glands and surrounding dark red petal

tissues of the four *Pseuduvaria* species (Fig. 2e). Drosophilid flies and nitidulid beetles were

observed to be the primary floral visitors of *Ps. glabrescens*, *Ps. hylandii* and *Ps.*

mulgraveana (Table 1). These insects were observed accessing the floral chamber and

consuming nectar, often positioning themselves underneath the sexual organs to access the

nectary. Pollen grains were observed to be deposited either onto the back of the insects (Fig.

2f), or onto the inner petal glands where the pollen grains were then picked up secondarily by

the flies and beetles on their legs. The drosophilid flies and nitidulid beetles were observed

moving across the stigmas and stamens with their legs touching the sexual organs, which may

assist pollen transfer. Some drosophilid flies are large enough to brush the back of their

thorax or abdomen against the sexual organs, which may additionally deposit pollen grains

onto the stigmas. Both drosophilid flies and nitidulid beetles were observed to access the

sexual organs of pistillate and staminate flowers for *Ps. glabrescens*, *Ps. hylandii* and *Ps.*

mulgraveana. While pollen grains were observed on the drosophilid flies and nitidulid beetles

found in pistillate-phase flowers of *Ps. hylandii*, pollen deposition was observed only on

nitidulid beetles from pistillate-phase flowers of *Ps. glabrescens* and *Ps. mulgraveana*. Other

dipteran visitors, namely lauxaniid and neriid flies, were also observed in *Ps. hylandii*, but

they are too large to access the floral chamber and thus unlikely to be pollinators.

Unfortunately, we failed to observe any floral visitors for *Ps. villosa*, though it has similar

floral colour and odour to the other *Pseuduvaria* species (Fig. 1j).

Floral biology and pollinator activities—*Monoon*

Monoon patinatum produced flowers with greenish petals (Fig. 1d). Only two staphylinid

beetle visitors from the subfamily Tachyporinae were observed during the pistillate phase

(Table 1). Pollen grains were not observed on their bodies.

Floral Colour

Various types of floral coloration were observed in the studied species, including red (*Uvaria*

concava), orange (*U. rufa*), yellow (*Meiogyne cylindrocarpa*, *Me. trichocarpa*, *Me.*

stenopetala, *Polyalthia hispida* and *Po. xanthocarpa*), green (*Monoon patinatum*), and a

mixtue of white and maroon (*Pseuduvaria hylandii*, *Ps. glabrescens*, *Ps. mulgraveana* and

Ps. villosa). The spectral reflectance of the petals of 12 Annonaceae species is shown in Fig.

3. The petals of all species are not UV-reflective from 300–350 nm; most of the reflectance

was observed in the visible spectrum (400–700 nm). Note that while yellow-flowered species

possess similar coloration to human perception, their spectral reflectance can be quite varied,

highlighting the importance of quantification of reflective spectra.

Floral Odour

The floral headspace composition of 11 Annonaceae species is summarised in Table 2. We

detected a total of 97 volatiles from seven classes of compounds. We did not detect any major

volatiles in the headspace of *Polyalthia hispida* against the ambient air control, and therefore

did not present the headspace data here.

Aliphatic ester was the most abundant class of volatiles in *Meiogyne cylindrocarpa*, *Me.*

trichocarpa, *Me. stenopetala*, *Polyalthia xanthocarpa*, *Uvaria concava* and *U. rufa*. The

headspaces of all *Meiogyne* species and *Po. xanthocarpa* contained a varying degree of

esters, including isobutyl acetate (*Me. cylindrocarpa*: $7.87 \pm 2.16\%$; *Me. trichocarpa*: 11.86

$\pm 4.37\%$; *Me. stenopetala*: $56.71 \pm 1.5\%$) and isoamyl acetate (*Me. cylindrocarpa*: $79.58 \pm$

7.65%; *Me. trichocarpa*: $84.07 \pm 4.01\%$; *Me. stenopetala*: $28.17 \pm 4.13\%$; *Po. xanthocarpa*:

$99.28 \pm 0.02\%$). For *U. concava*, the floral headspaces were dominated by two esters, namely

methyl-(*Z*)-4-octenoate ($25.36 \pm 0.32\%$), and methyl hexanoate ($54.69 \pm 0.32\%$), while for *U.*

rufa, the most abundant volatiles were hexyl acetate ($21.27 \pm 0.25\%$), butyl acetate ($20.65 \pm$

0.21%), and ethyl octanoate ($20.04 \pm 0.22\%$). In *Meiogyne* and *Polyalthia*, branched-chain

aliphatic esters dominated the floral odour, while in *Uvaria*, the floral odours primarily

consisted of straight-chain the aliphatic esters. For *Monoon patinatum*, the headspace mainly

comprised the monoterpenes β-ocimene ($45.50 \pm 0.61\%$) and allo-ocimene ($11.89 \pm 0.59\%$),

and 2-methyl-butanoic acid (16.48 ± 0.53%). The oxygenated aliphatic compound was the

most abundant volatile class for *Pseuduvaria hylandii*, *Ps. glabrescens*, *Ps. mulgraveana* and

Ps. villosa, consisting of the molecule 2,3-butanediol (84.83–96.98% of the headspaces).

Discussion

Floral phenology

Most species assessed in this study exhibited a 3–5 day anthesis period and none exhibited

circadian trapping, a time-dependent trapping mechanism in which pollinators are trapped

and released in accordance to the insects' peak circadian activities (Lau et al., 2017). The

duration of anthesis for most species in the current study is similar to the typical non-trapping

Annonaceae species (≥ 2 days) (Pang and Saunders, 2014; Lau et al., 2017). As the family is

self-compatible (Pang and Saunders, 2014), temporal separation of sexual phases is an

important adaptation to reduce autogamy. An interim phase was observed in *Uvaria rufa*,

during which the stigmas are no longer receptive and anther dehiscence has not begun,

potentially offering additional assurance to prevent autogamy (Pang and Saunders, 2014).

However, in some of the other species, including *Meiogyne* spp. and *U. concava*, an overlap

phase was observed, during which carpels and stamens are simultaneously functional within a

flower. This overlap phase could offer reproductive assurance by autogamy when pollinator

availability is scarce. It was previously reported that *U. concava* was pollinated by stingless

bees (Meliponinae) in its natural population (50 km from our site of study) (Silberbauer-

Gottsber et al., 2003). Because bee visitors to Annonaceae often collect pollen only as a food

reward, and seldom visit pistillate-phase flowers, the overlap phase has been suggested to be

essential for bee pollination (Gottsberger, 2014). It was, however, reported that meliponine

visitors also collect stigmatic exudate as a reward (Silberbauer-Gottsberger et al., 2013),

suggesting the flower offers reward to the meliponine visitors even during pistillate phase.

The role of the prolonged overlap phase for *U. concava* is therefore contentious.

Floral phenology also played a role in the sexual function of the flower, especially for

Pseuduvaria species. Previous studies have shown that *Ps. mulgraveana* is structurally

andromonoecious (Pang et al., 2013), meaning that staminate flowers and bisexual flowers

are borne on the same plant. The stamens of bisexual flowers, however, release functional

pollen grains only after petal abscission (Pang et al., 2013), rendering the plant functionally

monoecious, which is corroborated by our observations.

Inner petal corrugation and glands

Inner petal outgrowth was observed in *Meiogyne* spp. and *Pseuduvaria* spp. in the current

study. This outgrowth has independently evolved in around 20 genera, including *Annona* L.,

Alphonsea Hook.f. & Thomson, *Asimina* Adans., *Asteranthe* Engl. & Diels, *Duguetia* A.St.-

Hil., *Maasia* Mols, Kessler & Rogstad, *Meiogyne* Miq., *Miliusa* Lesch. ex A.DC., *Orophea*

Blume, *Porcelia* Ruiz & Pav., *Pseuduvaria* Miq., *Sapranthus* Seem., *Stenanona* Standl.,

Tetrameranthus R.E.Fr., *Tridimeris* Baill., *Uvaria* L., *Wangia* X.Guo & R.M.K.Saunders,

and *Xylopia* L. (van Heusden, 1992; Schatz and Maas, 2010; Xue et al., 2017). In

Pseuduvaria, the inner petal outgrowth has been reported to be a nectary gland (Silberbauer-

Gottsberger et al., 2003). In the current study, the four *Pseuduvaria* species also possessed

this floral tissue, which invariably functioned as nectary, corroborating the previous study.

Likewise, all three *Meiogyne* species assessed in our study possessed weakly folded to

longitudinally grooved basal growths on the adaxial surface of the inner petals. Although this

structure has previously been interpreted as a gland (van Heusden, 1992, 1994), histological

studies have failed to identify secretory ducts or hairs (Xue et al., 2021). The inner petal

corrugation may potentially offer tactile cues for the floral mimicry of aerial litter in

Meiogyne heteropetala (Liu et al., 2024), but the role of this tissue remains enigmatic for

most *Meiogyne* species. Our observations indicate that pollinators preferentially consume the

inner petals over other floral structures (Fig. 2), suggesting that the corrugation likely

functions as a food body in *Me. cylindrocarpa*, *Me. trichocarpa* and *Me. stenopetala*. In

Annonaceae, specialised nutritious petal tissues are common in species that are pollinated by

scarab beetles in the neotropics, including *Annona*, *Cymbopetalum*, *Duguetia*, and *Malmea*

species (Gottsberger and Webber, 2018). Among these lineages with specialised food body,

there is potential convergence in tissue structure. The inner petal corrugation of these species

is anatomically similar to those found in *Meiogyne*, lacking secretory openings, ducts or

trichromes, and comprising a mixture of cells full of starch granules and cells enriched with

tannins (Gottsberger and Webber, 2018; Xue et al., 2021).

Exploitation of pollinators associated with fruits

The fruity floral aromas and beetle pollination of Annonaceae have led some authors to

propose some Annonaceae flowers mimic fruits (Johnson and Schiestl, 2016; Goodrich and

Jürgens, 2018). In particular, fruit mimicry has been postulated in *Anaxagorea*, *Annona*,

Artabotrys, *Duguetia* and *Xylopia* and is likely to be prevalent in the Annonaceae (Johnson

and Schiestl, 2016; Goodrich and Jürgens, 2018; Chen et al., 2020). These flowers are

pollinated by small beetles such as Curculionidae, Nitidulidae and Staphylinidae that are

known to use fruits as a food source or oviposition site (Phelan and Lin, 1991; Racette et al.,

1992; Frank and Thomas, 1999; Mutinelli et al., 2015). Previous studies have recorded small-

beetle visitors in *Polyalthia xanthocarpa* flowers (as "*Haplostichanthus* spec. ("Cape

tribulation")"; Morawetz, 1988), and curculionid beetles have been observed visiting the

flowers in Cape Tribulation, Australia (pers. comm., Dr David Tng). For *Meiogyne*, nitidulid

and curculionid beetles have been reported to be the pollinators of *Me. cylindrocarpa* in

tropical rainforests in West Malesia (Momose, 2005; Roubik et al, 2005). Beetles from these

families were also observed in the current study.

In this study, floral aromas resembling ripe fruits were recorded in *Me. cylindrocarpa*, *Me.*

trichocarpa, *Me. stenopetala*, *Po. xanthocarpa*. Small beetle pollinators (families Nitidulidae,

Curculionidae and Staphylinidae) were observed as floral visitor of *Me. cylindrocarpa*, *Me.*

trichocarpa, *Me. stenopetala* and *Po. xanthocarpa* (Table 1). Their floral odour resembled

strong banana-like scent to human perception, and is largely composed of either isoamyl

acetate or isobutyl acetate (Table 2). These two molecules have been reported in various

fruits, including banana (Jordán et al., 2001), papaya (Katague and Kirch, 1965), pear (Zhang

et al., 2023) and wild soursop (Pino et al., 2002), and is implicated as the semiochemicals of

Curculionidae and Nitidulidae (Rochat et al., 2000; Torto et al., 2007). The spectral

reflectance of *Meiogyne* species, especially *Me. cylindrocarpa* and *Me. trichocarpa*,

exhibited low UV reflectance, with stronger reflectance from 500–700 nm (Fig. 3). The UV-

absorbing yellow spectral profile is a characteristic for pollen (Dötterl et al., 2014) and was

also reported in fruits such as bananas (Xie et al., 2018). Some nitidulid and curculionid

beetles are known to associate with fruits (Phelan and Lin, 1991; Racette et al., 1992;

Mutinelli et al., 2015), and nitidulidae has previously been demonstrated to be attracted to

various shade of yellow (Döring et al., 2012; Vuts et al., 2022). The current evidence

suggests that *Me. cylindrocarpa*, *Me. trichocarpa*, *Me. stenopetala* and *Po. xanthocarpa*

likely attract their pollinators at least by exploiting visual and olfactory cues that indicate

fruits.

Exploitation of pollinators associated with fermented substrates

Records of floral visitors to *Pseuduvaria* are more comprehensive than for most other

Annonaceae genera in Australia. In a previous pollination study on another Australian

Pseuduvaria species, *Ps. froggattii*, it was shown that the flowers are primarily pollinated by

drosophilid flies (Silberbauer-Gottsberger et al., 2003). Similarly, *Ps. glabrescens* was

previously reported to be visited primarily by small flies (as *Pseuduvaria* spec. ("Davies

Creek"); Morawetz, 1988). *Pseuduvaria mulgraveana*, on the other hand, was reported to be

pollinated by the nitidulid beetle, *Aethina australis*, and was only infrequently visited by

flies. Our observations are largely congruent with these studies: *Ps. hylandii*, *Ps. glabrescens*

and *Ps. mulgraveana* were observed to be visited by both nitidulid beetles and drosophilid

flies. Pollen deposition on the nitidulid beetles was observed in the pistillate-phase flowers of

all three species, while drosophilid flies appeared to be the pollinator only for *Ps. hylandii*. It

was previously postulated that flies could pollinate *Ps. villosa*, but floral visitors were not

observed in our study.

Floral coloration in the *Pseuduvaria* species varies across different areas of petal tissues

(Figs. 1 & 3). The abaxial surface of the inner petals in most species demonstrate largely

similar reflectance spectra with the adaxial surface of the outer petals, except for *Ps. hylandii*.

In contrast to these areas, the inner petal glands of all four *Pseuduvaria* species studied are

burgundy red with generally lower reflectance across all wavelengths. Additionally, all four

Pseuduvaria species emit an odour reminiscent of rotten fruits. Rotting fruits have been

reported to display a reduction in overall reflectance across the visible spectrum under fungal

infection (Liu et al., 2020). The contrast between the dark burgundy inner petal gland and the

remaining bright floral tissues may have evolved to imitate rotten patches on fruits.

Interestingly, in *Pseuduvaria*, nectar secretion is localised in the darkened area (Fig. 2),

suggesting there is co-localisation of gustatory cue and visual cue, which may help position

the pollinators underneath the reproductive organs and facilitate pollen transfer. The floral

odours of all four *Pseuduvaria* species are largely composed of 2,3-butanediol (Table 2),

which is a common volatile emitted by yeast and fermenting substrates (Goodrich et al.,

2006, 2023). This molecule is one of the major volatiles of *Asimina triloba* (Goodrich et al.,

2006), a maroon-petalled floral mimic of fermentation substrates (Goodrich et al., 2023). As

with *Ps. glabrescens*, *Ps. hylandii* and *Ps. mulgraveana*, *Asimina triloba* is pollinated by

small flies and small beetles (Martin, 2021; Goodrich et al., 2023). This suggests

convergence in floral colour and odour between *Asimina* and *Pseuduvaria* for attracting the

same pollinator guilds. The four *Pseuduvaria* species may therefore be potential candidates

for floral mimics of fermentation substrates.

Exploitation of bee and beetle pollinators

Uvaria concava was reported to be pollinated by stingless bees in the wild (Silberbauer-Gottsberger et al., 2003) but is here inferred to be pollinated by nitidulid beetles in cultivation (Table 1). This species possesses bright-red petals, reflecting the visible spectrum mostly at 600–700 nm (Fig. 3). However, neither the tribe Meliponini (stingless bees) nor the family Apidae have photoreceptors within this range (van der Kooi et al., 2021). Though the photosensitivity of Nitidulidae is unknown, most coleopteran families with known visual systems are unable to detect this part of the visible spectrum (van der Kooi et al., 2021). It is therefore unclear how floral visual cues assist pollinator attraction in *U. concava*. The floral odour of *U. concava* is largely composed of methyl hexanoate and methyl (*Z*)-4-octenoate. While there is little documentation how meliponine bees and nitidulid beetles respond to these molecules, methyl hexanoate is reported to be a pheromone for male bumblebees (Valterová et al., 2001). Among Annonaceae species that emit esters, branched-chain aliphatic esters are typically produced (Table 2; Jürgens et al, 2000), but *Uvaria* spp. primarily produce straight-chain aliphatic esters instead. Interestingly, the branched-chain aliphatic ester isoamyl acetate (the dominant floral scent component of *Meiogyne* spp. and *Polyalthia xanthocarpa*) is a bee alarm pheromone that reduces foraging in the honeybee species *Apis mellifera* and *Apis cerana* (Gong et al., 2017). Nonetheless, whether this structural change in floral esters contributes to the shift to bee pollination remains to be investigated. Future studies comparing the contrast between spectral reflectance of the petals and reproductive organs as well as colour-based and scent-based bioassays could help assess the role played by visual and olfactory cues in bee-pollinated Annonaceae species.

Conclusions

The basic floral biology, petal spectral reflectance and floral scent of selected species in five Annonaceae genera are characterised here in detail. Our study has identified multiple species that have floral traits resembling fruits (*Meiogyne* spp. and *Po. xanthocarpa*) and fermented substrates (*Meiogyne* spp. and *Po. xanthocarpa*). The specialised floral cues and adaptations identified in this study provide preliminary evidence for potential floral mimicry of fruits and fermented substrates, and offer insights into how floral cues may attract floral visitors. These findings serve as a prelude to future studies on specialised pollination systems, and provide baseline information on floral traits in the pantropical family Annonaceae. Future studies involving the characterisation of colour and odour of co-ocurring fruits and fermenting

substrates would be important to assess whether these species exploit their pollinators through floral mimicry.

Acknowledgements

We would like to express our gratitude to Garry and Nada Sankowsky, Graham Wood, Charles Clarke (Cairns Botanic Gardens), and Janelle Jung (Gardens by the Bay) for allowing us to work on the living collections of Annonaceae species in their arboreta or botanic gardens. We are grateful for Frank Zich from Australian Tropical Herbarium, and staff from Mount Tamborine National Park and the Queensland Department of Environment, Science and Innovation for their assistance, and Jessie Lai and Laura Wong for their technical support. This research is funded by the Hong Kong Research Grants Council (HKU17112616), awarded to R.M.K.S.

References

Chen, J. Liu, M.-F., and Saunders, R M. K. (2020). Contrasting floral biology of *Artabotrys* species (Annonaceae): Implications for the evolution of pollinator trapping. *Plant Species Biology*, *35*, 210–223.

Couvreur, T. L. P., Helmstetter, A. J., Koenen, E. J. M., Bethune, K., Brandão, R. D., Little, S. A., Sauquet, H. and Erkens, R. H. J. (2019). Phylogenomics of the major tropical plant family Annonaceae using targeted enrichment of nuclear genes. *Frontiers in Plant Science*, *9*, 431757.

Dafni, A., and Maués, M. M. (1998). A rapid and simple procedure to determine stigma receptivity. *Sexual Plant Reproduction*, *11*(3), 177–180.

Döring, T. F., Skellern, M., Watts, N., & Cook, S. M. (2012). Colour choice behaviour in the pollen beetle *Meligethes aeneus* (Coleoptera: Nitidulidae). *Physiological Entomology*, *37*(4), 360–378.

Dötterl, S., Glück, U., Jürgens, A., Woodring, J., and Aas, G. (2014). Floral reward,

advertisement and attractiveness to honey bees in dioecious *Salix caprea*. *PloS One*, *9*(3),

e93421.

Frank, J. H., and Thomas, M. C. (1999). Rove Beetles of Florida, Staphylinidae (Insecta:

Coleoptera: Staphylinidae). *DPI Entomology Circulars*, *343*, 1–11.

Gong, Z., Wang, C., Dong, S., Zhang, X., Wang, Y., Hu, Z., and Tan, K. (2017). High

concentrations of the alarm pheromone component, isopentyl acetate, reduces foraging and

dancing in *Apis mellifera* Ligustica and *Apis cerana* Cerana. *Journal of Insect Behavior*, *30*,

188–198.

Goodrich, K. R., and Jürgens, A. (2018). Pollination systems involving floral mimicry of

fruit: aspects of their ecology and evolution. *New Phytologist*, *217*(1), 74–81.

Goodrich, K. R., Zjhra, M. L., Ley, C. A., and Raguso, R. A. (2006). When flowers smell

fermented: the chemistry and ontogeny of yeasty floral scent in pawpaw (*Asimina triloba*:

Annonaceae). *International Journal of Plant Sciences*, *167*(1), 33–46.

Goodrich, K. R., Ellis, I., DeHaas, A., Senski, R., and Savage, J. (2023). False advertising

with fermented scents: floral mimicry in pawpaw (*Asimina* triloba: Annonaceae)

pollination. *International Journal of Plant Sciences*, *184*(6), 485–497.

Gottsberger, G. (1999). Pollination and evolution in neotropical Annonaceae. *Plant Species

Biology*, *14*(2), 143–152.

Gottsberger, G. (2012). How diverse are Annonaceae with regard to pollination? *Botanical

Journal of the Linnean Society*, *169*(1), 245–261.

Gottsberger, G. (2014). Evolutionary steps in the reproductive biology of

Annonaceae. *Revista Brasileira de Fruticultura*, *36*, 32–43.

Gottsberger, G., and Webber, A. C. (2018). Nutritious tissue in petals of Annonaceae and its

function in pollination by scarab beetles. *Acta Botanica Brasilica*, *32*, 279–286.

Johnson, S. D., and Schiestl, F. P. (2016). *Floral mimicry*. Oxford University Press.

Jordán, M. J., Tandon, K., Shaw, P. E., and Goodner, K. L. (2001). Aromatic profile of

aqueous banana essence and banana fruit by gas chromatography-mass spectrometry (GC-

MS) and gas chromatography-olfactometry (GC-O). *Journal of Agricultural and Food*

Chemistry, *49*(10), 4813–4817.

Jürgens, A., Webber, A. C., and Gottsberger, G. (2000). Floral scent compounds of

Amazonian Annonaceae species pollinated by small beetles and

thrips. *Phytochemistry*, *55*(6), 551–558.

Kantsa, A., Raguso, R. A., Dyer, A. G., Olesen, J. M., Tscheulin, T., and Petanidou, T.

(2018). Disentangling the role of floral sensory stimuli in pollination networks. *Nature*

Communications, *9*(1), 1041.

Katague, D.B. and Kirch, E.R. (1965), Chromatographic analysis of the volatile components

of papaya fruit. *Journal of Pharmaceutical Sciences*, *54*(6), 891–894.

Lau, J. Y., Guo, X., Pang, C. C., Tang, C. C., Thomas, D. C., and Saunders, R. M. K. (2017).

Time-dependent trapping of pollinators driven by the alignment of floral phenology with

insect circadian rhythms. *Frontiers in Plant Science*, *8*, 1119.

Liu, Q., Zhou, D., Tu, S., Xiao, H., Zhang, B., Sun, Y., Pan., L., and Tu, K. (2020).

Quantitative visualization of fungal contamination in peach fruit using hyperspectral

imaging. *Food Analytical Methods*, *13*, 1262–1270.

Liu, M. F., Chen, J., Goodrich, K. R., Chiu, S. K., Pang, C. C., Scharaschkin, T., and

Saunders, R. M. K. (2024). Aerial litter mimicry: a novel form of floral deception mediated

by a monoterpene synthase. *bioRxiv*, 2024.06.12.596753; doi:

https://doi.org/10.1101/2024.06.12.596753

Maia, A. C. D., Dötterl, S., Kaiser, R., Silberbauer-Gottsberger, I., Teichert, H., Gibernau,

M., do Amaral Ferraz Navarro, D. M., Schlindwein, C., and Gottsberger, G. (2012). The key

role of 4-methyl-5-vinylthiazole in the attraction of scarab beetle pollinators: a unique

olfactory floral signal shared by Annonaceae and Araceae. *Journal of Chemical Ecology*, *38*,

1072–1080.

Martin, K. R. (2021). *When flowers play Dead: microbes as architects of a 'deceptive' floral*

phenotype (Doctoral dissertation, Cornell University).

Momose, K. 2005. Beetle pollination in tropical rain forests. In Roubik, D.W., Sakai, S. and

Karim, A.A.H. (Eds.), *Pollination ecology and the rain forest* (pp. 104–110). New York:

Springer.

Morawetz, W. (1988). Karyosystematics and evolution of Australian Annonaceae as

compared with Eupomatiaceae, Himantandraceae, and Austrobaileyaceae. *Plant Systematics*

and Evolution, *159*, 49–79.

Mutinelli, F., Federico, G., Carlin, S., Montarsi, F., and Audisio, P. (2015). Preliminary

investigation on other Nitidulidae beetles species occurring on rotten fruit in Reggio Calabria

province (south-western Italy) infested with small hive beetle (*Aethina tumida*). *Journal of*

Apicultural Research, 54(3), 233–235.

Nagamitsu, T., and Inoue, T. (1997). Cockroach pollination and breeding system of *Uvaria*

elmeri (Annonaceae) in a lowland mixed☐dipterocarp forest in Sarawak. *American Journal*

of Botany, *84*(2), 208–213.

Pang, C. C., Scharaschkin, T., Su, Y. C., and Saunders, R. M. K. (2013). Functional monoecy

due to delayed anther dehiscence: a novel mechanism in *Pseuduvaria mulgraveana*

(Annonaceae). *PLoS One*, *8*(3), e59951.

Pang, C. C., and Saunders, R. M. K. (2014). The evolution of alternative mechanisms that

promote outcrossing in Annonaceae, a self-compatible family of early-divergent

angiosperms. *Botanical Journal of the Linnean Society*, *174*(1), 93–109.

Phelan, P. L., and Lin, H. (1991). Chemical characterization of fruit and fungal volatiles

attractive to dried-fruit beetle, *Carpophilus hemipterus* (L.)(Coleoptera: Nitidulidae). *Journal*

of Chemical Ecology, *17*, 1253–1272.

Pino, J. A., Marbot, R., and Agüero, J. (2002). Volatile components of wild soursop (*Annona*

montana Macf.) fruit. *Journal of Essential Oil Research*, *14*(4), 257–258.

Racette, G., Chouinard, G., Vincent, C., and Hill, S. B. (1992). Ecology and management of

plum curculio, *Conotrachelus nenuphar* [Coleoptera: Curculionidae], in apple

orchards. *Phytoprotection*, *73*(3), 85–100.

Rochat, D., Meillour, P. N. L., Esteban-Duran, J. R., Malosse, C., Perthuis, B., Morin, J. P.,

and Descoins, C. (2000). Identification of pheromone synergists in American palm weevil,

Rhynchophorus palmarum, and attraction of related *Dynamis borassi*. *Journal of Chemical*

Ecology, *26*, 155–187.

Roubik, D. W., Sakai, S. and Karim, A. A. H. (2005). Appendix A. In Roubik, D. W., Sakai,

S. and Karim, A. A. H. (Eds.), *Pollination ecology and the rain forest*: *Sarawak studies* (pp.

223–245). New York: Springer.

Saunders, R. M. K. (2012). The diversity and evolution of pollination systems in

Annonaceae. *Botanical Journal of the Linnean Society*, *169*(1), 222–244.

Saunders, R. M. K. (2020). The evolution of key functional floral traits in the early divergent

angiosperm family Annonaceae. *Journal of Systematics and Evolution*, *58*(4), 369–392.

Sauquet, H., Von Balthazar, M., Magallón, S., Doyle, J. A., Endress, P. K., Bailes, E. J., de

Morais, E. B. et al. (2017). The ancestral flower of angiosperms and its early

diversification. *Nature Communications*, *8*(1), 16047.

Schaefer, H. M., and Ruxton, G. D. (2011). Plant-animal communication. OUP Oxford.

Schatz, G. E., and Maas, P. J. M. (2010). Synoptic revision of *Stenanona*

(Annonaceae). *Blumea*, *55*(3), 205–223.

Silberbauer-Gottsberger, I., Gottsberger, G., and Webber, A. C. (2003). Morphological and

functional flower characteristics of new and old world Annonaceae with respect to their mode

of pollination. *Taxon*, *52*(4), 701–718.

Sletvold, N., Trunschke, J., Smit, M., Verbeek, J., and Ågren, J. (2016). Strong pollinator-

mediated selection for increased flower brightness and contrast in a deceptive

orchid. *Evolution*, *70*(3), 716–724.

Sosef, M. S., Dauby, G., Blach-Overgaard, A., van Der Burgt, X., Catarino, L., Damen, T.,

Deblauwe, V., Dessein, S., Dransfield, J., Droissart, V., Duarte, M. C., Engledow, H., Fadeur,

G., Figueira, R., Gereau, R. E., Hardy, O. J., Harris, D. J., de Heij, J., Janssens, S., Klomberg,

Y., Ley, A. C., Mackinder, B. A., Meerts, P., van de Poel, J. L., Sonké, B., Stévart, T.,

Stoffelen, P., Svenning, J.-C., Sepulchre, P., Zaiss, R., Wieringa, J. J. and Couvreur, T. L.

(2017). Exploring the floristic diversity of tropical Africa. *BMC Biology*, *15*, 1–23.

Teichert, H., Dötterl, S., Zimma, B., Ayasse, M., and Gottsberger, G. (2009).

Perfume□collecting male euglossine bees as pollinators of a basal angiosperm: the case of

Unonopsis stipitata (Annonaceae). *Plant Biology*, *11*(1), 29–37.

Teichert, H., Dötterl, S., Frame, D., Kirejtshuk, A., and Gottsberger, G. (2012). A novel

pollination mode, saprocantharophily, in *Duguetia cadaverica* (Annonaceae): a stinkhorn

(Phallales) flower mimic. *Flora-Morphology, Distribution, Functional Ecology of*

Plants, *207*(7), 522–529.

Torto, B., Arbogast, R. T., Alborn, H., Suazo, A., Van Engelsdorp, D., Boucias, D.,

Tumlinson, J. H., and Teal, P. E. (2007). Composition of volatiles from fermenting pollen

dough and attractiveness to the small hive beetle *Aethina tumida*, a parasite of the honeybee

Apis mellifera. Apidologie, *38*(4), 380–389.

Valterová, I., Urbanová, K., Hovorka, O., and Kindl, J. (2001). Composition of the labial

gland secretion of the bumblebee males *Bombus pomorum. Zeitschrift für Naturforschung*

C, *56*(5–6), 430–436.

Van Heusden, E. C. H. (1992). Flowers of Annonaceae: morphology, classification, and

evolution. *Blumea. Supplement*, *7*(1), 1–218.

Van Heusden, E. C. H. (1994). Revision of *Meiogyne* (Annonaceae). *Blumea: Biodiversity,*

Evolution and Biogeography of Plants, *38*(2), 487–511.

Van Der Kooi, C. J., Stavenga, D. G., Arikawa, K., Belušič, G., and Kelber, A. (2021).

Evolution of insect color vision: from spectral sensitivity to visual ecology. *Annual Review of*

Entomology, *66*, 43–5461.

Vuts, J., Szarukán, I., Marczali, Z., Csonka, É. B., Szilágyi, A., Imrei, Z., Nagy, A. and Tóth,

M. (2022). Differences in colour preference among pollen beetle species (Coleoptera:

Nitidulidae). *Journal of Applied Entomology*, *146*(3), 301–309.

Xue, B., Shao, Y. Y., Saunders, R. M., and Tan, Y. H. (2017). *Alphonsea glandulosa*

(Annonaceae), a new species from Yunnan, China. *PLoS One*, *12*(2), e0170107.

Xie, C., Chu, B., and He, Y. (2018). Prediction of banana color and firmness using a novel

wavelengths selection method of hyperspectral imaging. *Food Chemistry*, *245*, 132–140.

Xue, B., Shao, Y. Y., Xiao, C. F., Liu, M. F., Li, Y., and Tan, Y. H. (2021). *Meiogyne*

oligocarpa (Annonaceae), a new species from Yunnan, China. *PeerJ*, *9*, e10999.

Zhang, W., Yan, M., Zheng, X., Chen, Z., Li, H., Mao, J., Qin, H., Zhu, C., Du, H., and Abd

El-Aty, A. M. (2023). Exploring the aroma fingerprint of various Chinese pear cultivars

through qualitative and quantitative analysis of volatile compounds using HS-SPME and

GC× GC-TOFMS. *Molecules*, *28*(12), 4794.

753
Fig. 1. Flowers of Annonaceae species studied. (a) *Meiogyne cylindrocarpa*. (b) *Meiogyne*
trichocarpa with curculionid and nitidulid visitors. (c) *Meiogyne stenopetala* with a
staphylinid visitor. (d) *Monoon patinatum*. (e) *Polyalthia hispida*. (f) *Polyalthia xanthocarpa*.
(g) *Pseuduvaria glabrescens* with nitidulid visitors. (h) *Pseuduvaria hylandii* with
drosophilid visitors. (i) *Pseuduvaria mulgraveana*. (j) *Pseuduvaria villosa*. (k) *Uvaria*
concava. (l) *Uvaria rufa* at interim phase with a fly visitor. Photo credit: Photo credits: (a, b,
d, e, g–l): Ming-Fai Liu; (c): Chun-Chiu Pang; (f): Garry & Nada Sankowsky.

Fig. 2. Inner petal outgrowths. (a) Adaxial surface of inner petal of *Meiogyne cylindrocarpa*.
(b) Gnaw marks (magenta arrows) created by curculionid beetles on the inner petals of
Meiogyne trichocarpa. (c) Gnaw marks (magenta arrows) created by staphylinid beetles on
the inner petals of *Meiogyne stenopetala*: longitudinal ridges of the basal corrugation have
been gnawed away. (d) Pair of nectary glands on the adaxial surface of the inner petal of
Pseuduvaria hylandii. (e) Nectar secreted by *Pseuduvaria mulgraveana* (yellow arrows). (f)
Drosophilid fly consuming nectar in a *Pseuduvaria hylandii* flower. Photo credit: (a–f):
Ming-Fai Liu.

Fig. 3. Spectral reflectance of 12 Annonaceae species. (a) *Meiogyne cylindrocarpa*. (b) *Meiogyne trichocarpa*. (c) *Meiogyne stenopetala*. (d) *Monoon patinatum*. (e) *Polyalthia hispida*. (f) *Polyalthia xanthocarpa*. (g) *Pseuduvaria glabrescens*. (h) *Pseuduvaria hylandii*. (i) *Pseuduvaria mulgraveana*. (j) *Pseuduvaria villosa*. (k) *Uvaria concava*. (l) *Uvaria rufa*. For species with drooping inner petals and no colour differentiation between outer and inner petal whorls, the inner petal abaxial surface was measured (a–f; i.e. *Meiogyne*, *Monoon* and *Polyalthia*). For species with spreading inner petals and no colour differentiation between outer and inner petal whorls, the inner petal adaxial surface (k–l; i.e. *Uvaria*) was measured. For species with dissimilar petal whorls (g–j; i.e. *Pseuduvaria*), three perianth areas were measured (solid line: inner petal abaxial surface; dotted line: outer petal adaxial surface; dashed line: inner petal gland); the sample size (*n*) for these species refer to the sample size for these three perianth areas in the aforementioned order. Photo credits: (a, b, d, e, g–l): Ming-Fai Liu; (c): Chun-Chiu Pang; (f): Garry & Nada Sankowsky.

Table 1. Floral biology of *Meiogyne*, *Monoon*, *Polyalthia*, *Pseuduvaria* and *Uvaria* species. Abbreviations: CBG: Cairns Botanic Gardens, NE Queensland, Australia; GBB: Gardens by the Bay, Singapore; MT: Mt. Tamborine, SE Queensland, Australia; SA: Sankowsky Arboretum, Tolga, NE Queensland, Australia; WA: Graham Wood Arboretum, Wondecla, NE Queensland, Australia; +: present, -: absent, [†]: presence on both pistillate-phase and staminate-phase flowers, [*]: contact with reproductive organs, [‡]: pistillate-phase visitors were pollen-laden.

	Meiogyne cylindrocarpa	*Meiogyne trichocarpa*	*Meiogyne stenopetala*	*Monoon patinatum*	*Polyalthia hispida*	*Polyalthia xanthocarpa*	*Pseuduvaria hylandii*	*Pseuduvaria glabrescens*	*Pseuduvaria mulgraveana*	*Pseuduvaria villosa*	*Uvaria concava*	*Uvaria rufa*
Site of study (*n*= individual studied)	SA (*n*=8), GBB (*n*=3), WA (*n*=4)	SA (*n*=8)	SA (*n*=1), WA (*n*=1), MT (*n*=20)	SA (*n*=1)	SA (*n*=2)	SA (*n*=1)	SA (*n*=5)	SA (*n*=1), CBG (*n*=1)	SA (*n*=3)	SA (*n*=1)	SA (*n*=2)	SA (*n*=2)
Habit	treelet/shrub	Treelet	treelet	tree	treelet	treelet	treelet	treelet	treelet	treelet	climber	climber
Flower position	axillary/apical	axillary/apical	axillary/apical	ramiflorous	axillary/apical	axillary/apical/ ramiflorous	cauliflorous	axillary/apical	axillary/apical	ramiflorous	axillary/apical	axillary/apical/ ramiflorous
Sexual system	bisexual flower	bisexual flower	bisexual flower	bisexual flower	bisexual flower	bisexual flower	unisexual flower, dioecious	andromonoecious	andromonoecious	? (only seen staminate treelet)	bisexual flower	bisexual flower
Pistillate phase duration	2d	2d	2d	not assessed	3d	2–3d	>6d	not assessed	2d	not assessed	2–3d	1d
Staminate phase duration	2d	1d	3d	not assessed	2d	1d	>6d	not assessed	2d	not assessed	0–1d	4d
Overlap phase duration	*ca.* 0–3h	*ca.* 5h	*ca.* 1–3h	not assessed	?	?	not applicable	not applicable	-	not assessed	1–2d	-
Interim phase duration	-	-	-	not assessed	?	?	not applicable	not applicable	2d	not assessed	-	1d
Protogyny	+	+	+	not assessed	+	+	not applicable	not applicable	+	not assessed	+	+
Pollinator trapping	-	-	-	-	-	-	-	-	-	-	-	-
Thermogenesis	-	-	-	-	-	-	-	-	-	-	-	-
Floral odour	banana-like	banana-like	banana-like	odorous	odourless	banana-like	rotten fruit	rotten fruit	rotten fruit	rotten fruit	fruity	fruity
Inner petal growth/gland	+, smooth–weakly longitudinally grooved	+, smooth–weakly longitudinally grooved	+, longitudinally grooved	-	-	-	+, smooth & raised	+, smooth & raised	+, smooth & raised	+, smooth & raised	-	-
Inner petal corrugation function	food body	food body	food body	corrugation absent	corrugation absent	corrugation absent	nectary	nectary	nectary	nectary	corrugation absent	corrugation absent
Floral visitors (*n*=total number of floral visitors observed)	*n*=42 (sites: SA+WA) **Coleoptera:** Curculionidae: Ochyromerini (14.3%)[†,*,‡] Curculionidae: *Carpophilus* sp. (4.8%)[†,*] Nitidulidae: Nitidulinae (81%)[†,*,‡]	*n*=29 **Coleoptera:** Curculionidae: Ochyromerini (17.2%)[†,*,‡] Nitidulidae: Nitidulinae (82.8%)[†,*,‡]	*n*=14 (site: MT+WA) **Coleoptera:** Staphylinidae: Omaliinae (64.3%)[†,*,‡] Nitidulidae: Epuraeinae (35.7%)[†,*,‡]	*n*=2 **Coleoptera:** Staphylinidae: Tachyporinae (100%)*	not observed	*n*=8 **Coleoptera:** Curculionidae: Ochyromerini (100%)[†,*]	*n*=58 **Diptera:** Drosophilidae: Drosophilinae (60.3%)[†,*,‡] Lauxaniidae (3.4%)[†] Neriidae (3.4%)[†] **Coleoptera:** Nitidulidae: Nitidulinae (31.0%)[†,*,‡] **Hymenoptera:** Formicidae (1.7%)	*n*=19 (site: CBG) **Diptera:** Drosophilidae: Drosophilinae (21%)[†,*] **Coleoptera:** Nitidulidae: Nitidulinae (79%)[†,*,‡]	*n*=33 **Diptera:** Drosophilidae: Drosophilinae (21.2%)[†,*] **Coleoptera:** [†,*]Nitidulidae: Nitidulinae (72.7%)[†,*,‡] **Hymenoptera:** Formicidae (6.1%)	not observed	*n*=13 **Coleoptera:** Nitidulidae: Nitidulinae (92.3%)[†,*,‡] **Hymenoptera:** *Apis* sp. (7.7%)[†,*]	*n*=1 **Diptera:** unknown sp. (100%)

Table 2. Floral scent composition of *Meiogyne*, *Monoon*, *Polyalthia*, *Pseuduvaria* and *Uvaria* species. The average percentage of the relative peak area for each volatile and its standard error are presented below. Italicised molecules were verified by co-injecting with commercially available standards. Unknown molecules were denoted by the top ten most abundant ion fragments, followed by their relative intensity in parentheses. *n* refers to the number of technical replicates performed, followed by the number of individuals the odour was collected from.

	standard retention index (DB-WAX)	Relative content (%)										
		Meiogyne cylindrocarpa (*n*=15;15)	*Meiogyne trichocarpa* (*n*=8;8)	*Meiogyne stenopetala* (*n*=5;2)	*Monoon patinatum* (*n*=3;1)	*Polyalthia xanthocarpa* (*n*=5;1)	*Pseuduvaria hylandii.* (*n*=3;3)	*Pseuduvaria glabrescens* (*n*=4;2)	*Pseuduvaria mulgraveana* (*n*=4;3)	*Pseuduvaria villosa* (*n*=4;1)	*Uvaria concava* (*n*=3;2)	*Uvaria rufa* (*n*=4;2)
Aliphatic hydrocarbon												
1-Tetradecene	1348	—	0.19 ± 0.19	—	—	—	—	—	—	—	—	—
Tetradecane	1401	—	0.33 ± 0.22	—	—	—	—	—	—	—	—	—
Oxygenated Aliphatic Compound												
2,3-Butanediol	1589	—	—	—	—	—	95.97 ± 0.91	84.83 ± 0.80	96.98 ± 0.21	95.32 ± 0.15	—	—
2-Methyl-butanoic acid	1678	—	—	—	16.48 ± 0.53	—	—	—	—	—	—	—
2-Methyl-(*E*)-7-hexadecene	1725	—	—	—	—	—	—	—	—	—	—	0.05 ± 0.00
Ester												
Isobutyl acetate	1014	7.87 ± 2.16	11.86 ± 4.37	56.71 ± 1.50	—	—	—	—	—	—	—	—
Ethyl butanoate	1039	—	—	—	—	—	—	—	—	—	—	7.35 ± 0.38
Butyl acetate	1073	0.22 ± 0.24	—	—	—	—	—	—	—	—	—	20.65 ± 0.21
Isoamyl acetate	1109	79.58 ± 7.65	84.07 ± 4.01	28.17 ± 4.13	—	99.28 ± 0.02	—	—	—	—	—	—
Methyl hexanoate	1191	—	—	—	—	—	—	—	—	—	54.69 ± 0.32	4.70 ± 0.08
Butyl butanoate	1223	—	—	—	—	—	—	—	—	—	—	1.94 ± 0.07
2-methyl-(2*E*)-butenyl acetate	1225	0.82 ± 0.70	—	—	—	—	—	—	—	—	—	—
Ethyl hexanoate	1238	—	—	—	—	—	—	—	—	—	—	12.58 ± 0.29
2-methyl-(2*Z*)-butenyl acetate	1254	1.16 ± 0.70	—	—	—	—	—	—	—	—	—	—
Hexyl acetate	1277	0.07 ± 0.87	—	0.87 ± 0.12	—	—	—	—	—	—	—	21.27 ± 0.25
Isoamyl isovalerate	1316	1.09 ± 0.85	—	—	—	—	—	—	—	—	—	—
Methyl octanoate	1396	—	—	—	—	—	—	—	—	—	4.30 ± 0.29	0.56 ± 0.02
Hexyl butanoate	1423	—	—	—	—	—	—	—	—	—	—	6.38 ± 0.10
Methyl (*Z*)-4-octenoate	1438	—	—	—	—	—	—	—	—	—	25.36 ± 0.32	—

Compound	RI											
Ethyl octanoate	1442	—	—	—	—	—	—	—	—	—	—	20.04 ± 0.22
Hexyl 2-methylbutanoate	1452	—	—	0.44 ± 0.07	—	—	—	—	—	—	—	—
Octyl acetate	1482	—	—	—	—	—	—	—	—	—	—	1.48 ± 0.06
trans-2-Hexenyl isovalerate	1508	—	—	2.34 ± 0.12	—	—	—	—	—	—	—	—
2,3-Butanediyl diacetate	1530	1.26 ± 1.37	—	0.09 ± 0.05	—	—	—	—	—	—	—	—
linalyl anthranilate	1565	—	0.01 ± 0.01	—	—	—	—	—	—	—	—	—
Hexyl hexanoate	1618	—	—	—	—	—	—	—	—	—	—	0.86 ± 0.02
Octyl butanoate	1625	—	—	—	—	—	—	—	—	—	—	0.66 ± 0.03
Hexyl tiglate	1632	—	—	—	—	—	—	—	—	—	—	0.16 ± 0.02
Ethyl decanoate	1646	—	—	—	—	—	—	—	—	—	—	0.23 ± 0.01
Ethyl trans-4-decenoate	1675	—	—	—	—	—	—	—	—	—	—	0.09 ± 0.00
6-Methyl-4-heptenyl pentanoate	1678	—	—	—	—	—	—	—	—	—	—	0.20 ± 0.00
Ethyl benzoate	1685	—	—	—	—	—	—	—	—	—	—	0.08 ± 0.01
(E)-Methylgeranate	1706	—	—	—	—	—	—	—	—	—	—	0.30 ± 0.00
Geranyl 2-methylbutanoate	1735	—	0.03 ± 0.01	—	—	—	—	—	—	—	—	—
Geranyl acetate	1766	1.56 ± 1.97	1.24 ± 0.59	—	—	—	—	—	—	—	—	—
2-Phenylethyl acetate	1835	—	—	0.52 ± 0.01	—	—	—	—	—	—	—	—
Phenylethyl isovalerate	2011	—	—	0.54 ± 0.27	—	—	—	—	—	—	—	—
hexyl benzoate	2098	—	—	—	—	—	—	—	—	—	—	0.05 ± 0.01
Phenylpropanoid												
Methyl eugenol	2026	—	—	—	—	—	—	1.06 ± 0.09	—	—	—	—
Monoterpene												
β-Myrcene	1170	—	0.15 ± 0.11	—	—	—	—	—	—	—	—	—
Limonene	1205	—	1.24 ± 0.61	—	—	—	—	—	—	—	—	—
1,8-cineole	1210	—	0.01 ± 0.01	—	—	—	2.42 ± 1.22	11.78 ± 0.38	—	—	—	—
trans-β-Ocimene	1241	—	0.06 ± 0.04	—	—	—	—	—	—	—	—	—
cis-β-Ocimene	1259	—	0.14 ± 0.10	—	45.50 ± 0.61	—	—	—	—	—	—	—
terpinolene	1283	—	0.05 ± 0.03	—	—	—	—	—	—	—	—	—
allo-Ocimene	1384	—	0.18 ± 0.13	—	11.89 ± 0.59	—	—	—	—	—	—	—
Menthone	1456	—	0.03 ± 0.02	—	—	—	—	—	—	—	1.04 ± 0.07	—
α-Terpineol	1711	—	—	—	—	—	—	—	—	—	—	—
camphor	1532	—	0.02 ± 0.02	—	—	—	—	—	—	—	—	—
menthacamphor	1654	—	0.06 ± 0.04	—	—	—	—	—	—	—	—	—
geranylacetone	1868	—	—	0.18 ± 0.04	3.01 ± 0.12	—	—	—	—	—	—	—

Sesquiterpene

Compound	RI											
α-Cubebene	1471	1.11 ± 1.31	—	0.06 ± 0.06	—	—	—	—	—	—	—	—
δ-elemene	1485	1.14 ± 1.36	—	—	—	—	—	—	—	—	—	—
Copaene	1511	—	—	0.83 ± 0.35	—	0.32 ± 0.01	0.26 ± 0.26	1.12 ± 0.05	0.65 ± 0.05	1.05 ± 0.10	—	—
β-bourbonene	1518	1.25 ± 1.45	—	—	—	—	—	—	—	—	—	—
β-Cubebene	1554	1.25 ± 1.50	—	—	—	0.04 ± 0.00	—	—	—	—	—	—
Cyperene	1549	—	—	—	—	—	0.42 ± 0.42	—	—	—	—	—
β-ylangene	1578	1.32 ± 1.57	—	—	—	—	—	—	—	—	—	—
Calarene	1602	1.48 ± 1.60	—	—	—	—	—	—	—	—	—	—
(*E)-β-caryophyllene*	1608	2.87 ± 1.51	—	—	6.08 ± 0.16	0.14 ± 0.01	—	—	—	—	1.86 ± 0.05	—
γ-elemene	1647	1.42 ± 1.72	—	—	—	—	—	—	—	—	—	—
cis-β-Farnesene	1672	—	0.02 ± 0.01	—	—	—	—	—	—	—	3.39 ± 0.00	—
humulene	1681	2.26 ± 1.70	—	—	2.83 ± 0.27	0.04 ± 0.00	—	—	—	—	—	—
γ-Muurolene	1709	—	—	—	—	0.07 ± 0.00	—	—	—	—	—	—
δ-Selinene	1724	—	—	0.08 ± 0.03	—	—	—	—	—	—	—	—
Germacrene D	1732	1.86 ± 1.83	—	—	3.87 ± 0.14	0.05 ± 0.00	—	—	—	—	0.31 ± 0.16	—
Copaene/ γ-Muurolene	1732	—	—	0.10 ± 0.03	—	—	—	—	—	—	—	—
α-Muurolene	1744	—	—	0.10 ± 0.03	—	—	—	—	—	—	0.79 ± 0.02	—
α-Farnesene	1756	—	—	—	7.40 ± 0.07	—	—	—	—	—	—	0.03 ± 0.00
Bicylogermacrene	1758	1.59 ± 1.91	—	0.10 ± 0.04	—	—	—	—	—	—	—	—
δ-Cadinene	1777	1.65 ± 1.96	—	0.37 ± 0.10	1.83 ± 0.18	0.06 ± 0.00	—	—	—	—	—	0.04 ± 0.01
Cada-1,4-diene	1805	—	—	0.08 ± 0.02	—	—	—	—	—	—	—	—
Miscellaneous Compound												
Naphthalene	1768	—	—	0.06 ± 0.02	1.11 ± 0.09	—	—	—	—	—	0.21 ± 0.11	0.02 ± 0.00
β-Ionene	1872	—	—	—	—	—	—	—	—	—	0.49 ± 0.03	—
Unknown												
43(24), 57(20), 41(14), 56(11), 42(8), 70(7), 55(6), 86(4), 61(3), 69(3)	1176	0.56 ± 0.51	—	0.09 ± 0.06	—	—	—	—	—	—	—	—
57(26), 43(19), 41(12), 56(11), 70(7), 42(7), 71(6), 55(5), 85(4), 86(3)	1201	0.16 ± 0.72	—	1.95 ± 0.54	—	—	—	—	—	—	—	—
57(23), 43(22), 41(13), 56(12), 71(8), 42(6), 85(5), 69(4), 55(4), 68(4)	1244	—	—	1.55 ± 0.33	—	—	—	—	—	—	—	—
57(20), 70(15), 43(14), 85(12),	1299	—	—	1.17 ± 0.34	—	—	—	—	—	—	—	—

41(10), 71(9), 56(7), 55(6), 42(5), 86(3)													
73(29), 57(17), 41(11), 43(11), 56(9), 147(7), 42(5), 32(5), 86(4), 207(3)	1316	—	—	—	—	—	—	0.61 ± 0.21	—	—	—	—	—
43(24), 67(16), 57(15), 82(12), 41(11), 56(7), 29(4), 42(4), 55(3), 39(3)	1319	—	—	0.37 ± 0.14	—	—	—	—	—	—	—	—	—
57(19), 43(18), 41(15), 56(11), 105.1(8), 42(7), 119(7), 29(6), 32(6), 86(4)	1344	—	—	—	—	—	0.47 ± 0.24	—	1.09 ± 0.08	1.90 ± 0.10	—	—	—
43(21), 57(18), 56(15), 41(13), 70(7), 42(6), 55(6), 32(5), 69(4), 86(4)	1381	—	—	0.01 ± 0.01	—	—	—	—	—	—	—	—	—
43(52), 87(11), 57(9), 70(6), 41(5), 56(5), 42(3), 55(3), 86(3), 45(3)	1387	—	—	2.11 ± 0.46	—	—	0.12 ± 0.06	—	—	—	—	—	—
57(28), 43(15), 41(13), 56(12), 85(6), 71(6), 42(6), 55(6), 32(4), 86(3)	1405	0.19 ± 0.85	—	0.61 ± 0.15	—	—	0.19 ± 0.10	0.59 ± 0.20	1.12 ± 0.05	1.74 ± 0.07	—	—	—
57(20), 82(15), 67(14), 41(11), 43(10), 85(9), 56(7), 55(5), 32(4), 42(4)	1475	1.66 ± 1.25	—	0.08 ± 0.08	—	—	0.09 ± 0.05	—	0.17 ± 0.10	—	—	1.09 ± 0.19	0.02 ± 0.02
57(20), 43(15), 41(15), 56(10), 80(9), 32(8), 29(6), 79.1(6), 42(6), 71(5)	1493	—	—	—	—	—	—	—	—	—	—	0.48 ± 0.11	—
57(21),43(17),56(12),85(11),41(11),103(9),84(6),42(5),55(5),32(5)	1499	1.81 ± 1.36	—	0.27 ± 0.05	—	—	—	—	—	—	—	0.44 ± 0.10	—
57(22), 43(16), 41(16), 56(12), 32(8), 55(7), 42(6), 71(5), 119(5), 69(5)	1510	—	—	—	—	—	—	—	—	—	—	1.34 ± 0.11	—
57(23), 85(14), 43(14), 41(11), 70(10), 56(8), 71(5), 112(5), 42(5), 55(5)	1555	—	—	0.15 ± 0.08	—	—	0.06 ± 0.06	—	—	—	—	—	—
111(26), 81(13), 154(12),	1607	0.14 ± 1.73	—	—	—	—	—	—	—	—	—	3.59 ± 0.07	—

95(10), 112(8), 123(7), 79(7), 41(6), 94(6), 53(5)												
84(15), 57(13), 123(13), 41(12), 43(11), 42(8), 56(8), 81(8), 95(6), 32(6)	1614	—	—	—	—	—	—	—	—	—	—	0.05 ± 0.00
79(16), 108(13), 57(13), 41(12), 43(11), 150(8), 67(8), 56(7), 32(7), 29(6)	1736	—	—	—	—	—	—	—	—	—	0.39 ± 0.02	0.06 ± 0.00
57(19), 43(16), 41(12), 32(11), 56(11), 108(8), 42(6), 45(6), 91(6), 44(5)	1745	—	—	—	—	—	—	—	—	—	0.21 ± 0.12	0.01 ± 0.00
43(16), 57.1(13), 56(11), 41(11), 117.1(10), 84.1(10), 71(8), 55(8), 145.1(7), 69(7)	1821	—	—	—	—	—	—	—	—	—	—	0.15 ± 0.01