1 **A chronic murine model of pulmonary *Acinetobacter baumannii* infection enabling the**
2 **investigation of late virulence factors, long-term antibiotic treatments, and polymicrobial**
3 **infections**

Clay D. Jackson-Litteken[1], Gisela Di Venanzio[1], Manon Janet-Maitre[1], Ítalo A. Castro[1,2], Joseph
J. Mackel[3], David A. Rosen[1,3], Carolina B. López[1,2], Mario F. Feldman[1,]*

[1]Department of Molecular Microbiology, Washington University School of Medicine, Saint Louis,
Missouri, USA.

[2]Center for Women's Infectious Diseases Research, Washington University School of Medicine,
Saint Louis, Missouri, USA.

[3]Department of Pediatrics, Division of Infectious Diseases, Washington University School of
Medicine, Saint Louis, Missouri, USA.

*Corresponding author

Keywords: *Acinetobacter baumannii*; Murine model, Virulence factors, Antibiotic treatments,
Polymicrobial infections, Pathogenesis

Abstract

Acinetobacter baumannii can cause prolonged infections that disproportionately affect immunocompromised populations. Our understanding of A. baumannii respiratory pathogenesis relies on an acute murine infection model with limited clinical relevance that employs an unnaturally high number of bacteria and requires the assessment of bacterial load at 24-36 hours post-infection. Here, we demonstrate that low intranasal inoculums in immunocompromised mice with a tlr4 mutation leads to reduced inflammation, allowing for persistent infections lasting at least 3 weeks. Using this "chronic infection model," we determined the adhesin InvL is an imperative virulence factor required during later stages of infection, despite being dispensable in the early phase. We also demonstrate that the chronic model enables the distinction between antibiotics that, although initially reduce bacterial burden, either lead to complete clearance or result in the formation of bacterial persisters. To illustrate how our model can be applied to study polymicrobial infections, we inoculated mice with an active A. baumannii infection with Staphylococcus aureus or Klebsiella pneumoniae. We found that S. aureus exacerbates the infection, while K. pneumoniae enhances A. baumannii clearance. In all, the chronic model overcomes some limitations of the acute pulmonary model, expanding our capabilities to study of A. baumannii pathogenesis and lays the groundwork for the development of similar models for other important opportunistic pathogens.

Introduction

Acinetobacter baumannii is a Gram-negative opportunistic pathogen that causes diverse infections including pneumonia, urinary tract infection (UTI), bone and soft tissue infection, and septicemia (1–5). While becoming an increasingly more common cause of community-acquired infections, A. baumannii still primarily causes hospital-acquired infections in critically ill and immunocompromised patients, ~25% of which are polymicrobial (6–11). These infections are associated with an alarming mortality rate, up to 80% in some populations, largely owing to extremely high rates of multi-drug resistance (8, 12, 13). Notably, A. baumannii isolates exhibit the highest rates of multi-drug resistance of all Gram-negative pathogens, leading the World Health Organization to classify the bacterium at its highest priority for research and development of new treatments (13, 14). There is consequently an urgent need to better understand the virulence mechanisms employed by A. baumannii to guide the development of novel therapeutic approaches to combat infections.

While A. baumannii can cause a variety of infections, it is most commonly associated with pneumonia (4, 15). In fact, A. baumannii causes up to 10% of all hospital-acquired pneumonia (HAP) cases in the United States, highlighting its importance in clinical settings (16, 17). Despite this, little is known regarding the pathogenesis of this bacterium in the respiratory tract (18). A major hindrance in the ability to investigate A. baumannii pneumonia is the lack of available clinically-relevant murine infection models. This is, in large part, due to the low virulence of most strains in immunocompetent mice. This is a shared feature among many pathogens that commonly cause HAP, including Pseudomonas aeruginosa and Staphylococcus aureus, for which animal models closely mimicking human infection are not available (19, 20). An acute infection model requiring a very high, and rather artificial, inoculum of 10^8-10^9 bacteria introduced intranasally or intratracheally is most often used to investigate these pathogens (10, 20, 21). Wild-type (WT) mice will typically either succumb to infection or clear the organism by 72 h, thus

requiring early readouts of infection such as bacterial pulmonary titers at 24-36 h. While this

model may serve as a useful tool to study pathogenesis early during infection, the quick bacterial

clearance does not allow for the study of bacterial virulence mechanisms at later timepoints.

Importantly, *A. baumannii* respiratory infection in humans results in an average length of hospital

stay of ~30 days, and this number is much higher in cases caused by multi-drug resistant strains,

highlighting the need for a long-term infection model (22, 23). In this pursuit, some laboratories

have used antibody or cyclophosphamide treatments to render mice neutropenic (24–30). These

treatments initially make mice more susceptible to *A. baumannii* infection, enabling the study of

bacterial pathogenesis up to 7 d̲ p̲ost-i̲nfection (dpi) using lower inoculums (~10^7 bacteria).

However, these models do not achieve stable neutropenia in mice which leads to clearance of

infection. To maintain neutropenia over longer periods, multiple injections would be necessary,

which can lead to fluctuating neutrophil levels, thereby altering the overall course of disease. A

notable caveat to many reports using these currently available immunocompetent and

immunocompromised infection models is that older, lab-domesticated strains and non-lung

isolates, such as *Ab*19606 and *Ab*17978, are employed, despite the extensive literature

demonstrating numerous genotypic and phenotypic differences between these and modern

respiratory isolates (31–35). In all, there is an urgent need for alternative infection models to

study bacterial pathogenesis during long-term infection by relevant clinical isolates.

77   Previous reports have used genetically immunocompromised mice to study the role of the

host immune response to *A. baumannii* infection. One example is mice carrying a mutation in

t̲oll-l̲ike r̲eceptor 4 (TLR4). TLR4 recognizes the lipid A moiety of bacterial l̲ip̲op̲oly̲s̲accharide

(LPS) and l̲ip̲oo̲lig̲os̲accharide (LOS), the main component of the outer membrane of most Gram-

negative bacteria (36–38). The recognition of lipid A by TLR4 triggers a signaling cascade through

MyD88- or TRIF-dependent pathways, resulting in increased inflammatory cytokine and type 1

interferon production, respectively (39). The role of TLR4 during *A. baumannii* infection has been

examined in murine septicemia, acute pneumonia, UTI, and catheter-associated UTI (CAUTI)

models (40–42). In the acute pneumonia model, Knapp et al. showed that *tlr4* mutant mice had

increased *A. baumannii* CFU in the lungs with reduced inflammatory cytokines compared to WT

mice (41, 43). Using a bloodstream infection model, Lin et al. demonstrated that WT C3H/FeJ

and *tlr4* mutant C3H/HeJ mice had similar bacterial burdens (40, 44). However, all WT mice

succumbed to infection by day 4, whereas all *tlr4* mutant mice survived. This could be attributed

to WT mice experiencing septic shock associated with increased inflammatory cytokines. Finally,

in a UTI model, our laboratory found that *tlr4* mutant C3H/HeJ mice were more susceptible to

infection than WT C3H/HeN mice (42). Moreover, we found that C3H/HeJ mice in the UTI model

formed small intracellular populations in urothelial cells referred to as *Acinetobacter baumannii*

intracellular reservoirs (ABIRs), which could seed a recurrent infection upon catheterization at

higher rates relative to WT mice. In addition to playing a significant role during murine infection,

TLR4 is relevant in clinical settings as well. In fact, numerous studies have identified links

between *tlr4* polymorphisms and infection outcomes from *A. baumannii* pneumonia (45–47). In

all, these studies demonstrate the key role of TLR4 in controlling *A. baumannii* infection and

disease progression and highlight the clinical relevance of the associated signaling cascade.

In this work we describe a novel murine model of *A. baumannii* pneumonia that employs

tlr4 mutant mice and low bacterial inoculums (10^5 bacteria). Using this model, we show that

clinically-relevant *A. baumannii* strains can establish chronic infection.   We additionally

demonstrate that our model enables the discovery of virulence factors not detectable in the acute

infection model. Finally, we illustrate how our model can be employed to assess the efficacy of

antibiotics over the course of infection and investigate polymicrobial infections.

Results

tlr4 mutant mice are susceptible to chronic infection at low inoculums.

To assess if *tlr4* mutant mice could serve as permissive hosts for long-term respiratory

infection, we performed intranasal inoculations of WT (C3H/HeN) and *tlr4* mutant (C3H/HeJ) mice

with high (10^8) and low (10^5) inoculums of a modern *A. baumannii* respiratory isolate, G636, and

sacrificed groups of mice every 3 days starting at 24 hours post-infection (hpi). At the higher

inoculum, WT mice cleared infection by day 4, consistent with previously published results using

the acute pulmonary infection model (**Fig. 1A**) (21). *tlr4* mutant mice infected with the higher

inoculum also cleared infection relatively early after inoculation, with most mice having no

detectable bacteria in the lungs by ~7 dpi. Strikingly, while WT mice infected with the lower dose

of 10^5 bacteria cleared infection after 1 day, *tlr4* mutant mice maintained detectable bacteria in

the lungs out to the latest timepoint tested, 19 dpi (**Fig. 1D**). Despite this long infection course,

dissemination to distal organs was rarely detected.   This is consistent with the clinical

manifestations of non-ventilator *A. baumannii* pneumonia, as less than 20% of patients will

develop subsequent bacteremia (48). Notably, by employing confocal microscopy, we were able

to visualize bacteria in *tlr4* mutant mice with the low inoculum; at early timepoints (4 hpi and 2

dpi), bacteria were identified inside cells in the bronchoalveolar lavage fluid (BALF), as well as

extracellularly, consistent with our previously published results in the acute infection model (**Fig.**

S1) (31).

We then assessed if a second modern *A. baumannii* respiratory isolate, G654, behaves

similarly to G636 (**Fig. 1E**). Again, detectable levels of bacteria were present in the lungs out to

19 dpi with the low inoculum in *tlr4* mutant mice, while WT mice cleared this inoculum within 1

day. At the higher inoculum, G654 was cleared soon after inoculation regardless of TLR4

functionality (**Fig. 1B**). On the contrary, when we tested an older, lab-domesticated urinary isolate

that is commonly used to study *A. baumannii* respiratory pathogenesis, *Ab*19606, we found that

WT and *tlr4* mutant mice cleared infection after 1 day regardless of inoculum size or mouse

background (**Fig. 1C and 1F**). In all, these results indicate that modern *A. baumannii* respiratory

isolates can cause infection out to nearly 3 weeks at lower and likely more clinically-relevant

inoculums than previously used in the literature. The finding that *Ab*19606 was unable to establish

long-term infection further highlights the differences between modern, infection site-specific

isolates and lab-domesticated strains (31–35).   Importantly, this infection duration with low

inoculums of modern respiratory isolates in *tlr4* mutant mice is the longest reported for *A.*

baumannii in any animal model to date.   We therefore chose to further characterize these

conditions as a model to study pulmonary pathogenesis, referred to hereafter as the "chronic

respiratory infection model."

Lower A. baumannii inoculums result in a decreased immune response in tlr4 mutant mice.

Given the unexpected result that *tlr4* mutant mice exhibit chronic infection at lower

inoculums, while WT and *tlr4* mutant mice clear infection at higher inoculums, we sought to

characterize the host immune response in these different conditions. We intranasally infected

groups of WT and *tlr4* mutant mice with 10^5 or 10^8 bacteria or mock infected them with phosphate-

buffered saline (PBS). Then, at early timepoints of 4 hpi and 2 dpi and a later timepoint of 7 dpi,

BALF was collected for immune cell quantification (**Fig. 2**). Regardless of timepoint, inoculum, or

mouse background, few significant changes were observed in number of alveolar macrophages

(AMs) (**Fig. 2A-2C**).   In WT mice, the number of polymorphonuclear leukocytes (PMNs) was

increased with the higher inoculum relative to lower and mock inoculums at every timepoint (**Fig.**

2D-2F). Additionally, at the higher inoculum, WT mice had increased PMNs relative to *tlr4* mutant

mice at every time point in line with previous results (41). Interestingly, at the lower inoculum,

while PMN counts trended higher at 4 hpi for WT mice relative to *tlr4* mutant mice, no significant

differences were noted between these groups at any timepoint. This result suggests that, despite

neutrophil influx being a predominant mechanism of *A. baumannii* clearance, PMN numbers alone

may not account for differences in clearance between WT and *tlr4* mutant mice at the lower

inoculum (49–55).

To further evaluate the host response, we quantified 13 common inflammatory cytokines

in the BALF (**Table S1**). At the higher inoculum, WT and *tlr4* mutant mice exhibited significantly

increased levels of IL-1α, IFN-γ, TNF-α, MCP-1, IL-1β, IL-6, and IL-17A early during infection

relative to the lower inoculum while levels dissipated by 7 dpi, consistent with bacterial clearance

(See **Fig. 1A**). WT mice infected with the high inoculum had significantly increased levels of IFN-

β relative to *tlr4* mutant mice at 4 hpi and increased levels of IL-1α, IFN-γ, TNF-α, MCP-1, IL12-

p70, IL-1β, IL-6, IL-27, and IL17A at 2 dpi, likely leading to the earlier clearance observed. At the

lower inoculum, although WT mice clear infection within nearly 24 h and *tlr4* mutant mice maintain

infection out to at least three weeks, minimal significant differences in inflammatory cytokines

were observed (See **Fig. 1D**). In fact, the only significant difference noted was the increased

levels of GM-CSF at 4 hpi in WT mice relative to *tlr4* mutant mice. Other inflammatory cytokines

that trended higher at early timepoints in WT mice at the lower inoculum include the

inflammasome-associated cytokines IL-1α and IL-1β, as well as TNF-α and IL-6. These elevated

levels of inflammatory cytokines early during infection in WT mice could possibly account for the

earlier clearance. Later during infection, however, *tlr4* mutant mice had elevated, albeit not

significantly higher, amounts of TNF-α, IL-1α, and IL-6 relative to WT mice, consistent with

persistent infection.

The chronic respiratory infection model results in lung pathology.

We next assessed if the chronic respiratory infection model is associated with lung

pathology. *tlr4* mutant mice were infected with 10^5 G636 or mock infected with PBS, and mice

were sacrificed at 4 hpi, 2 dpi, 7 dpi, 14 dpi, and 21 dpi. Lungs were then sectioned, stained with

hematoxylin and eosin (H&E), and scored for pathological changes as previously described (**Fig.**

3, **Fig. S2**, and **Fig. S3**) (56). The chronic model resulted in significant increases in alveolitis,

peribronchiolitis, smooth muscle hypertrophy, squamous epithelium metaplasia, and formation of

bronchus-associated lymphoid tissue (BALT) relative to mock-treated mice (**Fig. 3** and **Fig S2**).

Significant changes in goblet cell hyperplasia and fibrosis were not detected (**Fig. S3**). Of note,

significant signs of disease were detected out to 14 dpi, and, even at 21 dpi, infected mice showed

trends toward increased lung damage relative to mock-infected mice. In all, histopathological

analyses revealed that the chronic respiratory infection model results in sustained lung damage,

indicative of chronic infection.

InvL is a critical virulence factor for long-term infection.

The acute pulmonary infection model has been widely used to characterize *A. baumannii*

virulence factors (21). While the acute model is valuable for identifying bacterial proteins required

at early timepoints, these mice clear infection within 3-4 dpi, not allowing for the identification of

factors required for prolonged infection. As a proof of principle, we sought to determine if the

chronic respiratory infection model could identify proteins required for bacterial persistence in the

lungs. We hypothesized that prolonged adherence to respiratory epithelium would be required

for persistence, so we first tested individual mutants lacking previously identified *A. baumannii*

adhesins (Bap, Ata, FhaBC, and InvL) for attenuation in the chronic infection model (**Fig. S4**) (57–

66). *tlr4* mutant mice were infected with 10^5 G636 WT or mutant bacteria, and mice were

sacrificed at 1 or 14 dpi for lung colony-forming units (CFU) quantification. This experiment

indicated a possible role for InvL in long-term infection as mice began to clear bacteria in the lungs

by 14 dpi. To further confirm the importance of InvL for bacterial persistence, we performed more

extensive analyses with G636 WT, *invL* mutant (Δ*invL*), and complemented *invL* mutant (*invL*⁺)

strains in the chronic infection model, sacrificing mice at 1, 7, 14, and 21 dpi to quantify CFU in

the lungs (**Fig. 4A-D**). Early during infection, the Δ*invL* mutant exhibited only a modest defect.

However, at later timepoints the infection defect became more pronounced, as some mice cleared

the bacteria as early as 7 dpi. By 21 dpi, all but two mice had cleared the Δ*invL* mutant, while the

majority of mice infected with the WT strain still had detectable bacteria in their lungs. Genetic

complementation partially rescued this defect at these later timepoints, as no significant difference

was detected between WT and complemented strains.

We next compared results from the chronic respiratory infection model to the acute

infection model. We infected C57BL/6 mice with 10^9 G636 WT, Δ*invL*, or *invL*+ strains. 24 hpi,

mice were sacrificed, and CFU in the lungs, spleens, and kidneys were quantified (**Fig. 4E-F**). As

opposed to results seen in the chronic infection model, the Δ*invL* mutant had no significant defect

in bacterial load in the lungs. Additionally, no defect was noted in dissemination to the spleen and

kidneys, indicating that InvL is dispensable in the acute infection model. In all, these results

highlight the differences in required bacterial genes between these disparate pulmonary infection

models and show the importance of continuing to explore models that can better approximate

clinical disease. Additionally, these experiments establish InvL as the first known *A. baumannii*

virulence factor required for long-term infection.

The chronic infection model can be used to study the outcome of antibiotic treatment.

The acute pulmonary infection model has been employed extensively to assess effects of

antibiotic treatment (21). However, this model only allows us to estimate the efficacy of antibiotics

by measuring the initial reduction in the bacterial burden at 24-36 hpi due to rapid bacterial

clearance by the host. A clear limitation of this model is that it does not inform if bacterial infection

is cleared, or if persistent bacteria remain in the lung. The chronic respiratory infection model

therefore represents a novel platform that could be used to track the kinetics of *A. baumannii*

clearance due to antibiotic treatment. As a proof of principle, we assessed the effect of tigecycline,

colistin, and imipenem in the chronic model with strains G636 and G654 at antibiotic

concentrations similar to those previously used to determine treatment efficacy in mice (**Fig. 5A-**

B and **Fig. S5A-D**) (67–73). We additionally assessed the effect of apramycin, a drug with

demonstrated efficacy and safety in mice that is currently in Phase I clinical trials for use in

humans (**Fig. 5C-D**) (74–76). Minimum inhibitory concentrations (MICs) for G636 and G654 for

these and other commonly used antibiotics are listed in **Table S2**. Colistin was ineffective for both

strains in the acute infection model, while initial reductions in CFU were noted in the chronic

infection model (**Fig. S5A-B**). However, bacterial numbers appeared to stabilize over time in the

chronic infection model, consistent with the development of bacterial persisters (discussed

below). Imipenem showed limited efficacy against both strains in both models (**Fig. S5C-D**), as

expected given the strains' resistance *in vitro* (**Table S2**). At 24 hpi, tigecycline and apramycin

treatment resulted in initial reductions in CFUs in both the chronic and the acute infection models

relative to PBS-treated mice (**Fig. 5A-D**). However, the chronic model enabled us to differentiate

the efficacy of both antibiotics at later times. Apramycin treatment ultimately led to clearance after

3-5 days, demonstrating the efficacy of this antibiotic. However, with tigecycline treatment,

although there were initial reductions in CFU, bacterial numbers leveled out over time indicative

of treatment failure. The behavior of bacteria in presence of tigecycline over time is consistent

with the development of persisters. Notably, the efficacy of tigecycline and apramycin against *A.*

baumannii cannot be distinguished at 24 hpi. These results indicate that the chronic model can

be used to determine outcome of infection with therapeutic intervention, a significant advantage

over the currently employed acute infection model.

Use of the chronic infection model to study bacterial co-infections reveals that Staphylococcus

aureus exacerbates ongoing A. baumannii infection while Klebsiella pneumoniae leads to earlier

clearance.

Approximately 25% of *A. baumannii* pulmonary infections are polymicrobial, and two of

the most commonly co-infecting pathogens are *Staphylococcus aureus* and *Klebsiella*

pneumoniae (9). We thus sought to assess the impact of secondary infections with these two

bacteria on the outcome of *A. baumannii* infection in the context of the chronic respiratory infection

model. For these experiments, we first established a primary *A. baumannii* infection by

inoculating *tlr4* mutant mice with 10^5 CFU of strain G636. Following 14 days of *A. baumannii*

infection, we inoculated mice with 5 x 10^7 CFU of *S. aureus* strain Newman or *K. pneumoniae*

strain TOP52, mock-treated mice with PBS, or left mice untreated. One and two days post-

secondary infection, mice were sacrificed, and bacterial CFU were quantified in the lungs,

spleens, and kidneys (**Fig. 6**, **Fig. S6**, and **Fig. S7**). Secondary infection with *S. aureus* led to a

resurgence of *A. baumannii* CFU in the lungs of many mice, though the overall mean CFU in

these mice were not significantly different from mock-infected and untreated groups (**Fig. 6A** and

Fig. 6D-E). *A. baumannii* were also identified in the spleens and kidneys of some mice that

received the secondary *S. aureus* infection, even though *A. baumannii* bacteremia rarely occurs

in the context of this chronic respiratory infection model (**Fig. S6**). Additionally, *S. aureus* trended

toward increased numbers in the lungs, spleens, and kidneys in the context of polymicrobial

infection with *A. baumannii* relative to monomicrobial infection (**Fig. S7A-C**). Notably, two mice

succumbed to *A. baumannii-S. aureus* polymicrobial infection ~24 hpi following the secondary

inoculation, an outcome that did not occur with monomicrobial infection with either bacterium.

Contrarily, secondary infection with *K. pneumoniae* significantly decreased *A. baumannii* CFU in

the lungs relative to mock-infected and untreated groups (**Fig. 6A** and **Fig. 6D-E**). Additionally,

polymicrobial infection with *A. baumannii* and *K. pneumoniae* resulted in significantly reduced *K.*

pneumoniae CFU recovered in the lungs, spleens, and kidneys of mice relative to *K. pneumoniae*

monomicrobial infection (**Fig. S7D-E**). Although understanding the interactions between these

bacteria is beyond the scope of this work, these experiments indicate that *S. aureus* exacerbates

A. baumannii infection, while *K. pneumoniae* attenuates infection in the context of the chronic

infection model. Additionally, these results demonstrate the ability of the model to be used to

study longer-term aspects of polymicrobial interactions that were not previously able to be done

with the acute infection model.

Discussion

A. baumannii has emerged as a significant cause of nosocomial pneumonia and is of

major clinical importance due to its extremely high rates of multidrug resistance (8, 12, 13).

Despite this, our understanding of *A. baumannii* respiratory pathogenesis is hindered by a

shortage of clinically relevant infection models. Here, we aimed to address this significant gap in

the field by developing a novel respiratory infection model. In this pursuit, we found that, at likely

more clinically-relevant inoculums, *tlr4* mutant mice maintain long-term respiratory infections by

A. baumannii. We then demonstrate the versatility of this model which enabled i) the identification

of a bacterial virulence factor required for long-term respiratory infection, which is not required in

acute models, ii) the study of kinetics of bacterial clearance upon treatment with clinically-relevant

antibiotics, and iii) the exploration of the impact of secondary infections with two commonly co-

isolated respiratory pathogens.

In this study, we found that InvL is required for chronic infection, and, more importantly, at

the later stages of infection. However, InvL was dispensable in the context of the acute infection

model. There are multiple possible reasons for this discrepancy. First, the massive bacterial dose

required for the acute infection model may mask potential defects that can now be detected with

a smaller, more clinically-relevant inoculum. This is unlikely, as WT and *invL* mutant bacteria

behave similarly at early time points in our model. An alternative reason could be that adhesins

required early during infection/interaction with the healthy airway differ from those required during

persistent interaction with a more inflamed or damaged airway. It is well-established that the

airway e̲xtra̲cellular m̲atrix (ECM) is altered by bacterial infection, lung damage, and/or

inflammation (91, 92). Long-term lung damage and inflammation results in increased fibronectin,

collagen, laminin, and fibrinogen in the ECM (93–98). Moreover, specific pathogens elicit different

inflammatory responses, resulting in distinct changes to the lung ECM. For example, in an acute

mouse model of pneumonia, *Pseudomonas aeruginosa* induces versican deposition in the lungs,

while *Escherichia coli* induces robust versican and hyaluronan deposition (99, 100). We

previously showed that InvL can bind α5β1 integrin, collagen V, and fibrinogen (66). However,

whether *A. baumannii* infection or the associated inflammation induces production of these

protein(s) during pulmonary infection is unknown. Future work will investigate this possibility, as

well as assess which InvL-host protein interactions are essential for chronic infection.

Herein, we demonstrate the potential to use the chronic respiratory infection model to

study the efficacy of antibiotic treatments over time. One intriguing finding from these experiments

is that with antibiotics such as colistin and tigecycline, an initial decrease in CFU (~10-100 fold)

recovered from the lungs at 1 dpi was observed. However, following this decrease, the number

of bacteria in the lungs appeared to stabilize over time. It is tempting to speculate that this is the

result of the formation of bacterial persisters, defined as bacterial cells that become tolerant to

antibiotics despite undergoing no genetic changes (101–103). Importantly, the commonly used

acute infection model does not allow for the study of bacterial persisters due to the short time

course of the model. Given that persister cells represent a major cause of treatment failure and

chronic infection, the chronic infection model presented here represents a unique platform that is

desperately needed to understand this aspect of *A. baumannii* pathogenesis. Furthermore, our

model offers new possibilities to study efficacy of novel antibiotics in murine models before

committing to expensive clinical trials.

While a significant portion of *A. baumannii* infections are polymicrobial, the acute infection

model has limitations for use with polymicrobial infections. First the quick clearance of the

bacteria usually only allows inoculation at a single timepoint, thus not enabling investigation of

secondary infections. Second, the high required infectious dose often means that typical

inoculums for bacteria used in these experiments must be adjusted, so mice do not succumb to

infections at early timepoints. Here, we applied the chronic respiratory infection model to assess

the result of secondary infection with two pathogens commonly co-isolated with *A. baumannii*, *S.*

aureus and *K. pneumoniae*. We found opposite results with these different bacteria; *S. aureus*

secondary infection trended toward exacerbation of *A. baumannii* infection, while *K. pneumoniae*

secondary infection led to reduced *A. baumannii* numbers. The potential synergism of *A.*

baumannii and *S. aureus* in the chronic infection model aligns with previous reports. For example,

using a Tn-Seq-based approach, Li et al. demonstrated that the 49% of genes required by *S.*

aureus for monomicrobial infection in a murine systemic infection model became non-essential

upon *A. baumannii* co-infection (104). Another recent report showed that *S. aureus* can support

A. baumannii growth *in vitro* by providing acetoin as a carbon source (105).

Although we found that *K. pneumoniae* secondary infection led to reduced *A. baumannii*

numbers in the lungs in the chronic infection model, one study has shown that *K. pneumoniae*

could cross-feed *A. baumannii* through products of sugar fermentation *in vitro* and demonstrated

that co-infection led to reduced survival of *Galleria mellonella* relative to monomicrobial infection

with either pathogen (106). This, in part, shows that these two bacteria can have beneficial

interactions. There are two potential reasons however for the reduction of CFU for both bacteria

in the context of the chronic infection model reported here; i) bacterial competition or ii) the host

response to the secondary infection. Regarding bacterial competition, there have been several

lines of evidence pointing to direct bacterial killing between diverse *A. baumannii* and *K.*

pneumoniae strains mediated by the type VI secretion system (107–110). In addition to direct

killing, this bacterial competition could be indirect as well, as both *A. baumannii* and *K.*

pneumoniae may be competing for similar nutrients in the lung microenvironment. With respect

to the immune response, a difference between this work and the above study is that the

microenvironment encountered in the mammalian lung is not perfectly modeled by the wax moth

(111). Our results may therefore be the result of TLR4-independent host response elicited by the

combination of both bacteria that is not recapitulated by a *G. mellonella* model. While

understanding the precise mechanism behind the *in vivo* interactions between *A. baumannii* and

commonly co-isolated pathogens is outside the scope of the current study, these results highlight

the practicality of applying the chronic respiratory infection model to better understand

polymicrobial infections.

In this study, we have validated several different uses for the chronic respiratory infection

model. However, there are also other potential uses for this model that were not previously

investigable. For example, we can now perform experiments differentiating between virulence

factors required for establishment of infection and factors required for maintenance of infection,

assessing bacterial evolution during long-term infection, investigating changes in the pulmonary

microbiome due to infection over time, and analyzing the long-term outcomes of novel therapies

such as newly developed phage cocktails. Additionally, while this model was initially developed

to study *Acinetobacter* respiratory infections, it has the potential to be applied to research with

other respiratory pathogens in cases where suitable animal models are lacking. In all, this work

describes the longest-term infection model available to investigate *A. baumannii* host-pathogen

interactions to date, which will ultimately aid in the development of novel therapeutics to combat

infection by this increasingly multidrug-resistant bacterium.

Materials and Methods

Bacterial plasmids, strains, and growth conditions.

Plasmids and strains used in this study are detailed in **Table S3**. Bacterial cultures were grown at 37°C in Lennox broth/agar supplemented with 10 µg/mL chloramphenicol, 50 µg/mL apramycin, 100 µg/mL ampicillin, 50 µg/mL kanamycin, 10 µg/mL tetracycline, or 10% sucrose when appropriate.

Murine pneumonia models.

All animal experiments were approved by the Washington University Animal Care and Use Committee, and we have complied with all relevant ethical regulations. The acute pneumonia model was performed similar to previously described experiments (21, 112). Briefly, overnight cultures were subcultured at a 1:200 dilution and grown shaking at 37°C for 3 h to mid-exponential growth phase. Six- to eight-week-old female C57BL/6 mice (Charles River Laboratories, Wilmington, MA) anesthetized with 4% isoflurane were intranasally inoculated with 10^9 CFU that were twice-washed in PBS. At 24 hpi, mice were sacrificed, and CFU in the lungs, spleen, and kidneys were quantified by serial dilution plating the homogenized organs. For experiments with C3H/HeN (Envigo International Holdings, Indianapolis, IN) and C3H/HeJ (Jackson Laboratory, Bar Harbor, ME) mice, *A. baumannii*, *S. aureus*, and *K. pneumoniae* inoculums were prepared and mice were intranasally inoculated as described above, with the exception that inoculums of 10^5 and 10^8 CFU were used for *A. baumannii*, and 5 x 10^7 CFU was used for *S. aureus* and *K. pneumoniae*. Following, at the indicated timepoints, mice were sacrificed and bacteria in the lungs, spleen, and kidneys were quantified as described above. For co-infections, *A. baumannii* was distinguished from *S. aureus* and *K. pneumoniae* by plating on LB agar supplemented with 10 µg/mL chloramphenicol. For antibiotic treatment experiments the indicated mice were treated

intraperitoneally with PBS or 100 mg/kg tigecycline every 12 h, PBS or 5 mg/kg colistin every 8

h, PBS or 500 mg/kg apramycin every 12 h, or PBS or 100 mg/kg imipenem every 12 h with all

treatments beginning 4 hpi. Antibiotics for intraperitoneal treatments were dissolved in PBS, and

the injection volume was 100 µl.

Flow cytometry.

Flow cytometry was performed similarly to previously described methods (31). Briefly,

BALF samples were collected in PBS supplemented with 1 mM EDTA, and cells were collected

by centrifugation at 300 x *g* for 5 min. Cells were then resuspended in Pharm Lyse Buffer (BD

Biosciences, Franklin Lakes, NJ) and incubated for 3 min at room temperature to lyse red blood

cells. Cells were subsequently washed in fluorescence-activated cell sorting (FACS) buffer (PBS

supplemented with 1% heat inactivated fetal bovine serum and 0.1% sodium azide) and blocked

with TruStain FcX PLUS (BioLegend, San Diego, CA) for 15 min at 4°C. Samples were then

stained with anti-CD45-BV605 (BioLegend), anti-CD11c-APC (BioLegend), anti-SiglecF-

PerCP5.5 (BioLegend), and anti-Ly6G-BV421 (Biolegend) for 30 min at 4°C. Following, cells

were washed in FACS buffer and fixed in 2% paraformaldehyde (PFA). Samples were read on a

LSR II Fortessa cytometer (BD Biosciences) or an Aurora cytometer (Cytek Biosciences, Fremont,

CA). Total cell counts in the BALF were calculated using Precision Count Beads (BioLegend)

according to the manufacturer's instructions.

Antibiotic protection assays.

Antibiotic protection assays were performed as previously described (31). To determine

the number of total and intracellular bacteria present in BALF from *A. baumannii* infected mice,

two 500 µl aliquots of lavage fluid were centrifuged at 4100 x *g* for 5 min. Pelleted cells were

resuspended in warm Dulbecco's Modified Eagle Medium (DMEM) (total bacteria) or DMEM with

colistin (50 µg/mL) (intracellular bacteria) and incubated for 1h at 37°C. Samples were then

washed three times with PBS and lysed with 500 µL of Triton X-100 (0.05%). CFUs were

determined by serial dilutions of the bacterial suspensions. The remaining lungs following BALF

collection were also homogenized, and CFUs were quantified by serial dilution plating.

Cytospin of BALF cells.

Cytospin of BALF cells was performed similar to previously described work (31). BALF

samples were centrifugated at 300 x *g* for 5 min, and the pellets were resuspended in 1 mL Pharm

Lyse Buffer (BD Biosciences) and incubated for 5 min on ice to lyse red blood cells. 9 mL of PBS

was added to stop the lysis, viability was determined using Trypan Blue solution (Sigma-Aldrich,

St. Louis, MO), and cells were counted using the TC20 Automated Cell Counter (Bio-Rad

Laboratories, Hercules, CA). Samples were centrifugated at 300 x *g* for 6 min onto CytoPro Poly-

L-Lysine Coated Microscope Slides (ELITechGroup Inc., Logan, UT) using a Cytospin

Cytocentrifugue (Fisher Scientific, Hampton, NH). The slides were air-dried overnight at 4°C and

fixed in 4% PFA for 30 min at room temperature. Samples were incubated with permeabilizing

and blocking solution (PBS supplemented with 0.1% saponin, 0.5% bovine serum albumin, and

10% heat inactivated fetal bovine serum). Cells were stained with Alexa Fluor 555 Phalloidin (Cell

Signaling Technology, Danvers, MA) and 4',6-Diamidino-2-phenylindole dihydrochloride (DAPI)

solution (Invitrogen) for 1 h at 37°C. After staining, the samples were rinsed with washing solution

[PBS supplemented with 0.1% saponin and 0.5% bovine serum albumin (BSA)], and then rinsed

with water and mounted on a coverslip in ProLong Gold Antifade Mountant (Invitrogen).

Confocal microscopy.

Confocal microscopy was performed as previously described (31, 66). Microscopy slides

were analyzed with a Zeiss LSM880 laser scanning confocal microscope (Carl Zeiss AG,

Oberkochen, Germany) equipped with 405nm diode, 488nm Argon, 543nm HeNe, and 633nm

HeNe lasers. A Plan-Apochromat 63X DIC objective and ZEN black 2.1 SP3 software were used

for image acquisition. Images were analyzed using ImageJ software (National Institutes of Health,

Bethesda, MD) (113).

Cytokine analysis.

BALF was collected and centrifuged at 300 x *g* for 5 min. Supernatant containing

cytokines was then collected and frozen at -20°C until the analysis was performed. Cytokine

levels were determined using the LEGENDplex Mouse Inflammation Panel (13-plex) with V-

bottom Plate (BioLegend) according to the manufacturer's instructions. Samples were read using

an Aurora cytometer (Cytek Biosciences).

Histopathology of lung slices.

Lung slices were prepared, stained, and scored as previously described by Castro et al.

(71). Briefly, lungs were perfused with PBS, inflated with optimal cutting temperature (OCT)

compound (Fisher Scientific) diluted in 4% PFA at a 1:1 ratio, snap-frozen, and stored at -80°C

until sectioning. For histology imaging, 4 µm tissue sections were stained with H&E and imaged

with a ZEISS Axioscan 7 Microscope Slide Scanner (Carl Zeiss AG). Lung tissues were blindly

scored on a scale of 0 to 3 for alveolitis, peribronchiolitis, smooth muscle hypertrophy, squamous

epithelium metaplasia, BALT formation, goblet cell hyperplasia, and fibrosis. Area affected was

quantified, multiplied by previously defined intensity scores, and the resulting weighted scores are

reported.

Generation of constructs and strains used in this study.

Primers used in this study are listed in **Table S4**. DNA fragments were assembled using

either the In-Fusion HD EcoDry Cloning Kit (TaKaRa Bio, Mountain View, CA) or NEBuilder HiFi

DNA Assembly Master Mix (New England Biolabs, Ipswich, MA). To generate the vector for

generation of the *invL* mutational construct, pEX18Tc was amplified without the tetracycline

resistance cassette (primers: 5' pEX18 marker swap and 3' pEX18 marker swap), the apramycin

resistance cassette was amplified from pKD4-Apr (primers: 5' Apr for pEX18Ap and 3' Apr for

pEX18Ap), and the amplicons were assembled, generating pEX18Ap (114, 115). The pEX18Ap

mutational constructs were then made by amplifying the pEX18Ap vector (primers: 5' pEX18Tc

and 3' pEX18Tc), a ~1000 bp region upstream of the genes of interest (*invL*KO primers: 5' F1

G636 *invL*KO and 3' F1 G636 *invL*KO; *bap*KO primers: 5' F1 G636 *bap*KO and 3' F1 G636 *bap*KO;

*ata*KO primers: 5' F1 G636 *ata*KO and 3' F1 G636 *ata*KO; *fhaBC*KO primers: 5' F1 G636 *fhaBC*KO

and 3' F1 G636 *fhaBC*KO), and a ~1000 bp region downstream of the genes of interest (*invL*KO

primers: 5' F2 G636 *invL*KO and 3' F2 G636 *invL*KO; *bap*KO primers: 5' F2 G636 *bap*KO and 3'

F2 G636 *bap*KO; *ata*KO primers: 5' F2 G636 *ata*KO and 3' F2 G636 *ata*KO; *fhaBC*KO primers: 5'

F2 G636 *fhaBC*KO and 3' F2 G636 *fhaBC*KO), followed by assembly of these amplicons.

Mutational constructs were then transformed into G636, and strains with the integrated plasmid

were selected for by apramycin treatment. Counterselection for double crossover was performed

by plating these strains on LB agar without NaCl supplemented with 10% sucrose. Mutants were

then confirmed by PCR analyses and whole-genome sequencing.

The *invL* complementation construct was generated by amplifying the putative promoter

region (~300 bp upstream) along with the *invL* open reading frame (primers: 5' G636 *fdeC*KO

Comp and 3' G636 *fdeC*KO Comp-His6 v2) and the pUC18T-miniTn7T-Apr vector (primers: Tn7

linear Fwd-His6 and Tn7 liner Rev) (116). These amplicons were then assembled, generating

pUC18T-miniTn7T-Apr::G636 *invL*KO comp. To generate the *gfp* integration construct, the *gfp*

cassette was amplified from PB-FLuc+GFPd2 (primers: 5' d2EGFP for pUC18T-mTn7 and 3'

d2EGFP for pUC18T-mTn7) and pUC18T-mTn7-Apr was amplified (primers: 5' pUC18T-mTn7 for

d2EGFP and 3' pUC18T-mTn7 for d2EGFP). These fragments were then assembled, generating

pUC18T-miniTn7T-Apr::*gfpd2*.   pUC18T-miniTn7T-Apr::G636 *invL*KO comp and pUC18T-

miniTn7T-Apr::*gfpd2* were introduced into G636 Δ*invL* and G636, respectively, using a four-

parental conjugation technique, as previously described (116–119). Selection was achieved using

LB supplemented with apramycin and chloramphenicol, and insertion of the respective fragments

at the mTn7 site in the resulting G636 *invL*$^+$ and G636-*gfp* strains was confirmed by PCR

analyses.

Antibiotic susceptibility assays.

MIC analyses were performed using a two-fold broth dilution microtiter assay similar to

previously described protocols (112, 120, 121). Briefly, overnight cultures were sub-cultured at

0.05 Abs$_{600}$ and grown for 3 h shaking at 37 hpi. Mid-exponential growth phase cultures were

then inoculated at 0.01 Abs$_{600}$ into a 96-well microtiter plate (Corning Inc, Corning, NY) containing

two-fold decreasing dilutions of the indicated antibiotics. Plates were then incubated at 37°C with

shaking for 24 h. The MIC was defined as less than 10% of the Abs$_{600}$ of an untreated control.

Statistical methods.

All statistical analyses were performed using GraphPad Prism version 9, and *P* values of

<0.05 were considered statistically significant. When normally distributed, data sets were

analyzed with Unpaired Student's *t*-tests (comparing two samples), one-way analysis of variance

(ANOVA) with Tukey's test for multiple comparisons (comparing more than two samples), or two-

way ANOVA with Tukey's test for multiple comparisons (comparing more than two samples with

two independent variables). For non-normally distributed data sets, the Mann-Whitney U test

(comparing two samples) or the Kruskal Wallis H test with Dunn's test for multiple comparisons

(comparing more than two samples) was used.

Acknowledgements

This work was supported by funding to M.F.F. (R01AI166359 - CHECK), C.J.L.

(T32AI007172), and C.B.L (R01AI137062) through the National Institute of Allergy and Infectious

Diseases of the National Institutes of Health. JM was supported through The American

Association of Immunologists Careers in Immunology Fellowship Program and The Pediatric

Cardiovascular and Pulmonary Research Training Program (5T32HL125241-07). The modern

respiratory isolates used in this study, G636 (strain 3689) and G654 (strain 6919), were collected

by the CDC-funded Georgia Emerging Infections Program's (EIP) Multi-site Gram-Negative

Surveillance Initiative (MuGSI) and kindly provided by Sarah Satola. We also acknowledge

Jennifer Philips, Jacco Boon, and Gayan Bamunuarachchi for thoughtful discussion about the

manuscript. We thank Wandy Beatty and the Washington University School of Medicine

Molecular Microbiology Imaging Facility for microscopy assistance, Alma Johnson of the

Washington University Center for Reproductive Health Sciences Histocore for lung tissue slide

mounting and staining assistance, and De Chen of the Washington University Center for Cellular

Imaging for assistance with the Zeiss AxioScan Z1. Finally, we thank Dakota Hall for technical

assistance with experiments.

543 Tables

544 Table S1. Cytokine analysis at 4 h, 2 d, and 7 d post-intranasal infection with G636.

Cytokine[a,b]	WT			*tlr4* mutant		
4 h	10^5 G636	10^8 G636	Mock	10^5 G636	10^8 G636	Mock
IL-23	23.26 (10.14)	38.53 (6.36)	10.67 (7.09)	22.57 (13.26)	32.61 (3.61)	15.90 (15.90)
IL-1α	58.57 (18.27)	323.11 (75.42)[#,$]	2.5 (0.43)	2.77 (0.47)	429.19 (90.22)[#,$]	1.49 (0.13)
IFN-γ	0.77 (0.36)	5.13 (1.30)[#,$]	0.00 (0.00)	0.23 (0.23)	4.34 (0.39)[#,$]	0.00 (0.00)
TNF-α	2219.55 (214.72)	14888.73 (311.27)[#,$]	108.38 (66.66)	49.51 (6.47)	12802.47 (3073.56)[#,$]	12.82 (3.58)
MCP-1	0.00 (0.00)	65.73 (15.07)[#,$]	0.00 (0.00)	0.00 (0.00)	65.57 (8.23)[#,$]	0.00 (0.00)
IL-12p70	0.00 (0.00)	1.43 (1.43)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-1β	9.50 (1.95)	46.26 (10.54)[#,$]	0.00 (0.00)	0.00 (0.00)	56.58 (12.58)[#,$]	0.00 (0.00)
IL-10	0.00 (0.00)	4.18 (4.18)	0.00 (0.00)	0.00 (0.00)	2.04 (2.04)	0.00 (0.00)
IL-6	910.41 (172.71)	7885.57 (1645.56)[#,$]	18.19 (5.67)	8.09 (2.69)	6334.72 (1218.67)[#,$]	2.70 (2.70)
IL-27	0.00 (0.00)	85.90 (13.54)[#,$]	0.00 (0.00)	0.00 (0.00)	42.77 (24.83)[$]	0.00 (0.00)
IL-17A	1.05 (0.74)	8.36 (2.46)[#,$]	0.00 (0.00)	0.00 (0.00)	4.66 (0.39)[#,$]	0.00 (0.00)
IFN-β	0.00 (0.00)	29.75 (17.23)[*,#,$]	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
GM-CSF	96.65 (13.37)[*,#]	55.42 (7.19)[#,$]	0.00 (0.00)	0.00 (0.00)	47.41 (6.96)[#,$]	0.00 (0.00)
2 d						
IL-23	19.93 (9.61)	27.05 (8.76)	44.09 (21.87)	26.04 (11.37)	11.35 (5.56)	72.50 (36.30)
IL-1α	1.14 (0.19)	157.28 (58.45)[*,#,$]	1.13 (0.20)	2.00 (0.50)	25.68 (6.15)	19.64 (18.02)
IFN-γ	2.20 (0.98)	212.26 (54.96)[*,#,$]	0.00 (0.00)	3.70 (2.48)	4.96 (2.48)	0.00 (0.00)
TNF-α	1.85 (0.82)	592.26 (154.11)[*,#,$]	1.03 (0.60)	19.79 (6.79)	52.20 (8.86)	5.76 (4.31)
MCP-1	0.00 (0.00)	160.30 (14.27)[*,#,$]	0.00 (0.00)	0.00 (0.00)	33.52 (6.37)[#,$]	0.00 (0.00)
IL-12p70	0.00 (0.00)	46.17 (17.63)[*,#,$]	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-1β	1.06 (1.06)	12.35 (2.80)[*,#,$]	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-10	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-6	0.00 (0.00)	1179.63 (323.29)[*,#,$]	0.00 (0.00)	0.00 (0.00)	20.45 (3.83)	4.29 (4.29)
IL-27	0.00 (0.00)	137.76 (81.43)[*,#,$]	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-17A	0.42 (0.42)	25.31 (10.22)[*,#,$]	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IFN-β	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
GM-CSF	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
7 d						
IL-23	41.43 (14.43)	30.14 (19.51)	20.66 (20.66)	9.07 (5.32)	29.77 (12.87)	12.69 (12.69)
IL-1α	1.05 (0.18)	41.50 (23.66)	1.62 (0.92)	5.54 (4.27)	4.60 (2.50)	1.79 (0.85)
IFN-γ	0.00 (0.00)	53.36 (40.45)[$]	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
TNF-α	0.00 (0.00)	93.32 (36.14)[*,#,$]	0.00 (0.00)	3.17 (3.17)	2.41 (1.54)	0.00 (0.00)
MCP-1	0.00 (0.00)	14.83 (14.83)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-12p70	0.00 (0.00)	1.54 (1.54)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-1β	0.00 (0.00)	1.43 (1.43)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-10	0.00 (0.00)	6.84 (6.84)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-6	0.00 (0.00)	232.41 (182.45)	0.00 (0.00)	5.19 (5.19)	16.72 (16.72)	0.00 (0.00)
IL-27	0.00 (0.00)	26.87 (26.87)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IL-17A	0.00 (0.00)	13.67 (11.48)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
IFN-β	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
GM-CSF	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)

545 [a]Mean pg/ml (SEM) from two independent experiments at each timepoint is displayed.

546 [b]*$P < 0.05$ relative to *tlr4* mutant at same inoculum; #$P < 0.05$ relative to mock in same mouse

547 strain. $$P < 0.05$ relative to 10^5 inoculum in same mouse strain. Two-way ANOVA, Tukey's test

548 for multiple comparisons. Significant differences are also highlighted in green.

550 **Table S2. MICs for *A. baumannii* strains G636 and G654.**

Antibiotic	G636 (Resistant/Sensitive)	G654 (Resistant/Sensitive)	Clinical Breakpoint[a]
Imipenem	>256 µg/ml (Resistant)	>256 µg/ml (Resistant)	2 µg/ml
Ampicillin	>256 µg/ml (N/A)	>256 µg/ml (N/A)	N/A[c]
Ciprofloxacin	>256 µg/ml (Resistant)	>256 µg/ml (Resistant)	1 µg/ml
Levofloxacin	32 µg/ml (Resistant)	128 µg/ml (Resistant)	2 µg/ml
Colistin	1 µg/ml (Intermediate)[b]	8 µg/ml (Resistant)	2 µg/ml[b]
Polymyxin B	2 µg/ml (Intermediate)[b]	4 µg/ml (Resistant)	2 µg/ml[b]
Tigecycline	2 µg/ml (N/A)[c]	1 µg/ml (N/A)[c]	N/A[c]
Gentamicin	>256 µg/ml (Resistant)	2-4 µg/ml (Sensitive)	4 µg/ml
Apramycin	16 µg/ml (N/A)[c]	16 µg/ml (N/A)[c]	N/A[c]

[a]Clinical breakpoints are according to the Clinical and Laboratory Standard Institute (CLSI) M100
Performance Standards for Antimicrobial Susceptibility Testing 30th Edition (122).
[b] A "sensitive" breakpoint is not available for colistin or polymyxin B from the CLSI. Strains with
MICs of less than or equal to 2 µg/ml are considered to have "intermediate resistance."
[c]The clinical breakpoint has not been defined for ampicillin, tigecycline, and apramycin by the
CLSI.

Table S3. Plasmids and strains used in this study.

Plasmid or Strain	Description[a]	Source[b]
Plasmids		
pEX18Tc	Precursor plasmid used for generation of pEX18Ap; Tet^r	(114)
pKD4-Apr	Source for apramycin cassette for mutant generation; Apr^r	(115)
pEX18Ap	Plasmid background used for generation of *A. baumannii* mutants; Apr^r	This study
pEX18Ap::G636 *invL*KO	Plasmid used for mutation of *invL* in G636; Apr^r	This study
pUC18T-miniTn7T-Apr	Vector used for genetic complementation at the mTn7 site; Apr^r	(116)
pUC18T-miniTn7T-Apr::G636 *invL*KO comp	Plasmid used for complementation of the Δ*invL* mutant; Apr^r	This study
PB-FLuc+GFPd2	Plasmid source for *gfp* cassette; Amp^r	b
pUC18T-miniTn7T-Apr::*gfpd2*	Expression vector; Apr^r	This study
pRK2013	Helper plasmid for mobilization of non-self-transmissible plasmids; Kan^r	(123)
pTNS2	T7 transposase expression vector; Amp^r	(124)
Strains		
E. coli		
Stellar	*mrr-hsdRMS-mcrBC* and *mcrA*; Host strain for cloning	TaKaRa
HB101	F- *mcrB mrr hsdS*20(rB- mB-) *recA*13 *leuB*6 *ara*-14 *proA*2 *lacY*1 *galK*2 *xyl*-5 *mtl*-1 *rpsL*20 *glnV*44 λ-; Host strain for pRK2013	Promega
EC100D	F - mcrA Δ(*mrr-hsdRMS-mcrBC*) φ80*dlacZ*ΔM15 Δ*lacX*74 *recA*1 *endA*1 *araD*139 Δ(*ara, leu*)7697 *galU galK* λ- *rpsL nupG pir* +(DHFR); Host strain for pTNS2	Fisher
A. baumannii		
G636	2018 *A. baumannii* respiratory isolate (Strain 3689)	c
G636 Δ*invL*	G636 *invL* mutant	This study
G636 *invL*$^+$	G636 *invL* mutant complemented	This study
G636 Δ*bap*	G636 *bap* mutant	This study
G636 Δ*ata*	G636 *ata* mutant	This study
G636 Δ*fhaBC*	G636 *fhaBC* mutant	This study
G636-*gfp*	G636 expressing *gfpd2*	This study
G654	2020 *A. baumannii* respiratory isolate (Strain 6919)	c
*Ab*19606	1948 *A. baumannii* urinary isolate	(125)
S. aureus		
Newman	1952 osteomyelitis isolate	(126)
K. pneumoniae		
TOP52	2006 cystitis isolate	(127)

559 [a]Tet, tetracycline; Apr, apramycin; Amp, ampicillin; Kan, kanamycin.

567 **Table S4. Primers used in this study.**

Primer	Sequence
5' pEX18 marker swap	ACACGGTGCCTGACTGCGTTAGC
3' pEX18 marker swap	ATGGAAGCCGGCGGCACC
5' Apr for pEX18Ap	GAGGTGCCGCCGGCTTCCATGATCCTCAGCCAATCGACTGGC
3' Apr for pEX18Ap	AACGCAGTCAGGCACCGTGTGATTCCCTTTGTCAACAGCAATGG
5' pEX18Tc	ATGCCTGCAGGTCGACTCTAGAGG
3' pEX18Tc	GCAAGCTTGGCACTGGCCGT
5' F1 G636 *invL*KO	ACGGCCAGTGCCAAGCTTGCGGCAATGTCTCAAATAAAAAATTTAACTC
3' F1 G636 *invL*KO	TGAGATCCGCTATTATTACTTCCAG
5' F2 G636 *invL*KO	AGTAATAATAGCGGATCTCATGCTTCTTTTTTTAGAGTTGTGTTCC
3' F2 G636 *invL*KO	TAGAGTCGACCTGCAGGCATAAAATAACCGCATAGCCAGCTTGAGC
5' G636 *fdeC*KO Comp	GCATGAGCTCACTAGTGGATCCGAGATTAAGACTTTACTTGGCATACACC
5' F1 G636 *bap*KO	ACGGCCAGTGCCAAGCTTGCAGAAGCGGCTGGCAATGTCACG
3' F1 G636 *bap*KO	TCAAGCACCGGTGCATACTGACC
5' F2 G636 *bap*KO	CAGTATGCACCGGTGCTTGAGGTGGTAACACTACAATTCAGATTGACC
3' F2 G636 *bap*KO	TAGAGTCGACCTGCAGGCATTCCATAAATGAATTTGCCATTTTCTTGAATCTG
5' F1 G636 *ata*KO	ACGGCCAGTGCCAAGCTTGCTAAGTCGGTCTGGCTATTCGCC
3' F1 G636 *ata*KO	TGATGACGTTGAGAAAAAGCTAATGCAGG
5' F2 G636 *ata*KO	CTTTTTTCTCAACGTCATCAAAAACTTCTCAGACAAATACCGAACTCAACG
3' F2 G636 *ata*KO	TAGAGTCGACCTGCAGGCATCGTGATCAATTTCTTCTGTAAGCGAATCTTTTTGC
5' F1 G636 *fhaBC*KO	ACGGCCAGTGCCAAGCTTGCTTAAAATTTTAAAGCAGTTTGATGAGCC
3' F1 G636 *fhaBC*KO	CAGAATTGTACGTATAAGAACTTTATTTTACAC
5' F2 G636 *fhaBC*KO	TTCTTATACGTACAATTCTGTAACAATGAAAAATGCACATGCGG
3' F2 G636 *fhaBC*KO	TAGAGTCGACCTGCAGGCATCTTTATTGGTACCCTGATTGCG
3' G636 *fdeC*KO Comp-His6 v2	TCAGTGGTGATGGTGATGATGATTACCATTTGAACAGTTTGGATCTATTCC
Tn7 linear Fwd-His6	CATCATCACCATCACCACTGAAAGCTTGGGCCCGGTACCTC
Tn7 linear Rev	GGATCCACTAGTGAGCTCATGC

5' d2EGFP for pUC18T-mTn7	AGAAAGAGGAGAAATACTAGATGGTGAGCAAGGGCGAGG
3' d2EGFP for pUC18T-mTn7	GAGGTACCGGGCCCAAGCTTCTACACATTGATCCTAGCAGAAGC
5' pUC18T-mTn7 for d2EGFP	AAGCTTGGGCCCGGTACCTCG
3' pUC18T-mTn7 for d2EGFP	CTAGTATTTCTCCTCTTTCTCTAGTAATTGTTATCC

Figure Legends

Figure 1. Low inoculums of modern respiratory *A. baumannii* clinical isolates result in chronic lung infection in *tlr4* mutant mice. Groups of C3H/HeN (WT) or C3H/HeJ (*tlr4* mutant) mice were intranasally inoculated with 10^8 G636 (A), 10^8 G654 (B), 10^8 *Ab*19606 (C), 10^5 G636 (D), 10^5 G654 (E), or 10^5 *Ab*19606 (F). Beginning at 24 hpi, groups of mice were sacrificed every three days, and bacteria in the lungs were quantified. Each data point indicates and individual mouse.

Figure 2. Lower intranasal *A. baumannii* inoculums result in reduced lung neutrophil influx. Groups of C3H/HeN (WT) or C3H/HeJ (*tlr4* mutant) mice were intranasally inoculated with 10^5 G636, 10^8 G636, or mock inoculated with PBS. At 4 h (A and D), 2 d (B and E), and 7 d (C and F) pi, alveolar macrophages (AMs) (A-C) and polymorphonuclear leukocytes (PMNs) (D-F) in the BALF were enumerated by flow cytometry. Shown are pooled results from at least two independent experiments, and each data point represents an individual mouse. The horizontal line represents the mean, and the standard error of the mean (SEM) is indicated by error bars. *P < 0.05; two-way analysis of variance (ANOVA), Tukey's test for multiple comparisons.

Figure 3. The chronic respiratory infection model results in lung pathology. Groups of C3H/HeJ (*tlr4* mutant) mice were inoculated with 10^5 G636 or mock-inoculated with PBS, and at 4 hpi, 2 dpi, 7 dpi, 14 dpi, and 21 dpi, lungs slices were prepared, H&E stained, and scored for alveolitis (A), peribronchiolitis (B), smooth muscle hypertrophy (C), squamous epithelium metaplasia (D), and BALT formation (E). The mean is shown on the graph, and the SEM is indicated by error bars. *P < 0.05, Unpaired Student's *t*-test.

Figure 4. InvL is a critical virulence factor for long-term respiratory infection, but dispensable in the acute infection model. C3H/HeJ (*tlr4* mutant) mice were infected with 10^5 G636, G636 Δ*invL*, or G636 *invL*$^+$. Groups of mice were then sacrificed at 1 dpi (A), 7 dpi (B), 14 dpi (C), and 21 dpi (D), and CFU in the lungs were quantified. Shown are the results from 3 independent experiments. For the acute infection model, groups of C57BL/6 mice were infected with 10^9 G636, G636 Δ*invL*, or G636 *invL*$^+$. 24 hpi, mice were sacrificed, and CFU in the lungs (A), spleen (B), and kidneys (C) were enumerated. Each data point represents an individual mouse, the horizontal line represents the mean, and the SEM is indicated by error bars. Shown are the results of 2-3 independent experiments. *P < 0.05; Kruskal-Wallis *H* test with Dunn's test for multiple comparisons; ns = not significant.

Figure 5. The chronic respiratory infection model can be used to study outcomes of antibiotic treatment. Groups of C3H/HeJ (*tlr4* mutant) mice were infected with 10^5 G636 (A, C) or 10^5 G654 (B, D) and sacrificed at 1, 3, and 5 dpi (long-term). Additionally, groups of C57Bl/6 mice were infected with 10^9 G636 (A, C) or 10^9 G654 (B, D) and sacrificed at 24 hpi (acute). Mice in both infection models were treated intraperitoneally treated with PBS or 100 mg/kg tigecycline (tig) every 12 h (A, B) or PBS or 500 mg/kg apramycin (apr) every 12 h (C, D) with all treatments beginning 4 hpi. At each timepoint, CFU were quantified in the lungs. Shown are the results from at least two independent experiments, each data point represents an individual mouse, the horizontal line represents the mean, and the SEM is represented by error bars. *P < 0.05; Mann-Whitney *U* test.

Figure 6. Bacterial secondary infection alters the course of chronic *A. baumannii* pneumonia. C3H/HeJ (*tlr4* mutant) mice were intranasally inoculated with 10^5 G636, and groups of mice were sacrificed at 1, 7, and 14 dpi. At 14 days post-*A. baumannii* infection, groups of mice were either not inoculated (untreated), inoculated with PBS (mock-infected), infected with *S. aureus*, or

infected on *K. pneumoniae*. Subsequently, on days 15 and 16 post-*A. baumannii* infection (1 and
2 days post-secondary infection), mice were sacrificed, and *A. baumannii* CFU were quantified in
the lungs (A, D, E), spleen (B), and kidneys (C). In panels A, B, and C, each data point represents
the mean, the SEM is represented by error bars, and the limit of detection is indicated by the
dashed line. In panels D and E, each data point represents an individual mouse, the horizontal
line represents the mean, and the SEM is indicated by error bars. Shown are results from at least
2 independent experiments. *$P < 0.05$; Kruskal-Wallis *H* test with Dunn's test for multiple
comparisons.

Figure S1. Intracellular *A. baumannii* are detectable in BALF at early timepoints in the chronic
respiratory infection model. Groups of C3H/HeJ (*tlr4* mutant) mice were intranasally inoculated
with 10^5 G636, and BALF was collected at 4 hpi (A), 2 dpi (B), and 7 dpi (C) and either treated
with 50 µg/ml colistin or mock-treated. Following, bacterial CFU in the treated (intracellular; IC)
and mock treated (total) BALF, as well as in the remaining lungs following BALF collection, were
enumerated by serial dilution plating. The horizontal line represents the mean, and the SEM is
indicated by error bars. Shown are the results from at least two independent experiments.
C3H/HeJ (*tlr4* mutant) mice were infected with G636 expressing *gfp*, and, at these same
timepoints, BALF was collected, and host cells were isolated and stained with DAPI (blue) and
phalloidin (red). Intracellular bacteria were identified by microscopy at 4 hpi (D) and 2 dpi (E).
Shown are representative images from independent samples from at least two biological
replicates. Scale bar = 5 µm.

Figure S2. The chronic respiratory infection model results in lung pathology during infection.
Groups of C3H/HeJ (*tlr4* mutant) mice were inoculated with 10^5 G636 or mock-inoculated with
PBS, and at 4 hpi (A), 2 dpi (B), 7 dpi (C), 14 dpi (D), and 21 dpi (E), lungs slices were prepared
and H&E stained. Shown are representative images from each timepoint. Lung slice scale bar:
1000 µm; Inset scale bar: 100 µm.

Figure S3. The chronic respiratory infection model does not cause goblet cell hyperplasia or
fibrosis. Groups of C3H/HeJ (*tlr4* mutant) mice were inoculated with 10^5 G636 or mock-inoculated
with PBS, and at 4 hpi, 2 dpi, 7 dpi, 14 dpi, and 21 dpi, lungs slices were prepared, H&E stained,
and scored for goblet cell hyperplasia (A) and fibrosis (B). The mean is shown on the graph, and
the SEM is indicated by error bars. Unpaired Student's *t*-test.

Figure S4. Testing of G636 adhesin mutants in the chronic respiratory infection model reveals a
potential role for InvL in bacterial persistence. Groups of C3H/HeJ (*tlr4* mutant) mice were
intranasally inoculated with 10^5 G636, G636 Δ*bap*, G636 Δ*ata*, G636 Δ*fhaBC*, and G636 Δ*invL*.
1 (A) and 14 (B) dpi, mice were sacrificed, and CFU in the lungs were quantified. Each data point
represents an individual mouse, the horizontal line represents the mean, and the SEM is indicated
by error bars. Shown are results from single experiments for each strain.

Figure S5. The chronic respiratory infection model can be used to study outcomes of antibiotic
treatment. Groups of C3H/HeJ (*tlr4* mutant) mice were infected with 10^5 G636 (A, C) or 10^5 G654
(B, D) and sacrificed at 1, 3, and 5 dpi (long-term). Additionally, groups of C57Bl/6 mice were
infected with 10^9 G636 (A, C) or 10^9 G654 (B, D) and sacrificed at 24 hpi (acute). Mice in both
infection models were treated intraperitoneally treated with PBS or 5 mg/kg colistin (col) every 8
h (A, B) or PBS or 100 mg/kg imipenem (im) every 12 h (C, D) with all treatments beginning 4 hpi.
At each timepoint, CFU were quantified in the lungs. Shown are the results from at least two

independent experiments, each data point represents an individual mouse, the horizontal line
represents the mean, and the SEM is represented by error bars. *P < 0.05; Mann-Whitney U test.

Figure S6. *S. aureus* secondary infection sometimes causes *A. baumannii* dissemination to the
spleen and kidneys in the chronic respiratory infection model. C3H/HeJ (*tlr4* mutant) mice were
intranasally inoculated with 10^5 G636. At 14 days post-*A. baumannii* infection, groups of mice
were either not inoculated (untreated), inoculated with PBS (mock-infected), infected with *S.*
aureus, or infected with *K. pneumoniae*. Subsequently, on days 15 (A and B) and 16 (C and D)
post-*A. baumannii* infection (1 and 2 days post-secondary infection), groups of mice were
sacrificed, and *A. baumannii* CFU were quantified in the spleen (A and C), and kidneys (C and
D). Each data point represents an individual mouse, the horizontal line represents the mean, and
the SEM is indicated by error bars. Shown are results from at least 2 independent experiments.
Significant differences were not detected; Kruskal-Wallis H test with Dunn's test for multiple
comparisons.

Figure S7. Ongoing *A. baumannii* pneumonia alters bacterial numbers following secondary
infection with *S. aureus* and *K. pneumoniae*. C3H/HeJ (*tlr4* mutant) mice were either intranasally
inoculated with 10^5 G636 14 days prior to infection with *S. aureus* or *K. pneumoniae* (+*Ab*) or not
infected prior to *S. aureus* or *K. pneumoniae* infection (-*Ab*). At 14 days post-*A. baumannii*
infection, groups of mice were infected with *S. aureus* or *K. pneumoniae*. 1 and 2 dpi with *S.*
aureus (A, B, and C) or *K. pneumoniae* (D, E, and F), mice were sacrificed, and these bacteria
were quantified in the lung (A and D), spleen (B and E), and kidneys (C and F). Each data point
represents an individual mouse, the horizontal line represents the mean, and the SEM is indicated
by error bars. Shown are results from at least 2 independent experiments. *P < 0.05; Mann-
Whitney U test.

Figure 1:

    Figure 2:

687   Figure 3:

688
689

690   Figure 4:

693   Figure 5:

696   Figure 6:

699 Figure S1:

700
701

702   Figure S2:

705   Figure S3

708   Figure S4:

711   Figure S5

714 Figure S6

References

720   **References**

721   1.   Vijayakumar S, Biswas I, Veeraraghavan B. 2019. Accurate identification of clinically
722   important Acinetobacter spp.: an update. Future Sci OA 5.
723
724   2.   Cerqueira GM, Peleg AY. 2011. Insights into Acinetobacter baumannii pathogenicity.
725   IUBMB Life 63:1055–1060.
726
727   3.   Sarshar M, Behzadi P, Scribano D, Palamara AT, Ambrosi C. 2021. Acinetobacter
728   baumannii: An Ancient Commensal with Weapons of a Pathogen. Pathogens 10.
729
730   4.   Di Venanzio G, Flores-Mireles AL, Calix JJ, Haurat MF, Scott NE, Palmer LD, Potter RF,
731   Hibbing ME, Friedman L, Wang B, Dantas G, Skaar EP, Hultgren SJ, Feldman MF. 2019.
732   Urinary tract colonization is enhanced by a plasmid that regulates uropathogenic
733   Acinetobacter baumannii chromosomal genes. Nat Commun 10.
734
5.   Murray CJ, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Gray A, Han C,
Bisignano C, Rao P, Wool E, Johnson SC, Browne AJ, Chipeta MG, Fell F, Hackett S,
Haines-Woodhouse G, Kashef Hamadani BH, Kumaran EAP, McManigal B, Agarwal R,
Akech S, Albertson S, Amuasi J, Andrews J, Aravkin A, Ashley E, Bailey F, Baker S,
Basnyat B, Bekker A, Bender R, Bethou A, Bielicki J, Boonkasidecha S, Bukosia J,
Carvalheiro C, Castañeda-Orjuela C, Chansamouth V, Chaurasia S, Chiurchiù S,
Chowdhury F, Cook AJ, Cooper B, Cressey TR, Criollo-Mora E, Cunningham M, Darboe
S, Day NPJ, De Luca M, Dokova K, Dramowski A, Dunachie SJ, Eckmanns T, Eibach D,
Emami A, Feasey N, Fisher-Pearson N, Forrest K, Garrett D, Gastmeier P, Giref AZ,
Greer RC, Gupta V, Haller S, Haselbeck A, Hay SI, Holm M, Hopkins S, Iregbu KC,
Jacobs J, Jarovsky D, Javanmardi F, Khorana M, Kissoon N, Kobeissi E, Kostyanev T,
Krapp F, Krumkamp R, Kumar A, Kyu HH, Lim C, Limmathurotsakul D, Loftus MJ, Lunn
M, Ma J, Mturi N, Munera-Huertas T, Musicha P, Mussi-Pinhata MM, Nakamura T,
Nanavati R, Nangia S, Newton P, Ngoun C, Novotney A, Nwakanma D, Obiero CW,
Olivas-Martinez A, Olliaro P, Ooko E, Ortiz-Brizuela E, Peleg AY, Perrone C, Plakkal N,
Ponce-de-Leon A, Raad M, Ramdin T, Riddell A, Roberts T, Robotham JV, Roca A, Rudd
KE, Russell N, Schnall J, Scott JAG, Shivamallappa M, Sifuentes-Osornio J, Steenkeste
N, Stewardson AJ, Stoeva T, Tasak N, Thaiprakong A, Thwaites G, Turner C, Turner P,
van Doorn HR, Velaphi S, Vongpradith A, Vu H, Walsh T, Waner S, Wangrangsimakul T,
Wozniak T, Zheng P, Sartorius B, Lopez AD, Stergachis A, Moore C, Dolecek C, Naghavi
M. 2022. Global burden of bacterial antimicrobial resistance in 2019: a systematic
analysis. The Lancet 399:629–655.
757
758   6.   Dexter C, Murray GL, Paulsen IT, Peleg AY. 2015. Community-acquired Acinetobacter
759   baumannii: clinical characteristics, epidemiology and pathogenesis. Expert Rev Anti Infect
760   Ther 13:567–573.
761
762   7.   Cisneros JM, Rodriguez-Baño J. 2002. Nosocomial bacteremia due to Acinetobacter
763   baumannii: epidemiology, clinical features and treatment. Clin Microbiol Infect 8:687–693.
764
765   8.   Ibrahim S, Al-Saryi N, Al-Kadmy IMS, Aziz SN. 2021. Multidrug-resistant Acinetobacter
766   baumannii as an emerging concern in hospitals. Mol Biol Rep 48:6987–6998.
767

768   9.   Karakonstantis S, Kritsotakis EI. 2021. Systematic review and meta-analysis of the
769   proportion and associated mortality of polymicrobial (vs monomicrobial) pulmonary and
770   bloodstream infections by Acinetobacter baumannii complex. Infection 49:1149–1161.
771
772   10.   Xiao D, Wang L, Zhang D, Xiang D, Liu Q, Xing X. 2017. Prognosis of patients with
773   Acinetobacter baumannii infection in the intensive care unit: A retrospective analysis. Exp
774   Ther Med 13:1630–1633.
775
776   11.   Sengstock DM, Thyagarajan R, Apalara J, Mira A, Chopra T, Kaye KS. 2010. Multidrug-
777   resistant Acinetobacter baumannii: an emerging pathogen among older adults in
778   community hospitals and nursing homes. Clin Infect Dis 50:1611–1616.
779
780   12.   Inchai J, Pothirat C, Bumroongkit C, Limsukon A, Khositsakulchai W, Liwsrisakun C.
781   2015. Prognostic factors associated with mortality of drug-resistant Acinetobacter
782   baumannii ventilator-associated pneumonia. J Intensive Care 3.
783
784   13.   Giammanco A, Calà C, Fasciana T, Dowzicky MJ. 2017. Global Assessment of the
785   Activity of Tigecycline against Multidrug-Resistant Gram-Negative Pathogens between
786   2004 and 2014 as Part of the Tigecycline Evaluation and Surveillance Trial. mSphere 2.
787
788   14.   Tacconelli E, Carrara E, Savoldi A, Kattula D, Burkert F. 2018. Discovery, research, and
789   development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and
790   tuberculosis. Lancet Infect Dis 18:318–327.
791
792   15.   Kollef MH, Torres A, Shorr AF, Martin-Loeches I, Micek ST. 2021. Nosocomial Infection.
793   Crit Care Med 49:169–187.
794
795   16.   Jones RN. 2010. Microbial etiologies of hospital-acquired bacterial pneumonia and
796   ventilator-associated bacterial pneumonia. Clin Infect Dis 51 Suppl 1.
797
798   17.   Sader HS, Castanheira M, Mendes RE, Flamm RK. 2018. Frequency and antimicrobial
799   susceptibility of Gram-negative bacteria isolated from patients with pneumonia
800   hospitalized in ICUs of US medical centres (2015-17). J Antimicrob Chemother 73:3053–
801   3059.
802
803   18.   Weber BS, Harding CM, Feldman MF. 2015. Pathogenic Acinetobacter: from the Cell
804   Surface to Infinity and Beyond. J Bacteriol 198:880–887.
805
806   19.   Kebaier C, Chamberland RR, Allen IC, Gao X, Broglie PM, Hall JD, Jania C, Doerschuk
807   CM, Tilley SL, Duncan JA. 2012. Staphylococcus aureus α-Hemolysin Mediates Virulence
808   in a Murine Model of Severe Pneumonia Through Activation of the NLRP3
809   Inflammasome. J Infect Dis 205:807–817.
810
811   20.   Hoffmann N, Rasmussen TB, Jensen PØ, Stub C, Hentzer M, Molin S, Ciofu O, Givskov
812   M, Johansen HK, Høiby N. 2005. Novel Mouse Model of Chronic Pseudomonas
813   aeruginosa Lung Infection Mimicking Cystic Fibrosis. Infect Immun 73:2504.
814
815   21.   Palmer LD, Green ER, Sheldon JR, Skaar EP. 2019. Assessing Acinetobacter baumannii
816   Virulence and Persistence in a Murine Model of Lung Infection. Methods Mol Biol
817   1946:289–305.
818

819 22. Pogue JM, Zhou Y, Kanakamedala H, Cai B. 2022. Burden of illness in carbapenem-
820   resistant Acinetobacter baumannii infections in US hospitals between 2014 and 2019.
821   BMC Infect Dis 22.
822

823 23. Alotaibi T, Abuhaimed A, Alshahrani M, Albdelhady A, Almubarak Y, Almasari O. 2021.
824   Prevalence of multidrug-resistant Acinetobacter baumannii in a critical care setting: A
825   tertiary teaching hospital experience. SAGE Open Med 9.
826

827 24. Koomanachai P, Kim A, Nicolau DP. 2009. Pharmacodynamic evaluation of tigecycline
828   against Acinetobacter baumannii in a murine pneumonia model. Journal of Antimicrobial
829   Chemotherapy 63:982–987.
830

831 25. Koomanachai P, Kim A, Nicolau DP. 2009. Pharmacodynamic evaluation of tigecycline
832   against Acinetobacter baumannii in a murine pneumonia model. Journal of Antimicrobial
833   Chemotherapy 63:982–987.
834

835 26. Crandon JL, Kim A, Nicolau DP. 2009. Comparison of tigecycline penetration into the
836   epithelial lining fluid of infected and uninfected murine lungs. Journal of Antimicrobial
837   Chemotherapy 64:837–839.
838

839 27. Braunstein A, Papo N, Shai Y. 2004. In Vitro Activity and Potency of an Intravenously
840   Injected Antimicrobial Peptide and Its dl Amino Acid Analog in Mice Infected with Bacteria.
841   Antimicrob Agents Chemother 48:3127.
842

843 28. Joly-Guillou ML, Wolff M, Pocidalo JJ, Walker F, Carbon C. 1997. Use of a new mouse
844   model of Acinetobacter baumannii pneumonia to evaluate the postantibiotic effect of
845   imipenem. Antimicrob Agents Chemother 41:345.
846

847 29. Song JY, Cheong HJ, Lee J, Sung AK, Kim WJ. 2009. Efficacy of monotherapy and
848   combined antibiotic therapy for carbapenem-resistant Acinetobacter baumannii
849   pneumonia in an immunosuppressed mouse model. Int J Antimicrob Agents 33:33–39.
850

851 30. Manepalli S, Gandhi JA, Ekhar V V., Asplund MB, Coelho C, Martinez LR. 2013.
852   Characterization of a cyclophosphamide-induced murine model of immunosuppression to
853   study Acinetobacter baumannii pathogenesis. J Med Microbiol 62:1747–1754.
854

855 31. Distel JS, Di Venanzio G, Mackel JJ, Rosen DA, Feldman MF. 2023. Replicative
856   Acinetobacter baumannii strains interfere with phagosomal maturation by modulating the
857   vacuolar pH. PLoS Pathog 19.
858

859 32. Sycz G, Venanzio G Di, Distel JS, Sartorio MG, Le NH, Scott NE, Beatty WL, Feldman
860   MF. 2021. Modern Acinetobacter baumannii clinical isolates replicate inside spacious
861   vacuoles and egress from macrophages. PLoS Pathog 17.
862

863 33. Valcek A, Philippe C, Whiteway C, Robino E, Nesporova K, Bové M, Coenye T, De Pooter
864   T, De Coster W, Strazisar M, Van der Henst C. 2023. Phenotypic Characterization and
865   Heterogeneity among Modern Clinical Isolates of Acinetobacter baumannii. Microbiol
866   Spectr 11.
867

868 34. Rubio T, Gagné S, Debruyne C, Dias C, Cluzel C, Mongellaz D, Rousselle P, Göttig S,
869   Seifert H, Higgins PG, Salcedo SP. 2022. Incidence of an Intracellular Multiplication Niche
870   among Acinetobacter baumannii Clinical Isolates. mSystems 7.
871
872 35. Sato Y, Unno Y, Miyazaki C, Ubagai T, Ono Y. 2019. Multidrug-resistant Acinetobacter
873   baumannii resists reactive oxygen species and survives in macrophages. Sci Rep 9.
874
875 36. Cochet F, Peri F. 2017. The Role of Carbohydrates in the Lipopolysaccharide (LPS)/Toll-
876   Like Receptor 4 (TLR4) Signalling. Int J Mol Sci 18.
877
878 37. Akira S, Yamamoto M. 2010. Lipid A receptor TLR4-mediated signaling pathways. Adv
879   Exp Med Biol 667:59–68.
880
881 38. Homma JY, Matsuura M, Kanegasaki S, Kawakubo Y, Kojima Y, Shibukawa N,
882   Kumazawa Y, Yamamoto A, Tanamoto K ichi, Yasuda T, Imoto M, Yoshimura H,
883   Kusumoto S, Shiba T. 1985. Structural requirements of lipid A responsible for the
884   functions: a study with chemically synthesized lipid A and its analogues. J Biochem
885   98:395–406.
886
887 39. Kawasaki T, Kawai T. 2014. Toll-like receptor signaling pathways. Front Immunol
888   5:112681.
889
890 40. Lin L, Tan B, Pantapalangkoor P, Ho T, Baquir B, Tomaras A, Montgomery JI, Reilly U,
891   Barbacci EG, Hujer K, Bonomo RA, Fernandez L, Hancock REW, Adams MD, French
892   SW, Buslon VS, Spellberg B. 2012. Inhibition of LpxC protects mice from resistant
893   Acinetobacter baumannii by modulating inflammation and enhancing phagocytosis. mBio
894   3.
895
896 41. Knapp S, Wieland CW, Florquin S, Pantophlet R, Dijkshoorn L, Tshimbalanga N, Akira S,
897   Van Der Poll T. 2006. Differential roles of CD14 and toll-like receptors 4 and 2 in murine
898   Acinetobacter pneumonia. Am J Respir Crit Care Med 173:122–129.
899
900 42. Hazen JE, Di Venanzio G, Hultgren SJ, Feldman MF. 2023. Catheterization triggers
901   resurgent infection seeded by host Acinetobacter baumannii reservoirs. Sci Transl Med
902   15:eabn8134.
903
904 43. Dijkshoorn L, Aucken HM, Gerner-Smidt P, Kaufmann ME, Ursing J, Pitt TL. 1993.
905   Correlation of typing methods for Acinetobacter isolates from hospital outbreaks. J Clin
906   Microbiol 31:702–705.
907
908 44. Luo G, Lin L, Ibrahim AS, Baquir B, Pantapalangkoor P, Bonomo RA, Doi Y, Adams MD,
909   Russo TA, Spellberg B. 2012. Active and passive immunization protects against lethal,
910   extreme drug resistant-Acinetobacter baumannii infection. PLoS One 7.
911
912 45. Behairy MY, Abdelrahman AA, Toraih EA, Ibrahim EEDA, Azab MM, Sayed AA, Hashem
913   HR. 2022. Investigation of TLR2 and TLR4 Polymorphisms and Sepsis Susceptibility:
914   Computational and Experimental Approaches. Int J Mol Sci 23.
915
916 46. Huang WH, Nie LH, Zhang LJ, Jing LP, Dong F, Wang M, Zhang N, Liu Y, Zhang BH,
917   Chen C, Lin HS, Wei XC, Yang G, Jing CX. 2015. Association of TLR2 and TLR4 non-
918   missense single nucleotide polymorphisms with type 2 diabetes risk in a Southern

919　　　　Chinese population: A case-control study. Genetics and Molecular Research 14:8694–
920　　　　8705.
921
922　47.　Chatzi M, Papanikolaou J, Makris D, Papathanasiou I, Tsezou A, Karvouniaris M,
923　　　　Zakynthinos E. 2018. Toll-like receptor 2, 4 and 9 polymorphisms and their association
924　　　　with ICU-acquired infections in Central Greece. J Crit Care 47:1–8.
925
926　48.　Wang S-H, Teng C-K, Chan M-C, Yang K-Y, Sheu C-C, Liang S-J, Huang W-H, Feng J-Y,
927　　　　Chen C-M, Weng Z-X, Peng C-K. 2024. The impact and risk factors for developing
928　　　　pneumogenic bacteremia in carbapenem-resistant Acinetobacter baumannii nosocomial
929　　　　pneumonia in the intensive care unit: A multicenter retrospective study. Int J Infect Dis
930　　　　146:107128.
931
932　49.　Van Faassen H, KuoLee R, Harris G, Zhao X, Conlan JW, Chen W. 2007. Neutrophils
933　　　　play an important role in host resistance to respiratory infection with Acinetobacter
934　　　　baumannii in mice. Infect Immun 75:5597–5608.
935
936　50.　Liu Z, Xu W. 2022. Neutrophil and Macrophage Response in Acinetobacter Baumannii
937　　　　Infection and Their Relationship to Lung Injury. Front Cell Infect Microbiol 12.
938
939　51.　Grguric-Smith LM, Lee HH, Gandhi JA, Brennan MB, DeLeon-Rodriguez CM, Coelho C,
940　　　　Han G, Martinez LR. 2015. Neutropenia exacerbates infection by Acinetobacter
941　　　　baumannii clinical isolates in a murine wound model. Front Microbiol 6.
942
943　52.　Bruhn KW, Pantapalangkoor P, Nielsen T, Tan B, Junus J, Hujer KM, Wright MS, Bonomo
944　　　　RA, Adams MD, Chen W, Spellberg B. 2015. Host fate is rapidly determined by innate
945　　　　effector-microbial interactions during Acinetobacter baumannii bacteremia. J Infect Dis
946　　　　211:1296–1305.
947
948　53.　Tsuchiya T, Nakao N, Yamamoto S, Hirai Y, Miyamoto K, Tsujibo H. 2012. NK1.1(+) cells
949　　　　regulate neutrophil migration in mice with Acinetobacter baumannii pneumonia. Microbiol
950　　　　Immunol 56:107–116.
951
952　54.　Qiu H, KuoLee R, Harris G, Chen W. 2009. High susceptibility to respiratory
953　　　　Acinetobacter baumannii infection in A/J mice is associated with a delay in early
954　　　　pulmonary recruitment of neutrophils. Microbes Infect 11:946–955.
955
956　55.　Breslow JM, Meissler J, Hartzell RR, Spence PB, Truant A, Gaughan J, Eisenstein TK.
957　　　　2011. Innate immune responses to systemic acinetobacter baumannii infection in mice:
958　　　　Neutrophils, but not interleukin-17, mediate host resistance. Infect Immun 79:3317–3327.
959
960　56.　Castro ÍA, Yang Y, Gnazzo V, Kim D-H, Van Dyken SJ, López CB. 2023. Murine
961　　　　Parainfluenza Virus Persists in Lung Innate Immune Cells Sustaining Chronic Lung
962　　　　Pathology. bioRxiv https://doi.org/10.1101/2023.11.07.566103.
963
964　57.　Brossard KA, Campagnari AA. 2012. The Acinetobacter baumannii biofilm-associated
965　　　　protein plays a role in adherence to human epithelial cells. Infect Immun 80:228–233.
966
967　58.　Pérez A, Merino M, Rumbo-Feal S, Álvarez-Fraga L, Vallejo JA, Beceiro A, Ohneck EJ,
968　　　　Mateos J, Fernández-Puente P, Actis LA, Poza M, Bou G. 2017. The FhaB/FhaC two-

partner secretion system is involved in adhesion of Acinetobacter baumannii AbH12O-A2 strain. Virulence 8:959–974.

59. Astaneh SDA, Rasooli I, Gargari SLM. 2017. Filamentous hemagglutinin adhesin FhaB limits A.baumannii biofilm formation. Front Biosci (Elite Ed) 9:266–275.

60. Darvish Alipour Astaneh S, Rasooli I, Mousavi Gargari SL. 2014. The role of filamentous hemagglutinin adhesin in adherence and biofilm formation in Acinetobacter baumannii ATCC19606(T). Microb Pathog 74:42–49.

61. Bentancor L V., Camacho-Peiro A, Bozkurt-Guzel C, Pier GB, Maira-Litrán T. 2012. Identification of Ata, a multifunctional trimeric autotransporter of Acinetobacter baumannii. J Bacteriol 194:3950–3960.

62. Weidensdorfer M, Ishikawa M, Hori K, Linke D, Djahanschiri B, Iruegas R, Ebersberger I, Riedel-Christ S, Enders G, Leukert L, Kraiczy P, Rothweiler F, Cinatl J, Berger J, Hipp K, Kempf VAJ, Göttig S. 2019. The Acinetobacter trimeric autotransporter adhesin Ata controls key virulence traits of Acinetobacter baumannii. Virulence 10:68–81.

63. Hatefi Oskuei R, Darvish Alipour Astaneh S, Rasooli I. 2021. A conserved region of Acinetobacter trimeric autotransporter adhesion, Ata, provokes suppression of Acinetobacter baumannii virulence. Arch Microbiol 203:3483–3493.

64. Tram G, Poole J, Adams FG, Jennings MP, Eijkelkamp BA, Atack JM. 2021. The Acinetobacter baumannii Autotransporter Adhesin Ata Recognizes Host Glycans as High-Affinity Receptors. ACS Infect Dis 7:2352–2361.

65. Ishikawa M, Nakatani H, Hori K. 2012. AtaA, a new member of the trimeric autotransporter adhesins from Acinetobacter sp. Tol 5 mediating high adhesiveness to various abiotic surfaces. PLoS One 7.

66. Jackson-Litteken CD, Venanzio G Di, Le NH, Scott NE, Djahanschiri B, Distel JS, Pardue EJ, Ebersberger I, Feldman MF. 2022. InvL, an Invasin-Like Adhesin, Is a Type II Secretion System Substrate Required for Acinetobacter baumannii Uropathogenesis. mBio 13.

67. Yesil C, Yalcin AN, Ogunc D, Ongut G, Ozhak B, Colak D, Er H, Sarltas ZE. 2022. Use of colistin with rifampicin, trimethoprim-sulfamethoxazole and teicoplanin in Acinetobacter mouse infection model. Future Microbiol 17:665–671.

68. İzci F, Ture Z, Dinc G, Yay AH, Eren EE, Bolat D, Gönen ZB, Ünüvar GK, Yıldız O, Aygen B. 2023. The efficacy of mesenchymal stem cell treatment and colistin-fosfomycin combination on colistin-resistant Acinetobacter baumannii sepsis model. European Journal of Clinical Microbiology and Infectious Diseases 42:1365–1372.

69. Dinc G, Demiraslan H, Elmali F, Ahmed SS, Metan G, Alp E, Doganay M. 2014. Efficacy of Sulbactam and Its Combination with Imipenem, Colistin and Tigecycline in an Experimental Model of Carbapenem-Resistant Acinetobacter baumannii Sepsis. Chemotherapy 59:325–329.

1019   70.   Nicasio AM, Crandon JL, Nicolau DP. 2009. In Vivo Pharmacodynamic Profile of
1020   Tigecycline against Phenotypically Diverse Escherichia coli and Klebsiella pneumoniae
1021   Isolates. Antimicrob Agents Chemother 53:2756.
1022

1023   71.   Joly-Guillou ML, Wolff M, Farinotti R, Bryskier A, Carbon C. 2000. In vivo activity of
1024   levofloxacin alone or in combination with imipenem or amikacin in a mouse model of
1025   Acinetobacter baumannii pneumonia. Journal of Antimicrobial Chemotherapy 46:827–
1026   830.
1027

1028   72.   Queenan AM, Davies TA, He W, Lynch AS. 2013. Assessment of the combination of
1029   doripenem plus a fluoroquinolone against non-susceptible Acinetobacter baumannii
1030   isolates from nosocomial pneumonia patients. Journal of Chemotherapy 25:141–147.
1031

1032   73.   Al-Madboly LA. 2022. A Novel Triple Combination To Combat Serious Infections with
1033   Carbapenem-Resistant Acinetobacter baumannii in a Mouse Pneumonia Model. Microbiol
1034   Spectr 10.
1035

1036   74.   Kang AD, Smith KP, Berg AH, Truelson KA, Eliopoulos GM, McCoy C, Kirby JE. 2018.
1037   Efficacy of apramycin against multidrug-resistant Acinetobacter baumannii in the murine
1038   neutropenic thigh model. Antimicrob Agents Chemother 62.
1039

1040   75.   Becker K, Aranzana-Climent V, Cao S, Nilsson A, Shariatgorji R, Haldimann K, Platzack
1041   B, Hughes D, Andrén PE, Böttger EC, Friberg LE, Hobbie SN. 2021. Efficacy of EBL-
1042   1003 (apramycin) against Acinetobacter baumannii lung infections in mice. Clinical
1043   Microbiology and Infection 27:1315–1321.
1044

1045   76.   Zhao C, Chirkova A, Rosenborg S, Palma Villar R, Lindberg J, Hobbie SN, Friberg LE.
1046   2022. Population pharmacokinetics of apramycin from first-in-human plasma and urine
1047   data to support prediction of efficacious dose. Journal of Antimicrobial Chemotherapy
1048   77:2718.
1049

1050   77.   Ambrosi C, Scribano D, Sarshar M, Zagaglia C, Singer BB, Palamara AT. 2020.
1051   Acinetobacter baumannii Targets Human Carcinoembryonic Antigen-Related Cell
1052   Adhesion Molecules (CEACAMs) for Invasion of Pneumocytes. mSystems 5.
1053

1054   78.   An Z, Huang X, Zheng C, Ding W. 2019. Acinetobacter baumannii outer membrane
1055   protein A induces HeLa cell autophagy via MAPK/JNK signaling pathway. Int J Med
1056   Microbiol 309:97–107.
1057

1058   79.   Wang Y, Zhang K, Shi X, Wang C, Wang F, Fan J, Shen F, Xu J, Bao W, Liu M, Yu L.
1059   2016. Critical role of bacterial isochorismatase in the autophagic process induced by
1060   Acinetobacter baumannii in mammalian cells. FASEB J 30:3563–3577.
1061

1062   80.   Bist P, Dikshit N, Koh TH, Mortellaro A, Tan TT, Sukumaran B. 2014. The Nod1, Nod2,
1063   and Rip2 axis contributes to host immune defense against intracellular Acinetobacter
1064   baumannii infection. Infect Immun 82:1112–1122.
1065

1066   81.   Parra-Millán R, Guerrero-Gómez D, Ayerbe-Algaba R, Pachón-Ibáñez ME, Miranda-
1067   Vizuete A, Pachón J, Smani Y. 2018. Intracellular Trafficking and Persistence of
1068   Acinetobacter baumannii Requires Transcription Factor EB. mSphere 3.
1069

82. Maure A, Robino E, Van der Henst C. 2023. The intracellular life of Acinetobacter baumannii. Trends Microbiol 31:1238–1250.

83. Asensio NC, Rendón JM, Burgas MT. 2021. Time-Resolved Transcriptional Profiling of Epithelial Cells Infected by Intracellular Acinetobacter baumannii. Microorganisms 9:1–14.

84. Rumbo C, Tomás M, Moreira EF, Soares NC, Carvajal M, Santillana E, Beceiro A, Romero A, Bou G. 2014. The Acinetobacter baumannii Omp33-36 porin is a virulence factor that induces apoptosis and modulates autophagy in human cells. Infect Immun 82:4666–4680.

85. An Z, Ding W. 2021. Acinetobacter baumannii up-regulates LncRNA-GAS5 and promotes the degradation of STX17 by blocking the activation of YY1. Virulence 12:1965–1979.

86. Kho ZY, Azad MAK, Han ML, Zhu Y, Huang C, Schittenhelm RB, Naderer T, Velkov T, Selkrig J, Zhou Q, Li J. 2022. Correlative proteomics identify the key roles of stress tolerance strategies in Acinetobacter baumannii in response to polymyxin and human macrophages. PLoS Pathog 18.

87. Smani Y, Docobo-Pе́rez F, Loṕez-Rojas R, Domińguez-Herrera J, Ibáñez-Martínez J, Pachón J. 2012. Platelet-activating factor receptor initiates contact of Acinetobacter baumannii expressing phosphorylcholine with host cells. J Biol Chem 287:26901–26910.

88. Jacobs AC, Hood I, Boyd KL, Olson PD, Morrison JM, Carson S, Sayood K, Iwen PC, Skaar EP, Dunman PM. 2010. Inactivation of phospholipase D diminishes Acinetobacter baumannii pathogenesis. Infect Immun 78:1952–1962.

89. Zhao D, Li Y, Peng C, Lin J, Yu F, Zhao Y, Zhang X, Zhao D. 2021. Outer membrane protein a in Acinetobacter baumannii induces autophagy through mTOR signalling pathways in the lung of SD rats. Biomedicine & Pharmacotherapy 135:111034.

90. Stahl J, Bergmann H, Göttig S, Ebersberger I, Averhoff B. 2015. Acinetobacter baumannii Virulence Is Mediated by the Concerted Action of Three Phospholipases D. PLoS One 10.

91. Tomlin H, Piccinini AM. 2018. A complex interplay between the extracellular matrix and the innate immune response to microbial pathogens. Immunology 155:186–201.

92. Wight TN, Frevert CW, Debley JS, Reeves SR, Parks WC, Ziegler SF. 2017. Interplay of Extracellular Matrix and Leukocytes in Lung Inflammation. Cell Immunol 312:1.

93. Onishi Y, Kawamura T, Higashino T, Mimura R, Tsukamoto H, Sasaki S. 2021. Clinical features of acute fibrinous and organizing pneumonia: An early histologic pattern of various acute inflammatory lung diseases. PLoS One 16:e0249300.

94. Chapman HA. 2012. Epithelial Responses to Lung Injury. https://doi.org/101513/pats201112-053AW 9:89–95.

95. Åhrman E, Hallgren O, Malmström L, Hedström U, Malmström A, Bjermer L, Zhou XH, Westergren-Thorsson G, Malmström J. 2018. Quantitative proteomic characterization of

the lung extracellular matrix in chronic obstructive pulmonary disease and idiopathic pulmonary fibrosis. J Proteomics 189:23–33.

96. Booth AJ, Hadley R, Cornett AM, Dreffs AA, Matthes SA, Tsui JL, Weiss K, Horowitz JC, Fiore VF, Barker TH, Moore BB, Martinez FJ, Niklason LE, White ES. 2012. Acellular normal and fibrotic human lung matrices as a culture system for in vitro investigation. Am J Respir Crit Care Med 186:866–876.

97. Annoni R, Lancąs T, Tanigawa RY, Matsushita MDM, Fernezlian SDM, Bruno A, Da Silva LFF, Roughley PJ, Battaglia S, Dolhnikoff M, Hiemstra PS, Sterk PJ, Rabe KF, Mauad T. 2012. Extracellular matrix composition in COPD. Eur Respir J 40:1362–1373.

98. White ES. 2015. Lung extracellular matrix and fibroblast function. Ann Am Thorac Soc 12:S30–S33.

99. Chang MY, Tanino Y, Vidova V, Kinsella MG, Chan CK, Johnson PY, Wight TN, Frevert CW. 2014. A rapid increase in macrophage-derived versican and hyaluronan in infectious lung disease. Matrix Biol 34:1–12.

100. Snyder JM, Washington IM, Birkland T, Chang MY, Frevert CW. 2015. Correlation of Versican Expression, Accumulation, and Degradation during Embryonic Development by Quantitative Immunohistochemistry. J Histochem Cytochem 63:952–967.

101. Niu H, Gu J, Zhang Y. 2024. Bacterial persisters: molecular mechanisms and therapeutic development. Signal Transduction and Targeted Therapy 2024 9:1 9:1–32.

102. Wood TK, Knabel SJ, Kwan BW. 2013. Bacterial Persister Cell Formation and Dormancy. Appl Environ Microbiol 79:7116.

103. Kunnath AP, Suodha Suoodh M, Chellappan DK, Chellian J, Palaniveloo K. 2024. Bacterial Persister Cells and Development of Antibiotic Resistance in Chronic Infections: An Update. Br J Biomed Sci 81.

104. Li G, Shen W, Gong Y, Li M, Rao X, Liu Q, Yu Y, Zhou J, Zhu K, Yuan M, Shang W, Yang Y, Lu S, Wang J, Zhao Y. 2022. Essential Fitness Repertoire of Staphylococcus aureus during Co-infection with Acinetobacter baumannii In Vivo. mSystems 7.

105. Timme S, Wendler S, Klassert TE, Saraiva JP, da Rocha UN, Wittchen M, Schramm S, Ehricht R, Monecke S, Edel B, Rödel J, Löffler B, Ramirez MS, Slevogt H, Figge MT, Tuchscherr L. 2024. Competitive inhibition and mutualistic growth in co-infections: deciphering Staphylococcus aureus-Acinetobacter baumannii interaction dynamics. ISME communications 4.

106. Semenec L, Cain AK, Dawson CJ, Liu Q, Dinh H, Lott H, Penesyan A, Maharjan R, Short FL, Hassan KA, Paulsen IT. 2023. Cross-protection and cross-feeding between Klebsiella pneumoniae and Acinetobacter baumannii promotes their co-existence. Nat Commun 14.

107. Krasauskas R, Skerniškytė J, Armalytė J, Sužiedėlienė E. 2019. The role of Acinetobacter baumannii response regulator BfmR in pellicle formation and competitiveness via contact-dependent inhibition system. BMC Microbiol 19.

1171 108. Repizo GD, Gagné S, Foucault-Grunenwald ML, Borges V, Charpentier X, Limansky AS,
1172   Gomes JP, Viale AM, Salcedo SP. 2015. Differential Role of the T6SS in Acinetobacter
1173   baumannii Virulence. PLoS One 10.
1174
1175 109. Weber BS, Ly PM, Irwin JN, Pukatzki S, Feldman MF. 2015. A multidrug resistance
1176   plasmid contains the molecular switch for type VI secretion in Acinetobacter baumannii.
1177   Proc Natl Acad Sci U S A 112:9442–9447.
1178
1179 110. Storey D, McNally A, Åstrand M, Santos JPG, Rodriguez-Escudero I, Elmore B, Palacios
1180   L, Marshall H, Hobley L, Molina M, Cid VJ, Salminen TA, Bengoechea JA. 2020.
1181   Klebsiella pneumoniae type VI secretion system-mediated microbial competition is
1182   PhoPQ controlled and reactive oxygen species dependent. PLoS Pathog 16:e1007969.
1183
1184 111. Tsai CJY, Loh JMS, Proft T. 2016. Galleria mellonella infection models for the study of
1185   bacterial diseases and for antimicrobial drug testing. Virulence 7:214–229.
1186
1187 112. McGuffey JC, Jackson-Litteken CD, Venanzio G Di, Zimmer AA, Lewis JM, Distel JS, Kim
1188   KQ, Zaher HS, Alfonzo J, Scott NE, Feldman MF. 2023. The tRNA methyltransferase
1189   TrmB is critical for Acinetobacter baumannii stress responses and pulmonary infection.
1190   mBio 14.
1191
1192 113. Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image
1193   analysis. Nat Methods 9:671–675.
1194
1195 114. Hoang TT, Karkhoff-Schweizer RR, Kutchma AJ, Schweizer HP. 1998. A broad-host-
1196   range Flp-FRT recombination system for site-specific excision of chromosomally-located
1197   DNA sequences: application for isolation of unmarked Pseudomonas aeruginosa
1198   mutants. Gene 212:77–86.
1199
1200 115. Pontes MH, Groisman EA. 2019. Slow growth dictates non-heritable antibiotic resistance
1201   in Salmonella enterica. Sci Signal 12.
1202
1203 116. Ducas-Mowchun K, De Silva PM, Crisostomo L, Fernando DM, Chao TC, Pelka P,
1204   Schweizer HP, Kumar A. 2019. Next Generation of Tn 7-Based Single-Copy Insertion
1205   Elements for Use in Multi- and Pan-Drug-Resistant Strains of Acinetobacter baumannii.
1206   Appl Environ Microbiol 85.
1207
1208 117. Harding CM, Tracy EN, Carruthers MD, Rather PN, Actis LA, Munson RS. 2013.
1209   Acinetobacter baumannii strain M2 produces type IV pili which play a role in natural
1210   transformation and twitching motility but not surface-associated motility. mBio 4.
1211
1212 118. Carruthers MD, Nicholson PA, Tracy EN, Munson RS. 2013. Acinetobacter baumannii
1213   utilizes a type VI secretion system for bacterial competition. PLoS One 8.
1214
1215 119. Kumar A, Dalton C, Cortez-Cordova J, Schweizer HP. 2010. Mini-Tn7 vectors as genetic
1216   tools for single copy gene cloning in Acinetobacter baumannii. J Microbiol Methods
1217   82:296–300.
1218
1219 120. Leus I V., Adamiak J, Trinh AN, Smith RD, Smith L, Richardson S, Ernst RK, Zgurskaya
1220   HI. 2020. Inactivation of AdeABC and AdeIJK efflux pumps elicits specific nonoverlapping

1221   transcriptional and phenotypic responses in Acinetobacter baumannii. Mol Microbiol
1222   114:1049–1065.
1223

1224 121. Leus I V., Weeks JW, Bonifay V, Smith L, Richardson S, Zgurskaya HI. 2018. Substrate
1225   Specificities and Efflux Efficiencies of RND Efflux Pumps of Acinetobacter baumannii. J
1226   Bacteriol 200.
1227

1228 122.   M100 Performance Standards for Antimicrobial Susceptibility Testing A CLSI supplement
1229   for global application.
1230

1231 123. Figurski DH, Helinski DR. 1979. Replication of an origin-containing derivative of plasmid
1232   RK2 dependent on a plasmid function provided in trans. Proc Natl Acad Sci U S A
1233   76:1648–1652.
1234

1235 124. Choi KH, Gaynor JB, White KG, Lopez C, Bosio CM, Karkhoff-Schweizer RAR,
1236   Schweizer HP. 2005. A Tn7-based broad-range bacterial cloning and expression system.
1237   Nat Methods 2:443–448.
1238

1239 125. HUGH R, REESE R. 1968. A comparison of 120 strains of Bacterium anitratum Schaub
1240   and Hauber with the type strain of this species. Int J Syst Evol Microbiol 18:207–229.
1241

1242 126. DUTHIE ES, LORENZ LL. 1952. Staphylococcal coagulase; mode of action and
1243   antigenicity. J Gen Microbiol 6:95–107.
1244

1245 127. Rosen DA, Hooton TM, Stamm WE, Humphrey PA, Hultgren SJ. 2007. Detection of
1246   Intracellular Bacterial Communities in Human Urinary Tract Infection. PLoS Med 4:1949–
1247   1958.
1248