Cross-expression meta-analysis of 695 brain samples reveals coordinated gene expression across spatially adjacent cells

Ameer Sarwar[1] (ameer.sarwar@mail.utoronto.ca), Mara Rue[2] (mara.rue@alleninstitute.org)

Leon French[3] (leon.french@utoronto.ca), Helen Cross[2] (crossh@carleton.edu), Sarah Choi[3]

(sr.choi@mail.utoronto.ca), Xiaoyin Chen[2] (xiaoyin.chen@alleninstitute.org) and Jesse Gillis[3]*

(jesse.gillis@utoronto.ca)

[1]Department of Cell and Systems Biology and Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto, ON, Canada

[2]Allen Institute for Brain Science, Seattle, WA, USA

[3]Department of Physiology and Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto, ON, Canada

*Corresponding Author

Abstract

Spatial transcriptomics promises to transform our understanding of tissue biology by molecularly

profiling individual cells *in situ*. A fundamental question they allow us to ask is how nearby cells

orchestrate their gene expression. Rather than focus on how these cells (samples) communicate

with each other, we reframe the problem to investigate how genes (features) coordinate their

expression between neighboring cells. To study these phenomena – called cross-expression –

we compare all genes to find pairs that coordinate their expression between adjacent cells,

thereby avoiding curating gene lists or annotating cell types. Our end-to-end method recovers

ligand-receptor pairs as cross-expressing genes and finds gene combinations that mark

anatomical regions, complementing marker gene-based region annotation. Leveraging the

overlapping genes across different panels, we use multiple atlas-scale adult mouse brain datasets

(~25 million cells, 695 samples, 8 technologies) to create an integrated, meta-analytic cross-

expression network, whose communities are enriched in spatial processes such as synaptic

signaling and G protein coupled receptor activity. Highlighting cross-expression's biological utility,

our network shows that genes *Drd1* and *Gpr6*, which are individually implicated in Parkinson's

disease (PD) and are being pursued as therapeutic targets, are cross-expressed within the

striatum, hinting at their joint role in PD pathophysiology. We provide an efficient R package

(https://github.com/gillislab/CrossExpression/) to computationally analyze and visually explore

cross-expression patterns, which allow us to better understand how genes coordinate their

expression in space to perform tissue-level functions.

Keywords

spatial transcriptomics; cross-expression; cell-neighbor relations; gene expression coordination;

spatial signaling; meta-analytic network

Background

Spatial transcriptomic technologies record cells' gene expression alongside their physical locations, enabling us to understand how they influence one another within the tissue [1]. Focusing on select genes, imaging-based platforms profile expression at the single cell level, giving us a high-resolution snapshot of spatial gene expression [2–8]. They have facilitated numerous studies on defining local spatial patterns [9–12], finding gene covariation in spatial niches [13–19], elucidating cell-cell interactions using ligand-receptor expression [20–28], and determining spatial cell type heterogeneity and tissue structure [29–31]. These efforts have resulted in a greater understanding of tissue biology, culminating in the generation and exploration of reference atlases [32–35].

Imaging-based platforms can now profile over a thousand genes in millions of cells [2–8], generating large amounts of data ripe for biological discovery. These data can be analyzed at a high resolution, where we can investigate individual genes, single cells, and their spatial relations. The dominant framework in this space is the cell-cell communication methods [20–28], which infer co-localized ligands and receptors in neighboring cells. These approaches typically rely on curated ligand-receptor databases to infer interactions between cell types. Despite rapid progress in this area, these methods have important limitations. First, the reliance on extant databases limits novel discoveries to well-known ligands and receptors, thereby overlooking potential interaction partners present in the gene panel. Second, the use of cell types requires that the annotations are sufficiently accurate, but this is currently an open problem owing to the limited size of the gene panels, often requiring 'label transfer' from matched single-cell RNA-seq datasets. Indeed, the communication inference is made at the aggregate, cell type level when the underlying data allows a fully 'bottom-up' analysis at the single cell level. Third, the cell-centric perspective makes it difficult to integrate datasets across experiments, limiting the inferences to single studies. Because large-scale datasets are rapidly being collected, an approach is needed to develop a 'common map' between them to discover reproducible biological signals.

The gene-centric framework is an alternative to the cell-cell inference approaches. By

prioritizing genes (features) over cells (samples), it allows comparisons across datasets using

genes shared between their panels. Moreover, when the cell-cell interaction methods use ligand-

receptor co-localization, they implicitly leverage the gene-centric perspective, where inferences

are made if the relevant genes are expressed in neighboring cells more frequently than expected

by chance. A number of studies have explicitly used the gene-centric approach. For example,

Haviv et al. (2024) model how gene-gene co-expression within the same cells changes across

spatial niches [14]. Studies by Miller et al. (2021) and Li et al. (2023) consider gene-gene

coordination between spatially adjacent cells using the global bivariate Moran's I statistic [18,19].

Whereas Haviv et al. (2024) ignore gene-gene coordination between neighboring cells, Miller et

al. (2021) and Li et al. (2023) require ligand-receptor databases, thus limiting their scope.

Additionally, the spatial correlation methods, such as the global bivariate Moran's I statistic, are

known to increase false positives because they do not account for "within location" association.

For example, if genes A and B are both co-expressed in neighboring cells, then they would appear

correlated across neighbors even when the two cells function fully autonomously.

Here, we introduce "cross-expression", which models the degree to which genes

coordinate their expression across neighboring cells. Avoiding the use of curated databases and

cell type labels, our fully end-to-end and bottom-up method uses the raw gene expression and

cell location matrices to determine which gene pairs, among all possible pairs, coordinate their

expression between neighboring cells. We explicitly account for spurious associations induced by

"within cell" co-expression patterns, thereby revealing genuine coordination once cell intrinsic

processes have been controlled. A consequence of our method is that it recovers ligand-receptor

pairs as cross-expressed genes, e.g., *Sst* and *Sstr2*, thus extending the well-developed cell-cell

communication framework. Its gene-centric approach facilitates integrative analysis, where we

create a cross-expression meta-analytic network using 13 datasets spanning ~25 million cells,

695 brain samples, and 8 technological platforms. Our network shows highly modular structure,

with communities enriched for gene ontology (GO) terms like synaptic signaling, neurotransmitter

regulation, cell adhesions, G protein receptor activity, and other spatial processes. Highlighting

its biological utility, the network reveals that across numerous samples the genes *Drd1* and *Gpr6*,

which are individually known to play roles in Parkinson's disease (PD) and are being pursued as

therapeutic targets, are cross-expressed in the striatum, hinting at their joint function in the central

anatomical locus in PD pathophysiology. We also show that the cross-expression patterns in one

dataset reliably predict those in other datasets as well as that similar anatomical regions have

similar cross-expression profiles, indicating that our method reliably detects subtle variations in

gene expression across batches and anatomical space. To facilitate the analysis and exploration

of cross-expression patterns, we provide a fully vectorized R package that processes hundreds

of thousands of gene pairs across millions of cells within minutes on a standard personal laptop,

allowing in-depth analyses of how genes coordinate their expression in space to perform tissue-

level functions.

Results

Cross-expression overview

Cross-expression is defined as the degree to which two genes coordinate their expression across

neighboring cells (Fig. 1a). Specifically, it asks whether the expression of gene A in a cell is

associated with the expression of gene B in the cell's (nearest) neighbor. To properly capture

these patterns, we control for co-expression or, more generally, the association induced between

neighboring locations due to the association present within the same location. In particular, if

genes A and B are co-expressed in both the neighboring cells, then they would appear correlated

across these cells even when the cells function completely independently. We control for this by

defining cross-expression mutually exclusively, namely gene A is expressed in the cell without

gene B and gene B is expressed in the neighbor without gene A.

We quantify cross-expression in three statistically robust and highly interpretable ways.

First, we use the hypergeometric test to compute p-values (Fig. 1b). We begin by finding each

cell's nearest neighbor to create a cell-to-neighbor 'mapping'. While cells almost never have two

nearest (equidistant) neighbors, two or more cells can have the same neighbor. The

hypergeometric test is performed on these cell-neighbor pairs. For genes A and B, we count the

number of pairs where the cell expresses gene A (sample size) and the number of pairs where

the neighbor expresses gene B (possible successes). We then compute the intersection

(observed successes), namely the number of pairs where the cell expresses A and the neighbor

expresses B. Together with the total number of pairs (population size), we use the hypergeometric

cumulative distribution function (CDF) to analytically compute the probability of observing as many

or more successes, giving us a p-value for the cross-expression of genes A and B.

Second, we quantify the effect size by comparing the number of cross-expressing pairs to

the number of co-expressing cells (Fig. 1c). The central idea is that if gene B is randomly assigned

to cells while gene A is held constant, then we expect some co-expression and some cross-

expression simply by chance, where their expected ratio is 1. This ratio is our effect size, with

values greater than 1 indicating more cross- than co-expression, and we represent it till the n-th

neighbor using concentric rings in the bullseye plot.

Third, we compute a Pearson correlation between gene A in each A-expressing cell and

gene B in its nearest neighbor. This yields a directed set of cell-neighbor pairs, capturing how B

varies across the local context of A. Some neighbors are shared by multiple cells, reflecting spatial

centrality; this weighting mirrors the biological structure of the tissue, where certain cells may

influence more of their surroundings. Although this induces non-uniform contributions, each pair

is treated independently from the target cell's perspective, preserving statistical validity for

correlation.

Although we focus on individual cells, groups of cells may form spatial niches and gene

expression may be coordinated between niches. To assess cross-expression at this coarser

resolution, we average a gene's expression in a cell with its expression in the neighbors (Fig. 1d),

thus smoothing it within a spatial niche, with the number of neighbors forming the niche size.

Accordingly, cross-expression can be compared across niches by, for example, finding

associations between smoothed niche-specific gene expression profiles.

To enable these analyses, we provide an efficient software package in R that requires the

gene expression and cell location matrices as inputs, and outputs a gene-gene p-value matrix

that facilitates downstream analyses, such as the creation of cross-expression networks (Fig. 1e).

The package also contains functions for computing effect sizes, making bullseye plots, smoothing

gene expression, viewing cross-expressing cells *in situ*, and assessing if cross-expression is

spatially enriched. Collectively, the cross-expression framework uses spatial information to

discover how genes coordinate their expression across neighboring cells, thereby providing a

useful analytical framework for deeply exploring spatial transcriptomic data.

Fig. 1 | Cross-expression analysis. a, Cross-expression is the mutually exclusive expression of
genes between neighboring cells. If either cell expresses both genes, the cell pair is not
considered to cross-express. **b**, The probability that two genes cross-express is modeled by the
hypergeometric distribution, where the 'successful trials' are the cell-neighbor pairs where the cell
expresses gene A and the neighbor expresses gene B. **c**, Cross-expression is compared to co-
expression to quantify the effect size, where the number of neighbors with gene B is compared to
the number of cells co-expressing genes A and B. 'Sampled cells' (center) are those expressing
gene A and neighbors are concentric rings, with the order indicating the *n*-th neighbor. **d**,
Averaging gene expression between cells and their neighbors smooths it, extending cross-
expression analysis from cell pairs to regions. Number of neighbors is the kernel size. **e**, Software
inputs are the gene expression and cell location matrices, and the output is a p-value matrix,
which enables downstream analyses, such as the creation of cross-expression networks. Created
with BioRender.com.
181

Cross-expression recovers ligand-receptor pairs and reveals coordinated gene expression

profiles across the tissue

To study cross-expression, we used BARseq (barcoded anatomy resolved by sequencing) to

collect data from a whole mouse brain. This dataset profiled expression in 1 million cells across

16 sagittal slices, using a gene panel of 104 cortical cell type markers and 25 ligands and

receptors, including neuropeptides, their receptors, and monoamine neuromodulatory receptors.

Because receptors and the enzymes that synthesize the corresponding ligands are often

expressed in nearby cells [20–28], we reasoned that these genes should show cross-expression.

As an example, we find that across the cortical somatosensory nose region and visceral areas

(Fig. 2a), the neuropeptide somatostatin *Sst* and its cognate receptor *Sstr2* are cross-expressed

(Fig. 2b, left, p-values ≤ 0.01 and 0.05, respectively). Indeed, these genes are consistently

expressed across neighboring cells (Fig. 2b, right), a pattern that is otherwise difficult to discover

without prior knowledge. While cross-expression recovers ligand-receptor genes, it does not imply

that they form synaptic or endocrine connections, since spatial transcriptomic measurements are

restricted to the cell body.

Next, we explore the bullseye plots, which allow us to quantify the effect size by comparing

cross-expression with co-expression. For *Sst* and *Sstr2* in the somatosensory nose (2,015 cells)

and visceral regions (1,603 cells), we see a bullseye pattern with low co-expression and high

cross-expression that decreases for distant neighbors (Fig. 2c). Specifically, for these regions the

bullseye score ratio between the first neighbor and the central cell is 1.8 and 1.6, respectively,

whereas the ratio between the averaged second-to-tenth neighbor and the central cell is 1.3 and

1.2. These findings suggest that for central cells expressing one gene in a pair, a higher proportion

of adjacent neighbors, but not the more distant ones, express the cognate gene within the local

spatial niche, underscoring the specificity and resolution with which patterns of coordinated gene

expression can be recovered. We next compare the bullseye plots for gene pairs with and without

cross-expression (Fig. 2d), finding that the former match the patterns just described. To quantify

this, we compare the bullseye scores of the nearest neighbors with those of cells expressing gene

A, discovering that this ratio is much greater for genes that cross-express than for those that do

not (Fig. 2d, inset, Mann-Whitney U test, p-value $\leq$ 0.001, median ratios: 1.5 and 0.9,

respectively). Notably, this ratio is approximately 1 for genes that do not cross-express,

suggesting that here gene B is expressed in neighbors and cells alike. Hence, the bullseye

approach intuitively visualizes and quantifies the effect size, making it suitable for downstream

analysis, such as comparing cross-expression between different regions.

We next conducted brain-wide analysis and found that 20% of possible ligand-receptor

gene pairs and 4% of non-signaling gene pairs are cross-expressed, thus generating novel

candidates that potentially encode functionally relevant interactions. In fact, these patterns are

spatially enriched, where most gene pairs cross-express in a few slices and some cross-express

in multiple slices (Extended Data Fig. 1a). We now highlight some notable examples of cross-

expression for both signaling and non-signaling genes. The dopamine receptor D_1 (*Drd1*) and

proenkephalin (*Penk*) are strongly cross-expressed (Extended Data Fig. 1b), with discernible

spatial enrichment in the striatal regions. *Drd1* is involved in the reward system [36,37] while *Penk*

generates opioids that modulate fear response [38] and nociception [39,40], suggesting that these

genes may be involved in avoidance behavior. Indeed, Penk is strongly co-expressed with the

dopamine receptor D_2 (*Drd2*) (Pearson's *R* = 0.72 in scRNA-seq striatal data; *Drd2* is not in our

gene panel), indicating that the D1 and D2 neurons are spatially intermingled, allowing them to

play interrelated roles in motor control [41]. Additionally, we find that the somatostatin receptor

Sstr2 cross-expresses with vasoactive intestinal polypeptide receptor 1 (*Vipr1*/*VPAC1*) in the

cortex (Extended Data Fig. 1c), suggesting a potential complementary interaction in modulating

local neuronal circuits and influencing neuroendocrine signaling pathways [42]. Beyond the

signaling genes, we note that the fibril-associated *Col19a1* (collagen type XIX alpha 1 chain), a

gene involved in maintaining the extracellular matrix (ECM) integrity [43,44], cross-expresses with

C1ql3 (complement C1q like 3) (Extended Data Fig. 1d), whose secretion in the ECM facilitates

synapse homeostasis and the formation of cell-cell adhesion complexes [45,46]. Finally, our

analysis reveals that *Marcksl1* (myristoylated alanine-rich C-kinase substrate), which is involved

in adherens junctions and cytoskeletal processes [47,48], cross-expresses with actin beta (*Actb*)

(Extended Data Fig. 1e), hinting at their involvement in local tissue architecture [49]. Taken

together, the cross-expression analysis not only reveals expected relationships between signaling

molecules, but it also discovers genes implicated in the tissue microenvironment. Accordingly,

cross-expression is an unbiased, data-driven framework for finding genes with orchestrated

spatial expression profiles, with potential for novel discovery increasing with varying sizes and

compositions of the gene panels.

We have thus far investigated cross-expression between cells and their neighbors. Yet,

gene expression may be coordinated between more distant neighbors or between large spatial

niches. The former is facilitated by changing the rank of the nearest neighbor tested. The latter is

enabled by smoothing a gene's expression in a cell by averaging it with its expression in nearby

cells, as shown for cortical layer 4 marker *Rorb* (Fig. 2e) and layer 6 marker *Foxp2* (Extended

Data Fig. 1f) in the auditory cortex [32].

Although cross-expression may appear at varying length scales, we focus our analyses at

the cellular level to investigate its signature between individual cells.

Fig. 2 | Cross-expression analysis reveals coordinated gene expression between neighboring cells. a, Sagittal brain slices showing cortical somatosensory nose region (top) and visceral area (bottom) as randomly selected regions of interest. **b**, Neuropeptide somatostatin *Sst* and its cognate receptor *Sstr2* cross-express in regions shown in (a). Points indicate cells and colors indicate gene expression (left), with cross-expressing cell pairs highlighted (right). **c**, Bullseye scores for *Sst* and *Sstr2* in the regions shown in (a, b). The scores are reported as ratio of cross- to co-expression. **d**, Bullseye scores for cross-expressing (significant) and non-cross-expressing (not significant) gene pairs in the somatosensory nose region. 'Cell' corresponds to the central ring in (c), and the red rectangle highlights the first neighbor/ ring. Inset, ratio of bullseye scores for the first neighbor to the central cells for cross-expressing and non-cross-expressing genes. Central line, median; box limits, first and third quartiles; whiskers, ±1.5x interquartile range; points, outliers. **e**, Smoothed gene expression for different numbers of neighbors for the auditory cortical layer 4 marker gene *Rorb*. Created with BioRender.com.

Cross-expression is driven by subtle and consistent cell subtype compositional differences

Having seen that cross-expression recovers coordinated spatial gene expression, we now explore its relationship with cell type heterogeneity. For this purpose, we use another BARseq dataset [35] that was recently used to create a mouse cortical cell type atlas using the same 104 excitatory marker genes as before. Here, we observe that genes cross-express between cells of the same and of different types. For example, *Gfra1* and *Foxp2* are cross-expressed within the same cell type L4/5 IT (intratelencephalic) and between different cell types Car3 or CT (corticothalamic) and L4/5 IT (Fig. 3a). In general, genes vary greatly in terms of the cell type labels of cross-expressing cell pairs (Fig. 3b). For instance, for some gene pairs 40% of the cell pairs have the same cell type label while in others as many as 90% of the cell pairs belong to different cell types (Extended Data Fig. 2a). Moreover, some genes involve many while others involve few cross-expressing cells. For example, in the analyzed data the median number of cross-expressing cell pairs is 2,378, and 27% of genes involve over 4,000 while only 5% involve 400 or fewer pairs (Extended Data Fig. 2b), indicating that the density of gene cross-expression is highly variable. Interestingly, cell type purity – the proportion of cell pairs with the same type – decreases as more cell pairs cross-express (Fig. 3c, Spearman's $\rho = -0.46$), highlighting a potential role for spatially intermingled cell types in patterns of cross-expression.

To assess the influence of spatial cell type composition more broadly, we use our hierarchical cell type taxonomy [35], where types at a higher-level divide into subtypes at a lower level. Using cross-expressing glutamatergic cells, we find that 64% of the pairs consist of different cell subtypes (Fig. 3d, right-tailed Wilcoxon signed-rank test, different labels ≥ same labels, p-value ≤ 0.0001, Extended Data Fig. 2c), suggesting that subtle cell type differences drive cross-expression. However, for cross-expressing GABAergic cells, we find that only 44% of the pairs have different cell subtype labels (Extended Data Fig. 2d-e right-tailed Wilcoxon signed-rank test, different labels ≥ same labels, p-value = 1), reflecting the fact that our gene panel is optimized to

292   detect cell subtype differences between excitatory, but not inhibitory, neurons. Crucially, we

293   observe that cells of one type consistently cross-express with cells of another type (Fig. 3e,

294   Extended Data Fig. 2f), indicating that cross-expression recapitulates patterns of cell type

295   composition. Since cell type labels are assigned based on the expression of many genes,

296   repeated spatial proximity of cell types results in the cross-expression of their marker genes.

Fig. 3 | Cross-expression patterns are discovered independently of cell type labels but are driven by cell type heterogeneity. a, Cells of the same (yellow) and different (green) types cross-express genes *Gfra1* and *Foxp2* in the auditory cortex. Discovering cross-expression relations between this or any other gene pair does not require cell type labels. **b**, Genes are cross-expressed across numerous cells, with the dot size indicating the number of cell-neighbor pairs and the color showing the proportion of pairs with the same label (cell type purity). **c**, Cell type purity against the number of cross-expressing cell-neighbor pairs. Each point is a gene pair from (b), and shaded area is 95% confidence interval. **d**, Number of cell-neighbor pairs with the same or different cell subtype labels given that they were both labeled 'glutamatergic' at the higher level of the cell type hierarchy. Each point is a cross-expressing gene pair. **e**, Heatmap showing the normalized frequencies of cell type label combinations between cross-expressing cells. Created with BioRender.com.

Cross-expression offers a common framework for analyzing multiple studies and the meta-analytic network reveals uniquely spatial biological processes

After analyzing ligand-receptor genes and cell type variability in terms of cross-expression, we asked whether its gene-centric approach can be used to simultaneously analyze multiple studies. To this end, we downloaded 13 adult mouse brain datasets (see methods), which were collected using 8 different platforms (4 commercial and 4 laboratory-based), spanning ~25 million cells (after dataset-specific quality control; median 724,790 cells) across 695 large samples/ slices (hemi-coronal, coronal, or sagittal) (Fig. 4a). Although the gene panels had varying sizes and compositions, they were sufficiently overlapping to allow integrative analyses (Fig. 4b; median pairwise overlap is 77 genes), reflecting that well-studied and informative genes are enriched across panels [50].

We first investigated cross-expression replicability within datasets. Here, we predicted the p-values of a held-out slice using the averaged correlations of the remaining slices. Our cross-validation-based classification revealed that the predictions (AUROCs) improved and then plateaued as more samples were included (Fig. 4c), suggesting that the sample ensemble progressively became more representative, e.g., due to the inclusion of anatomically adjacent slices. While the improvements were consistent across datasets, the AUROC scores varied between them, indicating the presence of potential batch effects, including differences in data quality between samples, enrichment of spatially variable or localized genes in the panels, and

large gaps between the slices, etc. In general, the cross-expression patterns within datasets are

broadly replicable, where including more samples improves performance, possibly due to better

representation of the underlying anatomical regions.

Next, we examined whether cross-expression correlations in one dataset can predict the

p-values in another dataset using the genes shared between their panels. We found high

replicability for some datasets and modest performance for others (Fig. 4d), indicating that

integrative meta-analysis is possible using at least a subset of these data. Although our results

here are reported as median AUROCs, the performance can suffer if (say) in one dataset most

slices are anterior while in the other most are posterior. Consequently, our subsequent analyses

focus on replicability of cross-expression between (anatomically homologous) slices across

datasets.

Finding that cross-expression is broadly replicable between datasets, we created a meta-

analytic network, where the nodes represent genes and the edges indicate cross-expression (Fig.

4e). Our network includes gene pairs only when they are cross-expressed in two or more datasets,

thus excluding those cross-expressed in just one dataset. The network shows high modularity,

with communities enriched for gene ontology (GO) terms representing spatially-mediated

biological processes. For example, the community 'synapse organization' (red) is connected with

the 'neurotransmitter regulation' community (pink), recapitulating ligand-receptor interactions

present in these data. Likewise, the 'cell adhesion' community (light green) is connected to the

'extracellular and vascular' community (dark green), implicating genes potentially involved in

structural and supportive processes. The 'G protein coupled receptor signaling' community (dark

blue) is diffusely connected to multiple other communities, reflecting these genes' diverse

physiological roles, such as in hormonal and metabolic processes. Similarly, the 'CNS

development' community shows distributed connectivity, likely revealing the cross-expression of

cell type markers with multiple other marker and non-marker genes, e.g., due to transcription

factors (indirectly) specifying cell type identity [51]. Lastly, the network contains a small 'amyloid

precursor protein' community (purple) composed of Alzheimer's disease (AD) genes cross-

expressed with *ApoE*, which helps form amyloid plaques and neurofibrillary tangles [52]. Together,

the cross-expression framework allows us to use the shared genes to perform integrative

analyses across multiple studies, with our meta-analytic network highlighting genes involved in

distinctively spatial biological processes.

Fig. 4 | Cross-expression facilitates integrative meta-analysis of numerous samples across 13 studies. a, Number of cells in each dataset after applying dataset-specific quality control metrics. Red line is the median. **b**, Upset plot showing the overlaps between the gene panels of different datasets. **c**, Predictions of held-out slices' p-values (significant or non-significant) using the average correlation of the remaining slices. Performance (AUROC) is reported as the function of the number of slices used to average the correlation (cross-validation). **d**, Median performance (AUROC) when predicting the p-values of a slice in one dataset using the correlation of slice(s) from another dataset using the genes shared between their panels. **e**, Cross-expression meta-analytic network where the nodes are genes, with edges present if the genes are cross-expressed in two or more datasets. The network is clustered into communities, with the colors and labels indicating gene ontology (GO) annotations, where slash ('/') separates biological process (BP), cellular components (CC), and molecular function (MF). Created with BioRender.com.

Parkinson's disease (PD) genes *Drd1* and *Gpr6* show highly reproducible and localized cross-expression patterns in the striatum

Our meta-analytic network revealed that cross-expressing genes are involved in spatially-mediated biological processes. Since genes were included in the network if they cross-expressed in two or more datasets, we sought to identify an example gene pair that is both highly reproducible and biologically informative. This process is inherently dependent on the researchers' interests, so we focused on *Drd1* and *Gpr6*, which are implicated in Parkinson's disease (PD) due to their hypo- and hyperactivity, respectively [53–60]. Our datasets are registered to the Allen Institute's CCFv3 mouse brain region atlas [61], which allowed us to arrange samples from multiple brains and different datasets in the same anatomical space. Looking at coronal and sagittal sections, we found these genes' cross-expression highly concentrated within a restricted range and effectively absent outside it, suggesting that it is localized within specific anatomical regions (Fig. 5a). Next, we overlaid the Allen CCFv3 atlas on the combined cross-expression signal, finding that it is located within the striatum (Fig. 5b, protruded region), the central locus in PD pathophysiology [62]. Consistent with this observation, we found that their cross-expression within slices is located in the striatal regions but not elsewhere, with its patterns following gross neuroanatomy in the anterior-to-posterior and lateral-to-medial directions (Fig. 5b, panels 1-9 and P-S). Accordingly, the genes *Drd1* and *Gpr6* show

coordinated expression between cells in the striatum, a property observed in multiple slices

across different datasets.

The cross-expression of these genes can help facilitate subsequent studies. Briefly, *Drd1*

is the dopamine receptor D1, whose stimulation by dopamine initiates movement, whereas *Gpr6*

is the G protein coupled receptor 6, whose constitutive activity prevents movement initiation. In

PD, the dopaminergic neurons in the substantia nigra insufficiently activate *Drd1*, and *Gpr6*

exhibits higher baseline activity, leading to severe difficulty in starting movement, the cardinal

symptom of PD [53–60]. While extant therapies increase dopamine (L-DOPA as Levodopa) to

stimulate *Drd1* [63,64], where the drug's effectiveness decreases over time and causes sides

effects, recent clinical trials have explored the inverse agonist CVN424 to inhibit *Gpr6* [55–60].

Although at present these approaches have not been pursued in tandem, their localized cross-

expression suggests that the drugs' staggered or co-delivery might offer complementary,

potentially amplified, effects. In general, once reproducible cross-expression signatures are

discovered, ideally across different samples and studies, one can further investigate gene pairs

of interest, making cross-expression a basic tool for biological research using spatial

transcriptomic data.

Fig. 5 | Cross-expression of genes *Drd1* and *Gpr6* in the striatum across multiple brains and different studies. a, Cross-expression signature (–log10 p-value) of *Drd1* and *Gpr6* genes in the coronal sections ordered in the anterior-to-posterior direction using the Allen Institute CCFv3 mouse brain atlas coordinates (top). Bottom, same as top but with sagittal slices ordered in the medial-to-lateral direction. **b**, Cross-expression score (outer product of coronal and sagittal slices' –log10 p-values) overlaid with the Allen Institute CCFv3 brain region annotations, with the striatum as the protruded (blue) area. Different colors represent the source brain of each slice. Numbers (1-9) and letters (P-S) indicate the position of example coronal and sagittal slices, respectively, within the overall gross neuroanatomy. Created with BioRender.com.

Cross-expression network reveals *Gpr20* as a central gene and discovers possible interaction partners between astrocytes and the brain microvasculature

Having assessed multiple datasets, we now analyze a single study to show how cross-expression

patterns can be used alongside additional information, such as cell type labels. We first

supplemented our network formalism by including (second-order) edges between two genes if

they independently cross-expressed with a third same gene (Fig. 6a). Using the Vizgen MERFISH

data, we created a cross-expression network (Fig. 6b), which contains 200 genes with 382 first-

order, 217 second-order, and 107 dual-order edges. We observe that *Gpr20*, a G protein-coupled

receptor, is a central gene with a high node degree of 40 while the other genes form a median of

4.8 edges (Extended Data Fig. 4a). We performed gene ontology (GO) enrichment for genes

cross-expressed with *Gpr20*, finding functional groups like 'regulation of macromolecule

biosynthetic process', 'regulation of gene expression', and 'regulation of metabolic process'

(Extended Data Fig. 4b, all p-values ≤ 0.05). While some of these genes are co-expressed with

astrocytic and microglial cell type markers (Extended Data Fig. 4c), their global co-expression

with the endothelial marker is higher, where the co-expression profiles were computed using

neighbors cross-expressed with *Gpr20* rather than the entire dataset (Extended Data Fig. 4d,

Mann-Whitney U test, endothelial vs others, all p-values ≤ 0.01; remaining pairwise comparisons,

all p-values > 0.05).

Noting that the neighbors of *Gpr20*-positive cells are involved in the microvasculature, we

next viewed the spatial distribution of cells expressing *Gpr20*, finding that they form contiguous

linear streaks resembling blood vessels (Fig. 6c; anterior slice from mouse brain 1 shown). To test

this observation, we looked at whether the neighbors of *Gpr20*-positive cells also express this

gene and compared it to randomly selected cells, which constitute the expectation that *Gpr20* is

uniformly expressed across space. Consistent with the visualization, we find that cells with *Gpr20*

are surrounded by neighbors that also express this gene, a pattern that disappears for neighbor

order of 50 or more cells (Fig. 6d-e, area under curve (AUC), neighbors vs random cells, 0.69 vs

0.49; right-tailed Wilcoxon signed-rank test, neighbors vs random cells, p-value ≤ 0.0001). Having

seen that cells with *Gpr20* possibly reflect blood vessels, we asked whether these cells are

themselves vascular or whether they line the vasculature, especially since the cells that cross-

express with *Gpr20* are endothelial. We find *Gpr20* is poorly co-expressed with *Igfr1* (Pearson's

R = 0.0024), the vascular/endothelial marker [65–67] in our gene panel, suggesting that it lines

but does not mark the blood vessels. Moreover, it is lowly co-expressed with other cell type

markers (average Pearson's *R*, astrocytes = –0.0027, microglia = 0.0018, oligodendrocytes = –

0.022, neurons = –0.0025), eschewing cell type characterization. Taken together, *Gpr20*, a salient

topological feature of our cross-expression network, seems to be expressed in diverse cell types

that line the blood vessels, reflecting its possible role in the microvasculature.

Cross-expression driven by cell types might be particularly common when two genes

which cross-express with a third gene are co-expressed together, reflecting some common

transcriptional program jointly cross-expressing with neighboring cells. To investigate this, we

reduced co-expression further by specifying that cross-expressing genes must show lack of

significant co-expression, a procedure that yielded a subnetwork, which we further curated by

removing genes with fewer than two edges. Indeed, we find that two genes that independently

cross-express with another gene tend to be co-expressed (Fig. 5f, Extended Data Fig. 5a) and,

as expected, belong to the same cell types, as revealed by their co-expression with cell type

marker genes (Extended Data Fig. 5b). Confirming these results, the subnetwork genes are

enriched in GO groups like 'endothelial cell proliferation', 'positive regulation of vascular

endothelial growth factor production', and 'regulation of endothelial cell migration' (Fig. 6g, all p-

values ≤ 0.05). These results indicate that while cross-expressing genes are present in specific

cell types, the relations between them are functionally suggestive as opposed to simply reflecting

cell type compositional differences, especially since the cell type markers are not cross-

expressed. For example, the astrocytic *EGFR* (epidermal growth factor receptor) cross-expresses

with the vascular *Flt4*/*VEGFR-3* (FMS-like tyrosine kinase 4), *Tek*/*Tie2* (TEK tyrosine kinase/

angiopoietin-1), and *Tie1* (tyrosine kinase with immunoglobulin-like and EGF-like domain 1).

These three vascular receptors promote angiogenesis via the *VEGF* (vascular epidermal growth

factor) ligand [68,69], prevent endothelial cell apoptosis [70,71], and negatively regulate

angiogenesis [72], respectively, thus reflecting their potential role in the brain microvasculature in

coordination with the astrocytes, whose endfeet ensheathe the blood microvessels to constitute

the blood-brain barrier (BBB).

Within the same subnetwork, the astrocytic gene *Ppp1r3g* (protein phosphatase 1

regulatory subunit 3G), which helps convert glucose to glycogen [73], cross-expresses with *Epha2*

(ephrin type-A receptor 2), whose activity makes the BBB more permeable [74], likely enabling

glucose's transport and eventual conversion into glycogen, thereby making this cross-expression

relation relevant for energy metabolism. Indeed, this observation can be used to generate

hypotheses about the (directional) relationship between energy needs within a local

microenvironment and remodeling of the microvasculature, making cross-expression a powerful

approach with which to form testable hypotheses.

Next, we asked whether cross-expression networks change across the brain. Because

gene expression is regional, slices from various areas should show cross-expression between

distinct genes. We assessed this by forming networks for each slice in our sagittal BARseq data.

As expected, we find that adjacent slices have similar networks than distant slices (Fig. 6h,

Spearman's ρ = –0.9), a trend also seen in our BARseq coronal data (Extended Data Fig. 6a,

Spearman's ρ = –0.87) but not when the two datasets are mixed and the "distance" reflects the

difference in the order of slices (Extended Data Fig. 6b, Spearman's ρ = 0.094). Hence, cross-

expression is sensitive to broad spatial variation in gene expression.

Fig. 6 | Networks of cross-expression. a, Cross-expression (edges) between genes (nodes) forms a network (left), where second-order edges (right) between genes share a node (first-order). **b**, Example cross-expression network, with first-order node degree represented by size and edge color showing first-, second-, or dual-order (first-order and second-order) connections. Threshold for the second-order edges is 4. Node color shows community membership assigned by Louvain clustering the second-order network. **c**, Cells are colored based on *Gpr20* expression. Numbered rectangles in the central figure correspond to zoomed-in versions. **d**, Number of neighbors with *Gpr20* given that the source cells also express this gene. **e**, Cumulative sums (after L1 normalization) from (d) for true and randomly selected neighbors. Identity line is dashed. **f**,

Subnetwork created from (b) by pruning edges with significant co-expression and then removing
nodes with degree 1. Nodes are colored by cell types based on their co-expression with marker
genes. **g**, GO functional groups for genes in the subnetwork in (f). **h**, Similarity in the network
structures of nearby and distant brain slices. Shaded area is 95% confidence interval. Created
with BioRender.com.

Cross-expression discovers anatomical marker gene combinations that delineate the

thalamus and refine cortical layer VI boundaries

A key goal in biological research is finding marker genes, such as those that identify cell types,

e.g., *Olig1* for oligodendrocytes [32], anatomical regions, e.g., *Rorb* for cortical layer IV [35], and

functional modules, e.g., *Trpc4* for lateral septum in social behaviors [75]. In addition to using

individual markers, one can use co-expression to discover marker gene combinations, an

approach that uses single-cell or single-nucleus RNA-seq databases [76]. Although co-expression

provides more combinations than individual markers, it relies on measurements from the same

cells, thereby underutilizing the spatial relations between cells in spatial transcriptomics datasets.

Leveraging the spatial dimension to discover marker gene combinations, we asked

whether cross-expressing genes can delineate anatomical regions, including putative functional

modules. As an example, we found that cross-expression between *Lgr6* and *Adra2b* delineates

the thalamus even though these genes are expressed throughout the brain (Fig. 7a). Specifically,

while 48% of *Lgr6*- and 57% of *Adra2b*-expressing cells are thalamic, 91% of their cross-

expressing cell-neighbor pairs are in the thalamus (Extended Data Fig. 3a), underscoring the

spatial enrichment of their cross-expression signature (Extended Data Fig. 3b). We find that *Lgr6*

also cross-expresses with *Ret* in the thalamus despite brain-wide expression of both genes (Fig.

7b, Extended Data Fig. 3c). Next, we examined whether *Adra2b* and *Ret*, both of which cross-

express with *Lgr6*, show enriched co-expression in the thalamus. We find that they are indeed co-

expressed within the thalamus but not in the rest of the brain (Fig. 7c), e.g., 89% of their co-

expressing cells are in the thalamus, thus serving as robust combinatorial markers.

To evaluate whether the combinatorial marker-based approach is reliable, we asked

whether single gene markers, when assessed for cross-expression, rediscover the anatomical

locations. Using the BARseq cortical cell type atlas data [35], we assessed cross-expression

between cortical layer 6 marker *Foxp2* and ubiquitously expressed gene *Cdh13*. We discover that

cross-expression between these genes delineates layer 6 boundary (Fig. 7d), further supporting

the view that combinatorial anatomical markers can be discovered using cross-expression.

Indeed, the layer 6 boundary recovered by cross-expression captures additional L6 IT neurons

whereas *Foxp2*-based boundary overlooks these cells, indicating that combinatorial markers can

refine extant anatomical regions. More generally, this process leverages the spatial enrichment

of cross-expression, where we assess whether cross-expressing pairs are closer to other cross-

expressing pairs than to randomly selected cells. Our framework therefore discovers gene pairs

that annotate anatomical regions and refine extant boundaries, thus extending the single gene

and co-expression-based approaches.

Fig. 7 | Cross-expression can discover combinatorial anatomical markers. a, Comparing the thalamus (left) to the rest of the brain, genes *Lgr6* and *Adra2b* are widely expressed across multiple brain regions (middle) but are preferentially cross-expressed in the thalamus (right). **b**, Same as in (a) but for genes *Lgr6* and *Ret*. **c**, Genes cross-expressing with *Lgr6* in (a) and (b) co-express in the thalamus. **d**, Cross-expression of *Cdh13* with cortical layer 6 marker *Foxp2* (middle) recapitulates layer 6 boundaries (right, cf. left). Created with BioRender.com.

Cross-expression signal is replicable across datasets, and global co-expression between

spatial and single cell datasets indicates reliable cell segmentation

Two sources of non-biological variation in spatial transcriptomics [2–8] are batch effects, which

result from technical differences between experimental runs, and cell segmentation, which draws

boundaries around and assigns transcripts to cells, a process that can alter gene expression

profiles and affect downstream analysis, including cross-expression.

We assess batch effects by comparing cross-expression between corresponding slices

across biological replicates. The MERFISH data contains three replicates with three slices each,

where the slices are sampled from roughly the same position. We find that the cross-expression

signature is highly similar across replicates. For example, the average correlation for the anterior

slices between the three replicates is 0.83 (Fig. 8a), with similar findings for the middle and

posterior slices (Extended Data Fig. 7a-b, Spearman's ρ = 0.81 and 0.8, respectively).

We next assessed the degree to which cross-expression within the BARseq sagittal or

coronal experiments [35] is similar to that between experiments. To this end, we compared cross-

expression patterns between brain slices. As expected, the cross-expression profiles are more

similar within brains than between brains (Fig. 8b, Mann-Whitney U tests, FDR-corrected, coronal

vs sagittal, p-value = 0.2, coronal vs mixed, p-value ≤ 0.001, and sagittal vs mixed, p-value ≤

0.001), suggesting that the sectioning procedure samples different brain regions and therefore

reveals distinct underlying gene expression profiles. Supporting this result, we find that the same

anatomical regions (per Allen CCFv3 brain atlas [61]) across brains have more similar cross-

expression profiles than do different regions within or between brains (Fig. 8c, Mann-Whitney U

test, different regions vs same regions, p-value ≤ 0.001). Noting that the sagittal and coronal

brains contain the same regions in the dorsal to ventral directions, we asked whether the cross-

expression is similar in this shared dimension. Here, we computed the density of cross-expressing

cells in the dorsal to ventral direction and compared these distributions across the two brains,

finding that 99% (without FDR correction) of the genes did not have significantly different density

profiles (Fig. 8d), suggesting that the cross-expression patterns are highly similar across batches

at the whole-brain level.

Having found that the cross-expression profiles are generally robust, we assessed cell

segmentation at a global level by comparing gene co-expression between the single cell RNA-

sequencing (scRNA-seq) [32] and spatial transcriptomic data. We reasoned that scRNA-seq does

not require segmentation and therefore captures genes co-expressed within the cell's boundaries

(Fig. 8e). Because cell segmentation alters transcript assignment, it could change co-expression

in spatial transcriptomic data. Reassuringly, we find a strong association between gene co-

expression in the scRNA-seq and spatial transcriptomic data (Fig. 8f, Pearson's R = 0.83). We

further examine whether this correlation is sufficiently strong by comparing co-expression

between scRNA-seq and single-nucleus RNA-sequencing (snRNA-seq) [77] (Fig. 8g, Pearson's

R = 0.86), finding agreement between the two comparisons (R = 0.83 vs. R = 0.86). These results

imply similar levels of technical variability between platforms while suggesting that gene co-

expression is congruent between scRNA-seq and spatial transcriptomic data.

The data in our work was processed using CellPose [78], a deep learning-based cell

segmentation algorithm. A recent benchmarking study showed that it outperforms other methods

on a variety of metrics [79]. In fact, it uses the nuclear stain DAPI as a cell landmark and forms

boundaries using cytoplasmic signal, such as the transcript distributions, making it the state-of-

the-art segmentation algorithm on a variety of assessments. Further, the cell segmentation

algorithms are continuously being improved [80], allowing users to re-segment and reanalyze

their data. Most importantly, the analysis conducted using the cross-expression framework may

suffer if segmentation is performed poorly, but the validity of the concept and the soundness of its

statistics do not rely on this potential artefact and, with rapid improvements in data quality, the

inferences drawn from it will become increasingly more reliable.

Moreover, we assessed the relationship between cross-expression and noise in gene

expression measurement. Since the algorithm requires binarizing the expression matrix, an

appropriate threshold needs to be applied prior to analysis. To count a gene as expressed in a

cell, we applied thresholds of 1 to 10 molecules, finding that the cross-expression patterns are

generally concordant across these noise levels (Extended Data Fig. 8a-b, median Pearson's R =

0.88). Importantly, our framework is agnostic to and compatible with multiple models of gene

expression noise [81], and once an appropriate threshold has been applied, the resultant

expression matrix can be used for cross-expression analysis.

Finally, we explored the patterns of cell-neighbor relations and found that over 60% of cells

are the nearest neighbors of exactly one cell but the remaining cells are the nearest neighbors of

two or more cells (Extended Data Fig. 9a). Patterns such as these may be biologically important

if the 'neighbor' cell plays a central role in the local microenvironment, so deviations from one-to-

one mappings should be captured by statistical analyses. To investigate that our results are

consistent across these patterns, we compared cross-expression in one-to-one against many-to-

one mappings and with the full dataset, finding an average Pearson's correlation of 0.96

(Extended Data Fig. 9b). Importantly, our procedure is consistent with the assumption of

independent sampling because while a cell may be the nearest neighbor of multiple cells, each

cell-neighbor pair is statistically independent.

We facilitate these analyses by providing a highly efficient R package. A laptop with 16 GB

RAM can test for cross-expression in large datasets containing hundreds of thousands of cells

and thousands of genes within minutes (Fig. 8h). At present, most (commercial) imaging-based

platforms cannot profile gene panels of this magnitude [2–8], though such capabilities are

anticipated. Our software's performance makes it well-suited for analyzing current and future

spatial transcriptomic datasets.

Fig. 8 | Assessing batch effects, cell segmentation, and software runtime. a, Correlation between cross-expression signatures across three biological replicates. **b**, Correlation between cross-expression signatures within (sagittal or coronal) and between (mixed) brains. Positive signal between brains likely reflects the fact that the sagittal and coronal brains both contain regions in the dorsal to ventral direction. **c**, Correlation between cross-expression signatures between the same anatomical regions across brains or between different anatomical regions across or within brains. **d**, Density of cross-expressing cells in the dorsal to ventral directions is compared across the sagittal and coronal brains. Significant p-values (without FDR correction) indicate that a cross-expressing gene pair has different densities across the two brains. Red dotted line is the significance threshold at alpha = 0.05. **e**, Single cell RNA-sequencing (scRNA-seq) profiles cells' gene expression without cell segmentation. Co-expression between scRNA-seq and spatial transcriptomic data helps diagnose segmentation artifacts. **f**, Gene co-expression in spatial transcriptomic and in scRNA-seq data. Each point is a gene pair. **g**, Gene co-expression in single-nucleus RNA-sequencing (snRNA-seq) and in scRNA-seq data. Same gene panel is used in (f) and (g). **h**, Software runtime for varying numbers of genes and cells using a personal laptop with 16 GB RAM. Created with BioRender.com.

Discussion

Cross-expression allows us to study gene-gene networks that reflect how nearby cells influence each other by coordinating their gene expression. Using this framework, we recapitulated known ligand-receptor interactions at the single cell level, revealing biologically meaningful tissue phenotypes. We further showed that cross-expression is discovered without cell type labels but often reflects cell subtype compositional differences in the form of marker gene cross-expression. We also revealed that its gene-centric perspective enables integrative meta-analysis, where many studies can be combined to find robust biological signals, such as the cross-expression of *Drd1* and *Gpr6* in the striatum. Moreover, it helps us perform deeper analyses of individual studies, revealing the relationships between astrocytes and the brain microvasculature, and discovers paired markers for anatomical region annotation. Together, cross-expression is a powerful way of analyzing spatial transcriptomic data and allows us to study gene-gene relations between adjacent cells, thereby fully harnessing the high-throughput of these technologies.

The cross-expression framework complements current approaches analyzing spatial transcriptomic data, such as those exploring niche-specific co-expression patterns [13–19]. Specifically, niche-specific cross-expression networks may be compared with co-expression networks to examine if inter-cellular relations are associated with intra-cellular gene programs and vice versa. This may be approached at different, potentially hierarchical spatial scales to reveal spatial gene expression programs within the tissue. Moreover, the cross-expression patterns can be quantified in multiple ways, such as using mutual information or graphlets, allowing investigations into the best approaches that capture the signal of interest. For example, just as co-expression relations can be measured using the Pearson's correlation coefficient, cross-expression patterns may be investigated from numerous perspectives to discover the most robust formalism. In this sense, the cross-expression framework introduced here is primarily a way of conceptualizing gene-gene relations within spatial transcriptomics data, thereby serving as a powerful framework for research in tissue biology. For instance, it can be used to study

cancer [82], where tumor progresses via signaling with the stromal tissue, as well as

neurodegenerative diseases like Alzheimer's [83] or senescence [84], where the progression of

pathology is spatially structured, making it a broadly useful approach for numerous problems.

Cross-expression is not restricted to imaging-based spatial transcriptomics. Instead, it can

be applied to any biological assay that provides cell-by-features and cell-by-coordinate matrices.

For example, it can be extended to spatial proteomics [85], with potential to discover ligand-

receptor interactions. Likewise, it may be applied to spatial translatomics [86] to focus on

translating mRNAs that are more likely to form functional proteins, making conclusions about cell-

cell relations more robust. In fact, with the increasing resolution of spatially barcoded RNA capture

based methods [87,88], the framework may be extended transcriptome-wide to understand

relations between spots at near single-cell resolution.

A key challenge in imaging-based spatial transcriptomics [2–8], including the datasets

used in this work, is the size and constitution of the gene panels, which sets an upper limit on

biological discovery. Although our framework will become more powerful as the quality of spatial

transcriptomic data, especially the gene panel, increases, care must be taken to not interpret the

results in mechanistic terms. Instead, the coordinated gene expression between neighboring cells

should be viewed as a target for further investigation. In this sense, the cross-expression

framework substantially narrows the space of gene-gene relations by identifying pairs that are

potentially biologically meaningful, making the problem experimentally tractable. Overall, cross-

expression offers a unique and powerful perspective on using spatial transcriptomic data for

driving biological discovery.

Conclusions

Cross-expression is a useful conceptual and analytical framework, which compares all genes and

identifies pairs that coordinate their expression between neighboring cells. The accompanying R

software efficiently facilitates these and other analyses as well as provides new set of

visualizations to deeply explore spatial transcriptomics data by leveraging coordinated gene

expression at the single-cell resolution.

The package is available at https://github.com/gillislab/CrossExpression.

Extended Data Figures

Extended Data Fig. 1 | Cross-expression across tissue slices and regions for ligand-
receptor and non-signaling genes. a, Distribution of the number of gene pairs cross-expressed
in different slices. Dataset has 16 slices sampled sagittally from the left hemisphere of a mouse
brain. **b-e**, Cells are colored by gene expression (left) and cross-expressing cells are highlighted
(center). Right, distances between cross-expressing cells are compared with those between
cross-expressing and randomly selected cells. Smaller distances mean that cross-expressing
cells are nearer each other (spatial enrichment) than expected by chance (p-values ≤ 0.01, left-
tailed Mann-Whitney U test). Genes include ligands and receptors (b, c) and non-ligands and non-
receptors (d, e). **f**, Smoothed gene expression for different numbers of neighbors for the auditory
cortical layer 6 marker gene *Foxp2*. Created with BioRender.com.

Extended Data Fig. 2 | Relationship between cross-expression and cell type heterogeneity.
a, Proportion of cross-expressing cell pairs belonging to the same cell type label. **b**, Number of cell-neighbor pairs involved in cross-expression. **c**, Proportion of cell-neighbor pairs with different cell subtype labels given that both were labeled 'glutamatergic' at the higher level in the cell type hierarchy. **d**, Number of cell-neighbor pairs with the same or different cell subtype label given that both were labeled 'GABAergic' at the higher level in the cell type hierarchy. Each point is a cross-expressing gene pair. **e**, Same as in (c) but for 'GABAergic' cells. **f**, Proportion of neighbor cell types against which cell types cross-express. Created with BioRender.com.

Extended Data Fig. 3 | Combinatorial anatomical markers discovered using spatially
enriched cross-expression. a, Proportion of *Adra2b-* and *Ret*-expressing cells in the thalamus
(red) and the proportion of cell pairs in the thalamus (blue) when cross-expressing with *Lgr6*. **b-**
c, Distances between cross-expressing cells versus those between cross-expressing and
randomly chosen cells for genes *Lgr6* and *Adra2b* (c) and for *Lgr6* and *Ret* (d). Smaller distances
mean that cross-expressing cells are nearer each other (spatial enrichment) than expected by
chance (p-values ≤ 0.01, left-tailed Mann-Whitney U test). Created with BioRender.com.

Extended Data Fig. 4 | Exploration of *Gpr20* and its cross-expressing genes. a, Distribution of node degree, with *Gpr20* highlighted. **b**, Gene ontology (GO) functional groups for genes cross-expressed with *Gpr20*. **c**, Co-expression of genes cross-expressed with *Gpr20* (right) against cell type marker genes (bottom). For each gene, co-expression was computed using cells involved in cross-expression and not the entire dataset. **d**, Distribution of cell type marker genes' co-expression across the genes in (c). Created with BioRender.com.

Extended Data Fig. 5 | Exploration of the MERFISH cross-expression (sub)network. a, Co-expression of genes in the subnetwork. **b**, Co-expression between genes in the subnetwork (right) and cell type marker genes (bottom). Created with BioRender.com.

Extended Data Fig. 6 | Cross-expression network similarity between slices. a, Slice-specific cross-expression networks compared and shown as a function of distance between slices. **b**, Same as in (a) but slice-specific networks compared between sagittal and coronal datasets, where the "distance" is the difference in slice ID's. Shaded areas are 95% confidence intervals. Created with BioRender.com.

Extended Data Fig. 7 | Cross-expression network similarity between replicates. a-b, Cross-expression networks compared between three replicates for the middle (a) and posterior (b) slices. Created with BioRender.com.

Extended Data Fig. 8 | Cross-expression network similarity at different levels of gene expression noise thresholds. a, Cross-expression networks compared after applying different noise thresholds, which are the minimum number of molecules a gene must express within a cell to be considered as detected. **b**, Distribution of the network similarities across noise levels, with the median indicated using the dotted line. Created with BioRender.com.

Extended Data Fig. 9 | Patterns of cell-neighbor mappings and their relationship with cross-expression. a, Cells considered as 'nearest neighbor' by other cells reported as a proportion of total cell-neighbor relations. '1' is one-to-one mapping and '2-4' is many-to-one mapping. **b**, Cross-expression networks computed using one-to-one mappings, many-to-one mappings, and the full dataset (both mappings). Created with BioRender.com.

Online Methods

765 We first explain the theoretical underpinnings of our approach and outline the features of the

766 associated R package. We then specify how these are used in various analyses.

Statistics of cross-expression between a gene pair

769 Cross-expression is the mutually exclusive expression of a gene pair across neighboring cells. To

770 assess whether gene A's expression in cells and gene B's expression in their spatial neighbors is

771 significant, we calculate the probability using the hypergeometric approach

$$P(X = k) = \frac{\binom{K}{k}\binom{N-K}{n-k}}{\binom{N}{n}} \tag{1}$$

where N is the population size, K is the number of possible successes, n is the number of samples

or draws, k is the number of observed successes, and the form $\binom{a}{b}$ is the binomial coefficient.

Equation 1 outlines all the ways in which success can be observed—$\binom{K}{k}$—*and* (product

rule) all the ways in which failure can be obtained—$\binom{N-K}{n-k}$—normalized by all possible ways of

generating our sample $\binom{N}{n}$, making the outcome probabilistic by bounding it between   [0,1].

Traditionally, the *n* samples are assessed for the presence of some property *k*. Here, we sample

cell-neighbor *pairs* and ask whether the cell expresses gene A while the neighbor expresses gene

B. Thus, the sample size *n* is the number of pairs where the cells express gene A, the number of

observed successes *k* is the number of pairs where the cells express gene A and the neighbors

express gene B, and the number of success states *K* is the number of pairs where the neighbors

express gene B. The population size *N* is the total number of cell-neighbor pairs, including those

that co-express A and B and those that express neither gene. To calculate the probability of *k* or

more successes, we use the hypergeometric cumulative distribution function (CDF)

$$P(X \geq k) = 1 - P(X < k) = 1 - \sum_{i=0}^{k-1} \frac{\binom{K}{i}\binom{N-K}{n-i}}{\binom{N}{n}} \tag{2}$$

A value lower than alpha α indicates an unusually large number of pairs where the neighbors

express gene B and the cells express gene A, making their cross-expression significant.

Statistics of cross-expression between all gene pairs

We need to assess cross-expression across all gene pairs, which rise quadratically by $\binom{N}{2}$ or

$\frac{N(N-1)}{2}$ for *N* genes. To efficiently explore this space, we implement the procedure above using

matrix operations and specialized packages in R.

We begin with a cells-by-genes expression matrix **E** and a cells-by-coordinates location

matrix **L**, where the coordinates in our data are cell centroids on two-dimensional slices. We input

L into `RANN` package's function `nn2` with search type as `standard`, which implements a kd-tree

algorithm to explore data subspaces and efficiently finds the *n*-th neighbors [89,90]. Using the

neighbor indices, we re-order the expression matrix **E** to generate the neighbors-by-genes matrix

E′. The value of *n* can be changed to generate paired gene expression matrices, where the

corresponding rows represent cells and their *n*-th nearest neighbors. (To use a distance-based

approach, one can use the *n*-th neighbor insofar as the average distance to this neighbor is

smaller than the threshold.)

Our aim is to use **E** and **E′** to compute *N* (population), *K* (neighbors with B), *n* (cells with

A), and *k* (neighbors with B when their corresponding cells express gene A) for each gene pair.

These four values can be inputted into R's `phyper` function for all gene pairs to calculate

probabilities. The population size *N* is the total number of cells and is the same across all pairs.

To compute *n*, we binarize **E** based on if the genes are expressed, and compute co-occurrences

using the dot product

$$\mathbf{C} = \mathbf{E}^T \cdot \mathbf{E} \tag{3.1}$$

807   where $\mathbf{C}_{ii}$ is the number of cells expressing gene i and $\mathbf{C}_{ij}$ is the number of cells co-expressing

808   genes i and j. We perform

$$\mathbf{U}_{ij} = \mathbf{C}_{ii} - \mathbf{C}_{ij} \tag{3.2}$$

where $\mathbf{U}_{ij}$ is the number of cells uniquely expressing gene i. We implement this by extracting the

diagonal of **C**, and "broadcast" it against its off-diagonal entries, thus aligning the corresponding

values before subtraction. For each pair, this gives us the number of cells *n* uniquely expressing

each gene. We perform an analogous calculation for *K* using **E′** instead of **E**, giving us the number

of neighbors uniquely expressing each gene within a gene pair.

We now turn to *k*, the number of neighbors observed with gene B given that their

corresponding cells express gene A. Using binarized matrices **E** and **E′**, we compute the number

of cell-neighbor pairs such that the cells express gene A without gene B and the neighbors express

gene B without gene A

$$\mathbf{X} = (\mathbf{E} \odot (1 - \mathbf{E}')) \tag{4.1}$$

$$\mathbf{Y} = ((1 - \mathbf{E}) \odot \mathbf{E}') \tag{4.2}$$

$$\mathbf{Q} = \mathbf{X}^T \cdot \mathbf{Y} \qquad\qquad (5)$$

where $\odot$ is the Hadamard (elementwise) product and $\mathbf{Q}_{ij}$ is the number of cell-neighbor pairs with

mutually exclusive expression. In $\mathbf{X}$, $\mathbf{E}$ contains '1' in cells where a gene is expressed and $1 - \mathbf{E}'$

contains '1' in neighbors where a gene is *not* expressed. Their elementwise product $\mathbf{X}$ has '1' to

indicate genes' presence in cells and their absence in neighbors. $\mathbf{Y}$ shows the analogous

procedure for genes' presence in the neighbor and their absence in cells. Hence, the dot product

of $\mathbf{X}$ and $\mathbf{Y}$ gives $\mathbf{Q}$, a genes-by-genes asymmetric matrix, whose entries show the number of cell-

neighbor pairs with mutually exclusive expression. ($\mathbf{Q}$ is asymmetric because the number of cell-

neighbor pairs in the A-to-B and B-to-A directions are not always identical.) This is *k* or observed

successes. These steps generate four number – *N*, *K*, *n*, and *k* – per gene pair. We input these

into R's `phyper` function in accordance with equation (2), giving us corresponding p-values.

Since $\mathbf{Q}$ is asymmetric, we obtain two p-values per gene pair, one in the A-to-B and the

other in the B-to-A direction. We perform Benjamini-Hochberg false discovery rate (FDR)

correction on the entire p-value distribution [91]. For each gene pair, we then use the lower FDR-

corrected p-value as the final output $\mathbf{P}$ to indicate whether or not these genes cross-express.

Cross-expression networks

We can threshold and binarize $\mathbf{P}$ at a pre-selected alpha α to form an adjacency matrix $\mathbf{N}$, where

'1' indicates cross-expression (edges) between genes (nodes)

$$\mathbf{N}_{ij} = \begin{cases} 1 & \mathbf{P}_{ij} \leq \alpha \text{ and } i \neq j \\ 0 & \text{otherwise} \end{cases} \qquad\qquad (6.1)$$

836   This allows us to perform cross-expression network analysis, where higher-order

837   community structure is discovered using shared connections between genes

$$\mathbf{S}^R = \mathbf{N}^1 \cdot \mathbf{N}^2 \cdot \ldots \cdot \mathbf{N}^{R-1} \cdot \mathbf{N}^R \qquad\qquad (6.2)$$

838   where we restrict *R* = 2 to discover second-order connections between genes.

Cross-expression correlations

To compute correlations between genes, we first use the "masks" $\mathbf{X}$ and $\mathbf{Y}$ (eq. 4.1 and 4.2) and perform elementwise multiplication with non-binarized $\mathbf{E}$ and $\mathbf{E}'$, respectively. This makes the gene counts "0" if cell-neighbor pairs co-express the gene pair of interest. We then compute the correlation between all pairs of columns (genes) of $\mathbf{E}$ and $\mathbf{E}'$, yielding a gene-by-gene correlation matrix. We report the results as the average of this matrix with its transpose, giving a symmetric gene-gene correlation matrix as the output.

Cross-expression at multiple length scales

Cross-expression is coordinated gene expression between neighboring cells. Yet, these patterns may be present at larger length scales, requiring us to understand associations between regions. To facilitate this, we smooth the expression of each gene in a cell by averaging it with its expression in n nearby cells. Using the RANN package [89,90], we find the indices of each cell's n nearest neighbors, and make the corresponding values '1' in the cells-by-cells matrix $\mathbf{C}$

$$\mathbf{C}_{ij} = \sum_{k=1}^{s} \mathbb{I}\{(i,j) = (i_k, j_k)\} \tag{7.1}$$

where the indicator function $\mathbb{I}$ specifies

$$\begin{cases} 1 & \text{if } (i,j) = (i_k, j_k) \\ 0 & \text{otherwise} \end{cases}$$

and

$$s = c + (c \times n) \tag{7.2}$$

where c is the number of cells and n is the number of neighbors. Here, s is the total number of row-column indices i-j that k iterates over. We perform averaging using the expression matrix $\mathbf{E}$

$$\mathbf{S} = \frac{1}{n}(\mathbf{C} \cdot \mathbf{E}) \tag{8}$$

where $\mathbf{S}_{ij}$ is the j-th gene's average value in i-th cell across n neighbors. The smoothed gene expression matrix $\mathbf{S}$ can be used for downstream analysis.

861 **Bullseye scores as effect size**

862 The bullseye scores quantify the effect size by comparing cross-expression with co-expression.

863 Here, the number of neighbors with gene B is compared to the number of cells co-expressing

864 genes A and B. We use binarized cell and neighbor expression matrices $\mathbf{E}$ and $\mathbf{E'}$, respectively

$$\mathbf{B}_n = \mathbf{E}^T \cdot \mathbf{E}'_n \tag{9.1}$$

865 where n is the n-th neighbor, giving us n gene-by-gene asymmetric matrices $\mathbf{B}_n$. The i-th and j-th

866 entries of $\mathbf{B}_n$ indicate the number of n-th nearest neighbors expressing gene B when cells in $\mathbf{E}$

867 express gene A. $\mathbf{B}_n$ is a co-occurrence matrix when $n = 0$. Viewing $\mathbf{B}_n$ as a tensor with dimensions

868 i, j, and n, for each gene pair we take the cumulative sum and normalize across the neighbors

$$\mathbf{B}_{ijn} = \frac{1}{n} \sum_{n'=1}^{n} \mathbf{B}_{ijn'} \quad \text{for } n \geq 1 \tag{9.2}$$

These matrices can be compared with $\mathbf{B}_{n=0}$ to find the ratio of cross-expression to co-

expression and/or $\log_2$-transformed for further analysis. The output is provided as an array of

matrices (tensor) or as an edge list, where columns represent different n neighbors.

Expression of gene pairs on tissue

A powerful way of viewing cross-expression is to plot the cells and color them by their gene

expression. For a gene pair, a cell can express genes A, B, both, or neither. We make these plots

for user-selected gene pairs using the expression matrix $\mathbf{E}$ and the cell coordinates matrix $\mathbf{L}$. We

can also exclusively highlight cross-expressing cell-neighbor pairs. Finally, the tissue sections are

often not upright, partly due to their misorientation with respect to the glass slide, making it difficult

to interpret the results. Accordingly, we rotate them using user-defined n-degrees

$$\theta = n_{degrees} \times \frac{\pi}{180} \tag{10.1}$$

$$x' = \cos(\theta) \cdot x - \sin(\theta) \cdot y \tag{10.2}$$

$$y' = \sin(\theta) \cdot x + \cos(\theta) \cdot y \tag{10.3}$$

where *x′* and *y′* are the cell coordinates after counterclockwise rotation. Rotation does not change

the distances between cells, so *x′* and *y′* can be used for downstream analysis.

Spatial enrichment of cross-expression

Cross-expressing cells may be distributed across the tissue or show spatial localization. To

quantify their enrichment, we first average the distance between cell-neighbor pairs. We next

compare the distances between all cross-expressing cells to the distances between cross-

expressing and randomly selected cells. If the former distance is significantly smaller than the

latter distance, then cross-expression is spatially enriched.

Data acquisition and preprocessing

BARseq datasets

We collected a sagittal mouse brain data (P56 male) from the left hemisphere (20μm thick

sections, 300μm distance between slices) and measured 133 genes, including ligand-receptor

pairs (neuropeptides, neuropeptide receptors, monoamine receptors such as cholinergic,

adrenergic, serotonergic, and dopaminergic). This dataset includes 16 slices, which were assayed

as previously described [35]. We removed cells expressing fewer than 5 genes or with less than

20 counts.

We downloaded the BARseq coronal mouse brain dataset [35], which contains 9 brains,

with 4 control and binocular enucleation pairs as well as a pilot brain. These data contain 289

coronal slices and 109 genes. They were quality controlled just like the sagittal dataset.

10x Genomics Xenium datasets

We obtained three datasets from the Xenium platform https://www.10xgenomics.com/datasets.

The   first   dataset   contains   1   coronal   slice   with   5006   genes   obtained   from

https://www.10xgenomics.com/datasets/xenium-prime-fresh-frozen-mouse-brain. We removed

cells with fewer than 50 counts.

The second dataset contains 3 coronal age-matched AD-control FFPE brain pairs, where

the ages are 2.5 months, 5.7 months, and 13.4 (control) versus 17.9 months (AD). This data

includes 346 genes and was obtained from https://www.10xgenomics.com/datasets/xenium-in-

situ-analysis-of-alzheimers-disease-mouse-model-brain-coronal-sections-from-one-hemisphere-

over-a-time-course-1-standard. We removed cells with fewer than 50 counts.

The third dataset includes 4 coronal replicates with 248 genes. Three of these were

obtained from https://www.10xgenomics.com/datasets/fresh-frozen-mouse-brain-replicates-1-

standard and the remaining from https://www.10xgenomics.com/datasets/fresh-frozen-mouse-

brain-for-xenium-explorer-demo-1-standard. We removed cells with fewer than 30 counts.

EEL FISH dataset

We obtained a sagittal brain section data assaying 440 genes [92]. We removed cells with fewer

than 15 counts.

NanoString CosMx dataset

We obtained a coronal brain section data from https://nanostring.com/products/cosmx-spatial-

molecular-imager/ffpe-dataset/cosmx-smi-mouse-brain-ffpe-dataset/. This dataset assays 950

genes. We removed cells with fewer than 100 counts.

Resolve Biosciences Molecular Cartography dataset

We downloaded one hemi-coronal slice data from https://resolvebiosciences.com/open-

dataset/?dataset=mouse-brain-2021, which assayed 99 genes. We removed cells with fewer than

30 counts.

Vizgen MERFISH brain receptor map datasets

We obtained Vizgen MERSCOPE's mouse brain receptor map from

https://info.vizgen.com/mouse-brain-data. This data contains three coronal slices from three

replicates, with the middle slice covering the center of the brain. It assays 483 genes. We filtered

cells with fewer than 50 counts and those lacking brain region annotations (see below). We

registered the slices to the Allen CCFv3 (Common Coordinate Framework version 3) brain region

atlas [61]. To facilitate this, we annotated the cells using Seurat [93]. Here, we created a Seurat

object and used `SCTransform` with the `clip.range` between -10 and 10. We then ran Principal

Component Analysis (PCA), setting the number of components to 30 and specifying the features

as genes. Next, we used `FindNeighbors` and `FindClusters` with the resolution set to 0.3.

The clusters are cell type labels, which help us identify brain structures during registration. For

registration, we used QuickNii (v3 2017) [94] to linearly align the slice to the Allen CCFv3 atlas

using discernible regions like the hippocampus and the ventricles as anchors. We then used

VisuAlign (v0.9) [94] to non-linearly transform the slice to improve alignment with the atlas. This

procedure assigns a brain region annotation to every cell.

STARmap dataset

The STARmap dataset contains 20 coronal, sagittal, or spinal sections, where 1022 genes were

assayed [34]. We removed cells with fewer than 30 counts.

Allen Institute MERSCOPE atlas dataset

We obtained MERSCOPE dataset from the Allen Institute brain atlas, which contains 500 genes

measured across 59 coronal slices [32]. We removed cells with fewer than 15 unique genes or

lower than 40 counts.

MERFISH brain atlas datasets

The first dataset assays 1122 genes across 239 coronal and sagittal slices [33]. We removed cells

in the top and bottom 1% by total counts.

The second dataset assays 374 genes across age-matched (4 weeks, 24 weeks, 90

weeks) controls and LPS-infected coronal brains, with 49 total slices [95]. (LPS is

lipopolysaccharide bacterial infection.) We removed cells expressing fewer than 5 genes or with

less than 20 counts.

Single-cell RNA-seq (scRNA-seq) and single-nucleus RNA-seq (snRNA-seq) datasets

We obtained the scRNA-seq [32] and snRNA-seq [77] data from the Brain Initiative Cell Census

Network (BICAN) cell type atlases. These data were collected from dissected tissue regions,

giving us the cells' coarse anatomical origin. We removed cells with a doublet score of 30 or above

and randomly selected 10,000 cells from each region for subsequent analysis.

Ligand-receptor cross-expression

We aimed to find cross-expression between known ligand-receptor pairs. In our sagittal data, we

selected two slices and within each slice we chose a cortical region. These choices were made

randomly. In practice, we chose the visceral area (VISC) in slice 3 and the somatosensory nose

region (SSp-n) in slice 5. Next, we selected the well-known neuropeptide somatostatin *Sst* and

its cognate receptor *Sstr2* as the candidate pair. Finding their cross-expression significant, we

show their expression on tissue and highlight cross-expressing cells. We also compute their

bullseye scores and report them as a ratio of cross- to co-expression across 10 neighbors.

Cross-expression and cell type heterogeneity

We explore the relationship between cross-expression and cell type heterogeneity using the

BARseq coronal data [35]. First, we use *Gfra1* and *Foxp2* to highlight cross-expressing cells and

map different cell types to distinct shapes. Second, we count the number of cross-expressing cell-

neighbor pairs for numerous genes. Since each cell has a cell type label, we compute cell type

purity as the proportion of pairs with the same label. Third, we use the cell type hierarchy to assess

if cell-neighbor pairs with the same H1 label have the same H3 label. We first find cross-

expressing gene pairs using the entire dataset. Next, using cell pairs with the 'glutamatergic' H1

label, we compute the number of pairs with the same or different H3 labels. We perform a similar

analysis using cells labelled as 'GABAergic' at the H1 level. Finally, we compute the frequencies

with which cell type label combinations are associated between neighboring cells and normalize

this by the expected frequencies of those cell type pairs in the population.

Replicability of cross-expression

We evaluate the replicability of our data using the hypergeometric and correlation-based

approaches, which provide independent ways of quantifying cross-expression. We predict the

binary p-values (significant or not, alpha = 0.05) using correlations and report the performance as

AUROC. For performance within datasets, we predict the labels of a held-out slice using the

average correlation of the remaining slices using a leave-one-out cross-validation procedure, with

the results reported as AUROC against the number of slices/ samples used. For performance

between datasets, we predict the labels of slices in one study using correlations of slices from

another study and report the performance as median AUROC between study pairs. This process

requires using gene pairs shared between studies' panels.

1002

Creation and annotation of meta-analytic cross-expression network

Our cross-expression meta-analytic network included connections (edges) between genes

(nodes) if they were cross-expressed in two or more studies. We used the `igraph` package [96]

to create a network and Leiden clustered it [97], with "modularity" as the objective function ran for

100 iterations. Since Leiden clustering is somewhat stochastic, we performed this process 100

times, giving us 100 distinct partitions/ community assignments per gene. To find consensus

communities, we calculated the adjusted Rand index (ARI) [98], which tells us the degree to which

two clustering results agree with each other, between all pairs of partitions. Next, we found the

average ARI for each partition and choose the one with the highest ARI, which gives us the

'consensus' communities. Importantly, we observed that multiple partitions had the same highest

average ARI, and upon further inspection, we discovered that these partitions were identical,

giving credence to our final community assignments.

After assigning genes to communities, we performed gene ontology (GO) [99,100]

enrichment per community using the union of the studies' gene panels as the background. We

merged significant GO terms together using Revigo [101], which gave us the biological processes

(BP), cellular components (CC), and molecular functions (MF) of each community. We performed

minor manual curation of these terms to generate our final GO annotations. The meta-analytic

network was visualized using Cytoscape (v3.10.1) [102,103].

1021

Networks of cross-expression when studying *Gpr20*

Using the MERFISH data, we computed cross-expression p-values between all genes and

binarized the matrix at $\alpha \leq 0.05$ to create an adjacency matrix. We calculate the node degree as

the number of edges formed by each gene and create a network with second-order edges (shared

connections) as outlined in equation 6.2. We set the threshold for second-order edges to 4,

meaning that two genes are connected if they share at least 4 first-order edges, ensuring that the

higher-order network is robust. Next, we use the `igraph` package [96] to perform Louvain

clustering [104] (with default parameters) on the second-order network and thus assign genes to

communities.

We visualize the network using Cytoscape (v3.10.1) [102,103], mapping node size to

degree, color to node community, and edge color to edge type (first-order, second-order, or both).

We use the "organic" layout and apply "remove overlaps" from the `yFiles` app [103] and tweak

the network to further reduce overlaps. Finally, we use the `Legend Creator` app [103] to render

a legend with node degree size and community assignment.

Because our network revealed *Gpr20* as topologically salient, we performed gene

ontology (GO) [99,100] enrichment analysis on genes that cross-expressed with it ('test set').

Here, we used the entire gene panel (except *Gpr20*) as the background set and used the

hypergeometric test to determine if it significantly overlapped with the test set, giving us p-values

for each GO functional group. We report FDR-corrected p-values. Additionally, for each gene

cross-expressed with *Gpr20*, we used the cells involved in cross-expression, rather than the entire

dataset, to compute co-expression with cell type marker genes and compared these global

profiles between marker types.

Since the cells expressing *Gpr20* visually showed spatial autocorrelation, we assessed

their neighbors as well as randomly chosen cells for the expression of *Gpr20*. We L1-normalized

the counts for both groups, rendering them into probability distributions, and computed cumulative

sums. To calculate the area under curve (AUC), we scaled the neighbor order between 0 and 1

and used the `trapz` function from R's `pracma` package to calculate the AUC [105].

Within the main network, we introduce a further constraint that cross-expressing genes

must lack significant co-expression. We curate the subnetwork by removing genes with node

degree of 1 and assign cell type labels based on genes' co-expression with marker genes. Like

before, we perform GO enrichment using the subnetwork genes as the test set and the gene

panel as the background set, and report FDR-corrected p-values.

To assess whether cross-expression networks are more similar between adjacent slices

than between distant slices, we compute slice-specific cross-expression networks and calculate

Spearman's correlation between these networks. The correlations are plotted against distances

between slices, where the "distance" is the difference in the slice order. As a control, we compute

the Spearman's correlations between slice-specific networks obtained from different brains and

plot this against the "distance" between the slice ID's.

Discovering combinatorial anatomical marker genes

1060

We observed that cross-expression discovers anatomical marker genes that delineate the

thalamus. To quantitatively assess this, we made a mask by combining the following regions:

anterior group of the dorsal thalamus (ATN), intralaminar nuclei of the dorsal thalamus (ILM),

lateral group of the dorsal thalamus (LAT), medial group of the dorsal thalamus (MED), midline

group of the dorsal thalamus (MTN), ventral group of the dorsal thalamus (VENT), and ventral

posterior complex of the thalamus (VP). Importantly, we compared every brain region annotation

in our data with Allen CCFv3 atlas [61] and judged the ones presented here to best mark the

thalamic regions. This allowed us to calculate the number of cells expressing each gene within or

outside the thalamus. For cross-expressing cells, we considered a pair as thalamic if both cells

were part of the regional mask. More generally, potential combinatorial markers can be discovered

by assessing if their cross-expression is spatially enriched.

Our second exploration involved well-known genes *Foxp2* and *Cdh13*, which mark cortical

layer 6 and show pan-layer expression in the cortex, respectively. These genes exhibited

significant cross-expression, which was spatially enriched in layer 6, whose boundaries we

identified using H2 cell type annotation. The spatial enrichment was viewed by comparing tissue

plots with and without highlighting cross-expressing cells.

1077

Robustness of cross-expression to batch effects

1078

To assess the replicability of the cross-expression signature, we used the MERFISH dataset

containing 3 biological replicates (mouse brains) with 3 slices each, where the slices are sampled

from approximately the same location across the brains. We compared the slice-specific networks

between corresponding slices. Moreover, for slice-specific and brain region-specific networks, we

performed comparisons within the BARseq sagittal data and within the coronal data [35] as well

as between these two datasets. Finally, observing that the dorsal to ventral direction is sampled

in both the coronal and the sagittal brains, we compared the densities of cross-expressing cells

in the dorsal to ventral directions across these datasets.

1087

Cell segmentation quality control assessment

We assessed the quality of cell segmentation at a global level by comparing co-expression

between the scRNA-seq [32] and MERFISH spatial transcriptomic data. Since the scRNA-seq

was obtained from dissected brain regions, we established correspondence between these and

the brain region annotations in the MERFISH data. The regions used in both data are reported in

Supplementary Table 1. We included only those genes – and in the same order – as present in

the MERFISH data. We calculated gene co-expression using Pearson's correlation and compared

these across the two datasets.

To quantify variability between platforms, we compared gene co-expression between

scRNA-seq and snRNA-seq [77] for the same genes as above. Because the snRNA-seq was

obtained from dissected brain regions, we established correspondence between these and the

scRNA-seq data. The regions used in these data are reported in Supplementary Table 2. Like

before, we quantified co-expression using Pearson's correlation and compared it across the two

datasets.

1102

Gene expression noise thresholds and cell-neighbor relations

Because gene expression measurement is noisy, we applied thresholds of 1 to 10 molecules,

thus specifying the minimum number of counts per cell a gene must have to be considered

expressed. We then compared cross-expression networks across these thresholds.

Additionally, a cell might be the nearest neighbor of one or more cells. To ensure that our

framework captures this variability, we compare cross-expression networks for the one-to-one

and many-to-one mappings with each other and with that of the full dataset.

1110

Benchmarking the algorithm's speed

We assessed the speed of the cross-expression algorithm by duplicating our BARseq coronal data, where the gene panel ranged from 2,000 to 8,000 and the number of cells ranged from 20,000 to 200,000. We ran the cross-expression algorithm and calculated the time on a 16 GB Apple M1 Pro macOS Sonoma 14.5 laptop.

Declarations

Ethics approval and consent to participate

The experimental work in this study involved the collection of a mouse sagittal brain dataset using BARseq. All experimental procedures were carried out in accordance with the Institutional Animal Care and Use Committee at the Allen Institute for Brain Science.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used in this study are available in the repositories or articles outlined in the "data acquisition and preprocessing" section. The BARseq sagittal data's sequencing images are being deposited to Brain Image Library (BIL). While it is being approved, we stored the cell-level gene expression and cell metadata at

https://drive.google.com/drive/folders/1fk5JDeVJcE71iH1AaICT0il9PN9DTJJm?usp=drive_link.

Competing interests

L.F. owns shares in Quince Therapeutics and has received consulting fees from PeopleBio Co., GC Therapeutics Inc., Cortexyme Inc., and Keystone Bio. The remaining authors declare no competing interests.

Funding

A.S. received funding from University of Toronto's FAST Doctoral Fellowship, Ontario Graduate Scholarship, and NSERC's Canada Graduate Scholarship. X.C. was supported by grants R01MH133181 and DP2MH132940. L.F. received support from R01MH133181 and J.G. from both R01MH133181 and R01MH113005. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Authors' contributions

J.G. conceived the project. A.S. conducted the analyses, developed the software, and wrote the manuscript under supervision from J.G. M.R and H.C. collected the sagittal brain data under supervision from X.C. L.F. managed, curated, and parsed the datasets. S.C. created the R package. All authors interpreted the results and edited the manuscript.

Acknowledgements

Not applicable.

Authors' information (optional)

Not applicable.

Code availability

The R package is available at https://github.com/gillislab/CrossExpression/, including installation guide via Bioconductor and a tutorial with an example expression matrix and metadata.

References

1. Marx V. Method of the Year: spatially resolved transcriptomics. Nat Methods. 2021;18:9–14.

2. Zhuang X. Spatially resolved single-cell genomics and transcriptomics by imaging. Nat
Methods. 2021;18:18–22.

3. Moffitt JR, Lundberg E, Heyn H. The emerging landscape of spatial profiling technologies. Nat
Rev Genet. 2022;23:741–59.

4. Tian L, Chen F, Macosko EZ. The expanding vistas of spatial transcriptomics. Nat Biotechnol.
2023;41:773–82.

5. Bressan D, Battistoni G, Hannon GJ. The dawn of spatial omics. Science.
2023;381:eabq4964.

6. Liu L, Chen A, Li Y, Mulder J, Heyn H, Xu X. Spatiotemporal omics for biology and medicine.
Cell. 2024;187:4488–519.

7. Moses L, Pachter L. Museum of spatial transcriptomics. Nat Methods. 2022;19:534–46.

8. Ren J, Luo S, Shi H, Wang X. Spatial omics advances for in situ RNA biology. Mol Cell.
2024;84:3737–57.

9. Ma Y, Zhou X. Accurate and efficient integrative reference-informed spatial domain detection
for spatial transcriptomics. Nat Methods. 2024;21:1231–44.

10. Kim J, Rustam S, Mosquera JM, Randell SH, Shaykhiev R, Rendeiro AF, et al.
Unsupervised discovery of tissue architecture in multiplexed imaging. Nat Methods.
2022;19:1653–61.

11. Chidester B, Zhou T, Alam S, Ma J. SpiceMix enables integrative single-cell spatial modeling
of cell identity. Nat Genet. 2023;55:78–88.

12. Birk S, Bonafonte-Pardàs I, Feriz AM, Boxall A, Agirre E, Memi F, et al. Quantitative
characterization of cell niches in spatially resolved omics data. Nat Genet. 2025;57:897–909.

13. Jerby-Arnon L, Regev A. DIALOGUE maps multicellular programs in tissue from single-cell
or spatial transcriptomics data. Nat Biotechnol. 2022;40:1467–77.

14. Haviv D, Remšík J, Gatie M, Snopkowski C, Takizawa M, Pereira N, et al. The covariance
environment defines cellular niches for spatial inference. Nat Biotechnol. 2024;1–12.

15. Sun S, Zhu J, Zhou X. Statistical analysis of spatial expression patterns for spatially
resolved transcriptomic studies. Nat Methods. 2020;17:193–200.

16. Ghazanfar S, Lin Y, Su X, Lin DM, Patrick E, Han Z-G, et al. Investigating higher-order
interactions in single-cell data with scHOT. Nat Methods. 2020;17:799–806.

17. Bernstein MN, Ni Z, Prasad A, Brown J, Mohanty C, Stewart R, et al. SpatialCorr identifies
gene sets with spatially varying correlation structure. Cell Rep Methods [Internet]. 2022 [cited
2024 Aug 19];2. Available from: https://www.cell.com/cell-reports-methods/abstract/S2667-
2375(22)00271-5

1196 18. Miller BF, Bambah-Mukku D, Dulac C, Zhuang X, Fan J. Characterizing spatial gene
1197 expression heterogeneity in spatially resolved single-cell transcriptomic data with nonuniform
1198 cellular densities. Genome Res. 2021;31:1843–55.

1199 19. Li Z, Wang T, Liu P, Huang Y. SpatialDM for rapid identification of spatially co-expressed
1200 ligand–receptor and revealing cell–cell communication patterns. Nat Commun. 2023;14:3995.

1201 20. Armingol E, Baghdassarian HM, Lewis NE. The diversification of methods for studying cell–
1202 cell interactions and communication. Nat Rev Genet. 2024;25:381–400.

1203 21. Almet AA, Tsai Y-C, Watanabe M, Nie Q. Inferring pattern-driving intercellular flows from
1204 single-cell and spatial transcriptomics. Nat Methods. 2024;1–12.

1205 22. Yang W, Wang P, Xu S, Wang T, Luo M, Cai Y, et al. Deciphering cell–cell communication at
1206 single-cell resolution for spatial transcriptomics with subgraph-based graph attention network.
1207 Nat Commun. 2024;15:7101.

1208 23. Cang Z, Zhao Y, Almet AA, Stabell A, Ramos R, Plikus MV, et al. Screening cell–cell
1209 communication in spatial transcriptomics via collective optimal transport. Nat Methods.
1210 2023;20:218–28.

1211 24. Dries R, Zhu Q, Dong R, Eng C-HL, Li H, Liu K, et al. Giotto: a toolbox for integrative
1212 analysis and visualization of spatial expression data. Genome Biol. 2021;22:78.

1213 25. Troulé K, Petryszak R, Cakir B, Cranley J, Harasty A, Prete M, et al. CellPhoneDB v5:
1214 inferring cell–cell communication from single-cell multiomics data. Nat Protoc. 2025;1–29.

1215 26. Liu J, Manabe H, Qian W, Wang Y, Gu Y, Chu AKY, et al. CytoSignal Detects Locations and
1216 Dynamics of Ligand-Receptor Signaling at Cellular Resolution from Spatial Transcriptomic Data
1217 [Internet]. bioRxiv; 2024 [cited 2024 Aug 19]. p. 2024.03.08.584153. Available from:
1218 https://www.biorxiv.org/content/10.1101/2024.03.08.584153v1

1219 27. Jin S, Plikus MV, Nie Q. CellChat for systematic analysis of cell–cell communication from
1220 single-cell transcriptomics. Nat Protoc. 2025;20:180–219.

1221 28. Yuan Y, Bar-Joseph Z. GCNG: graph convolutional networks for inferring gene interaction
1222 from spatial transcriptomics data. Genome Biol. 2020;21:300.

1223 29. Singhal V, Chou N, Lee J, Yue Y, Liu J, Chock WK, et al. BANKSY unifies cell typing and
1224 tissue domain segmentation for scalable spatial omics data analysis. Nat Genet. 2024;56:431–
1225 41.

1226 30. Hu J, Li X, Coleman K, Schroeder A, Ma N, Irwin DJ, et al. SpaGCN: Integrating gene
1227 expression, spatial location and histology to identify spatial domains and spatially variable
1228 genes by graph convolutional network. Nat Methods. 2021;18:1342–51.

1229 31. Zubair A, Chapple RH, Natarajan S, Wright WC, Pan M, Lee H-M, et al. Cell type
1230 identification in spatial transcriptomics data can be improved by leveraging cell-type-informative
1231 paired tissue images using a Bayesian probabilistic model. Nucleic Acids Res. 2022;50:e80.

32. Yao Z, van Velthoven CTJ, Kunst M, Zhang M, McMillen D, Lee C, et al. A high-resolution
transcriptomic and spatial atlas of cell types in the whole mouse brain. Nature. 2023;624:317–
32.

33. Zhang M, Pan X, Jung W, Halpern AR, Eichhorn SW, Lei Z, et al. Molecularly defined and
spatially resolved cell atlas of the whole mouse brain. Nature. 2023;624:343–54.

34. Shi H, He Y, Zhou Y, Huang J, Maher K, Wang B, et al. Spatial atlas of the mouse central
nervous system at molecular resolution. Nature. 2023;622:552–61.

35. Chen X, Fischer S, Rue MCP, Zhang A, Mukherjee D, Kanold PO, et al. Whole-cortex in situ
sequencing reveals input-dependent area identity. Nature. 2024;1–10.

36. Wu X-B, Zhu Q, Gao M-H, Yan S-X, Gu P-Y, Zhang P-F, et al. Excitatory Projections from
the Prefrontal Cortex to Nucleus Accumbens Core D1-MSNs and κ Opioid Receptor Modulate
Itch-Related Scratching Behaviors. J Neurosci. 2023;43:1334–47.

37. Xiao P, Yan W, Gou L, Zhong Y-N, Kong L, Wu C, et al. Ligand recognition and allosteric
regulation of DRD1-Gs signaling complexes. Cell. 2021;184:943-956.e18.

38. Szklarczyk K, Korostynski M, Cieslak PE, Wawrczak-Bargiela A, Przewlocki R. Opioid-
dependent regulation of high and low fear responses in two inbred mouse strains. Behav Brain
Res. 2015;292:95–101.

39. Minett MS, Pereira V, Sikandar S, Matsuyama A, Lolignier S, Kanellopoulos AH, et al.
Endogenous opioids contribute to insensitivity to pain in humans and mice lacking sodium
channel Nav1.7. Nat Commun. 2015;6:8967.

40. Wawrczak-Bargieła A, Ziółkowska B, Piotrowska A, Starnowska-Sokół J, Rojewska E, Mika
J, et al. Neuropathic Pain Dysregulates Gene Expression of the Forebrain Opioid and Dopamine
Systems. Neurotox Res. 2020;37:800–14.

41. Beaulieu J-M, Gainetdinov RR. The Physiology, Signaling, and Pharmacology of Dopamine
Receptors. Sibley DR, editor. Pharmacol Rev. 2011;63:182–217.

42. Fabricius D, Karacay B, Shutt D, Leverich W, Schafer B, Takle E, et al. Characterization of
Intestinal and Pancreatic Dysfunction in VPAC1-Null Mutant Mouse. Pancreas. 2011;40:861.

43. Khaleduzzaman M, Sumiyoshi H, Ueki Y, Inoguchi K, Ninomiya Y, Yoshioka H. Structure of
the Human Type XIX Collagen (COL19A1) Gene, Which Suggests It Has Arisen from an
Ancestor Gene of the FACIT Family. Genomics. 1997;45:304–12.

44. Calvo AC, Moreno L, Moreno L, Toivonen JM, Manzano R, Molina N, et al. Type XIX
collagen: a promising biomarker from the basement membranes. Neural Regen Res.
2020;15:988.

45. Sticco MJ, Peña Palomino PA, Lukacsovich D, Thompson BL, Földy C, Ressl S, et al.
C1QL3 promotes cell-cell adhesion by mediating complex formation between ADGRB3/BAI3
and neuronal pentraxins. FASEB J. 2021;35:e21194.

46. Martinelli DC, Chew KS, Rohlmann A, Lum MY, Ressl S, Hattar S, et al. Expression of C1ql3
in Discrete Neuronal Populations Controls Efferent Synapse Numbers and Diverse Behaviors.
Neuron. 2016;91:1034–51.

47. Van Itallie CM, Tietgens AJ, Aponte A, Gucek M, Cartagena-Rivera AX, Chadwick RS, et al.
MARCKS-related protein regulates cytoskeletal organization at cell–cell and cell–substrate
contacts in epithelial cells. J Cell Sci. 2018;131:jcs210237.

48. Kondrychyn I, Kelly DJ, Carretero NT, Nomori A, Kato K, Chong J, et al. Marcksl1 modulates
endothelial cell mechanoresponse to haemodynamic forces to control blood vessel shape and
size. Nat Commun. 2020;11:5476.

49. Shestakova EA, Singer RH, Condeelis J. The physiological significance of β-actin mRNA
localization in determining cell polarity and directional motility. Proc Natl Acad Sci.
2001;98:7045–50.

50. Crow M, Gillis J. Co-expression in Single-Cell Analysis: Saving Grace or Original Sin?
Trends Genet. 2018;34:823–31.

51. Xu M, Bai X, Ai B, Zhang G, Song C, Zhao J, et al. TF-Marker: a comprehensive manually
curated database for transcription factors and related markers in specific cell and tissue types in
human. Nucleic Acids Res. 2022;50:D402–12.

52. Jackson RJ, Hyman BT, Serrano-Pozo A. Multifaceted roles of APOE in Alzheimer disease.
Nat Rev Neurol. 2024;20:457–74.

53. Jones-Tabah J, Mohammad H, Paulus EG, Clarke PBS, Hébert TE. The Signaling and
Pharmacology of the Dopamine D1 Receptor. Front Cell Neurosci [Internet]. 2022 [cited 2025
Apr 17];15. Available from:
https://www.frontiersin.orghttps://www.frontiersin.org/journals/cellular-
neuroscience/articles/10.3389/fncel.2021.806618/full

54. Jones-Tabah J, Mohammad H, Hadj-Youssef S, Kim LEH, Martin RD, Benaliouad F, et al.
Dopamine D1 receptor signalling in dyskinetic Parkinsonian rats revealed by fiber photometry
using FRET-based biosensors. Sci Rep. 2020;10:14426.

55. Sun H, Monenschein H, Schiffer HH, Reichard HA, Kikuchi S, Hopkins M, et al. First-Time
Disclosure of CVN424, a Potent and Selective GPR6 Inverse Agonist for the Treatment of
Parkinson's Disease: Discovery, Pharmacological Validation, and Identification of a Clinical
Candidate. J Med Chem. 2021;64:9875–90.

56. Brice NL, Schiffer HH, Monenschein H, Mulligan VJ, Page K, Powell J, et al. Development of
CVN424: A Selective and Novel GPR6 Inverse Agonist Effective in Models of Parkinson
Disease. J Pharmacol Exp Ther. 2021;377:407–16.

57. Barekatain M, Johansson LC, Lam JH, Chang H, Sadybekov AV, Han GW, et al. Structural
insights into the high basal activity and inverse agonism of the orphan receptor GPR6 implicated
in Parkinson's disease. Sci Signal. 2024;17:eado8741.

58. Margolin DH, Brice NL, Davidson AM, Matthews KL, Carlton MBL. A Phase I, First-in-
Human, Healthy Volunteer Study to Investigate the Safety, Tolerability, and Pharmacokinetics of

CVN424, a Novel G Protein-Coupled Receptor 6 Inverse Agonist for Parkinson's Disease. J Pharmacol Exp Ther. 2022;381:33–41.

59. Kieburtz K, Brice N, Carlton M, Thompson C, Dubow J. CVN424, A novel GPR6 inverse agonist for the treatment of Parkinson disease (PD): Phase 2 clinical trial. Parkinsonism Relat Disord [Internet]. 2024 [cited 2025 Apr 20];122. Available from: https://www.prd-journal.com/article/S1353-8020%2824%2900136-6/fulltext?utm_source=chatgpt.com

60. Brice NL, Carlton M, Margolin DH, Bexon M, Matthews KL, Dawson LA, et al. CVN424, a GPR6 inverse agonist, for Parkinson's disease and motor fluctuations: a double-blind, randomized, phase 2 trial. eClinicalMedicine [Internet]. 2024 [cited 2025 Apr 17];77. Available from: https://www.thelancet.com/journals/eclinm/article/PIIS2589-5370%2824%2900461-9/fulltext?utm_source=chatgpt.com

61. The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas. Cell. 2020;181:936-953.e20.

62. Heng N, Malek N, Lawton MA, Nodehi A, Pitz V, Grosset KA, et al. Striatal Dopamine Loss in Early Parkinson's Disease: Systematic Review and Novel Analysis of Dopamine Transporter Imaging. Mov Disord Clin Pract. 2023;10:539–46.

63. van Vliet EF, Knol MJ, Schiffelers RM, Caiazzo M, Fens MHAM. Levodopa-loaded nanoparticles for the treatment of Parkinson's disease. J Controlled Release. 2023;360:212–24.

64. Bogetofte H, Alamyar A, Blaabjerg M, Meyer M. Levodopa Therapy for Parkinson's Disease: History, Current Status and Perspectives. CNS Neurol Disord - Drug Targets Former Curr Drug Targets - CNS Neurol Disord. 2020;19:572–83.

65. Delafontaine P, Song Y-H, Li Y. Expression, Regulation, and Function of IGF-1, IGF-1R, and IGF-1 Binding Proteins in Blood Vessels. Arterioscler Thromb Vasc Biol. 2004;24:435–44.

66. Dallinga MG, Yetkin-Arik B, Kayser RP, Vogels IMC, Nowak-Sliwinska P, Griffioen AW, et al. IGF2 and IGF1R identified as novel tip cell genes in primary microvascular endothelial cell monolayers. Angiogenesis. 2018;21:823–36.

67. Imrie H, Viswambharan H, Sukumar P, Abbas A, Cubbon RM, Yuldasheva N, et al. Novel Role of the IGF-1 Receptor in Endothelial Function and Repair: Studies in Endothelium-Targeted IGF-1 Receptor Transgenic Mice. Diabetes. 2012;61:2359–68.

68. Liang W, Zheng Y, Zhang J, Sun X. Multiscale modeling reveals angiogenesis-induced drug resistance in brain tumors and predicts a synergistic drug combination targeting EGFR and VEGFR pathways. BMC Bioinformatics. 2019;20:203.

69. Le X, Nilsson M, Goldman J, Reck M, Nakagawa K, Kato T, et al. Dual EGFR-VEGF Pathway Inhibition: A Promising Strategy for Patients With *EGFR*-Mutant NSCLC. J Thorac Oncol. 2021;16:205–15.

70. Milner CS, Hansen TM, Singh H, Brindle NPJ. Roles of the receptor tyrosine kinases Tie1 and Tie2 in mediating the effects of angiopoietin-1 on endothelial permeability and apoptosis. Microvasc Res. 2009;77:187–91.

71. Niu Q, Perruzi C, Voskas D, Lawler J, Dumont D, Benjamin LE. Inhibition of Tie-2 Signaling
Induces Endothelial Cell Apoptosis, Decreases Akt Signaling and Induces Endothelial Cell
Expression of the Endogenous Anti-Angiogenic Molecule, Thrombospondin-1. Cancer Biol Ther.
2004;3:402–5.

72. Savant S, La Porta S, Budnik A, Busch K, Hu J, Tisch N, et al. The Orphan Receptor Tie1
Controls Angiogenesis and Vascular Remodeling by Differentially Regulating Tie2 in Tip and
Stalk Cells. Cell Rep. 2015;12:1761–73.

73. Zhang Y, Xu D, Huang H, Chen S, Wang L, Zhu L, et al. Regulation of Glucose Homeostasis
and Lipid Metabolism by PPP1R3G-mediated Hepatic Glycogenesis. Mol Endocrinol.
2014;28:116–26.

74. Aaron PA, Jamklang M, Uhrig JP, Gelli A. The blood–brain barrier internalises via the
EphA2-tyrosine kinase receptor. Cell Microbiol. 2018;20:e12811.

75. Phillips RA, Oh S, Bach SV, Du Y, Miller RA, Kleinman JE, et al. Transcriptomic
characterization of human lateral septum neurons reveals conserved and divergent marker
genes across species. iScience. 2025;28:111820.

76. Chi Y, Qi R, Zhou Y, Tong H, Jin H, Turck CW, et al. scBrainMap: a landscape for cell types
and associated genetic markers in the brain. Database. 2023;2023:baad035.

77. Langlieb J, Sachdev NS, Balderrama KS, Nadaf NM, Raj M, Murray E, et al. The molecular
cytoarchitecture of the adult mouse brain. Nature. 2023;624:333–42.

78. Stringer C, Wang T, Michaelos M, Pachitariu M. Cellpose: a generalist algorithm for cellular
segmentation. Nat Methods. 2021;18:100–6.

79. Fu X, Lin Y, Lin DM, Mechtersheimer D, Wang C, Ameen F, et al. BIDCell: Biologically-
informed self-supervised learning for segmentation of subcellular spatial transcriptomics data.
Nat Commun. 2024;15:509.

80. Ma J, Xie R, Ayyadhury S, Ge C, Gupta A, Gupta R, et al. The multimodality cell
segmentation challenge: toward universal solutions. Nat Methods. 2024;21:1103–13.

81. Choudhary S, Satija R. Comparison and evaluation of statistical error models for scRNA-
seq. Genome Biol. 2022;23:27.

82. Yu Q, Jiang M, Wu L. Spatial transcriptomics technology in cancer research. Front Oncol
[Internet]. 2022 [cited 2024 Aug 21];12. Available from:
https://www.frontiersin.org/journals/oncology/articles/10.3389/fonc.2022.1019111/full

83. Chen W-T, Lu A, Craessaerts K, Pavie B, Frigerio CS, Corthout N, et al. Spatial
Transcriptomics and In Situ Sequencing to Study Alzheimer's Disease. Cell. 2020;182:976-
991.e19.

84. Gurkar AU, Gerencser AA, Mora AL, Nelson AC, Zhang AR, Lagnado AB, et al. Spatial
mapping of cellular senescence: emerging challenges and opportunities. Nat Aging.
2023;3:776–90.

85. Lundberg E, Borner GHH. Spatial proteomics: a powerful discovery tool for cell biology. Nat
Rev Mol Cell Biol. 2019;20:285–302.

86. Zeng H, Huang J, Ren J, Wang CK, Tang Z, Zhou H, et al. Spatially resolved single-cell
translatomics at molecular resolution. Science. 2023;380:eadd3067.

87. Russell AJC, Weir JA, Nadaf NM, Shabet M, Kumar V, Kambhampati S, et al. Slide-tags
enables single-nucleus barcoding for multimodal spatial genomics. Nature. 2024;625:101–9.

88. Hu C, Borji M, Marrero GJ, Kumar V, Weir JA, Kammula SV, et al. Scalable spatial
transcriptomics through computational array reconstruction. Nat Biotechnol. 2025;1–7.

89. Bentley JL. Multidimensional binary search trees used for associative searching. Commun
ACM. 1975;18:509–17.

90. Arya S, Mount DM, Netanyahu NS, Silverman R, Wu AY. An optimal algorithm for
approximate nearest neighbor searching fixed dimensions. J ACM. 1998;45:891–923.

91. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. J R Stat Soc Ser B Methodol. 1995;57:289–300.

92. Borm LE, Mossi Albiach A, Mannens CCA, Janusauskas J, Özgün C, Fernández-García D,
et al. Scalable in situ single-cell profiling by electrophoretic capture of mRNA using EEL FISH.
Nat Biotechnol. 2023;41:222–31.

93. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al. Dictionary
learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol.
2024;42:293–304.

94. Puchades MA, Csucs G, Ledergerber D, Leergaard TB, Bjaalie JG. Spatial registration of
serial microscopic brain images to three-dimensional reference atlases with the QuickNII tool.
PLOS ONE. 2019;14:e0216796.

95. Allen WE, Blosser TR, Sullivan ZA, Dulac C, Zhuang X. Molecular and spatial signatures of
mouse brain aging at single-cell resolution. Cell. 2023;186:194-208.e18.

96. Csárdi G, Nepusz T, Müller K, Horvát S, Traag V, Zanini F, et al. igraph for R: R interface of
the igraph library for graph theory and network analysis [Internet]. Zenodo; 2024 [cited 2024 Aug
21]. Available from: https://zenodo.org/records/10681749

97. Traag VA, Waltman L, van Eck NJ. From Louvain to Leiden: guaranteeing well-connected
communities. Sci Rep. 2019;9:5233.

98. Hubert L, Arabie P. Comparing partitions. J Classif. 1985;2:193–218.

99. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene Ontology: tool
for the unification of biology. Nat Genet. 2000;25:25–9.

100. The Gene Ontology Consortium, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin
HJ, et al. The Gene Ontology knowledgebase in 2023. Genetics. 2023;224:iyad031.

101. Supek F, Bošnjak M, Škunca N, Šmuc T. REVIGO Summarizes and Visualizes Long Lists
of Gene Ontology Terms. PLOS ONE. 2011;6:e21800.

102. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: A
Software Environment for Integrated Models of Biomolecular Interaction Networks. Genome
Res. 2003;13:2498–504.

103. Morris JH, Kuchinsky A, Ferrin TE, Pico AR. enhancedGraphics: a Cytoscape app for
enhanced node graphics [Internet]. F1000Research; 2014 [cited 2024 Aug 21]. Available from:
https://f1000research.com/articles/3-147

104. Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E. Fast unfolding of communities in
large networks. J Stat Mech Theory Exp. 2008;2008:P10008.

105. Borchers HW. pracma: Practical Numerical Math Functions [Internet]. 2023 [cited 2025 Apr
20]. Available from: https://cran.r-project.org/web/packages/pracma/index.html

1429