Contrasting Cognitive, Behavioral, and Physiological Responses to Breathwork vs. Naturalistic Stimuli in Reflective Chamber and VR Headset Environments: Randomized Control Study

Ninette Simonian[1], Micah-Alan Johnson, PhD[1,2], Caitlin Lynch[1], Geena Wang[1], Velu Kumaravel[1], Taylor Kuhn, PhD, MS[3], Félix Schoeller, PhD[1], Nicco Reggente, PhD[1]

[1] Institute for Advanced Consciousness Studies, Santa Monica, CA, USA
[2] Department of Biostatistics, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA
[3] Department of Psychiatry and Behavioral Sciences, Semel Institute of Neuroscience and Human Behavior, University of California, Los Angeles, Los Angeles, CA, USA

Corresponding Author:
Nicco Reggente, PhD
Institute for Advanced Consciousness Studies
2811 Wilshire Blvd.
Santa Monica, CA 90403
United States
Email: nicco@advancedconsciousness.org

Abstract

Background: Virtual reality (VR) and immersive technologies offer mental health benefits, including anxiety reduction, mood enhancement, and cognitive function improvement. Previous research has demonstrated the anxiolytic effects of breathwork and exposure to natural environments, leading to enhanced well-being.

Objective: This study examines the effectiveness of rain stimuli and guided breathwork delivered through two distinct systems: MindGym, a reflective chamber, and a traditional VR headset.

Methods: Physiological measures, cognitive, and trait and state assessments were collected for 126 participants, randomly assigned to VR (Breathwork), VR (Rain), MindGym (Breathwork), or MindGym (Rain) conditions.

Results: Significant improvements were observed across all groups in cognitive performance, anxiety, and mood. The Trail Making Test (RTACC) decreased (ΔM=-30.4, W=5443.00, Z=4.32, $P.FDR$<.001, RBC=.45), Architex Total Speed improved (ΔM=-76.4, W=6044.00, Z=7.82, $P.FDR$<.001, RBC=.84), and anxiety (STAI) reduced (ΔM=-3.78, W=5156.50, Z=4.86,

P.FDR<.001, *RBC*=.52). Breathwork conditions showed a greater decrease in breath rate compared to Rain conditions (*F*(2.48, 208.49)=28.49, *P.FDR*=.002). Individual differences moderated treatment responses, with MODTAS predicting Awe (*b*=0.37, *SE*=0.11, *Z*=3.28, *P.FDR*=.004) and Ego Dissolution (*b*=0.03, *SE*=0.01, *Z*=2.98, *P.FDR*=.015). Openness interacted with stimuli type on anxiety reduction (*F*(1, 111)=9.58, *P.FDR*=.038), with higher openness associated with greater anxiety reduction in Breathwork conditions. No significant differences were found between MindGym and VR in effectiveness or immersion.

Conclusions: MindGym and VR demonstrated promise as acute anxiolytics, with MindGym-generated content maintaining effectiveness when translated to VR. This highlights MindGym's potential as a versatile content creation platform for immersive, anxiety-reducing experiences. Individual differences moderated treatment responses, suggesting opportunities for personalized interventions. Future research should explore complex MindGym experiences and adaptation to various delivery systems for accessible, effective anxiety management tools.

KEYWORDS
Immersive technology; Virtual reality; Presence; Immersion; Anxiolytic intervention; Guided breathwork; Naturalistic stimuli

Introduction

Experience shapes human cognition and behavior, yet its variability limits prescriptive outcomes. Experiential technology, leveraging audiovisual stimuli, offers potential for targeted interventions with predictable effects on mood and cognition [1–4]. Immersive virtual reality (VR) experiences can elicit specific emotional responses (e.g., awe) and enhance well-being through increased felt connectedness [5].

Anxiety disorders, affecting 19.1% of adults globally, significantly impact healthcare systems and individual well-being, with even subclinical levels diminishing quality of life [6,7][8]. First-line treatments like cognitive-behavioral therapy and medication offer benefits but face limitations such as side effects and relapse potential [9,10]. Alternative approaches (mindfulness, exercise, pharmacotherapies) show variable efficacy [6,10,11], while emerging interventions (vagal nerve stimulation, cannabidiol) require further safety and efficacy research [11,12]. This landscape necessitates innovative approaches to anxiety management.

Two promising avenues for acute anxiety reduction are exposure to naturalistic stimuli and guided breathwork. Nature exposure and related therapies have been shown to facilitate stress recovery, mood enhancement, improved cognitive functions, and overall well-being while alleviating anxiety symptoms and boosting health and self-efficacy [13–21]. Breathwork techniques demonstrate comparable efficacy through distinct mechanisms, inducing parasympathetic states that promote stress reduction and enhance emotional regulation [22–28].

Despite their potential, these approaches face accessibility challenges, particularly in urban environments with limited access to natural settings [29] and due to the lack of integration of breathwork and somatic therapies into standard mental health care [11]. Immersive technologies offer a solution by creating scalable, accessible virtual experiences, enabling well-controlled,

"plug-and-play" interventions with optimized effectiveness and expanded reach. Innovative platforms for generating immersive content, such as the MindGym, hold promise for creating highly effective experiences that could potentially be adapted to various delivery systems, including more portable VR formats. Research shows combined audiovisual nature scenes offer enhanced psychological benefits over single-modality presentations [1,3], aligning with Attention Restoration and Stress Reduction theories [13,30]. 360-degree nature videos demonstrate rapid mood improvements [21,31–36], while natural sounds enhance physiological recovery [37] and cognitive performance [38].

The efficacy of these virtual interventions hinges on immersion—the technological capability to create a sense of "being there"—and presence, the subjective experience of feeling transported to the virtual environment. These factors are crucial for eliciting powerful psychological and physiological responses [39,40]. Crucially, research shows the cognitive benefits of VR depend on users truly feeling "there," emphasizing the fundamental role of high immersion and presence [41]. Both technological capabilities and subjective experiences like perceived realism and enjoyment influence these factors [42–44]. Immersion levels vary across VR systems, from moderate levels in consumer-grade headsets to heightened experiences in advanced setups [45,46]. These findings underscore the potential of immersive technology to create targeted, accessible interventions that leverage nature's restorative properties.

The current study compared the effectiveness of two immersive technologies in delivering anxiolytic content: a "reflective chamber" (MindGym) and a traditional VR headset. Both incorporated naturalistic rain sounds or guided 4-7-8 breathwork [24,47,48]. We sought to investigate not only the relative efficacy of these delivery systems but also the potential for translating experiences designed for the MindGym to more widely accessible VR platforms. We hypothesized that MindGym, designed for enhanced immersion and presence, would outperform VR by amplifying nature and breathwork effects while minimizing VR-associated discomfort [49–52]. This comparison sought to inform the development of novel, scalable anxiety-reducing interventions, bridging the gap between traditional therapies and the need for accessible mental health solutions.

Methods

Participants

A total of 126 participants (63 females; age range 18-74 years; mean 41, SD 15.19) participated in the study. Participants were randomly assigned to one of four groups, as shown in Table 1.

Table 1. Participant characteristics by group and analysis type

Group	N	Gender (M/F)	Age Range	M	SD
VR (Breathwork)	33	16/17	20-71	42.61	15.18
VR (Rain)	29	16/13	18-69	41.30	15.54
MindGym (Breathwork)	31	15/16	22-74	38.45	15.93
MindGym (Rain)	32	16/16	19-68	41.53	14.50

F = female; M = male. Age is reported in years. M and SD represent the mean and standard deviation of age, respectively.

10 participants were excluded from the physiological analysis due to equipment failure or excessive movement resulting in a total of 116 participants for statistical analysis (61 females; age range 18-74 years; mean 40.54, SD 14.56). Respiration was analyzed for 88 participants, ECG was analyzed for 80 participants, and EDA was analyzed for 96 participants (see Table 2). All participants passed quality assurance checks, with respiration rates below 20 breaths per minute following data cleaning and processing (see Methods for further explanation).

Table 2. Participant characteristics for physiological measures by group

Measure	Group	N	Gender (F/M)	Age Range	M	SD
Respiration						
	VR (Breathwork)	22	12/10	20-65	39.91	14.36
	VR (Rain)	19	8/11	18-69	43.15	16.56
	MindGym (Breathwork)	28	14/14	23-74	36.96	15.17
	MindGym (Rain)	19	11/8	19-63	41.89	14.11
ECG						
	VR (Breathwork)	18	9/9	20-65	41.17	14.31
	VR (Rain)	16	9/7	18-68	41.44	17.00
	MindGym (Breathwork)	26	15/11	23-74	37.04	15.84
	MindGym (Rain)	20	12/8	19-68	41.80	13.33
EDA						
	VR (Breathwork)	23	14/9	20-65	41.61	13.26
	VR (Rain)	22	9/13	18-69	40.77	15.72
	MindGym (Breathwork)	28	14/14	23-68	36.82	14.65
	MindGym (Rain)	23	14/9	20-63	41.43	13.03

F = female; M = male. Age is reported in years. M and SD represent the mean and standard deviation of age, respectively. ECG = Electrocardiogram; EDA = Electrodermal Activity.

Recruitment

Participants were recruited for this study through newsletters sent to individuals with prior engagement in our previous research studies and targeted advertising on Facebook and Instagram, aimed at adults residing within a 50-mile radius of Santa Monica, California. Participants were compensated $30 per hour via cash or Venmo payments, and parking fees were validated. The total participation duration, measured from arrival to departure, was rounded up to the nearest 15-minute increment for compensation purposes.

Eligibility

Inclusion criteria for study participation were as follows: (1) absence of neurological conditions including epilepsy and migraine; (2) no history of claustrophobia; (3) absence of photosensitivity or ophthalmological conditions such as cataracts, corneal abrasions, keratitis, or uveitis; (4) normal or corrected-to-normal auditory and visual acuity; (5) no current use of medications known to induce photophobia or alter auditory perception; and (6) sufficient mobility to enter the MindGym apparatus without assistive devices (e.g., wheelchairs, walkers, or canes).

Prior to enrollment, potential participants underwent screening via a standardized online questionnaire administered through the Google Forms platform. This pre-screening process ensured adherence to all inclusion and exclusion criteria before formal study initiation.

Materials

All participants sat on an OMEGA Gaming Chair (SecretLab, Inc.) and completed the behavioral questionnaires (administered through Google Forms) and cognitive tasks on a 27" 2022 iMac (Apple, Inc.) regardless of group.

Physiological Monitoring

The CGX Aim II Physiological Monitoring device (CGX, Inc.), using ECG electrodes (Skintact Inc.), measured the following bio peripherals: Electromyography (EMG; 2 electrodes on the L/R base of the neck on the sternocleidomastoid muscle), Bio-Impedance-Based Respiration Rate (2 paddles with 2 electrodes each on the L/R pectoralis major), Heart Rate, and Galvanic Skin Response (GSR; 2 electrodes on the palm of the non-dominant hand).

Lab Recorder

Lab Streaming Layer with LabRecorder [6,10,11] was utilized to temporally synchronize our peripheral and experimental time series (experience started and experience ended) within an XDF file format.

Behavioral Questionnaires and Tasks

Trait Measures

a. Dispositional Positive Emotion Scale (D-PES;[53]) – The Awe Subscale of D-PES measures an individual's dispositional propensity to feel awe toward the world, consisting of 6 items.
b. Modified Tellegen Absorption Scale (MODTAS; [54]) – Adapted from the Tellegen Absorption Scale (Tellegen & Atkinson, 1974), uses a Likert scale to assess the imaginative involvement and the tendency to become mentally absorbed in everyday activities.
c. NEO Five-Factor Inventory (NEO-FFI-3; [55]) – Measures five broad personality dimensions – neuroticism, extraversion, openness to experience, agreeableness, and conscientiousness.

State Measures

a. State-Trait Anxiety Inventory (STAI;[56]) – Assesses state anxiety (temporary/situational) and trait anxiety (enduring propensity).
b. Profile of Mood States 2nd Edition (POMS 2; [57]) – Evaluates current mood state affective dimensions like anger, confusion, depression, fatigue, tension, and vigor.

c. Valence and Arousal – Measures emotional state on two Likert scales from 0 to 10: Valence (0= very unpleasant,10 = very pleasant) and Arousal (0 = totally calm, 10 = totally excited/agitated).

Outcome Measures

a. Toronto Mindfulness Scale (TMS; [58]) – A 13-item administered post-meditation to assess state mindfulness, while differentiating between reflective awareness and ruminative attention. The TMS contains two subscales: Curiosity, gauging the interest in one's experiences, and Decentering, reflecting the ability to view thoughts and feelings as transient mental events.
b. Awe Scale (Awe-S; [59]) – An 8-item self-report scale measuring individual differences in the tendency to experience awe, using a 7-point Likert scale to rate their agreement with statements regarding feelings of awe and perceptions of vastness.
c. Body Size Estimation (BSE; [60]) – Assesses body image perception, from a selection of nine progressive figural drawings that depict varying body sizes to estimate their own perceived body size, both in height and width. This measure has been linked to the intensity of awe and immersion in non-ordinary experiences [61].
d. Ego-Dissolution Inventory (EDI; [62]) – Consists of 16 items assessing altered states of ego-consciousness, using a visual analog scale ranging from 0% to 100%. It includes 8 items relating to ego-dissolution, representing the core of the inventory, and 8 to ego-inflation, contrasting experience of heightened self-assuredness.
e. Igroup Presence Questionnaire (IPQ; [63]) – Measures sense of presence, immersion, engagement, and connection in virtual environments.
f. Modified Immersion Experience Questionnaire (adapted per [64]) – Measures depth and quality of immersion/engagement in a particular context or virtual environment.
g. Motion Sickness Assessment Questionnaire [65] – A 16-item scale assessing multiple dimensions of motion sickness (gastrointestinal, central, peripheral, and sopite-related symptoms).

Behavioral Tasks

a. Trail-Making Test (TMT; [66]) – A neuropsychological assessment, taken on the computer, evaluating visual attention, scanning, processing speed, and cognitive flexibility. Participants were instructed to connect a series of 25 dots in sequential order, alternating between numbers and letters (1-A-2-B, and so on), aiming for speed and accuracy.
b. Architex [67] – A validated, computerized neuropsychological assessment tool designed to objectively measure multiple cognitive domains and their interactions, including attention, working memory, processing speed, affect recognition, planning, organization, impulsivity, and verbal and nonverbal abstract reasoning.

Content Delivery Systems

The first immersive content-delivery system was MetaQuest 2 (Meta Platforms, Inc.) VR headset, equipped with an Elite Strap (see Figure 1), presenting stimuli through its native video

player at a resolution of 5376x2688, scaled to the maximum allowed resolution of 1920x3664, and a refresh rate of 120Hz. All videos were captured using a 360 camera (GoPro Max 360) placed inside the MindGym during the programmed experience, with the audio content subsequently overlaid. During the experiment, the VR is set to cast through an iPad Air (iPadOS 16.6.1) and navigated using a wired mouse (Razer Inc.).

The second system was the MindGym (Lumena, Inc.), a 7' isotropic MindGym lined with reflective mylar on its interior walls, while its floor and ceiling are equipped with mirrors. (see Supplemental Information). The MindGym incorporates WS2815 LEDs, with 121 pixels per edge between the vertices, totaling 1452 pixels across 12 edges. It utilizes SMD5050 RGB LED chip, offering RGB color with 256 grayscale levels and an output of 990-1080 lumens per meter. The LEDs are operated at less than half of their maximum capacity. Additionally, the system features a color temperature of 5500K. Participants in the MindGym group were provided over-ear wireless, noise-canceling headphones (Sony Group Corp.) to wear during the experience (see Figure 1).

Figure 1   **Figure 2**

Figure 1: Participant in VR setup; **Figure 2:** Participant in MindGym setup

Stimuli

The first piece of content was "4-8 Breathwork" (see Figure 3 and Figure 5), led by Dr. Jannell MacAulay's guided audio and utilizing two primary lighting setups. The beginning featured the "Mindful Minute" mindfulness exercise with multiple colors and slowly upward-moving LEDs, designed to encourage mind wandering among participants. MacAulay then introduced the breathwork technique and its underlying rationale, aided by "Galaxy" visuals of blue LEDS, around 6000K, embedded in the walls, giving the sensation of being surrounded by stars. The guided breathwork audio was complemented by upward and downward cascades of vertically aligned LEDs, synchronized with the inhalation and exhalation periods (see Supplemental Information).

The second piece of content was "Rain" (see Figure 4 and Figure 6), an immersive environment created for reflection. The audio begins with complete darkness as the sound of rain starts to trickle in. This is complemented by a downward cascade of vertically aligned blue LEDs, around 6500K, against a black background, simulating rain visuals falling from each vertex. The participant remains immersed in the light and sound environment for the remainder of the reflective experience.

Figure 3   Figure 4

Figure 5   Figure 6

Figure 3. MindGym (Breathwork); **Figure 4.** MindGym (Rain); **Figure 5.** VR (Breathwork); **Figure 6.** VR (Rain)

Procedure

Overview

Before each session, all participants provided signed consent forms. They began the session by filling out pre-experience questionnaires and completing cognitive tasks, followed by CGX setup. Subsequently, each group underwent a 10-minute experiential phase, after which they completed post-experience questionnaires and tasks (see Figure 7).

Figure 7. Procedure overview. Total participation time was roughly 80 minutes.

Randomization

Subjects were iteratively assigned to one of two groups using a MATLAB-based randomization algorithm that maintained gender balance, with recruitment ceasing for each gender category upon reaching the predetermined quota. The experimenter remained blind to group assignment until launching the Python script, which initiated the group-appropriate content delivery system and stimuli by entering the participant's ID number. Participants were randomly assigned to one of four content delivery systems and stimuli pairings: (1) MindGym (Breathwork), (2) MindGym (Rain), (3) VR (Breathwork), and (4) VR (Rain).

Procedure

Upon arrival, participants were seated on an office chair facing a 35.5" x 55" desk and given an overview of the session (pre-questionnaires, equipment setup, 10-minute experience, post-questionnaires). They were informed their "experience" would involve sitting for 10 minutes in an immersive audiovisual environment.

For approximately the first 30 minutes, participants used the iMac to complete a set of questionnaires (NEO-FFI-3, D-PES, MODTAS, STAI, POMS-2, Valence, and Arousal) on Google Forms, followed by a brief neurophysiological assessment via the TMT and Architex. Participants were instructed to silence their phones and remove jewelry and bulky items for comfort, before being told which experiential group they were assigned to. Using a cotton pad and 91% isopropyl alcohol, the experimenter cleansed the relevant areas (see CGX materials above) before outfitting the participant with biosensors and peripherals.

Participants assigned to VR sat in the same room, with the room shades lowered. The VR headset was positioned and adjusted with the tension dial on the Elite strap for clear vision and comfort. The experimenter cast the VR to the iPad and adjusted the settings, such that the audio

was played through the headset. Upon connecting to Lab Recorder, the stimuli are launched concurrently with the initiation of the LSL trigger, by the Python script.

If assigned to the MindGym, participants entered the MindGym and sat centrally, while holding the CGX device and wearing headphones. The Lab Recorder was connected, and the experience began within 15 seconds of closing the MindGym door.

Across groups, participants were told to sit still and relax during the 10-minute experience without falling asleep. Participants were shown how to remove the VR or exit the MindGym if they decided to end the experience early, otherwise, the experimenter would return to assist them once the experience ended.

After the experience, biosensors were removed and participants used the restroom if needed before completing post-experience behavioral assessments (STAI, POMS-2, Valence, Arousal, IPQ, MSAQ, TMS, Awe-S, BSE, and EDI) and cognitive tasks (TMT and Architex). Finally, participants were compensated and dismissed.

Behavioral Statistical Analysis

Dependent Variables

Seven primary outcome measures collected at both timepoints were used as dependent variables (DV) in separate models. There were four mood state measures: Anxiety (STAI), Arousal, Valence, and Total Mood Disturbance (POMS). There were also three cognitive measures: two that tracked performance on the Architex task (Total Speed and Total Score) and one that tracked performance on the TMT using the ratio of reaction time divided by accuracy, here called TMT RTACC, which decreases with increasing performance due to increased accuracy and/or decreased reaction time).

Eleven DVs were collected at the Post timepoint only, which included Awe, BSE, EDI, Immersion (based on IEQ), overall Motion, Curiosity and Decentering (both from Toronto), and four IPQ variables (General, Involvement, Spatial Presence, Experienced Realism).

There were also four physiological variables – Breath Rate, EDA, HR, and HRV – that were collected throughout the duration of each experience. These measures were averaged or computed in either two 5-minute timepoints (for HRV), five 2-minute timepoints (for EDA and Breath Rate), or ten 1-minute timepoints (for HR).

Independent Variables

The primary independent variables (IV) were Time or Timepoint (Pre, Post) as a categorical, within-subjects factor, for any variables collected at both timepoints, as well as both Tech (MindGym, VR) and Stim (Breathwork, Rain) as categorical, between-subjects group factors. Several continuous or ordinal IVs were tested for potential moderation or mediation of the Tech

and Stim effects. There were eight psychological variables: DPES, Immersion, IPQ (General), IPQ (Involvement), IPQ (Spatial Presence), IPQ (Experienced Realism), MODTAS, and Openness. There were seven physiological variables: Motion (Overall), initial breath rate (Breath Rate, first 2 min), breath rate change (Breath Rate change, last 2 min – first 2 min), initial heart rate (HR, first 1 min), heart rate change (HR change, last 1 min – first 1 min), initial heart rate variability (HRV, first 5 min), and change in heart rate variability (HRV change, last 5 min – first 5 min). The demographic variables of age and sex were also included in all moderation models to control for their potential effects.

Nonparametric tests

For all DVs collected at both timepoints (pre, post), changes over time were tested with nonparametric t-tests using Wilcoxon signed-rank tests in JASP [68]. Effect sizes were reported as the matched rank biserial correlation (RBC), an appropriate measure for nonparametric t-tests which ranges between –1 to 1 and indicates the relative proportion of positive and negative rank scores while ignoring ties (e.g., RBC = 0.60 means that 60% more subjects showed an increase instead of a decrease)**.** Timepoint tests were performed separately for each of the four experimental groups (MindGym (Breathwork), MindGym (Rain), VR (Breathwork), VR (Rain)) as well as the combined MindGym and VR groups, the combined Breathwork and Rain groups, and all groups combined.

For all DVs, differences between groups were tested with nonparametric Mann-Whitney U tests in JASP, which included RBC effect sizes. Six separate group comparisons were tested: MindGym (Breathwork) vs MindGym (Rain), VR (Breathwork) vs VR (Rain), MindGym (Breathwork) vs VR (Breathwork), MindGym (Rain) vs VR (Rain), MindGym vs VR, and Rain vs Breathwork. For any DV collected at both timepoints, the difference score (Post – Pre) was used to compare group differences in changes over time. For the physiological measures, the first timepoint, last timepoint, and the difference between first and last timepoints were separately used as DVs to compare group differences in initial, final, or change in physiological response.

Moderation

Moderation of the experimental effects on mood or cognition was tested as interactions between each covariate and the Tech and Stim group factors in separate models for the three-way interaction (Tech x Stim x Moderator) and two-way interactions (Tech x Moderator, Stim x Moderator), as well as a separate model with only main effects. Demographic covariates (age and sex) were also included as main effects in all models. For any outcome measure collected at both timepoints, the difference score (Post – Pre) was used as the DV. Because we did not have any a priori hypotheses about differential moderation effects between nested levels of Tech and Stim, we conducted post-hoc t-tests for only the interaction terms that were deemed significant.

Moderations were analyzed with generalized linear models (GLM), using the *glmmTMB* [69] and *emtrends* [70] packages in RStudio [71]. Based on extensive model diagnostics, which were

conducted with the *DHARMa* [72] package, it was determined that either gaussian or t-family distributions (both with the identity link) were optimal for model fitting to ensure no substantial violations or problems with linearity, independence of errors, homoscedasticity, dispersion, zero-inflation, outliers, or within-group normality of residuals.

Separate models were tested for the hypothesized moderators, which included eight psychological variables (DPES, Immersion, IPQ General, IPQ Involvement, IPQ Spatial Presence, IPQ Experienced Realism, MODTAS, and Openness) and seven physiological variables (overall motion, initial Breath Rate, Breath Rate change, initial HR, HR change, initial HRV, HRV change). All moderation models were tested separately for 10 outcome variables as DVs: Arousal, Anxiety (STAI), Awe, BSE, EDI, Total Mood Disturbance (POMS), Valence, Architex (Total Score), Architex (Total Speed), and TMT (RTACC).

Repeated measures ANOVA

For the physiological measures collected during the experiences, repeated measures ANOVAs (rmANOVA) were performed in JASP to assess changes over time as a main effect of Timepoint (two timepoints for HRV, five timepoints for Breath Rate and EDA, ten timepoints for HR) and to assess experimental effects on changes over time, with separate three-way (Tech x Stim x Timepoint) or two-way interactions (Tech x Timepoint, Stim x Timepoint). Levene's test was used to ensure no violations of homogeneity. The assumption of sphericity was tested (using Mauchly's test) and corrected (with Greenhouse-Geisser) if necessary.

Multiple Comparison Correction

We used the false discovery rate (FDR) method of correcting *P* values for multiple comparisons (significance threshold chosen as *a*=.05, *P*<.05) to control Type I and Type II errors [73,74]. FDR correction was performed across DVs and separately for 1) the nonparametric tests of timepoint effects for each group (single or combined), 2) the nonparametric tests of group differences, separately for each group comparison and separately for the measures collected at both timepoints, the measures collected at only the Post timepoint, and the physiological measures collected during the experience, as well as 3) the moderation tests, separately for each interaction or main effect. FDR correction of the rmANOVAs was performed across the different models testing interaction or main effects and separately for each physiological measure.

Mediation

Mediation analysis was conducted in JASP based on the *lavaan* [75] package in R. Missing values were excluded listwise. Separate models were tested for each potential mediator, which included the same variables that were used as potential moderators in the moderation analysis, and for each DV that was also used in the moderation analysis.

Neurophysiological Analysis

Data Acquisition and Preprocessing

Physiological data, including electrocardiogram (ECG), electrodermal activity (EDA), electromyography (EMG), and respiration, were recorded using the CGX AIM Phys. Mon. AIM-0106 device. The data were stored in XDF format, which is a standardized format for multi-channel time series data [76]. The XDF files were loaded using the pyxdf library, and the relevant data streams were extracted based on their names and labels.

ECG Processing

The ECG data were first sanitized to remove any invalid or missing data points [77]. A notch filter was then applied to remove powerline interference at 60 Hz. The ECG signal was further cleaned by applying a bandpass filter with a low cut-off frequency of 0.5 Hz and a high cut-off frequency of 45 Hz using a 4^{th}-order Butterworth filter [78]. The filtered ECG data were then filtered again to remove any values exceeding a predefined amplitude threshold of 1500 mV.

Baseline wandering, a low-frequency artifact in the ECG signal, was computed by measuring the peak-to-peak amplitude of the ECG signal. If the baseline wandering exceeded a predefined threshold of 1700 mV, the ECG data were considered unacceptable for further analysis.

EDA Processing

The EDA data were processed in conjunction with the EMG data to identify and remove motion artifacts. First, the EMG signal was normalized. A sliding window approach was then used to identify windows of EMG data that contained a high proportion of outliers, which were considered to be indicative of motion artifacts. The Local Outlier Factor (LOF) algorithm [79] was used to detect outliers within each window (implemented using the scikit-learn library) [80].

The EDA signal was smoothed using a median filter with a kernel size of 8 seconds. A low-pass Butterworth filter with a cut-off frequency of 5 Hz was then applied to remove high-frequency noise. The contaminated windows identified in the EMG signal were used to guide the interpolation of the EDA signal. Linear interpolation was performed to replace the artifact-contaminated segments of the EDA signal with values interpolated from the neighboring clean segments.

The quality of the cleaned EDA signal was assessed by checking for values outside the acceptable range (0.05 - 60 μS) and for rapid changes exceeding ±10 μS/sec within a 1-second interval [81]. The proportion of data points violating these criteria was used to compute a quality index ranging from 0 to 100, with higher values indicating better signal quality. If the quality index was below 50, the EDA data were considered unacceptable for further analysis.

Respiration Processing

The respiration data were preprocessed following the methods of [82]. We first applied a high-pass Butterworth filter with a cut-off frequency of 0.1 Hz to remove low-frequency drift. The filtered signal was then resampled to a fixed sampling rate of 100 Hz. A sliding window approach was used to estimate the respiration period within each window using the Fast Fourier Transform (FFT). The estimated respiration period was used to guide the application of a moving average filter to smooth the respiration signal within each window. The smoothed signal was then resampled back to the original sampling rate.

Inhalation and exhalation onsets were detected by identifying the intercepts between the smoothed respiration signal and its moving mean. The intercepts were then classified as inhalation or exhalation onsets based on the direction of the slope at each intercept. A minimum amplitude threshold of 0.25 was used to filter out small fluctuations that do not correspond to true respiratory events.

The number of breaths within the first and last 5 minutes of the experiment was computed using the inhalation and exhalation onsets to define individual breaths.

Heart Rate Variability Analysis

Heart rate variability (HRV) analysis was performed by computing the low-frequency (LF) and high-frequency (HF) components of the ECG signal using the Welch's method for power spectral density estimation [83]. The LF component was defined as the power in the frequency range of 0.04-0.15 Hz, while the HF component was defined as the power in the range of 0.15-0.4 Hz. The ratio of LF to HF power (LF/HF ratio) was computed as a measure of sympathovagal balance [84]. The LF/HF ratio was computed separately for the first and last 5 minutes of the experiment.

EDA Tonic Component Analysis

The tonic component of the EDA signal, which reflects the slow-varying level of skin conductance, was extracted by applying a low-pass filter to the EDA signal [77]. The mean value of the tonic component was computed within non-overlapping 2-minute intervals. These mean values were used as a measure of the overall level of arousal throughout the experiment.

Ethics

IRB

The Advarra (Columbia, MD) Institutional Review Board approved all recruitment and testing procedures prior to initiating enrollment (Pro00070581). Following ethical standards and legal requirements, all study participants provided informed consent using DropboxSign. They were afforded ample time to seek clarification from the Principal Investigator and study personnel.

The informed consent process incorporated the California Experimental Research Bill of Rights as mandated by Health and Safety Code Section 24172. Additionally, the study adhered to the principles outlined in the Declaration of Helsinki. Furthermore, all laboratory personnel possessed and maintained up-to-date certifications in Good Clinical Practice and the Protection of Human Research Participants through online training.

Results

Outcome Measures as Function of Group

No statistically significant differences were observed between the group comparisons for the post-intervention measures (all $P.FDR>.05$) (see Supplemental Information: Appendix C). The closest result to reaching statistical significance with a notable effect size was found for IPQ (Involvement) when comparing Breathwork to Rain ($W=2501.50$, $P.FDR=.12$, $RBC=.26$), suggesting a potential trend towards higher involvement in the Breathwork condition compared to the Rain condition.

Pre-Post Changes as a Function of Group

Timepoint effects within group

Analysis of pre-post changes revealed significant improvements across various measures when considering all groups combined and within specific conditions (see Table 3 and Supplemental Pages: Appendix A for detailed results).

Table 3. Summary of results from the nonparametric (Wilcoxon Signed-Rank) t-tests of timepoint (pre vs post) differences for the different groups or group combinations.

DV	MindGym (Breathwork)			MindGym (Rain)			VR (Breathwork)			VR (Rain)		
	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig
TMT (RTACC)	2.68	0.45	~	-42.64	0.66	***	-43.21	0.33		-35.42	0.33	
Architex (Total Speed)	-51.77	0.70	***	-72.88	0.94	***	-114.97	0.88	***	-69.26	0.87	***
Architex (Total Score)	-0.93	0.37		0.65	-0.34		-0.37	0.25		-0.96	0.22	
Anxiety (STAI)	-3.68	0.49	~	-4.09	0.63	**	-5.70	0.74	***	-1.43	0.16	
Arousal	-0.19	0.13		-0.38	0.11		0.70	-0.29		-0.63	0.28	
Valence	1.03	-0.51	~	0.41	-0.45	~	-0.12	-0.07		0.03	-0.03	
POMS	-9.90	0.63	*	-10.50	0.78	***	-10.58	0.64	**	-7.17	0.31	

DV	Breathwork			Rain			MindGym			VR			All		
	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig
TMT (RTACC)	-21.6	0.391	*	-39.1	0.51	***	-20.35	0.56	***	-39.9	0.33	*	-30.4	0.45	***
Architex (Total Speed)	-81.7	0.791	***	-71.1	0.9	***	-62.33	0.82	***	-91.7	0.87	***	-76.4	0.84	***
Architex (Total Score)	-0.67	0.312	~	-0.13	-0		-0.141	0.06		-0.67	0.23		-0.4	0.14	
Anxiety (STAI)	-4.72	0.621	***	-2.81	0.4	*	-3.889	0.55	***	-3.67	0.47	**	-3.78	0.52	***
Arousal	0.266	-0.12		-0.5	0.22		-0.285	0.12		0.063	0.01		-0.11	0.06	
Valence	0.437	-0.26		0.226	-0.2		0.715	-0.5	**	-0.05	-0.1		0.333	-0.3	*
POMS	-10.3	0.636	***	-8.89	0.59	***	-10.21	0.71	***	-8.95	0.5	***	-9.58	0.61	***

'DV' refers to the dependent variable. 'ΔM' refers to the mean difference (post – pre). 'RBC' refers to rank biserial correlation as a measure of effect size (ranging from -1 to 1) and reflects the relative proportions of positive or negative ranks. 'sig' refers to the FDR-corrected P-value where '~' indicates $.05<P.FDR<.10$, '*' indicates $P.FDR<.05$, '**' indicates $P.FDR<.01$, and '***' indicates $P.FDR<.001$.

When considering all groups combined, there was evidence of a decrease in TMT (RTACC) (W=5443.00, Z=4.32, $P.FDR<.001$, RBC=.45), Architex (Total Speed) (W=6044.00, Z=7.82, $P.FDR<.001$, RBC=.84), Anxiety (STAI) (W=5156.50, Z=4.86, $P.FDR<.001$, RBC = .52), Valence (W=1626.00, Z=-2.14, $P.FDR$=.042, RBC=-.26), and POMS (W=5659.00, Z=5.77, $P.FDR<.001$, RBC=.61)

Examining specific conditions, the MindGym (Breathwork) group demonstrated improvements in Architex (Total Speed) (W=396.00, Z=3.36, $P.FDR<.001$, RBC=.70) and POMS (W=331.00, Z=2.92, $P.FDR$=.014, RBC=.63). The MindGym (Rain) condition showed enhancements for TMT (RTACC) (W=411.00, Z=3.19, $P.FDR<.001$, RBC=.66), Architex (Total Speed) (W=421.00, Z=4.40, $P.FDR<.001$, RBC=.94), STAI (W=353.50, Z=2.94, $P.FDR$=.005, RBC=.63), and POMS (W=442.00, Z=3.80, $P.FDR<.001$, RBC=.78)

The VR (Breathwork) group exhibited improvements in Architex (Total Speed) (W=355.00, Z=3.99, $P.FDR<.001$, RBC=.88), STAI (W=405.00, Z=3.55, $P.FDR<.001$, RBC=.74), and POMS (W=406.00, Z=3.10, $P.FDR$=.005, RBC=.64) (see Supplemental Pages: Appendix A3), while the VR (Rain) condition only showed a significant improvement in Architex (Total Speed) (W=379.00, Z=4.01, $P.FDR<.001$, RBC=.87)

When comparing the combined Breathwork and Rain conditions, both showed significant improvements across multiple measures, with slight variations in effect sizes. Similarly, the combined MindGym and VR conditions significantly enhanced cognitive performance, anxiety reduction, and mood improvement.

Group differences in timepoint effects

Analysis of group differences in timepoint effects revealed no statistically significant differences between experimental conditions after FDR correction for multiple comparisons (all $P.FDR>.05$) (see Supplemental Pages: Appendix B). While all interventions yielded positive outcomes across various measures, including improvements in cognitive performance, anxiety levels, and mood, the magnitude of these changes did not differ significantly between experimental conditions when accounting for multiple comparisons. This lack of significant between-group differences indicates that no single condition demonstrated superior efficacy over the others in terms of the measured variables.

Physiology Measures
Group Differences

Analysis of physiological measures revealed significant differences between experimental conditions, particularly in breath rate patterns, the most notable being between the Breathwork and Rain conditions (see Supplemental Pages: Appendix D for all results).

At the end of the session, Breathwork participants had significantly lower final Breath Rate compared to Rain (*W*=269.50, *P.FDR*<.001, *RBC*=-.70). The overall change in Breath Rate from beginning to end of the experience was significantly different between the two conditions (*W*=198.50, *P.FDR*<.001, *RBC*=-.78), indicating a more substantial decrease in breath rate for the Breathwork group.

Similar patterns were observed when comparing the MindGym (Breathwork) and MindGym (Rain) conditions. MindGym (Breathwork) had a significantly lower final Breath Rate (*W* =77.00, *P.FDR*<.001, *RBC*=-.68) compared to MindGym (Rain), and the overall change in Breath Rate showed greater decrease, with a significantly different Breath Rate (*W*=68.50, *P.FDR*<.001, *RBC*=-.72) with MindGym (Breathwork).

VR (Breathwork) and VR (Rain) showed significant differences in breath rate patterns, with significantly lower Breath Rate at the end of the experience (*W*=55.00, *P.FDR*<.001, *RBC*=-.73) compared to VR (Rain). The overall change in Breath Rate was significantly different (*W*=37.00, *P.FDR*<.001, *RBC*= -.82), with VR (Breathwork) demonstrating a more substantial decrease.

These results consistently demonstrate that participants in the Breathwork conditions, regardless of the delivery method (MindGym or VR), experienced a decrease in breath rate throughout the session compared to those in the Rain conditions. This pattern suggests that the Breathwork interventions effectively modulated participants' breathing patterns, potentially leading to a more relaxed physiological state by the end of the session.

No other physiological measures (EDA, HR, HRV) showed significant differences between conditions after FDR correction.

Changes Over Time

We employed repeated measures ANOVA (rmANOVA) to examine changes over time and the potential effects of experimental conditions. We observed significant temporal changes in HR, Breath Rate, and EDA levels, with only Breath Rate showing interactions between time and technology. No significant main effects of time or interactions with groups were observed for HRV after FDR correction, likely assessed using repeated measures ANOVA.

The results revealed a significant main effect of time on HR ($F_{5.17, 398.38}$=8.78, *P.FDR*=.004), indicating a gradual increase throughout the experiment. EDA levels revealed a significant main effect of time ($F_{2.35, 162.37}$=17.11, *P.FDR*=.004), characterized by an initial increase followed by a plateau around the third timepoint.

Breath Rate showed a significant main effect of time ($F_{2.35, 162.37}$=17.11, *P.FDR*=.002), characterized by an initial decrease followed by a plateau around the third timepoint. A significant interaction between time and Stimuli was also observed ($F_{2.48, 208.49}$=28.49, *P.FDR*=.002). Post hoc paired-samples t-tests (with Holm correction of *P* values) were conducted between timepoints within Stimuli groups. Within the Rain groups combined, breath rate did not reliably change between any timepoints (all *P.HOLM*>.05). Within the Breathwork groups combined, breath rate significantly decreased from the first to second timepoint (*ΔM*=-3.55, *SE*=.49, *t*=7.19, *P.HOLM*<.001, *d*=1.08), remained similar between second and third timepoints (*ΔM*=-1.12, *SE*=0.49, *t*=2.26, *P.HOLM*=.416, *d*=.034), decreased again from third to

fourth timepoints (ΔM=-2.28, SE=.49, t = 4.62, $P.HOLM$<.001, d=.69), without any further reliable decrease from fourth to fifth timepoints (ΔM=.77, SE=.49, t=1.56, $P.HOLM$=1.000, d=.23).

Mediation

Mediation analyses were conducted to test if hypothesized variables mediated the group effects on the outcome scores. No mediation analyses passed the threshold for significance after FDR correction (all $P.FDR$>.05).

DV	Tech x Stim x Moderator			Tech x Moderator			Stim x Moderator			Moderator				
Awe	F	P.RAW	P.FDR	F	P.RAW	P.FDR	F	P.RAW	P.FDR	b	SE	Z	P.RAW	P.FDR
Psych Moderator														
DPES	1.74	0.190	0.824	0.14	0.711	0.887	0.20	0.657	0.957	13.89	3.48	4.00	< .0001	< .001
Immersion	0.03	0.870	0.892	0.08	0.776	0.887	0.02	0.896	0.957	1.16	0.22	5.17	< .0001	< .001
IPQ (General)	0.02	0.892	0.892	4.70	0.032	0.362	1.92	0.169	0.853	6.18	1.72	3.59	< .001	0.002
IPQ (Involvement)	1.08	0.301	0.892	0.12	0.727	0.887	0.82	0.367	0.853	2.24	0.76	2.96	0.003	0.009
IPQ (Spatial Presence)	0.38	0.538	0.892	0.00	0.957	0.957	0.77	0.381	0.853	1.83	0.93	1.98	0.048	0.104
IPQ (Experienced Realism)	1.52	0.220	0.824	0.34	0.560	0.887	3.98	0.048	0.725	0.68	0.87	0.78	0.436	0.620
MODTAS	0.56	0.458	0.892	0.05	0.828	0.887	0.38	0.539	0.899	0.37	0.11	3.28	0.001	0.004
Openness	0.36	0.553	0.892	0.85	0.359	0.796	0.03	0.871	0.957	1.22	0.81	1.51	0.131	0.218
Physio Moderator														
Motion (overall)	2.69	0.104	0.824	0.14	0.711	0.887	1.41	0.238	0.853	0.79	0.50	1.59	0.112	0.210
Breath Rate (first 2 min)	0.13	0.724	0.892	1.08	0.301	0.796	0.06	0.805	0.957	0.15	1.02	0.15	0.883	0.883
HR (first 1 min)	2.45	0.122	0.824	0.59	0.446	0.837	0.01	0.908	0.957	-0.70	0.29	-2.39	0.017	0.042
HRV (first 5 min)	0.03	0.853	0.892	4.04	0.048	0.362	0.00	0.957	0.957	42.30	65.70	0.64	0.520	0.650
Breath Rate change (last - first)	0.02	0.876	0.892	0.81	0.371	0.796	0.82	0.367	0.853	-0.19	0.82	-0.24	0.814	0.872
HR change (last - first)	0.17	0.686	0.892	2.44	0.123	0.508	0.72	0.398	0.853	-0.60	0.80	-0.75	0.455	0.620
HRV change (last - first)	0.68	0.412	0.892	2.28	0.135	0.508	0.55	0.461	0.865	9.92	38.69	0.26	0.798	0.872

Moderation

Our moderation analysis revealed significant effects of both psychological and physiological factors on participants' experiences and performance during the immersive interventions (see Supplemental Pages: Appendix E for detailed results). Among the psychological trait measures, MODTAS emerged as a significant predictor of both Awe (b=.37, SE=.11, Z=3.28, $P.FDR$=.004) (see Table 4) and EDI (b=.03, SE=.01, Z=2.98, $P.FDR$=.015), with higher MODTAS scores associated with greater reported Awe and EDI. DPES also showed a significant main effect on Awe (b=13.89, SE=3.48, Z=4.00, $P.FDR$<.001), indicating that individuals with higher dispositional positive emotions were more likely to experience awe during the intervention.

Table 4. Results from GLM tests of moderation.
'DV' refers to the dependent variable of Awe from the post timepoint. 'Psych Moderator' and 'Physio Moderator' refer to either the psychological or physiological variable defined as the

moderator. 'Tech x Stim x Moderator' refers to the model of a three-way interaction between the moderator and the group factors of Tech (MindGym vs VR) and Stimuli (Breathwork vs Rain). 'Tech x Moderator' refers to the model of a two-way interaction between the moderator and the group factor of Tech (MindGym vs VR). 'Stim x Moderator' refers to the model of a two-way interaction between the moderator and the group factor of Stimuli (Breathwork vs Rain). 'Moderator' refers to the effect of the moderator from the model of only main effects. 'F' refers to F-test from the GLM model. 'b' and 'SE' refer to the unstandardized beta coefficient and its standard error. 'Z' refers to the Z-test from the model. 'p.raw' and 'P.FDR' refer to the uncorrected p-value and FDR-corrected p-value.

The trait of Openness demonstrated an interaction effect with the type of stimuli on anxiety levels as measured by STAI ($F_{1,111}$=9.58, $P.FDR$=.038). Specifically, higher levels of openness were associated with a greater reduction in anxiety for participants in Breathwork, but not for those in Rain.

Psychological state measures also played a significant role in moderating participants' experiences. Immersion showed strong main effects on both Awe (b=1.16, SE=0.22, Z=5.17, $P.FDR$<.001) and EDI (b=0.07, SE=0.02, Z=4.55, $P.FDR$< .001), with higher levels of immersion consistently associated with greater reported awe and emotional depth. IPQ (General) was also positively associated with Awe (b=6.18, SE=1.72, Z=3.59, $P.FDR$=.002) and EDI (b=0.32, SE=0.13, Z=2.49, $P.FDR$=.048). Only IPQ (Involvement) followed this same pattern with Awe (b=2.24, SE=0.76, Z=2.96, $P.FDR$=.009) and EDI (b=0.18, SE=0.05, Z=3.24, $P.FDR$=.009).

Turning to physiological moderators, we found the change in heart rate throughout the intervention (HR-diff) showed a significant interaction effect with the type of technology used on Architex (Total Score) (F_{df1}=10, $P.FDR$=.023). Specifically, an increase in heart rate was associated with decreased accuracy for participants in the MindGym groups, but not for those in the VR.

Discussion

This study aimed to investigate the impact of content delivery systems on two different anxiolytic stimuli: guided breathwork and naturalistic rain. We compared the efficacy of a novel "reflective chamber" (MindGym) with a traditional virtual reality (VR) headset in delivering these interventions. While our initial hypothesis anticipated clear superiority of one technology over the other, our results instead demonstrated the robust effectiveness of MindGym-generated content across both delivery systems. This finding highlights the MindGym's potential as a powerful content generation platform, capable of producing experiences that maintain their efficacy even when translated to more portable VR formats.

Our findings revealed significant reductions in anxiety measures across all conditions, with the STAI showing significant improvements in all groups except for the breathwork condition in the MindGym. These results may be of clinical relevance and are comparable to benchmarks from other acute anxiety interventions. Our study demonstrated an overall STAI reduction of 11.67%

across all groups, with breathwork showing the highest reduction at 15.10%. These findings align with recent studies on non-pharmacological interventions for acute anxiety reduction (e.g., Johnson et al. [4] reported a 15% reduction in STAI scores in non-anxious populations exposed to a psychedelic-esque stroboscopic lights and binaural beats experience, and a 13% reduction in response to breath-focused meditation; Sriboonlert et al. [85] found a 17% reduction in anxiety scores following aromatherapy interventions;   et al. [86] observed a 7% reduction from exercise in lower trait anxious individuals and a 17% reduction for those with higher trait anxiety).

Our results suggest that both the MindGym and VR delivery systems, combined with either rain or breathwork stimuli, can produce anxiety reductions comparable to these established interventions. The fact that such basic interventions, which don't fully showcase the complex experiences unique to the MindGym, were equally effective in both systems highlights the MindGym's potential as a content creation tool. It suggests that even more sophisticated MindGym-generated experiences could be successfully adapted for VR, potentially enhancing their accessibility without compromising efficacy. However, it's important to note that these effects are still considerably lower than the 30-50% reduction typically observed with pharmacological interventions [87]. This comparison highlights the potential for non-pharmacological approaches while also acknowledging the current limitations in their efficacy compared to medication-based treatments. Interestingly, we observed no significant changes in arousal levels, while valence only improved significantly in the MindGym condition. The interpretation of pre-post results centered around task performance (e.g., Architex) is challenging without a control condition, as some improvements may be attributable to practice effects. For instance, the TMT typically shows improvement with repeated administrations [88,89].

Contrary to our initial expectations, we found no main effects of either stimulus type or technology on the outcome measures. This lack of differentiation between the MindGym and VR conditions is particularly noteworthy, as it suggests that the MindGym-generated content retained its effectiveness when translated to a VR format. While there were numerical trends suggesting potentially better affective outcomes in the MindGym (e.g., greater reduction in STAI scores and mood disturbance), these differences did not reach statistical significance. We observed a consistent pattern of "more" significant effects in the MindGym across most measures, but we caution against over-interpreting these trends. Surprisingly, we did not observe the expected differences in immersion between the two technologies. This lack of differentiation may suggest that both systems were equally effective in creating an immersive experience, or that our measures were not sensitive enough to detect subtle differences. It also points to the successful adaptation of MindGym-generated content for VR platforms.

The most notable physiological finding was the decrease in breath rate in the breathwork condition compared to the rain condition. This result serves as a sanity check, confirming that participants engaged with the breathwork instructions as intended. However, we found no significant differences in electrodermal activity (EDA), heart rate (HR), or heart rate variability (HRV) as a function of technology or stimulus type.

We explored whether individual differences could account for variations in response to the interventions. Several interesting moderation effects emerged. DPES predicted the amount of

awe experienced, though this finding is somewhat expected given the overlap between DPES and awe measures. MODTAS, which has previously been shown to predict audiovisual-induced chills [90], had a main effect on both awe and ego dissolution. MODTAS also moderated the effects of stimuli on anxiety reduction, with higher trait absorption associated with greater anxiety reduction in the breathwork condition across both technologies. Openness to experience moderated the impact on STAI scores, with more open individuals experiencing greater anxiety reduction in the breathwork condition, but not in the rain condition.

These moderation effects exclusive to breathwork may reflect the importance of explicit instructions and participants' willingness to fully engage with the experience. The breathwork condition may have benefited individuals who are more prone to absorption and open to new experiences, possibly due to its more structured nature compared to the passive rain condition. Immersion and presence, while not differing between conditions, were potent predictors of awe and ego dissolution across all groups. This aligns with previous research showing that the degree of immersion in virtual environments impacts memory encoding and neural representations [41]. Interestingly, motion sickness negatively impacted ego dissolution, suggesting that physical discomfort may anchor individuals in their bodily sensations, hindering the experience of ego transcendence.

In conclusion, this study contributes to the growing body of research on non-pharmacological interventions for acute anxiety reduction while also demonstrating the potential of the MindGym as a versatile content creation platform. The comparable efficacy of both the MindGym and VR systems, using content originally generated for the MindGym, suggests that experiences designed in the MindGym can be successfully translated to more portable VR formats without significant loss of effectiveness. This finding is particularly promising given that the interventions used in this study were relatively basic and did not fully leverage the unique capabilities of the MindGym. Future research should focus on developing more complex, immersive experiences using the MindGym's full potential, and exploring how these can be effectively adapted for VR and other delivery systems. This approach could lead to the development of highly effective, accessible, and scalable interventions for anxiety management, potentially bridging the gap with pharmacological treatments while offering the benefits of non-pharmacological approaches.

Limitations

A significant limitation of this study is that the "VR" condition was not a true virtual reality experience, but rather a 3D video displayed in a VR headset. A more equitable comparison would have involved creating a virtual environment that mimicked the MindGym chamber and responded to head movements, potentially enhancing immersion and presence. The current design may have inadvertently positioned the MindGym experience as a "steel man" argument, with many of its immersive properties carried over to the VR condition.

However, this limitation also highlights the potency of the MindGym programming, demonstrating that its effects can be successfully replicated using a VR device. This finding suggests that the content itself, rather than the specific delivery system, may be the primary driver of the observed anxiolytic effects.

Future Directions

Future research should focus on conducting experiments with more visually diverse content to identify where these content delivery systems meaningfully diverge. This could involve developing true VR environments that more closely mimic the MindGym experience, allowing for a more direct comparison of the technologies. Investigating the impact of these delivery systems on different types of anxiety (e.g., social anxiety, specific phobias) could determine if certain technologies are more effective for particular anxiety subtypes. Exploring the long-term effects of repeated exposures to these interventions would assess their potential as ongoing anxiety management tools. Examining the role of individual differences (e.g., absorption, openness) in treatment response could lead to the development of more personalized anxiety interventions.

The lack of significant differences in these physiological measures across conditions may suggest that the anxiolytic effects are mediated through cognitive or emotional pathways rather than direct physiological changes. Therefore, investigating the neural correlates of the anxiolytic effects observed in this study would better our understanding of the mechanisms underlying the effectiveness of these interventions. Additionally, this might account for some individual differences in responses that weren't seen by self-reported trait variables.

Conclusion

In conclusion, while our study did not find significant differences between the MindGym and VR delivery systems, it demonstrated the potential of both technologies as platforms for acute anxiety reduction. The observed moderation effects highlight the importance of individual differences in treatment response and suggest that tailoring interventions to personal characteristics may enhance their effectiveness. As immersive technologies continue to evolve, further research in this area holds promise for developing more accessible and effective anxiety management tools.

Acknowledgments

We warmly thank the research participants. We also thank Asli Sara Bilgili who assisted N.S., G.W., and C.L. with data collection. We thank Lumena, Inc. for their troubleshooting assistance throughout the course of the experiment. We also extend gratitude to Leonardo Christov-Moore who provided commentary on the study design in its early phases.

Competing Interests

This research was financially supported under a Research Services agreement executed between IACS and Lumena, Inc. It is imperative to note that the terms of this agreement did not incorporate any provisions or incentives contingent upon the attainment of specific outcomes or successful results but were solely predicated on underwriting the requisite expenses for the genuine and unbiased execution of the study. Lumena, Inc. had no input on the design nor analysis of this study, apart from supplying MindGym's content, delivery system hardware and software, and the 360 videos utilized in the VR.

Author Contributions Statement

Study conceptualization and design: N.R. Study implementation and logistics: N.R., N.S. Data collection: N.S., G.W., C.L. Data analysis: M.J., V.K., F.S., V.K., N.R., N.S. Manuscript writing and editing (all drafts): N.S., N.R., G.W., C.L., F.S., M.J. All authors approved the final version for submission.

References

1. Naef AC, Jeitziner MM, Knobel SEJ, et al. Investigating the role of auditory and visual sensory inputs for inducing relaxation during virtual reality stimulation. *Sci Rep*. 2022;12(1):17073. doi:10.1038/s41598-022-21575-9

2. Weller R, Cepok J, Arzaroli R, et al. Effects of immersion and navigation agency in virtual environments on emotions and behavioral intentions. *Frontiers in Virtual Reality*. 2022;3. Accessed December 12, 2023. https://www.frontiersin.org/articles/10.3389/frvir.2022.893052

3. Browning MHEM, Shin S, Drong G, et al. Daily exposure to virtual nature reduces symptoms of anxiety in college students. *Sci Rep*. 2023;13(1):1239. doi:10.1038/s41598-023-28070-9

4. Johnson M, Simonian N, Reggente N. Lightening the Mind with Audiovisual Stimulation as an Accessible Alternative to Breath-Focused Meditation for Mood and Cognitive Enhancement. Published online January 1, 2024. doi:10.31234/osf.io/hb7wx

5. Miller N, Stepanova ER, Desnoyers-Stewart J, et al. Awedyssey: Design Tensions in Eliciting Self-transcendent Emotions in Virtual Reality to Support Mental Well-being and Connection. In: *Proceedings of the 2023 ACM Designing Interactive Systems Conference*. ACM; 2023:189-211. doi:10.1145/3563657.3595998

6. Haller H, Cramer H, Lauche R, Gass F, Dobos GJ. The prevalence and burden of subthreshold generalized anxiety disorder: a systematic review. *BMC Psychiatry*. 2014;14:128. doi:10.1186/1471-244X-14-128

7. National Institute of Mental Health (NIMH) - Transforming the understanding and treatment of mental illnesses. Accessed September 12, 2024. https://www.nimh.nih.gov/

8. Any Anxiety Disorder - National Institute of Mental Health (NIMH). Accessed September 12, 2024. https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder

9. Hofmann SG, Smits JAJ. Cognitive-behavioral therapy for adult anxiety disorders: a meta-analysis of randomized placebo-controlled trials. *J Clin Psychiatry*. 2008;69(4):621-632. doi:10.4088/jcp.v69n0415

10. Bandelow B, Reitt M, Röver C, Michaelis S, Görlich Y, Wedekind D. Efficacy of treatments for anxiety disorders: a meta-analysis. *Int Clin Psychopharmacol*. 2015;30(4):183-192. doi:10.1097/YIC.0000000000000078

11. Crippa JA, Guimarães FS, Campos AC, Zuardi AW. Translational Investigation of the Therapeutic Potential of Cannabidiol (CBD): Toward a New Age. *Front Immunol*. 2018;9. doi:10.3389/fimmu.2018.02009

12. George MS, Ward HE, Ninan PT, et al. A pilot study of vagus nerve stimulation (VNS) for treatment-resistant anxiety disorders. *Brain Stimulation*. 2008;1(2):112-121. doi:10.1016/j.brs.2008.02.001

13. Kaplan S. The restorative benefits of nature: Toward an integrative framework. *Journal of Environmental Psychology*. 1995;15(3):169-182. doi:10.1016/0272-4944(95)90001-2

14. Maller C, Townsend M, Pryor A, Brown P, St Leger L. Healthy nature healthy people: "contact with nature" as an upstream health promotion intervention for populations. *Health Promot Int*. 2006;21(1):45-54. doi:10.1093/heapro/dai032

15. Alvarsson JJ, Wiens S, Nilsson ME. Stress Recovery during Exposure to Nature Sound and Environmental Noise. *Int J Environ Res Public Health*. 2010;7(3):1036-1046. doi:10.3390/ijerph7031036

16. Bratman GN, Hamilton JP, Daily GC. The impacts of nature experience on human cognitive function and mental health. *Ann N Y Acad Sci*. 2012;1249:118-136. doi:10.1111/j.1749-6632.2011.06400.x

17. Jo H, Song C, Miyazaki Y. Physiological Benefits of Viewing Nature: A Systematic Review of Indoor Experiments. *Int J Environ Res Public Health*. 2019;16(23):4739. doi:10.3390/ijerph16234739

18. Van Hedger SC, Nusbaum HC, Clohisy L, Jaeggi SM, Buschkuehl M, Berman MG. Of cricket chirps and car horns: The effect of nature sounds on cognitive performance. *Psychon Bull Rev*. 2019;26(2):522-530. doi:10.3758/s13423-018-1539-1

19. Hu C, Zhu K, Huang K, et al. Using natural intervention to promote subjective well-being of essential workers during public-health crises: A Study during COVID-19 pandemic. *J Environ Psychol*. 2022;79:101745. doi:10.1016/j.jenvp.2021.101745

20. Johansson G, Juuso P, Engström Å. Nature-based interventions to promote health for people with stress-related illness: An integrative review. *Scand J Caring Sci*. 2022;36(4):910-925. doi:10.1111/scs.13089

21. Song I, Baek K, Kim C, Song C. Effects of nature sounds on the attention and physiological and psychological relaxation. *Urban Forestry & Urban Greening*. 2023;86:127987. doi:10.1016/j.ufug.2023.127987

22. Chinagudi S, Badami S, Herur A, Patil S, Gv S, Ankad R. Immediate effect of short duration of slow deep breathing on heart rate variability in healthy adults. *Natl J Physiol Pharm Pharmacol*. 2014;4(3):233. doi:10.5455/njppp.2014.4.060520141

23. Jerath R, Crawford MW, Barnes VA, Harden K. Self-Regulation of Breathing as a Primary Treatment for Anxiety. *Appl Psychophysiol Biofeedback*. 2015;40(2):107-115. doi:10.1007/s10484-015-9279-8

24. Russo MA, Santarelli DM, O'Rourke D. The physiological effects of slow breathing in the healthy human. *Breathe (Sheff)*. 2017;13(4):298-309. doi:10.1183/20734735.009817

25. Zaccaro A, Piarulli A, Laurino M, et al. How Breath-Control Can Change Your Life: A Systematic Review on Psycho-Physiological Correlates of Slow Breathing. *Front Hum Neurosci*. 2018;12:353. doi:10.3389/fnhum.2018.00353

26. Aideyan B, Martin GC, Beeson ET. A Practitioner's Guide to Breathwork in Clinical Mental Health Counseling. *Journal of Mental Health Counseling*. 2020;42(1):78-94. doi:10.17744/mehc.42.1.06

27. Magnon V, Dutheil F, Vallet GT. Benefits from one session of deep and slow breathing on vagal tone and anxiety in young and older adults. *Sci Rep*. 2021;11:19267. doi:10.1038/s41598-021-98736-9

28. Balban MY, Neri E, Kogon MM, et al. Brief structured respiration practices enhance mood and reduce physiological arousal. *CR Med*. 2023;0(0). doi:10.1016/j.xcrm.2022.100895

29. World Urbanization Prospects The 2018 Revision.

30. Ulrich RS. Aesthetic and Affective Response to Natural Environment. In: Altman I, Wohlwill JF, eds. *Behavior and the Natural Environment*. Springer US; 1983:85-125. doi:10.1007/978-1-4613-3539-9_4

31. Anderson AP, Mayer MD, Fellows AM, Cowan DR, Hegel MT, Buckey JC. Relaxation with Immersive Natural Scenes Presented Using Virtual Reality. *Aerospace Medicine and Human Performance*. 2017;88(6):520-526. doi:10.3357/AMHP.4747.2017

32. Maples-Keller JL, Bunnell BE, Kim SJ, Rothbaum BO. The Use of Virtual Reality Technology in the Treatment of Anxiety and Other Psychiatric Disorders. *Harv Rev Psychiatry*. 2017;25(3):103-113. doi:10.1097/HRP.0000000000000138

33. Schutte NS, Bhullar N, Stilinović EJ, Richardson K. The Impact of Virtual Environments on Restorativeness and Affect. *Ecopsychology*. 2017;9(1):1-7. doi:10.1089/eco.2016.0042

34. Jerdan SW, Grindle M, van Woerden HC, Kamel Boulos MN. Head-Mounted Virtual Reality and Mental Health: Critical Review of Current Research. *JMIR Serious Games*. 2018;6(3):e14. doi:10.2196/games.9226

35. White MP, Yeo NL, Vassiljev P, et al. A prescription for "nature" – the potential of using virtual nature in therapeutics. *Neuropsychiatr Dis Treat*. 2018;14:3001-3013. doi:10.2147/NDT.S179038

36. Yu CP, Lee HY, Lu WH, Huang YC, Browning MHEM. Restorative effects of virtual natural settings on middle-aged and elderly adults. *Urban Forestry & Urban Greening*. 2020;56:126863. doi:10.1016/j.ufug.2020.126863

37. Annerstedt M, Jönsson P, Wallergård M, et al. Inducing physiological stress recovery with sounds of nature in a virtual reality forest--results from a pilot study. *Physiol Behav*. 2013;118:240-250. doi:10.1016/j.physbeh.2013.05.023

38. Proverbio AM, De Benedetto F, Ferrari MV, Ferrarini G. When listening to rain sounds boosts arithmetic ability. *PLoS One*. 2018;13(2):e0192296. doi:10.1371/journal.pone.0192296

39. Villani D, Riva F, Riva G. New technologies for relaxation: The role of presence. *International Journal of Stress Management*. 2007;14(3):260-274. doi:10.1037/1072-5245.14.3.260

40. Slobounov SM, Ray W, Johnson B, Slobounov E, Newell KM. Modulation of cortical activity in 2D versus 3D virtual reality environments: An EEG study. *International Journal of Psychophysiology*. 2015;95(3):254-260. doi:10.1016/j.ijpsycho.2014.11.003

41. Essoe JKY, Reggente N, Ohno AA, Baek YH, Dell'Italia J, Rissman J. Enhancing learning and retention with distinctive virtual reality environments and mental context reinstatement. *npj Sci Learn*. 2022;7(1):1-14. doi:10.1038/s41539-022-00147-6

42. Lessiter J, Freeman J, Keogh E, Davidoff J. A Cross-Media Presence Questionnaire: The ITC-Sense of Presence Inventory. *Presence: Teleoperators and Virtual Environments*. 2001;10(3):282-297. doi:10.1162/105474601300343612

43. Dede C. Immersive Interfaces for Engagement and Learning. *Science*. 2009;323(5910):66-69. doi:10.1126/science.1167311

44. Reggente N. VR for Cognition and Memory. *Curr Top Behav Neurosci*. Published online July 14, 2023. doi:10.1007/7854_2023_425

45. Manjrekar S, Sandilya S, Bhosale D, Kanchi S, Pitkar A, Gondhalekar M. CAVE: An emerging immersive technology -a review. *Proceedings - UKSim-AMSS 16th International Conference on Computer Modelling and Simulation, UKSim 2014*. Published online March 1, 2014:131-136. doi:10.1109/UKSim.2014.20

46. Reggente N, Essoe JKY, Aghajan ZM, et al. Enhancing the Ecological Validity of fMRI Memory Research Using Virtual Reality. *Frontiers in Neuroscience*. 2018;12. Accessed October 8, 2023. https://www.frontiersin.org/articles/10.3389/fnins.2018.00408

47. Weil A. *Breathing: The Master Key to Self Healing (The Self Healing Series)*. Sounds True; 1999.

48. Pandekar PP, Thangavelu PD. Effect of 4-7-8 Breathing Technique on Anxiety and Depression in Moderate Chronic Obstructive Pulmonary Disease Patients. *International Journal of Health Sciences*. 2019;(5).

49. Riva G, Baños RM, Botella C, Mantovani F, Gaggioli A. Transforming Experience: The Potential of Augmented Reality and Virtual Reality for Enhancing Personal and Clinical Change. *Front Psychiatry*. 2016;7. doi:10.3389/fpsyt.2016.00164

50. Chirico A, Ferrise F, Cordella L, Gaggioli A. Designing Awe in Virtual Reality: An Experimental Study. *Front Psychol*. 2017;8:2351. doi:10.3389/fpsyg.2017.02351

51. Chirico A, Cipresso P, Yaden DB, Biassoni F, Riva G, Gaggioli A. Effectiveness of Immersive Videos in Inducing Awe: An Experimental Study. *Sci Rep*. 2017;7(1):1218. doi:10.1038/s41598-017-01242-0

52. Adrien V, Bui E, Baubet T, et al. Beyond Virtual Reality: Towards Screen-Free Interfaces for Post-traumatic Stress Disorder Interventions. Published online November 16, 2023. doi:10.31234/osf.io/24beh

53. Shiota MN, Keltner D, John OP. Positive emotion dispositions differentially associated with Big Five personality and attachment style. *The Journal of Positive Psychology*. 2006;1(2):61-71. doi:10.1080/17439760500510833

54. Jamieson G. The Modified Tellegen Absorption Scale: a clearer window on the structure and meaning of absorption. *Australian Journal of Clinical and Experimental Hypnosis*. 2005;Vol. 33:119-139.

55. Costa PT, McCrae RR. The Five-Factor Model of Personality and Its Relevance to Personality Disorders. *Journal of Personality Disorders*. 1992;6(4):343-359. doi:10.1521/pedi.1992.6.4.343

56. Spielberger C, Gorsuch R, Lushene R, Vagg P, Jacobs G. *Manual for the State-Trait Anxiety Inventory (Form Y1 – Y2)*. Vol IV.; 1983.

57. Heuchert JP, McNair DM. Profile of Mood States 2nd Edition™. Published online June 11, 2012. doi:10.1037/t05057-000

58. Lau MA, Bishop SR, Segal ZV, et al. The Toronto Mindfulness Scale: Development and Validation. *Journal of Clinical Psychology*. 2006;62(12):1445-1467. doi:10.1002/jclp.20326

59. Yaden D, Kaufman S, Hyde E, et al. The development of the Awe Experience Scale (AWE-S): A multifactorial measure for a complex emotion. *The Journal of Positive Psychology*. 2018;14. doi:10.1080/17439760.2018.1484940

60. Swami V, Salem N, Furnham A, Tovée MJ. Initial examination of the validity and reliability of the female photographic figure rating scale for body image assessment. *Personality and Individual Differences*. 2008;44(8):1752-1761. doi:10.1016/j.paid.2008.02.002

61. Elk M, Karinen A, Specker E, Stamkou E, Baas M. 'Standing in Awe': The Effects of Awe on Body Perception and the Relation with Absorption. *Collabra*. 2016;2:4. doi:10.1525/collabra.36

62. Nour MM, Evans L, Nutt D, Carhart-Harris RL. Ego-Dissolution and Psychedelics: Validation of the Ego-Dissolution Inventory (EDI). *Front Hum Neurosci*. 2016;10:269. doi:10.3389/fnhum.2016.00269

63. Schubert T, Friedmann F, Regenbrecht H. The Experience of Presence: Factor Analytic Insights. *Presence: Teleoperators & Virtual Environments*. 2001;10(3):266-281. doi:10.1162/105474601300343603

64. Jacobs C, M Rigby J. Developing measures of immersion and motivation for learning technologies in healthcare simulation: a pilot study. *J Adv Med Educ Prof*. 2022;10(3):163-171. doi:10.30476/JAMP.2022.95226.1632

65. Gianaros PJ, Muth ER, Mordkoff JT, Levine ME, Stern RM. A Questionnaire for the Assessment of the Multiple Dimensions of Motion Sickness. *Aviat Space Environ Med*. 2001;72(2):115-119.

66. Reitan RM. Validity of the Trail Making Test as an indicator of organic brain damage. *Perceptual and Motor Skills*. 1958;8:271-276. doi:10.2466/PMS.8.7.271-276

67. Kuhn T, Gottlieb L. Architex. Published online 2022. https://www.myarchitex.com/

68. JASP Team. JASP. Published online 2024. https://jasp-stats.org/

69. Brooks ME, Kristensen K, Benthem KJ van, et al. glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*. 2017;9(2):378-400.

70. Lenth RV. _emmeans: Estimated Marginal Means, aka Least Squares Means. Published online 2022. https://CRAN.R-project.org/package=emmeans

71. RStudio Team. RStudio: Integrated Development for R. RStudio, PBC. Published online 2020. http://www.rstudio.com/

72. Hartig F. DHARMa. GitHub. Accessed September 9, 2024. http://florianhartig.github.io/DHARMa/

73. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*. 1995;57(1):289-300. doi:10.1111/j.2517-6161.1995.tb02031.x

74. Lindquist MA, Mejia A. Zen and the art of multiple comparisons. *Psychosom Med*. 2015;77(2):114-125. doi:10.1097/PSY.0000000000000148

75. Rosseel Y. lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical Software*. 2012;48:1-36. doi:10.18637/jss.v048.i02

76. Kothe C, Shirazi SY, Stenner T, et al. The Lab Streaming Layer for Synchronized Multimodal Recording. Published online February 14, 2024:2024.02.13.580071. doi:10.1101/2024.02.13.580071

77. Makowski D, Pham T, Lau ZJ, et al. NeuroKit2: A Python toolbox for neurophysiological signal processing. *Behav Res Methods*. 2021;53(4):1689-1696. doi:10.3758/s13428-020-01516-y

78. Butterworth S. On the Theory of Filter Amplifiers. *Experimental Wireless \& the Wireless Engineer*. 1930;7:536-541.

79. Breunig MM, Kriegel HP, Ng RT, Sander J. LOF: identifying density-based local outliers. *SIGMOD Rec*. 2000;29(2):93-104. doi:10.1145/335191.335388

80. Pedregosa F, Varoquaux G, Gramfort A, et al. Scikit-learn: Machine Learning in Python. *MACHINE LEARNING IN PYTHON*. Published online 2011.

81. Kleckner IR, Jones RM, Wilder-Smith O, et al. Simple, Transparent, and Flexible Automated Quality Assessment Procedures for Ambulatory Electrodermal Activity Data. *IEEE Trans Biomed Eng*. 2018;65(7):1460-1467. doi:10.1109/TBME.2017.2758643

82. Lu W, Nystrom MM, Parikh PJ, et al. A semi-automatic method for peak and valley detection in free-breathing respiratory waveforms. *Med Phys*. 2006;33(10):3634-3636. doi:10.1118/1.2348764

83. Welch P. The use of fast Fourier transform for the estimation of power spectra: A method based on time averaging over short, modified periodograms. *IEEE Transactions on Audio and Electroacoustics*. 1967;15(2):70-73. doi:10.1109/TAU.1967.1161901

84. Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation*. 1996;93(5):1043-1065.

85. Sriboonlert J, Munkong W, Rintawut S, Paladkhua S, Suwongsa R, Kirisattayakul W. Cananga odorata Aromatherapy Reduces Anxiety in Unexperienced Patients Hospitalized for Interventional Neuroradiology Procedures: A Randomized Control Trial. *J Evid Based Complementary Altern Med*. 2023;28:2515690X221150527. doi:10.1177/2515690X221150527

86. Petruzzello SJ, Snook EM, Gliottoni RC, Motl RW. Anxiety and mood changes associated with acute cycling in persons with multiple sclerosis. *Anxiety, Stress, & Coping*. 2009;22(3):297-307. doi:10.1080/10615800802441245

87. Mudgalkar N, Kandi V, Baviskar A, Kasturi RR, Bandurapalli B. Preoperative Anxiety Among Cardiac Surgery Patients and Its Impact on Major Adverse Cardiac Events and Mortality– A Randomized, Parallel-Group Study. *Annals of Cardiac Anaesthesia*. 2022;25(3):293. doi:10.4103/aca.aca_80_21

88. Craddick RA, Stern MR. Practice Effects on the Trail Making Test. *Percept Mot Skills*. 1963;17(3):651-653. doi:10.2466/pms.1963.17.3.651

89. McKeown K, Richards E, Centre for Innovative Ageing, Swansea University, Swansea, et al. The Trails Making Test. Does a Single Trial Reflect Performance Capability? *obm neurobiol*. 2021;05(02):1-1. doi:10.21926/obm.neurobiol.2102100

90. Schoeller F, Christov-Moore L, Lynch C, Diot T, Reggente N. Predicting individual differences in peak emotional response. *PNAS Nexus*. 2024;3(3):pgae066. doi:10.1093/pnasnexus/pgae066

Abbreviations
VR: Virtual Reality
ECG: Electrocardiogram
EDA: Electrodermal Activity
HR: Heart Rate
HRV: Heart Rate Variability
RR: Respiration Rate