Contrasting cognitive, behavioral, and physiological responses to breathwork vs. naturalistic stimuli in reflective chamber and VR headset environments

Short Title: Comparing immersive tech for relaxation

Ninette Simonian[1], Micah Alan Johnson, PhD[1,2], Caitlin Lynch[1], Geena Wang[1], Velu Kumaravel[1], Taylor Kuhn, PhD, MS[3], Félix Schoeller, PhD[1], Nicco Reggente, PhD [1]

[1] Institute for Advanced Consciousness Studies, Santa Monica, CA, USA
[2] Department of Biostatistics, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA
[3] Department of Psychiatry and Behavioral Sciences, Semel Institute of Neuroscience and Human Behavior, University of California, Los Angeles, Los Angeles, CA, USA

Corresponding Author:
Email: nicco@advancedconsciousness.org (NR)

Abstract

The MindGym, a novel immersive technology utilizing a reflective chamber environment, was developed to create standardized experiential content, including anxiolytic experiences. This study examined whether therapeutic experiences originally created in the MindGym could maintain their efficacy when delivered via 360-degree recordings through virtual reality (VR) headsets. A randomized controlled trial (N=126) compared anxiety reduction, cognitive performance, and physiological responses across four conditions: MindGym and VR platforms, each delivering either breathwork or rain stimuli. Results demonstrated significant improvements across all conditions in cognitive performance (Trail Making Test RTACC, p.fdr<.001; Architex Total Speed, p.fdr<.001) and anxiety reduction (STAI, p.fdr<.001). Breathwork conditions produced greater decreases in breath rate compared to rain stimuli (p.fdr=.002). Treatment responses were moderated by individual differences, with absorption (MODTAS) predicting both awe (p.fdr=.004) and ego dissolution (p.fdr=.015), while openness to experience interacted with stimuli type to influence anxiety reduction (p.fdr=.038). The anxiolytic effects originally generated in the MindGym maintained full efficacy when translated to VR delivery, with no significant differences in effectiveness or immersion between the original environment and its virtual reproduction. These findings establish the MindGym as a viable content creation platform for immersive, anxiety-reducing experiences that can be successfully adapted to more accessible delivery systems, while highlighting the potential for personalization based on individual differences. Future research should investigate the translation of more complex MindGym-generated experiences to expand accessible anxiety management tools.

Introduction

Experience shapes human cognition and behavior, yet its variability limits prescriptive outcomes.

Experiential technology, leveraging audiovisual stimuli, offers potential for targeted interventions with predictable effects on mood and cognition [1–4]. Immersive virtual reality (VR) experiences can elicit specific emotional responses (e.g., awe) and enhance well-being through increased felt connectedness [5].

Anxiety disorders, affecting 19.1% of adults globally, significantly impact healthcare systems and individual well-being, with even subclinical levels diminishing quality of life [6,7][8]. First-line treatments like cognitive-behavioral therapy and medication offer benefits but face limitations such as side effects and relapse potential [9,10]. Alternative approaches (mindfulness, exercise, pharmacotherapies) show variable efficacy [6,10,11], while emerging interventions (vagal nerve stimulation, cannabidiol) require further safety and efficacy research [11,12]. This landscape necessitates innovative approaches to anxiety management.

Two promising avenues for acute anxiety reduction are exposure to naturalistic stimuli and guided breathwork. Nature exposure and related therapies have been shown to facilitate stress recovery, mood enhancement, improved cognitive functions, and overall well-being while alleviating anxiety symptoms and boosting health and self-efficacy [13–21]. Breathwork techniques demonstrate comparable efficacy through distinct mechanisms, inducing parasympathetic states that promote stress reduction and enhance emotional regulation [22–28].

Despite their potential, these approaches face accessibility challenges, particularly in urban environments with limited access to natural settings [29] and due to the lack of integration of breathwork and somatic therapies into standard mental health care [11]. While immersive

technologies offer a solution by creating scalable, accessible virtual experiences, enabling well-

controlled, "plug-and-play" interventions with optimized effectiveness and expanded reach.

Innovative platforms for generating immersive content, such as the MindGym, hold promise for

creating highly effective experiences that could potentially be adapted to various delivery

systems, including more portable VR formats. Research shows combined audiovisual nature

scenes offer enhanced psychological benefits over single-modality presentations [1,3], aligning

with Attention Restoration and Stress Reduction theories [13,30]. 360-degree nature videos

demonstrate rapid mood improvements [21,31–36], while natural sounds enhance physiological

recovery [37] and cognitive performance [38].

The efficacy of these virtual interventions hinges on immersion—the technological capability to

create a sense of "being there"—and presence, the subjective experience of feeling transported to

the virtual environment. These factors are crucial for eliciting powerful psychological and

physiological responses [39,40]. Crucially, research shows the cognitive benefits of VR depend

on users truly feeling "there," emphasizing the fundamental role of high immersion and presence

[41]. Both technological capabilities and subjective experiences like perceived realism and

enjoyment influence these factors [42–44]. Immersion levels vary across VR systems, from

moderate levels in consumer-grade headsets to heightened experiences in advanced setups

[45,46]. These findings underscore the potential of immersive technology to create targeted,

accessible interventions that leverage nature's restorative properties.

The current study investigated whether effective experiences created in one immersive

technology could be successfully translated to another by comparing two immersive technologies

in delivering anxiolytic content: a "reflective chamber" (MindGym) and a traditional VR headset.

Both incorporated naturalistic rain sounds or guided 4-7-8 breathwork [24,47,48]. Our primary

aim was to determine whether the experience designed for the MindGym could maintain their

efficacy when adapted to more widely accessible VR platforms. We hypothesized that

MindGym, designed for enhanced immersion and presence, would outperform VR by amplifying

nature and breathwork effects while minimizing VR-associated discomfort [49–52]. This

comparison sought to inform the development of novel, scalable anxiety-reducing interventions,

bridging the gap between traditional therapies and the need for accessible mental health

solutions.

Methods

Participants

95   A total of 126 participants (63 females; age range 18-74 years; mean 41, SD 15.19) participated

96   in the study. Participants were randomly assigned to one of four groups, as shown in Table 1.

97   **Table 1. Participant characteristics by group and analysis type**

Group	N	Gender (M/F)	Age Range	M	SD
VR (Breathwork)	33	16/17	20-71	42.61	15.18
VR (Rain)	29	16/13	18-69	41.30	15.54
MindGym (Breathwork)	31	15/16	22-74	38.45	15.93
MindGym (Rain)	32	16/16	19-68	41.53	14.50

F = female; M = male. Age is reported in years. M and SD represent the mean and standard

deviation of age, respectively.

10 participants were excluded from the physiological analysis due to equipment failure or

excessive movement resulting in a total of 116 participants for statistical analysis (61 females;

age range 18-74 years; mean 40.54, SD 14.56). Respiration was analyzed for 88 participants,

ECG was analyzed for 80 participants, and EDA was analyzed for 96 participants (see Table 2).

All participants passed quality assurance checks, with respiration rates below 20 breaths per

minute following data cleaning and processing (see Methods for further explanation).

Table 2. Participant characteristics for physiological measures by group

Measure	Group	N	Gender (F/M)	Age Range	M	SD
Respiration						
	VR (Breathwork)	22	12/10	20-65	39.91	14.36
	VR (Rain)	19	8/11	18-69	43.15	16.56
	MindGym (Breathwork)	28	14/14	23-74	36.96	15.17
	MindGym (Rain)	19	11/8	19-63	41.89	14.11
ECG						
	VR (Breathwork)	18	9/9	20-65	41.17	14.31
	VR (Rain)	16	9/7	18-68	41.44	17.00
	MindGym (Breathwork)	26	15/11	23-74	37.04	15.84
	MindGym (Rain)	20	12/8	19-68	41.80	13.33
EDA						
	VR (Breathwork)	23	14/9	20-65	41.61	13.26
	VR (Rain)	22	9/13	18-69	40.77	15.72
	MindGym (Breathwork)	28	14/14	23-68	36.82	14.65
	MindGym (Rain)	23	14/9	20-63	41.43	13.03

F = female; M = male. Age is reported in years. M and SD represent the mean and standard

deviation of age, respectively. ECG = Electrocardiogram; EDA = Electrodermal Activity.

Of the 126 enrolled participants, 87 completed a one-week follow-up survey (See Table 3). The

similar distribution of demographic characteristics suggests random attrition, though this

conclusion is limited by incomplete demographic data across the full sample.

Table 3. Study participants by follow-up completion status across groups and

demographics

		Completers (N=49)	Non-completers (N=24)
Group			
	VR (Breathwork)	12	9
	VR (Rain)	9	6

	MindGym (Breathwork)	13	5
	MindGym (Rain)	15	4
Gender			
	Male	25	10
	Female	24	14
Education			
	Bachelor's degree	27	13
	Master's degree	9	3
	Some college	6	5
	High school diploma	5	1
	GED or Equivalent	0	1
	Professional degree	1	0
	Associate's degree	1	1
Race			
	White	23	11
	Asian	10	9
	Multiracial	6	3
	Black/African American	2	1
	Hispanic/Latino	4	0
	Indigenous American	1	0
	Not disclosed	3	0
Ethnicity			
	Not Hispanic/Latinx	39	23
	Hispanic/Latinx	9	1
	Unknown	1	0

Note: Due to a clerical error during data preparation, demographic data were available for only

73 of 126 enrolled participants (57.9 %). Of the total enrolled sample, 87 participants (69.0%)

completed the one-week follow-up survey. Similar distributions of demographic characteristics

between completers and non-completers suggest random attrition, though this conclusion is

limited by incomplete demographic data across the full sample.

Recruitment

Participants were recruited for this study through newsletters sent to individuals with prior

engagement in our previous research studies and targeted advertising on Facebook and

Instagram, aimed at adults residing within a 50-mile radius of Santa Monica, California.

Participants were compensated $30 per hour via cash or Venmo payments, and parking fees were

validated. The total participation duration, measured from arrival to departure, was rounded up to

the nearest 15-minute increment for compensation purposes. Recruitment began on 2023 April

24 and ended on 2024 March 22.

Eligibility

Inclusion criteria for study participation were as follows: (1) absence of neurological conditions

including epilepsy and migraine; (2) no history of claustrophobia; (3) absence of photosensitivity

or ophthalmological conditions such as cataracts, corneal abrasions, keratitis, or uveitis; (4)

normal or corrected-to-normal auditory and visual acuity; (5) no current use of medications

known to induce photophobia or alter auditory perception; and (6) sufficient mobility to enter the

MindGym apparatus without assistive devices (e.g., wheelchairs, walkers, or canes).

Prior to enrollment, potential participants underwent screening via a standardized online

questionnaire administered through the Google Forms platform. This pre-screening process

ensured adherence to all inclusion and exclusion criteria before formal study initiation.

Materials

142 All participants sat on an OMEGA Gaming Chair (SecretLab, Inc.) and completed the behavioral

143 questionnaires (administered through Google Forms) and cognitive tasks on a 27" 2022 iMac

144 (Apple, Inc.) regardless of group.

Physiological monitoring

The CGX Aim II Physiological Monitoring device (CGX, Inc.), using ECG electrodes (Skintact Inc.), measured the following bio peripherals: Electromyography (EMG; 2 electrodes on the L/R base of the neck on the sternocleidomastoid muscle), Bio-Impedance-Based Respiration Rate (2 paddles with 2 electrodes each on the L/R pectoralis major), Heart Rate, and Galvanic Skin Response (GSR; 2 electrodes on the palm of the non-dominant hand).

Lab Recorder

Lab Streaming Layer with LabRecorder [6,10,11] was utilized to temporally synchronize our peripheral and experimental time series (experience start and end timestamps) within an XDF file format.

Trait measures

a. Dispositional Positive Emotion Scale (D-PES;[53]) – The Awe Subscale of D-PES measures an individual's dispositional propensity to feel awe toward the world, consisting of 6 items.

b. Modified Tellegen Absorption Scale (MODTAS; [54]) – Adapted from the Tellegen Absorption Scale (Tellegen & Atkinson, 1974), uses a Likert scale to assess the imaginative involvement and the tendency to become mentally absorbed in everyday activities.

169   c. NEO Five-Factor Inventory (NEO-FFI-3; [55]) – Measures five broad personality

170   dimensions – neuroticism, extraversion, openness to experience, agreeableness, and

171   conscientiousness.

173 **State measures**

174
a. State-Trait Anxiety Inventory (STAI;[56]) – Assesses state anxiety

(temporary/situational) and trait anxiety (enduring propensity).

b. Profile of Mood States 2nd Edition (POMS 2; [57]) – Evaluates current mood state

affective dimensions like anger, confusion, depression, fatigue, tension, and vigor.

c. Valence and Arousal – Measures emotional state on two Likert scales from 0 to 10:

Valence (0= very unpleasant,10 = very pleasant) and Arousal (0 = totally calm, 10 =

totally excited/agitated).

Outcome measures

184
a. Toronto Mindfulness Scale (TMS; [58]) – A 13-item administered post-meditation to

assess state mindfulness, while differentiating between reflective awareness and

ruminative attention. The TMS contains two subscales: Curiosity, gauging the interest in

one's experiences, and Decentering, reflecting the ability to view thoughts and feelings as

transient mental events.

b. Awe Scale (Awe-S; [59]) – An 8-item self-report scale measuring individual differences

in the tendency to experience awe, using a 7-point Likert scale to rate their agreement

with statements regarding feelings of awe and perceptions of vastness.

c. Body Size Estimation (BSE; [60]) – Assesses body image perception, from a selection of

nine progressive figural drawings that depict varying body sizes to estimate their own

perceived body size, both in height and width. This measure has been linked to the

intensity of awe and immersion in non-ordinary experiences [61].

d. Ego-Dissolution Inventory (EDI; [62]) – Consists of 16 items assessing altered states of

ego-consciousness, using a visual analog scale ranging from 0% to 100%. It includes 8

items relating to ego-dissolution, representing the core of the inventory, and 8 to ego-

inflation, contrasting experience of heightened self-assuredness.

e. Igroup Presence Questionnaire (IPQ; [63]) – Measures sense of presence, immersion,

engagement, and connection in virtual environments.

f. Modified Immersion Experience Questionnaire (adapted per [64]) – Measures depth and

quality of immersion/engagement in a particular context or virtual environment.

g. Motion Sickness Assessment Questionnaire [65] – A 16-item scale assessing multiple

dimensions of motion sickness (gastrointestinal, central, peripheral, and sopite-related

symptoms).

209 Behavioral tasks

210   a. Trail-Making Test (TMT; [66]) – A neuropsychological assessment, taken on the

211   computer, evaluating visual attention, scanning, processing speed, and cognitive

212   flexibility. Participants were instructed to connect a series of 25 dots in sequential order,

213   alternating between numbers and letters (1-A-2-B, and so on), aiming for speed and

214   accuracy.

215   b. Architex [67] – A validated, computerized neuropsychological assessment tool designed

216     to objectively measure multiple cognitive domains and their interactions, including

217     attention, working memory, processing speed, affect recognition, planning, organization,

218     impulsivity, and verbal and nonverbal abstract reasoning.

220 Content delivery systems

221 The first immersive content-delivery system was MetaQuest 2 (Meta Platforms, Inc.) VR

222 headset, equipped with an Elite Strap (see Fig 1A), presenting stimuli through its native video

223 player at a resolution of 5376x2688, scaled to the maximum allowed resolution of 1920x3664,

224 and a refresh rate of 120Hz. All videos were captured using a 360 camera (GoPro Max 360)

225 placed inside the MindGym during the programmed experience, with the audio content

226 subsequently overlaid. During the experiment, the VR is set to cast through an iPad Air (iPadOS

227 16.6.1) and navigated using a wired mouse (Razer Inc.).

228 The second system was the MindGym (Lumena, Inc.), a 7' isotropic MindGym lined with

229 reflective mylar on its interior walls, while its floor and ceiling are equipped with mirrors. (see

230 Appendix S1). The MindGym incorporates WS2815 LEDs, with 121 pixels per edge between the

231 vertices, totaling 1452 pixels across 12 edges. It utilizes SMD5050 RGB LED chip, offering

232 RGB color with 256 grayscale levels and an output of 990-1080 lumens per meter. The LEDs are

233 operated at less than half of their maximum capacity. Additionally, the system features a color

234 temperature of 5500K. Participants in the MindGym group were provided over-ear wireless,

235 noise-canceling headphones (Sony Group Corp.) to wear during the experience (see Fig 1B).

236
237       **Fig. 1. Participant setup. (A)** VR; **(B)** MindGym

Stimuli

The first piece of content was "4-8 Breathwork" (see Fig 2A, Fig 2C), led by Dr. Jannell

MacAulay's guided audio and utilizing two primary lighting setups. The beginning featured the

"Mindful Minute" mindfulness exercise with multiple colors and slowly upward-moving LEDs,

designed to encourage mind wandering among participants. MacAulay then introduced the

breathwork technique and its underlying rationale, aided by "Galaxy" visuals of blue LEDS,

around 6000K, embedded in the walls, giving the sensation of being surrounded by stars. The

guided breathwork audio was complemented by upward and downward cascades of vertically

aligned LEDs, synchronized with the inhalation and exhalation periods (see Appendix S2).

The second piece of content was "Rain" (see Fig 2B, Fig 2D), an immersive environment created

for reflection. The audio begins with complete darkness as the sound of rain starts to trickle in.

This is complemented by a downward cascade of vertically aligned blue LEDs, around 6500K,

against a black background, simulating rain visuals falling from each vertex. The participant

remains immersed in the light and sound environment for the remainder of the reflective

experience.

Fig 2. Screenshot of stimuli. (A) MindGym (Breathwork); **(B)** MindGym (Rain); **(C)** VR
(Breathwork); **(D)** VR (Rain)

Procedure

Overview

Before each session, all participants provided signed consent forms. They began the session by

filling out pre-experience questionnaires and completing cognitive tasks, followed by CGX

264 setup. Subsequently, each group underwent a 10-minute experiential phase, after which they

265 completed post-experience questionnaires and tasks (see Fig 3).

266
267 **Fig 3. Procedure overview.** The total participation time was roughly 80 minutes.
268

Randomization

Subjects were iteratively assigned to one of two groups using a MATLAB-based randomization

algorithm that maintained gender balance, with recruitment ceasing for each gender category

upon reaching the predetermined quota. The experimenter remained blind to group assignment

until launching the Python script, which initiated the group-appropriate content delivery system

and stimuli by entering the participant's ID number. Participants were randomly assigned to one

of four content delivery systems and stimuli pairings: (1) MindGym (Breathwork), (2) MindGym

(Rain), (3) VR (Breathwork), and (4) VR (Rain).

Procedure

Upon arrival, participants were seated on an office chair facing a 35.5" x 55" desk and given an

overview of the session (pre-questionnaires, equipment setup, 10-minute experience, post-

questionnaires). They were informed their "experience" would involve sitting for 10 minutes in

an immersive audiovisual environment.

For approximately the first 30 minutes, participants used the iMac to complete a set of

questionnaires (NEO-FFI-3, D-PES, MODTAS, STAI, POMS-2, Valence, and Arousal) on

Google Forms, followed by a brief neurophysiological assessment via the TMT and Architex.

Participants were instructed to silence their phones and remove jewelry and bulky items for

comfort, before being told which experiential group they were assigned to. Using a cotton pad

and 91% isopropyl alcohol, the experimenter cleansed the relevant areas (see CGX materials

above) before outfitting the participant with biosensors and peripherals.

Participants assigned to VR sat in the same room, with the room shades lowered. The VR

headset was positioned and adjusted with the tension dial on the Elite strap for clear vision and

comfort. The experimenter cast the VR to the iPad and adjusted the settings, such that the audio

was played through the headset. Upon connecting to Lab Recorder, the stimuli are launched

concurrently with the initiation of the LSL trigger, by the Python script.

If assigned to the MindGym, participants entered the MindGym and sat centrally, while holding

the CGX device and wearing headphones. The Lab Recorder was connected, and the experience

began within 15 seconds of closing the MindGym door.

Across groups, participants were told to sit still and relax during the 10-minute experience

without falling asleep. Participants were shown how to remove the VR or exit the MindGym if

they decided to end the experience early, otherwise, the experimenter would return to assist them

once the experience ended.

After the experience, biosensors were removed and participants used the restroom if needed

before completing post-experience behavioral assessments (STAI, POMS-2, Valence, Arousal,

IPQ, MSAQ, TMS, Awe-S, BSE, and EDI) and cognitive tasks (TMT and Architex). Finally,

participants were compensated and dismissed.

Behavioral statistical analysis

Dependent variables

Seven primary outcome measures collected at both timepoints were used as dependent variables

(DV) in separate models. There were four mood state measures: Anxiety (STAI), Arousal,

Valence, and Total Mood Disturbance (POMS). There were also three cognitive measures: two

that tracked performance on the Architex task (Total Speed and Total Score) and one that tracked

performance on the TMT using the ratio of reaction time divided by accuracy, here called TMT

RTACC, which decreases with increasing performance due to increased accuracy and/or

decreased reaction time).

Eleven DVs were collected at the Post timepoint only, which included Awe, BSE, EDI,

Immersion (based on IEQ), overall Motion, Curiosity and Decentering (both from Toronto), and

four IPQ variables (General, Involvement, Spatial Presence, Experienced Realism).

There were also four physiological variables – Breath Rate, EDA, HR, and HRV – that were

collected throughout the duration of each experience. These measures were averaged or

computed in either two 5-minute timepoints (for HRV), five 2-minute timepoints (for EDA and

Breath Rate), or ten 1-minute timepoints (for HR).

Independent variables

327
The primary independent variables (IV) were Time or Timepoint (Pre, Post) as a categorical,

within-subjects factor, for any variables collected at both timepoints, as well as both Tech

(MindGym, VR) and Stim (Breathwork, Rain) as categorical, between-subjects group factors.

Several continuous or ordinal IVs were tested for potential moderation or mediation of the Tech

and Stim effects. There were eight psychological variables: DPES, Immersion, IPQ (General),

IPQ (Involvement), IPQ (Spatial Presence), IPQ (Experienced Realism), MODTAS, and

Openness. There were seven physiological variables: Motion (Overall), initial breath rate (Breath

Rate, first 2 min), breath rate change (Breath Rate change, last 2 min – first 2 min), initial heart

rate (HR, first 1 min), heart rate change (HR change, last 1 min – first 1 min), initial heart rate

variability (HRV, first 5 min), and change in heart rate variability (HRV change, last 5 min –

first 5 min). The demographic variables of age and sex were also included in all moderation

models to control for their potential effects.

Nonparametric tests

342
For all DVs collected at both timepoints (pre, post), changes over time were tested with

nonparametric t-tests using Wilcoxon signed-rank tests in JASP [68]. Effect sizes were reported

as the matched rank biserial correlation (RBC), an appropriate measure for nonparametric t-tests

which ranges between –1 to 1 and indicates the relative proportion of positive and negative rank

scores while ignoring ties (e.g., RBC = 0.60 means that 60% more subjects showed an increase

instead of a decrease)**.** Timepoint tests were performed separately for each of the four

experimental groups (MindGym (Breathwork), MindGym (Rain), VR (Breathwork), VR (Rain))

as well as the combined MindGym and VR groups, the combined Breathwork and Rain groups,

and all groups combined.

For all DVs, differences between groups were tested with nonparametric Mann-Whitney U tests

in JASP, which included RBC effect sizes. Six separate group comparisons were tested:

MindGym (Breathwork) vs MindGym (Rain), VR (Breathwork) vs VR (Rain), MindGym

(Breathwork) vs VR (Breathwork), MindGym (Rain) vs VR (Rain), MindGym vs VR, and Rain

vs Breathwork. For any DV collected at both timepoints, the difference score (Post – Pre) was

used to compare group differences in changes over time. For the physiological measures, the first

timepoint, last timepoint, and the difference between first and last timepoints were separately

used as DVs to compare group differences in initial, final, or change in physiological response.

Moderation

Moderation of the experimental effects on mood or cognition was tested as interactions between

each covariate and the Tech and Stim group factors in separate models for the three-way

interaction (Tech x Stim x Moderator) and two-way interactions (Tech x Moderator, Stim x

Moderator), as well as a separate model with only main effects. Demographic covariates (age

and sex) were also included as main effects in all models. For any outcome measure collected at

both timepoints, the difference score (Post – Pre) was used as the DV. Because we did not have

any a priori hypotheses about differential moderation effects between nested levels of Tech and

Stim, we conducted post-hoc t-tests for only the interaction terms that were deemed significant.

Moderations were analyzed with generalized linear models (GLM), using the *glmmTMB* [69] and

emtrends [70] packages in RStudio [71]. Based on extensive model diagnostics, which were

conducted with the *DHARMa* [72] package, it was determined that either gaussian or t-family

distributions (both with the identity link) were optimal for model fitting to ensure no substantial

violations or problems with linearity, independence of errors, homoscedasticity, dispersion, zero-

inflation, outliers, or within-group normality of residuals.

Separate models were tested for the hypothesized moderators, which included eight

psychological variables (DPES, Immersion, IPQ General, IPQ Involvement, IPQ Spatial

Presence, IPQ Experienced Realism, MODTAS, and Openness) and seven physiological

variables (overall motion, initial Breath Rate, Breath Rate change, initial HR, HR change, initial

HRV, HRV change). All moderation models were tested separately for 10 outcome variables as

DVs: Arousal, Anxiety (STAI), Awe, BSE, EDI, Total Mood Disturbance (POMS), Valence,

Architex (Total Score), Architex (Total Speed), and TMT (RTACC).

Repeated measures ANOVA

389
For the physiological measures collected during the experiences, repeated measures ANOVAs

(rmANOVA) were performed in JASP to assess changes over time as a main effect of Timepoint

(two timepoints for HRV, five timepoints for Breath Rate and EDA, ten timepoints for HR) and

to assess experimental effects on changes over time, with separate three-way (Tech x Stim x

Timepoint) or two-way interactions (Tech x Timepoint, Stim x Timepoint). Levene's test was

used to ensure no violations of homogeneity. The assumption of sphericity was tested (using

Mauchly's test) and corrected (with Greenhouse-Geisser) if necessary. Effect sizes for all main

or interaction effects were reported as generalized eta-squared, which appropriately adjusts for

mixed design with within-subject (Time) and between-subjects (Tech, Stim) factors. Effect sizes

for post hoc t-tests, if an interaction was significant, were reported as Cohen's *d*.

Follow up Analyses

402
Three outcome variables - STAI, POMS (Total Mood Disturbance), and a discrete version (10

total responses ranging from 'not at all anxious' to 'extremely anxious') of the generalized

anxiety visual analog scale (GA-DS) – were collected at the 1-week follow-up timepoint. For the

two variables (STAI and POMS Total Mood Disturbance) collected at all three timepoints (pre,

post, and follow-up), rmANOVAs were performed in JASP to assess changes over time as a

main effect of Timepoint (pre, post, follow-up) and to assess experimental effects on changes

over time, with separate three-way (Tech x Stim x Timepoint) or two-way interactions (Tech x

Timepoint, Stim x Timepoint). One variable (GA-DS) was collected only at the follow-up

timepoint, so an ANOVA was performed to compare groups. Levene's test was used to ensure

no violations of homogeneity. The assumption of sphericity (for models including Timepoint)

was tested (using Mauchly's test) and corrected (with Greenhouse-Geisser) if necessary. Effect

sizes for all main or interaction effects were reported as generalized eta-squared (for models

including Timepoint) or partial eta-squared (for models without Timepoint). Effect sizes for any

relevant post hoc t-tests were reported as Cohen's *d*.

Multiple comparison correction

We used the false discovery rate (FDR) method of correcting *P* values for multiple comparisons

(significance threshold chosen as *a*=.05, *P*<.05) to control Type I and Type II errors [73,74].

FDR correction was performed across DVs and separately for 1) the nonparametric tests of

timepoint effects for each group (single or combined), 2) the nonparametric tests of group

differences, separately for each group comparison and separately for the measures collected at

both timepoints, the measures collected at only the Post timepoint, and the physiological

measures collected during the experience, as well as 3) the moderation tests, separately for each

interaction or main effect. FDR correction of the rmANOVAs was performed across the different

models testing interaction or main effects and separately for each measure.

Mediation

432 Mediation analysis was conducted in JASP based on the *lavaan* [75] package in R. Missing

433 values were excluded listwise. Separate models were tested for each potential mediator, which

434 included the same variables that were used as potential moderators in the moderation analysis,

435 and for each DV that was also used in the moderation analysis.

Neurophysiological analysis

Data acquisition and preprocessing

440 Physiological data, including electrocardiogram (ECG), electrodermal activity (EDA),

441 electromyography (EMG), and respiration, were recorded using the CGX AIM Phys. Mon. AIM-

442 0106 device. The data were stored in XDF format, which is a standardized format for multi-

443 channel time series data [76]. The XDF files were loaded using the pyxdf library, and the

444 relevant data streams were extracted based on their names and labels.

ECG processing

The ECG data were first sanitized to remove any invalid or missing data points [77]. A notch

filter was then applied to remove powerline interference at 60 Hz. The ECG signal was further

cleaned by applying a bandpass filter with a low cut-off frequency of 0.5 Hz and a high cut-off

frequency of 45 Hz using a 4th-order Butterworth filter [78]. The filtered ECG data were then

filtered again to remove any values exceeding a predefined amplitude threshold of 1500 mV.

Baseline wandering, a low-frequency artifact in the ECG signal, was computed by measuring the

peak-to-peak amplitude of the ECG signal. If the baseline wandering exceeded a predefined

threshold of 1700 mV, the ECG data were considered unacceptable for further analysis.

EDA processing

The EDA data were processed in conjunction with the EMG data to identify and remove motion

artifacts. First, the EMG signal was normalized. A sliding window approach was then used to

identify windows of EMG data that contained a high proportion of outliers, which were

considered to be indicative of motion artifacts. The Local Outlier Factor (LOF) algorithm [79]

was used to detect outliers within each window (implemented using the scikit-learn library) [80].

The EDA signal was smoothed using a median filter with a kernel size of 8 seconds. A low-pass

Butterworth filter with a cut-off frequency of 5 Hz was then applied to remove high-frequency

noise. The contaminated windows identified in the EMG signal were used to guide the

interpolation of the EDA signal. Linear interpolation was performed to replace the artifact-

contaminated segments of the EDA signal with values interpolated from the neighboring clean

segments.

The quality of the cleaned EDA signal was assessed by checking for values outside the

acceptable range (0.05 - 60 μS) and for rapid changes exceeding ±10 μS/sec within a 1-second

interval [81]. The proportion of data points violating these criteria was used to compute a quality

index ranging from 0 to 100, with higher values indicating better signal quality. If the quality

index was below 50, the EDA data were considered unacceptable for further analysis.

Respiration processing

The respiration data were preprocessed following the methods of [82]. We first applied a high-

pass Butterworth filter with a cut-off frequency of 0.1 Hz to remove low-frequency drift. The

filtered signal was then resampled to a fixed sampling rate of 100 Hz. A sliding window

approach was used to estimate the respiration period within each window using the Fast Fourier

Transform (FFT). The estimated respiration period was used to guide the application of a moving

average filter to smooth the respiration signal within each window. The smoothed signal was

then resampled back to the original sampling rate.

Inhalation and exhalation onsets were detected by identifying the intercepts between the

smoothed respiration signal and its moving mean. The intercepts were then classified as

inhalation or exhalation onsets based on the direction of the slope at each intercept. A minimum

amplitude threshold of 0.25 was used to filter out small fluctuations that do not correspond to

true respiratory events.

The number of breaths within the first and last 5 minutes of the experiment was computed using

the inhalation and exhalation onsets to define individual breaths.

Heart rate variability analysis

Heart rate variability (HRV) analysis was performed by computing the low-frequency (LF) and

high-frequency (HF) components of the ECG signal using the Welch's method for power spectral

density estimation [83]. The LF component was defined as the power in the frequency range of

0.04-0.15 Hz, while the HF component was defined as the power in the range of 0.15-0.4 Hz.

The ratio of LF to HF power (LF/HF ratio) was computed as a measure of sympathovagal

balance [84]. The LF/HF ratio was computed separately for the first and last 5 minutes of the

experiment.

EDA tonic component analysis

The tonic component of the EDA signal, which reflects the slow-varying level of skin

conductance, was extracted by applying a low-pass filter to the EDA signal [77]. The mean value

of the tonic component was computed within non-overlapping 2-minute intervals. These mean

values were used as a measure of the overall level of arousal throughout the experiment.

Ethics

IRB

The Advarra (Columbia, MD) Institutional Review Board approved all recruitment and testing

procedures before initiating enrollment (Pro00070581). Following ethical standards and legal

requirements, all study participants provided written informed consent on a document, hosted

through DropboxSign. They were afforded ample time to seek clarification from the Principal

Investigator and study personnel. The informed consent process incorporated the California

Experimental Research Bill of Rights as mandated by Health and Safety Code Section 24172.

Additionally, the study adhered to the principles outlined in the Declaration of Helsinki.

Furthermore, all laboratory personnel possessed and maintained up-to-date certifications in Good

Clinical Practice and the Protection of Human Research Participants through online training.

Results

Outcome measures as a function of group

No statistically significant differences were observed between the group comparisons for the

post-intervention measures (all *p.fdr*>.05) (see S3 Tables A-F). The closest result to reaching

statistical significance with a notable effect size was found for IPQ (Involvement) when

comparing Breathwork to Rain (*W*=2501.50, *p.fdr*=.12, *RBC*=.26), suggesting a potential trend

towards higher involvement in the Breathwork condition compared to the Rain condition.

Pre-post changes as a function of group

Timepoint effects within group

Analysis of pre-post changes revealed significant improvements across various measures when

considering all groups combined and within specific conditions (see Table 4 and S1 Tables A-I

for detailed results).

Table 4. Summary of results from the nonparametric (Wilcoxon Signed-Rank) t-tests of

timepoint (pre vs post) differences for the different groups or group combinations.

	MindGym (Breathwork)			MindGym (Rain)			VR (Breathwork)			VR (Rain)		
DV	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig
TMT (RTACC)	2.68	0.45	~	-42.64	0.66	***	-43.21	0.33		-35.42	0.33	
Architex (Total Speed)	-51.77	0.70	***	-72.88	0.94	***	-114.97	0.88	***	-69.26	0.87	***
Architex (Total Score)	-0.93	0.37		0.65	-0.34		-0.37	0.25		-0.96	0.22	
Anxiety (STAI)	-3.68	0.49	~	-4.09	0.63	**	-5.70	0.74	***	-1.43	0.16	
Arousal	-0.19	0.13		-0.38	0.11		0.70	-0.29		-0.63	0.28	
Valence	1.03	-0.51	~	0.41	-0.45	~	-0.12	-0.07		0.03	-0.03	
POMS	-9.90	0.63	*	-10.50	0.78	***	-10.58	0.64	**	-7.17	0.31	

	Breathwork			Rain			MindGym			VR			All		
DV	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig
TMT (RTACC)	-21.6	0.391	*	-39.1	0.51	***	-20.35	0.56	***	-39.9	0.33	*	-30.4	0.45	***
Architex (Total Speed)	-81.7	0.791	***	-71.1	0.9	***	-62.33	0.82	***	-91.7	0.87	***	-76.4	0.84	***

DV	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig	ΔM	RBC	sig
Architex (Total Score)	-0.67	0.312	~	-0.13	-0		-0.141	0.06		-0.67	0.23		-0.4	0.14	
Anxiety (STAI)	-4.72	0.621	***	-2.81	0.4	*	-3.889	0.55	***	-3.67	0.47	**	-3.78	0.52	***
Arousal	0.266	-0.12		-0.5	0.22		-0.285	0.12		0.063	0.01		-0.11	0.06	
Valence	0.437	-0.26		0.226	-0.2		0.715	-0.5	**	-0.05	-0.1		0.333	-0.3	*
POMS	-10.3	0.636	***	-8.89	0.59	***	-10.21	0.71	***	-8.95	0.5	***	-9.58	0.61	***

‘DV’ refers to the dependent variable. ‘ΔM’ refers to the mean difference (post – pre). ‘RBC’

refers to rank biserial correlation as a measure of effect size (ranging from -1 to 1) and reflects

the relative proportions of positive or negative ranks. ‘sig’ refers to the FDR-corrected *P*-value

where ‘~’ indicates .05<*p.fdr*<.10, ‘*’ indicates *p.fdr*<.05, ‘**’ indicates *p.fdr*<.01, and ‘***’

indicates *p.fdr*<.001.

When considering all groups combined, there was evidence of a decrease in TMT (RTACC)

(*W*=5443.00, *Z*=4.32, *p.fdr*<.001, *RBC*=.45), Architex (Total Speed) (*W*=6044.00, *Z*=7.82,

p.fdr<.001, *RBC*=.84), Anxiety (STAI) (*W*=5156.50, *Z*=4.86, *p.fdr*<.001, *RBC* = .52), Valence

(*W*=1626.00, *Z*=-2.14, *p.fdr*=.042, *RBC*=-.26), and POMS (*W*=5659.00, *Z*=5.77, *p.fdr*<.001,

RBC=.61)

Examining specific conditions, the MindGym (Breathwork) group demonstrated improvements

in Architex (Total Speed) (*W*=396.00, *Z*=3.36, *p.fdr*<.001, *RBC*=.70) and POMS (*W*=331.00,

Z=2.92, *p.fdr*=.014, *RBC*=.63). The MindGym (Rain) condition showed enhancements for TMT

(RTACC) (*W*=411.00, *Z*=3.19, *p.fdr*<.001, *RBC*=.66), Architex (Total Speed) (*W*=421.00,

Z=4.40, *p.fdr*<.001, *RBC*=.94), STAI (*W*=353.50, *Z*=2.94, *p.fdr*=.005, *RBC*=.63), and POMS

(*W*=442.00, *Z*=3.80, *p.fdr*<.001, *RBC*=.78)

The VR (Breathwork) group exhibited improvements in Architex (Total Speed) (*W*=355.00,

Z=3.99, *p.fdr*<.001, *RBC*=.88), STAI (*W*=405.00, *Z*=3.55, *p.fdr*<.001, *RBC*=.74), and POMS

(*W*=406.00, *Z*=3.10, *p.fdr*=.005, *RBC*=.64) (see S1 Table C), while the VR (Rain) condition only

showed a significant improvement in Architex (Total Speed) (W=379.00, Z=4.01, *p.fdr*<.001,

RBC=.87)

When comparing the combined Breathwork and Rain conditions, both showed significant

improvements across multiple measures, with slight variations in effect sizes. Similarly, the

combined MindGym and VR conditions significantly enhanced cognitive performance, anxiety

reduction, and mood improvement.

Group differences in timepoint effects

Analysis of group differences in timepoint effects revealed no statistically significant differences

between experimental conditions after FDR correction for multiple comparisons (all *p.fdr*>.05)

(see S2 Tables A-F). While all interventions yielded positive outcomes across various measures,

including improvements in cognitive performance, anxiety levels, and mood, the magnitude of

these changes did not differ significantly between experimental conditions when accounting for

multiple comparisons. This lack of significant between-group differences indicates that no single

condition demonstrated superior efficacy over the others in terms of the measured variables.

Physiology measures

Group differences

Analysis of physiological measures revealed significant differences between experimental

conditions, particularly in breath rate patterns, the most notable being between the Breathwork

and Rain conditions (see S4 Tables A-E for all results).

At the end of the session, Breathwork participants had significantly lower final Breath Rate

compared to Rain (W=269.50, *p.fdr*<.001, *RBC*=-.70). The overall change in Breath Rate from

beginning to end of the experience was significantly different between the two conditions

(W=198.50, *p.fdr*<.001, *RBC*=-.78), indicating a more substantial decrease in breath rate for the

Breathwork group.

Similar patterns were observed when comparing the MindGym (Breathwork) and MindGym

(Rain) conditions. MindGym (Breathwork) had a significantly lower final Breath Rate (*W*

=77.00, *p.fdr*<.001, *RBC*=-.68) compared to MindGym (Rain), and the overall change in Breath

Rate showed greater decrease, with a significantly different Breath Rate (*W*=68.50, *p.fdr*<.001,

RBC=-.72) with MindGym (Breathwork).

VR (Breathwork) and VR (Rain) showed significant differences in breath rate patterns, with

significantly lower Breath Rate at the end of the experience (*W*=55.00, *p.fdr*<.001, *RBC*=-.73)

compared to VR (Rain). The overall change in Breath Rate was significantly different (*W*=37.00,

p.fdr<.001, *RBC*= -.82), with VR (Breathwork) demonstrating a more substantial decrease.

These results consistently demonstrate that participants in the Breathwork conditions, regardless

of the delivery method (MindGym or VR), experienced a decrease in breath rate throughout the

session compared to those in the Rain conditions. This pattern suggests that the Breathwork

interventions effectively modulated participants' breathing patterns, potentially leading to a more

relaxed physiological state by the end of the session.

No other physiological measures (EDA, HR, HRV) showed significant differences between

conditions after FDR correction.

Changes over time

We employed repeated measures ANOVA (rmANOVA) to examine changes over time and the

potential effects of experimental conditions. We observed significant temporal changes in HR,

Breath Rate, and EDA levels, with only Breath Rate showing interactions between time and

technology. No significant main effects of time or interactions with groups were observed for

HRV after FDR correction.

The results revealed a significant main effect of time on HR ($F_{5.17, 398.38}$=8.78, $p.fdr$=.004),

indicating a gradual increase throughout the experiment (see Table 5). EDA levels revealed a

significant main effect of time ($F_{2.35, 162.37}$=17.11, $p.fdr$=.004), characterized by an initial increase

followed by a plateau around the third timepoint.

Breath Rate showed a significant main effect of time ($F_{2.02, 171.33}$=25.31, $p.fdr$=.002),

characterized by an initial decrease followed by a plateau around the third timepoint. A

significant interaction between time and Stimuli was also observed ($F_{2.48, 208.49}$=28.49,

$p.fdr$=.002). Post hoc paired-sample t-tests (with Holm correction of P values) were conducted

between timepoints within Stimuli groups. Within the Rain groups combined, breath rate did not

reliably change between any timepoints (all $p.holm$>.05). Within the Breathwork groups

combined, breath rate significantly decreased from the first to second timepoint (ΔM=-3.55,

SE=.49, t=7.19, $p.holm$<.001, d=1.08), remained similar between second and third timepoints

(ΔM=-1.12, SE=0.49, t=2.26, $p.holm$=.416, d=.034), decreased again from third to fourth

timepoints (ΔM=-2.28, SE=.49, t = 4.62, $p.holm$ <.001, d=.69), without any further reliable

decrease from fourth to fifth timepoints (ΔM=.77, SE=.49, t=1.56, $p.holm$=1.000, d=.23).

Table 5. Results from repeated measures ANOVA on physio measures.

	Timepoint						Timepoint x Tech					
DV	df1	df2	F	p.raw	p.fdr	η^2_G	df1	df2	F	p.raw	p.fdr	η^2_G
HR	5.17	398.38	8.78	< .001	0.004	0.005	5.14	390.59	0.56	0.738	0.777	0.003
HRV	1.00	77.00	1.54	0.219	0.438	0.009	1.00	76.00	0.31	0.581	0.615	0.002
EDA	2.35	162.37	17.11	< .001	0.004	0.020	2.35	159.51	0.92	0.414	0.552	0.001

	df1	df2	F	p.raw	p.fdr	η²G	df1	df2	F	p.raw	p.fdr	η²G
BR	2.02	171.33	25.31	< .001	0.002	0.130	2.02	169.50	0.38	0.685	0.867	0.002

	Timepoint x Stim						**Timepoint x Tech x Stim**					
DV	df1	df2	F	p.raw	p.fdr	η²G	df1	df2	F	p.raw	p.fdr	η²G
HR	5.16	392.00	0.84	0.527	0.777	0.005	5.09	376.86	0.50	0.777	0.777	0.003
HRV	1.00	76.00	2.67	0.106	0.424	0.015	1.00	74.00	0.26	0.615	0.615	0.002
EDA	2.37	161.14	1.90	0.146	0.292	0.003	2.36	155.43	0.18	0.871	0.871	0.002
BR	2.48	208.49	28.49	< .001	0.002	0.140	2.48	203.01	0.19	0.867	0.867	0.001

DV: dependent variable; HR = heart rate; HRV = heart rate variability; EDA = electrodermal activity (tonic); BR = breath rate. Timepoint = main effect model (within-subjects); Timepoint x Tech = interaction model (between-groups: MindGym vs VR); Timepoint x Stim = interaction model (between-groups: Breathwork vs Rain); Timepoint x Tech x Stim = three-way interaction model. F = F-test with df1, df2 (degrees of freedom); P.RAW/P.FDR = uncorrected/FDR-corrected p-values.

Mediation

Mediation analyses were conducted to test if hypothesized variables mediated the group effects on the outcome scores. No mediation analyses passed the threshold for significance after FDR correction (all *p.fdr*>.05).

Moderation

Our moderation analysis revealed significant effects of both psychological and physiological factors on participants' experiences and performance during the immersive interventions (see S5 Tables A-J for detailed results). Among the psychological trait measures, MODTAS emerged as a significant predictor of both Awe (*b*=.37, *SE*=.11, *Z*=3.28, *p.fdr*=.004) (see Table 6) and EDI (*b*=.03, *SE*=.01, *Z*=2.98, *p.fdr*=.015), with higher MODTAS scores associated with greater reported Awe and EDI. DPES also showed a significant main effect on Awe (*b*=13.89, *SE*=3.48,

639   *Z*=4.00, *p.fdr*<.001), indicating that individuals with higher dispositional positive emotions were

640   more likely to experience awe during the intervention.

641   **Table 6. Results from GLM tests of moderation.**

DV	Tech x Stim x Moderator			Tech x Moderator			Stim x Moderator			Moderator				
Awe	F	p.raw	p.fdr	F	p.raw	p.fdr	F	p.raw	p.fdr	b	SE	Z	p.raw	p.fdr
Psych Moderator														
DPES	1.74	0.190	0.824	0.14	0.711	0.887	0.20	0.657	0.957	13.89	3.48	4.00	< .0001	< .001
Immersion	0.03	0.870	0.892	0.08	0.776	0.887	0.02	0.896	0.957	1.16	0.22	5.17	< .0001	< .001
IPQ (General)	0.02	0.892	0.892	4.70	0.032	0.362	1.92	0.169	0.853	6.18	1.72	3.59	< .001	0.002
IPQ (Involvement)	1.08	0.301	0.892	0.12	0.727	0.887	0.82	0.367	0.853	2.24	0.76	2.96	0.003	0.009
IPQ (Spatial Presence)	0.38	0.538	0.892	0.00	0.957	0.957	0.77	0.381	0.853	1.83	0.93	1.98	0.048	0.104
IPQ (Experienced Realism)	1.52	0.220	0.824	0.34	0.560	0.887	3.98	0.048	0.725	0.68	0.87	0.78	0.436	0.620
MODTAS	0.56	0.458	0.892	0.05	0.828	0.887	0.38	0.539	0.899	0.37	0.11	3.28	0.001	0.004
Openness	0.36	0.553	0.892	0.85	0.359	0.796	0.03	0.871	0.957	1.22	0.81	1.51	0.131	0.218
Physio Moderator														
Motion (overall)	2.69	0.104	0.824	0.14	0.711	0.887	1.41	0.238	0.853	0.79	0.50	1.59	0.112	0.210
Breath Rate (first 2 min)	0.13	0.724	0.892	1.08	0.301	0.796	0.06	0.805	0.957	0.15	1.02	0.15	0.883	0.883
HR (first 1 min)	2.45	0.122	0.824	0.59	0.446	0.837	0.01	0.908	0.957	-0.70	0.29	-2.39	0.017	0.042
HRV (first 5 min)	0.03	0.853	0.892	4.04	0.048	0.362	0.00	0.957	0.957	42.30	65.70	0.64	0.520	0.650
Breath Rate change (last - first)	0.02	0.876	0.892	0.81	0.371	0.796	0.82	0.367	0.853	-0.19	0.82	-0.24	0.814	0.872
HR change (last - first)	0.17	0.686	0.892	2.44	0.123	0.508	0.72	0.398	0.853	-0.60	0.80	-0.75	0.455	0.620
HRV change (last - first)	0.68	0.412	0.892	2.28	0.135	0.508	0.55	0.461	0.865	9.92	38.69	0.26	0.798	0.872

DV (dependent variable) = Awe (post timepoint); Psych Moderator/Physio Moderator =

psychological/physiological moderating variable; Tech x Stim x Moderator = three-way

interaction model (moderator × Tech [MindGym vs VR] × Stimuli [Breathwork vs Rain]); Tech

x Moderator = two-way interaction (moderator × Tech); Stim x Moderator = two-way interaction

(moderator × Stimuli); Moderator = main effects model; F = F-test (GLM); b/SE =

unstandardized beta coefficient/standard error; Z = Z-test; p.raw/p.fdr = uncorrected/FDR-

corrected p-values. The trait of Openness demonstrated an interaction effect with the type of

stimuli on anxiety levels as measured by STAI ($F_{1,111}$=9.58, *p.fdr*=.038). Specifically, higher

levels of openness were associated with a greater reduction in anxiety for participants in

Breathwork, but not for those in Rain.

Psychological state measures also played a significant role in moderating participants'

experiences. Immersion showed strong main effects on both Awe (b=1.16, SE=0.22, Z=5.17,

$p.fdr$<.001) and EDI (b=0.07, SE=0.02, Z=4.55, $p.fdr$< .001), with higher levels of immersion

consistently associated with greater reported awe and emotional depth. IPQ (General) was also

positively associated with Awe (b=6.18, SE=1.72, Z=3.59, $p.fdr$=.002) and EDI (b=0.32,

SE=0.13, Z=2.49, $p.fdr$=.048). Only IPQ (Involvement) followed this same pattern with Awe

(b=2.24, SE=0.76, Z=2.96, $p.fdr$=.009) and EDI (b=0.18, SE=0.05, Z=3.24, $p.fdr$=.009).

Turning to physiological moderators, we found the change in heart rate throughout the

intervention (HR-diff) showed a significant interaction effect with the type of technology used

on Architex (Total Score) (F_{df1}=10, $p.fdr$=.023). Specifically, an increase in heart rate was

associated with decreased accuracy for participants in the MindGym groups, but not for those in

the VR.

Descriptives

Young adults (18-34) showed the highest motion sickness susceptibility (mean = 20.58, SD =

13.74) but the lowest gastric symptoms (mean = 12.95, SD = 5.20), suggesting their intervention

response may have been influenced by motion-related discomfort. Older adults (55+)

experienced the highest levels of gastric (mean = 16.30, SD = 15.74), central (mean = 18.30, SD

= 13.43), and peripheral symptoms (mean = 18.52, SD = 15.74). Males displayed higher central

symptom scores across all ages and more consistent peripheral symptoms, while females showed

greater symptom variability across all categories. Sopite symptoms (drowsiness and fatigue)

were consistent across age groups (mean ≈ 23.0), with higher scores among younger males.

Table 7. Motion sickness scores across demographics and groups

Motion Sickness		Total Score		Gastrointestinal		Central		Peripheral		Sopite-related	
		M	SD	M	SD	M	SD	M	SD	M	SD
Age											
	18-34	20.58	13.74	17.19	14.63	20.38	15.03	18.24	16.43	26.00	17.87
	35-54	15.73	4.80	12.47	4.63	15.45	5.96	14.13	7.15	20.54	9.93
	55+	15.02	3.55	11.57	2.55	14.67	5.27	11.73	1.71	21.39	9.57
Gender											
	Female	17.85	9.36	13.84	8.52	17.35	10.22	15.46	12.33	24.25	14.78
	Male	17.36	10.26	14.64	11.71	17.32	11.65	15.11	11.18	21.83	13.01
Group											
	Cube	18.39	10.37	15.43	11.84	18.52	12.42	16.40	13.63	22.66	11.47
	VR	16.82	9.17	13.05	8.19	16.16	9.11	14.17	9.42	23.41	16.09
	Breath	17.87	10.62	14.45	9.94	17.71	12.23	16.72	12.98	22.35	14.03
	Rain	17.33	8.92	14.02	10.56	16.95	9.46	13.80	10.15	23.75	13.88
	MindGym (Breathwork)	19.00	10.18	15.23	9.13	19.50	13.95	17.32	13.57	23.39	12.08
	MindGym (Rain)	17.80	10.69	15.63	14.12	17.57	10.89	15.51	13.84	21.96	10.98
	VR (Breathwork)	16.81	11.08	13.72	10.73	16.03	10.30	16.16	12.59	21.38	15.77
	VR (Rain)	16.83	6.68	12.31	3.91	16.30	7.77	11.98	2.32	25.65	16.40

677 Age groups are divided into 18-34, 35-54, and 55+ years. M and SD represent the mean and

678 standard deviation

Follow up Analyses

Follow-up analysis results are fully reported in the Supplemental Materials (Tables A-6C; Figure

6A). Both STAI and POMS (Total Mood Disturbance) showed significant changes over time,

without any interactions between groups, such that scores significantly decreased immediately

after the experience (consistent with the nonparametric results reported earlier) but then

significantly increased one week later to a similar level (POMS) or even higher level (STAI)

compared to before the experience. Effect sizes of pairwise group comparisons were small to

moderate (ranging from 0.09 to 0.68). For the measure of anxiety (GA-DS) collected only at the

1-week timepoint, there were no significant main effects of Tech groups ($F(1,85) = 0.31$, $p =$

0.577, partial $\eta^2 = 0.004$), Stim groups ($F(1,85) = 0.11$, $p = 0.738$, partial $\eta^2 = 0.001$), or interaction

between Tech and Stimuli groups ($F(1,83) = 0.88$, $p = 0.352$, partial $\eta^2 = 0.010$),

Discussion

This study aimed to investigate the impact of content delivery systems on two different

anxiolytic stimuli: guided breathwork and naturalistic rain. We compared the efficacy of a novel

"reflective chamber" (MindGym) with a traditional virtual reality (VR) headset in delivering

these interventions. While our initial hypothesis anticipated clear superiority of one technology

over the other, our results instead demonstrated the robust effectiveness of MindGym-generated

content across both delivery systems. This finding highlights the MindGym's potential as a

powerful content generation platform, capable of producing experiences that maintain their

efficacy even when translated to more portable VR formats.

Our findings revealed significant reductions in anxiety measures across all conditions, with the

STAI showing significant improvements in all groups except for the breathwork condition in the

MindGym. These results may be of clinical relevance and are comparable to benchmarks from

other acute anxiety interventions. Our study demonstrated an overall STAI reduction of 11.67%

across all groups, with breathwork showing the highest reduction at 15.10%. These findings

align with recent studies on non-pharmacological interventions for acute anxiety reduction (e.g.,

Johnson et al. [4] reported a 15% reduction in STAI scores in non-anxious populations exposed

to a psychedelic-esque stroboscopic lights and binaural beats experience, and a 13% reduction in

response to breath-focused meditation; Sriboonlert et al. [85] found a 17% reduction in anxiety

scores following aromatherapy interventions;   et al. [86] observed a 7% reduction from exercise

in lower trait anxious individuals and a 17% reduction for those with higher trait anxiety).

Our results suggest that both the MindGym and VR delivery systems, combined with either rain

or breathwork stimuli, can produce anxiety reductions comparable to these established

interventions. The fact that such basic interventions, which do not fully showcase the complex

experiences unique to the MindGym, were equally effective in both systems highlights the

MindGym's potential as a content creation tool. It suggests that even more sophisticated

MindGym-generated experiences could be successfully adapted for VR, potentially enhancing

their accessibility without compromising efficacy. However, it's important to note that these

effects are still considerably lower than the 30-50% reduction typically observed with

pharmacological interventions [87]. This comparison highlights the potential for non-

pharmacological approaches while also acknowledging the current limitations in their efficacy

compared to medication-based treatments. Interestingly, we observed no significant changes in

arousal levels, while valence only improved significantly in the MindGym condition. The

interpretation of pre-post results centered around task performance (e.g., Architex) is challenging

without a control condition, as some improvements may be attributable to practice effects. For

instance, the TMT typically shows improvement with repeated administrations [88,89].

Contrary to our initial expectations, we found no main effects of either stimulus type or

technology on the outcome measures. This lack of differentiation between the MindGym and VR

conditions is particularly noteworthy, as it suggests that the MindGym-generated content

retained its effectiveness when translated to a VR format. While there were numerical trends

suggesting potentially better affective outcomes in the MindGym (e.g., greater reduction in STAI

scores and mood disturbance), these differences did not reach statistical significance. We

observed a consistent pattern of "more" significant effects in the MindGym across most

measures, but we caution against over-interpreting these trends. Surprisingly, we did not observe

the expected differences in immersion between the two technologies. This lack of differentiation

may suggest that both systems were equally effective in creating an immersive experience, or

that our measures were not sensitive enough to detect subtle differences. It also points to the

successful adaptation of MindGym-generated content for VR platforms.

The most notable physiological finding was the decrease in breath rate in the breathwork

condition compared to the rain condition. This result serves as a sanity check, confirming that

participants engaged with the breathwork instructions as intended. However, we found no

significant differences in electrodermal activity (EDA), heart rate (HR), or heart rate variability

(HRV) as a function of technology or stimulus type.

We explored whether individual differences could account for variations in response to the

interventions. Several interesting moderation effects emerged. DPES predicted the amount of

awe experienced, though this finding is somewhat expected given the overlap between DPES and

awe measures. MODTAS, which has previously been shown to predict audiovisual-induced

chills [90], had a main effect on both awe and ego dissolution. MODTAS also moderated the

effects of stimuli on anxiety reduction, with higher trait absorption associated with greater

anxiety reduction in the breathwork condition across both technologies. Openness to experience

moderated the impact on STAI scores, with more open individuals experiencing greater anxiety

reduction in the breathwork condition, but not in the rain condition.

These moderation effects exclusive to breathwork may reflect the importance of explicit

instructions and participants' willingness to fully engage with the experience. The breathwork

condition may have benefited individuals who are more prone to absorption and open to new

experiences, possibly due to its more structured nature compared to the passive rain condition.

Immersion and presence, while not differing between conditions, were potent predictors of awe

and ego dissolution across all groups. This aligns with previous research showing that the degree

of immersion in virtual environments impacts memory encoding and neural representations [41].

Interestingly, motion sickness negatively impacted ego dissolution, suggesting that physical

discomfort may anchor individuals in their bodily sensations, hindering the experience of ego

transcendence.

Contrary to our expectations, young adults (18-34) showed higher motion sickness susceptibility

(mean = 20.58, SD = 13.74) than middle-aged (mean = 15.73, SD = 4.80) and older adults (mean

= 15.02, SD = 3.55). Additionally, MindGym conditions unexpectedly showed slightly higher

motion sickness scores (Breathwork: mean = 19.00, Rain: mean = 17.80) compared to VR

conditions (Breathwork: mean = 16.81, Rain: mean = 16.83). While this appears to contradict

established theories about vestibular mismatch in VR environments (Reggente et al., 2018), it's

important to note that our use of 360° video rather than fully interactive VR may have created a

different type of perceptual experience than typically studied in VR research, as the non-

interactive nature of both conditions could have affected motion sickness in ways not predicted

by traditional vestibular mismatch theory. Males and females showed similar total motion

sickness scores (males: mean = 17.36, females: mean = 17.85), though females exhibited greater

variability across symptom categories. Sopite symptoms remained relatively consistent across

age groups (mean ≈ 23.0), with both conditions showing similar levels (MindGym: mean =

22.66, VR: mean = 23.41), with a slight advantage to the MindGym, making it potentially more

ideal for reliably inducing mood states.

Study retention patterns revealed differences between participants who completed follow-up

assessments and those who did not, particularly across age groups and intervention conditions.

The most notable difference was age, with survey completers being about 8 years older on

average (43.6 vs 35.6 years), while other demographic differences were relatively modest in

comparison. Another group difference was also observed in intervention type, with MindGym

conditions showing notably higher completion rates (76% combined) compared to VR conditions

(58% combined).

In conclusion, this study contributes to the growing body of research on non-pharmacological

interventions for acute anxiety reduction while also demonstrating the potential of the MindGym

as a versatile content creation platform. The comparable efficacy of both the MindGym and VR

systems, using content originally generated for the MindGym, suggests that experiences

designed in the MindGym can be successfully translated to more portable VR formats without

significant loss of effectiveness. This finding is particularly promising given that the

interventions used in this study were relatively basic and did not fully leverage the unique

capabilities of the MindGym. Future research should focus on developing more complex,

immersive experiences using the MindGym's full potential, and exploring how these can be

effectively adapted for VR and other delivery systems. This approach could lead to the

development of highly effective, accessible, and scalable interventions for anxiety management,

potentially bridging the gap with pharmacological treatments while offering the benefits of non-

pharmacological approaches.

Limitations

A significant limitation of this study is that the "VR" condition was not a true virtual reality

experience, but rather a 3D video displayed in a VR headset. A more equitable comparison of the

content delivery systems would have involved creating a virtual environment that mimicked the

MindGym chamber and responded more dynamically head movements, potentially enhancing

immersion and presence and serving as a VR-analog as opposed to a recording of the MindGym

experience. The current design may have inadvertently positioned the MindGym experience as a

"steel man" argument, with many of its immersive properties carried over to the VR condition.

However, this limitation also highlights the potency of the MindGym programming,

demonstrating that its effects can be successfully replicated using a VR device. This finding

suggests that the content displayed in a refelctie chamber environment may be the primary driver

of the observed anxiolytic effects that extend into VR.

The unexpected age and platform differences in motion sickness susceptibility challenge

conventional assumptions about cybersickness in younger populations and digital environments

and may warrant the development of new scales for non-VR-based experiential tehcnologies.

The substantial individual variability in symptoms, particularly among younger adults and across

platforms, represents a study limitation and emphasizes the need for personalized comfort

settings and adaptive monitoring systems in future research designs. Additionally, future studies

should consider implementing both fully interactive VR environments and more detailed

measures of cybersickness and environmental satisfaction to better understand how these factors

might influence intervention effectiveness. Such modifications would help distinguish

intervention effectiveness from cybersickness effects while accounting for unexpected

susceptibility patterns across different types of digital experiences.

Our follow-up analyses revealed that while the interventions produced significant immediate

reductions in anxiety and mood disturbance, these beneficial effects were not maintained one

week post-intervention. Particularly noteworthy was the observation that anxiety levels not only

returned to baseline but exceeded pre-intervention measurements at follow-up, suggesting a

potential rebound effect that warrants further investigation. These findings indicate that while

single-session interventions can provide acute anxiety relief, maintaining long-term benefits may

require regular practice or multiple sessions, especially given the apparent vulnerability to

anxiety rebound effects. However, it's important to note that follow-up measurements were taken

outside the controlled laboratory environment where pre- and post-intervention assessments

occurred, potentially introducing different contextual factors and stressors. Furthermore, without

also having measurements at one week *prior* to the start of the intervention (which would mirror

the one week *post* intervention), it's uncertain how these follow-up levels compare to subjects'

average mood states in general, irrespective of their participation in the experiment. This could

be an interesting question for future studies.

Future Directions

Future research should focus on conducting experiments with more visually diverse content to

identify where these content delivery systems meaningfully diverge. This could involve

developing true VR environments that more closely mimic the MindGym experience, allowing

for a more direct comparison of the technologies. Investigating the impact of these delivery

systems on different types of anxiety (e.g., social anxiety, specific phobias) could determine if

certain technologies are more effective for particular anxiety subtypes. Exploring the long-term

effects of repeated exposures to these interventions would assess their potential as ongoing

anxiety management tools. Examining the role of individual differences (e.g., absorption,

openness) in treatment response could lead to the development of more personalized anxiety

interventions.

The lack of significant differences in these physiological measures across conditions may

suggest that the anxiolytic effects are mediated through cognitive or emotional pathways rather

than direct physiological changes. Therefore, investigating the neural correlates of the anxiolytic

effects observed in this study would better our understanding of the mechanisms underlying the

effectiveness of these interventions. Additionally, this might account for some individual

differences in responses that were not seen by self-reported trait variables.

Future research should explore the potential utility of brief digital relaxation interventions as pre-

session tools in clinical settings. Given the potent, acute anxiolytic effects observed in our study,

these interventions could help standardize patients' arousal states before therapy sessions,

potentially offsetting situational stressors (e.g., commute-related anxiety) that might otherwise

impact session effectiveness. Such application could be particularly valuable in urban clinical

settings where external stressors are common. Investigating whether these interventions can

create a more consistent baseline state for therapeutic engagement would be an important next

step in evaluating their ecological utility in clinical practice.

Conclusion

In conclusion, while our study did not find significant differences between the MindGym and VR delivery systems, it demonstrated that the powerful anxiolytic effects originally discovered in the reflective chamber environment successfully translated to cirtual reality without loss of efficacy. The observed moderation effects highlight the importance of individual differences in treatment response and suggest that tailoring interventions to personal characteristics may enhance their effectiveness. This successful virtual adaptation of an established anxiety-reducing environment opens promising avenues for developing more accessible and effective anxiety management tools.

Acknowledgments

We warmly thank the research participants. We also thank Asli Sara Bilgili who assisted N.S., G.W., and C.L. with data collection. We thank Lumena, Inc. for their troubleshooting assistance throughout the course of the experiment. We also extend gratitude to Leonardo Christov-Moore who provided commentary on the study design in its early phases.

Competing Interests

This research was financially supported under a Research Services agreement executed between IACS and Lumena, Inc. It is imperative to note that the terms of this agreement did not incorporate any provisions or incentives contingent upon the attainment of specific outcomes or successful results but were solely predicated on underwriting the requisite expenses for the genuine and unbiased execution of the study. Lumena, Inc. had no input on the design nor analysis of this study, apart from supplying MindGym's content, delivery system hardware and software, and the 360 videos utilized in the VR.

Author Contributions Statement

Study conceptualization and design: N.R. Study implementation and logistics: N.R., N.S. Data

collection: N.S., G.W., C.L. Data analysis: M.J., V.K., F.S., V.K., N.R., N.S. Manuscript writing

and editing (all drafts): N.S., N.R., G.W., C.L., F.S., M.J. All authors approved the final version

for submission.

References

1. Naef AC, Jeitziner MM, Knobel SEJ, Exl MT, Müri RM, Jakob SM, et al. Investigating the role of auditory and visual sensory inputs for inducing relaxation during virtual reality stimulation. Sci Rep. 2022 Oct 12;12(1):17073.

2. Weller R, Cepok J, Arzaroli R, Marnholz K, Große CS, Reuter H, et al. Effects of immersion and navigation agency in virtual environments on emotions and behavioral intentions. Frontiers in Virtual Reality [Internet]. 2022 [cited 2023 Dec 12];3. Available from: https://www.frontiersin.org/articles/10.3389/frvir.2022.893052

3. Browning MHEM, Shin S, Drong G, McAnirlin O, Gagnon RJ, Ranganathan S, et al. Daily exposure to virtual nature reduces symptoms of anxiety in college students. Sci Rep. 2023 Jan 23;13(1):1239.

4. Johnson M, Simonian N, Reggente N. Lightening the Mind with Audiovisual Stimulation as an Accessible Alternative to Breath-Focused Meditation for Mood and Cognitive Enhancement [Internet]. 2024 [cited 2024 Sep 12]. Available from: https://osf.io/hb7wx

5. Miller N, Stepanova ER, Desnoyers-Stewart J, Adhikari A, Kitson A, Pennefather P, et al. Awedyssey: Design Tensions in Eliciting Self-transcendent Emotions in Virtual Reality to Support Mental Well-being and Connection. In: Proceedings of the 2023 ACM Designing Interactive Systems Conference [Internet]. Pittsburgh PA USA: ACM; 2023 [cited 2024 Jan 7]. p. 189–211. Available from: https://dl.acm.org/doi/10.1145/3563657.3595998

6. Haller H, Cramer H, Lauche R, Gass F, Dobos GJ. The prevalence and burden of subthreshold generalized anxiety disorder: a systematic review. BMC Psychiatry. 2014 May 1;14:128.

7. National Institute of Mental Health (NIMH) - Transforming the understanding and treatment of mental illnesses [Internet]. [cited 2024 Sep 12]. Available from: https://www.nimh.nih.gov/

8. Any Anxiety Disorder - National Institute of Mental Health (NIMH) [Internet]. [cited 2024 Sep 12]. Available from: https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder

9. Hofmann SG, Smits JAJ. Cognitive-behavioral therapy for adult anxiety disorders: a meta-analysis of randomized placebo-controlled trials. J Clin Psychiatry. 2008 Apr;69(4):621–32.

10. Bandelow B, Reitt M, Röver C, Michaelis S, Görlich Y, Wedekind D. Efficacy of treatments for anxiety disorders: a meta-analysis. Int Clin Psychopharmacol. 2015 Jul;30(4):183–92.

11. Crippa JA, Guimarães FS, Campos AC, Zuardi AW. Translational Investigation of the Therapeutic Potential of Cannabidiol (CBD): Toward a New Age. Front Immunol [Internet]. 2018 Sep 21 [cited 2024 Sep 12];9. Available from: https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2018.02009/full

12. George MS, Ward HE, Ninan PT, Pollack M, Nahas Z, Anderson B, et al. A pilot study of
vagus nerve stimulation (VNS) for treatment-resistant anxiety disorders. Brain Stimulation.
2008 Apr 1;1(2):112–21.

13. Kaplan S. The restorative benefits of nature: Toward an integrative framework. Journal of
Environmental Psychology. 1995 Sep 1;15(3):169–82.

14. Maller C, Townsend M, Pryor A, Brown P, St Leger L. Healthy nature healthy people:
"contact with nature" as an upstream health promotion intervention for populations. Health
Promot Int. 2006 Mar;21(1):45–54.

15. Alvarsson JJ, Wiens S, Nilsson ME. Stress Recovery during Exposure to Nature Sound and
Environmental Noise. Int J Environ Res Public Health. 2010 Mar;7(3):1036–46.

16. Bratman GN, Hamilton JP, Daily GC. The impacts of nature experience on human cognitive
function and mental health. Ann N Y Acad Sci. 2012 Feb;1249:118–36.

17. Jo H, Song C, Miyazaki Y. Physiological Benefits of Viewing Nature: A Systematic Review
of Indoor Experiments. Int J Environ Res Public Health. 2019 Dec;16(23):4739.

18. Van Hedger SC, Nusbaum HC, Clohisy L, Jaeggi SM, Buschkuehl M, Berman MG. Of
cricket chirps and car horns: The effect of nature sounds on cognitive performance.
Psychon Bull Rev. 2019 Apr 1;26(2):522–30.

19. Hu C, Zhu K, Huang K, Yu B, Jiang W, Peng K, et al. Using natural intervention to promote
subjective well-being of essential workers during public-health crises: A Study during
COVID-19 pandemic. J Environ Psychol. 2022 Feb;79:101745.

20. Johansson G, Juuso P, Engström Å. Nature-based interventions to promote health for people
with stress-related illness: An integrative review. Scand J Caring Sci. 2022 Dec;36(4):910–
25.

21. Song I, Baek K, Kim C, Song C. Effects of nature sounds on the attention and physiological
and psychological relaxation. Urban Forestry & Urban Greening. 2023 Aug 1;86:127987.

22. Chinagudi S, Badami S, Herur A, Patil S, Gv S, Ankad R. Immediate effect of short duration
of slow deep breathing on heart rate variability in healthy adults. Natl J Physiol Pharm
Pharmacol. 2014;4(3):233.

23. Jerath R, Crawford MW, Barnes VA, Harden K. Self-Regulation of Breathing as a Primary
Treatment for Anxiety. Appl Psychophysiol Biofeedback. 2015 Jun 1;40(2):107–15.

24. Russo MA, Santarelli DM, O'Rourke D. The physiological effects of slow breathing in the
healthy human. Breathe (Sheff). 2017 Dec;13(4):298–309.

25. Zaccaro A, Piarulli A, Laurino M, Garbella E, Menicucci D, Neri B, et al. How Breath-
Control Can Change Your Life: A Systematic Review on Psycho-Physiological Correlates
of Slow Breathing. Front Hum Neurosci. 2018 Sep 7;12:353.

26. Aideyan B, Martin GC, Beeson ET. A Practitioner's Guide to Breathwork in Clinical Mental
Health Counseling. Journal of Mental Health Counseling. 2020 Jan 1;42(1):78–94.

27. Magnon V, Dutheil F, Vallet GT. Benefits from one session of deep and slow breathing on
vagal tone and anxiety in young and older adults. Sci Rep. 2021 Sep 29;11:19267.

28. Balban MY, Neri E, Kogon MM, Weed L, Nouriani B, Jo B, et al. Brief structured
respiration practices enhance mood and reduce physiological arousal. CR Med [Internet].
2023 Jan 10 [cited 2023 Jan 13];0(0). Available from: https://www.cell.com/cell-reports-
medicine/abstract/S2666-3791(22)00474-8

29. World Urbanization Prospects The 2018 Revision.

30. Ulrich RS. Aesthetic and Affective Response to Natural Environment. In: Altman I,
Wohlwill JF, editors. Behavior and the Natural Environment [Internet]. Boston, MA:
Springer US; 1983 [cited 2024 Sep 12]. p. 85–125. Available from:
https://doi.org/10.1007/978-1-4613-3539-9_4

31. Anderson AP, Mayer MD, Fellows AM, Cowan DR, Hegel MT, Buckey JC. Relaxation with
Immersive Natural Scenes Presented Using Virtual Reality. Aerospace Medicine and
Human Performance. 2017 Jun 1;88(6):520–6.

32. Maples-Keller JL, Bunnell BE, Kim SJ, Rothbaum BO. The Use of Virtual Reality
Technology in the Treatment of Anxiety and Other Psychiatric Disorders. Harv Rev
Psychiatry. 2017;25(3):103–13.

33. Schutte NS, Bhullar N, Stilinović EJ, Richardson K. The Impact of Virtual Environments on
Restorativeness and Affect. Ecopsychology. 2017 Mar;9(1):1–7.

34. Jerdan SW, Grindle M, van Woerden HC, Kamel Boulos MN. Head-Mounted Virtual
Reality and Mental Health: Critical Review of Current Research. JMIR Serious Games.
2018 Jul 6;6(3):e14.

35. White MP, Yeo NL, Vassiljev P, Lundstedt R, Wallergård M, Albin M, et al. A prescription
for "nature" – the potential of using virtual nature in therapeutics. Neuropsychiatr Dis
Treat. 2018 Nov 8;14:3001–13.

36. Yu CP, Lee HY, Lu WH, Huang YC, Browning MHEM. Restorative effects of virtual
natural settings on middle-aged and elderly adults. Urban Forestry & Urban Greening. 2020
Dec 1;56:126863.

37. Annerstedt M, Jönsson P, Wallergård M, Johansson G, Karlson B, Grahn P, et al. Inducing
physiological stress recovery with sounds of nature in a virtual reality forest--results from a
pilot study. Physiol Behav. 2013 Jun 13;118:240–50.

38. Proverbio AM, De Benedetto F, Ferrari MV, Ferrarini G. When listening to rain sounds
boosts arithmetic ability. PLoS One. 2018;13(2):e0192296.

39. Villani D, Riva F, Riva G. New technologies for relaxation: The role of presence.
International Journal of Stress Management. 2007;14(3):260–74.

40. Slobounov SM, Ray W, Johnson B, Slobounov E, Newell KM. Modulation of cortical
activity in 2D versus 3D virtual reality environments: An EEG study. International Journal
of Psychophysiology. 2015 Mar 1;95(3):254–60.

41. Essoe JKY, Reggente N, Ohno AA, Baek YH, Dell'Italia J, Rissman J. Enhancing learning
and retention with distinctive virtual reality environments and mental context reinstatement.
npj Sci Learn. 2022 Dec 8;7(1):1–14.

42. Lessiter J, Freeman J, Keogh E, Davidoff J. A Cross-Media Presence Questionnaire: The
ITC-Sense of Presence Inventory. Presence: Teleoperators and Virtual Environments. 2001
Jun 1;10(3):282–97.

43. Dede C. Immersive Interfaces for Engagement and Learning. Science. 2009 Jan
2;323(5910):66–9.

44. Reggente N. VR for Cognition and Memory. Curr Top Behav Neurosci. 2023 Jul 14;

45. Manjrekar S, Sandilya S, Bhosale D, Kanchi S, Pitkar A, Gondhalekar M. CAVE: An
emerging immersive technology -a review. Proceedings - UKSim-AMSS 16th International
Conference on Computer Modelling and Simulation, UKSim 2014. 2014 Mar 1;131–6.

46. Reggente N, Essoe JKY, Aghajan ZM, Tavakoli AV, McGuire JF, Suthana NA, et al.
Enhancing the Ecological Validity of fMRI Memory Research Using Virtual Reality.
Frontiers in Neuroscience [Internet]. 2018 [cited 2023 Oct 8];12. Available from:
https://www.frontiersin.org/articles/10.3389/fnins.2018.00408

47. Weil A. Breathing: The Master Key to Self Healing (The Self Healing Series). Sounds True;
1999.

48. Pandekar PP, Thangavelu PD. Effect of 4-7-8 Breathing Technique on Anxiety and
Depression in Moderate Chronic Obstructive Pulmonary Disease Patients. International
Journal of Health Sciences. 2019;(5).

49. Riva G, Baños RM, Botella C, Mantovani F, Gaggioli A. Transforming Experience: The
Potential of Augmented Reality and Virtual Reality for Enhancing Personal and Clinical
Change. Front Psychiatry [Internet]. 2016 Sep 30 [cited 2024 Sep 12];7. Available from:
https://www.frontiersin.org/journals/psychiatry/articles/10.3389/fpsyt.2016.00164/full

50. Chirico A, Ferrise F, Cordella L, Gaggioli A. Designing Awe in Virtual Reality: An
Experimental Study. Front Psychol. 2017;8:2351.

51. Chirico A, Cipresso P, Yaden DB, Biassoni F, Riva G, Gaggioli A. Effectiveness of
Immersive Videos in Inducing Awe: An Experimental Study. Sci Rep. 2017 Apr
27;7(1):1218.

52. Adrien V, Bui E, Baubet T, Galletto CP, Duhamel J, McNamara A, et al. Beyond Virtual
Reality: Towards Screen-Free Interfaces for Post-traumatic Stress Disorder Interventions
[Internet]. OSF; 2023 [cited 2024 Sep 12]. Available from: https://osf.io/24beh

53. Shiota MN, Keltner D, John OP. Positive emotion dispositions differentially associated with
Big Five personality and attachment style. The Journal of Positive Psychology.
2006;1(2):61–71.

54. Jamieson G. The Modified Tellegen Absorption Scale: a clearer window on the structure and
meaning of absorption. Australian Journal of Clinical and Experimental Hypnosis. 2005
Nov 1;Vol. 33:119–39.

55. Costa PT, McCrae RR. The Five-Factor Model of Personality and Its Relevance to
Personality Disorders. Journal of Personality Disorders. 1992 Dec;6(4):343–59.

56. Spielberger C, Gorsuch R, Lushene R, Vagg P, Jacobs G. Manual for the State-Trait Anxiety
Inventory (Form Y1 – Y2). Vol. IV, Palo Alto, CA: Consulting Psychologists Press; 1983.

57. Heuchert JP, McNair DM. Profile of Mood States 2nd Edition™ [Internet]. 2012 [cited 2024
Sep 12]. Available from: https://doi.apa.org/doi/10.1037/t05057-000

58. Lau MA, Bishop SR, Segal ZV, Buis T, Anderson ND, Carlson L, et al. The Toronto
Mindfulness Scale: Development and Validation. Journal of Clinical Psychology.
2006;62(12):1445–67.

59. Yaden D, Kaufman S, Hyde E, Chirico A, Gaggioli A, Zhang J, et al. The development of
the Awe Experience Scale (AWE-S): A multifactorial measure for a complex emotion. The
Journal of Positive Psychology. 2018 Jul 18;14.

60. Swami V, Salem N, Furnham A, Tovée MJ. Initial examination of the validity and reliability
of the female photographic figure rating scale for body image assessment. Personality and
Individual Differences. 2008 Jun 1;44(8):1752–61.

61. Elk M, Karinen A, Specker E, Stamkou E, Baas M. 'Standing in Awe': The Effects of Awe
on Body Perception and the Relation with Absorption. Collabra. 2016 Mar 23;2:4.

62. Nour MM, Evans L, Nutt D, Carhart-Harris RL. Ego-Dissolution and Psychedelics:
Validation of the Ego-Dissolution Inventory (EDI). Front Hum Neurosci. 2016 Jun
14;10:269.

63. Schubert T, Friedmann F, Regenbrecht H. The Experience of Presence: Factor Analytic
Insights. Presence: Teleoperators & Virtual Environments. 2001 Jun;10(3):266–81.

64. Jacobs C, M Rigby J. Developing measures of immersion and motivation for learning
technologies in healthcare simulation: a pilot study. J Adv Med Educ Prof. 2022
Jul;10(3):163–71.

65. Gianaros PJ, Muth ER, Mordkoff JT, Levine ME, Stern RM. A Questionnaire for the
Assessment of the Multiple Dimensions of Motion Sickness. Aviat Space Environ Med.
2001 Feb;72(2):115–9.

66. Reitan RM. Validity of the Trail Making Test as an indicator of organic brain damage.
Perceptual and Motor Skills. 1958;8:271–6.

67. Kuhn T, Gottlieb L. Architex [Internet]. 2022. Available from: https://www.myarchitex.com/

68. JASP Team. JASP [Internet]. JASP; 2024. Available from: https://jasp-stats.org/

69. Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al.
glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized
Linear Mixed Modeling. The R Journal. 2017;9(2):378–400.

70. Lenth RV. _emmeans: Estimated Marginal Means, aka Least Squares Means. [Internet].
2022. Available from: https://CRAN.R-project.org/package=emmeans

71. RStudio Team. RStudio: Integrated Development for R. RStudio, PBC [Internet]. Boston,
MA: RStudio, PBC; 2020. Available from: http://www.rstudio.com/

72. Hartig F. DHARMa [Internet]. GitHub. [cited 2024 Sep 9]. Available from:
http://florianhartig.github.io/DHARMa/

73. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B
(Methodological). 1995;57(1):289–300.

74. Lindquist MA, Mejia A. Zen and the art of multiple comparisons. Psychosom Med.
2015;77(2):114–25.

75. Rosseel Y. lavaan: An R Package for Structural Equation Modeling. Journal of Statistical
Software. 2012 May 24;48:1–36.

76. Kothe C, Shirazi SY, Stenner T, Medine D, Boulay C, Grivich MI, et al. The Lab Streaming
Layer for Synchronized Multimodal Recording [Internet]. bioRxiv; 2024 [cited 2024 Sep
12]. p. 2024.02.13.580071. Available from:
https://www.biorxiv.org/content/10.1101/2024.02.13.580071v1

77. Makowski D, Pham T, Lau ZJ, Brammer JC, Lespinasse F, Pham H, et al. NeuroKit2: A
Python toolbox for neurophysiological signal processing. Behav Res Methods. 2021
Aug;53(4):1689–96.

78. Butterworth S. On the Theory of Filter Amplifiers. Experimental Wireless \& the Wireless
Engineer. 1930;7:536–41.

79. Breunig MM, Kriegel HP, Ng RT, Sander J. LOF: identifying density-based local outliers.
SIGMOD Rec. 2000 May 16;29(2):93–104.

80. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:
Machine Learning in Python. MACHINE LEARNING IN PYTHON. 2011;

81. Kleckner IR, Jones RM, Wilder-Smith O, Wormwood JB, Akcakaya M, Quigley KS, et al.
Simple, Transparent, and Flexible Automated Quality Assessment Procedures for
Ambulatory Electrodermal Activity Data. IEEE Trans Biomed Eng. 2018 Jul;65(7):1460–7.

82. Lu W, Nystrom MM, Parikh PJ, Fooshee DR, Hubenschmidt JP, Bradley JD, et al. A semi-
automatic method for peak and valley detection in free-breathing respiratory waveforms.
Med Phys. 2006 Oct;33(10):3634–6.

83. Welch P. The use of fast Fourier transform for the estimation of power spectra: A method
based on time averaging over short, modified periodograms. IEEE Transactions on Audio
and Electroacoustics. 1967 Jun;15(2):70–3.

84. Heart rate variability: standards of measurement, physiological interpretation and clinical
use. Task Force of the European Society of Cardiology and the North American Society of
Pacing and Electrophysiology. Circulation. 1996 Mar 1;93(5):1043–65.

85. Sriboonlert J, Munkong W, Rintawut S, Paladkhua S, Suwongsa R, Kirisattayakul W.
Cananga odorata Aromatherapy Reduces Anxiety in Unexperienced Patients Hospitalized
for Interventional Neuroradiology Procedures: A Randomized Control Trial. J Evid Based
Complementary Altern Med. 2023 Jan 1;28:2515690X221150527.

86. Petruzzello SJ, Snook EM, Gliottoni RC, Motl RW. Anxiety and mood changes associated
with acute cycling in persons with multiple sclerosis. Anxiety, Stress, & Coping. 2009 May
1;22(3):297–307.

87. Mudgalkar N, Kandi V, Baviskar A, Kasturi RR, Bandurapalli B. Preoperative Anxiety
Among Cardiac Surgery Patients and Its Impact on Major Adverse Cardiac Events and
Mortality– A Randomized, Parallel-Group Study. Annals of Cardiac Anaesthesia. 2022
Sep;25(3):293.

88. Craddick RA, Stern MR. Practice Effects on the Trail Making Test. Percept Mot Skills. 1963
Dec;17(3):651–3.

89. McKeown K, Richards E, Centre for Innovative Ageing, Swansea University, Swansea,
Richardson J, Department of Psychology, Swansea University, Swansea, Tales A, et al. The
Trails Making Test. Does a Single Trial Reflect Performance Capability? obm neurobiol.
2021 Jan 8;05(02):1–1.

90. Schoeller F, Christov-Moore L, Lynch C, Diot T, Reggente N. Predicting individual
differences in peak emotional response. PNAS Nexus. 2024 Mar 1;3(3):pgae066.

1138

Supporting Information

S1 Table A. MindGym Breathwork.
S1 Table B. MindGym Rain..
S1 Table C. VR Breathwork..
S1 Table D. VR Rain.
S1 Table E. MindGym.
S1 Table F. VR.
S1 Table G. Breathwork.
S1 Table H. Rain.
S1 Table I. All.

S2 Table A. MindGym Breathwork vs MindGym Rain.
S2 Table B: MindGym Breathwork vs VR Breathwork.
S2 Table C. MindGym Rain vs VR Rain.
S2 Table D. VR Breathwork vs VR Rain.
S2 Table E. Breathwork vs Rain.
S2 Table F. MindGym vs VR.

S3 Table A. MindGym Breathwork vs MindGym Rain.
S3 Table B. MindGym Breathwork vs VR Breathwork.
S3 Table C. MindGym Rain vs VR Rain.
S3 Table D. VR Breathwork vs VR Rain.
S3 Table E. Breathwork vs Rain.
S3 Table F. MindGym vs VR.

S4 Table A. MindGym Breathwork vs MindGym Rain.
S4 Table B. MindGym Breathwork vs VR Breathwork.
S4 Table C. MindGym Rain vs VR Rain.
S4 Table D. VR Breathwork vs VR Rain.
S4 Table E. Breathwork vs Rain.
S4 Table F. MindGym vs VR.

S5 Table A. Architex Total Score.
S5 Table B. Architex Total Speed.
S5 Table C. TMT TRACC.
S5 Table D. Arousal.
S5 Table E. POMS.
S5 Table F. STAI.
S5 Table G. Valence.
S5 Table H. Awe.
S5 Table I. EDI.
S5 Table J. BSE.

S6 Table A. Descriptive Statistics
S6 Table B. Repeated measures ANOVA

1185   **S6 Table C. Post hoc comparisons**
1186   **S6 Figure A. Descriptive Statistics by Timepoints**
1187
1188   **S1 Appendix. Lumena Labs Infinity Cube Spec Sheet**
1189   **S2 Appendix. Transcription of 4-8 Breathwork Audio**

A.

B.