Distinct duration and diet dependent lipid profiles and renal complications

in GCK-MODY

Yadi Huang[1,#], Yuxin Fan[1,#], Yang Liu[1,#], Xuan Liu[1], Wei Li[1], Ying Yang[1], Ziyue Zhang[1],

Shifeng Ma[1], Shuhui Ji[1], Shanshan Chen[1], Hua Shu[1], Wenli Feng[1], Kunlin Wang[1], Qing He[1],

Wenjun Qi[2], Yu Fan[1,‡], Xin Li[1,‡], Ming Liu[1,‡]

[1]Department of Endocrinology and Metabolism, Tianjin Medical University General Hospital,

Tianjin, 300052, China

[2]NHC Key Laboratory of Hormones and Development, Chu Hsien-I Memorial Hospital and

Institute of Endocrinology, Tianjin Medical University, Tianjin, 300134，China

[#]Authors contribute equally to the work.

[‡]To whom correspondence may be addressed. Ming Liu, Department of Endocrinology and

Metabolism, Tianjin Medical University General Hospital, #154 Anshan Road, Heping District,

Tianjin, 300052, China. Email: mingliu@tmu.edu.cn; Yu Fan, Email: fanyu5@hotmail.com; Xin

Li, email: lixin920616@163.com.

Corresponding author:

Ming Liu[1,‡]   M.D., Ph.D.

Email: mingliu@tmu.edu.cn

Tel: +86-22-60817182

Address: Department of Endocrinology and Metabolism, Tianjin Medical University General

Hospital, 154 Anshan Road, Heping District, Tianjin, China 300052

Distinct duration and diet dependent lipid profiles and renal complications

in GCK-MODY

Yadi Huang[1,#], Yuxin Fan[1,#], Yang Liu[1,#], Xuan Liu[1], Wei Li[1], Ying Yang[1], Ziyue Zhang[1],

Shifeng Ma[1], Shuhui Ji[1], Shanshan Chen[1], Hua Shu[1], Wenli Feng[1], Kunlin Wang[1], Qing He[1],

Wenjun Qi[2], Yu Fan[1,‡], Xin Li[1,‡], Ming Liu[1,‡]

Abstract

Heterozygous inactivating mutations in the glucokinase (GCK) gene cause one of the most common types of maturity-onset diabetes of the young type 2 (MODY2), also named GCK-MODY. Studies suggest that, unlike other types of diabetes, patients with GCK-MODY do not have increased risk of diabetic complications and therefore do not typically require antihyperglycemic therapy. However, long-term outcomes of GCK-MODY on lipid profiles and chronic complications remain unclear. Herein, using a knock-in mouse model expressing a novel MODY causing mutation GCK-Q26L (GCK[Mut]), we examined age- and diet-related lipid profiles and diabetic complications. We found that, although GCK[Mut] mice exhibited mild elevated blood glucose, the lipid profiles, body fat composition, and diabetic kidney disease (DKD) were initially improved in high-fat-diet (HFD) fed mice at the age of 28 weeks, supporting potential beneficial effects of GCK[Mut] on lipid metabolism and kidney health. Unexpectedly however, those protective effects diminished by 40 weeks, and became more severe dyslipidemia and kidney injury associated with renal fibrosis and inflammation at 60-week-old mice fed with normal diet (ND) or HFD. Those findings revealed distinct duration- and diet-dependent effects

46 of inactivating GCK mutation on lipid profiles and DKD, highlighting previously unrealized

47 long-term chronic complications in GCK-MODY.

49 **Keywords:** Glucokinase, GCK-MODY, Lipid profiles, Diabetic kidney disease

1. Introduction

Glucokinase (GCK), a glycolytic enzyme belonging to the hexokinase family, plays a pivotal role in catalyzing the conversion of glucose into glucose-6-phosphate, a primary step in glucose metabolism within pancreatic β-cells and hepatocytes [1-3]. Heterozygous inactivating mutations in the GCK gene cause maturity-onset diabetes of the young type 2 (MODY2, also named GCK-MODY), which is one of the most common types of monogenic diabetes [4-6]. Patients with GCK-MODY typically present mild to moderate non-progressive elevated fasting blood glucose (FBG) without significant abnormalities or even improvement of lipid metabolism [7, 8]. More importantly, current evidence indicates that individuals with GCK-MODY are not associated with increased risks of chronic diabetic complications compared to the other types of diabetes [9]. Therefore, current consensuses suggest that such patients may not necessitate pharmacological interventions [10, 11].

However, due to the limited number of patients diagnosed with GCK-MODY, the perspective interventional or observational studies focusing long-term outcomes of GCK-MODY are scarce. Additionally, given the potential impact of persistent hyperglycemia on ocular, vascular, neurological and renal diseases [12-15], it is important to investigate long-term effects of inactivating GCK mutations on lipid metabolism and functional and structural alterations in microvascular complications associated with prolonged hyperglycemia.

Recently, we have established a knock-in mouse line expressing a novel GCK mutation (GCK-Q26L, GCK^{Mut}) identified in two members of a family with early onset diabetes, and demonstrated that this mutation indeed impaired insulin secretion causing mild elevated FBG

and impaired glucose tolerance that mirrors the phenotypes of patients with GCK-MODY (https://doi.org/10.1101/2024.08.13.24311668). In this study, using this knock-in mouse model, we examined age- and diet-related lipid profiles and diabetic complications in the GCK-MODY. We found that GCK mutant indeed presented beneficial effects on lipid metabolism, body fat composition, and renal injury in high-fat-diet (HFD) fed GCK[Mut] mice at the age of 28 weeks. However, those protective effects diminished with time, and more severe dyslipidemia and kidney injury associated with renal fibrosis and inflammation were observed in GCK[Mut] mice fed with HFD at the age of 60 weeks. Those findings revealed distinct duration- and diet-dependent effects of inactivating GCK mutation on body composition, lipid profiles, and diabetic kidney disease (DKD), highlighting previously unrealized long-term metabolic abnormalities and chronic diabetes complications in GCK-MODY. This is poised to offer valuable insights for the prevention and treatment of individuals diagnosed with GCK-MODY aiming to improve long-term reduction of chronic complications. Given the fact that decreased expression and function of GCK are also seen in type 2 diabetes [16], this study may also have clinical implications in the management of type 2 diabetes (T2D).

2. Methods and Materials

2.1 Animal models

Male WT mice were purchased from SPF Biotechnology, while a global GCK-Q26L (GCK[Mut]) knock-in mouse line was established as described in our previous study. Both GCK[Mut] mice and WT mice were housed in pathogen-free conditions within the the Laboratory Animal Center of Tianjin Medical University Chu Hsien-I Memorial Hospital. WT and GCK[Mut] male mice were randomly assigned to four groups and subjected to either a normal diet (ND; comprising 20%

protein, 4% fat, 5% fiber, SCXK(LU) 2018-0003) or a high-fat diet (HFD; comprising 20%

protein, 60% fat, 20% carbohydrates, D12492, Research Diets) from four weeks of age until they

reached 28, 40, and 60 weeks of age. Throughout this period, monitoring of body weight and

blood glucose levels was conducted every 4 weeks. All mice utilized in the study were

maintained on a C57BL/6J background and housed in a 12 h light/dark cycle in a temperature

(22–25 °C) and humidity (55 ± 5%) control room.

2.2 Biochemical measurements

Blood glucose levels were assessed using an automated glucometer (Roche, ACCU-CHEK,

Germany). Mice were placed in metabolic cages to collect 24-hour urine samples. Various

parameters including serum total cholesterol (TC), triglycerides (TG), low-density lipoprotein

cholesterol (LDL-c), high-density lipoprotein cholesterol (HDL-c), serum creatinine, 24-hour

urinary albumin excretion (UAE), and urinary creatinine were determined by kits provided from

Nanjing Jiancheng Institute of Bioengineering (Nanjing, China).

2.3 Intraperitoneal glucose tolerance test and insulin tolerance test

The intraperitoneal glucose tolerance test (IPGTT) involved the intraperitoneal administration of

1 g/kg body weight of glucose in mice following a 6-hour fasting period. The area under the

curve (AUC) was determined using the trapezoidal rule. Simultaneously, blood samples collected

at each time point were utilized to measure blood glucose and serum insulin levels employing an

insulin ELISA assay kit (EZassay, China), following the manufacturer's instructions. For the

intraperitoneal insulin tolerance test (IPITT), mice that fasted for 4 hours received an

intraperitoneal injection of 0.75 U/kg body weight of insulin. Tail vein glucose levels were

monitored at 0, 15, 30, 60 and 120 minutes post-injection. All experiments mentioned above

were conducted using age- and sex-matched cohorts.

2.4 Body composition measurements

Fat mass of was measured using a Bruker Minispec mq7.5 (Brucker).

2.5 Histologic evaluation and transmission electron microscopy anslysis

Following anesthesia, mice were euthanized subsequently the kidneys were harvested，which

were fixed in 4% paraformaldehyde, then paraffin-embedded and were cut into 5-μm-thick

sections. Subsequently, these sections were stained, and after sealing with neutral balsam, images

were captured using AxioImagerM2 (Carl Zeiss, Oberkochen, Germany). The extent of kidney

injury within each group was assessed utilizing Image-ProPlus 6.0 software. The sections

underwent staining procedures involving hematoxylin and eosin (H&E), Periodic acid-Schiff's

(PAS), and Masson's trichrome. For semi-quantitative analysis, twenty stained glomeruli per

section were randomly chosen (5 mice per group) for analyses. For transmission electron

microscopy (TEM), fresh isolated kidney tissues were sliced into small pieces and then fixed in

2.5% glutaraldehyde for a duration of 12 hours. Subsequently, the tissues underwent treatment

with 1% osmium tetroxide. Ultrathin sections were obtained utilizing an ultramicrotome

(LeicaEMUC7, Germany), followed by a dehydration process using gradient alcohol,

permeabilization with acetone, and final embedding in epoxy. The obtained ultrathin sections,

measuring between 70-90 nm, were stained with uranyl acetate and lead citrate. These sections

were examined using a transmission electron microscope (TEM) (HT7700, Japan).

2.6 RNA Extraction, cDNA Synthesis, and Quantitative real-time PCR (qRT-PCR)

Total RNA was extracted and purified from renal tissues using TRIzol reagent (Invitrogen, USA). Subsequently, cDNA was synthesized utilizing the PrimeScriptTM RT kit with gDNAEraser (Takara, Japan), following the manufacturer's instructions. qRT-PCR assays were conducted in a 20 μL reaction volume employing TB Green Premix Ex Taq^{TM} (Tli RNaseH Plus) (Takara, Japan). GAPDH served as the internal control. The primers utilized in this study were designed and chemically synthesized by ZTSINGKE Biological Technology (Beijing, China) Co. Ltd. Details of the primer sequences were provided in **Supplemental Table S1**.

2.7 Protein Extraction and Western Blot Analysis

Renal tissues were lysed using RIPA lysis buffer for protein extraction. Protein samples were then loaded onto SDS-PAGE, electrophoretically separated, and subsequently transferred to nitrocellulose membranes. These membranes were blocked using 5% skimmed milk followed by blotted with indicated antibodies, and visualized by enhanced chemiluminescence (CLiNX, Shanghai, China). The following antibodies were utilized in this study: anti-E-Cadherin (A3044; ABclonal), anti-Vimentin (A11423; ABclonal), anti-α-SMA (A7248; ABclonal), anti-IL-1β (A16288; ABclonal), anti-TNF-α (A11534; ABclonal), and anti-β-actin (KM9001T, Sungene).

2.8 Transcriptome analysis

Renal transcriptome analysis was conducted by BGI Group, encompassing RNA extraction, library preparation, and sequencing. Data processing adhered to the standard BGI mRNA analysis pipeline. Statistical analysis was executed utilizing the Deseq2 software package (v1.28.1). Differentially expressed genes (DEGs) were identified using a threshold of log2 (fold

change) absolute value $\geqslant$ 1 and false discovery rate $\leqslant 0.05$. For further insights, Gene

Ontology (GO) analysis was carried out using cluster/profiler (v3.16.1). Pathway visualization

was accomplished using GraphPad Prism 8.

2.9 Statistics

Statistical analyses were conducted utilizing GraphPad Prism 8 software. The data are presented

as mean ± standard deviation. Analysis involved unpaired Student's t-tests, with statistical

significance set at the traditional critical value of $P < 0.05$. Each experiment was independently

repeated at least three times.

3. Results

3.1 GCK^{Mut} impairs insulin secretion leading to elevated FBG but improves lipid profiles

without affecting body fat composition in mice fed with normal diet (ND)

We have monitored body weight (BW) and FBG of ND fed mice every 4 weeks until 60 weeks

of age, and found that GCK^{Mut} caused non-progressively mild elevated FBG without affecting

BW (**Fig. 1A-B**). To evaluate the effect of GCK^{Mut} on insulin secretion and glucose tolerance,

we performed intraperitoneal glucose tolerance tests (IPGTT) at the ages of 28, 40, and 60 weeks,

respectively. We found that GCK^{Mut} resulted in impaired glucose tolerance associated (**Fig. 1C-E,**

Fig. S1A-C) with decreased basal and glucose stimulated insulin secretion (**Fig. 1F-H, Fig.**

S1D-F). However, GCK^{Mut} did not appeared to affect insulin sensitivity during intraperitoneal

insulin tolerance tests (IPITT) (**Fig. 1I-K, Fig. S1G-I**). Next, we examined body composition

and lipid profiles of GCK^{Mut} mice, and found that although GCK^{Mut} did not change the body fat

composition, the weight of perirenal white adipose tissue (pWAT), and the weight of gonadal

white adipose tissue (gWAT) (**Fig. 1L-N**), it indeed decreased levels of serum triglyceride (TG)

and low-density protein cholesterol (LDL-c) in 28-week-old mice. However, those beneficial

effects on lipid metabolism diminished at 40 and 60 weeks of age (**Fig. 1O-R**). Those data

demonstrate that decreased activity of GCK causes mild non-progressive hyperglycemia with

improved lipid profiles at young age and this improvement waned over time.

3.2 Exacerbation of proteinuria and renal pathological changes in 60-week-old GCK^{Mut}

mice fed with ND

Hyperglycemia is associated with microvascular complications in patients with T2D [17, 18].

Given the fact that persistent elevated blood glucose in patients with GCK-MODY, it is

important to determine the long-term effects of GCK-MODY on chronic diabetes complications.

We therefore measured kidney weight (KW) and BW and found no significant difference in the

KW/BW ratio between GCK^{Mut} and control mice at 28, 40, and 60 weeks of age (**Fig. 2A**).

Additionally, in comparison to WT mice, GCK^{Mut} mice displayed no differences in the levels of

24-hour urinary albumin excretion (UAE), urinary albumin-to-creatinine ratio (UACR), and

serum creatinine at 28 and 40 weeks of age. However, UAE, UACR, and serum creatinine were

notably increased at 60 weeks of age in GCK^{mut} mice (**Fig. 2B-D**). Histological analyses

demonstrated relatively normal glomerular and tubular morphology and structure in GCK^{Mut}

mice at 28 and 40 weeks of age. PAS and Masson staining indicated marginal differences in

thylakoid dilation and renal fibrosis between GCK^{Mut} and control mice at 28 and 40 weeks,

which exacerbated notably at 60 weeks of age (**Fig. 2E-G and K-L**). Subsequently, we utilized

transmission electron microscopy to examine glomerular structure at 28, 40, and 60 weeks of age

in both groups of mice. The observations indicated a trend of increased thickness of glomerular

basement membrane (GBM) at 28 and 40 weeks of age, and a significant thickening of GBM

was evident at 60 weeks of age in GCK^{Mut} mice (**Fig. 2H-J and M**).

3.3 Renal fibrosis- and inflammation-associated genes exhibit high expression in aging

GCK^{Mut} mice fed with ND

To investigate pathways involved in long-term renal changes in GCK^{mut} mice, we performed

transcriptome analysis using kidneys from littermate controls and GCK^{Mut} mice. Gene Ontology

(GO) analysis of the two groups of mice at 60-week-old age revealed that the 940 differentially

expressed genes (DEGs, with │Log2│ fold change ≥1 and false discovery rate ≤0.05), which

were primarily enriched in collagen fibril organization, cell-matrix adhesion, and inflammatory

response (**Fig. 3A**), consistent with previous reports indicating that excessive accumulation of

extracellular mesenchymal tissue and pro-inflammatory responses contribute to the progression

of DKD [19-21]. We further analyzed the DEGs involved in biological progression of DKD, and

found that Vim (Vimentins, encoding intermediate filaments), Acta2 (α-SMA, encoding alpha-

smooth muscle actin), Col1a1 (COL1A1, encoding type I collagen) and Tnf exhibited significant

up-regulation in GCK^{Mut} mice at 60 weeks of age (**Fig. 3B**). The changes of the these genes were

further confirmed by quantitative real-time PCR (qRT-PCR) excepted Col1a1 that did not show a

statistic difference between two groups. The gene Cdh1 (cadherin 1, encoding E-cadherin),

which maintains stable cellular connections and cellular architecture integrity [22], was

significantly downregulated (**Fig. 3C**). Furthermore, Western blot analysis verified that the

protein levels of Vim, α-SMA, TNF-α and IL-1β were indeed significantly elevated in the

kidneys of 60-week-old GCK^{Mut} mice, whereas no significant changes in proteins levels and

mRNA levels were observed in 28- and 40-week-old mice (**Fig. 3D-G, Fig. S2**). These results

suggest that upregulation of the genes associated with renal fibrotic and inflammatory responses

may be attributed to long-term kidney injury in the old GCK^{mut} mice.

3.4 The body composition and lipid profile of 28-week-old GCK^{Mut} mice fed with HFD

improved, but these improvements disappeared at 40 weeks and worsened further at 60

weeks

A recent study has revealed that partial inactivation of GCK can yield beneficial effects on

hepatic lipid metabolism and inflammation [23]. These effects have been postulated as a

potential rationale behind the favorable lipid profile and reduced cardiovascular risk observed in

patients with GCK-MODY [24, 25]. Our data indicated that although improved circulating lipid

profiles were indeed validated at early time in GCK^{Mut} mutant mice, those improvements

diminished over the time in mice fed with ND (**Fig. 1O-R**). We therefore asked what effects of

additional metabolic stress inactivation of GCK on glucolipid metabolism and renal injury. We

randomly grouped 4-week-old littermate mutant and control mice for the HFD regimen while

maintaining the same experimental protocols as those used for mice fed with ND. Our findings

revealed a transient small but statistic decrease in body weight in 28-week-old GCK^{Mut} mice fed

with HFD. However, this divergence disappeared in the mice fed with HFD till 40- and 60-week-

old (**Fig. 4A**). Additionally, elevated fasting and postprandial blood glucose (**Fig. 4B-E, Fig.**

S3A-C), decreased circulating insulin (**Figs. 4F-H, Fig. S3D-F**), and insulin sensitivity (**Figs. 4I-**

K, Fig. S3G-I) had all similar trends as that we observed in mice fed with ND. These data

further demonstrate that although elevated FBG and impaired glucose intolerance persist, these

diabetes phenotypes are stable and are not aggravated even under HFD in GCK^{mut} mice.

Next, we analyzed body composition and revealed a significant reduction in fat mass in HFD fed

GCK^{Mut} mice at 28 weeks in comparison to the control group (**Fig. 4L**). Similarly, pWAT and

gWAT (**Fig. 4M-N**) as well as serum TC and LDL-c levels displayed noteworthy reductions

along with elevated HDL-c in HFD fed GCK^{Mut} mice at 28 weeks (**Fig. 4O-R**). However, at 40

to 60 weeks of age, the aforementioned indices were progressively worse, resulting in significant

deleterious changes of lipid profiles at the age of 60 weeks (**Fig. 4S-V**). These data suggest that

long-term exposure of hyperglycemia and HFD may override the potential beneficial effects of

GCK^{Mut} on lipid metabolism and body composition, manifesting more severe metabolism

disorders.

3.5 GCK^{Mut} attenuates HFD induced renal injury at 28 weeks of age, but aggravates renal

injury at 60 weeks of age

Previous study has indicated that ectopic lipid accumulation in the kidneys contributes to kidney

injury [26-28]. Analogous to alterations in lipid profiles, quantitative analysis of lipids extracted

from kidneys exhibited significantly reduced TC and TG levels along with decreased kidney

weight in GCK^{Mut} mice fed with HFD at the age of 28 weeks. However, with extended exposure

to the hyperglycemia and HFD, TC and TG contents in the kidneys of GCK^{Mut} mice gradually

increased in comparison to WT mice over time (**Fig. 5A-C, Fig. S4A-B**). To further evaluate

renal damage, we also examined 24-hour UAE, UACR, and serum creatinine levels in both

groups of mice. The findings revealed reduced UAE, UACR, and serum creatinine levels in

GCK^{Mut} mice fed with HFD at the age of 28 weeks, but increased alterations in 60-week-old

HFD fed GCK^{Mut} mice (**Fig. 5D-F, Fig. S4C-E**). Notably, histological staining indicated

significantly reduced renal mesangial matrix and collagen fiber deposition in GCK^{Mut} mice

compared to WT mice fed with HFD at the age of 28 weeks, but these indicators gradually

increased at 40 and 60 weeks (**Fig. 5G-I and M-N**). Furthermore, TEM analysis demonstrated a

significantly smaller degree of GBM thickness in GCK^Mut mice at 28 weeks compared to control

mice. However, this thickness significantly increased at 60 weeks (**Fig. 5J-L and O**). These

findings indicate that proteinuria progression and pathological damage induced by a HFD were

mitigated in GCK^Mut mice fed with HFD at early age, whereas long-term (60 weeks) HFD and

hyperglycemia worsened renal damage at 60 weeks in GCK^mut mice. Altogether, those findings

highlight distinct diet and duration dependent effects of GCK inactivating mutation on lipid

profiles and the kidney.

3.6 GCK^mut attenuates the expression of genes associated with renal fibrosis and

inflammation in 28-week-old HFD fed mice, whereas reversal effects GCK^mut on the

changes of above genes are observed after long-term HFD exposure

To investigate the mechanism of attenuated renal injury in GCK^Mut induced by HFD, we

conducted transcriptome analyses of renal samples from GCK^Mut mice fed with HFD at the age

of 28 weeks. GO enrichment analysis revealed that 462 DEGs were predominantly involved in

fatty acid metabolic process, extracellular matrix organization, immune system process, cell-

matrix adhesion, collagen fibril organization and inflammatory response (**Fig. 6A**). Further

analysis of genes enriched in GO highlighted that fibrosis-related genes such as Col1a1 and Vim,

and inflammatory cytokines like Tnf were significantly down-regulated. Additionally, genes

associated with lipid catabolic process, such as Ppara and Lpl, displayed significant up-

regulation (**Fig. 6B**). Subsequently, qRT-PCR confirmed that mRNA levels of Vim, Tnf were

significantly reduced and Cdh1, Ppara, whereas Lpl were significantly elevated in the kidneys of

GCK^{Mut} mice compared with WT mice (**Fig. 6C**). Western blot also showed that the expression

of E-Cadherin was increased, whereas the expression of Vim , α-SMA and TNF-α were

decreased in GCK^{Mut} mice fed with HFD at the age of 28 weeks. Importantly, transcriptome

analyses confirmed by qRT-PCR and western blotting showed that that the genes linked to renal

fibrosis and inflammatory of the above gene expression were reversed in GCK^{mut} mice fed with

HFD at the age of 60 weeks (**Fig. 6D-G, Fig. S5**). These findings suggest that the attenuation of

renal injury in GCK^{Mut} mice on a HFD at a young age may be correlated with the attenuation of

tissue lipid accumulation, fibrosis, and inflammatory responses.

4. Discussion

GCK-MODY is one of most common types of monogenic diabetes [29]. Patients with GCK-

MODY usually exhibit mild and asymptomatic fasting hyperglycemia, persisting from birth with

minimal progression over time [4-6]. Generally, the mild elevated blood glucose in GCK-MODY

does not require treatment with antidiabetic medications, except during pregnancy [30-32]. One

primary rationale for this lies in the markedly low incidence of microvascular and macrovascular

complications, nearly akin to those observed in healthy individuals [33, 34]. Previous studies

have revealed that patients with GCK-MODY showcase normal metabolic profiles, including

relatively lower serum TG with higher HDL-c levels, which may contribute to fewer

complications in these patients [35, 36]. However, given the fact that uncontrolled prolonged

hyperglycemia is strongly associated with diabetes complications and the lack of perspective

studies focusing on patients with GCK-MODY, the long-term effects of GCK inactivating

mutations on chronic complications remain to be further determined.

In this study, using a newly established mouse model carrying a novel MODY causing GCK

mutation, we delved into the impact of the GCK mutant on lipid profiles and diabetic

complications from youth to old age, paying particular attention to a commonly overlooked

factors: the combination of mild hyperglycemia and a high-fat diet. We founded a reduction of

abnormal lipid profiles and mitigated renal injuries in 28-week-old GCK^{Mut} mice fed with HFD.

However, over an extended duration, a tendency towards exacerbated abnormalities in lipid

metabolism and heightened renal damage was observed (**Figs. 4-6**). These data revealed

surprisingly distinct duration and diet dependent effects of GCK^{Mut} and uncover previously

unrecognized pathological significance of prolonged reduction of GCK activity in lipid

metabolism and kidney injuries in patients with GCK-MODY. These data may be more

informative as controlled experimental variables can minimize complex clinical factors including

genetic diversity and use of different medications. These observations warrant further studies in

humans.

One of clinical features of patients with GCK-MODY is optimal lipid profile characterizing as

significantly lower levels of serum LDL-c and TG along with a higher level of HDL-c compared

with nondiabetic control and patients with T2D [24, 36]. A recent study reveals an increased

recruitment of adipose triglyceride lipase and choline/ethanolamine phosphotransferase 1 onto

the surface of HDLs, resulting in enhanced production of plasmalogen phosphatidylcholines that

may confer vascular protection against chronic exposure to hyperglycemia [35]. In this study, we

found that although there were decreased levels of serum LDL-c and TG, there were no

differences in serum HDL-c between control and GCK^{mut} fed with ND (**Fig. 1**). However, under

a metabolic stress (fed with HFD), GCK^{mut} exhibits more beneficial effects not only on levels of

HDL-c (**Fig. 4**), but also on renal lipid deposition and injuries (**Fig. 5**) at the age of 28 weeks.

These observations suggest that beneficial effects GCK^{mut} on lipid metabolism may override the

negative impacts of hyperglycemia on the renal damage at the early stages of diabetes. However,

as prolonged hyperglycemia further extends especially under the HFD conditions, these

beneficial lipid metabolisms were gradually reversed along with increased proteinuria and renal

injuries (**Figs. 2-3 and 5-6**). It remains to be determined whether these paradoxical duration and

diet dependent effects of GCK^{Mut} were solely due to extra long-term exposure of hyperglycemia

that directly disallows the protective effects of GCK^{Mut} on kidneys, or further prolonged

hyperglycemia rejects beneficial effects of GCK^{Mut} on lipid metabolism, resulting more severe

abnormal lipid profiles that may in turn damage kidney.

Recent studies have underscored the pivotal role of inflammation in the pathogenesis of DKD,

indicating its significance in triggering metabolic, biochemical, and hemodynamic disturbances

in diabetes [37, 38]. DKD, recognized as a chronic, low-grade inflammatory disease, represents a

primary feature associated with T2D and its microvascular complications [39-41]. Renal

inflammation manifests through immune cell infiltration, cytokine and chemokine release, and

activation of inflammatory signaling pathways [42]. Our investigation aligns with these findings,

demonstrating that 60-week-old GCK^{Mut} mice, both on a ND or HFD, exhibited more severe

renal inflammatory infiltration compared to controls (**Figs. 3 and 6**). Conversely, the observed

reduction in renal proteinuria and pathological changes in 28-week-old GCK^{Mut} mice fed with

HFD may be associated with diminished inflammatory infiltration. The characteristic

pathological hallmark linked with DKD involves the accumulation of extracellular matrix (ECM)

components in the glomeruli, often leading to renal angio-glomerulosclerosis and functional

decline [19-22]. Our study suggests that GCK^{Mut} may mitigate renal fibrosis by curbing ECM

accumulation and expansion of the thylakoid matrix in the early stages of diabetic nephropathy,

thereby substantially preserving renal function in diabetic mice. Additionally, previous research

has indicated that kidneys of mice on a HFD exhibited notable glomerular and tubular injury,

inclusive of substantial defects in the glomerular filtration barrier and heightened tubular cell

apoptosis [43, 44]. These effects were possibly linked to increased TC and TG content, alongside

activation of the adipogenic pathway in the kidneys [45, 46]. Correspondingly, our observations

reinforce these findings, indicating that this protective effect on the kidney wanes or even

exacerbates renal injury as the lipid profile deteriorates with prolonged high-fat feeding in

GCK^{Mut} mice.

5. Conclusions

In summary, our study presents novel evidence emphasizing the pivotal role of GCK in

glucolipid metabolism and the progression of diabetic complications. MODY causing GCK

mutant benefits lipid metabolism, decreasing renal lipid deposition, inflammatory infiltration,

and fibrosis progression in the early stages, thus retarding disease advancement. However, GCK

mutant appeared to be associated with exacerbated abnormalities of lipid metabolism and renal

injury in later stages. These distinct duration-dependent effects of GCK mutant raise the question

regarding current consensus for the management of patients with GCK-MODY.

Declarations of interest

The authors declare that no conflicts of interest exist and have no financial interests associated

with the present study.

Data availability

The authors are willing to make the raw data generated in their study available to other

investigators upon reasonable request.

Funding

The work was supported by the National Key R&D Program (2022YFE0131400 and

2019YFA0802502), and also by the National Natural Science Foundation of China

(82220108014, 82100865, and 82301951). We acknowledge the support of Tianjin Key Medical

Discipline (Specialty) Construction Project (TJYXZDXK-030A) ，Tianjin Municipal Health

Commission (TJWJ2021ZD001), Tianjin Municipal Science and Technology Commission

(23JCQNJC00680), Tianjin Medical University General Hospital Clinical Research Program

(22ZYYLCZD02)，and Tianjin Medical University Clinical Special Disease Research Center -

Neuroendocrine Tumor Clinical Special Disease Research Center.

Author contributions

YH, YF, YF, XL, XL,WQ and YY conducted experiments and analyzed data; YL, WL, ZZ, SM,

SJ, SC, HS, KW and WF conducted experiments; QH discussed project and edited manuscript;

ML initiated and oversaw the project, designed the experiments, and wrote the manuscript.

References

[1] Matschinsky FM. Glucokinase as glucose sensor and metabolic signal generator in pancreatic

beta-cells and hepatocytes. Diabetes. 1990;39(6):647-52.

[2] Sternisha SM, and Miller BG. Molecular and cellular regulation of human glucokinase. Arch

Biochem Biophys. 2019;663:199-213.

[3] Bonnefond A, Unnikrishnan R, Doria A, Vaxillaire M, Kulkarni RN, Mohan V, et al.

Monogenic diabetes. Nat Rev Dis Primers. 2023;9(1):12.

[4] Pulst K, Arbo T, Kahre T, Peet A, and Tillmann V. MODY2 caused by a novel mutation of

GCK gene. J Pediatr Endocrinol Metab. 2012;25(7-8):801-3.

[5] Shields BM, Hicks S, Shepherd MH, Colclough K, Hattersley AT, and Ellard S. Maturity-

onset diabetes of the young (MODY): how many cases are we missing? Diabetologia.

2010;53(12):2504-8.

[6] Chakera AJ, Steele AM, Gloyn AL, Shepherd MH, Shields B, Ellard S, et al. Recognition and

Management of Individuals With Hyperglycemia Because of a Heterozygous Glucokinase

Mutation. Diabetes Care. 2015;38(7):1383-92.

[7] Hulín J, Škopková M, Valkovičová T, Mikulajová S, Rosoľanková M, Papcun P, et al.

Clinical implications of the glucokinase impaired function - GCK MODY today. Physiol Res.

2020;69(6):995-1011.

[8] Bishay RH, and Greenfield JR. A review of maturity onset diabetes of the young (MODY)

and challenges in the management of glucokinase-MODY. Med J Aust. 2016;205(10):480-5.

[9] Steele AM, Shields BM, Wensley KJ, Colclough K, Ellard S, and Hattersley AT. Prevalence

of vascular complications among patients with glucokinase mutations and prolonged, mild

hyperglycemia. Jama. 2014;311(3):279-86.

[10] Stride A, Shields B, Gill-Carey O, Chakera AJ, Colclough K, Ellard S, et al. Cross-sectional

and longitudinal studies suggest pharmacological treatment used in patients with glucokinase

mutations does not alter glycaemia. Diabetologia. 2014;57(1):54-6.

[11] Carmody D, Naylor RN, Bell CD, Berry S, Montgomery JT, Tadie EC, et al. GCK-MODY

in the US National Monogenic Diabetes Registry: frequently misdiagnosed and unnecessarily

treated. Acta Diabetol. 2016;53(5):703-8.

[12] Au SC, Tang SM, Rong SS, Chen LJ, and Yam JC. Association between hyperglycemia and

retinopathy of prematurity: a systemic review and meta-analysis. Sci Rep. 2015;5:9091.

[13] Nieves-Cintrón M, Flores-Tamez VA, Le T, Baudel MM, and Navedo MF. Cellular and

molecular effects of hyperglycemia on ion channels in vascular smooth muscle. Cell Mol Life

Sci. 2021;78(1):31-61.

[14] Tomlinson DR, and Gardiner NJ. Glucose neurotoxicity. Nat Rev Neurosci. 2008;9(1):36-45.

[15] Alicic RZ, Rooney MT, and Tuttle KR. Diabetic Kidney Disease: Challenges, Progress, and

Possibilities. Clin J Am Soc Nephrol. 2017;12(12):2032-45.

[16] Liu J, Fu H, Kang F, Ning G, Ni Q, Wang W, et al. β-Cell glucokinase expression was

increased in type 2 diabetes subjects with better glycemic control. J Diabetes. 2023;15(5):409-18.

[17] Beckman JA, and Creager MA. Vascular Complications of Diabetes. Circ Res.

2016;118(11):1771-85.

[18] Stumvoll M, Goldstein BJ, and van Haeften TW. Type 2 diabetes: principles of pathogenesis

and therapy. Lancet. 2005;365(9467):1333-46.

[19] Xue M, Sun H, Xu R, Wang Y, Guo J, Li X, et al. GADD45B Promotes Glucose-Induced

Renal Tubular Epithelial-Mesenchymal Transition and Apoptosis via the p38 MAPK and JNK

Signaling Pathways. Front Physiol. 2020;11:1074.

[20] Pan Y, Wang Y, Cai L, Cai Y, Hu J, Yu C, et al. Inhibition of high glucose-induced

inflammatory response and macrophage infiltration by a novel curcumin derivative prevents

renal injury in diabetic rats. Br J Pharmacol. 2012;166(3):1169-82.

[21] Wang S, Li Y, Fan J, Zhang X, Luan J, Bian Q, et al. Interleukin-22 ameliorated renal injury

and fibrosis in diabetic nephropathy through inhibition of NLRP3 inflammasome activation. Cell

Death Dis. 2017;8(7):e2937.

[22] Chang YP, Sun B, Han Z, Han F, Hu SL, Li XY, et al. Saxagliptin Attenuates Albuminuria

by Inhibiting Podocyte Epithelial- to-Mesenchymal Transition via SDF-1α in Diabetic

Nephropathy. Front Pharmacol. 2017;8:780.

[23] Xie Z, Xie T, Liu J, Zhang Q, and Xiao X. Glucokinase Inactivation Ameliorates Lipid

Accumulation and Exerts Favorable Effects on Lipid Metabolism in Hepatocytes. Int J Mol Sci.

2023;24(5).

[24] Ping F, Fu J, and Xiao X. Distinguishing the lipid profile of GCK-MODY patients and its

correlation with hsCRP levels. Front Endocrinol (Lausanne). 2022;13:1024431.

[25] Fendler W, Rizzo M, Borowiec M, Malachowska B, Antosik K, Szadkowska A, et al. Less

but better: cardioprotective lipid profile of patients with GCK-MODY despite lower HDL

cholesterol level. Acta Diabetol. 2014;51(4):625-32.

[26] D'Agati VD, Chagnac A, de Vries AP, Levi M, Porrini E, Herman-Edelstein M, et al.

Obesity-related glomerulopathy: clinical and pathologic characteristics and pathogenesis. Nat

Rev Nephrol. 2016;12(8):453-71.

[27] Xiong W, Xiong Z, Song A, Lei C, Ye C, and Zhang C. Relieving lipid accumulation

through UCP1 suppresses the progression of acute kidney injury by promoting the

AMPK/ULK1/autophagy pathway. Theranostics. 2021;11(10):4637-54.

[28] Chen J, Li T, Vladmir C, Yuan Y, and Sun Z. Renal lipid accumulation induced by high-fat

diet regulates glucose homeostasis via sodium-glucose cotransporter 2. Diabetes Res Clin Pract.

2021;179:109027.

[29] Anık A, Çatlı G, Abacı A, and Böber E. Maturity-onset diabetes of the young (MODY): an

update. J Pediatr Endocrinol Metab. 2015;28(3-4):251-63.

[30] Timsit J, Ciangura C, Dubois-Laforgue D, Saint-Martin C, and Bellanne-Chantelot C.

Pregnancy in Women With Monogenic Diabetes due to Pathogenic Variants of the Glucokinase

Gene: Lessons and Challenges. Front Endocrinol (Lausanne). 2021;12:802423.

[31] Urakami T. Maturity-onset diabetes of the young (MODY): current perspectives on

diagnosis and treatment. Diabetes Metab Syndr Obes. 2019;12:1047-56.

[32] Tosur M, and Philipson LH. Precision diabetes: Lessons learned from maturity-onset

diabetes of the young (MODY). J Diabetes Investig. 2022;13(9):1465-71.

[33] Hattersley AT, Greeley SAW, Polak M, Rubio-Cabezas O, Njølstad PR, Mlynarski W, et al.

ISPAD Clinical Practice Consensus Guidelines 2018: The diagnosis and management of

monogenic diabetes in children and adolescents. Pediatr Diabetes. 2018;19 Suppl 27:47-63.

[34] Gloyn AL. Glucokinase (GCK) mutations in hyper- and hypoglycemia: maturity-onset

diabetes of the young, permanent neonatal diabetes, and hyperinsulinemia of infancy. Hum

Mutat. 2003;22(5):353-62.

[35] Wang X, Lam SM, Cao M, Wang T, Wang Z, Yu M, et al. Localized increases in CEPT1 and

ATGL elevate plasmalogen phosphatidylcholines in HDLs contributing to atheroprotective lipid

profiles in hyperglycemic GCK-MODY. Redox Biol. 2021;40:101855.

[36] Spégel P, Ekholm E, Tuomi T, Groop L, Mulder H, and Filipsson K. Metabolite profiling

reveals normal metabolic control in carriers of mutations in the glucokinase gene (MODY2).

Diabetes. 2013;62(2):653-61.

[37] Pérez-Morales RE, Del Pino MD, Valdivielso JM, Ortiz A, Mora-Fernández C, and

Navarro-González JF. Inflammation in Diabetic Kidney Disease. Nephron. 2019;143(1):12-6.

[38] Rayego-Mateos S, Rodrigues-Diez RR, Fernandez-Fernandez B, Mora-Fernández C,

Marchant V, Donate-Correa J, et al. Targeting inflammation to treat diabetic kidney disease: the

road to 2030. Kidney Int. 2023;103(2):282-96.

[39] Bonner R, Albajrami O, Hudspeth J, and Upadhyay A. Diabetic Kidney Disease. Prim Care.

2020;47(4):645-59.

[40] Jung CY, and Yoo TH. Pathophysiologic Mechanisms and Potential Biomarkers in Diabetic

Kidney Disease. Diabetes Metab J. 2022;46(2):181-97.

[41] Lin YC, Chang YH, Yang SY, Wu KD, and Chu TS. Update of pathophysiology and

management of diabetic kidney disease. J Formos Med Assoc. 2018;117(8):662-75.

[42] Winiarska A, Knysak M, Nabrdalik K, Gumprecht J, and Stompór T. Inflammation and

Oxidative Stress in Diabetic Kidney Disease: The Targets for SGLT2 Inhibitors and GLP-1

Receptor Agonists. Int J Mol Sci. 2021;22(19).

[43] Yamamoto T, Takabatake Y, Takahashi A, Kimura T, Namba T, Matsuda J, et al. High-Fat

Diet-Induced Lysosomal Dysfunction and Impaired Autophagic Flux Contribute to Lipotoxicity

in the Kidney. J Am Soc Nephrol. 2017;28(5):1534-51.

[44] Nakamura J, Yamamoto T, Takabatake Y, Namba-Hamano T, Minami S, Takahashi A, et al.

TFEB-mediated lysosomal exocytosis alleviates high-fat diet-induced lipotoxicity in the kidney.

JCI Insight. 2023;8(4).

[45] de Vries AP, Ruggenenti P, Ruan XZ, Praga M, Cruzado JM, Bajema IM, et al. Fatty kidney: emerging role of ectopic lipid in obesity-related renal disease. Lancet Diabetes Endocrinol. 2014;2(5):417-26.

[46] Chen Y, Deb DK, Fu X, Yi B, Liang Y, Du J, et al. ATP-citrate lyase is an epigenetic regulator to promote obesity-related kidney injury. Faseb j. 2019;33(8):9602-15.

573   **Figure 1. GCK^Mut mice had elevated FBG but improved lipids.** Body weight (**A**) and fasting

574   blood glucose (**B**) were monitored every 4 weeks (n=7-8 mice/group). (**C-E**) Two groups of

mice were intraperitoneally injected with 1 g/kg for IPGTT at 28, 40, or 60 weeks of age (n=5mice/group). (**F-H**) Serum insulin levels were measured by ELISA during IPGTT in both groups of mice (n=5 mice/group). (**I-K**) Two groups of mice were subjected to IPITT (intraperitoneal injection of 0.75 U/kg, n=5mice/group) at 28, 40, or 60 weeks of age. (**L**) Body composition analysis of both groups of mice using a body fat analyzer. (**M-N**) Wet weights of fresh pWAT and gWAT of mice were measured at 28, 40, and 60 weeks of age. Blood samples were used to measure lipid levels including TC (**O**), TG (**P**), LDL-c (**Q**), and HDL-c (**R**), n=5-8mice/group. Values are expressed as mean±SEM. WT+ND vs. GCK^{Mut}+ND: *P<0.05, **P<0.01, ***P<0.001, ns, not significant.

Figure 2. Changes in UAE, serum creatinine and renal histology in GCK^Mut mice fed with ND. (**A**) Kidney and body weights were measured at 28, 40, 60-week old, the ratio of kidney and bodyweight (KW/BW) were calculated. 24-hour UAE (**B**), UACR (**C**), and blood creatinine

levels (**D**) were examined in two groups of mice fed with ND at 28, 40, and 60 weeks of age

(n=7-8 mice/group). (**E-G**) H&E staining, PAS staining, and Masson staining of kidney sections

were performed at 28, 40, and 60 weeks in mice fed with ND (n=5 mice/group). Original

magnification = 400. Scale bar = 50μm. (**H-J**) Changes in glomerular structure observed by

electron microscopy (n=3 mice/group). Original magnification is x8k. Semi-quantitative analysis

of the mesangial matrix (**K**) and the degree of renal fibrosis (**L**). (**M**) Quantification of GBM

thickness. Values are expressed as mean±SEM. WT+ND vs. GCK^{Mut}+ND: *P<0.05, ns, not

significant.

Figure 3. Increased renal fibrosis and inflammatory infiltration in 60-week-old GCK^{Mut}

mice fed with ND. (A) GO analysis of RNA-seq data showed significant changes of pathways in

the renal from 60-week-old WT control and GCK^{Mut} mice fed with ND (n=3mice/group). **(B)**

Volcano plots depicting transcriptome data, dashed line on the y-axis indicates Q-value=0.05 and

fold change on the x-axis is greater than 1. **(C)** The mRNA levels of indicated genes in the

kidneys of 60-week-old WT control and GCK^{Mut} mice were determined by qRT-PCR (n=4

mice/group). **(D-F)** At 28, 40, and 60 weeks of age, the expression of E-Cadherin, Vimentin, α-

SMA, TNF-α, and IL-1β in kidneys of control and GCK^{Mut} mice fed with ND were examined by

western blots (n=6 mice/group). **(G)** Quantification of western blots shown in D-F. Values are

expressed as mean±SEM. WT+ND vs. GCK^{Mut}+ND: *P<0.05, **P<0.01, ns, not significant.

Figure 4. Changes in FBG and lipid metabolism in GCK^{Mut} mice fed with HFD. GCK^{Mut} and

control mice were fed with HFD started from 4 weeks of age. The body weight (**A**) and FBG (**B**)

have been monthly monitored and recorded till 60 weeks of age. IPGTT were performed in mice

fed with HFD at 28 (**C**), 40 (**D**), and 60 (**E**) weeks of age. Serum insulin levels during IPGTT

were also measured at 28 (**F**), 40 (**G**), and 60 (**H**) weeks of age. IPITT were performed in mice

fed with HFD at 28 (**I**), 40 (**J**), and 60 (**K**) weeks of age (n=5-8 mice/group). (**L**) Body

compositions were analyzed in the mice fed with HFD at 28, 40, and 60 weeks of age. (**M-N**)

The weight of pWAT and gWAT were measured in the mice above. TC (**O**), TG (**P**), LDL-c (**Q**),

HDL-c (**R**) were measured in the mice fed with HFD at 28, 40, and 60 weeks of age. (**S-V**)

Changes of TC, TG, LDL-c, HDL-c in mice aged 40 weeks and 60 weeks compared with those

aged 28 weeks. n=5-8 mice/group. Values were expressed as mean±SEM. WT+HFD vs.

GCK^{Mut}+HFD: *P<0.05, **P<0.01, ***P<0.001, ns, not significant.

Figure 5. GCK^{Mut} attenuated HFD associated renal injury at 28 weeks of age, but aggravated renal injury at 60 weeks of age. Renal TC (**A**), Renal TG (**B**), KW/BW (**C**), 24-hour UAE (**D**), UACR (**E**), and serum creatinine levels (**F**) were examined at 28, 40, and 60 weeks of age in both groups of mice (n=7-8 mice/group). (**G-I**) H&E staining, PAS staining, and Masson staining of renal sections at 28, 40, and 60 weeks in both groups of mice (n=5 mice/group). Original magnification = 400. Scale bar = 50μm. (**J-L**) Electron microscopic observation of changes in glomerular structure in two groups of mice (n=3 mice/group). Original magnification is x8k. Semi-quantitative analysis of the mesangial matrix (**M**) and the degree of renal fibrosis (**N**). Quantification of GBM thickness (**O**). Values are expressed as mean±SEM. WT+HFD vs. GCK^{Mut}+HFD: *P<0.05, ns, not significant.

Figure 6

Figure 6. Renal fibrosis and inflammatory were attenuated in HFD GCK^{Mut} mice at 28

weeks of age and aggravated at 60 weeks of age. (**A**) The GO analysis of data obtained by

RNA-Seq showed significantly different gene enrichment (n=3 mice/group). (**B**) Volcano plots

depicting transcriptomics data, dashed line on the y-axis indicates Q-value=0.05 and fold change

on the x-axis is greater than 1. (**C**) The mRNA levels of the mentioned factors in the kidneys of

28-week-old HFD WT control and GCK^{Mut} mice were determined by qRT-PCR (n=4

mice/group). (**D-F**) Renal E-Cadherin, Vimentin, α-SMA, TNF-α, and IL-1β protein levels were

detected by Western blot analysis at 28, 40, and 60 weeks of age (n=6 mice/group). (**G**) Results

of quantitative immunoblot analysis for each group. Values are expressed as mean±SEM.

WT+HFD vs. GCK^{Mut}+HFD: *P<0.05, **P<0.01, ***P<0.001, ns, not significant.

731   **Supplemental Figure 1. The area under the curve (AUC) of IPGTT, serum insulin and**

732   **IPITT in two groups of mice on ND. (A-C)** The AUCs of blood glucose levels were calculated

733   according to Fig.1C-E. (**D-F**) The AUCs of serum insulin levels were calculated from Fig. 1F-H.

734   (**G-I**) The AUCs of blood glucose during IPITT were calculated from Fig.1I-K. n=5mice/group.

735   Values are expressed as mean±SEM. WT+ND vs. GCK^{Mut}+ND: *P<0.05, ns, not significant.

Supplemental Figure 2. The kidney transcriptome analysis at 28 and 40 weeks in two groups of mice on ND. (**A-B**) The number of significantly different genes up- and down-regulated and GO analysis in two groups of mice on regular diet at 28 weeks of age. (**C-D**) The number of significantly different genes up- and down-regulated and GO analysis in two groups of mice on ND at 40 weeks of age. n=3 mice/group.

Supplemental Figure 3. The area under the curve (AUC) of IPGTT, serum insulin, and IPITT in two groups of mice fed with HFD. (**A-C**) The AUC of IPGTT was calculated according to Figure.5C-E. (**D-F**) The AUC for serum insulin levels was calculated from Figure.5F-H. (**G-I**) The AUC values for IPITT were calculated from Figure.5I-K. n=5 mice/group. Values are expressed as mean±SEM. WT+HFD vs. GCK^{Mut}+HFD: *P<0.05, ns, not significant.

Supplemental Figure 4. Changes of Renal TC, Renal TG, 24-hour UAE, UACR, and serum creatinine in mice aged 40 weeks and 60 weeks compared with those aged 28 weeks fed with HFD. Changes of Renal TC (**A**), Renal TG (**B**), 24-hour UAE (**C**), UACR (**D**), and serum creatinine (**E**) in mice aged 40 weeks and 60 weeks compared with those aged 28 weeks. Values were expressed as mean±SEM. WT+HFD vs. GCK^{Mut}+HFD: *$P<0.05$, **$P<0.01$, ***$P<0.001$, ns, not significant.

Supplemental Figure 5. The kidney transcriptome analysis at 40 and 60 weeks in two groups of mice on HFD. (**A-B**) The number of significantly different genes up- and down-regulated and GO analysis in two groups of mice on HFD at 40 weeks of age. (**C-D**) The number of significantly different genes up- and down-regulated and GO analysis in two groups of mice

784   on HFD at 60 weeks of age. (**E**) The mRNA levels of relevant fibrosis and inflammatory factors

785   in the kidneys of two groups of mice on HFD at 60 weeks of age.

Supplemental Table 1. Primer sequences (m-Mus musculus) for qRT-PCR (5'-3')

Gene	Forward	Reverse
mCdh1	CAGGTCTCCTCATGGCTTTGC	CTTCCGAAAAGAAGGCTGTCC
mVim	CGGCTGCGAGAGAAATTGC	CCACTTTCCGTTCAAGGTCAAG
mCol1a1	ATGTTCAGCTTTGTGGACCTC	CAGAAAGCACAGCACTCGC
mActa2	GTCCCAGACATCAGGGAGTAA	TCGGATACTTCAGCGTCAGGA
mTNF-α	GCGACGTGGAACTGGCAGAAGAGG	GTACTTGGGCAGATTGACCTCAGC
mIL-1β	CACTACAGGCTCCGAGATGAACAAC	TGTCGTTGCTTGGTTCTCCTTGTAC
mSrebf1	GGAGCCATGGATTGCACATT	GGCCCGGGAAGTCACTGT
mPpara	AGAGCCCCATCTGTCCTCTC	ACTGGTAGTCTGCAAAACCAAA
mLpl	GGGAGTTTGGCTCCAGAGTTT	TGTGTCTTCAGGGGTCCTTAG
mGAPDH	ACCCACTCCTCCACCTTTGAC	TGTTGCTGTAGCCAAATTCGTT

789