What are the best practices for curating eDNA custom barcode reference libraries? A case study using Australian subterranean fauna.

Michelle T. Guzik[1], Danielle N. Stringer[1], Jake Thornhill[1], Peterson J. Coates[1], Mieke van der Heyde[2], Mia J. Hillyer[3], Nicole E White[2], Mattia Saccò[2], Perry Beasley-Hall[1], William F. Humphreys[4, 5], Mark S. Harvey[4, 5], Joel A. Huey[6], Nerida G. Wilson[4, 5], Jason Alexander [2], Garth Humphreys[4, 7]; Rachael A. King[1, 8], Steve J. B. Cooper[1, 8], Adrian Pinder[9], Giulia Perina[2, 5, 6], Paul Nevill[10]; Andy D. Austin[1, 9]

School of Biological Sciences, The University of Adelaide, Adelaide, South Australia, Australia

Subterranean Research and Groundwater Ecology (SuRGE) Group, Trace and Environmental DNA (TrEnD) Lab, School of Molecular and Life Sciences, Curtin University, Perth, Western Australia, Australia

School of Molecular and Life Sciences, Curtin University, Perth, Western Australia, Australia

Collections and Research, Western Australian Museum, Welshpool, Western Australia, Australia.

School of Biological Sciences, University of Western Australia, Crawley, Western Australia, Australia

Biologic Environmental Survey, East Perth, Western Australia, Australia

Biota Environmental Sciences, West Perth Western Australia, Australia

South Australian Museum, North Terrace, Adelaide, South Australia, Australia

Biodiversity and Conservation Science, Department of Biodiversity, Conservation and Attractions, Kensington, Western Australia, Australia

MBioMe - Mine Site Biomonitoring using eDNA Research Group, Trace and Environmental DNA (TrEnD) Laboratory, School of Molecular and Life Sciences, Curtin University, Perth, Western Australia, Australia

Scientific Significance Statement - 125 words: In threatened ecosystems, environmental

DNA (eDNA) metabarcoding for biodiversity discovery and assessment offers considerable

promise for improvement in the rate, efficiency and accuracy of species detection. For a

better understanding of the biodiversity and ecology of organisms detected using eDNA, a

custom library of known reference sequences with associated correct taxonomic metadata is

required. Minimal guidance is currently available on how an effective (i.e. shareable, multi-

sequence, permits metadata and provides a unified nomenclature) custom barcode reference

library (BRL) can be achieved for subterranean fauna. Here, we present a road map for sound

and reliable curation of a BRL using subterranean fauna from Australia as a case study.

Keywords: groundwater, stygofauna, troglofauna, barcode reference library, metabarcoding,

biomonitoring, biodiversity assessment, data infrastructure

Authors, Institutions, and Author Contribution Statement

MTG conceptualised, analysed and wrote the manuscript; DNS, JT, PJC, MVDH, MS

undertook analyses and designed figures; NEW, PB-H, WFH provided valuable expertise;

MJH contributed to laboratory analyses; JAH, NGW contributed background knowledge and

specimen access and DNA sequence data; RAK, GP and AP contributed to taxonomy; JA and

GH contributed invaluable field knowledge; MSH SJBC, PN, ADA contributed to funding.

All contributed to writing.

Abstract

Identification of species for environmental assessment and monitoring is essential for

understanding anthropogenic impacts on biodiversity, but for subterranean fauna this task is

frequently difficult and time consuming. The implementation of environmental DNA (eDNA)

metabarcoding for biodiversity discovery and assessment offers considerable promise for

improving the rate, accuracy and efficiency of species detection in ecosystems both above

and below the ground. Importantly, for a better understanding of the biodiversity and ecology

of organisms detected using eDNA, a custom library of known reference sequences with

associated correct taxonomic metadata—i.e., a barcode reference library (BRL)—is required.

Yet, minimal guidance is currently available on how an effective (i.e. shareable, multi-

sequence, that permits metadata and has a unified nomenclature) and accurate (i.e. verified)

custom BRL can be achieved. Here, we present a detailed roadmap for curation of a BRL for

subterranean fauna. To do this, we (1) curated a custom sequence database of subterranean

fauna at an environmentally sensitive location, Bungaroo Creek in the Pilbara region of

Western Australia, for four gene loci useful for eDNA metabarcoding (*COI*, *18S* rRNA, *12S*

rRNA and *16S* rRNA); (2) addressed major gaps in taxonomy and disparate nomenclature of

subterranean fauna by estimating 17–29 putative new species with standard delimitation

methods, including 34 Barcode Index Numbers (BINs) in BOLD, and (3) summarised a best

practice workflow for curation of a custom BRL that has broad applicability and can be

applied to any taxa.

Introduction

Groundwater is a high-value natural resource that is vulnerable to human-induced impacts and climate change (Kuang *et al.* 2024). In Australia, the economic value of groundwater resources is estimated to be $33.8 billion (Deloitte Access Economics 2013). Importantly, groundwater also supports groundwater-dependent ecosystems (GDEs), such as subterranean habitats, where a unique and ancient fauna provide essential ecosystem services that maintain underground water condition and quality (Boulton *et al.* 2008). Globally, considerable tension exists with respect to the use of groundwater for economic growth and environmental sustainability (Mammola *et al.* 2019; Saccò *et al.* 2022a; Mammola *et al.* 2024; Saccò *et al.* 2024). In certain countries groundwater fauna are surveyed during routine environmental impact assessments (EIAs) using biomonitoring methods (Environmental Protection Authority 2021), particularly in areas of potential resource development (Gibson *et al.* 2019). For all stakeholders reliant on such biomonitoring, four important issues frequently impact decision-making capabilities (Gibson *et al.* 2019; Mammola *et al.* 2021; Saccò *et al.* 2022b): 1) inconsistent metadata quality (Costello *et al.* 2013; Saccò *et al.* 2022b); 2) minimal data sharing (e.g., Meyer *et al.* (2015)); 3) uncertainty around sampling effort required to provide a robust understanding of the values and predicted impacts and 4) high numbers of undescribed species (Costello 2015; Ficetola *et al.* 2015) with insufficient associated taxonomic knowledge (Mammola *et al.* 2019; Mammola *et al.* 2021). Overcoming these impediments is complicated by reduced taxonomic capacity for many groups and difficulty in distinguishing species on morphological characteristics alone due to anatomical convergence and their often-cryptic nature (Abrams *et al.* 2012; King *et al.* 2022).

Environmental DNA (eDNA) metabarcoding has been identified as an approach with significant potential to alleviate the difficulties associated with traditional sampling (van der Heyde et al. 2023). Indeed, it has been demonstrated as effective for detection of subterranean fauna (West *et al.* 2020; White *et al.* 2020; van der Heyde *et al.* 2023a; van der Heyde *et al.* 2023b). To identify subterranean fauna using eDNA, sequences belonging to anonymous operational taxonomic units (OTUs) must be assigned taxonomic information from a reference database (Taberlet *et al.* 2012). However, this assignment is often patchy due to a lack of data in public DNA databases (e.g., BOLD and GenBank) for both genes and species as well as inconsistencies in taxonomic metadata availability (Costello 2015; Kvist 2013; Dormontt *et al.* 2018; DeWaard *et al.* 2019; Ficetola *et al.* 2019). For these reasons, much of the eDNA data being generated by metabarcoding studies are likely underutilised

(Korbel *et al.* 2017). While taxonomy-independent approaches to eDNA bioinformatics are

being developed (Brinkmann *et al.* 2023; Wilkinson *et al.* 2024), many legislative

frameworks for subterranean fauna require species-level identifications for bioassessment,

although more holistic approaches are being advocated (Koch *et al.* 2024). Therefore, custom

reference databases for querying OTUs are critical for the successful implementation of

eDNA metabarcoding methods (Saccò *et al.* 2022b) and resulting biodiversity assessments.

Guidance on how these databases should be curated is sorely needed.

Selection criteria for the curation of specimens and DNA sequences when submitting to

public databases are established for certain taxonomic groups of subterranean fauna

(pseudoscorpions (Harvey *et al.* 2023; Hlebec *et al.* 2023); isopods and myriapods (Rendoš *et*

al. 2023); asellid isopods (Saclier *et al.* 2024)). Such criteria include: 1) morphologically

verified specimens; 2) homologous DNA sequences corresponding to common

metabarcoding assay regions; 3) estimating levels of genetic variation within and between

species; and 4) metadata linking specimens and current taxonomic information per sequence.

Problematically, for understudied and biodiverse groups such as invertebrate fauna, sequence

records frequently lack associated taxonomic information as it often does not exist. Instead, a

unified system of nomenclature that does not solely rely on formally described species is

required. Various methods for delimiting species are already widely used (e.g., Automatic

Barcode Gap Discovery (ABGD; Puillandre *et al.* (2012)), Assemble Species by Automatic

Partitioning (ASAP; Puillandre *et al.* (2021)) and phylogenetic methods such as Bayesian

Poisson Tree Processes (bPTP; Zhang *et al.* (2018)). Such methods require specialist

knowledge to implement, and are frequently not linked to an existing database platform that

is user-friendly, and where the sequences are available in the same place. Whilst some

authors have suggested that these approaches be used as an alternative to formal taxonomic

approaches (Meierotto *et al.* 2019; Sharkey *et al.* 2021), most researchers recognise that a

major problem exists in providing up-to-date and shareable nomenclature in the absence of

taxonomic information.

To combat the taxonomic impediment facing subterranean fauna, delimited lineages may act

as a proxy for species until such time as these species are formally described. The Barcode of

Life Database (BOLD) applies a system of Barcode Index Numbers (BINs), which are an

index of unique identifiers based on the sequences of varying length and specimens that exist

in BOLD. The BIN pipeline analyses new sequence data for the cytochrome *c* oxidase

subunit I (*COI*) barcoding gene as they are uploaded to the database platform. The power of

the BIN system is that it provides a unified nomenclature for sequences entirely independent of the sequence submitter and their taxonomic capacity, and that information is retained as part of the database. Here, we present a roadmap for curation of a custom barcode reference library (BRL) for subterranean fauna using the BOLD system and build on the new and rapidly expanding area of eDNA metabarcoding research. Our roadmap focuses on an exemplar groundwater-dependent ecosystem at Bungaroo Creek, located in the resource-rich Pilbara region of Western Australia (WA). In future, this data infrastructure can be expanded across Australia and globally in subsequent research since it meets critical principles of a findable, accessible, interoperable and reusable (FAIR) system of data management outlined by Wilkinson *et al.* (2016).

Subterranean biodiversity is comprehensively understudied (Guzik *et al.* 2011), but depending on the region, subterranean fauna can be repeatedly monitored or remain completely unknown to science (Saccò *et al.* 2022a). Given the articulated need for a subterranean fauna BRL by multiple stakeholders (i.e. industry, regulators and academic researchers (Gibson *et al.* 2019)), our overarching aim was to develop a protocol that is easily accessible and usable for effective and efficient characterisation of subterranean biodiversity. Our aims were to:

1. Curate a *de novo* custom BRL of sequences for an exemplar subterranean faunal community of high biodiversity value.

2. Investigate a unified nomenclature that is stable and repeatable, even in the absence of formal description of species, by using species delimitation methods, such as BINs, to identify and name species proxies.

3. Develop and implement a best-practice workflow for a future dynamic subterranean fauna BRL. We outline a protocol for generating and uploading DNA sequence data and associated metadata for the exemplar location using preferred criteria (i.e. 1–2 listed above).

Methods

Bungaroo Creek, WA: a high biodiversity value subterranean community

The Pilbara region of WA is globally recognised for its significant subterranean fauna (Halse *et al.* 2014; Deharveng *et al.* 2024), with many distinct and isolated communities. The Robe River supports several subterranean communities of high ecological value, including that below the major tributary known as Bungaroo Creek (Clark *et al.* 2021). Featuring channel

iron deposits and alluvial aquifers, this section of the Robe River provides an ideal habitat for

subterranean fauna. Indeed, evolutionarily distinct taxon groups such as the blind cave eel

(*Ophisternon candidum*), hydrobiid snails, copepods, ostracods, bathynellaceans and unique

amphipod lineages make this environmentally complex location a significant subterranean

community (Clark *et al*. 2021) which is recognised as a Priority Ecological Community

(Priority 1) in Western Australia (Department of Biodiversity, Conservation and Attractions

2023). Due to the use of Bungaroo Creek groundwater to supply water to the West Pilbara

Water Supply Scheme, the subterranean community is frequently sampled, surveyed, and

monitored for assessment of environmental impact. Here, we used the subterranean

community present at Bungaroo Creek as an exemplar location for the curation of a custom

reference library of DNA sequences for future eDNA metabarcoding.

Specimen collection and curation

To curate a *de novo* custom reference library of subterranean fauna from Bungaroo Creek

(herein called Bungaroo BRL), whole specimens from 21 mining exploration bores were

obtained for Sanger sequencing (Figure 1). Initially, samples were obtained from archival

collections of subterranean fauna from surveys undertaken during 2014 and 2017 and

deposited at the Western Australian Museum (WAM). Collection methods included net

hauling (Saccò *et al.* 2022b) from aquifers using pre-existing bore holes. Morphological

identification was undertaken to the lowest possible taxonomic level, with some specimens

identified to morphospecies but most only to order level or higher.

In addition to the archived material from the WAM collection, invertebrates were also

sampled at the Bungaroo Creek borefield site during June 2017 (as above). Bores were

selected using previous survey data with the aim of capturing the greatest possible genetic

diversity within the constraints of time and logistical feasibility. The locations of bores

sampled are provided in Figure 1, Supplementary Material 1. Invertebrate specimens were

taken from each bore using stygofauna haul nets. Five consecutive haul net samples were

collected, each involving lowering of a net to the bottom of the bore, with nets being

sterilised using bleach between sites to prevent contamination. Bulk invertebrate samples

were stored in a portable refrigerator until the end of sampling each day and were then

transferred to a 4°C refrigerator. The samples for each bore were pooled, and specimens were

sorted *in situ* into major taxonomic groups using a dissecting microscope. Specimens of the

same order were each stored in glass vials, labelled with a single unique identifier, and stored

at –20°C in 100% ethanol until DNA extraction. Representative individuals were catalogued

at the WAM for reference (see Supplementary Material 1 for catalogue information

metadata).

200

Figure 1: Map showing the locations of sampled boreholes at the Bungaroo Creek site, with

bore locations and names. Map created using ArcMap v 10.3.1 (ESRI 2017), data layers

provided by Rio Tinto.

Barcode gene regions

Barcode gene regions are specific segments of DNA sequences that are utilised for molecular

taxonomic identification, particularly in the context of delineating species. This was first

proposed in 2003 (BOLD; Hebert *et al.* (2003)) where standardised DNA sequences were

employed to identify different species. The barcode gene regions selected for this purpose are

typically short (~600-700bp) containing highly conserved regions to allow for primer binding

which then amplify across a variable region to allow for the differentiation of species at the

molecular level. The most commonly used barcode region for animals is *COI*, but other genes

are also used (https://v3.boldsystems.org/index.php/resources/boldfaq). In order to curate a

useable custom library for Bungaroo Creek, we sequenced three mitochondrial genes (*COI*,

12S and *16S*) and one nuclear gene (*18S*). Therefore, we ensured our BRL would; 1) allow

for a strong match to existing metabarcoding assay primers and provide a reference for the

development of new metabarcoding primers (if required); 2) ensure accuracy and breadth of

species identification and documentation of biodiversity (Epp *et al.* 2012; Hebert *et al.* 2013;

Deagle *et al.* 2014); and 3) allow the consolidation of sequence data from other database

libraries, both old and new.

DNA extraction, amplification, and sequencing

DNA was extracted from 468 specimens in December 2017 using a Qiagen DNeasy Blood

and Tissue Kit. Established universal invertebrate primers (below) were used for PCR and

sequencing of specimens, but optimisation of PCR conditions were minimally adjusted

between specimens (Supplementary Material 2). Products were visualised using the eGel

Electrophoresis system (Life Technologies, Scoresby, Victoria, Australia) on 2% agarose gels

with ethidium bromide. Bidirectional long-read Sanger sequencing was carried out at the

Australian Genome Research Facility (Adelaide).

DNA sequence database submissions

For specimens from the Bungaroo BRL, haplotype sequences new to this study were

submitted to GenBank (accession numbers: OR524806-OR524929) and BOLD (public

BOLD Project: BUBRL

https://boldsystems.org/index.php/MAS_Management_DataConsole?codes=BUBRL)

(Supplementary Material 3) including near-complete taxonomy and provenance of the data.

Our rationale for curating and submitting data to both public platforms was that, while

GenBank is the most ubiquitously used to query metabarcoding data due to the BLAST

function, BOLD has many of the GUI features for curating and storing metadata needed for

the current study and any future work. Specifically, BOLD is a long-term funded, user-

friendly web interface with taxonomic, geographic, and photographic metadata available for

each sequence (Ratnasingham and Hebert 2007). In addition, BOLD hosts an *in silico*

delimitation algorithm that assigns identifiers (BINs) to *COI* lineages based on the genetic

distance between all sequences in its database. The BINs provide a centralised system for

identifying lineages that are potentially equivalent to species (Hajibabaei *et al.* 2007). While

BINs frequently do not exist for underrepresented taxonomic groups and regions (Lahaye *et*

al. 2008), we consider that this study fills in some of the gaps in nomenclature and taxonomy

that substantially impact data sharing and transferability between projects. GenBank lacks

most of these features and, because it does not require the association of sequence data with

specimen vouchers, the taxonomy associated with a sequence can never be verified.

Nucleotide analysis and species delimitation

Geneious v10.0.9 (https://www.geneious.com) was used to edit and align *COI*, *12S, 16S* and

18S sequence data for all individuals. Forward and reverse sequence fragments were aligned,

edited and trimmed. *COI* sequencers were translated into amino acids to ensure no stop

codons and a single consensus sequence was created. Consensus sequences of all individuals

for each gene were aligned using MUSCLE (Edgar 2004), trimmed to equal lengths, and

FaBox (Villesen 2007) was used to collapse the final sequence alignments into haplotypes

using its *DNA to haplotype collapser and converter* functionality. Automatic Barcode Gap

Discovery (ABGD) (Puillandre *et al.* 2012), a computationally efficient distance-based

method for species delimitation, was implemented to identify putative species (lineages

equivalents) (https://bioinfo.mnhn.fr/abi/public/abgd/abgdweb.html). This method seeks to

quantify the location of the barcode gap that separates intra- from inter-specific distances. It

performs well when compared to tree-based coalescent methods (Puillandre *et al.* 2012;

Kekkonen and Hebert 2014; Kapli *et al.* 2017) and other threshold techniques (Ratnasingham

and Hebert 2013).

Further to ABGD, we implemented the BOLD BIN method of lineage delimitation. BINs are

an index of unique identifiers based on a database of specimens belonging to each BIN with

their associated metadata as implemented by BOLD systems. The pipeline analyses new

sequence data for the *COI* barcoding region as they are uploaded to BOLD by applying a

two-stage algorithm (Refined Single Linkage) that couples single linkage and Markov

clustering to assign sequences to a sequence cluster and is subsequently assigned a unique

identifier termed a BIN (Ratnasingham and Hebert 2013). The Refined Single Linkage

algorithm matches the taxonomic performance of competing approaches, but couples this

with protocols that are simple enough to allow the automated assignment of all new barcode

records to a BIN. Sequences that establish a new BIN add an entry to the BIN index, while

sequences assigned to an existing BIN contribute their metadata to it.

Results

Summary of haplotypes and genetic distances

A total of 468 individuals were available for PCR and sequencing, but success rates for the

total suite of specimens were low (*COI*: $n = 103$ (22%), *18S*: $n = 137$ (29%), *12S*: $n = 88$

(19%) and *16S*: $n = 21$ (5%)) (see Discussion for explanation). Sequence data were obtained

from a total of 195 individuals (42%), including 660 base pairs (bp) for *COI* ($n = 103$ and $h =$

59), 1017 bp for *18S* ($n = 136$ and $h = 58$), 425 bp for *12S* ($n = 88$ and $h = 48$), and 511 bp for

16S ($n = 21$ and $h = 11$). Only seven individuals were successfully sequenced for all four

barcoding loci (see Supplementary Material 1). All 195 sequences were submitted to our

BOLD project ID BUBRL (Bungaroo BRL) for four barcode genes *COI*-5P (103 sequences),

12S (88), *16S* (21), and *18S* (136), (GenBank accession numbers *COI*: OR524806 -

OR524908; *18S*: OR524930 - OR525066; *12S*: OR525144 - OR525231; *16S*: OR524909-

OR524929) and BOLD (public BOLD Project:

https://boldsystems.org/index.php/MAS_Management_DataConsole?codes=BUBRL).

Species delimitation

Haplotype groups equivalent to lineages were delineated using ABGD (Figure 2 and

Supplementary Material 4) and compared to BOLD BINs as well as to morphospecies

identified by Bennelongia Ltd, an environmental consulting company that undertook routine

biomonitoring in the area during a 2017 survey at Bungaroo Creek (Table 1). Results of the

ABGD analysis revealed 29 lineages for *COI*, 16 lineages for *12S,* 9 lineages for *16S* and 17

lineages for *18S*. Thirty-four BINs were identified for *COI*. Delineated lineages based on

ABGD (for all four genes) and BOLD BINs (based on *COI* only) for major taxon groups

included amphipods (ABGD = 4–8 lineages; BINs = 9), bathynellaceans (ABGD = 1–4

lineages; BINs = 4), thermosbenaceans (ABGD = 1–2 lineages; BINs = 3), annelids (ABGD

= 1–6 lineages; BINs = 5), ostracods (ABGD = 1–3 lineages; BINs = 3) copepods (ABGD =

4–5 lineages; BINs = 8), and a single lineage of gastropod. It should be noted that Pauropoda

are most likely troglophilic/ edaphobitic but were retained here as representatives of the

subterranean ecosystem at Bungaroo Creek.

Figure 2: Phylogenetic tree of taxa from Bungaroo creek for the mitochondrial *COI* gene–
with species delimitation results represented by taxonomic group in colour: Amphipoda
(blue), Copepoda (purple), Bathynellacea (green), Thermosbaenacea (pink), Ostracoda (red),
Annelida (orange), Gastropoda (dark blue). Relationships between major taxonomic groups
are not representative of actual phylogenetic relationships.

Table 1: Morphospecies and delimited lineages of subterranean fauna from Bungaroo Creek, Western Australia (for lineages see Figure 2) using ABGD and BOLD BINs for all four barcoding genes examined in this study.

High level rank	Lineage	Morphological ID	COI Group ID		BIN	COI	18S	12S	16S
			ABGD	BIN		Number of ABGD groups			
Gastropoda (class)	Hydrobiidae		17	AEE7100	1	1	1	1	1
Pauropoda (class)	Pauropodidae		4	AEE8965	1	1	0	1	0
Annelida (superclass)	Phreodrilidae		12	AEE7562	5	5	2	3	6
	Phreodrilidae		11	AEE7131					
	Unknown lineage		10	AEE1055					
	Tubificidae (now Naididae)		8	AEE9005					
	Phreodrilidae		9	AEE7132					
Amphipoda (order)	Eriopisidae	*Nedsia* sp.	29	AEE9183	9	8	5	4	0
				AEE7196					
	Eriopisidae	*Nedsia* sp.	3	AEE7280					
	Unknown lineage		2	AEE5579					
	Eriopisidae	*Pilbarana* sp.	1	AEE8991					
	Paramelitidae		13	AEE8550					
	Paramelitidae		14	AEE9924					

	Unknown lineage		15	AEE2761					
	Bogidiellidae	*Bogidiella* sp.	16	AEE4462					
Bathynellacea (order)	Parabathynellidae	nr *Billibathynella*	23	AEE8197	4	4	1	3	0
	Parabathynellidae	nr *Billibathynella* sp.	22	AEE5164					
	Parabathynellidae	nr *Billibathynella*	21	AEE0507					
	Parabathynellidae	*Hexabathynella* sp.	20	AEE2039					
Thermosbaenacea (order)	Thermosbaenacea		19	AEF0046	3	2	1	2	0
				ADX1249					
	Thermosbaenacea	*Halosbaena* sp.	18	AEE4740					
Ostracoda (order)	Unknown lineage		7	AEE0613	3	3	1	0	2
	Unknown lineage		6	AEE1979					
	Unknown lineage		5	AEE0612					
Copepoda (subclass)	Cyclopoida	*Diacyclops* nr *humphreysi humphreysi*	26	AEE7353	8	5	4	0	0
	Cyclopoida	*Diacyclops* sp.	25	AEE2774					
	Cyclopoida	*Halicyclops* (*Rochacyclops*) *rochai*	28	AEE3241					

				AEE3240				
Harpacticoida		Elaphoidella humphreysi	27	AEE7339				
Calanoida		Stygoridgewayia trispinosa	24	AEE5020				
				AEE5021				
				AEE2775				
Isopoda (order)	Cirolanidae	Kagalana tonde	-		0	1	1	0
Polyxenida (order)	Lophoproctidae	Lophoturus madecassus	-		0	1	1	0
Total			**34**		**29**	**17**	**16**	**9**

Discussion

The use of eDNA metabarcoding to estimate the biodiversity of subterranean faunal

communities has the capacity to alleviate a bottleneck that morphological examination of

physical specimens, limited taxonomic expertise, low-quality or non-existent taxonomic keys,

and a poorly described fauna have on biodiversity assessment (Gibson 2019). The most

efficient way to identify species from anonymous eDNA metabarcoding OTUs is to employ a

curated reference library of sequences (barcodes), preferably with matching voucher

specimens and/or taxonomic information (i.e., a custom BRL). Here, we developed preferred

sequence data and metadata criteria for submission to BOLD for subterranean fauna. This

best-practice protocol addresses a suite of issues specific to subterranean fauna by applying a

unified nomenclature to the exceptional and under-described biodiversity that exists in

groundwater-dependent ecosystems, such as the exemplar site chosen for this study. Further,

our Bungaroo BRL forms the basis of a publicly available data resource that can be used in

eDNA based studies of subterranean fauna into the future.

Estimating community diversity at Bungaroo Creek

Combined, the four genes sequenced here for Bungaroo Creek subterranean fauna delineated

up to 29 genetic lineages as estimated with ABGD based on the *COI* data for 17 taxon

groups. The majority of these lineages were further supported by ABGD results for the other

three genes. These putative species included mites, amphipods, rotifers, copepods, isopods,

worms, snails, and millipedes, indicating a substantially speciose invertebrate community.

The four genes were effective for sequencing the key taxon groups present, but there was

some variation in the number of delineated lineages between genes. For instance, *COI*

delineated the highest number of lineages (29), followed by *18S* (17), *12S* (16) and *16S* (9).

Some of the variation in numbers of delineated species was attributable to sequencing success

or lack thereof (e.g. *16S*, see below) (e.g. Asmyhr and Cooper (2012)), but the delineation

methods used here may also reflect strong phylogeographic variation and structuring (Ortiz

and Francke 2017).

Compared to other studies, 29 lineages of subterranean fauna represent a significantly more

diverse community at Bungaroo than for other Australian aquifers, although this number is

consistent with biodiversity assessment surveys based on morphological examinations (e.g.

Bennelongia (2013)). For example, we observed between two and three lineages of ancient

Thermosbaenacea, whereas only one species is currently described from the Pilbara (Poore

and Humphreys 1992; Eberhard *et al.* 2005). Our results are supported by (Page *et al.* 2018) who also detected a suite of genetic lineages indicating a much higher species diversity of this tethyan crustacean than previously thought. In other Australian aquifers, estimates of 8– 16 putative species have been reported for eastern Australian alluvial aquifers (Asmyhr and Cooper 2012), and calcrete aquifers at Sturt Meadows (Bradford *et al.* 2014) and Lake Violet (Watts and Humphreys 2003; Eberhard *et al.* 2005) in the Yilgarn region WA, and the Millstream aquifer in the Pilbara (Eberhard *et al.* 2005). As such, Bungaroo Creek represents one of the highest putative species estimates for an Australian aquifer. Alongside our own study, recent studies by (Perina *et al.* 2018; Perina *et al.* 2019; Matthews *et al.* 2020) reinforce the view that the Pilbara harbours an exceptionally high and little-explored diversity of subterranean faunal species.

BINs

Barcode Index Numbers (BINs) from genetic databases (i.e., BOLD) have been promoted to offer a standardised and globally unique identifier for each species or species proxy, enhancing comparability and repeatability of studies, allowing researchers to pool data and conduct meta-analyses (Hajibabaei *et al.* 2007; Ratnasingham and Hebert 2007; Lahaye *et al.* 2008). In particular, BINs, based on the genetic distance between sequences independent of prior taxonomic knowledge, have the potential to help identify cryptic or overlooked species to provide an objective means for identifying species boundaries (Hajibabaei *et al.* 2007). Here, ABGD and BIN estimates of putative species for the Bungaroo BRL were generally consistent, although for the copepods three extra lineages were identified using the latter system. While in some cases BINs have been recognised to have some detrimental aspects in certain cases. For example, over-splitting and potential misidentification of lineages is recognised as problematic (Deagle *et al.* 2014) (i.e. BIN assignment has an 86% success rate (Hartop *et al.* 2022)). Further, for unsampled taxon groups and geographic locations where representation in BOLD is absent, existing BINs do not exist, leading to the creation of many new BINs when sampling increases (Lahaye *et al.* 2008). Finally, BIN assignments can change over time as new sequences and taxon groups are added. However, for large, well-represented BINs, the changes are often small and can be tracked by BOLD (Ratnasingham and Hebert 2013). An important advantage of BINS is their usefulness in providing a unified nomenclature that is publicly available and usable. If required, BINs can be verified using current taxonomy or alternative species delimitation methods (i.e., morphospecies, delimited species) and updated on BOLD if needed.

Recommended workflow for Barcode Reference Library subterranean fauna sequence/data

additions

Using Bungaroo Creek as an example, we have demonstrated how to create a dynamic subterranean

faunal reference library for ongoing and future research and biomonitoring that can be updated and,

importantly, follows FAIR Principles of Data Infrastructure (Wilkinson et al. 2016). Below we make

recommendations for a suitable workflow for addition of data to our BRL for subterranean fauna (i.e.

illustrative summary, Figure 3).

Figure 3: Recommended workflow for construction of a Barcode Reference Library.

SPECIMENS

Handling and sorting: in-field and laboratory

Animal sampling requires compliance with regional collection standards and permits. Tissues

should be handled according to those standards. Here, we address handling of invertebrates

and their tissue specifically since sampling of subterranean vertebrate specimens pose

substantially different requirements (Saccò et al. 2022b).

Development of a BRL first requires long-strand sequencing of DNA (Sanger or next

generation sequencing (NGS)) from reference specimens that are also suitable for

morphological examination either pre- or post-sequencing. Overall, the quality of specimen

preservation for DNA sequencing is critical to the successful development of a reference

library. Snap-freezing of specimens is the most optimal for long-term storage of DNA but

difficult to achieve in the field, especially at remote locations. Instead, under ideal conditions,

specimens should be kept alive and then killed by sorting into cold 100% undenatured

ethanol or propylene glycol with no other additives for preservation, and them immediately

refrigerated at a minimum of –20° C (Moreau *et al.* 2013). It should be noted that for small

specimens, especially microcrustaceans, sorting in a laboratory with good lab practises,

ergonomics and quality equipment can yield improved results. Storage of specimens at room

temperature is not ideal but may be unavoidable (Nichols *et al.* 2020). For many surveys, it is

not feasible to spend extended time in the field, and bulk samples are frequently transported

back to the laboratory for sorting. In these situations, it is recommended that the maximum

amount of water from bulk samples be removed prior to sorting, with animals sorted live

where possible, and the organic material is refrigerated to as cold as possible in 100% ethanol

and sorted as soon as possible (Hajibabaei *et al.* 2012). Clean specimen handling is

important, and should be undertaken with gloves and instruments cleaned in 100% ethanol to

control for cross contamination between specimens. It should be noted that NGS approaches

are significantly increasing the potential of using DNA information from degraded specimens

in biodiversity analysis (Hajibabaei *et al.* 2012).

Morphological identification

For critical taxonomic metadata and establishment of specimen vouchers, morphological

analysis of specimens should always be undertaken when adding data to a BRL. A total

evidence approach (with morphological and DNA sequence data) permits validation of

identified species (whether described or new) and can ultimately be used to establish formal

nomenclature for species. While it is important to remember that any species concept is a

working hypothesis, we specifically note that morphological characters alone do not always

reveal the full extent of species-level genetic diversity in a population or at a location (King

et al. 2012; King *et al.* 2022). Additionally, it may not always be feasible to identify

specimens to species or even genus level due to a lack of taxonomic expertise and/or ongoing

funding, the presence of cryptic species, inadequate taxon sampling, the collection of various

life-history stages or damaged specimens that make taxonomic identification difficult, and

taxonomic uncertainty and/or incompleteness, especially in poorly-explored geographic

locations. High-quality images of putative morphotypes taken prior to DNA extraction can be

used as a record of morphological features in the event of destructive sampling and can be

uploaded to BOLD. While new techniques are being developed to retain the exoskeleton of

specimens during DNA extraction and library preparation (e.g. acoustic droplet ejection (Cain-

Hom *et al.* 2016)), currently this technology is highly specialised and expensive to

implement. However, ongoing development by major consortia, such as the BIOSCAN

project (Park *et al.* 2023), has the potential to expand the application of these methods and decrease the cost of widespread implementation. In the interim, intact mature animals should ideally be morphologically identified where possible prior to sequencing to yield morphological and molecular data linked within the same specimen.

Voucher specimens

Retention of specimen vouchers for future morphological validation, and description is vital, and further DNA sequencing can aid with verification of genetic lineages. Archived specimens and tissues are valuable as a temporal genetic and morphological record of the status of a species. Specimen storage (Burrell *et al.* 2015) and age of specimens (Miller *et al.* 2013) are important parameters to successful DNA extraction and PCR amplification and can significantly affect the task of building a BRL (Baird *et al.* 2011; Hebert *et al.* 2013; Burrell *et al.* 2015). In order to retain specimens, dissection and extraction of DNA from a part of the body that will minimally interfere with morphological identification is ideal and usually feasible. However, for very small individuals, sequencing of one to five representatives from a group of morphologically identical individuals collected from the same sampling site allows remaining individuals to be treated as a voucher specimens. Recent technical developments have enabled whole genome sequencing of single individuals for tiny species using genome amplification approaches (Picogram input multimodal sequencing (PiMmS) method; Laumer (2023)), but these methods are yet to be widely employed by the biodiversity community. .

DNA SEQUENCE DATA

Reference sequences: specimen identification with DNA sequencing

DNA sequencing is a powerful tool for identifying specimens to species and genus levels, particularly when traditional morphological or biochemical methods are inadequate or inconclusive. Depending on the gene of interest, a minimum of one sequence (i.e. a single specimen) is recommended to define an OTU and sequencing multiple individuals from a population will provide an indication of diversity within a species. Obtaining sequences from multiple genes also increases confidence in the accuracy of the identification, as any sequencing errors (particularly relevant for Sanger sequencing since errors caused by nuclear copies of mitochondrial DNA (NUMTs are a possibility (Richly and Leister 2004)), or contamination can be detected and potentially corrected. Proper retention and storage of raw reads is important for verification and reproducibility of the data. Raw data, such as trace files, can be uploaded to BOLD alongside additional metadata (see below).

     Primers

Primer mismatch is known to be a major issue in amplification of barcode fragments.

Universal primers (LCOI490/HCO2198 (Folmer *et al.* 1994) and C1-J-1718 (Simon *et al.*

1994)) are effective for *COI* amplification for representatives of many animal phyla (Hebert

et al. 2003). However, in agreement with past studies, we observed that these standard

primers can amplify inconsistently across different groups in a variety of subterranean

crustacean taxa (e.g. Camacho *et al.* 2011; Karanovic and Cooper 2012; Asmyhr and Cooper

2012). The universal primers for the nuclear *18S* (Whiting 2002) and mitochondrial *12S*

rRNA (Wetzer 2001) genes are known to amplify well in many crustacean taxa (Asmyhr and

Cooper 2012) and worked well here to produce alternative barcodes (Corse *et al.* 2010;

Hardy *et al.* 2011). The standard *16S* primers (16sar/16Sbr Palumbi (1996)), whilst not

successful for many taxa, were extremely useful for gastropods and copepods in this study.

For these reasons, many studies develop *ad hoc* primers depending on the taxon group (e.g.,

Little *et al.* (2016). In future, it will become increasingly feasible and affordable to sequence

whole mitochondrial genomes employing next-generation sequencing (e.g., eDNA

metagenomics), obtaining many barcoding genes at once or, alternatively, using the entire

mitochondrial genome for species delineation or identification.

For a BRL to be useful for implementation into future eDNA metabarcoding studies, the

barcodes of voucher specimens should match the most commonly used assay regions. Due to

the historical technological impediment of the last 20 years where Sanger sequencing of

barcode loci was the most inexpensive, subterranean fauna have most commonly been

sequenced for loci including *COI*, *16S, 12S* and *18S,* in order of frequency. BRL DNA

sequences should be homologous to metabarcoding assay regions (i.e., standard barcodes).

Currently, *COI* and *18S* are most commonly implemented for metazoan metabarcoding

detection, and *16S* for microbial detection from eDNA. Commonly used primers for eDNA

metabarcoding assays of freshwater invertebrates include *18S* (Pochon *et al.* 2013), *16S* (such

as for crustaceans (Berry *et al.* 2017) and insects (Clarke *et al.* 2014)) and *COI* (Vamos *et al.*

2017). In future, genome skimming, including shotgun sequencing of mitochondrial genomes

and Ultra Conserved Elements, may effectively be used as a universal approach for

generating sequence data, especially for genuinely old and degraded specimens (Blaimer *et*

al. 2016; McCormack *et al.* 2016) but also for contemporary specimens.

METADATA

Linking specimen voucher information to DNA sequence and best available taxonomy

For an effective custom BRL database, the link between DNA sequences, morphologically identified specimens by a taxonomic practitioner (where possible), specimen metadata and the most up to date available taxonomy that can be linked to sequence data are critical to its usability. A comprehensive guide to curation of a BRL for aquatic organisms (eukaryotic and prokaryotic) is described by Rimet *et al.* (2021) who recommended BOLD for its ability to link varied metadata to a sequence. For BOLD, minimum information per DNA sequence is: Sample ID (collection code), Field ID and/or Museum voucher ID (voucher number), Institution Storing, Phylum, and Country. However, it is strongly recommended that latitude and longitude (sample locality), morphological identification to lowest taxonomic level (Rimet *et al.* 2021), the name of the taxonomist, and the year of the identification also be included. BOLD allows submission of trace files for sequences and images (https://boldsystems.org/index.php/databases), which enhances accessibility and reusability. However, required information for GenBank is minimal and the infrastructure needed for hosting additional data, such as chromatograms and images, is not available making it difficult to provide maximum metadata per specimen.

DATABASE

Unified nomenclature and species delimitation

As mentioned previously, BOLD has developed the BIN system for *COI* sequences. The validity of using BINs as equivalent to species has been debated in the literature (Kekkonen *et al.* 2015; Meier *et al.* 2022). This is especially the case since use of *COI* only in conjunction with the OTU designation method of BOLD (i.e. the Refined Single Linkage (RESL) algorithm (Ratnasingham and Hebert 2013)) there is a tendency for over-splitting of lineages (Hlebec *et al.* 2023). However, we consider, as in other studies (e.g. Kekkonen *et al.* 2015; Blagoev *et al*. 2016), the utility of BINS as a built-in clustering method for generating species hypotheses is a valid approach, particularly for use by non-specialists. We observed that BINs in conjunction with the ABGD species delimitation approach and gene fragment concordance to be consistent and in-line with the findings of other studies (Kekkonen *et al.* 2015). In future, the 29 lineages and 34 BINs we resolved here may be verified as undescribed species, with morphological examination of specimen vouchers.

GenBank, as the most widely used DNA sequence repository, can be used for querying

metabarcoding reads and OTUs obtained in eDNA studies. Whilst this database is useful,

especially given its centralization, the implementation and accessibility of metadata (e.g.,

taxonomic, geographic and photographic) associated with sequences means that there are

added advantages to BOLD. Designed principally for biodiversity studies, the GUI of BOLD

is uniquely configured to summarise geographic and taxonomic data. For example,

phylogenetic trees can be directly estimated from DNA sequences, and the data can be stored

in various projects and databases. Further, these aspects have the added benefit of being user-

friendly for non-specialists, and provide an accessible implementation of the subterranean

fauna BRL presented here. BOLD is also one of the most usable and collaborative databases

available for depositing and analysing eDNA data. Whilst other databases, such as GenBank,

have projects and databases (e.g. BioProjects, BioSamples, and the Sequence Read Archive),

there are distinct advantages to using BOLD since, in addition to its ongoing funding, it is

extremely attractive for the accessibility of custom BRLs for future research and, as such, we

recommend its use for accessing our subterranean fauna database.

The Darwin Core Standard (DwC) offers a stable and flexible framework for compiling

biodiversity data from varied and variable sources. Originally developed by the Biodiversity

Information Standards community, Darwin Core is an evolving platform providing

definitions and common terms that are shareable and searchable (Wieczorek *et al.* 2012). It

plays a fundamental role in the sharing, use and reuse of open-access biodiversity data. We

recommend that in curating data for custom BRL's, the DwC standard terms be used.

Importantly, BOLD implements the DwC standard (Ratnasingham and Hebert 2007).

Future taxonomic considerations

The usefulness of a subterranean fauna BRL rests in its ability to assign OTUs a unified

nomenclature and, where possible, species names. Reference sequences must be directly

linked to vouchered specimens of each species that are curated and maintained long-term

within museum collections. For conservation management, species names are also critical

(Costello *et al*., 2015), but as is the case for most invertebrate groups, especially subterranean

fauna, formal taxonomy is not progressing at a rapid enough pace compared to the

recognition of the burgeoning number of molecular OTUs (Perina *et al.* 2018; Perina *et al.*

2019; Matthews *et al.* 2020). The number of new lineages we recovered in this study at

Bungaroo Creek highlight the need for a unified nomenclature that is used consistently

among stakeholders and to which taxonomy can be later applied, when available, to alleviate

the well-documented taxonomic impediments (Riedel *et al.* 2013).

Conclusions

A robust BRL is essential for the effective translation of anonymous DNA sequence data to

meaningful taxonomic identification, including for the detection of fauna using

metabarcoding. However, shareable, reusable and publicly available datasets for poorly

known ecosystems are frequently unavailable for query. To address this lack of fundamental,

but crucially important research, we have presented a custom BRL for an Australian

subterranean fauna as a test case and, in doing so, outline best-practice standards for

establishing a BRL to invigorate future subterranean faunal community assessment and

monitoring. However, these recommendations also apply more broadly to all custom BRLs.

Priorities to transform our consensus of species delimitation results into a more complete

BRL in the future include morphological verification and taxonomic diagnoses of relevant

species. A custom BRL requires DNA sequences at both broad (order, family) and narrow

(genus to population level) taxonomic coverage to be useful in identifying unknown

individuals and sequences in genetic and taxonomic studies (Hebert *et al.* 2013). Additional

verification through more extensive sampling of subterranean fauna, barcoding of a much

broader range of genes, and further phylogenetic analyses will also be essential. Given the

unique subterranean biodiversity present at Bungaroo Creek and within the Pilbara region

more broadly, a coordinated effort will be required to advance this data infrastructure in the

future.

Acknowledgements

This research was part funded by Australian Research Council (ARC) Industry Linkage

Project Grant LP140100555 to A.D.A. S.J.B.C. and W.F.H. (linkage partners: South

Australian Museum, Western Australian Museum, the WA Department of Biodiversity,

Conservation and Attractions (DBCA) (formerly Department of Parks and Wildlife),

Bennelongia Pty Ltd, Biota Environmental Sciences Pty Ltd), ARC Industry Linkage Project

Grant LP190100555 to A.D.A. (linkage partners: Rio Tinto, BHP, Chevron, South Australian

Museum, Western Australian Museum, WA DBCA, WA Department for Water and

Environmental Regulation, Western Australian Biodiversity Science Institute (WABSI), Rio

Tinto consulting contract to N.W. and M.B., and a research contract with WABSI funded by

Rio Tinto and BHP. We thank S. Halse (Bennelongia) for collaboration and specimen

collection as part of faunal surveys for ecological impact assessments, E. Matthews (formerly

The University of Adelaide) for assistance collecting and sorting specimen samples in the

Robe River Valley, Caitlin O'Neill, Dean Main and Michael Curran (Rio Tinto) for field

support and valuable site information. Specimens were sampled under a Department of Parks

and Wildlife Regulation 17 Licence to take fauna for scientific purposes (Number: 08-

000700-1). This work was supported by resources provided by the Pawsey Supercomputing

Centre with funding from the Australian Government and the Government of Western

Australia.

References

597 Abrams K., Guzik M., Cooper S., Humphreys W., King R., Cho J.-L., Austin A. (2012) What lies
598 beneath: molecular phylogenetics and ancestral state reconstruction of the ancient subterranean
599 Australian Parabathynellidae (Syncarida, Crustacea). *Molecular Phylogenetics and Evolution* **64**(1),
600 130-144.

601
602 Asmyhr M. G., Cooper S. J. B. (2012) Difficulties barcoding in the dark: the case of crustacean
603 stygofauna from eastern Australia. *Invertebrate Systematics* **26**(6), 583-591.

604
605 Environmental Protection Authority (2021) Technical guidance – Subterranean fauna surveys for
606 environmental impact assessment: Western Australia.

607
608 Baird D. J., Pascoe T. J., Zhou X., Hajibabaei M. (2011) Building freshwater macroinvertebrate
609 DNA-barcode libraries from reference collection material: formalin preservation vs specimen age.
610 *Journal of the North American Benthological Society* **30**(1), 125-130.

611
612 Bennelongia (2013) Bungaroo South: Subterranean Fauna Assessment. Report 2013/189

613
614 Berry T. E., Osterrieder S. K., Murray D. C., Coghlan M. L., Richardson A. J., Grealy A. K., Stat M.,
615 Bejder L., Bunce M. (2017) DNA metabarcoding for diet analysis and biodiversity: A case study
616 using the endangered Australian sea lion (Neophoca cinerea). *Ecology and Evolution* **7**(14), 5435-
617 5453.

618
619 Blaimer B. B., Lloyd M. W., Guillory W. X., Brady S. G. (2016) Sequence capture and phylogenetic
620 utility of genomic ultraconserved elements obtained from pinned insect specimens. *PLOS ONE* **11**(8),
621 e0161531.

622
623 Blagoev, G. A., Dewaard, J. R., Ratnasingham, S., Dewaard, S. L., Lu, L., Robertson, J., Telfer, A. C.
624 and Hebert, P. D. (2016). Untangling taxonomy: a DNA barcode reference library for Canadian
625 spiders. *Molecular Ecology Resources*, **16**(1), 325-341.
626
627 Boulton A. J., Fenwick G. D., Hancock P. J., Harvey M. S. (2008) Biodiversity, functional roles and
628 ecosystem services of groundwater invertebrates. *Invertebrate Systematics* **22**(2), 103-116.

629
630 Bradford T. M., Humphreys W. F., Austin A. D., Cooper S. J. (2014) Identification of trophic niches
631 of subterranean diving beetles in a calcrete aquifer by DNA and stable isotope analyses. *Marine and
632 Freshwater Research* **65**(2), 95-104.

633
634 Brinkmann I., Schweizer M., Singer D., Quinchard S., Barras C., Bernhard J. M., Filipsson H. L.
635 (2023) Through the eDNA looking glass: Responses of fjord benthic foraminiferal communities to
636 contrasting environmental conditions. *Journal of Eukaryotic Microbiology* **70**(4), e12975.

637
638 Burrell A. S., Disotell T. R., Bergey C. M. (2015) The use of museum specimens with high-
639 throughput DNA sequencers. *Journal of human evolution* **79**, 35-44.

641 Cain-Hom C., Pabalate R., Pham A., Patel H. N., Wiler R., Cox J. C. (2016) Mammalian genotyping
642 using acoustic droplet ejection for enhanced data reproducibility, superior throughput, and minimized
643 cross-contamination. *Journal of Laboratory Automation* **21**(1), 37-48.

644
645 Camacho A. I., Dorda B. A., Rey I. (2011) Identifying cryptic speciation across groundwater
646 populations: first COI sequences of Bathynellidae (Crustacea, Syncarida). *Graellsia* **67**(1), 7-12.

647 Clark H. L., Buzatto B. A., Halse, S. A. (2021). A hotspot of arid zone subterranean biodiversity: The
648 Robe Valley in Western Australia. *Diversity*, **13**(10), 482.
649
650 Clarke L. J., Soubrier J., Weyrich L. S., Cooper A. (2014) Environmental metabarcodes for insects: in
651 silico PCR reveals potential for taxonomic bias. *Molecular Ecology Resources* **14**(6), 1160-70.

652
653 Corse E., Costedoat C., Chappaz R., Pech N., Martin J. F., Gilles A. (2010) A PCR□based method for
654 diet analysis in freshwater organisms using 18S rDNA barcoding on faeces. *Molecular Ecology*
655 *Resources* **10**(1), 96-108.

656
657 Costello M. J. (2015) Biodiversity: the known, unknown, and rates of extinction. *Current Biology*
658 **25**(9), R368-R371.

659
660 Costello M. J., Michener W. K., Gahegan M., Zhang Z.-Q., Bourne P. E. (2013) Biodiversity data
661 should be published, cited, and peer reviewed. *Trends in Ecology & Evolution* **28**(8), 454-461.

662
663 Deagle B. E., Jarman S. N., Coissac E., Pompanon F., Taberlet P. (2014) DNA metabarcoding and the
664 cytochrome c oxidase subunit I marker: not a perfect match. *Biology Letters* **10**(9).

665
666 Deharveng L., Bedos A., Pipan T., Culver D. C. (2024) Global Subterranean Biodiversity: A Unique
667 Pattern. Diversity, 16(3), 157.

668
669 Department of Biodiversity, Conservation and Attractions (2023) Priority Ecological Communities for
670 Western Australia. Version 35: Perth, Western Australia

671
672 DeWaard J. R., Ratnasingham S., Zakharov E. V., Borisenko A. V., Steinke D., Telfer A. C., Perez K.
673 H., Sones J. E., Young M. R., Levesque-Beaudin V. (2019) A reference library for Canadian
674 invertebrates with 1.5 million barcodes, voucher specimens, and DNA samples. *Scientific Data* **6**(1),
675 308.

676
677 Dormontt E. E., Van Dijk K.-J., Bell K. L., Biffin E., Breed M. F., Byrne M., Caddy-Retalic S.,
678 Encinas-Viso F., Nevill P. G., Shapcott A. (2018) Advancing DNA barcoding and metabarcoding
679 applications for plants requires systematic analysis of herbarium collections—an Australian
680 perspective. *Frontiers in Ecology and Evolution* **6**, 134.

681
682 Eberhard S. M., Halse S. A., Humphreys W. F. (2005) Stygofauna in the Pilbara region, north-west
683 Western Australia: a review. *Journal of the Royal Society of Western Australia* **88**, 167–176.

684
685 Deloitte Access Economics (2013) Economic Value of Groundwater in Australia. National Centre for
686 Groundwater Research and Training.

687
688 Edgar R. C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput.
689 *Nucleic Acids Research* **32** (5), 1792-97.

690
691 Epp L. S., Boessenkool S., Bellemain E. P., Haile J., Esposito A., Riaz T., Erseus C., Gusarov V. I.,
692 Edwards M. E., Johnsen A., Stenøien H. K. (2012) New environmental metabarcodes for analysing
693 soil DNA: potential for studying past and present ecosystems. *Molecular Ecology* **21**(8), 1821-1833.

694
695 Ficetola G. F., Canedoli C., Stoch F. (2019) The Racovitzan impediment and the hidden biodiversity
696 of unexplored environments. *Conservation Biology* **33**(1), 214-216.

697
698 Ficetola G. F., Pansu J., Bonin A., Coissac E., Giguet-Covex C., De Barba M., Gielly L., Lopes C.
699 M., Boyer F., Pompanon F., Raye G., Taberlet P. (2015) Replication levels, false presences and the
700 estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources*
701 **15**(3), 543-56.

702
703 Folmer O., Black M., Hoeh W., Lutz R., Vrijenoek R. (1994) DNA primers for amplification of
704 mitochondrial cytochrome c oxidase subunit 1 from diverse metazoan invertebrates. *Molecular*
705 *Marine Biology and Biotechnology* **3**(5), 294-299.

706
707 Gibson L., Humphreys W. F., Harvey M., Hyder B., Winzer A. (2019) Shedding light on the hidden
708 world of subterranean fauna: A transdisciplinary research approach. *Science of The Total Environment*
709 **684**, 381-389

710
711 Guzik M. T., Austin A. D., Cooper S. J., Harvey M. S., Humphreys W. F., Bradford T., Eberhard S.
712 M., King R. A., Leys R., Muirhead K. A., Tomlinson M. (2011) Is the Australian subterranean fauna
713 uniquely diverse? *Invertebrate Systematics* **24**(5), 407-418.

714
715 Hajibabaei M., Singer G. A., Hebert P. D., Hickey D. A. (2007) DNA barcoding: how it complements
716 taxonomy, molecular phylogenetics and population genetics. *Trends in Genetics* **23**(4), 167-72.

717
718 Hajibabaei M., Spall J. L., Shokralla S., van Konynenburg S. (2012) Assessing biodiversity of a
719 freshwater benthic macroinvertebrate community through non-destructive environmental barcoding of
720 DNA from preservative ethanol. *BMC Ecology* **12**, 1-10.

721
722 Halse S. A., Scanlon M. D., Cocking J. S., Barron H. J., Richardson J. B., Eberhard S. M. (2014)
723 Pilbara stygofauna: deep groundwater of an arid landscape contains globally significant radiation of
724 biodiversity. *Records of the Western Australian Museum* **78**, 443–483.

725
726 Hardy C. M., Adams M., Jerry D. R., Morgan M. J., Hartley D. M. (2011) DNA barcoding to support
727 conservation: species identification, genetic structure and biogeography of fishes in the Murray—
728 Darling River Basin, Australia. *Marine and Freshwater Research* **62**(8), 887-901.

729
730 Hartop E., Srivathsan A., Ronquist F., Meier R. (2022) Towards large-scale integrative taxonomy
731 (LIT): resolving the data conundrum for dark taxa. *Systematic Biology* **71**(6), 1404-1422.

732   Harvey, M.S., Burger, M.A.A., Abrams, K.M., Finston, T.L., Huey, J.A. and Perina, G. (2023). The
733   systematics of the pseudoscorpion genus *Indohya* (Pseudoscorpiones: Hyidae) in Australia. *Zootaxa*
734   **5342**, 1–119. https://doi.org/10.11646/zootaxa.5342.1
735
736   Hebert P. D., Cywinska A., Ball S. L., deWaard J. R. (2003) Biological identifications through DNA
737   barcodes. *Proceedings of the Royal Society of London, Series B* **270**, 313-322.
738
739   Hebert P. D., Dewaard J. R., Zakharov E. V., Prosser S. W., Sones J. E., McKeown J. T., Mantle B.,
740   La Salle J. (2013) A DNA 'barcode blitz': rapid digitization and sequencing of a natural history
741   collection. *PLOS ONE* **8**(7), e68535.
742
743   Hlebec D., Podnar M., Kučinić M., Harms D. (2023) Molecular analyses of pseudoscorpions in a
744   subterranean biodiversity hotspot reveal cryptic diversity and microendemism. *Scientific Reports*
745   **13**(1), 430.
746
747   Kapli P., Lutteropp S., Zhang J., Kobert K., Pavlidis P., Stamatakis A., T. F. (2017) Multi-rate
748   Poisson tree processes for single-locus species delimitation under maximum likelihood and Markov
749   chain Monte Carlo. *Bioinformatics* **33** (11), 1630-1638.
750
751   Karanovic T., Cooper S. J. (2012) Explosive radiation of the genus Schizopera on a small
752   subterranean island in Western Australia (Copepoda: Harpacticoida): unravelling the cases of cryptic
753   speciation, size differentiation and multiple invasions. *Invertebrate Systematics* **26**(2), 115-192.
754
755   Kekkonen M., Hebert P. D. N. (2014) DNA barcode-based delineation of putative species: efficient
756   start for taxonomic workflows. *Mol Ecol Resour* **14**(4), 706-15.
757
758   Kekkonen M., Mutanen M., Kaila L., Nieminen M., Hebert P. D. (2015) Delineating species with
759   DNA barcodes: a case of taxon dependent method performance in moths. *PLOS ONE* **10**(4),
760   e0122481.
761
762   King R. A., Bradford T., Austin A. D., Humphreys W. F., Cooper S. J. (2012) Divergent molecular
763   lineages and not-so-cryptic species: the first descriptions of stygobitic chiltoniid amphipods
764   (Talitroidea: Chiltoniidae) from Western Australia. *Journal of Crustacean Biology* **32**(3), 465-488.
765
766   King R. A., Fagan-Jeffries E. P., Bradford T. M., Stringer D. N., Finston T. L., Halse S. A., Eberhard
767   S. M., Humphreys G., Humphreys B. F., Austin A. D. (2022) Cryptic diversity down under: defining
768   species in the subterranean amphipod genus Nedsia Barnard & Williams, 1995 (Hadzioidea:
769   Eriopisidae) from the Pilbara, Western Australia. *Invertebrate Systematics* **36**(2), 113-159.
770
771   Koch F., Blum P., Korbel K., Menberg K. (2024) Global overview on groundwater fauna.
772   *Ecohydrology* **17**(1), e2607.
773
774   Korbel K., Chariton A., Stephenson S., Greenfield P., Hose G. C. (2017) Wells provide a distorted
775   view of life in the aquifer: implications for sampling, monitoring and assessment of groundwater
776   ecosystems. *Sci Rep* **7**, 40702.
777

Kuang X., Liu J., Scanlon B. R., Jiao J. J., Jasechko S., Lancia M., Biskaborn B. K., Wada Y., Li H.,
Zeng Z. (2024) The changing nature of groundwater in the global water cycle. *Science* **383**(6686),
eadf0630.

Kvist S. (2013) Barcoding in the dark?: a critical view of the sufficiency of zoological DNA
barcoding databases and a plea for broader integration of taxonomic knowledge. *Molecular*
Phylogenetics and Evolution **69**(1), 39-45.

Lahaye R., Van der Bank M., Bogarin D., Warner J., Pupulin F., Gigot G., Maurin O., Duthoit S.,
Barraclough T. G., Savolainen V. (2008) DNA barcoding the floras of biodiversity hotspots.
Proceedings of the National Academy of Sciences **105**(8), 2923-2928.

Laumer C. (2023) Picogram input multimodal sequencing (PiMmS) Version 1
dx.doi.org/10.17504/protocols.io.rm7vzywy5lx1/v1

Little J., Schmidt D. J., Cook B. D., Page T. J., Hughes J. M. (2016) Diversity and phylogeny of
south-east Queensland Bathynellacea. *Australian Journal of Zoology* **64**(1), 36-47.

Mammola S., Altermatt F., Alther R., Amorim I. R., Băncilă R. I., Borges P. A., Brad T., Brankovits
D., Cardoso P., Cerasoli F. (2024) Perspectives and pitfalls in preserving subterranean biodiversity
through protected areas. *npj Biodiversity* **3**(1), 2.

Mammola S., Cardoso P., Culver D. C., Deharveng L., Ferreira R. L., Fišer C., Galassi D. M.,
Griebler C., Halse S., Humphreys W. F. (2019) Scientists' warning on the conservation of
subterranean ecosystems. *BioScience* **69**(8), 641-650.

Mammola S., Lunghi E., Bilandžija H., Cardoso P., Grimm V., Schmidt S. I., Hesselberg T., Martínez
A. (2021) Collecting eco evolutionary data in the dark: Impediments to subterranean research and
how to overcome them. *Ecology and evolution* **11**(11), 5911-5926.

Matthews E. F., Abrams K. M., Cooper S. J., Huey J. A., Hillyer M. J., Humphreys W. F., Austin A.
D., Guzik M. T. (2020) Scratching the surface of subterranean biodiversity: molecular analysis reveals
a diverse and previously unknown fauna of Parabathynellidae (Crustacea: Bathynellacea) from the
Pilbara, Western Australia. *Molecular Phylogenetics and Evolution* **142**, 106643.

McCormack J. E., Tsai W. L., Faircloth B. C. (2016) Sequence capture of ultraconserved elements
from bird museum specimens. *Molecular Ecology Resources* **16**(5), 1189-1203.

Meier R., Blaimer B. B., Buenaventura E., Hartop E., von Rintelen T., Srivathsan A., Yeo D. (2022)
A re analysis of the data in Sharkey et al.'s (2021) minimalist revision reveals that BINs do not
deserve names, but BOLD Systems needs a stronger commitment to open science. *Cladistics* **38**(2),
264-275.

Meierotto, S., Sharkey, M. J., Janzen, D. H., Hallwachs, W., Hebert, P. D., Chapman, E. G., & Smith,
M. A. (2019). A revolutionary protocol to describe understudied hyperdiverse taxa and overcome the
taxonomic impediment. *Deutsche Entomologische Zeitschrift*, 66(2), 119-145.

Meyer C., Kreft H., Guralnick R., Jetz W. (2015) Global priorities for an effective information basis
of biodiversity distributions. *Nature Communications* **6**(1), 1-8.

Miller J. A., Beentjes K. K., van Helsdingen P., IJland S. (2013) Which specimens from a museum
collection will yield DNA barcodes? A time series study of spiders in alcohol. *Zookeys*(365), 245.

Moreau C. S., Wray B. D., Czekanski-Moir J. E., Rubin B. E. (2013) DNA preservation: a test of
commonly used preservatives for insects. *Invertebrate Systematics* **27**(1), 81-86.

Nichols S. J., Kefford B. J., Campbell C. D., Bylemans J., Chandler E., Bray J. P., Shackleton M.,
Robinson K. L., Carew M. E., Furlan E. M. (2020) Towards routine DNA metabarcoding of
macroinvertebrates using bulk samples for freshwater bioassessment: Effects of debris and storage
conditions on the recovery of target taxa. *Freshwater Biology* **65**(4), 607-620.

Ortiz D., Francke O. F. (2017) Reconciling morphological and molecular systematics in tarantulas
(Araneae: Theraphosidae): revision of the Mexican endemic genus Bonnetina. *Zoological Journal of*
the Linnean Society **180**(4), 819-886.

Page T. J., Hughes J. M., Real K. M., Stevens M. I., King R. A., Humphreys W. F. (2018) Allegory of
a cave crustacean: systematic and biogeographic reality of Halosbaena (Peracarida:
Thermosbaenacea) sought with molecular data at multiple scales. *Marine Biodiversity* **48**, 1185-1202.

Palumbi S. R. (1996) Nucleic acids II: the polymerase chain reaction. In 'Molecular Systematics.' (Eds
DM Hillis, C Moritz and BK Mable). (Sinauer Associates: Sunderland, MA)

Park N., Betteridge E., Thurston S., Tuameh A., Mosca M., da Conceicoa L. P., Johnston I.,
Lawniczak M. (2023) High-throughput DNA barcoding library construction and sequencing protocol
for BIOSCAN using unpurified non-destructively extracted DNA from arthropods.
dx.doi.org/10.17504/protocols.io.8epv5jzxdl1b/v1.

Perina G., Camacho A., Huey J., Horwitz P., Koenders A. (2018) Understanding subterranean
variability: the first genus of Bathynellidae (Bathynellacea, Crustacea) from Western Australia
described through a morphological and multigene approach. *Invertebrate Systematics* **32**(2), 423-447.

Perina G., Camacho A. I., Huey J., Horwitz P., Koenders A. (2019) New Bathynellidae (Crustacea)
taxa and their relationships in the Fortescue catchment aquifers of the Pilbara region, Western
Australia. *Systematics and Biodiversity* **17**(2), 148-164.

Pochon X., Bott N. J., Smith K. F., Wood S. A. (2013) Evaluating detection limits of next-generation
sequencing for the surveillance and monitoring of international marine pests. *PLOS ONE* **8**(9),
e73935.

Poore G., Humphreys W. (1992) First record of Thermosbaenacea (Crustacea) from the Southern
Hemisphere: a new species from a cave in tropical Western Australia. *Invertebrate Systematics* **6**(3),
719-725.

Puillandre N., Brouillet S., Achaz G. (2021) ASAP: assemble species by automatic partitioning.
Molecular Ecology Resources **21**(2), 609-620.

Puillandre N., Lambert A., Brouillet S., Achaz G. (2012) ABGD, Automatic Barcode Gap Discovery
for primary species delimitation. *Molecular Ecology* **21**(8), 1864-1877.

875
876 Ratnasingham S., Hebert P. D. (2007) BOLD: The Barcode of Life Data System (http://www.
877 barcodinglife. org). *Molecular Ecology Notes* **7**(3), 355-364.

878
879 Ratnasingham S., Hebert P. D. N. (2013) A DNA-based registry for all animal species: the Barcode
880 Index Number (BIN) system. *PLOS ONE* **8**(e66213).

881
882 Rendoš M., Parimuchová A., Hřívová D. K., Karpowicz M., Papáč V., Jabłońska A., Płóciennik M.,
883 Haviarová D., Grabowski M. (2023) First insight into molecular diversity and DNA barcode library of
884 epikarst-dwelling invertebrates in the Western Carpathians. *Ecohydrology & Hydrobiology*.

885
886 Richly E., Leister D. (2004) NUMTs in sequenced eukaryotic genomes. *Molecular Biology and*
887 *Evolution* **21**(6), 1081-1084.

888
889 Riedel A., Sagata K., Suhardjono Y. R., Tänzler R., Balke M. (2013) Integrative taxonomy on the fast
890 track-towards more sustainability in biodiversity research. *Frontiers in Zoology* **10**, 1-9.

891
892 Rimet F., Aylagas E., Borja Á., Bouchez A., Canino A., Chauvin C., Chonova T., Ciampor Jr F.,
893 Costa F. O., Ferrari B. J. (2021) Metadata standards and practical guidelines for specimen and DNA
894 curation when building barcode reference libraries for aquatic life. *Metabarcoding and Metagenomics*
895 5: e58056. https://doi.org/10.3897/mbmg.5.58056

896
897 Saccò M., Blyth A. J., Douglas G., Humphreys W. F., Hose G. C., Davis J., Guzik M. T., Martínez A.,
898 Eberhard S. M., Halse S. A. (2022a) Stygofaunal diversity and ecological sustainability of coastal
899 groundwater ecosystems in a changing climate: The Australian paradigm. *Freshwater Biology* **67**(12),
900 2007-2023.

901
902 Saccò M., Guzik M. T., van der Heyde M., Nevill P., Cooper S. J., Austin A. D., Coates P. J.,
903 Allentoft M. E., White N. E. (2022b) eDNA in subterranean ecosystems: Applications, technical
904 aspects, and future prospects. *Science of The Total Environment*, 153223.

905
906 Saccò M., Mammola S., Altermatt F., Alther R., Bolpagni R., Brancelj A., Brankovits D., Fišer C.,
907 Gerovasileiou V., Griebler C. (2024) Groundwater is a hidden global keystone ecosystem. *Global*
908 *Change Biology* **30**(1), e17066.

909
Saclier N., Duchemin L., Konecny-Dupré L., Grison P., Eme D., Martin C., Callou C., Lefébure T.,
François C., Issartel C., Lewis J. J., Stoch F., Sket B., Gottstein S., Delić T., Zagmajster M.,
Grabowski M., Weber D., Reboleira A. S. P. S., Palatov D., Paragamian K., Knight L. R. F. D.,
Michel G., Lefebvre F., Hosseini M.-J. M., Camacho A. I., De Bikuña B. G., Taleb A., Belaidi N.,
Tuekam Kayo R. P., Galassi D. M. P., Moldovan O. T., Douady C. J., Malard F. (2024) A
collaborative backbone resource for comparative studies of subterranean evolution: The World
Asellidae database. *Molecular Ecology Resources* **24** (1), e13882.

Sharkey, M.J., Janzen, D.H., Hallwachs, W., Chapman, E.G., Smith, M.A., Dapkey, T., Brown, A.,
Ratnasingham, S., Naik, S., Manjunath, R. and Perez, K., (2021). Minimalist revision and description
of 403 new species in 11 subfamilies of Costa Rican braconid parasitoid wasps, including host records
for 219 species. *ZooKeys* **1013**, 1.

921
922 Simon, C., Frati, F., Beckenbach, A., Crespi, B., Liu, H. and Flook, P., (1994) Evolution, weighting,
923 and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase
924 chain reaction primers. Annals of the entomological Society of America, 87(6), pp.651-701.
925
926 Taberlet P., Coissac E., Pompanon F., Brochmann C., Willerslev E. (2012) Towards next-generation
927 biodiversity assessment using DNA metabarcoding. *Molecular Ecology* **21**, 2045–2050.
928
929 Vamos E., Elbrecht V., Leese F. (2017) Short COI markers for freshwater macroinvertebrate
930 metabarcoding. *Metabarcoding and Metagenomics* **1**: e14625. https://doi.org/10.3897/mbmg.1.14625.
931
932 van der Heyde M., Alexander J., Nevill P., Austin A. D., Stevens N., Jones M., Guzik M. T. (2023a)
933 Rapid detection of subterranean fauna from passive sampling of groundwater eDNA. *Environmental*
934 *DNA* **5**(6), 1706-1719.
935
936 van der Heyde M., White N. E., Nevill P., Austin A. D., Stevens N., Jones M., Guzik M. T. (2023b)
937 Taking eDNA underground: factors affecting eDNA detection of subterranean fauna in groundwater.
938 *Molecular Ecology Resources* 23(6), 1257-1274.
939
940 Villesen P. (2007) FaBox: an online toolbox for fasta sequences. *Molecular Ecology Notes* **7**(6), 965-
941 968.
942
943 Watts C. H. S., Humphreys W. F. (2003) Twenty-five new Dytiscidae (Coleoptera) of the genera
944 *Tjirtudessus* Watts & Humphreys, *Nirripirti* Watts & Humphreys and *Bidessodes* Regimbart, from
945 underground waters in Australia. *Records of the South Australian Museum* **36**, 135-187.
946
947 West K. M., Richards Z. T., Harvey E. S., Susac R., Grealy A., Bunce M. (2020) Under the karst:
948 detecting hidden subterranean assemblages using eDNA metabarcoding in the caves of Christmas
949 Island, Australia. *Scientific Reports* **10**(1), 21479.
950
951 Wetzer R. (2001) Hierarchical analysis of mtDNA variation and the use of mtDNA for isopod
952 (Crustacea: Peracarida: Isopoda) systematics. *Contributions to Zoology* **70**(1), 23-39.
953
954 White N. E., Guzik M. T., Austin A. D., Moore G. I., Humphreys W. F., Alexander J., Bunce M.
955 (2020) Detection of the rare Australian endemic blind cave eel (*Ophisternon candidum*) with
956 environmental DNA: implications for threatened species management in subterranean environments.
957 *Hydrobiologia* **847**, 3201-3211.
958
959 Whiting M. F. (2002) Mecoptera is paraphyletic: multiple genes and phylogeny of Mecoptera and
960 Siphonaptera. *Zoologica Scripta* **31**(1), 93-104.
961
962 Wilkinson M. D., Dumontier M., Aalbersberg I. J., Appleton G., Axton M., Baak A., Blomberg N.,
963 Boiten J.-W., da Silva Santos L. B., Bourne P. E. (2016) The FAIR Guiding Principles for scientific
964 data management and stewardship. *Scientific data* **3**(1), 1-9.
965
966 Wilkinson S. P., Gault A. A., Welsh S. A., Smith J. P., David B. O., Hicks A. S., Fake D. R., Suren A.
967 M., Shaffer M. R., Jarman S. N. (2024) TICI: a taxon-independent community index for eDNA-based
968 ecological health assessment. *PeerJ* **12**, e16963.

969   Wieczorek J., Bloom D., Guralnick R., Blum S., Döring M., Giovanni R, et al. (2012) Darwin Core:
970   An Evolving Community-Developed Biodiversity Data Standard. PLOS ONE **7**(1): e29715.
971
972   Zhang F., Jantarit S., Nilsai A., Stevens M. I., Ding Y., Satasook C. (2018) Species delimitation in the
973   morphologically conserved Coecobrya (Collembola: Entomobryidae): A case study integrating
974   morphology and molecular traits to advance current taxonomy. *Zoologica Scripta* **47**(3), 342-356.