What are the best practices for curating eDNA custom barcode reference libraries? A case study using Australian subterranean fauna.

Michelle T. Guzik[1], Danielle N. Stringer[1], Jake Thornhill[1], Peterson J. Coates[1], Mieke van der Heyde[2], Mia J. Hillyer[3], Nicole E White[2], Mattia Saccò[2], Perry Beasley-Hall[1], William F. Humphreys[4, 5], Mark S. Harvey[4, 5], Joel A. Huey[6], Nerida G. Wilson[4, 5], Jason Alexander [2], Garth Humphreys[4, 7]; Rachael A. King[1, 8], Steve J. B. Cooper[1, 8], Adrian Pinder[9], Giulia Perina[2, 5, 6], Andrew M. Hosie[4], Lisa Kirkendale[4], Paul Nevill[10]; Andy D. Austin[1, 9]

School of Biological Sciences, The University of Adelaide, Adelaide, South Australia, Australia

Subterranean Research and Groundwater Ecology (SuRGE) Group, Trace and Environmental DNA (TrEnD) Lab, School of Molecular and Life Sciences, Curtin University, Perth, Western Australia, Australia

School of Molecular and Life Sciences, Curtin University, Perth, Western Australia, Australia

Collections and Research, Western Australian Museum, Welshpool, Western Australia, Australia.

School of Biological Sciences, University of Western Australia, Crawley, Western Australia, Australia

Biologic Environmental Survey, East Perth, Western Australia, Australia

Biota Environmental Sciences, West Perth Western Australia, Australia

South Australian Museum, North Terrace, Adelaide, South Australia, Australia

Biodiversity and Conservation Science, Department of Biodiversity, Conservation and Attractions, Kensington, Western Australia, Australia

MBioMe - Mine Site Biomonitoring using eDNA Research Group, Trace and Environmental DNA (TrEnD) Laboratory, School of Molecular and Life Sciences, Curtin University, Perth, Western Australia, Australia

Running head - 40 characters: Barcode reference library best practice for subterranean

fauna

Scientific Significance Statement - 125 words: In threatened ecosystems, environmental

DNA (eDNA) metabarcoding for biodiversity discovery and assessment offers considerable

promise for improvement in the rate, efficiency and accuracy of species detection. For a

better understanding of the biodiversity and ecology of organisms detected using eDNA, a

custom library of known reference sequences with associated correct taxonomic metadata is

required. Minimal guidance is currently available on how an effective (i.e. shareable, multi-

sequence, permits metadata and provides a unified nomenclature) custom barcode reference

library (BRL) can be achieved for subterranean fauna. Here, we present a road map for sound

and reliable curation of a BRL using subterranean fauna from Australia as a case study.

Keywords: groundwater, stygofauna, troglofauna, barcode reference library, metabarcoding,

biomonitoring, biodiversity assessment, data infrastructure

Authors, Institutions, and Author Contribution Statement

MTG conceptualised, analysed and wrote the manuscript; DNS, JT, PJC, MVDH, MS

undertook analyses and designed figures; NEW, PB-H, WFH provided valuable expertise;

MJH contributed to laboratory analyses; LK, AMH, JAH, NGW contributed background

knowledge, specimen selection, permissions and access and DNA sequence data; RAK, GP

and AP contributed to taxonomy; JA and GH contributed invaluable field knowledge; MSH

SJBC, PN, ADA contributed to funding. All contributed to writing.

Abstract

Identification of species for environmental assessment and monitoring is essential for

understanding anthropogenic impacts on biodiversity, but for subterranean fauna this task is

frequently difficult and time consuming. The implementation of environmental DNA (eDNA)

metabarcoding for biodiversity discovery and assessment offers considerable promise for

improving the rate, accuracy and efficiency of species detection in ecosystems both above

and below the ground. Importantly, for a better understanding of the biodiversity and ecology

of organisms detected using eDNA, a custom library of known reference sequences with

associated correct taxonomic metadata—i.e., a barcode reference library (BRL)—is required.

Yet, minimal guidance is currently available on how an effective (i.e. shareable, multi-

sequence, that permits metadata and has a unified nomenclature) and accurate (i.e. verified)

custom BRL can be achieved. Here, we present a detailed roadmap for curation of a BRL for

subterranean fauna. To do this, we (1) curated a custom sequence database of subterranean

fauna at an environmentally sensitive location, Bungaroo Creek in the Pilbara region of

Western Australia, for four gene loci useful for eDNA metabarcoding (*COI*, *18S* rRNA, *12S*

rRNA and *16S* rRNA); (2) addressed major gaps in taxonomy and disparate nomenclature of

subterranean fauna by estimating 17–29 putative new species with standard delimitation

methods, including 34 Barcode Index Numbers (BINs) in BOLD, and (3) summarised a best

practice workflow for curation of a custom BRL that has broad applicability and can be

applied to any taxa.

Introduction

Groundwater is a high-value natural resource that is vulnerable to human-induced impacts

and climate change (Kuang *et al.* 2024). In Australia, the economic value of groundwater

resources is estimated to be $33.8 billion (Deloitte Access Economics 2013). Importantly,

groundwater also supports groundwater-dependent ecosystems (GDEs), such as subterranean

habitats, where a unique and ancient fauna provide essential ecosystem services that maintain

underground water condition and quality (Boulton *et al.* 2008). Globally, considerable

tension exists with respect to the use of groundwater for economic growth and environmental

sustainability (Mammola *et al.* 2019; Saccò *et al.* 2022a; Mammola *et al.* 2024; Saccò *et al.*

2024). In certain countries, groundwater fauna are surveyed during routine environmental

impact assessments (EIAs) using biomonitoring methods (Environmental Protection

Authority 2021), particularly in areas of potential resource development (Gibson *et al.* 2019).

For all stakeholders reliant on such biomonitoring, four important issues frequently impact

decision-making capabilities (Gibson *et al.* 2019; Mammola *et al.* 2021; Saccò *et al.* 2022b):

1) inconsistent metadata quality (Costello *et al.* 2013; Saccò *et al.* 2022b); 2) minimal data

sharing (e.g., Meyer *et al.* (2015)); 3) uncertainty around sampling effort required to provide

a robust understanding of the values and predicted impacts and 4) high numbers of

undescribed species (Costello 2015; Ficetola *et al.* 2015) with insufficient associated

taxonomic knowledge (Mammola *et al.* 2019; Mammola *et al.* 2021). Overcoming these

impediments is complicated by reduced taxonomic capacity for many groups and difficulty in

distinguishing species on morphological characteristics alone due to anatomical convergence

and their often-cryptic nature (Abrams *et al.* 2012; King *et al.* 2022).

Environmental DNA (eDNA) metabarcoding has been identified as an approach with

significant potential to alleviate the difficulties associated with traditional sampling (van der

Heyde et al. 2023). Indeed, it has been demonstrated as effective for detection of

subterranean fauna (West *et al.* 2020; White *et al.* 2020; van der Heyde *et al.* 2023a; van der

Heyde *et al.* 2023b). To identify subterranean fauna using eDNA, sequences belonging to

anonymous operational taxonomic units (OTUs) must be assigned taxonomic information

from a reference database (Taberlet *et al.* 2012). However, this assignment is often patchy

due to a lack of data in public DNA databases (e.g., BOLD and GenBank) for both genes and

taxonomically obscure species as well as inconsistencies in taxonomic metadata availability

(Costello 2015; Kvist 2013; Dormontt *et al.* 2018; DeWaard *et al.* 2019; Ficetola *et al.* 2019).

For these reasons, much of the eDNA data being generated by metabarcoding studies are

likely underutilised (Korbel *et al.* 2017). While taxonomy-independent approaches to eDNA

bioinformatics are being developed (Brinkmann *et al.* 2023; Wilkinson *et al.* 2024), many

legislative frameworks for subterranean fauna require species-level identifications for

bioassessment, although more holistic approaches are being advocated (Koch *et al.* 2024).

Therefore, custom reference databases for querying OTUs are critical for the successful

implementation of eDNA metabarcoding methods (Saccò *et al.* 2022b) and resulting

biodiversity assessments. This is especially the case given the extent of short-range

endemism as well as the numerous threatening process that impact subterranean communities

(Harvey *et al.* 2011). Guidance on how these databases should be curated is sorely needed.

Selection criteria for the curation of specimens and DNA sequences when submitting to

public databases are established for certain taxonomic groups of subterranean fauna

(pseudoscorpions (Harvey *et al.* 2023; Hlebec *et al.* 2023); isopods and myriapods (Rendoš *et*

al. 2023); asellid isopods (Saclier *et al.* 2024)). Such criteria include: morphologically

verified specimens; homologous DNA sequences corresponding to common metabarcoding

assay regions; estimating levels of genetic variation within and between species; and detailed

metadata, linking specimens and current taxonomic information per sequence.

Problematically, for understudied and biodiverse groups such as invertebrate fauna, sequence

records frequently lack associated taxonomic information as it often does not exist. Instead, a

unified system of nomenclature that does not solely rely on formally described species is

required. Various methods for delimiting species are already widely used (e.g., Automatic

Barcode Gap Discovery (ABGD; Puillandre *et al.* (2012)), Assemble Species by Automatic

Partitioning (ASAP; Puillandre *et al.* (2021)) and phylogenetic methods such as Bayesian

Poisson Tree Processes (bPTP; Zhang *et al.* (2018)). Such methods require specialist

knowledge to implement, and are frequently not linked to an existing database platform that

is user-friendly, and where the sequences are available in the same place. Whilst some

authors have suggested that these approaches be used as an alternative to formal taxonomic

approaches (Meierotto *et al.* 2019; Sharkey *et al.* 2021), most researchers recognise that a

major problem exists in providing up-to-date and shareable nomenclature in the absence of

taxonomic information.

To combat the taxonomic impediment (Engel *et al.* 2021) facing subterranean fauna

(Mammola *et al.* 2021), delimited lineages may act as a proxy for species until such time as

these species are formally described. The Barcode of Life Database (BOLD) applies a system

of Barcode Index Numbers (BINs), which are an index of unique identifiers based on the

sequences of varying length and specimens that exist in BOLD. The BIN pipeline analyses

new sequence data for the cytochrome *c* oxidase subunit I (*COI*) barcoding gene as they are

uploaded to the database platform. The power of the BIN system is that it provides a unified

nomenclature for sequences entirely independent of the sequence submitter and their

taxonomic capacity, and that information is retained as part of the database. Here, we present

a roadmap for curation of a custom barcode reference library (BRL) for subterranean fauna

using the BOLD system and build on the new and rapidly expanding area of eDNA

metabarcoding research. Our roadmap focuses on an exemplar groundwater-dependent

ecosystem at Bungaroo Creek, located in the resource-rich Pilbara region of Western

Australia (WA). In future, this data infrastructure can be expanded across Australia and

globally in subsequent research since it meets critical principles of a findable, accessible,

interoperable and reusable (FAIR) system of data management outlined by Wilkinson *et al.*

(2016).

Subterranean biodiversity is comprehensively understudied (Guzik *et al.* 2011), but

depending on the region, subterranean fauna can be repeatedly monitored or remain

completely unknown to science (Saccò *et al.* 2022a). Given the articulated need for a

subterranean fauna BRL by multiple stakeholders (i.e. industry, regulators and academic

researchers (Gibson *et al.* 2019)), our overarching aim was to develop a protocol that is easily

accessible and usable for effective and efficient characterisation of subterranean biodiversity.

Our aims were to:

1. Curate a *de novo* custom BRL of sequences for an exemplar subterranean faunal

community of high biodiversity value.

2. Investigate a unified nomenclature that is stable and repeatable, even in the absence of

formal description of species, by using species delimitation methods, such as BINs, to

identify and name species proxies.

3. Develop and implement a best-practice workflow for a future dynamic subterranean fauna

BRL. We outline a protocol for generating and uploading DNA sequence data and associated

metadata for the exemplar location using preferred criteria (i.e. 1–2 listed above).

Methods

Bungaroo Creek, WA: a high biodiversity value subterranean community

The Pilbara region of WA is globally recognised for its significant subterranean fauna (Halse

et al. 2014; Deharveng *et al.* 2024), with many distinct and isolated communities. The Robe

River supports several subterranean communities of high ecological value, including that

below the major tributary known as Bungaroo Creek (Clark *et al.* 2021). Featuring channel

iron deposits and alluvial aquifers, this section of the Robe River provides an ideal habitat for

subterranean fauna. Indeed, evolutionarily distinct taxon groups such as the blind cave eel

(*Ophisternon candidum*), hydrobiid snails, copepods, ostracods, bathynellaceans and unique

amphipod lineages make this environmentally complex location a significant subterranean

community (Clark *et al.* 2021) which is recognised as a Priority Ecological Community

(Priority 1) in Western Australia (Department of Biodiversity, Conservation and Attractions

2023). Due to the use of Bungaroo Creek groundwater to supply water to the West Pilbara

Water Supply Scheme, the subterranean community is frequently sampled, surveyed, and

monitored for assessment of environmental impact. Here, we used the subterranean

community present at Bungaroo Creek as an exemplar location for the curation of a custom

reference library of DNA sequences for future eDNA metabarcoding.

Specimen collection and curation

To curate a *de novo* custom reference library of subterranean fauna from Bungaroo Creek

(herein called Bungaroo BRL), whole specimens from 21 mining exploration bores were

obtained for Sanger sequencing (Figure 1). Initially, samples were obtained from archival

collections of subterranean fauna from surveys undertaken during 2014 and 2017 and

deposited at the Western Australian Museum (WAM). Collection methods included net

hauling (Saccò *et al.* 2022b) from aquifers using pre-existing bore holes. Morphological

identification was undertaken to the lowest possible taxonomic level, with some specimens

identified to morphospecies but most only to order level or higher.

In addition to the archived material from the WAM collection, invertebrates were also

sampled at the Bungaroo Creek borefield site during June 2017 (as above). Bores were

selected using previous survey data with the aim of capturing the greatest possible genetic

diversity within the constraints of time and logistical feasibility. The locations of bores

sampled are provided in Figure 1, Supplementary Material 1. Invertebrate specimens were

taken from each bore using stygofauna haul nets. Five consecutive haul net samples were

collected, each involving lowering of a net to the bottom of the bore, with nets being

sterilised using bleach between sites to prevent contamination. Bulk invertebrate samples

were stored in a portable refrigerator until the end of sampling each day and were then

transferred to a 4°C refrigerator. The samples for each bore were pooled, and specimens were

sorted *in situ* into major taxonomic groups using a dissecting microscope. Specimens of the

same order were each stored in glass vials, labelled with a single unique identifier, and stored

at –20°C in 100% ethanol until DNA extraction. Representative individuals were catalogued

at the WAM for reference (see Supplementary Material 1 for catalogue information

metadata).

Figure 1: Map showing the locations of sampled boreholes at the Bungaroo Creek site, with

bore locations and names. Map created using ArcMap v 10.3.1 (ESRI 2017), data layers

provided by Rio Tinto.

Barcode gene regions

Barcode gene regions are specific segments of DNA sequences that are utilised for molecular

taxonomic identification, particularly in the context of delineating species. This was first

proposed in 2003 (BOLD; Hebert *et al.* (2003)) where standardised DNA sequences were

employed to identify different species. The barcode gene regions selected for this purpose are

typically short (~600-700bp) containing highly conserved regions to allow for primer

binding, which then amplify across a variable region to allow for the differentiation of

species at the molecular level. The most commonly used barcode region for animals is *COI*,

but other genes are also used (https://v3.boldsystems.org/index.php/resources/boldfaq). In

order to curate a useable custom library for Bungaroo Creek, we sequenced three

mitochondrial genes (*COI*, *12S* and *16S*) and one nuclear gene (*18S*). Therefore, we ensured

our BRL would; 1) allow for a strong match to existing metabarcoding assay primers and

provide a reference for the development of new metabarcoding primers (if required); 2)

ensure accuracy and breadth of species identification and documentation of biodiversity (Epp

et al. 2012; Hebert *et al.* 2013; Deagle *et al.* 2014); and 3) allow the consolidation of

sequence data from other database libraries, both old and new.

DNA extraction, amplification, and sequencing

DNA was extracted from 468 specimens in December 2017 using a Qiagen DNeasy Blood

and Tissue Kit. Established universal invertebrate primers (below) were used for PCR and

sequencing of specimens, but optimisation of PCR conditions were minimally adjusted

between specimens (Supplementary Material 2). Products were visualised using the eGel

Electrophoresis system (Life Technologies, Scoresby, Victoria, Australia) on 2% agarose gels

with ethidium bromide. Bidirectional long-read Sanger sequencing was carried out at the

Australian Genome Research Facility (Adelaide).

DNA sequence database submissions

For specimens from the Bungaroo BRL, haplotype sequences new to this study were

submitted to GenBank (accession numbers: OR524806-OR524929) and BOLD (public

BOLD Project: BUBRL

https://boldsystems.org/index.php/MAS_Management_DataConsole?codes=BUBRL)

(Supplementary Material 3) including near-complete taxonomy and provenance of the data.

Our rationale for curating and submitting data to both public platforms was that, while

GenBank is the most ubiquitously used to query metabarcoding data due to the BLAST

function, BOLD has many of the GUI features for curating and storing metadata needed for

the current study and any future work. Specifically, BOLD is a long-term funded, user-

friendly web interface with taxonomic, geographic, and photographic metadata available for

each sequence (Ratnasingham and Hebert 2007). In addition, BOLD hosts an *in silico*

delimitation algorithm that assigns identifiers (BINs) to *COI* lineages based on the genetic

distance between all sequences in its database. The BINs provide a centralised system for

identifying lineages that are potentially equivalent to species (Hajibabaei *et al.* 2007). While

BINs frequently do not exist for underrepresented taxonomic groups and regions (Lahaye *et*

al. 2008), we consider that this study fills in some of the gaps in nomenclature and taxonomy

that substantially impact data sharing and transferability between projects. GenBank lacks

most of these features and, because it does not require the association of sequence data with

specimen vouchers, the taxonomy associated with a sequence can never be verified.

Nucleotide analysis and species delimitation

Geneious v10.0.9 (https://www.geneious.com) was used to edit and align *COI*, *12S, 16S* and

18S sequence data for all individuals. Forward and reverse sequence fragments were aligned,

edited and trimmed. *COI* sequencers were translated into amino acids to ensure no stop

codons and a single consensus sequence was created. Consensus sequences of all individuals

for each gene were aligned using MUSCLE (Edgar 2004), trimmed to equal lengths, and

FaBox (Villesen 2007) was used to collapse the final sequence alignments into haplotypes

using its *DNA to haplotype collapser and converter* functionality. Automatic Barcode Gap

Discovery (ABGD) (Puillandre *et al.* 2012), a computationally efficient distance-based

method for species delimitation, was implemented to identify putative species (lineages

equivalents) (https://bioinfo.mnhn.fr/abi/public/abgd/abgdweb.html). This method seeks to

quantify the location of the barcode gap that separates intra- from inter-specific distances. It

performs well when compared to tree-based coalescent methods (Puillandre *et al.* 2012;

Kekkonen and Hebert 2014; Kapli *et al.* 2017) and other threshold techniques (Ratnasingham

and Hebert 2013).

Further to ABGD, we implemented the BOLD BIN method of lineage delimitation. BINs are

an index of unique identifiers based on a database of specimens belonging to each BIN with

their associated metadata as implemented by BOLD systems. The pipeline analyses new

sequence data for the *COI* barcoding region as they are uploaded to BOLD by applying a

two-stage algorithm (Refined Single Linkage) that couples single linkage and Markov

clustering to assign sequences to a sequence cluster and is subsequently assigned a unique

identifier termed a BIN (Ratnasingham and Hebert 2013). The Refined Single Linkage

algorithm matches the taxonomic performance of competing approaches, but couples this

with protocols that are simple enough to allow the automated assignment of all new barcode

records to a BIN. Sequences that establish a new BIN add an entry to the BIN index, while

sequences assigned to an existing BIN contribute their metadata to it.

Results

Summary of haplotypes and genetic distances

A total of 468 individuals were available for PCR and sequencing, but success rates for the

total suite of specimens were low (*COI*: $n = 103$ (22%), *18S*: $n = 137$ (29%), *12S*: $n = 88$

(19%) and *16S*: $n = 21$ (5%)) (see Discussion for explanation). Sequence data were obtained

from a total of 195 individuals (42%), including 660 base pairs (bp) for *COI* ($n = 103$ and $h =$

59), 1017 bp for *18S* ($n = 136$ and $h = 58$), 425 bp for *12S* ($n = 88$ and $h = 48$), and 511 bp for

16S ($n = 21$ and $h = 11$). Only seven individuals were successfully sequenced for all four

barcoding loci (see Supplementary Material 1). All 195 sequences were submitted to our

BOLD project ID BUBRL (Bungaroo BRL) for four barcode genes *COI*-5P (103 sequences),

12S (88), *16S* (21), and *18S* (136), (GenBank accession numbers *COI*: OR524806 -

OR524908; *18S*: OR524930 - OR525066; *12S*: OR525144 - OR525231; *16S*: OR524909-

OR524929) and BOLD (public BOLD Project:

https://boldsystems.org/index.php/MAS_Management_DataConsole?codes=BUBRL).

Species delimitation

Haplotype groups equivalent to lineages were delineated using ABGD (Figure 2 and

Supplementary Material 4) and compared to BOLD BINs as well as to morphospecies

identified by Bennelongia Ltd, an environmental consulting company that undertook routine

biomonitoring in the area during a 2017 survey at Bungaroo Creek (Table 1). Results of the

ABGD analysis revealed 29 lineages for *COI*, 16 lineages for *12S,* 9 lineages for *16S* and 17

lineages for *18S*. Thirty-four BINs were identified for *COI*. Delineated lineages based on

ABGD (for all four genes) and BOLD BINs (based on *COI* only) for major taxon groups

included amphipods (ABGD = 4–8 lineages; BINs = 9), bathynellaceans (ABGD = 1–4

lineages; BINs = 4), thermosbenaceans (ABGD = 1–2 lineages; BINs = 3), annelids (ABGD

= 1–6 lineages; BINs = 5), ostracods (ABGD = 1–3 lineages; BINs = 3) copepods (ABGD =

4–5 lineages; BINs = 8), and a single lineage of gastropod. It should be noted that Pauropoda

are most likely troglophilic/ edaphobitic but were retained here as representatives of the

subterranean ecosystem at Bungaroo Creek.

Figure 2: Phylogenetic tree of taxa from Bungaroo creek for the mitochondrial *COI* gene–
with species delimitation results represented by taxonomic group in colour: Amphipoda
(blue), Copepoda (purple), Bathynellacea (green), Thermosbaenacea (pink), Ostracoda (red),
Annelida (orange), Gastropoda (dark blue). Relationships between major taxonomic groups
are not representative of actual phylogenetic relationships.

1 Table 1: Morphospecies and delimited lineages of subterranean fauna from Bungaroo Creek, Western Australia (for lineages see Figure 2) using ABGD and
2 BOLD BINs for all four barcoding genes examined in this study.

High level rank	Lineage	Morphological ID	COI Group ID		BIN	Number of ABGD groups			
			ABGD	BIN	BIN	COI	18S	12S	16S
Gastropoda (class)	Hydrobiidae		17	AEE7100	1	1	1	1	1
Pauropoda (class)	Pauropodidae		4	AEE8965	1	1	0	1	0
Annelida (superclass)	Phreodrilidae		12	AEE7562	5	5	2	3	6
	Phreodrilidae		11	AEE7131					
	Unknown lineage		10	AEE1055					
	Tubificidae (now Naididae)		8	AEE9005					
	Phreodrilidae		9	AEE7132					
Amphipoda (order)	Eriopisidae	*Nedsia* sp.	29	AEE9183	9	8	5	4	0
				AEE7196					
	Eriopisidae	*Nedsia* sp.	3	AEE7280					
	Unknown lineage		2	AEE5579					
	Eriopisidae	*Pilbarana* sp.	1	AEE8991					
	Paramelitidae		13	AEE8550					
	Paramelitidae		14	AEE9924					

	Unknown lineage		15	AEE2761					
	Bogidiellidae	*Bogidiella* sp.	16	AEE4462					
Bathynellacea (order)	Parabathynellidae	nr *Billibathynella*	23	AEE8197	4	4	1	3	0
	Parabathynellidae	nr *Billibathynella* sp.	22	AEE5164					
	Parabathynellidae	nr *Billibathynella*	21	AEE0507					
	Parabathynellidae	*Hexabathynella* sp.	20	AEE2039					
Thermosbaenacea (order)	Thermosbaenacea		19	AEF0046	3	2	1	2	0
				ADX1249					
	Thermosbaenacea	*Halosbaena* sp.	18	AEE4740					
Ostracoda (order)	Unknown lineage		7	AEE0613	3	3	1	0	2
	Unknown lineage		6	AEE1979					
	Unknown lineage		5	AEE0612					
Copepoda (subclass)	Cyclopoida	*Diacyclops* nr *humphreysi humphreysi*	26	AEE7353	8	5	4	0	0
	Cyclopoida	*Diacyclops* sp.	25	AEE2774					
	Cyclopoida	*Halicyclops* (*Rochacyclops*) *rochai*	28	AEE3241					

				AEE3240				
Harpacticoida		*Elaphoidella humphreysi*	27	AEE7339				
Calanoida		*Stygoridgewayia trispinosa*	24	AEE5020				
				AEE5021				
				AEE2775				
Isopoda (order)	Cirolanidae	*Kagalana tonde*	-		0	1	1	0
Polyxenida (order)	Lophoproctidae	*Lophoturus madecassus*	-		0	1	1	0
Total					**34**	**29**	**17**	**16**	**9**

Discussion

The use of eDNA metabarcoding to estimate the biodiversity of subterranean faunal communities has the capacity to alleviate a bottleneck that morphological examination of physical specimens, limited taxonomic expertise, low-quality or non-existent taxonomic keys, and a poorly described fauna have on biodiversity assessment (Gibson 2019). The most efficient way to identify species from anonymous eDNA metabarcoding OTUs is to employ a curated reference library of sequences (barcodes), preferably with matching voucher specimens and/or taxonomic information (i.e., a custom BRL). Here, we developed preferred sequence data and metadata criteria for submission to BOLD for subterranean fauna. This best-practice protocol addresses a suite of issues specific to subterranean fauna by applying a unified nomenclature to the exceptional and under-described biodiversity that exists in groundwater-dependent ecosystems, such as the exemplar site chosen for this study. Further, our Bungaroo BRL forms the basis of a publicly available data resource that can be used in eDNA based studies of subterranean fauna into the future.

Estimating community diversity at Bungaroo Creek

Combined, the four genes sequenced here for Bungaroo Creek subterranean fauna delineated up to 29 genetic lineages as estimated with ABGD based on the *COI* data for 17 taxon groups. The majority of these lineages were further supported by ABGD results for the other three genes. These putative species included mites, amphipods, rotifers, copepods, isopods, worms, snails, and millipedes, indicating a substantially speciose invertebrate community. The four genes were effective for sequencing the key taxon groups present, but there was some variation in the number of delineated lineages between genes. For instance, *COI* delineated the highest number of lineages (29), followed by *18S* (17), *12S* (16) and *16S* (9). Some of the variation in numbers of delineated species was attributable to sequencing success or lack thereof (e.g. *16S*, see below) (e.g. Asmyhr and Cooper (2012)), as well as a wide range of resolution given the diversity of markers employed, but the delineation methods used here may also reflect strong phylogeographic variation and structuring (Ortiz and Francke 2017).

Compared to other studies, 29 lineages of subterranean fauna represent a significantly more diverse community at Bungaroo than for other Australian aquifers, although this number is consistent with biodiversity assessment surveys based on morphological examinations (e.g. Bennelongia (2013)). For example, we observed between two and three lineages of ancient

Thermosbaenacea, whereas only one species is currently described from the Pilbara (Poore

and Humphreys 1992; Eberhard *et al.* 2005). Our results are supported by (Page *et al.* 2018)

who also detected a suite of genetic lineages indicating a much higher species diversity of

this tethyan crustacean than previously thought. In other Australian aquifers, estimates of 8–

16 putative species have been reported for eastern Australian alluvial aquifers (Asmyhr and

Cooper 2012), and calcrete aquifers at Sturt Meadows (Bradford *et al.* 2014) and Lake Violet

(Watts and Humphreys 2003; Eberhard *et al.* 2005) in the Yilgarn region WA, and the

Millstream aquifer in the Pilbara (Eberhard *et al.* 2005). As such, Bungaroo Creek represents

one of the highest putative species estimates for an Australian aquifer. Alongside our own

study, recent studies by (Perina *et al.* 2018; Perina *et al.* 2019; Matthews *et al.* 2020)

reinforce the view that the Pilbara harbours an exceptionally high and little-explored diversity

of subterranean faunal species.

BINs

Barcode Index Numbers (BINs) from genetic databases (i.e., BOLD) have been promoted to

offer a standardised and globally unique identifier for each species or species proxy,

enhancing comparability and repeatability of studies, allowing researchers to pool data and

conduct meta-analyses (Hajibabaei *et al.* 2007; Ratnasingham and Hebert 2007; Lahaye *et al.*

2008). In particular, BINs, based on the genetic distance between sequences independent of

prior taxonomic knowledge, have the potential to help identify cryptic or overlooked species

to provide an objective means for identifying species boundaries (Hajibabaei *et al.* 2007).

Here, ABGD and BIN estimates of putative species for the Bungaroo BRL were generally

consistent, although for the copepods three extra lineages were identified using the latter

system. While in some cases BINs have been recognised to have some detrimental aspects in

certain cases. For example, over-splitting and potential misidentification of lineages is

recognised as problematic (Deagle *et al.* 2014) (i.e. BIN assignment has an 86% success rate

(Hartop *et al.* 2022)). Further, for unsampled taxon groups and geographic locations where

representation in BOLD is absent, existing BINs do not exist, leading to the creation of many

new BINs when sampling increases (Lahaye *et al.* 2008). Finally, BIN assignments can

change over time as new sequences and taxon groups are added. However, for large, well-

represented BINs, the changes are often small and can be tracked by BOLD (Ratnasingham

and Hebert 2013). An important advantage of BINs is their usefulness in providing a unified

nomenclature that is publicly available and usable. If required, BINs can be verified using

current taxonomy or alternative species delimitation methods (i.e., via morphological and/or

molecular methods) and updated on BOLD if needed.

Recommended workflow for Barcode Reference Library sequence/data additions

Using Bungaroo Creek as an example, we have demonstrated how to create a dynamic subterranean

faunal reference library for ongoing and future research and biomonitoring that can be updated and,

importantly, follows FAIR Principles of Data Infrastructure (Wilkinson et al. 2016). Below we make

recommendations for a suitable workflow for addition of data to our BRL for subterranean fauna (i.e.

illustrative summary, Figure 3).

Figure 3: Recommended workflow for construction of a Barcode Reference Library.

SPECIMENS

Handling and sorting: in-field and laboratory

Animal sampling requires compliance with regional collection standards and permits. Tissues

should be handled according to those standards. Here, we address handling of invertebrates

and their tissue specifically since sampling of subterranean vertebrate specimens pose

substantially different requirements (Saccò et al. 2022b).

Development of a BRL first requires long-strand sequencing of DNA (Sanger or next

generation sequencing (NGS)) from reference specimens that are also suitable for

morphological examination either pre- or post-sequencing. Overall, the quality of specimen

preservation for DNA sequencing is critical to the successful development of a reference

library. Snap-freezing of specimens is the most optimal for long-term storage of DNA but

difficult to achieve in the field, especially at remote locations. Instead, under ideal conditions,

specimens should be kept alive and then killed by sorting into cold 100% undenatured

ethanol or propylene glycol with no other additives for preservation, and them immediately

refrigerated at a minimum of –20° C (Moreau *et al.* 2013). It should be noted that for small

specimens, especially microcrustaceans, sorting in a laboratory with good lab practises,

ergonomic consideration to prolonged microscopy and quality equipment can yield improved

results. Storage of specimens at room temperature is not ideal but may be unavoidable

(Nichols *et al.* 2020). For many surveys, it is not feasible to spend extended time in the field,

and bulk samples are frequently transported back to the laboratory for sorting. In these

situations, it is recommended that the maximum amount of water from bulk samples be

removed prior to sorting, with animals sorted live where possible, and the organic material is

refrigerated to the lowest temperature possible in 100% ethanol and sorted as soon as possible

(Hajibabaei *et al.* 2012). Clean specimen handling is important, and should be undertaken

with gloves and instruments cleaned in 100% ethanol to control for cross contamination

between specimens. It should be noted that NGS approaches are significantly increasing the

potential of using DNA information from degraded specimens in biodiversity analysis

(Hajibabaei *et al.* 2012).

Morphological identification

For critical taxonomic metadata and establishment of specimen vouchers, morphological

analysis of specimens should always be undertaken when adding data to a BRL. A total

evidence approach (with morphological and DNA sequence data) permits validation of

identified species (whether described or new) and can ultimately be used to establish formal

nomenclature for species. While it is important to remember that any species concept is a

working hypothesis, we specifically note that morphological characters alone do not always

reveal the full extent of species-level genetic diversity in a population or at a location (King

et al. 2012; King *et al.* 2022). Additionally, it may not always be feasible to identify

specimens to species or even genus level due to a lack of taxonomic expertise and/or ongoing

funding, the presence of cryptic species, inadequate taxon sampling, the collection of various

life-history stages or damaged specimens that make taxonomic identification difficult, and

taxonomic uncertainty and/or incompleteness, especially in poorly-explored geographic

locations. High-quality images of putative morphotypes taken prior to DNA extraction can be

used as a record of morphological features in the event of destructive sampling and can be

uploaded to BOLD. While new techniques are being developed to retain the exoskeleton of

specimens during DNA extraction and library preparation (e.g. acoustic droplet ejection (Cain-

Hom *et al.* 2016)), currently this technology is highly specialised and expensive to
implement. However, ongoing development by major consortia, such as the BIOSCAN
project (Park *et al.* 2023), has the potential to expand the application of these methods and
decrease the cost of widespread implementation. In the interim, intact mature animals should
ideally be morphologically identified where possible prior to sequencing to yield
morphological and molecular data linked within the same specimen.

Voucher specimens

Retention of specimen vouchers for future morphological validation, and description is vital,
and further DNA sequencing can aid with verification of genetic lineages. Archived
specimens and tissues are valuable as a temporal genetic and morphological record of the
status of a species. Specimen storage (Burrell *et al.* 2015) and age of specimens (Miller *et al.*
2013) are important parameters to successful DNA extraction and PCR amplification and can
significantly affect the task of building a BRL (Baird *et al.* 2011; Hebert *et al.* 2013; Burrell
et al. 2015). In order to retain specimens, dissection and extraction of DNA from a part of the
body that will minimally interfere with morphological identification is ideal and usually
feasible. However, for very small individuals, sequencing of one to five representatives from
a group of morphologically identical individuals collected from the same sampling site allows
remaining individuals to be treated as a voucher specimen. Recent technical developments
have enabled whole genome sequencing of single individuals for tiny species using genome
amplification approaches (Picogram input multimodal sequencing (PiMmS) method; Laumer
(2023)), but these methods are yet to be widely employed by the biodiversity community.

DNA SEQUENCE DATA
Reference sequences: specimen identification with DNA sequencing

DNA sequencing is a powerful tool for identifying specimens to species and genus levels,
particularly when traditional morphological or biochemical methods are inadequate or
inconclusive. Depending on the gene of interest, a minimum of one sequence (i.e. a single
specimen) is recommended to define an OTU and sequencing multiple individuals from a
population will provide an indication of diversity within a species. Obtaining sequences from
multiple genes also increases confidence in the accuracy of the identification, as any
sequencing errors (particularly relevant for Sanger sequencing since errors caused by nuclear
copies of mitochondrial DNA (NUMTs are a possibility (Richly and Leister 2004)), or
contamination can be detected and potentially corrected. Proper retention and storage of raw

reads is important for verification and reproducibility of the data. Raw data, such as trace

files, can be uploaded to BOLD alongside additional metadata (see below).

Primers

Primer mismatch is known to be a major issue in amplification of barcode fragments.

Universal primers (LCOI490/HCO2198 (Folmer *et al.* 1994) and C1-J-1718 (Simon *et al.*

1994)) are effective for *COI* amplification for representatives of many animal phyla (Hebert

et al. 2003). However, in agreement with past studies, we observed that these standard

primers can amplify inconsistently across different groups in a variety of subterranean

crustacean taxa (e.g. Camacho *et al.* 2011; Karanovic and Cooper 2012; Asmyhr and Cooper

2012). The universal primers for the nuclear *18S* (Whiting 2002) and mitochondrial *12S*

rRNA (Wetzer 2001) genes are known to amplify well in many crustacean taxa (Asmyhr and

Cooper 2012) and worked well here to produce alternative barcodes (Corse *et al.* 2010;

Hardy *et al.* 2011). The standard *16S* primers (16sar/16Sbr Palumbi (1996)), whilst not

successful for many taxa, were extremely useful for gastropods and copepods in this study.

For these reasons, many studies develop *ad hoc* primers depending on the taxon group (e.g.,

Little *et al.* (2016)). In future, it will become increasingly feasible and affordable to sequence

whole mitochondrial genomes employing next-generation sequencing (e.g., eDNA

metagenomics), obtaining many barcoding genes at once or, alternatively, using the entire

mitochondrial genome for species delineation or identification.

For a BRL to be useful for implementation into future eDNA metabarcoding studies, the

barcodes of voucher specimens should match the most commonly used assay regions. Due to

the historical technological impediment of the last 20 years where Sanger sequencing of

barcode loci was the most inexpensive, subterranean fauna have most commonly been

sequenced for loci including *COI*, *16S, 12S* and *18S,* in order of frequency. BRL DNA

sequences should be homologous to metabarcoding assay regions (i.e., standard barcodes).

Currently, *COI* and *18S* are most commonly implemented for metazoan metabarcoding

detection, and *16S* for microbial detection from eDNA. Commonly used primers for eDNA

metabarcoding assays of freshwater invertebrates include *18S* (Pochon *et al.* 2013), *16S* (such

as for crustaceans (Berry *et al.* 2017) and insects (Clarke *et al.* 2014)) and *COI* (Vamos *et al.*

2017). In future, genome skimming, including shotgun sequencing of mitochondrial genomes

and Ultra Conserved Elements, may effectively be used as a universal approach for

generating sequence data, especially for genuinely old and degraded specimens (Blaimer *et al.* 2016; McCormack *et al.* 2016) but also for contemporary specimens.

METADATA

Linking specimen voucher information to DNA sequence and best available taxonomy

For an effective custom BRL database, the link between DNA sequences, morphologically identified specimens by a taxonomic practitioner (where possible), specimen metadata and the most up to date available taxonomy, which can be linked to sequence data are critical to its usability. A comprehensive guide to curation of a BRL for aquatic organisms (eukaryotic and prokaryotic) is described by Rimet *et al.* (2021) who recommended BOLD for its ability to link varied metadata to a sequence. For BOLD, minimum information per DNA sequence is: Sample ID (collection code), Field ID and/or Museum voucher ID (voucher number), Institution Storing, Phylum, and Country. However, it is strongly recommended that latitude and longitude (sample locality), morphological identification to lowest taxonomic level (Rimet *et al.* 2021), the name of the taxonomist, and the year of the identification also be included. BOLD allows submission of trace files for sequences and images (https://boldsystems.org/index.php/databases), which enhances accessibility and reusability. However, required information for GenBank is minimal and the infrastructure needed for hosting additional data, such as chromatograms and images, is not available, making it difficult to provide comprehensive metadata per specimen.

DATABASE

Unified nomenclature and species delimitation

As mentioned previously, BOLD has developed the BIN system for *COI* sequences. The validity of using BINs as equivalent to species has been debated in the literature (Kekkonen *et al.* 2015; Meier *et al.* 2022). This is especially the case since use of *COI* only in conjunction with the OTU designation method of BOLD (i.e. the Refined Single Linkage (RESL) algorithm (Ratnasingham and Hebert 2013)), there is a tendency for over-splitting of lineages (Hlebec *et al.* 2023). However, we consider, as in other studies (e.g. Kekkonen *et al.* 2015; Blagoev *et al*. 2016), the utility of BINs as a built-in clustering method for generating species hypotheses is a valid approach, particularly for use by non-specialists. We observed that BINs, in conjunction with the ABGD species delimitation approach and gene fragment concordance to be consistent and in-line with the findings of other studies (Kekkonen *et al.*

2015). In future, the 29 lineages and 34 BINs we resolved here may be verified as

undescribed species, with morphological examination of specimen vouchers.

GenBank, as the most widely used DNA sequence repository, can be used for querying

metabarcoding reads and OTUs obtained in eDNA studies. Whilst this database is useful,

especially given its centralization, the implementation and accessibility of metadata (e.g.,

taxonomic, geographic and photographic) associated with sequences means that there are

added advantages to BOLD. Designed principally for biodiversity studies, the GUI of BOLD

is uniquely configured to summarise geographic and taxonomic data. For example,

phylogenetic trees can be directly estimated from DNA sequences, and the data can be stored

in various projects and databases. Further, these aspects have the added benefit of being user-

friendly for non-specialists, and provide an accessible implementation of the subterranean

fauna BRL presented here. BOLD is also one of the most usable and collaborative databases

available for depositing and analysing eDNA data. Whilst other databases, such as GenBank,

have projects and databases (e.g. BioProjects, BioSamples, and the Sequence Read Archive),

there are distinct advantages to using BOLD since, in addition to its ongoing funding, it is

extremely attractive for the accessibility of custom BRLs for future research and, as such, we

recommend its ongoing use for accessing our subterranean fauna database.

The Darwin Core Standard (DwC) offers a stable and flexible framework for compiling

biodiversity data from varied and variable sources. Originally developed by the Biodiversity

Information Standards community, Darwin Core is an evolving platform providing

definitions and common terms that are shareable and searchable (Wieczorek *et al.* 2012). It

plays a fundamental role in the sharing, use and reuse of open-access biodiversity data. We

recommend that in curating data for custom BRL's, the DwC standard terms be used.

Importantly, BOLD , but not GenBank, implements the DwC standard (Ratnasingham and

Hebert 2007).

Future taxonomic considerations

The usefulness of a subterranean fauna BRL rests in its ability to assign OTUs a unified

nomenclature and, where possible, species names. Reference sequences must be directly

linked to vouchered specimens of each species that are curated and maintained long-term

within museum collections. For conservation management, species names are also critical

(Costello *et al.*, 2015), but as is the case for most invertebrate groups, especially subterranean

fauna, formal taxonomy is not progressing rapidly enough compared to the recognition of the

burgeoning number of molecular OTUs (Perina *et al.* 2018; Perina *et al.* 2019; Matthews *et*

al. 2020). The number of new lineages we recovered in this study at Bungaroo Creek

highlight the need for a unified nomenclature that is used consistently among stakeholders

and to which taxonomy can be later applied, when available, to alleviate the well-documented

taxonomic impediments (Riedel *et al.* 2013).

Conclusions

A robust BRL is essential for the effective translation of anonymous DNA sequence data to

meaningful taxonomic identification, including for the detection of fauna using

metabarcoding. However, shareable, reusable and publicly available datasets for poorly

known ecosystems are frequently unavailable for query. To address this lack of fundamental,

but crucially important research, we have presented a custom BRL for an Australian

subterranean fauna as a test case and, in doing so, outline best-practice standards for

establishing a BRL to invigorate future subterranean faunal community assessment and

monitoring. However, these recommendations also apply more broadly to all custom BRLs.

Priorities to transform our consensus of species delimitation results into a more complete

BRL in the future include morphological verification and taxonomic diagnoses of relevant

species. A custom BRL requires DNA sequences at both broad (order, family) and narrow

(genus to population level) taxonomic coverage to be useful in identifying unknown

individuals and sequences in genetic and taxonomic studies (Hebert *et al.* 2013). Additional

verification through more extensive sampling of subterranean fauna, barcoding of a much

broader range of genes, and further phylogenetic analyses will also be essential. Given the

uniquely diverse subterranean fauna present at Bungaroo Creek, and within the Pilbara region

more broadly, a coordinated effort will be required to advance this data infrastructure in the

future.

Acknowledgements

This research was part funded by Australian Research Council (ARC) Industry Linkage

Project Grant LP140100555 to A.D.A. S.J.B.C. and W.F.H. (linkage partners: South

Australian Museum, Western Australian Museum, the WA Department of Biodiversity,

Conservation and Attractions (DBCA) (formerly Department of Parks and Wildlife),

Bennelongia Pty Ltd, Biota Environmental Sciences Pty Ltd), ARC Industry Linkage Project

Grant LP190100555 to A.D.A. (linkage partners: Rio Tinto, BHP, Chevron, South Australian

Museum, Western Australian Museum, WA DBCA, WA Department for Water and

Environmental Regulation, Western Australian Biodiversity Science Institute (WABSI), Rio

Tinto consulting contract to N.W. and M.B., and a research contract with WABSI funded by

Rio Tinto and BHP. We thank S. Halse (Bennelongia) for collaboration and specimen

collection as part of faunal surveys for ecological impact assessments, E. Matthews (formerly

The University of Adelaide) for assistance collecting and sorting specimen samples in the

Robe River Valley, Caitlin O'Neill, Dean Main and Michael Curran (Rio Tinto) for field

support and valuable site information. Specimen processing and handling was undertaken at

WAM by Ana Hara, Corey Whisson and Cathy Carr. Specimens were sampled under a

Department of Parks and Wildlife Regulation 17 Licence to take fauna for scientific purposes

(Number: 08-000700-1). This work was supported by resources provided by the Pawsey

Supercomputing Centre with funding from the Australian Government and the Government

of Western Australia.

References

603 Abrams K., Guzik M., Cooper S., Humphreys W., King R., Cho J.-L., Austin A. (2012) What lies
604 beneath: molecular phylogenetics and ancestral state reconstruction of the ancient subterranean
605 Australian Parabathynellidae (Syncarida, Crustacea). *Molecular Phylogenetics and Evolution* **64**(1),
606 130-144.

607
608 Asmyhr M. G., Cooper S. J. B. (2012) Difficulties barcoding in the dark: the case of crustacean
609 stygofauna from eastern Australia. *Invertebrate Systematics* **26**(6), 583-591.

610
611 Environmental Protection Authority (2021) Technical guidance – Subterranean fauna surveys for
612 environmental impact assessment: Western Australia.

613
614 Baird D. J., Pascoe T. J., Zhou X., Hajibabaei M. (2011) Building freshwater macroinvertebrate
615 DNA-barcode libraries from reference collection material: formalin preservation vs specimen age.
616 *Journal of the North American Benthological Society* **30**(1), 125-130.

617
618 Bennelongia (2013) Bungaroo South: Subterranean Fauna Assessment. Report 2013/189

619
620 Berry T. E., Osterrieder S. K., Murray D. C., Coghlan M. L., Richardson A. J., Grealy A. K., Stat M.,
621 Bejder L., Bunce M. (2017) DNA metabarcoding for diet analysis and biodiversity: A case study
622 using the endangered Australian sea lion (Neophoca cinerea). *Ecology and Evolution* **7**(14), 5435-
623 5453.

624
625 Blaimer B. B., Lloyd M. W., Guillory W. X., Brady S. G. (2016) Sequence capture and phylogenetic
626 utility of genomic ultraconserved elements obtained from pinned insect specimens. *PLOS ONE* **11**(8),
627 e0161531.

628
629 Blagoev, G. A., Dewaard, J. R., Ratnasingham, S., Dewaard, S. L., Lu, L., Robertson, J., Telfer, A. C.
630 and Hebert, P. D. (2016). Untangling taxonomy: a DNA barcode reference library for Canadian
631 spiders. *Molecular Ecology Resources*, **16**(1), 325-341.
632
633 Boulton A. J., Fenwick G. D., Hancock P. J., Harvey M. S. (2008) Biodiversity, functional roles and
634 ecosystem services of groundwater invertebrates. *Invertebrate Systematics* **22**(2), 103-116.

635
636 Bradford T. M., Humphreys W. F., Austin A. D., Cooper S. J. (2014) Identification of trophic niches
637 of subterranean diving beetles in a calcrete aquifer by DNA and stable isotope analyses. *Marine and
638 Freshwater Research* **65**(2), 95-104.

639
640 Brinkmann I., Schweizer M., Singer D., Quinchard S., Barras C., Bernhard J. M., Filipsson H. L.
641 (2023) Through the eDNA looking glass: Responses of fjord benthic foraminiferal communities to
642 contrasting environmental conditions. *Journal of Eukaryotic Microbiology* **70**(4), e12975.

643
644 Burrell A. S., Disotell T. R., Bergey C. M. (2015) The use of museum specimens with high-
645 throughput DNA sequencers. *Journal of human evolution* **79**, 35-44.

647 Cain-Hom C., Pabalate R., Pham A., Patel H. N., Wiler R., Cox J. C. (2016) Mammalian genotyping
648 using acoustic droplet ejection for enhanced data reproducibility, superior throughput, and minimized
649 cross-contamination. *Journal of Laboratory Automation* **21**(1), 37-48.

650
651 Camacho A. I., Dorda B. A., Rey I. (2011) Identifying cryptic speciation across groundwater
652 populations: first COI sequences of Bathynellidae (Crustacea, Syncarida). *Graellsia* **67**(1), 7-12.

653 Clark H. L., Buzatto B. A., Halse, S. A. (2021). A hotspot of arid zone subterranean biodiversity: The
654 Robe Valley in Western Australia. *Diversity*, **13**(10), 482.
655
656 Clarke L. J., Soubrier J., Weyrich L. S., Cooper A. (2014) Environmental metabarcodes for insects: in
657 silico PCR reveals potential for taxonomic bias. *Molecular Ecology Resources* **14**(6), 1160-70.

658
659 Corse E., Costedoat C., Chappaz R., Pech N., Martin J. F., Gilles A. (2010) A PCR‐based method for
660 diet analysis in freshwater organisms using 18S rDNA barcoding on faeces. *Molecular Ecology*
661 *Resources* **10**(1), 96-108.

662
663 Costello M. J. (2015) Biodiversity: the known, unknown, and rates of extinction. *Current Biology*
664 **25**(9), R368-R371.

665
666 Costello M. J., Michener W. K., Gahegan M., Zhang Z.-Q., Bourne P. E. (2013) Biodiversity data
667 should be published, cited, and peer reviewed. *Trends in Ecology & Evolution* **28**(8), 454-461.

668
669 Deagle B. E., Jarman S. N., Coissac E., Pompanon F., Taberlet P. (2014) DNA metabarcoding and the
670 cytochrome c oxidase subunit I marker: not a perfect match. *Biology Letters* **10**(9).

671
672 Deharveng L., Bedos A., Pipan T., Culver D. C. (2024) Global Subterranean Biodiversity: A Unique
673 Pattern. Diversity, **16**(3), 157.

674
675 Department of Biodiversity, Conservation and Attractions (2023) Priority Ecological Communities for
676 Western Australia. Version 35: Perth, Western Australia

677
678 Deloitte Access Economics (2013) Economic Value of Groundwater in Australia. National Centre for
679 Groundwater Research and Training.

680
681 DeWaard J. R., Ratnasingham S., Zakharov E. V., Borisenko A. V., Steinke D., Telfer A. C., Perez K.
682 H., Sones J. E., Young M. R., Levesque-Beaudin V. (2019) A reference library for Canadian
683 invertebrates with 1.5 million barcodes, voucher specimens, and DNA samples. *Scientific Data* **6**(1),
684 308.

685
686 Dormontt E. E., Van Dijk K.-J., Bell K. L., Biffin E., Breed M. F., Byrne M., Caddy-Retalic S.,
687 Encinas-Viso F., Nevill P. G., Shapcott A. (2018) Advancing DNA barcoding and metabarcoding
688 applications for plants requires systematic analysis of herbarium collections—an Australian
689 perspective. *Frontiers in Ecology and Evolution* **6**, 134.

690
691 Eberhard S. M., Halse S. A., Humphreys W. F. (2005) Stygofauna in the Pilbara region, north-west
692 Western Australia: a review. *Journal of the Royal Society of Western Australia* **88**, 167–176.

693
694 Edgar R. C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput.
695 *Nucleic Acids Research* **32** (5), 1792-97.

696
697 Engel, M.S., Ceríaco, L.M., Daniel, G.M., Dellapé, P.M., Löbl, I., Marinov, M., Reis, R.E., Young,
698 M.T., Dubois, A., Agarwal, I. and Lehmann A, P., 2021. The taxonomic impediment: a shortage of
699 taxonomists, not the lack of technical approaches. *Zoological Journal of the Linnean Society* **193**(2),
700 381-387.

701
702 Epp L. S., Boessenkool S., Bellemain E. P., Haile J., Esposito A., Riaz T., Erseus C., Gusarov V. I.,
703 Edwards M. E., Johnsen A., Stenøien H. K. (2012) New environmental metabarcodes for analysing
704 soil DNA: potential for studying past and present ecosystems. *Molecular Ecology* **21**(8), 1821-1833.

705
706 Ficetola G. F., Canedoli C., Stoch F. (2019) The Racovitzan impediment and the hidden biodiversity
707 of unexplored environments. *Conservation Biology* **33**(1), 214-216.

708
709 Ficetola G. F., Pansu J., Bonin A., Coissac E., Giguet-Covex C., De Barba M., Gielly L., Lopes C.
710 M., Boyer F., Pompanon F., Raye G., Taberlet P. (2015) Replication levels, false presences and the
711 estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources*
712 **15**(3), 543-56.

713
714 Folmer O., Black M., Hoeh W., Lutz R., Vrijenoek R. (1994) DNA primers for amplification of
715 mitochondrial cytochrome c oxidase subunit 1 from diverse metazoan invertebrates. *Molecular*
716 *Marine Biology and Biotechnology* **3**(5), 294-299.

717
718 Gibson L., Humphreys W. F., Harvey M., Hyder B., Winzer A. (2019) Shedding light on the hidden
719 world of subterranean fauna: A transdisciplinary research approach. *Science of The Total Environment*
720 **684**, 381-389

721
722 Guzik M. T., Austin A. D., Cooper S. J., Harvey M. S., Humphreys W. F., Bradford T., Eberhard S.
723 M., King R. A., Leys R., Muirhead K. A., Tomlinson M. (2011) Is the Australian subterranean fauna
724 uniquely diverse? *Invertebrate Systematics* **24**(5), 407-418.

725
726 Hajibabaei M., Singer G. A., Hebert P. D., Hickey D. A. (2007) DNA barcoding: how it complements
727 taxonomy, molecular phylogenetics and population genetics. *Trends in Genetics* **23**(4), 167-72.

728
729 Hajibabaei M., Spall J. L., Shokralla S., van Konynenburg S. (2012) Assessing biodiversity of a
730 freshwater benthic macroinvertebrate community through non-destructive environmental barcoding of
731 DNA from preservative ethanol. *BMC Ecology* **12**, 1-10.

732
733 Halse S. A., Scanlon M. D., Cocking J. S., Barron H. J., Richardson J. B., Eberhard S. M. (2014)
734 Pilbara stygofauna: deep groundwater of an arid landscape contains globally significant radiation of
735 biodiversity. *Records of the Western Australian Museum* **78**, 443–483.

736
Hardy C. M., Adams M., Jerry D. R., Morgan M. J., Hartley D. M. (2011) DNA barcoding to support
conservation: species identification, genetic structure and biogeography of fishes in the Murray—
Darling River Basin, Australia. *Marine and Freshwater Research* **62**(8), 887-901.

Hartop E., Srivathsan A., Ronquist F., Meier R. (2022) Towards large-scale integrative taxonomy
(LIT): resolving the data conundrum for dark taxa. *Systematic Biology* **71**(6), 1404-1422.

Harvey, M.S., Rix, M.G., Framenau, V.W., Hamilton, Z.R., Johnson, M.S., Teale, R.J., Humphreys,
G. and Humphreys, W.F., 2011. Protecting the innocent: studying short-range endemic taxa enhances
conservation outcomes. Invertebrate Systematics, 25(1), pp.1-10.

Harvey, M.S., Burger, M.A.A., Abrams, K.M., Finston, T.L., Huey, J.A. and Perina, G. (2023). The
systematics of the pseudoscorpion genus *Indohya* (Pseudoscorpiones: Hyidae) in Australia. *Zootaxa*
5342, 1–119. https://doi.org/10.11646/zootaxa.5342.1

751
752   Hebert P. D., Cywinska A., Ball S. L., deWaard J. R. (2003) Biological identifications through DNA
753   barcodes. *Proceedings of the Royal Society of London, Series B* **270**, 313-322.

754
755   Hebert P. D., Dewaard J. R., Zakharov E. V., Prosser S. W., Sones J. E., McKeown J. T., Mantle B.,
756   La Salle J. (2013) A DNA 'barcode blitz': rapid digitization and sequencing of a natural history
757   collection. *PLOS ONE* **8**(7), e68535.

758
759   Hlebec D., Podnar M., Kučinić M., Harms D. (2023) Molecular analyses of pseudoscorpions in a
760   subterranean biodiversity hotspot reveal cryptic diversity and microendemism. *Scientific Reports*
761   **13**(1), 430.

762
763   Kapli P., Lutteropp S., Zhang J., Kobert K., Pavlidis P., Stamatakis A., T. F. (2017) Multi-rate
764   Poisson tree processes for single-locus species delimitation under maximum likelihood and Markov
765   chain Monte Carlo. *Bioinformatics* **33** (11), 1630-1638.

766
767   Karanovic T., Cooper S. J. (2012) Explosive radiation of the genus Schizopera on a small
768   subterranean island in Western Australia (Copepoda: Harpacticoida): unravelling the cases of cryptic
769   speciation, size differentiation and multiple invasions. *Invertebrate Systematics* **26**(2), 115-192.

770
771   Kekkonen M., Hebert P. D. N. (2014) DNA barcode-based delineation of putative species: efficient
772   start for taxonomic workflows. *Mol Ecol Resour* **14**(4), 706-15.

773
774   Kekkonen M., Mutanen M., Kaila L., Nieminen M., Hebert P. D. (2015) Delineating species with
775   DNA barcodes: a case of taxon dependent method performance in moths. *PLOS ONE* **10**(4),
776   e0122481.

777
778   King R. A., Bradford T., Austin A. D., Humphreys W. F., Cooper S. J. (2012) Divergent molecular
779   lineages and not-so-cryptic species: the first descriptions of stygobitic chiltoniid amphipods
780   (Talitroidea: Chiltoniidae) from Western Australia. *Journal of Crustacean Biology* **32**(3), 465-488.

781
King R. A., Fagan-Jeffries E. P., Bradford T. M., Stringer D. N., Finston T. L., Halse S. A., Eberhard
S. M., Humphreys G., Humphreys B. F., Austin A. D. (2022) Cryptic diversity down under: defining
species in the subterranean amphipod genus Nedsia Barnard & Williams, 1995 (Hadzioidea:
Eriopisidae) from the Pilbara, Western Australia. *Invertebrate Systematics* **36**(2), 113-159.

Koch F., Blum P., Korbel K., Menberg K. (2024) Global overview on groundwater fauna.
Ecohydrology **17**(1), e2607.

789
790   Korbel K., Chariton A., Stephenson S., Greenfield P., Hose G. C. (2017) Wells provide a distorted
791   view of life in the aquifer: implications for sampling, monitoring and assessment of groundwater
792   ecosystems. *Sci Rep* **7**, 40702.

793
794   Kuang X., Liu J., Scanlon B. R., Jiao J. J., Jasechko S., Lancia M., Biskaborn B. K., Wada Y., Li H.,
795   Zeng Z. (2024) The changing nature of groundwater in the global water cycle. *Science* **383**(6686),
796   eadf0630.

797
798   Kvist S. (2013) Barcoding in the dark?: a critical view of the sufficiency of zoological DNA
799   barcoding databases and a plea for broader integration of taxonomic knowledge. *Molecular*
800   *Phylogenetics and Evolution* **69**(1), 39-45.

801
802   Lahaye R., Van der Bank M., Bogarin D., Warner J., Pupulin F., Gigot G., Maurin O., Duthoit S.,
803   Barraclough T. G., Savolainen V. (2008) DNA barcoding the floras of biodiversity hotspots.
804   *Proceedings of the National Academy of Sciences* **105**(8), 2923-2928.

805
806   Laumer C. (2023) Picogram input multimodal sequencing (PiMmS) Version 1
807   dx.doi.org/10.17504/protocols.io.rm7vzywy5lx1/v1

808
809   Little J., Schmidt D. J., Cook B. D., Page T. J., Hughes J. M. (2016) Diversity and phylogeny of
810   south-east Queensland Bathynellacea. *Australian Journal of Zoology* **64**(1), 36-47.

811
812   Mammola S., Altermatt F., Alther R., Amorim I. R., Băncilă R. I., Borges P. A., Brad T., Brankovits
813   D., Cardoso P., Cerasoli F. (2024) Perspectives and pitfalls in preserving subterranean biodiversity
814   through protected areas. *npj Biodiversity* **3**(1), 2.

815
816   Mammola S., Cardoso P., Culver D. C., Deharveng L., Ferreira R. L., Fišer C., Galassi D. M.,
817   Griebler C., Halse S., Humphreys W. F. (2019) Scientists' warning on the conservation of
818   subterranean ecosystems. *BioScience* **69**(8), 641-650.

819
820   Mammola S., Lunghi E., Bilandžija H., Cardoso P., Grimm V., Schmidt S. I., Hesselberg T., Martínez
821   A. (2021) Collecting eco□evolutionary data in the dark: Impediments to subterranean research and
822   how to overcome them. *Ecology and evolution* **11**(11), 5911-5926.

823
824   Matthews E. F., Abrams K. M., Cooper S. J., Huey J. A., Hillyer M. J., Humphreys W. F., Austin A.
825   D., Guzik M. T. (2020) Scratching the surface of subterranean biodiversity: molecular analysis reveals
826   a diverse and previously unknown fauna of Parabathynellidae (Crustacea: Bathynellacea) from the
827   Pilbara, Western Australia. *Molecular Phylogenetics and Evolution* **142**, 106643.

828
829   McCormack J. E., Tsai W. L., Faircloth B. C. (2016) Sequence capture of ultraconserved elements
830   from bird museum specimens. *Molecular Ecology Resources* **16**(5), 1189-1203.

831
Meier R., Blaimer B. B., Buenaventura E., Hartop E., von Rintelen T., Srivathsan A., Yeo D. (2022)
A re□analysis of the data in Sharkey et al.'s (2021) minimalist revision reveals that BINs do not

deserve names, but BOLD Systems needs a stronger commitment to open science. *Cladistics* **38**(2),
264-275.

Meierotto, S., Sharkey, M. J., Janzen, D. H., Hallwachs, W., Hebert, P. D., Chapman, E. G., & Smith,
M. A. (2019). A revolutionary protocol to describe understudied hyperdiverse taxa and overcome the
taxonomic impediment. *Deutsche Entomologische Zeitschrift*, 66(2), 119-145.
839
840 Meyer C., Kreft H., Guralnick R., Jetz W. (2015) Global priorities for an effective information basis
841 of biodiversity distributions. *Nature Communications* **6**(1), 1-8.

842
843 Miller J. A., Beentjes K. K., van Helsdingen P., IJland S. (2013) Which specimens from a museum
844 collection will yield DNA barcodes? A time series study of spiders in alcohol. *Zookeys*(365), 245.

845
846 Moreau C. S., Wray B. D., Czekanski-Moir J. E., Rubin B. E. (2013) DNA preservation: a test of
847 commonly used preservatives for insects. *Invertebrate Systematics* **27**(1), 81-86.

848
849 Nichols S. J., Kefford B. J., Campbell C. D., Bylemans J., Chandler E., Bray J. P., Shackleton M.,
850 Robinson K. L., Carew M. E., Furlan E. M. (2020) Towards routine DNA metabarcoding of
851 macroinvertebrates using bulk samples for freshwater bioassessment: Effects of debris and storage
852 conditions on the recovery of target taxa. *Freshwater Biology* **65**(4), 607-620.

853
854 Ortiz D., Francke O. F. (2017) Reconciling morphological and molecular systematics in tarantulas
855 (Araneae: Theraphosidae): revision of the Mexican endemic genus Bonnetina. *Zoological Journal of*
856 *the Linnean Society* **180**(4), 819-886.

857
858 Page T. J., Hughes J. M., Real K. M., Stevens M. I., King R. A., Humphreys W. F. (2018) Allegory of
859 a cave crustacean: systematic and biogeographic reality of Halosbaena (Peracarida:
860 Thermosbaenacea) sought with molecular data at multiple scales. *Marine Biodiversity* **48**, 1185-1202.

861
862 Palumbi S. R. (1996) Nucleic acids II: the polymerase chain reaction. In 'Molecular Systematics.' (Eds
863 DM Hillis, C Moritz and BK Mable). (Sinauer Associates: Sunderland, MA)

864
865 Park N., Betteridge E., Thurston S., Tuameh A., Mosca M., da Conceicoa L. P., Johnston I.,
866 Lawniczak M. (2023) High-throughput DNA barcoding library construction and sequencing protocol
867 for BIOSCAN using unpurified non-destructively extracted DNA from arthropods.
868 dx.doi.org/10.17504/protocols.io.8epv5jzxdl1b/v1.

869
870 Perina G., Camacho A., Huey J., Horwitz P., Koenders A. (2018) Understanding subterranean
871 variability: the first genus of Bathynellidae (Bathynellacea, Crustacea) from Western Australia
872 described through a morphological and multigene approach. *Invertebrate Systematics* **32**(2), 423-447.

873
874 Perina G., Camacho A. I., Huey J., Horwitz P., Koenders A. (2019) New Bathynellidae (Crustacea)
875 taxa and their relationships in the Fortescue catchment aquifers of the Pilbara region, Western
876 Australia. *Systematics and Biodiversity* **17**(2), 148-164.

877
878 Pochon X., Bott N. J., Smith K. F., Wood S. A. (2013) Evaluating detection limits of next-generation
879 sequencing for the surveillance and monitoring of international marine pests. *PLOS ONE* **8**(9),
880 e73935.

Poore G., Humphreys W. (1992) First record of Thermosbaenacea (Crustacea) from the Southern Hemisphere: a new species from a cave in tropical Western Australia. *Invertebrate Systematics* **6**(3), 719-725.

Puillandre N., Brouillet S., Achaz G. (2021) ASAP: assemble species by automatic partitioning. *Molecular Ecology Resources* **21**(2), 609-620.

Puillandre N., Lambert A., Brouillet S., Achaz G. (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Molecular Ecology* **21**(8), 1864-1877.

Ratnasingham S., Hebert P. D. (2007) BOLD: The Barcode of Life Data System (http://www. barcodinglife. org). *Molecular Ecology Notes* **7**(3), 355-364.

Ratnasingham S., Hebert P. D. N. (2013) A DNA-based registry for all animal species: the Barcode Index Number (BIN) system. *PLOS ONE* **8**(e66213).

Rendoš M., Parimuchová A., Hřívová D. K., Karpowicz M., Papáč V., Jabłońska A., Płóciennik M., Haviarová D., Grabowski M. (2023) First insight into molecular diversity and DNA barcode library of epikarst-dwelling invertebrates in the Western Carpathians. *Ecohydrology & Hydrobiology*.

Richly E., Leister D. (2004) NUMTs in sequenced eukaryotic genomes. *Molecular Biology and Evolution* **21**(6), 1081-1084.

Riedel A., Sagata K., Suhardjono Y. R., Tänzler R., Balke M. (2013) Integrative taxonomy on the fast track-towards more sustainability in biodiversity research. *Frontiers in Zoology* **10**, 1-9.

Rimet F., Aylagas E., Borja Á., Bouchez A., Canino A., Chauvin C., Chonova T., Ciampor Jr F., Costa F. O., Ferrari B. J. (2021) Metadata standards and practical guidelines for specimen and DNA curation when building barcode reference libraries for aquatic life. *Metabarcoding and Metagenomics* 5: e58056. https://doi.org/10.3897/mbmg.5.58056

Saccò M., Blyth A. J., Douglas G., Humphreys W. F., Hose G. C., Davis J., Guzik M. T., Martínez A., Eberhard S. M., Halse S. A. (2022a) Stygofaunal diversity and ecological sustainability of coastal groundwater ecosystems in a changing climate: The Australian paradigm. *Freshwater Biology* **67**(12), 2007-2023.

Saccò M., Guzik M. T., van der Heyde M., Nevill P., Cooper S. J., Austin A. D., Coates P. J., Allentoft M. E., White N. E. (2022b) eDNA in subterranean ecosystems: Applications, technical aspects, and future prospects. *Science of The Total Environment*, 153223.

Saccò M., Mammola S., Altermatt F., Alther R., Bolpagni R., Brancelj A., Brankovits D., Fišer C., Gerovasileiou V., Griebler C. (2024) Groundwater is a hidden global keystone ecosystem. *Global Change Biology* **30**(1), e17066.

Saclier N., Duchemin L., Konecny-Dupré L., Grison P., Eme D., Martin C., Callou C., Lefébure T., François C., Issartel C., Lewis J. J., Stoch F., Sket B., Gottstein S., Delić T., Zagmajster M.,

Grabowski M., Weber D., Reboleira A. S. P. S., Palatov D., Paragamian K., Knight L. R. F. D.,
Michel G., Lefebvre F., Hosseini M.-J. M., Camacho A. I., De Bikuña B. G., Taleb A., Belaidi N.,
Tuekam Kayo R. P., Galassi D. M. P., Moldovan O. T., Douady C. J., Malard F. (2024) A
collaborative backbone resource for comparative studies of subterranean evolution: The World
Asellidae database. *Molecular Ecology Resources* **24** (1), e13882.

Sharkey, M.J., Janzen, D.H., Hallwachs, W., Chapman, E.G., Smith, M.A., Dapkey, T., Brown, A.,
Ratnasingham, S., Naik, S., Manjunath, R. and Perez, K., (2021). Minimalist revision and description
of 403 new species in 11 subfamilies of Costa Rican braconid parasitoid wasps, including host records
for 219 species. *ZooKeys* **1013**, 1.
937
938 Simon, C., Frati, F., Beckenbach, A., Crespi, B., Liu, H. and Flook, P., (1994) Evolution, weighting,
939 and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase
940 chain reaction primers. Annals of the entomological Society of America, 87(6), pp.651-701.
941
942 Taberlet P., Coissac E., Pompanon F., Brochmann C., Willerslev E. (2012) Towards next-generation
943 biodiversity assessment using DNA metabarcoding. *Molecular Ecology* **21**, 2045–2050.

944
945 Vamos E., Elbrecht V., Leese F. (2017) Short COI markers for freshwater macroinvertebrate
946 metabarcoding. *Metabarcoding and Metagenomics* **1**: e14625. https://doi.org/10.3897/mbmg.1.14625.

947
948 van der Heyde M., Alexander J., Nevill P., Austin A. D., Stevens N., Jones M., Guzik M. T. (2023a)
949 Rapid detection of subterranean fauna from passive sampling of groundwater eDNA. *Environmental*
950 *DNA* **5**(6), 1706-1719.

951
952 van der Heyde M., White N. E., Nevill P., Austin A. D., Stevens N., Jones M., Guzik M. T. (2023b)
953 Taking eDNA underground: factors affecting eDNA detection of subterranean fauna in groundwater.
954 *Molecular Ecology Resources* 23(6), 1257-1274.

955
956 Villesen P. (2007) FaBox: an online toolbox for fasta sequences. *Molecular Ecology Notes* **7**(6), 965-
957 968.

958
959 Watts C. H. S., Humphreys W. F. (2003) Twenty-five new Dytiscidae (Coleoptera) of the genera
960 *Tjirtudessus* Watts & Humphreys, *Nirripirti* Watts & Humphreys and *Bidessodes* Regimbart, from
961 underground waters in Australia. *Records of the South Australian Museum* **36**, 135-187.

962
963 West K. M., Richards Z. T., Harvey E. S., Susac R., Grealy A., Bunce M. (2020) Under the karst:
964 detecting hidden subterranean assemblages using eDNA metabarcoding in the caves of Christmas
965 Island, Australia. *Scientific Reports* **10**(1), 21479.

966
967 Wetzer R. (2001) Hierarchical analysis of mtDNA variation and the use of mtDNA for isopod
968 (Crustacea: Peracarida: Isopoda) systematics. *Contributions to Zoology* **70**(1), 23-39.

969
White N. E., Guzik M. T., Austin A. D., Moore G. I., Humphreys W. F., Alexander J., Bunce M.
(2020) Detection of the rare Australian endemic blind cave eel (*Ophisternon candidum*) with
environmental DNA: implications for threatened species management in subterranean environments.
Hydrobiologia **847**, 3201-3211.

Whiting M. F. (2002) Mecoptera is paraphyletic: multiple genes and phylogeny of Mecoptera and
Siphonaptera. *Zoologica Scripta* **31**(1), 93-104.

977
978 Wilkinson M. D., Dumontier M., Aalbersberg I. J., Appleton G., Axton M., Baak A., Blomberg N.,
979 Boiten J.-W., da Silva Santos L. B., Bourne P. E. (2016) The FAIR Guiding Principles for scientific
980 data management and stewardship. *Scientific data* **3**(1), 1-9.

981
982 Wilkinson S. P., Gault A. A., Welsh S. A., Smith J. P., David B. O., Hicks A. S., Fake D. R., Suren A.
983 M., Shaffer M. R., Jarman S. N. (2024) TICI: a taxon-independent community index for eDNA-based
984 ecological health assessment. *PeerJ* **12**, e16963.

985 Wieczorek J., Bloom D., Guralnick R., Blum S., Döring M., Giovanni R, et al. (2012) Darwin Core:
986 An Evolving Community-Developed Biodiversity Data Standard. PLOS ONE **7**(1): e29715.
987
988 Zhang F., Jantarit S., Nilsai A., Stevens M. I., Ding Y., Satasook C. (2018) Species delimitation in the
989 morphologically conserved Coecobrya (Collembola: Entomobryidae): A case study integrating
990 morphology and molecular traits to advance current taxonomy. *Zoologica Scripta* **47**(3), 342-356.