Title: APEX2 proximity labeling of RNA in bacteria

Authors: Hadi Yassine[1,2], Elizabeta Sirotkin[3], Omer Goldberger[3], Vincent Lawal[1], Daniel
B. Kearns[1], Orna Amster-Choder[3], and Jared M. Schrader[1,2]*

[1]Department of Biology, Indiana University, Bloomington, IN, 47405, USA

[2]Departments of Chemistry and Biological Sciences, Wayne State University, Detroit, MI
48202, USA

[3]Department of Microbiology and Molecular Genetics, IMRIC, The Hebrew University
Faculty of Medicine, Jerusalem 91120, Israel

*Corresponding Author: jaschrad@iu.edu

Highlights:

11   ● APEX2 proximity labeling can be applied to RNA in bacteria
12   ● APEX2 RNA labeling reactions occur on the sub-minute timescale.
13   ● APEX2 workflow requires less material and time than current methods.
14   ● Alkyne-Phenol APEX2 substrate provides flexibility with click-chemistry.

Motivation:

Studies over the past several years have shown that distinct RNAs can be targeted
to subcellular locations in bacterial cells. The ability to investigate localized RNAs in
bacteria is currently limited to imaging-based approaches or to laborious procedures to
isolate ribonucleoprotein complexes by grad-seq, HITS-CLIP, or Rloc-seq. However, a
major challenge in studying mRNA localization in bacterial cells is that bacterial mRNAs
typically last for only a few minutes in the cell, while experiments to investigate their
localization or interaction partners can take much longer. Therefore, rapid methods of
studying RNA localization are needed to bridge this technical challenge.

Summary:

Rapid spatially controlled methods are needed to investigate RNA localization in
bacterial cells. APEX2 proximity labeling was shown to be adaptable to rapid RNA labeling
in eukaryotic cells, and through the fusion of APEX2 to different proteins targeted to
different subcellular locations, has been useful to identify RNA localization in these cells.
Therefore, we adapted APEX2 proximity labeling of RNA to bacterial cells by generating an
APEX2 fusion to the RNase E gene, which is necessary and sufficient for BR-body
formation. APEX2 fusion is minimally perturbative and RNA can be rapidly labeled on the
sub-minute timescale with Alkyne-Phenol, outpacing the rapid speed of mRNA decay in

bacteria. Alkyne-Phenol provides flexibility in the overall downstream application with
copper catalyzed click-chemistry for downstream applications, such as fluorescent dye-
azides or biotin-azides for purification. Altogether, APEX2 proximity labeling of RNA
provides a useful method for studying RNA localization in bacteria.

Introduction:

Bacterial mRNAs have been found to localize to distinct subcellular locations. At a
global level, many mRNAs have been found to be associated with the nucleoid,
membrane, or cell poles[1–4] and it has been hypothesized that mRNA localization can be
important for proper gene expression. For example, Flagellin mRNA in *Campylobacter*
jejuni localizes to the cell poles which may facilitate cotranslational flagellar assembly[5]. In
addition, biomolecular condensates, which are non-membrane bound organelles often
assembled through phase separation, have been rapidly expanding in bacterial cells, with
multiple involved in RNA metabolism[4,6]. Yet the functional significance of localization of
RNAs to biomolecular condensates in bacteria has just started to be explored.

GRAD-seq, HITS-CLIP, and Rloc-seq methods[1,7,8] have been successful at
identifying RNAs associated with specific RNPs or localized to certain subcellular
locations. However, a major difficulty is that bacterial mRNAs are very short lived, and
these methods require labeling or isolation times much longer than a typical mRNAs half-
life, leading to the potential for false negatives. Therefore, rapid methods are needed to
isolate the associated labile bacterial mRNA before they are degraded.

APEX2 proximity labeling allows for spatially controlled and rapid RNA labeling[9–12],
making it a useful method to assay for localized bacterial RNAs. In this manuscript, we
show that APEX2 proximity labeling can be adapted to bacterial condensates via fusion of
the core BR-body scaffold RNase E to APEX2. RNA can be labeled robustly by APEX2, as
many bacteria natively produce heme, including *E. coli*, *B. subtilis*, and *C. crescentus*.
With the addition of H_2O_2 and Alkyne-Phenol triggering labeling, allowing copper catalyzed
click chemistry of Alkyne-Phenol labeled RNA with a variety of azides such as Cy5-azide.
This protocol can be used to rapidly label bacterial RNAs by APEX2 on the sub-minute
timescale, and which can be rapidly purified via copper catalyzed click chemistry
conjugation of biotin-peg3-azide followed by streptavidin purification. Through the fusion
of APEX2 to different localized RNA binding proteins in bacteria, APEX2 proximity labeling
has the potential to improve our knowledge of RNA localization in bacteria.

Materials and Methods:

REAGENT or RESOURCE	SOURCE	IDENTIFIER

Antibodies		
DYKDDDDK Tag (D6W5B) Rabbit mAb (Binds to same epitope as Sigma-Aldrich Anti-FLAG M2 antibody) #14793	Cell Signaling Technology	Cat. No.:14793S
Goat anti-Rabbit IgG (H+L) Secondary Antibody, HRP	Invitrogen	Cat. No.: 31460
Bacterial and virus strains		
Caulobacter crescentus NA1000	Lucy Shapiro's Lab	N/A
Js767 (C. crescentus RNE-Apex2FlgC-2 kan^R)	This study	N/A
Js768 (C. crescentus RNE-Apex2Flg-EGFPC-2 kan^R)	This study	N/A
Js87 (C. crescentus RNE-msfGFPC-4 gent^R)	Al-Husini *et al.* Mol Cell 2018[13]	N/A
LS4379 (C. crescentus hfq-m2)	Lucy Shapiro's Lab	N/A
Js801 (RNEΔCTD-Apex2FlgC-2 kan^R)	This study	N/A
B. subtilis DK1042 comIQ12L	Konkol *et al.* J Bact. 2013[14]	N/A
DB2579 (B. subtilis amyE::rny-APEX2 spec^R amp^R)	This study	N/A
Escherichia coli MG1655	Orna Amster-Choder's Lab	N/A
Kti162 (E. coli rne-mCherry)	Strahl *et al.*, PLOS Genetics 2015[15]	N/A
ES413 (E. coli rne-mcherry-Apex2 kan^R)	This study	N/A
Chemicals, peptides, and recombinant proteins		
Alkyne-Phenol	MedChemExpress	Cat. No.: HY-131442

HYDROGEN PEROXIDE SOLN., 50%, ACS RGT.	Sigmaaldrich	Cat. No.: 7722-84-1
Sodium ascorbate	Sigmaaldrich	Cat. No.: 134-03-2
TRIzol	Thermofisher scientific	Cat. No.: 15596018
Chloroform	Thermofisher scientific	Cat. No.: AC423550010
2-PROPANOL, ANHYDROUS	Sigmaaldrich	Cat. No.: 67-63-0
Glycogen, RNA grade	Thermofisher scientific	Cat. No.: RO551
ETHANOL-D6 (D, 99%), ANHYD.	Sigmaaldrich	Cat. No.: 1516-08-1
Sodium Acetate (3 M), pH 5.5, RNase-free	Thermofisher scientific	Cat. No.: AM9740
Biotin-peg3-Azide	Vector Laboratories	CAS Number: CCT-AZ104
Cy5 Azide	Clickchemistrytools	Cat. No.: AZ118-1
BrightStar™-Plus Positively Charged Nylon Membrane, 30 cm x 45 cm	Thermofisher scientific	Cat. No.: AM10102
UltraPure™ SSC, 20X	Thermofisher scientific	Cat. No.: 15557044
SODIUM HYDROXIDE, PELLETS	Sigmaaldrich	Cat. No.: 1310-73-2
EDTA (0.5 M), pH 8.0, RNase-free	Thermofisher scientific	Cat. No.: AM9261

Sodium Dodecyl Sulfate (98.5%)	Sigmaaldrich	Cat. No.: L3771-500G
Bromophenol Blue	sigmaaldrich	Cat. No.: B0126-25G
Glycerol	sigmaaldrich	Cat. No.: 56-81-5
THPTA	Vectorslab	SKU: CCT-1010-100
Copper(II) sulfate pentahydrate	sigmaaldrich	Cat. No.: 7758-99-8
Bovine Serum Albumin	sigmaaldrich	Cat. No.: 9048-46-8
Thermo Scientific™ Pierce™ ECL Western Blotting Substrate	fisher scientific	Cat. No.: PI32109
Heparin sodium salt from porcine intestinal mucosa	sigmaaldrich	Cat. No.: 9041-08-1
Tris (1 M), pH 7.0, RNase-free	Thermofisher scientific	Cat. No.: AM9851
Tris (Tris Base)	GoldBio	Cat. No.: T-400-500
Hydrochloric acid	Carolina	Cat. No.: 867792
SODIUM CHLORIDE, CRYSTAL	Sigmaaldrich	Cat. No.: 7647-14-5
Tween-20	sigmaaldrich	Cat. No.: 9005-64-5
PageRuler™ Plus Prestained Protein Ladder, 10 to 250 kDa	Thermofisher scientific	Cat. No.: 26620
Urea (ACS)	Fisherchemicals	Cat. No.: U15-3

PBS - Phosphate-Buffered Saline (10X) pH 7.4, RNase-free	Thermofisher scientific	Cat. No.: AM9624
Agar	Fisherchemicals	Cat. No.: DF0001-17-0
Bactopeptone	Thermofisher scientific	Cat. No.: 211677
Yeast extract	sigmaaldrich	Cat. No.: 92144-500G-F
UltraPure™ Ethidium Bromide, 10 mg/mL	Thermofisher scientific	Cat. No.: 15585011
Thermo Scientific™ TriTrack DNA Loading Dye (6X)	Thermofisher scientific	Cat. No.: FERR1161
Thermo Scientific™ GeneRuler 1 kb Plus DNA Ladder, ready-to-use	Thermofisher scientific	Cat. No.: FERSM1334
Magnesium sulfate	sigmaaldrich	Cat. No.: M7506-1KG
Calcium chloride (97%)	sigmaaldrich	Cat. No.: 746495-500G
Kanamycin	Thermofisher scientific	Cat. No.: 11815032
Potassium phosphate dibasic anhydrous	Fisherchemicals	Cat. No.: P288-500
Ammonium Chloride (99.5%)	sigmaaldrich	Cat. No.: A9434-1KG
Sodium phosphate dibasic	sigmaaldrich	Cat. No.: S7907-1KG
Glucose (99.5%)	sigmaaldrich	Cat. No.: G8270-1KG

Luria Broth Base	Thermofisher scientific	Cat. No.: 12795084
Iron (II) sulfate heptahydrate	sigmaaldrich	Cat. No.: F8633-250G
FD Kpn1	Thermofisher scientific	Cat. No.: FD0524
FD EcoR1	Thermofisher scientific	Cat. No.: FD0274
FD EcoRV	Thermofisher scientific	Cat. No.: FD0303
FD Dpn1	Thermofisher scientific	Cat. No.: ER1701
Hifi DNA Assembly Master Mix	NEB	Cat. No.: E2621L
T4 DNA Ligase	Thermofisher scientific	Cat. No.: EL0011
Nalidixic Acid	Fisher scientific	Cat. No.: 50488853
Gentamycin Sulfate	sigmaaldrich	Cat. No.: 345814-1GM
Agarose	Alkali	Cat. No.: A7705
RNAprotect Bacteria Reagent	Qiagen	Cat. No.: 76506
Rifampicin	Sigmaaldrich	Cat. No.: 557303-1GM
Pierce™ ECL Western Blotting Substrate	Thermofisher scientific	Cat. No.: 32106
GeneJET Gel Extraction Kit	Thermofisher scientific	Cat. No.: K0692
GeneJET Plasmid Miniprep Kit	Thermofisher scientific	Cat. No.: K0503

GeneJET PCR Purification Kit	Thermofisher scientific	Cat. No.: K0702
Glycine (99%)	sigmaaldrich	Cat. No.: G8898
Methanol (NF), Fisher Chemical™	Fisher scientific	Cat. No.: 67-56-1
PVDF Transfer Membranes, 0.2 µm	Thermofisher scientific	Cat. No.: 88520
DTT	sigmaaldrich	Cat. No.: 3483-12-3
Phenol Solution	sigmaaldrich	Cat. No.: 108-95-2
Glass Beads, acid-washed 106µm (~140 U.S. sieve)	sigmaaldrich	Cat. No.: G4649
Phenol:Chloroform 5:1	sigmaladrich	Cat. No.: P1944
Lithium Chloride Solution	sigmaaldrich	Cat. No.: 7447-41-8
Experimental models: Organisms/strains		
Caulobacter crescentus NA1000	T00841	N/A
Escherichia coli K-12 MG1655	T00007	N/A
Bacillus subtilis DK1042 comIQ12L	T00010[14]	N/A
Oligonucleotides		
HY1F: tgctgatgccgactacaaggatgacgatg	Invitrogen	N/A
HY1R: actttcccatacgcgtaacgttcgaattc	Invitrogen	N/A
HY2F: cgttacgcgtatgggaaagtcttacccaac	Invitrogen	N/A
HY2R: ccttgtagtcggcatcagcaaacccaag	Invitrogen	N/A
HY3F: gtacaagtgagctagctgcagcccggggg	Invitrogen	N/A
HY3R: gcagctgcagcttgtcatcgtcatccttgtagtcggc	Invitrogen	N/A

HY4F: cgatgacaagctgcagctgcctcccctg	Invitrogen	N/A
HY4R: tgcagctagctcacttgtacagctcgtccatgc	Invitrogen	N/A
HY5F: ggtcgatgacgagaattcgaacgttacg	Invitrogen	N/A
HY5R: tcgcgcgtggggtaccatgcatattaattaag	Invitrogen	N/A
HY6F: gcatggtaccccacgcgcgagcgcaaca	Invitrogen	N/A
HY6R: tcgaattctcgtcatcgaccacgaccgacacg	Invitrogen	N/A
CtrA forward: actgatgctgaagtctgaagg	Al-Husini *et al.* Mol Cell. 2020[16]	N/A
CtrA reverse: gattgaggtcgagcaggataag	Al-Husini *et al.* Mol Cell. 2020[16]	N/A
5s forward: ccattccgaactcggttcgttaag	Al-Husini *et al.* Mol Cell. 2020[16]	N/A
5s reverse: tggcggcgacctactct	Al-Husini *et al.* Mol Cell. 2020[16]	N/A
8688: aggaggtcgaccatagcaagaggaggtgaaagtatg	IDT	N/A
8689: ctcctgctagcgccagaaccagcagcggagccagcggattttg catactctacggctcgag	IDT	N/A
8692: aggaggctagcatgggaaagtcttacccaactgtg	IDT	N/A
8693: ctcctgcatgcttacttgtcatcgtcatccttgtag	IDT	N/A
F-rne-mch: aacatcaagttggacatcacc	Sigma-Aldrich	N/A
R-rne: ctcttccagctcagc	Sigma-Aldrich	N/A
Recombinant DNA		
pFlgC-2	Thanbichler *et al.* NAR 2011[17]	N/A
pRNE-YFPC-1	Al-Husini *et al.* Mol Cell 2018[13]	N/A
Apex2FlgC-2	This paper	N/A

RNE-Apex2FlgC-2	This paper	N/A
RNE-Apex2Flg-EGFPC-2	This paper	N/A
RNEΔCTD-Apex2FlgC-2	This paper	N/A
pcDNA5/FRT/TO APEX2-GFP	Addgene	Cat. No.: 129640
pDP647 (amyE::Phypsank-rny-apex2 spec amp)	This paper	N/A
pDR111 (amyE::Physpank spec amp)	Gift from David Rudner	N/A

| rne-Apex2-KmR (gctgctgctgactacaaggatgacgacgataagGGAAAGTCTTACCCAACTGTGAGTGCTGATTACCAGGACGCCGTTGAGAAGGCGAAGAAGAAGCTCAGAGGCTTCATCGCTGAGAAGAGATGCGCTCCTCTAATGCTCCGTTTGGCATTCCACTCTGCTGGAACCTTTGACAAGGGCACGAAGACCGGTGGACCCTTCGGAACCATCAAGCACCCTGCCGAACTGGCTCACAGCGCTAACAACGGTCTTGACATCGCTGTTAGGCTTTTGGAGCCACTCAAGGCGGAGTTCCCTATTTTGAGCTACGCCGATTTCTACCAGTTGGCTGGCGTTGTTGCCGTTGAGGTCACGGGTGGACCTAAGGTTCCATTCCACCCTGGAAGAGAGGACAAGCCTGAGCCACCACCAGAGGGTCGCTTGCCCGATCCCACTAAGGGTTCTGACCATTTGAGAGATGTGTTTGGCAAAGCTATGGGGCTTACTGACCAAGATATCGTTGCTCTATCTGGGGGTCACACTATTGGAGCTGCACACAAGGAGCGTTCTGGATTTGAGGGTCCCTGGACCTCTAATCCTCTTATTTTCGACAACTCATACTTCACGGAGTTGTTGAGTGGTGAGAAGGAAGGTCTCCTTCAGCTACCTTCTGACAAGGCTCTTTTGTCTGACCCTGTATTCCGCCCTCTCGTTGACAAATATGCAGCGGACGAAGATGCCTTCTTTGCTGATTACGCTGAGGCTCACCAAAAGCTTTCCGAGCTTGGGTTTGCTGATGCCtaaattacacgtcttgagcgattgtgtaggctggagctgcttcgaagttcctatactttctagagaataggaacttcggaataggaacttcaagatcccctcacgctgccgcaagcactcagggcgcaagggctgctaaaggaagcggaacacgtagaaagccagtccgcagaaacggtgctgacccccggatgaatgtcagctactgggctatctggacaagggaaaacgcaagcgcaaagagaaagcaggtagcttgcagtgggcttacatggcgatagctagactgggcggttttatggacagcaagcgaaccggaattgccagctggggcgccctctggtaaggttgggaagccctgcaaagtaaactggatggctttcttgccgccaaggatctgatggcgcaggggatcaagatctgatcaagagacaggatgaggatcgtttcgcatgattgaacaagatggattgcacgcaggttctccggccgcttgggtggagaggctattcggctatgactgggcacaacagacaatcggctgctctgatgccgccgtgttccggctgtcagcgcaggggcgcccggttcttttttgtcaagaccgacctgtccggtgccctgaatgaactgcaggacgaggcagcgcggctatcgtggctggccacgacgggcgttccttgcgcagctgtgctcgacgttgtcactgaagcgggaagggactggctgctattgggcgaagtgccggggcaggatctcctgtcatctcaccttgctcctg | Twist Bioscience | N/A |

Software and algorithms		
ImageJ	Imagej	https://imagej.net/software/fiji/downloads
Ibright analysis software	Thermofisher scientific	https://www.thermofisher.com/us/en/home/technical-resources/software-downloads/ibright-western-imager.html

In vivo RNA proximity labeling

<u>APEX2, Alkyne-Phenol, and H_2O_2 are needed for RNA proximity labeling</u>

<u>Gram negative bacteria</u>

Gram negative bacteria were grown overnight in liquid media at their optimal growth
temperature. Once the cultures reached an optical density (OD) of ~0.3-0.6, 2.5 mM of
alkyne-phenol was added and incubated for 30 minutes in a shaker incubator at 200 rpm. A
100 mM H_2O_2 solution (994 µL H_2O + 6 µL H_2O_2, 50%) was then freshly prepared. 1 mM of
H_2O_2 was then added for 45 seconds as the cultures were still shaking. The labeling
experiment was carried out again with specific omissions as controls: no alkyne-phenol
added, no H_2O_2 added, no alkyne-phenol and H_2O_2 added, different concentrations of
alkyne-phenol added (0.5 mM and 1 mM), and different H_2O_2 exposures (15 seconds, 30
seconds, and 60 seconds). 4 mL of cell culture were then quickly transferred into two 2 mL
Eppendorf tubes and centrifuged at 14,000 rpm for 10-15 seconds. The supernatant was
quickly discarded, and the cell pellets were resuspended in 1 mL of 65°C pre-warmed
TRIzol. Because proximity labeling requires the folded APEX2 protein, its denaturation in
TRIzol can be used to quench the proximity labeling reactions. RNA extraction followed by
ethanol precipitation was then carried out. 55 µL of a 10 mM Tris-HCl pH 7.0 (no EDTA)
solution was added to dissolve the air-dried RNA pellets. The samples were vortexed
briefly to aid resuspension. 5 µL were taken from the samples and placed in new tubes for
Agilent Tapestation analyses and cell lysate RNA sequencing.

Gram positive bacteria

Gram positive bacteria were grown overnight in liquid media at their optimal growth
temperature. Once the cultures reached an optical density (OD) of ~0.3-0.6, the cultures
were re-inoculated into fresh media at an OD of ~0.05 supplemented with 1 mM isopropyl
β-D-thiogalactopyranoside (IPTG) to induce the expression of RNY-APEX2. Once the
cultures reached an OD of ~0.4-0.5, 2.5 mM of alkyne-phenol was added and incubated for
30 minutes in a shaker incubator at 200 rpm. A 100 mM H_2O_2 solution was then freshly
prepared. 1 mM of H_2O_2 was then added for 45 seconds as the cultures were still shaking.
An equal amount of cold methanol (pre-stored at -80 oC) was then added to the culture
solution. The mixture was then quickly vortexed for 5 seconds and centrifuged at 14,000
rpm for 10-15 seconds. The cell pellets were resuspended in LETS buffer (1M Tris-HCl, 1M
LiCl, 0.5M EDTA, and 10% SDS; for a 4 mL of gram positive cultures, the cell pellets were
dissolved in 550 uL of LETS buffer). Both the LETS buffer and bead:phenol solutions were
pre-warmed at 75 oC before being used. The solution was then transferred to a tube
containing glass beads + phenol (for a 4 mL of gram positive culture, 320 uL of glass beads
+ 380 uL of phenol were used). The mixture was vortexed for 3 minutes. Chloroform was
then added to the mixture and the solution was vortexed for 30 seconds (for a 4 mL of gram
positive culture, 400 uL of chloroform was added to the cells + beads solution). The
samples were then centrifuged at 3200 x g for 10 minutes. The top aqueous layer was then
transferred to a fresh tube containing a phenol:chloroform solution (for a 4 mL of gram
positive culture, 550 uL of phenol:chloroform was used). The phenol:chloroform solution
was pre-warmed at 75 oC before being used. The mixture was then vortexed for 3 minutes,
and the sample was centrifuged at 3200 x g for 10 minutes. The top aqueous layer was then
transferred to a fresh tube and RNA precipitation was carried out. 55 µL of a 10 mM Tris-
HCl pH 7.0 (no EDTA) solution was added to dissolve the air-dried RNA pellets. The
samples were vortexed briefly to aid resuspension.

Copper-catalyzed click chemistry reactions

Copper-catalyzed click chemistry reactions were then conducted on the labeled RNA. The
reactions have a 300 µL total volume and contained: 50 µL of 100 ng/uL RNA (5 µg) + 175 µL
H_2O + 12 µL of 125 mM sodium ascorbate + 3 µL of 10 mM Cy5-Azide + 60 µL of a mix of 2.5
mM Cu(II)SO_4/12.5 mM THPTA (tris-hydroxypropyltriazolylmethylamine). The reagents were
added in the order they are listed. The click reaction was vortexed for 5 seconds and
incubated for 10 minutes at room temperature away from light exposure. Afterward, RNA
precipitation was conducted and the RNA pellets were resuspended in 50 µL of 10 mM
Tris-HCl pH 7.0 and 0.1 mM EDTA. The samples were then subjected to dot blot filtration
on a nylon membrane following the manual's instructions of BIO RAD's Bio-Dot® and Bio-

Dot SF Microfiltration Apparatus. A nylon membrane was pre-wetted in a 6X SSC solution
for at least 10 minutes. The pre-wetted nylon membrane was then assembled into the dot
blot apparatus. Vacuuming was applied to the apparatus and tighter sealing was
implemented. The wells were then washed 3 times with 1 mL of an ice-cold mixture of 10
mM NaOH/1 mM EDTA. Afterward, the 5 µg RNA samples were directly applied to the wells
(no RNA dissolution was performed). The wells were washed 3 times with 1 mL of an ice-
cold mixture of 10 mM NaOH/1 mM EDTA. The blotted membrane was washed 3 times for
5 minutes with a 30 mL solution containing 2X SSC and 0.1% SDS. Fluorescent blot
imaging was then visualized using an ibright imaging system at a fluorescent light exposure
of 50 ms under the Cy5 channel.

RNase A & DNase I treatments

5 µg RNA (50 µL of 100 ng/µL of RNA) collected from a proximity labeling reaction (4 mL of
$Js767$ cells grown in M2G media with 2.5 mM alkyne phenol, 45 seconds H_2O_2) were
subjected to either an RNase A (5 µL of 10 mg/mL RNase A) or DNase I (5 µL of 1 U/µL
DNase I) treatment. The treated and mock samples were incubated for 2 hours at 37°C.
Following the enzyme or mock treatments, the RNA was precipitated and resuspended in
50 µL of 10 mM Tris-HCl pH 7.0 and then subjected to the Copper catalyzed click-
chemistry reaction with Cy5-azide. Following the 10-minute click reaction incubation, the
RNA was then precipitated and subjected to dot blot filtration as previously described.

Streptavidin capture of biotin labeled RNA

To enrich the transcriptome of BR-bodies, RNA labeled with alkyne-phenol was subjected
to the following click chemistry reaction: 50 µL of RNA (total RNA volume) + 175 µL H2O +
12 µL of 125 mM sodium ascorbate + 3 µL of 10 mM Biotin-peg3-azide + 60 µL of a mix of
2.5 mM Cu(II)SO$_4$/12.5 mM THPTA. The reagents were added in order as they are listed. The
click reaction was vortexed briefly and incubated for 10 minutes at room temperature away
from light exposure. RNA precipitation was then conducted and the air-dried RNA pellets
were resuspended in 100 µL B&W buffer (5 mM Tris-HCl, pH 7.5, 1 M NaCl, 0.5 mM EDTA,
0.1% v/v tween-20). The RNA samples were then immediately used for biotinylated RNA
enrichment.

High-capacity magnetic streptavidin beads (Vectorslab) were washed three times with 1
mL B&W buffer, twice with 1 mL of a mix of 0.1 M NaOH/0.05 M NaCl, and once with 1 mL
0.1 M NaCl at room temperature. The beads were then blocked using a blocking buffer (1
mg/mL BSA + 1 mg/ml heparin salt, dissolved in B&W buffer) at room temperature for 2
hours. The beads were then washed three times with 1 mL of B&W buffer under gently
vortexing. Afterward, the 100 µL RNA solution was added to the beads along with an extra

100 µL of B&W buffer. The RNA and beads were allowed to mix at room temperature for
one hour. The RNA-loaded beads were washed three times with 1 mL of B&W buffer, twice
with 1 mL of a PBS solution containing 4 M urea + 0.1% SDS, and twice with 1 mL of just
PBS at room temperature. To elute the enriched RNA, 900 uL of 65°C pre-warmed TRIzol
was added to the beads and the samples were incubated at 65°C for 10 minutes.
Afterward, 200 µL of chloroform was added and the samples were inverted 6-7 times and
then were incubated for 5 minutes at room temperature. The samples were centrifuged at
14,000 rpm for 10 minutes at 4°C. The upper aqueous layer was transferred to a new
Eppendorf tube and the enriched RNA was precipitated as previously described. The air-
dried RNA pellets were resuspended in 10 µL of 10 mM Tris-HCl pH 7.0 and 0.1 mM EDTA. 4
µL were then used for Agilent Tapestation analysis. The remaining volume could be utilized
for the follow-up RNA-seq experiments.

<u>RNA quality assessment</u>

Js767 (RNE-Apex2FlgC-2) was grown in PYE at 28°C overnight and re-inoculated into 5 mL
M2G the next day. Once the cultures reached an OD of 1.2, 2.5 mM of alkyne-phenol was
added and incubated for 30 minutes in a shaker incubator at 28°C at 200 rpm. 1 mM of
H_2O_2 was added to the culture for 45 seconds. Total RNA was then collected and
precipitated from 4 mL of culture as previously described using TRIzol. Afterward, 50 µL of
100 ng/µL of alkylated RNA were subjected to the following copper catalyzed click-
chemistry reaction: 50 µL of 100 ng/µL RNA (5 µg) + 175 µL H_2O + 12 µL of 125 mM sodium
ascorbate + 3 µL of 10 mM Cy5-Azide + 15 µL of 100 mM aminoguanidine hydrochloric acid
+ 60 µL of a mix of 2.5 mM Cu(II)SO_4/12.5 mM THPTA. The click reaction was incubated for
10 minutes at room temperature away from light exposure. The RNA from the click reaction
was then precipitated and the air-dried RNA pellet was resuspended in 50 µL of 10 mM
Tris-HCl pH 7.0 and 0.1 mM EDTA. 250 ng of RNA from the *Js767* RNA lysate and the click
reaction RNA were sent for Agilent Tapestation analysis.

RNA precipitation

RNA precipitation was carried out by adding, in order, 2 µL of 1 mg/mL glycogen, 1/10
sample volume of 3 M sodium acetate pH 5.5, and 1 sample volume of isopropanol. The
sample mixtures were vigorously vortexed and stored at -80°C overnight. The samples were
spun for 1 hour at 14,000 rpm. The supernatant was then decanted and 1 mL of ice-cold
80% ethanol was added. The samples were spun for 15 minutes at 14,000 rpm. The
ethanol was pipetted out and the samples were then spun for 1 minute at 14,000 rpm. Any
remaining ethanol was pipetted out and the pellets were allowed to air-dry until they
became translucent. The RNA was then resuspended in the buffer of choice.

Western Blot

The following buffers were initially prepared:

1) 10X transfering buffer: 144 grams of glycine and 30.2 grams of Tris-Base were
195
196 dissolved in 900 mL of ddH2O. The buffer volume was adjusted to 1 L of ddH2O and
197 then filtered utilizing a Nalgene Rapid-flow sterile disposable filter unit.
2) 10X TBS: 24 grams of Tris-Base and 88 grams of NaCl were dissolved in 900 mL of
198
199 ddH2O. The pH of the buffer was adjusted to 7.6 utilizing hydrochloric acid. The
200 buffer volume was adjusted to 1 L of ddH2O and then sterilized utilizing a Nalgene
201 Rapid-flow sterile disposable filter unit.
3) 1X TBST: 10X TBS was diluted 10-fold with ddH2O to make 1X TBS. 0.1% Tween-20
202
was then added to the 1X TBS solution to make 1X TBST.

NA1000, LS4379 (hfq-m2), and Js767(RNE-Apex2FlgC-2) were inoculated in PYE and
incubated overnight at 28°C until they reached log phase (OD ~0.3-0.6). 1 mL was taken
from each culture, placed in 1.5 mL Eppendorf tubes, and centrifuged for 2 minutes at
6,000 rpm. The liquid cultures were discarded, and the cells were resuspended in 4X SDS
loading dye. 125 µL of 4X SDS loading dye was added to cultures with an OD~0.5. The
samples were boiled at 95 °C for 5 minutes then vortexed vigorously. The vortexed samples
were quickly spun down and placed on ice. 5 µL of a pre-stained PageRuler marker and 20
µL of lysate from each sample were loaded onto an SDS-PAGE gel. 1X transferring buffer
was freshly prepared (20 mL of 10X transfer buffer + 20 mL of methanol in 160 mL ddH2O)
while the samples were running on the gel. A PVDF transferring membrane was wetted in
methanol and 6 blotting papers were wetted using 1X transferring buffer for at least 10
minutes. Once the lysate samples on the SDS-PAGE were properly resolved, the gel was
placed on top of 3 pre-wetted blotting papers, and 2 mL of 1X transferring buffer was
poured on the gel. The PVDF membrane was briefly plunged into the 1X transferring buffer
and then placed on top of the gel. The remaining 3 blotting papers were placed on top of
the PVDF membrane. The transferring of the lysates from the SDS-PAGE gel to the PVDF
membrane occurred by utilizing the BIO-RAD Trans-Blot Turbo Transfer system (1 amps,
2.5 volts, 15 minutes). Once the transfer was completed, the PVDF membrane was placed
in a new container containing 5% BSA (blocking solution). The container was nutated for 1
hour at room temperature. Afterward, the 5% BSA blocking solution was discarded and the
PVDF membrane was washed 5 times with ddH2O for 5 minutes per wash. The primary
antibody (DYKDDDDK anti-flag tag) was then diluted (1:10000) into a fresh 5% BSA
solution. Once the last ddH2O wash was completed, the primary antibody solution was
poured on the PVDF membrane, and the container was nutated at 4°C overnight. The
following day, the primary antibody solution was discarded and the PVDF membrane was

washed 5 times with TBST for 5 minutes per wash. The secondary antibody (goat anti-rabbit
secondary antibody, HRP conjugated) was then diluted (1:10000) into a fresh 5% BSA
solution. Once the last TBST wash was completed, the secondary antibody solution was
poured on the PVDF membrane, and the container was nutated at room temperature for 1
hour avoiding light exposure. The secondary antibody solution was then discarded and the
PVDF membrane was washed 5 times with TBST for 5 minutes per wash. Following the final
washing procedure, a pierce ECL western blotting substrate solution was prepared (1 mL
of reagent 1 + 1 mL of reagent 2 were mixed and vigorously vortexed). The PVDF membrane
was placed on Seram wrap and the 2 mL substrate solution was poured on top of the
membrane. The substrate was incubated on the PVDF membrane at room temperature for
5 minutes avoiding light exposure. The membrane was then placed into an iBright imaging
system and the signal was detected under the Chemi-blot filter.

Dilution Plates

NA1000 & Js767 (RNE-Apex2FlgC-2) were grown in PYE overnight at 28°C. The overnight
cultures were diluted into fresh media and incubated at 28°C until the cells reached an OD
of ~0.3-0.6. Cells were then diluted to an OD=0.05 in PYE and 4 10-fold serial dilutions
were made. 5 μL were spotted from each dilution onto PYE plates. The plates were
incubated at 28°C for two days, and images were taken using an ibright imaging system.

Fluorescent Cell Imaging

Js87 (RNE-msfGFP) & Js768 (RNE-Apex2FlgC-EGFPC-2) were grown in PYE + gent (0.5
μg/mL) and PYE + Kan, respectively, overnight at 28°C. Cells were then fixed on M2G +
1.5% agarose pads placed on microscope slides (3051, Thermofisher scientific). Imaging
was performed on an epifluorescence microscope with a 100X objective. Specifically,
Nikon elements software was used to control a Nikon Eclipse NI-E equipped with a
CoolSNAP MYO-CCD camera and a 100x Oil CFI Plan Fluor (Nikon) objective to capture the
images.
255
mRNA half-lives measurement

NA1000 and *Js767 (RNE-Apex2FlgC-2)* were grown in liquid PYE at 28°C overnight. Cells
were then serially re-inoculated into liquid M2G and incubated until overnight log phase
(OD ~0.3-0.6) was reached. Log phase cultures were then re-inoculated to an OD = 0.05 in
25 mL liquid M2G. Before adding rifampicin, at time point 0, 1 mL of cells were added to 2
mL of RNAprotect Bacterial reagent (Qiagen) and vortexed for 5 seconds. 200 μg/mL of
Rifampicin was administered to the cultures and 1 mL of cells were extracted and added to
2 mL of RNAprotect Bacterial reagent at each of the following time points (followed by 5
seconds vortexing): 1, 2, 4, and 8 minutes. Cells were incubated at room temperature in
the RNAprotect Bacterial reagent for 5 minutes before being spun at 5000 rpm for 5

minutes. The bacterial pellets were resuspended in 1 mL of 65°C pre-heated TRizol
(Ambion) and incubated at 65°C for 10 minutes. 200 µL of chloroform was then added and
the samples were incubated at room temperature for 5 minutes. Subsequently, the
samples were centrifuged at 14,000 rpm for 10 minutes at 4°C. The aqueous layer was
removed and placed in a new 1.5 mL Eppendorf tube. The RNA was then precipitated and
the pellets were resuspended in 50 µL elution buffer (10 mM Tris-HCl, pH=7.0, 0.1 mM
EDTA). PCR tubes were filled with a master mix that contained 0.4 µM of *ctrA* forward
primer and 0.4 µM of *ctrA* reverse primer, 1X Luna Universal One-Step Reaction Mix, 1X
Luna WarmStart RT Enzyme Mix, and water. 100 ng/µL of RNA template was additionally
aliquoted into each of the PCR tubes. The samples were mixed well and quickly spun
down. A QuantStudio Real-Time PCR apparatus was utilized to conduct and examine the
qRT-PCR experiments. The same qRT-PCR experiment was done using instead the 5S
forward and reverse primers. To determine the mRNA-decay rates, we fitted a linear curve
to the ln (fraction RNA remaining) at each time point. Using a standard curve, the Ct was
converted into the quantity of RNA. Each time point's 5S rRNA amount was divided by the
amount of 5S rRNA at time point 0. The number obtained at each time point was then used
to normalize the amount of *ctrA* at each time point. The natural log of % RNA remaining
found in each sample was divided by the natural log of RNA at time point 0. The slopes of
the linear curve fit were then converted into mRNA half-life using the following equation:
mRNA half-life=$-\ln(2)$/slope.

RNA-seq

Duplicate biological samples were prepared for RNA-seq of the Full-length RNase E-APEX2
fusion (JS767) and the RNase E-NTD-APEX2 (JS801). Lysates from each sample, as well as
the eluted RNA from affinity purification were prepared for RNA-seq. RNA-seq was
performed by IU-Bloomington CGB genomics service facility using the Illumina Tru-seq
stranded HT kit. Raw sequencing reads were deposited to the NCBI GEO database with
accession number GSE297938. Data analysis was performed similar to Al-Husini *et al.* Mol
Cell 2020[16]. RNA reads were stripped of adapters, and tRNA and rRNA reads were aligned
to unique file with duplicate copies deleted using bowtie[18], and reads not aligning to tRNA
and rRNA were aligned to the *C. crescentus* transcriptional units[19] using bowtie. Genes
with fewer than 50 reads mapping to them were omitted from analysis, and the RPKM
values were calculated for each sample. The log2 ratio between the RPKM values in the
elution and lysate samples were then calculated on all genes with >=50 reads in both
elution and lysate samples. Samples with log2(elution/lysate) measurements in all four
biological samples were used for comparison for BR-body proximity labeling enrichment.
To calculate which RNA types were enriched in BR-bodies, the distribution of enrichment
measurements log2(elution RPKM/lysate RPKM) were subjected to a T-test with unequal

variance between the BR-body + (JS767) and BR-body − (JS801) strains and the resulting p-
values were reported. Raw read count and processed proximity labeling RNA-seq data can
be found in Table S3.

Plasmid Construction

pAPEX2-FlgC-2 *Kan^R*

The pFlgC-2 vector[17] was PCR amplified using primers HY1F & HY1R. APEX2 was PCR
amplified using primers HY2F & HY2R from the addgene template #129640. The amplicons
were run on a 1% agarose gel and gel extracted using a GeneJET Gel Extraction Kit. The
purified vector was Dpn1 treated then column purified using the GeneJET PCR Purification
Kit. The Apex2FlgC-2 *Kan^R* plasmid was assembled via Gibson assembly (NEB) and
transformed into chemically competent DHbeta10 *E. coli* and plated on LB + Kan (50
µg/mL) plates. The resulting KanR colonies were minipreped using the GeneJET Plasmid
Miniprep Kit and screened via restriction digestion (EcoR1) and the insert sequence was
verified by Sanger sequencing (Genewiz).

pRNE-Apex2-FlgC-2 *Kan^R*

The last 534 RNase E (RNE) base pairs were obtained by digesting pRNE-YFPC-1[17] with
Nde1 and Kpn1. APEX2FlgC-2 was digested with Nde1 and Kpn1. The digestion reactions
were run on a 1% agarose gel and the RNE fragment and Apex2-FlgC-2 vector were gel
extracted using a GeneJET Gel Extraction Kit. The RNE fragment and Apex2FlgC-2 vector
were ligated using T4 ligase, transformed into chemically competent DHbeta10 *E. coli*, and
selected on LB + Kan plates. The resulting KanR colonies were minipreped using the
GeneJET Plasmid Miniprep Kit and screened via restriction digestion (EcoRV) and the insert
sequence was verified by Sanger sequencing (Genewiz).

pRNE-Apex2FlgC-EGFPC-2 *Kan^R*

The RNE-Apex2FlgC-2 vector was PCR amplified using primers HY3F & HY3R. The EGFP
insert was PCR amplified using primers HY4F & HY4R from the addgene template #129640.
The amplicons were run on a 1% agarose gel and gel extracted using a GeneJET Gel
Extraction Kit. The purified vector was Dpn1 treated then column purified using the
GeneJET PCR Purification Kit. The RNE-Apex2FlgC-EGFPC-2 plasmid was assembled via
Gibson assembly (NEB) and transformed into chemically competent DHbeta10 E. coli and
selected on LB + Kan plates. The resulting KanR colonies were minipreped using the
GeneJET Plasmid Miniprep Kit and screened via restriction digestion (EcoRV) and the insert
sequence was verified by Sanger sequencing (Genewiz).

pRNEΔCTD-Apex2FlgC-2 Kan[R]

The pApex2-FlgC-2 vector[17] was PCR amplified using primers HY5F & HY5R. RNE-NTD was
PCR amplified using primers HY6F & HY6R from NA1000 cells. The amplicons were run on
a 1% agarose gel and gel extracted using a GeneJET Gel Extraction Kit. The purified vector
was Dpn1 treated then column purified using the GeneJET PCR Purification Kit. The

pRNE(NTD)-Apex2FlgC-2 *Kan[R]* plasmid was assembled via Gibson assembly (NEB) and
transformed into chemically competent DHbeta10 *E. coli* and plated on LB + Kan (50
µg/mL) plates. The resulting KanR colonies were minipreped using the GeneJET Plasmid
Miniprep Kit and screened via restriction digestion (Kpn1) and the insert sequence was
verified by Sanger sequencing (Genewiz).

pDP647 amyE::Phypsank-rny-apex2 spec[R] amp[R]

To generate the *B. subtilis* inducible Rny-APEX2 construct pDP647, the *rny* gene was
amplified from wild type DK1042 DNA with primers 8688/8689 and *apex2* was amplified
from plasmid pAPEXC-2 with primers 8692/8693. Next, the *rny* amplicon was digested
with SalI and NheI, the *apex2* amplicon was digested with NheI and SphI and the two
fragments were simultaneously ligated into the SalI and SphI sites of pDR111 that carries a
polylinker downstream of the IPTG-inducible $P_{hyspank}$ promoter, the gene encoding the LacI
repressor, and a spectinomycin resistance cassette between the arms of the amyE gene
(generous gift of David Rudner, Harvard Medical School).

Strain construction

Strain construction

Js767: NA1000 rne::rne-apex2-flg Kan[R]

The RNE-Apex2FlgC2 plasmid was recombined into the *rne* locus in NA1000 via mating and
the selection was carried out on PYE + Nal (20 µg/mL) + Kan (25 µg/ mL) plates. The
resulting KanR colonies were first grown in PYE + Kan (5 µg/mL) cultures and then screened
by PCR.

Js801: NA1000 rne::rneΔCTD-apex2-flg Kan[R]

The RNEΔCTD-Apex2FlgC2 plasmid was recombined into the *rne* locus in NA1000 via
mating and the selection was carried out on PYE + Nal (20 µg/mL) + Kan (25 µg/ mL) plates.
The resulting KanR colonies were first grown in PYE + Kan (5 µg/mL) cultures and then
screened by PCR.

Js768: NA1000 rne::rne-apex2-flg-egfp Kan[R]

The RNE-Apex2FlgC-EGFPC-2 plasmid was recombined into the *rne* locus in NA1000 via
mating and the selection was carried out on PYE + Kan plates. The resulting KanR colonies
were first grown in PYE + Kan cultures and then screened by PCR.

DB2579 amyE::rny-APEX2 spec[R] amp[R]

The pDP647 plasmid was transformed into DK1042[14] cells, and selected on plates
containing LB and 100 µg/mL Spectinomycin. The resulting colonies were then screened
for the integration by PCR.

ES413: rne::rne-mcherry-flg-Apex2 kan[R]

The *in vitro* synthesized sequence rne-Apex2-KmR was PCR amplified using F-rne-mch and
R-rne primers and recombined into the *rne* locus in Kti162[15] by lambda red system[20], the
selection was carried out on LB + Kan (30 µg/ mL) plates. The *rne-mcherry-flag-Apex2 kan[R]*
was then moved to MG1655 by P1 transduction[21]. The resulting Kan[R] colonies were grown
in LB + Kan, screened by PCR and verified by Sanger sequencing (The Center for Genomic
Technologies – Huji).

Results:

RNaseE-APEX2 proximity labeling of RNA requires Alkyne-Phenol, H_2O_2, and APEX2

APEX2 is an engineered ascorbate peroxidase which can catalyze the creation of a
free radical on the oxygen of alkyne-phenol, and this highly reactive species can react with
molecules within a 10-20 nm radius[22,23] (Fig 1A). It was shown that APEX2 can label RNA in
eukaryotic cells, making this a useful experimental system for identifying localized
RNAs[10,12,24,25]. In order to perform RNA proximity labeling in bacteria, we first generated a
gene fusion between RNase E and APEX2. RNase E is known to phase-separate into
bacterial ribonucleoprotein bodies (BR-bodies), biomolecular condensates containing
mRNA and promoting the mRNA decay process[13,16]. To determine whether APEX2 fusions
are tolerated in bacteria, we examined the subcellular localization, cell fitness, and mRNA
decay activity of RNase E-APEX2 fusions in the bacterium *Caulobacter crescentus* (Fig 1B).
We observed that fusing APEX2 to RNase E led to proper expression of RNase E-APEX2 (Fig
S1) and did not alter its ability to phase separate into BR-bodies (Fig 1B). In addition, since
RNase E's ability to degrade mRNAs is essential for cell growth[16,26], we also examined the
cellular fitness of the RNase E-APEX2 fusion and found that both CFUs and colony size
were indistinguishable from wild-type (Fig 1B). Finally, we compared the mRNA decay
activity of the RNase E-APEX2 fusion and found that it has similar rates of mRNA decay as
compared to wild-type (Fig 1B), while 5S rRNA remained stable. Altogether, we could not

detect any measurable differences in RNase E function when fused to APEX2. To examine whether the APEX2 fusion can be used to label RNA we added all combinations of proximity labeling reactants (H_2O_2 and Alkyne-Phenol) in combination with or without the RNase E-APEX2 fusion (Fig 1C). Each of the proximity labeling conditions were performed, and then we extracted the RNA and used copper click chemistry to conjugate a Cy5-azide to the RNA. The RNA was then deposited on a positively charged nylon membrane using a dot blot apparatus, and the Cy5 signal was measured in a fluorescent gel imager. We find that only in the presence of APEX2, H_2O_2, and alkyne-phenol do we observe RNA labeling (Fig 1C), suggesting that APEX2 proximity labeling reactions occurred. To determine whether the Cy5 signal was a result of RNA, or contaminating DNA we performed RNase A or DNase I digestions on our samples before spotting them on the dot blot (Fig 1C). Here we see that RNase A treatment leads to a complete loss of Cy5-fluorescence, while DNase I treatment led to no difference in fluorescence signal, suggesting that our Cy5 fluorescence is from RNA.

Rapid labeling of cellular RNA with Alkyne-Phenol

One of the major challenges of studying bacterial mRNA decay is the very short lifetimes of cellular mRNAs. Most bacterial mRNAs in rapidly growing species have half-lives between 1-4 minutes[27,28], making it hard to harvest the RNA before they are degraded. To address this technical limitation, we first optimized the Alkyne-phenol concentration used for labeling cells and found that 2.5 mM Alkyne-Phenol robustly labels cellular RNA (Fig 2). Next, we performed a time-course of H_2O_2 exposure to identify the timescale of RNA labeling, ranging from 15 seconds reaction times to 1 minute reaction times. We found that robust labeling could be achieved in as little as 15 seconds, while labeling increased upon longer proximity labeling reaction times (Fig 2). Importantly, the labeling reaction occurs on the sub-minutes timescale, making this method well-suited to the use in exponentially growing bacterial cells.

RNA proximity labeling works across bacteria

To increase the applicability of this method, we sought to determine whether APEX2 can label RNA in other species of bacteria. As a key requirement of APEX2 activity is heme, we chose to fuse APEX2 to RNA degradosome scaffold proteins in *E. coli* (gram -) and *B. subtilis* (gram +) as both organisms have heme biosynthesis pathways encoded in their genomes. In *E. coli*, we fused APEX2 to RNase E, which scaffolds BR-bodies in this species[15,29], localizing into membrane anchored RNP foci[15]. In *B. subtilis*, we fused APEX2 to RNase Y, which scaffolds BR-bodies in this species and also localizes into RNP foci on the inner membrane of the cell[29,30]. In both of these species, we observed RNA labeling after 45 seconds of H_2O_2 incubation that required the APEX2 fusion (Fig 3), suggesting that

438 APEX2 proximity labeling of RNA can be used in diverse species of bacteria which encode
439 heme biosynthesis pathways.

Copper click-chemistry of Azide-Biotin and streptavidin purification of labeled RNA.

APEX2 proximity labeling of RNA is rather flexible due to the diversity of azides
commercially available for copper catalyzed click chemistry. To isolate cellular RNA for
RNA-sequencing, we altered the azide from azide-Cy5 dye to azide-biotin to allow for
streptavidin-mediated RNA purification from the cell (Fig 4A). In this approach, labeled
RNA is extracted from cells and conjugated to azide-biotin using copper catalyzed click
chemistry, which allows for the subsequent purification of the labeled RNA using
streptavidin-resin. While copper can cleave RNA, using a short incubation time for the
click reaction minimizes RNA cleavage by copper (Fig S2). After click chemistry, the RNA
can be purified under stringent conditions and eluted from the resin under denaturing
conditions. When applied to the RNase E-APEX2 fusion presented earlier, we found that
the eluted RNA profile indeed matched that of BR-body RNA isolated via density
centrifugation (Fig 4B)[16,31]. To compare the specificity of APEX RNA proximity labeling, we
prepared eluted RNAs from BR-body + cells (JS767) and BR-Body – cells (RNaseEΔCTD-
APEX2, JS801) and prepared them for RNA-seq. As a control for the initial amount of each
RNA in the cell, we also performed RNA-seq on the total RNA lysates of each strain. The
overall level of proximity labeling was calculated as the log2 enrichment (eluted/lysate) of
each RNA measured at >50 reads/sample. Overall, we found that BR-body + samples
(JS767) contained higher overall enrichment levels than BR-body – (JS801) (Fig S3A) whose
mean enrichment was near 0, with high reproducibility between biological replicates (Fig
S3B). Prior density centrifugation of BR-bodies comparing BR-body + and BR-body –
strains found they were enriched in mRNAs, small non-coding RNAs, and antisense RNAs
while depleted of tRNAs and rRNAs[16]. Our proximity labeled RNA samples showed mRNAs
and small non-coding RNAs were both enriched, while antisense appeared to be enriched,
however, the p-value was below a 95% confidence interval (p=0.1), while tRNAs and rRNA
were not enriched (Fig 4C). Interestingly, while 16S and 23S rRNA were not enriched, 5S
rRNA was enriched, in-line with RNase E's known role of 5S rRNA processing[32,33] and the
colocalization of BR-bodies adjacent to rRNA loci *in vivo*[34]. Importantly though, while
centrifugation-based isolation of BR-bodies required 200 mL of cell culture and 5 days of
hands-on work to isolate the BR-body enriched RNAs[31], APEX2 proximity labeling required
only 4 mL of cell culture and 3 days of hands-on work, making it higher throughput.

Discussion:

RNP complexes are important regulators of RNA biology, including transcription, RNA processing, transport, translation, and decay[4,35–37]. In addition to RNP complex formation, these complexes have been increasingly found to localize into biomolecular condensate structures, which help to facilitate the spatial organization of the stages in the mRNA life cycle via phase-separation[4,13,16,38–41]. As realizations that RNA localization has become increasingly important in bacteria, methods to identify the population of localized RNAs have been developed, yet these methods have been limited due to the long timescale of the procedures compared to the short mRNA lifetimes in bacteria. APEX2 proximity labeling can rapidly label RNA and can be easily genetically fused to genes whose proteins have known patterns of subcellular localization. APEX2 proximity labeling of RNA requires small amounts of cells, can be completed in a few hours by a single scientist, and yields sufficient RNA for downstream analysis. The reactivity radius of APEX2 is estimated to be 10-20nm[42], suggesting it gives high spatial precision. Genetic fusion of APEX2 is easy to generate, APEX2 RNA labeling works across heme-producing bacteria (*C. crescentus*, *B. subtilis*, and *E. coli*), and APEX2 is similar in size to a fluorescent protein and has no observable functional impacts when fused to RNase E, so we anticipate that this is unlikely to disrupt target protein function. The flexibility of click chemistry allows the conjugation of a wide array of clickable fluorophores or clickable affinity substrates as biotin or azide-resin. Altogether, APEX2 proximity labeling of RNA will help to accelerate the discovery of localized RNAs in bacteria.

Author contributions:

HY performed RNA labeling experiments. JMS and HY designed the study and wrote the paper. HY generated *C. crescentus* APEX2 fusions strains. DK generated *B. subtilis* APEX2 fusions strains. ES, OG, and OAC generated *E. coli* APEX2 fusion strains.

Acknowledgements:

We thank Tamara Hendrickson for equipment, reagents, and hosting the Schrader lab after it was destroyed in a fire. We thank Dr. Anat Nussbaum-Shochat for technical help in the construction of *ES413 E. coli* strain. NIH grants R35GM124733 to JMS and R35GM131783 to DBK. WSU Career Chair Award to JMS. IU startup funds to JMS. NIH T32GM142519-03 to HY. Research in the OAC lab was supported by the Israel Science Foundation (ISF) founded by the Israel Academy of Sciences and Humanities (grant no. 1274/19). OAC is an incumbent of the Dr. Jacob Grunbaum Chair in Medical Sciences.

Figure legends:

Figure 1. APEX2 proximity labeling of RNA in bacterial cells with minimal perturbation.
A) Schematic of APEX2 labeling of RNA protocol. The APEX2 protein was fused to RNase E,
the major protein that scaffolds BR-bodies. Cells are incubated in media containing
alkyne-phenol, and labeled was initiated with H_2O_2. After a brief labeling reaction, RNA was
extracted from the cells, and Cy5-azide was conjugated to the RNA by copper catalyzed
click chemistry. B) APEX2 fusion does not dramatically impact localization or function
(growth rate). Left: *In vivo* localization of RNase E-msfGFP vs RNase E APEX2-msfGFP
shows that fusion does not impact the formation of BR-bodies. Scale bar is 1µm. Middle:
Growth of RNase E-APEX2 fusion is similar to wild type *Caulobacter* cells. Right: mRNA
half-life measurements by qRT-PCR show that RNase E-APEX2 degrades mRNAs with a
similar half-life to wild-type. Data are from three biological and technical replicates. C)
APEX2 labeling requires H_2O_2 and alkyne-phenol to label RNAs. RNA labeling reactions
were placed on a nylon membrane to bind to the RNA in a dot blot apparatus and scanned
for Cy5 fluorescence in a gel imager. As a control, the RNA samples were incubated for 2
hours with DNase I and RNase A. The RNA was then precipitated before being subjected to
the azide-Cy5 click chemistry reaction and was re-precipitated before being filtered on the
nylon membrane in the dot blot apparatus.

Figure 2. APEX2 proximity labeling of RNA works rapidly. Optimization of APEX2 labeling
of RNA with Alkyne-Phenol. Top: Alkyne-phenol titration reveals that peak labeling occurs
with 2.5 mM Alkyne-Phenol. RNA was labeled in the scheme shown in Fig 1A, and the Cy5
intensity was measured in a gel imager. Bottom: Time course of APEX2 labeling. RNA was
labeled in the scheme shown in Fig 1A, and RNA labeling is apparent in as short as 15
seconds of H_2O_2 incubation, while peak labeling efficiency is observed at 45 seconds of
H_2O_2 incubation.

Figure 3. APEX2 proximity labeling of RNA works across species. APEX2 requires heme
for activity, so APEX2 fusions were generated in two additional species with heme
biosynthesis pathways, *B. subtilis* (gram +) and *E. coli* (gram –). Both species were and
subjected to RNA proximity labeling in mid-exponential phase of growth in LB media at
37°C, pre-incubated with alkyne-phenol for 30 minutes, and proximity labeling was
triggered by a 45 second incubation with H_2O_2. RNA labeling reactions were subjected to
azide-Cy5 click chemistry reactions, and 5 µg RNA placed on a nylon membrane to bind to
the RNA in a dot blot apparatus and scanned for Cy5 fluorescence in a gel imager.

Figure 4. APEX2 proximity labeled RNA can be isolated by streptavidin purification. A)
Schematic of the conjugation of biotin-azide to clicked alkyne-phenol and the resulting
streptavidin purification. B) Streptavidin purification of biotinylated RNAs. Tapestation

RNA profiles of the lysates and elution fractions of biotinylated proximity labeled RNAs. The bottom two bands are tRNAs, which are known to be highly depleted in BR-bodies from differential centrifugation-based isolation of BR-bodies[16,31]. C) RNA-seq enrichment analysis of proximity labeled RNA from BR-body + (RNaseE-APEX2, JS767) and BR-body – (RNaseEΔCTD-APEX2, JS801) cells. Log2 ratio of elution/lysate from each sample is plotted for each RNA quantified. Colored horizontal bars indicate the median values for each distribution. T-test with uneven variance was used to compare the distributions of each RNA type, and RNA types with $p<0.05$ are shown.

Figure S1. RNase E-APEX2-FLAG Western Blot. Protein lysates generated from *NA1000, Hfq-FLAG (LS4379)* and *RNase E-APEX2-FLAG (JS767)* were run on an SDS PAGE gel, transferred to PVDF, and probed with an anti-FLAG antibody.

Figure S2. Effects of copper catalyzed click reaction on RNA integrity. RNA extracted from APEX2 proximity labeling reactions were ran on an Agilent Tapestation from cells harboring RNase E-APEX2 fusion. The APEX2 labeled RNA was also incubated with Copper Sulfate under similar conditions for click chemistry reactions where we observe minimal RNA degradation from the copper.

Figure S3. RNA-seq data. A) Log2 enrichment (elution/lysate) for all four RNA proximity labeling biological samples. Mean values are shown in the horizontal bars. B) Correlation plots of the RPKM values between individual biological samples. R^2 values are shown between each sample for reference.

References:

1. Kannaiah, S., Livny, J., and Amster-Choder, O. (2019). Spatiotemporal Organization of the E. coli Transcriptome: Translation Independence and Engagement in Regulation. Mol. Cell *76*, 574-589.e7. https://doi.org/10.1016/j.molcel.2019.08.013.

2. Irastortza-Olaziregi, M., and Amster-Choder, O. (2021). RNA localization in prokaryotes: Where, when, how, and why. Wiley Interdiscip. Rev. RNA *12*, e1615. https://doi.org/10.1002/wrna.1615.

3. Campos, M., and Jacobs-Wagner, C. (2013). Cellular organization of the transfer of genetic information. Curr. Opin. Microbiol. *16*, 171–176. https://doi.org/10.1016/j.mib.2013.01.007.

4. Nandana, V., and Schrader, J.M. (2021). Roles of liquid–liquid phase separation in bacterial RNA metabolism. Curr. Opin. Microbiol. *61*, 91–98. https://doi.org/10.1016/j.mib.2021.03.005.

5.　Dugar, G., Svensson, S.L., Bischler, T., Wäldchen, S., Reinhardt, R., Sauer, M., and
Sharma, C.M. (2016). The CsrA-FliW network controls polar localization of the dual-
function flagellin mRNA in Campylobacter jejuni. Nat. Commun. *7*, 11667.
https://doi.org/10.1038/ncomms11667.

6.　Azaldegui, C.A., Vecchiarelli, A.G., and Biteen, J.S. (2021). The emergence of phase
separation as an organizing principle in bacteria. Biophys. J. *120*, 1123–1138.
https://doi.org/10.1016/j.bpj.2020.09.023.

7.　Darnell, R.B. (2010). HITS-CLIP: panoramic views of protein-RNA regulation in living
cells. Wiley Interdiscip. Rev. RNA *1*, 266–286. https://doi.org/10.1002/wrna.31.

8.　Hör, J., Di Giorgio, S., Gerovac, M., Venturini, E., Förstner, K.U., and Vogel, J. (2020).
Grad-seq shines light on unrecognized RNA and protein complexes in the model
bacterium Escherichia coli. Nucleic Acids Res. *48*, 9301–9319.
https://doi.org/10.1093/nar/gkaa676.

9.　Li, Y., Tian, C., Liu, K., Zhou, Y., Yang, J., and Zou, P. (2020). A Clickable APEX Probe for
Proximity-Dependent Proteomic Profiling in Yeast. Cell Chem. Biol. *27*, 858-865.e8.
https://doi.org/10.1016/j.chembiol.2020.05.006.

10.　Padron, A., Iwasaki, S., and Ingolia, N.T. (2019). Proximity RNA Labeling by APEX-Seq
Reveals the Organization of Translation Initiation Complexes and Repressive RNA
Granules. Mol Cell *75*, 875-887 e5. https://doi.org/10.1016/j.molcel.2019.07.030.

11.　Fazal, F.M., Han, S., Parker, K.R., Kaewsapsak, P., Xu, J., Boettiger, A.N., Chang, H.Y.,
and Ting, A.Y. (2019). Atlas of Subcellular RNA Localization Revealed by APEX-Seq. Cell
178, 473-490.e26. https://doi.org/10.1016/j.cell.2019.05.027.

12.　Zhou, Y., Wang, G., Wang, P., Li, Z., Yue, T., Wang, J., and Zou, P. (2019). Expanding
APEX2 Substrates for Proximity-Dependent Labeling of Nucleic Acids and Proteins in
Living Cells. Angew. Chem. Int. Ed Engl. *58*, 11763–11767.
https://doi.org/10.1002/anie.201905949.

13.　Al-Husini, N., Tomares, D.T., Bitar, O., Childers, W.S., and Schrader, J.M. (2018). α-
Proteobacterial RNA Degradosomes Assemble Liquid-Liquid Phase-Separated RNP
Bodies. Mol. Cell *71*, 1027-1039.e14. https://doi.org/10.1016/j.molcel.2018.08.003.

14.　Konkol, M.A., Blair, K.M., and Kearns, D.B. (2013). Plasmid-encoded ComI inhibits
competence in the ancestral 3610 strain of Bacillus subtilis. J. Bacteriol. *195*, 4085–
4093. https://doi.org/10.1128/JB.00696-13.

15.　Strahl, H., Turlan, C., Khalid, S., Bond, P.J., Kebalo, J.M., Peyron, P., Poljak, L., Bouvier,
M., Hamoen, L., Luisi, B.F., et al. (2015). Membrane recognition and dynamics of the
RNA degradosome. PLoS Genet *11*, e1004961.
https://doi.org/10.1371/journal.pgen.1004961.

16. Al-Husini, N., Tomares, D.T., Pfaffenberger, Z.J., Muthunayake, N.S., Samad, M.A., Zuo,
T., Bitar, O., Aretakis, J.R., Bharmal, M.-H.M., Gega, A., et al. (2020). BR-Bodies Provide
Selectively Permeable Condensates that Stimulate mRNA Decay and Prevent Release
of Decay Intermediates. Mol. Cell *78*, 670-682.e8.
https://doi.org/10.1016/j.molcel.2020.04.001.

17. Thanbichler, M., Iniesta, A.A., and Shapiro, L. (2007). A comprehensive set of plasmids
for vanillate- and xylose-inducible gene expression in Caulobacter crescentus. Nucleic
Acids Res. *35*, e137. https://doi.org/10.1093/nar/gkm818.

18. Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. Genome Biol. *10*,
R25. https://doi.org/10.1186/gb-2009-10-3-r25.

19. Bharmal, M.-H., Aretakis, J.R., and Schrader, J.M. (2020). An Improved Caulobacter
crescentus Operon Annotation Based on Transcriptome Data. Microbiol. Resour.
Announc. *9*. https://doi.org/10.1128/MRA.01025-20.

20. Murphy, K.C. (1998). Use of bacteriophage lambda recombination functions to
promote gene replacement in Escherichia coli. J. Bacteriol. *180*, 2063–2071.
https://doi.org/10.1128/JB.180.8.2063-2071.1998.

21. Thomason, L.C., Costantino, N., and Court, D.L. (2007). E. coli genome manipulation
by P1 transduction. Curr. Protoc. Mol. Biol. *Chapter 1*, 1.17.1-1.17.8.
https://doi.org/10.1002/0471142727.mb0117s79.

22. Kalocsay, M. (2019). APEX Peroxidase-Catalyzed Proximity Labeling and Multiplexed
Quantitative Proteomics. In Proximity Labeling: Methods and Protocols, M. Sunbul and
A. Jäschke, eds. (Springer), pp. 41–55. https://doi.org/10.1007/978-1-4939-9537-0_4.

23. Hung, V., Udeshi, N.D., Lam, S.S., Loh, K.H., Cox, K.J., Pedram, K., Carr, S.A., and Ting,
A.Y. (2016). Spatially resolved proteomic mapping in living cells with the engineered
peroxidase APEX2. Nat. Protoc. *11*, 456–475. https://doi.org/10.1038/nprot.2016.018.

24. Padrón, A., and Ingolia, N. (2022). Analyzing the Composition and Organization of
Ribonucleoprotein Complexes by APEX-Seq. Methods Mol. Biol. Clifton NJ *2428*, 277–
289. https://doi.org/10.1007/978-1-0716-1975-9_17.

25. Weissinger, R., Heinold, L., Akram, S., Jansen, R.-P., and Hermesh, O. (2021). RNA
Proximity Labeling: A New Detection Tool for RNA–Protein Interactions. Molecules *26*,
2270. https://doi.org/10.3390/molecules26082270.

26. Christen, B., Abeliuk, E., Collier, J.M., Kalogeraki, V.S., Passarelli, B., Coller, J.A., Fero,
M.J., McAdams, H.H., and Shapiro, L. (2011). The essential genome of a bacterium. Mol
Syst Biol *7*, 528–528. https://doi.org/10.1038/msb.2011.58.

27. Laalami, S., Zig, L., and Putzer, H. (2014). Initiation of mRNA decay in bacteria. Cell Mol
Life Sci *71*, 1799–1828. https://doi.org/10.1007/s00018-013-1472-4.

28. Jenniches, L., Michaux, C., Popella, L., Reichardt, S., Vogel, J., Westermann, A.J., and
Barquist, L. (2024). Improved RNA stability estimation through Bayesian modeling
reveals most Salmonella transcripts have subminute half-lives. Proc. Natl. Acad. Sci.
U. S. A. *121*, e2308814121. https://doi.org/10.1073/pnas.2308814121.

29. Muthunayake, N.S., Tomares, D.T., Childers, W.S., and Schrader, J.M. (2020). Phase-
separated bacterial ribonucleoprotein bodies organize mRNA decay. Wiley Interdiscip.
Rev. RNA *11*, e1599. https://doi.org/10.1002/wrna.1599.

30. Hamouche, L., Billaudeau, C., Rocca, A., Chastanet, A., Ngo, S., Laalami, S., and
Putzer, H. (2020). Dynamic Membrane Localization of RNase Y in Bacillus subtilis.
mBio *11*, e03337-19. https://doi.org/10.1128/mBio.03337-19.

31. Muthunayake, N.S., Al-Husini, N., and Schrader, J.M. (2020). Differential Centrifugation
to Enrich Bacterial Ribonucleoprotein Bodies (BR bodies) from Caulobacter
crescentus. STAR Protoc. *1*, 100205. https://doi.org/10.1016/j.xpro.2020.100205.

32. Nandana, V., Al-Husini, N., Vaishnav, A., Dilrangi, K.H., and Schrader, J.M. (2024).
Caulobacter crescentus RNase E condensation contributes to autoregulation and
fitness. Mol. Biol. Cell *35*, ar104. https://doi.org/10.1091/mbc.E23-12-0493.

33. Mackie, G.A. (2013). RNase E: at the interface of bacterial RNA processing and decay.
Nat. Rev. Microbiol. *11*, 45–57. https://doi.org/10.1038/nrmicro2930.

34. Bayas, C.A., Wang, J., Lee, M.K., Schrader, J.M., Shapiro, L., and Moerner, W.E. (2018).
Spatial organization and dynamics of RNase E and ribosomes in Caulobacter
crescentus. Proc Natl Acad Sci U A *115*, E3712–E3721.
https://doi.org/10.1073/pnas.1721648115.

35. Waters, L.S., and Storz, G. (2009). Regulatory RNAs in bacteria. Cell *136*, 615–628.
https://doi.org/10.1016/j.cell.2009.01.043.

36. Olejniczak, M., Jiang, X., Basczok, M.M., and Storz, G. (2022). KH domain proteins:
Another family of bacterial RNA matchmakers? Mol. Microbiol. *117*, 10–19.
https://doi.org/10.1111/mmi.14842.

37. Holmqvist, E., and Vogel, J. (2018). RNA-binding proteins in bacteria. Nat. Rev.
Microbiol. *16*, 601–615. https://doi.org/10.1038/s41579-018-0049-5.

38. Goldberger, O., Szoke, T., Nussbaum-Shochat, A., and Amster-Choder, O. (2022).
Heterotypic phase separation of Hfq is linked to its roles as an RNA chaperone. Cell
Rep. *41*, 111881. https://doi.org/10.1016/j.celrep.2022.111881.

39. Nandana, V., Rathnayaka-Mudiyanselage, I.W., Muthunayake, N.S., Hatami, A.,
Mousseau, C.B., Ortiz-Rodríguez, L.A., Vaishnav, J., Collins, M., Gega, A.,
Mallikaarachchi, K.S., et al. (2023). The BR-body proteome contains a complex network
of protein-protein and protein-RNA interactions. Cell Rep. *42*.
https://doi.org/10.1016/j.celrep.2023.113229.

40. Hondele, M., Sachdev, R., Heinrich, S., Wang, J., Vallotton, P., Fontoura, B.M.A., and
Weis, K. (2019). DEAD-box ATPases are global regulators of phase-separated
organelles. Nature *573*, 144–148. https://doi.org/10.1038/s41586-019-1502-y.

41. Krypotou, E., Townsend, G.E., Gao, X., Tachiyama, S., Liu, J., Pokorzynski, N.D.,
Goodman, A.L., and Groisman, E.A. (2023). Bacteria require phase separation for
fitness in the mammalian gut. Science *379*, 1149–1156.
https://doi.org/10.1126/science.abn7229.

42. Martell, J.D., Deerinck, T.J., Sancak, Y., Poulos, T.L., Mootha, V.K., Sosinsky, G.E.,
Ellisman, M.H., and Ting, A.Y. (2012). Engineered ascorbate peroxidase as a genetically
encoded reporter for electron microscopy. Nat. Biotechnol. *30*, 1143–1148.
https://doi.org/10.1038/nbt.2375.

A. Flow Chart of APEX2 Proximity Labeling of RNA

+H2O2 → RNA extraction → +CuSO4 + Sodium Ascorbate +Dye-azide →

Apex2 Alkyne-Phenol
RNase E RNA

Apex2 Alkyne-Phenol
RNase E RNA

Alkylated RNA

Fluorescently clicked RNA

B. APEX2 doesn't impact localization or protein function

RNase E-APEX2-GFP
1μm

RNase E-GFP

Cell Dilution

RNase E

RNase E-APEX2

RNase E RNase E-APEX2

5S rRNA

ctrA mRNA

Percent RNA remaining

Time (minutes)

C. APEX2 Proximity Labeling of RNA

APEX2:	–	–	–	–	+	+	+	+
Alk-Ph:	–	+	–	+	–	+	–	+
H₂O₂:	–	–	+	+	–	–	+	+

Mock   +RNase A   +DNase I

Optimized APEX2 RNA labeling occurs rapidly

Alk-Ph:

| 0 mM Alk-Ph | 0.5 mM Alk-Ph | 1 mM Alk-Ph | 2.5 mM Alk-Ph |

H_2O_2:

| 0 s | 15 s | 30 s | 45 s | 60 s |

A. APEX2 Proximity Labeling of RNA, clicked with biotin-azide for streptavidin purification

+CuSO$_4$
+Sodium ascorbate
+Biotin-peg3-azide

Mix

Stringent Washing → RNA Elution → RNA-seq

Alkylated RNA

Labeled and biotin clicked RNA

Streptavidin beads interact with biotinylated RNA

B. Streptavidin purification

Lysate | Streptavidin Elution

Ladder | NA1000 | RNaseE-APEX2 | NA1000 | RNaseE-APEX2

6000
4000
2000
1000 — 23s
500 — 16s
200
tRNA
25

C. RNA-seq enrichment

5S

p=5x10^{-302} p=5x10^{-3}

Log2 Enrichment (Elution/Lysate)

tRNA   rRNA   mRNA   sRNA   asRNA
BR-body + -   + -   + -   + -   + -