Overcoming immune evasion from post-translational modification of a mutant KRAS epitope to achieve TCR-engineered T cell-mediated antitumor activity

Jihoon William Lee[1,2], Emily Y. Chen[1], Taylor Hu[1], Rachel Perret[1], Mary E. Chaffee[1], Tijana Martinov[1], Shwetha Mureli[1], Clara L. McCurdy[1], Lisa A. Jones[3], Philip R. Gafken[3], Pritha Chanana[4], Yapeng Su[1,5], Aude G. Chapuis[1,6], Philip Bradley[5], Thomas M. Schmitt[1†*], Philip D. Greenberg[1,6,7†*].

[1] Program in Immunology, Division of Translational Science and Therapeutics, Fred Hutchinson Cancer Center, Seattle, WA, USA

[2] Medical Scientist Training Program, University of Washington, Seattle, WA, USA

[3] Proteomics and Metabolomics Shared Resource, Fred Hutchinson Cancer Center, Seattle, WA, USA

[4] Genomics and Bioinformatics Shared Resource, Fred Hutchinson Cancer Center, Seattle, WA, USA

[5] Herbold Computational Biology Program, Fred Hutchinson Cancer Center, Seattle, WA

[6] Department of Medicine, Division of Hematology and Oncology, University of Washington, Seattle, Washington, USA

[7] Department of Immunology, University of Washington, Seattle, WA

† Corresponding author(s): Philip Greenberg, MD, Thomas Schmitt, PhD
* Co-senior authors who contributed equally.
Email: pgreen@uw.edu, tschmitt@fredhutch.org

Author Contributions: Conceptualization: J.W.L., R.P., T.M., P.B., A.G.C., T.M.S., P.D.G.; Methodology: J.W.L., E.Y.C., R.P., M.E.C., T.M., Y.S., P.B., T.M.S., P.D.G.; Investigation: J.W.L., E.Y.C., T.H., R.P., M.E.C., T.M., S.M., C.L.M., L.N.J., P.R.G., P.C., P.B.; Formal analysis: J.W.L., E.Y.C., R.P., M.E.C., T.M., P.B., T.M.S.; Resources: A.G.C., P.B., T.M.S., P.D.G.; Writing – original draft: J.W.L., E.Y.C.; Writing – review and editing: J.W.L., E.Y.C., T.H., R.P., M.E.C., T.M., L.N.J., P.R.G., P.C., Y.S., A.G.C., P.B., T.M.S., P.D.G.; Visualization:

J.W.L., E.Y.C., T.M.S.; Supervision: P.B., T.M.S., P.D.G.; Funding acquisition: J.W.L., R.P., T.M., A.G.C., P.B., T.M.S., P.D.G.

Competing Interest Statement:

P.D.G. is on the Scientific Advisory Board of Celsius, Earli, Elpiscience, Immunoscape, Rapt, Catalio, and Nextech, was a scientific founder of Juno Therapeutics, and receives research support from Lonza. R.P., T.M., A.G.C., T.M.S., and P.D.G. are co-inventors on a patent associated with technology described in this work that is licensed to Affini-T. A.G.C., T.M.S., and P.D.G. are co-founders of, have equity in, and receive research support from Affini-T.

Abstract

T cell receptor (TCR)-T cell immunotherapy, in which T cells are engineered to express a TCR specific for a tumor epitope, is a form of adoptive cell therapy (ACT) that has demonstrated promise against various tumor types. Mutants of oncoprotein KRAS, particularly at glycine-12 (G12), are frequent drivers of tumorigenicity, but also attractive targets for TCR-T cell therapy. However, MHC class I-restricted TCRs specifically targeting G12-mutant KRAS epitopes in the context of tumors expressing HLA-A2, the most common human HLA-A allele, have remained elusive despite evidence that an epitope encompassing the mutation can bind HLA-A2 and induce T cell responses. We report that post-translational modifications of the protein on this epitope may allow tumor cells to evade immunologic pressure from TCR-T cells. A lysine side chain-methylated KRAS$_{G12V}$ peptide, rather than unmodified epitope, may be presented in HLA-A2 by tumor cells and impact recognition by TCRs. Using a novel computationally guided approach to design TCRs, we developed by mutagenesis TCRs that recognize this methylated peptide, enhancing tumor recognition and destruction. Additionally, we identified TCRs with similar functional activity in normal repertoires from rare primary T cells by stimulation with modified peptide, clonal expansion, and selection. Mechanistically, a gene knockout screen to identify mechanism(s) by which tumor cells methylate or demethylate this epitope unveiled SPT6 as a demethylating protein that could be targeted to improve effectiveness of these TCRs. These findings highlight the role of post-translational modifications in immune evasion and suggest that identifying and targeting such modifications should facilitate development of more effective TCR-T cell therapies.

Introduction

T cell receptors (TCRs) specific for a tumor-derived MHC class I epitope can be transduced into patient T cells for adoptive cell therapy (ACT) to achieve promising antitumor effects[1, 2], and a subset of such TCRs have been advanced into clinical trials[3, 4, 5]. The most desirable TCR-T cell therapies target antigens that are tumor-specific, contribute to and/or are essential to the malignant phenotype, and are well-presented by the tumor that is being targeted. Mutant KRAS is the most common driver oncogene in human malignancies, with 85% of gain-of-function KRAS mutations resulting from a change in the glycine amino acid at the 12th residue[6]. In particular, mutation of glycine to valine, which is frequently detected in pancreatic (34%), colon (21%), and lung (20%) cancers[6, 7] as well as other tumors, has been postulated to create an epitope (KRAS$_{G12V}$, residues 5-14) presented on the tumor cells by HLA-A2[8], the most

common Class I allele worldwide[9]. As tumors often become oncogene-addicted to G12-mutant

KRAS[10], these mutations are unlikely to be lost or revert to wild type, making epitopes

containing G12 mutations of KRAS highly sought after as targets for tumor-specific TCR-T cell

therapy.

TCRs targeting KRAS epitopes containing a mutation at G12 presented by several HLA

alleles including HLA-A3[11], HLA-A11[4, 12], and HLA-C8[13] have been reported, which not only

recognize tumors *in vitro* but also exhibit therapeutic efficacy *in vivo*[4, 14]. However, the

accessibility of this therapeutic strategy is limited by the low frequency of these alleles in the US

(e.g., 20–24% for HLA-A3 and less frequent for HLA-A11 and HLA-C8)[15]. In the US, HLA-A2 is

the most common allele (47.6%[16]); thus, targeting an A2-restricted KRAS epitope would enable

therapy for a far more substantial fraction of the population. However, defining an A2-restricted

KRAS epitope has proven problematic, including difficulties confirming within the sensitivity of

mass spectrometry (MS) that an A2-restricted epitope spanning the G12 position is present in

the immunopeptidome[17, 18]. However, $KRAS_{G12V}$ does have epitopes predicted to bind well to

HLA-A2, particularly residues 5-14 (KLVVVGAVGV)[8, 19]. As MS has been able to isolate G12V-

containing epitopes and other mutant KRAS epitopes from other HLA alleles[18], it has been

presumed that the HLA-A2 epitope may not be naturally processed and/or efficiently presented

by HLA-A2[17, 18].

Several mechanisms may cause defective processing or presentation of a predicted

epitope, leading to immune evasion. Examples include differential expression of proteasome

subunits that can alter processing of a protein into an epitope[20], defective transportation of a

processed epitope to the endoplasmic reticulum for binding to MHC I[21], and deficiencies in MHC

I expression or structure[22]. Additionally, when an epitope is presented, it may contain post-

translational modifications (PTMs), resulting in complete[23] or partial abrogation of epitope

binding or recognition by a TCR reactive to the unmodified epitope. Such changes include

oxidation[23], nitrosylation[23], methylation[24], phosphorylation[25], and glycosylation[26]; some PTMs

may also be more prevalent in tumors as a consequence of malignant transformation and

enzyme dysregulation[27, 28]. PTMs do occur in mutated KRAS[29], and could interfere with

detection of an HLA-A2-presented KRAS epitope by MS due to an altered migration pattern,

contributing to the lack of consensus regarding an A2 epitope.

Identifying a targetable epitope with a PTM for $KRAS_{G12V}$ is challenging due to

experimental and analytical limitations of MS in detecting PTMs[30, 31, 32], such as introduction of

confounding chemical artifacts[32], and PTMs have generally been overlooked when predicting

epitopes for therapeutic TCR development[33]. For the KRAS protein, PTMs are known to occur in

both the C-terminal region, which are well-characterized for their signaling functions[28, 34], and

the N-terminal region with methylation of the lysine-5 side chain[29], which is included in the

epitopes predicted to be presented by HLA-A2.

In this investigation, we demonstrate that methylation of the lysine-5 side chain in the

$KRAS_{G12V}$ epitope that binds to HLA-A2 may interfere with targeting of $KRAS_{G12V}$ tumor lines by

TCRs specific for the unmodified epitope. Combining machine learning-based computational

approaches to model TCR:pMHC interfaces with high-throughput experimental screens, we

developed TCRs that target methylated and unmodified epitopes, thereby providing broader

anti-tumor activity against HLA-A2$^+$, $KRAS_{G12V}^+$ tumor cell lines. Finally, we reveal the potential

role of a protein canonically known to diminish histone methylation, SPT6[35], in reducing

methylation of the $KRAS_{G12V}$ epitope. This may be targetable in tumor cells as a method to

enhance immunologic recognition by methylation-targeting TCR-T cells and exemplifies a

potential synergistic strategy to inhibit tumor evasion of T cell immunity.

Results

<u>Post-translational modifications of the HLA-A2-presented $KRAS_{G12V}$ epitope alter its recognition</u>

<u>by MHC class I-restricted TCRs and prevent tumor cell killing.</u>

Computational prediction, by NetMHCPan 4.1[36], of HLA-A2-presented $KRAS_{G12V}$

identified several epitopes that specifically contain the G12V mutation (**Supplementary Figure**

1a). Among the top ten predicted epitopes encompassing the full mutated KRAS protein, three

contained the G12V mutation, and the epitope consisting of residues 5-14 (i.e., KLVVVGAVGV)

was predicted to bind with the highest avidity.

For our initial effort to develop HLA-A2-restricted TCRs targeting this peptide, we

isolated primary CD8$^+$ T cells from peripheral blood mononuclear cells (PBMCs) from HLA-A2$^+$

healthy human donors and expanded epitope-reactive clonotypes by stimulation with

autologous antigen-presenting cells (APCs) pulsed with a synthetic $KRAS_{G12V}$ 5-14 peptide that

lacked any PTMs (**Materials and Methods**). We sorted cells with potentially high affinity for the

$KRAS_{G12V}$ peptide as measured by binding limiting concentrations of HLA-A2:$KRAS_{G12V(5-14)}$

tetramers. From the sorted cells, we then quantitated TCR clonotypes using the ImmunoSEQ

assay (Adaptive Biotech) and identified paired TCRα/TCRβ sequences using single cell RNA-

seq. TCR clonotypes preferentially enriched within the sorted populations were synthesized as

codon-optimized P2A-linked constructs of TCRα and TCRβ in a lentiviral backbone and used to

transduce CD8$^+$ T cells. The cells were stimulated with titrated concentrations of $KRAS_{G12V(5-14)}$

peptide, and the resulting IFNγ expression measured to assess functional avidities for this

peptide (**Figure 1a; Supplementary Figure 1b**). Two particularly high affinity TCRs were

identified, TCR_{19} and TCR_2, with the former exhibiting the highest functional avidity. However, in

a coculture with the HLA-A2$^+$, $KRAS_{G12V}{}^+$ pancreatic adenocarcinoma cell line CFPAC1, CD8$^+$ T

cells transduced with TCR_{19} unexpectedly demonstrated poorer inhibition of tumor growth

compared to T cells transduced with TCR_2 (**Figure 1b**), whereas TCR_{19} conversely

demonstrated better inhibition than TCR_2 of DAN-G tumor cell growth, which are also HLA-A2$^+$

and $KRAS_{G12V}{}^+$.

The reasons for this discrepancy between TCR functional avidities for $KRAS_{G12V(5\text{-}14)}$ and

killing of different tumor cells were unclear. One explanation could be alloreactivity rather than

tumor antigen specificity. For killing of CFPAC1 cells, alloreactivity to this cell line by TCR_2

would not necessarily have been predicted by measuring the dose response of cells with this

TCR to peptides presented by another antigen-presenting cell. However, TCR_2 did not induce a

reaction against any of the Class I HLA alleles found on CFPAC1 cells on other targets not

expressing mutant KRAS (HLA-A2, A3, B35, B73, C4, C15; **Supplementary Figure 1c**).

Alternatively, TCR_{19} killing of DAN-G cells could reflect allorecognition, but similar studies did

not detect recognition of the alleles on DAN-G by TCR_{19}. Thus, alloreactivity was an unlikely

explanation. We therefore hypothesized that, while both of these HLA-A2$^+$ $KRAS_{G12V}{}^+$ tumor

cells may be presenting the epitope, one may be presenting it in a modified form, and that there

was differential recognition of the modified and unmodified peptide by each TCR.

Before more closely examining the presence of PTMs, we wanted to confirm that the

predicted primary sequence, $KRAS_{G12V}$ 5-14, is in fact efficiently generated by proteasomal

cleavage and presented by HLA-A2. To accomplish this, we used the ARTEMIS technology,

which entails MS-based identification of MHC I epitopes bound to a secreted single-chain dimer

(SCD) of the selected HLA I allele[37] (**Materials and Methods**). This provides an abundant

source of easily purified peptide-bound Class I molecules, can detect the majority of peptides

that get endogenously processed and loaded on Class I in the ER, and overcomes the

challenge of identifying the limited amount of peptide specifically presented on the cell surface

by an individual Class I allele using MS[37]. We established an ARTEMIS system in the 293F cell

line, which supports exceptionally abundant protein expression compared to other mammalian

cell lines[38]. 293F cells were lentivirally transduced to express a secreted HLA-A2 SCD as well

as a truncated version of the $KRAS_{G12V}$ protein (amino acids 1–100 to render the protein

nonfunctional but preserve the N-terminal portion of the protein that contains the G12V

mutation), to increase the availability of $KRAS_{G12V}$-originated peptides that contain the G12V

mutation for binding to HLA-A2. Three HLA-A2-presented KRAS peptides were identified by

ARTEMIS, including the KRAS$_{G12V}$ 5-14 peptide, which was the only HLA-A2-presented peptide

containing the mutated G12V residue (**Figure 1c**). These results confirmed that the 5-14

epitope is a dominant product of KRAS$_{G12V}$ antigen processing and presentation associated with

HLA-A2 and is the correct peptide sequence to target with TCRs.

Given the known dysregulation of methylation processes in mutant KRAS-driven

cancers[27, 28, 39, 40], and the previously detected lysine-5 side chain methylation of Ras family

proteins[29], we considered the possibility that the KRAS$_{G12V}$ 5-14 epitope presented by tumor

cells contains some level of methylation or other post-translational modification. Having

confirmed the primary sequence of the epitope, we hypothesized that DAN-G cells were

presenting KRAS$_{G12V}$ 5-14 epitope such that some combination of lysine-5-methylated and

unmodified epitope was likely being presented. In turn, the presence of the weakly recognized,

modified epitope may have limited the recognition of these tumor cells by TCR$_2$ and TCR$_{19}$,

thereby limiting tumor killing. However, as 293F cells may not have the same protein

methylation machinery active in DAN-G, we attempted to design ARTEMIS in DAN-G cells.

Unfortunately, due to the low cell density of cultured DAN-G cells and lower level of protein

expression compared to 293-based ARTEMIS cells, we did not detect sufficient levels of peptide

to identify the methylated epitope. Therefore, to search for differences in methylation activities

that can lead to increased methylation within KRAS in DAN-G cells, we also performed

immunoprecipitation of whole-protein KRAS from DAN-G and CFPAC1 cells, followed by liquid

chromatography-MS. This revealed that in DAN-G cells, but not in CFPAC1 cells, lysine side

chain trimethylation was present at lysine-16 (**Supplementary Figure 1d**), representing full

replacement of all hydrogens of the amine group by methyl groups. No peptides containing

lysine-5 of KRAS were able to be recovered or detected in this data set, likely due to the

cleavage of the KRAS protein by trypsin immediately after lysine-5 during sample processing for

MS[41], producing a short 5-mer peptide that would be difficult to detect with our MS technology.

Thus, while we could not directly confirm the presence of a lysine-5-methylated peptide, a

pattern of increased KRAS lysine methylation was evident in DAN-G cells. Additionally, previous

reports clearly demonstrated methylation of KRAS at lysine residues[28, 29] including dimethylation

at lysine-5. Lysine side chain methylation may exist as mono-, di-, or trimethylation, with the

former two states being intermediate degrees of full methylation.

To determine whether the KRAS$_{G12V(5-14)}$ peptide with lysine-5 methylation binds to HLA-

A2 with comparable affinity as the unmodified peptide, we assessed the stability of HLA-

A2:KRAS$_{G12V}$ pMHC complexes with or without lysine-5 side chain methylation. T2 cells are

deficient in Transporter Associated with Antigen Processing (TAP) protein and express few

properly folded Class I molecules on the cell surface[42]. Pulsing T2 cells with exogenous peptide

can stabilize the conformation of HLA-A2 molecules on the cell surface, resulting in enhanced

surface expression that correlates with peptide affinity for HLA-A2[42]. We pulsed T2 cells with

$KRAS_{G12V(5-14)}$ with or without methylation and measured the resulting surface expression of

HLA-A2 (**Materials and Methods; Supplementary Figure 1e**). Compared to untreated control

cells, HLA-A2 surface expression was higher when T2 cells were pulsed with the $KRAS_{G12V(5-14)}$

peptide. Moreover, pulsing with the same peptide containing different degrees of lysine-5

methylation further increased A2 expression, indicating that methylation may enhance the stable

presentation of the epitope. Although the unmodified peptide was recognized well by both

TCRs, there were large differences in the abilities of the TCRs to recognize the methylated

epitope (**Figure 1d**). Therefore, differential lysine 5-methylation may explain the disparities in

the killing of different tumor cells observed with these TCRs.

In addition to the 10-mer $KRAS_{G12V}$ 5-14 sequence detected from ARTEMIS, a 9-mer

(LVVVGAVGV) is another predicted epitope containing the G12V mutation that bind HLA-A2

(**Supplementary Figure 1a**), and a previous study has also reported peptide splicing to

produce an HLA-A2-presented epitope containing residues 5-6/8-14 of $KRAS_{G12V}$[19]. To rule out

recognition of these alternative epitopes by the TCRs and to test recognition of methylated

epitope, we synthesized these epitope variants as well as $KRAS_{G12V(5-14)}$, including this peptide

post-translationally modified at lysine-5. To determine the functional avidities of the TCRs for

these variant epitopes, we stimulated T cells expressing TCR_2 or TCR_{19} with these epitopes as

presented by HLA-A2 and measured resulting IFNγ expression (**Figure 1d**). Indeed, while both

TCR_2 and TCR_{19} recognized unmethylated $KRAS_{G12V}$ 5-14 to similar degrees, only TCR_{19}

produced a response to the peptide with methylation of the amine group on the side chain of

lysine-5. Other forms of the G12V-containing epitope which had lower predicted binding ability

from NetMHCPan 4.1 (residues 6-14 or 5-12), and an alternatively spliced peptide $KRAS_{G12V}$ 5-

6/8-14 previously suggested to be processed[19], also did not elicit a response from the TCRs.

Thus, the DAN-G cell line may be modifying a fraction of $KRAS_{G12V}$ protein or its resulting

epitopes with lysine-5 side chain methylation, such that the epitope presented by this tumor cell

could not be recognized by TCR_2. The presence of lysine-5 methylation is consistent with prior

findings examining methylation of RAS family proteins, including KRAS[29], and is consistent with

the known dysregulation of protein methylation in mutant KRAS-driven cancers[27, 28].

Based on the evidence supporting contribution of a lysine-5 methylated peptide to TCR

recognition of the HLA-A2-presented $KRAS_{G12V(5-14)}$ epitope, we sought to identify structural

changes that may occur as a result of such methylation. Using the Rosetta[43] protein structure

modeling tool, we introduced monomethylation, dimethylation, and trimethylation to the lysine-5

side chain and evaluated the resulting changes in peptide positioning in the MHC cleft. This

computational pipeline indicated that methylation of the lysine-5 side chain within HLA-A2-

presented $KRAS_{G12V(5-14)}$ induces shifts of the positioning of the amine group of this side chain

that could affect TCR recognition of the epitope (**Figure 1e**): in the unmodified peptide, the

amine group of the lysine-5 side chain protrudes out of the MHC I cleft potentially into close

proximity to the TCR chain, suggesting it may serve as a contact point between the TCR and

pMHC complex. Monomethylation leads to displacement of the amine group deeper into the

MHC I cleft by the methyl group, which is further pronounced by dimethylation or trimethylation,

potentially explaining the observed impact of methylation on the recognition of this pMHC

complex by TCR_2 or TCR_{19}.

To evaluate the relative contribution of the lysine-5 residue to binding of each TCR to the

HLA-A2-presented $KRAS_{G12V}$ epitope, we performed alanine scans, in which each residue of the

HLA-A2-presented epitope is sequentially substituted with the small, neutral alanine amino acid.

Each alanine mono-substituted epitope was then used to pulse antigen-presenting cells to

stimulate CD8$^+$ T cells expressing TCR_2 or TCR_{19}, and IFNγ was then measured to determine if

recognition of the whole epitope is impacted by the alanine substitution[44], which would imply

that the substituted amino acid is a relevant contact point for the TCR. Residue 11 of the 5-14

epitope, which is already alanine, was alternatively substituted with threonine or glycine. At

peptide concentrations of 0.1 µg/ml, the analysis revealed that substitution of lysine-5 to alanine

reduced the response to the epitope from 67.9% responsive cells to 2.45% for TCR_2 (**Figure**

1f), suggesting that TCR_2 is sensitive to changes at that position of the epitope. The pMHC

complex with $KRAS_{G12V}$ epitope, as modeled by Rosetta, also indicates protrusion of the lysine-

5 side chain out of the groove of HLA-A2 in the direction where it could interact with a TCR

(**Supplementary Figure 1f**, which uses the same models as **Figure 1e** but is rotated for display

from a perspective showing this protrusion more clearly). By contrast, the alanine scan for

TCR_{19} revealed high tolerance for amino acid substitutions at most positions of the $KRAS_{G12V}$

epitope, which may explain its recognition of methylated lysine-5 but suggesting it would be

promiscuous and less suitable as a potentially therapeutic TCR to modify for pursuing

improvements (**Figure 1f**).

A rationalized mutagenesis approach can be used to generate TCRs that recognize methylated

$KRAS_{G12V}$ epitope to improve targeting of tumor cells.

Identifying TCRs that can recognize lysine-5 side chain mono-, di-, and/or trimethylation

of the KRAS$_{G12V}$ 5-14 peptide alone or in addition to the unmodified version of that peptide could

potentially increase the ability to therapeutically target HLA-A2$^+$ tumors with the KRAS$_{G12V}$

mutation. To generate new TCRs that can target the peptide with the three different degrees of

lysine-5 methylation, we first explored a strategy to mutagenize the CDR3α region of an existing

TCR, TCR$_2$, which recognizes the unmodified peptide with high functional avidity and specificity

for the KRAS$_{G12V}$ epitope, to determine if we could broaden peptide recognition to include

methylated variants without acquisition of more general promiscuity.

To determine the region of this TCR most likely to interact with the methylation site, we

developed a new computational structural prediction workflow (**Supplementary Figure 1g**). We

first used Alphafold[45], which utilizes deep learning-based approaches to predict the structure of

the TCR:pMHC complex without any PTMs, as the latter are not modeled well by Alphafold. We

fed the output into Rosetta, which utilizes physics-based energy functions to allow for modeling

additional molecular changes to that structure, such as introducing PTMs to the epitope, and

then readjusting the structural prediction. Our workflow, performed on the existing TCR$_2$ that

recognized the unmodified KRAS$_{G12V}$ 5-14 peptide presented by HLA-A2, revealed a core

sequence within the CDR3α region most likely to be in close proximity to the lysine-5 side chain

and any potential methylation modifications to this residue (**Figure 2a**).

Using the structural model as the guide to the most relevant region, we created a

rationalized mutagenesis library of this computationally identified core CDR3α sequence.

Library-transduced CD8$^+$ Jurkat cells were stimulated to detect TCR clonotypes that recognized

differently modified variants of the KRAS$_{G12V}$ 5-14 peptide (**Materials and Methods**). Library-

transduced T cells were analyzed for binding to unmethylated or lysine-5-methylated

KRAS$_{G12V(5-14)}$ peptide. A broad spectrum of reactivity towards one or both of these peptides

was observed, confirming that the computationally determined CDR3α sequence plays a key

role in recognition of the putative lysine-5 methylation site by TCR$_2$. We identified several

distinct clonotypes able to bind to only the unmethylated peptide, to a combination of

unmethylated and mono/di/trimethylated epitopes, or to only methylated peptides (**Figure 2b–**

c), which were sequenced to identify the corresponding mutant variants. New constructs were

then generated expressing these mutant variant TCRs. A dose response assay with CD8$^+$ T

cells transduced with each mutant variant was performed using the unmethylated and

methylated KRAS$_{G12V}$ epitopes individually. Some mutagenized TCRs, when compared to TCR$_2$,

had improved recognition of the methylated epitope (**Figure 2d**). A dose response assay

revealed no significant acquisition of recognition of wild type KRAS 5-14 or the alternative

KRAS$_{G12D}$ 5-14 epitope by the mutagenized TCRs compared to TCR$_2$ (**Supplementary Figure**

2a). Thus, changes in amino acids within only this computationally identified core CDR3α

sequence were able to alter the TCR's pattern of recognition for peptides with PTMs

(**Supplementary Figure 3a–b**) without altering recognition of non-G12V KRAS epitopes. For

TCR$_{EDST}$, labeled as such for the mutagenized 'core' CDR3α sequence, the amino acids in

closest proximity to the lysine-5 side chain of the HLA-A2-presented KRAS$_{G12V}$ epitope were

aspartate and serine. These amino acids are negatively charged and polar, respectively,

allowing for electrostatic or hydrogen bonding interactions with the lysine-5 side chain amine

group of the KRAS$_{G12V}$ epitope. While methylation introduces a hydrophobic methyl group, the

polar nature of the amino acids on this TCR may still allow for binding to the lysine side chain.

We then tested whether CD8$^+$ T cells expressing the TCR variants with enhanced

binding to methylated peptides could efficiently recognize and kill KRAS$_{G12V}$-expressing tumor

cells. In coculture with KRAS$_{G12V}$$^+$ HLA-A2$^+$ tumor cells, CD8$^+$ T cells expressing TCR$_{EDST}$ or

TCR$_{EDRS}$, which recognize unmethylated and methylated KRAS$_{G12V}$ epitopes, demonstrated

improved killing of DAN-G cells, which likely presents methylated and unmethylated KRAS$_{G12V(5-}$

$_{14)}$, and retained similar killing of CFPAC-1 cells compared to TCR$_2$ (**Figure 2e**). Consistent with

the lack of recognition of wild type KRAS$_{5-14}$ or KRAS$_{G12D(5-14)}$ in the peptide dose response

assay, these TCRs did not inhibit growth of HeLa cells, which express only wild type KRAS 5-

14, or of Panc1 cells, which express KRAS$_{G12D}$ (**Supplementary Figure 2b**). Taken together,

mutagenesis of the originally naturally isolated TCR$_2$ yielded new TCRs recognizing different

combinations of unmodified and varyingly methylated KRAS$_{G12V}$ epitopes without increasing "off-

target" recognition of G12D or wild type epitopes of KRAS.

A critical concern with TCRs deployed for ACT is that specificity needs to be limited only

to the intended targets to avoid off-target toxicities, which is of increased concern after a

naturally isolated TCR is mutagenized. To further evaluate the specificity of the selected TCRs,

we again performed alanine scans of the target epitope for recognition by the mutagenized

TCRs in Jurkat reporter cells expressing CD8αβ, TCR$_2$ or TCR$_{19}$, and the Nur77-GFP reporter

gene as a specific marker of TCR signaling[46]. At 0.1 µg/ml peptide concentrations, with

tolerance to a change defined as retaining >35% of responding cells positive for Nur77-GFP

reporter expression, there were distinct positions of the epitope that tolerated amino acid

replacements and remained stimulatory (**Supplementary Figure 2c**). The overall tolerance

pattern was K-x-x-V-V-x-A-x-x-x, with x signifying tolerance for an alanine substitution at that

position. Notably, while some TCRs could recognize both unmethylated and methylated

versions of the lysine-5 side chain, replacement of lysine-5 with alanine led to a loss of

recognition of the presented peptide. The diminution of recognition was greater than observed

with substitutions at other peptide positions, suggesting this lysine amino acid remained a major

contact point for the mutagenized TCRs as it did for TCR_2. A scan of the human peptidome for

epitopes that accommodated any substitution at these permissive residues yielded $PLD3_{396-405}$,

$MIGA1_{74-83}$, $NAGPA_{276-285}$, K/N/$HRAS_{5-14}$, $RSLBB_{35-44}$, $CFA61_{23-32/607-616/667-676}$, $CUL9_{635-644/745-754}$,

$CLCN6_{80-89}$, and $MACOI_{33-42}$ as potential off-target hits. These peptides were synthesized and

used to stimulate T cells expressing the TCR. Relative to HLA-A2-presented $KRAS_{G12V}$ 5-14, all

of the selected TCRs exhibited reduced responses against these peptides at 0.1 µg/ml

concentration (**Supplementary Figure 2d**), suggesting the mutagenized TCRs had not

acquired significant promiscuity compared to the original TCR_2. Additionally, for TCR_2 and

TCR_{EDST}, at more physiologic concentrations of 0.01 µg/ml, these peptides elicited response

rates of <6% compared to 18–19% for the $KRAS_{G12V}$ epitope, as measured by percentage of

cells expressing IFNγ (**Supplementary Figure 2e**).

Stimulation of primary T cells with selected peptides followed by clonal expansion reveals TCRs

from healthy donor repertoires that recognize methylated $KRAS_{G12V}$ epitope and can target

tumor cells.

The TCRs generated from the mutagenesis approach were able to specifically target

and effectively kill HLA-A2 tumor cell lines expressing $KRAS_{G12V}$ with PTMs. However, such

TCRs have not been naturally vetted by the thymic selection process, and thus may pose a risk

for cross-reactivity with other peptides from the human proteome despite the safety testing

based on the alanine scan. Thus, to identify TCRs from normal peripheral repertoires potentially

capable of binding to the methylated KRAS epitopes, primary human T cells from healthy donor

PBMCs were stimulated with $KRAS_{G12V}$ peptides including unmethylated, mono-, di-, and tri-

methylated lysine-5, and responding T cells were sort purified based on specific binding to

individual HLA-A2:PTM-containing $KRAS_{G12V}$ 5-14 tetramers at limiting peptide concentrations.

Quantitative TCR repertoire analysis was performed on the purified populations, and highly

enriched TCR clonotypes were synthesized and expressed in primary $CD8^+$ T cells (**Materials**

and Methods; Supplementary Figure 4a, b).

As measured by IFNγ expression by TCR-transduced T cells after exposure to

unmethylated and methylated $KRAS_{G12V}$ epitopes presented by HLA-A2, these TCRs exhibited

dose response patterns specific to the G12V mutation, responded to methylated and/or

unmethylated peptides, and did not recognize wild type or G12D mutant epitopes (**Figure 3a**).

The functional avidities of these TCRs, $TCR_{A2UoM1-1}$, $TCR_{A2UoM1-2}$, and $TCR_{A2UoM1-4}$, for the

unmethylated and monomethylated peptides were comparable to those of the mutagenesis-

derived TCRs, in particular TCR_{EDST} which exhibited superior killing of DAN-G cells and strong

recognition of unmethylated and methylated $KRAS_{G12V}$ epitopes. Among the TCRs that exhibited

the strongest recognition of methylated and unmodified peptides, coculture with tumor cells of

$CD8^+$ T cells expressing $TCR_{A2UoM1-1}$ demonstrated superior killing compared to T cells

expressing TCR_2 of the HLA-A2$^+$ $KRAS_{G12V}{}^+$ DAN-G tumor cell line, inferred to present

methylated and unmethylated $KRAS_{G12V}$ 5-14 (**Figure 3b**). $TCR_{A2UoM1-1}$ also exhibited similar

killing as the other TCRs against CFPAC1, which presumably presents unmethylated $KRAS_{G12V}$

5-14. $TCR_{A2UoM1-1}$ did not recognize titrated doses of HLA-A2-presented wild type KRAS 5-14 or

$KRAS_{G12D}$ 5-14 (**Supplementary Figure 4c**) but did exhibit some inhibition of HLA-A2$^+$ tumor

cell lines expressing either wild type or G12D KRAS (**Supplementary Figure 4d**), suggesting

that there may be some broader targeting. However, an alanine scan with $TCR_{A2UoM1-1}$

demonstrated very limited tolerance for amino acid substitutions within the $KRAS_{G12V}$ 5-14

epitope (**Supplementary Figure 4e**), such that the only potential match in the human

peptidome (using the same criteria as our previous alanine scans) was $EPIPL_{1948-1957}$. A dose

response assay performed with this peptide revealed no recognition by this TCR

(**Supplementary Figure 4f**), affirming its highly selective specificity.

Structural modeling of $TCR_{A2UoM1-1}$ revealed a CDR3α sequence that had amino acids

aspartate and serine in closest proximity to the lysine-5 side chain including the differently

methylated versions (**Figure 3c**). These TCR amino acids are identical to what was identified in

the mutagenized TCR_{EDST}, and thus may allow for close interaction with both methylated and

unmethylated lysine-5 of the $KRAS_{G12V}$ epitope by both TCRs.

<u>SPT6 is involved in the mechanism for demethylation of lysine-5 of the $KRAS_{G12V}$ epitope in</u>

<u>tumor cells and may be a target for creating presentation of a stable epitope with a methylation-</u>

<u>specific TCR.</u>

We next sought to determine potential genes involved in $KRAS_{G12V}$ epitope methylation.

To address this, we performed a CRISPR-Cas9 library screen targeting genes canonically

known to be involved in DNA or protein methylation (**Materials and Methods; Figure 4a**). The

library was transduced into the DAN-G cell line, which is inferred to express the methylated

$KRAS_{G12V(5-14)}$ epitope and is recognized by TCRs that have specificity for the methylated

epitope but not effectively by TCRs only specific for the unmodified epitope. Tumor cells were

collected before and after co-culture with methylation-responsive TCR-T cells (TCR_{EDST}) and

TCR-T cells that preferentially target unmodified KRAS (TCR_2), and the frequencies of each

gRNA in surviving tumor cells were determined. TCR_{EDST} was chosen as the methylation-

sensitive TCR for this comparison because it differs from TCR_2 only in its ability to recognize

methylated lysine-5. This analysis revealed reduced recovery of certain gene knockouts when

cocultured with TCR_{EDST}, but not when cocultured with TCR_2, suggesting that the knocked-out

gene conferred a survival disadvantage for tumor cells that might reflect increased presentation

of the methylated epitope. In contrast, a higher frequency of gRNAs targeting a given gene in

the same comparison of cocultures would mean that the knockout conferred a survival

advantage, thereby providing a means of immunological escape from the TCR_{EDST}-T cells by

diminishing presentation of the methylated epitope.

In the gene knockout screen, DAN-G cells with *SUPT6H* knockout (encoding the SPT6

protein) were the single most depleted knockout when comparing tumor cells cocultured with

the methylation-targeting TCR_{EDST}-T cell to the non-methylation-targeting TCR_2-T cell coculture

(**Figure 4b**). This suggests that SPT6 expression decreased methylation of this epitope. SPT6

is canonically known as a negative regulator of histone lysine methylation[35]. From our data,

SPT6 appears to have a non-canonical role as an inhibitor of lysine methylation on the

$KRAS_{G12V}$ epitope.

To further investigate the enhanced effectiveness of methylation-targeting TCR-T cells

specifically against SPT6-deficient DAN-G cells, we created a *SUPT6H* knockout DAN-G cell

line. Successful knockout was verified via long-read sequencing of the targeted genomic region

(**Supplementary Figure 5a**). Coculture of the *SUPT6H*-knockout DAN-G cells with $CD8^+$ T cells

transduced either with TCR_{EDST}, which targets both unmethylated and methylated $KRAS_{G12V}$

epitope, or with TCR_2, which only effectively targets the unmethylated $KRAS_{G12V}$ epitope,

affirmed that knockout of *SUPT6H* increased susceptibility to killing by T cells expressing either

TCR_{EDST} or $TCR_{A2UoM1-1}$ which also targets the methylated epitope (**Figure 4c**). Additionally, the

knockout did not increase killing by T cells expressing the methylation-unresponsive TCR_2.

Relative to the original DAN-G cells, *SUPT6H* knockout did not increase HLA-A2 expression

(**Supplementary Figure 5b**) and did not change KRAS expression levels (**Supplementary**

Figure 5c), indicating that the effect of *SUPT6H* knockout was not due to changes in total

epitope presentation levels but rather due to changes in the PTM composition of the epitope.

Data from The Cancer Genome Atlas (TCGA) show that loss-of-function mutations of

SUPT6H specifically within the $KRAS_{G12V}$ mutant cancer patient population lead to increased

survival compared to the *SUPT6H*-mutant patient population as a whole (**Figure 4d**). This

mutation-specific behavior may suggest that the methylated $KRAS_{G12V}$ epitope is more

spontaneously immunogenic in the context of a naturally occurring TCR repertoire, which would

be consistent with the more stabilized cell surface expression of the HLA-A2:KRAS$_{G12V}$ pMHC

complex when the epitope is methylated compared to unmethylated, as we described earlier

(**Supplementary Figure 1e**).

Discussion

Development of effective adoptive cell therapy using TCRs requires accurate

identification of the epitopes specifically presented by the targeted tumor, which can become

increasingly difficult in the context of PTMs. In this study, we have demonstrated, using two

approaches, the ability to isolate/generate TCRs that recognize and target PTMs that can alter

the immunogenicity of a tumor epitope. Deep learning-guided mutagenesis made it feasible to

refine a potentially promising TCR targeting an HLA-A2-presented KRAS$_{G12V}$ epitope into one

able to target both post-translationally modified and unmodified variants of the epitope, thereby

achieving improved killing of HLA-A2$^+$, KRAS$_{G12V}$ tumor cell types. We also generated TCRs

that can achieve similar activity by selectively expanding rare clonotypes reactive against

modified and unmodified KRAS$_{G12V}$ peptide variants from healthy donor-derived primary T cell

populations, thereby selecting for TCRs that have undergone the thymic central tolerance

process, meaning they are less likely to react against self-antigens and may offer a safer

therapeutic profile. For tumor cells that might present the unmodified KRAS$_{G12V}$ epitope together

with a mix of variants with different degrees of a lysine-5 methylation, our data suggests that

therapeutic TCRs designed or selected to target a wide range of these variants should prove

more effective at killing a larger subset of KRAS$_{G12V}$-expressing tumor cells, and preclude

immune evasion driven by altered or variable KRAS lysine methylation.

Computational prediction of TCR specificity against pMHC complexes has recently

experienced significant improvements[47], but purely computational approaches still require

experimental validation to determine whether TCR specificity is sufficiently high to be safe and

effective for antitumor therapy. Predicting the three-dimensional architecture between a pMHC

complex and TCR has the potential to provide insights for experimentally modifying and

improving the affinity/specificity of the TCR for the pMHC based on proposed contact points,

orientation of a peptide within the pMHC, and orientation of the TCR to that complex. However,

the structure of the TCR:pMHC complex has historically been difficult to accurately determine

computationally due to the still somewhat limited number of experimentally validated structures

of TCR:pMHC interactions from which to base new predictions, the bias of prediction models to

overfit based on existing common epitopes, and the lack of negative control data[48, 49]. Recent

innovations in computational protein structure prediction, including Alphafold, have allowed for

improved accuracy[48]. Nonetheless, Alphafold does not yet support the structural prediction of

changes resulting from PTMs due to the lack of available structural data for training that

includes PTMs[50]. Alternatively, Rosetta utilizes physics-based energy functions to optimize

protein structures, which better accommodates integration of PTMs into protein structure

prediction[51]. Thus, in this investigation, we combined these two pipelines into one prediction to

yield a more precise guide that accurately identified pMHC contact sites within the TCR CDR3

sequences that could subsequently be modified through saturation mutagenesis to specifically

enhance TCR association with post-translationally modified peptide within the pMHC complex.

This strategy allowed us to develop new TCRs that more effectively target HLA-A2$^+$ KRAS$_{G12V}^+$

tumors, while maintaining TCR specificity for mutant KRAS.

Developments in MS technology, including newer technology with greater sensitivity,

have allowed for its use to identify MHC class I-restricted T cell epitopes[30, 37]. We used such MS

to characterize the primary sequence of the HLA-A2-presented KRAS$_{G12V}$-containing epitope in

this study, analogous to previous studies that identified mutant KRAS epitopes presented by

other MHC I alleles[4, 8, 14]. These MS-based studies have been important for the development of

potentially therapeutic TCRs for mutant KRAS tumors. However, MS can remain unreliable for

detecting PTMs. The large size of the immunopeptidome presented by the multiple MHC I and II

alleles expressed by a tumor, the often-reduced level of MHC expression by tumor cells, the low

yield fraction of protein capture, and low expression of some proteins of interest can result in

low and inadequate sensitivity for detection of a specific epitope and any associated PTMs[30]. As

a result, MS-based workflows can benefit from maximizing protein expression and MHC capture

as we did with ARTEMIS in this study. Additionally, the introduction of chemical artifacts during

sample processing, such as oxidation[32], may confound whether a detected PTM is genuine or

not, and PTMs that are not suspected prior to analysis to be on a peptide may result in missing

the detection of that peptide[31]. To bypass these difficulties associated with epitope detection

and probe for a specific PTM on a presented epitope, TCRs with distinct PTM-peptide

recognition patterns may be used as an alternative to MS, as these TCRs will elicit T cell

activation in response to a target presenting distinct PTMs and can serve as a very sensitive

assay for the presence or absence of the PTM.

Based on the results of our efforts to generate TCRs targeting HLA-A2-presented

KRAS$_{G12V}$, TCR$_{EDST}$ and TCR$_{A2UoM1-1}$ appear to be two promising candidates for further

development as therapeutic TCRs. The former was generated by mutagenesis of an existing

TCR that naturally recognized an unmodified epitope derived from mutant KRAS, whereas the

latter was identified by selection from a healthy donor's TCR repertoire. The mutagenesis
approach allows for generation of TCRs that may not be present in normal repertoires and that
have increased affinity for the target. However, because these TCR sequences are mutated
from a TCR naturally existing in the repertoire, they have not been naturally screened during
development against the self-immunopeptidome, and pose an elevated risk of off-target activity
in a therapeutic setting. Thorough testing for TCR recognition of cross-reactive targets *in vitro*,
as we have initiated with alanine scans followed by identification and testing of potential non-
$KRAS_{G12V}$ targets in the proteome, should help to mitigate this problem. Additionally, such TCRs
may be useful as probes to identify the presence of particular PTMs on epitopes presented by
different tumors. TCR discovery from the repertoire of a healthy donor that has already been
centrally thymically tolerated is theoretically safer than TCRs generated by mutagenesis, but
can be limited if the desired TCR clonotypes are particularly rare within the repertoire or only of
low affinity. Overall, we show that either approach is feasible to generate potentially effective,
therapeutic TCRs able to target a modified and/or unmodified tumor antigen of interest.

Our findings with the histone methylation inhibitor SPT6 (*SUPT6H*) provide further
evidence that PTMs may be introduced by potentially non-canonical processes[27, 52], analogous
to the non-canonical methylation activity of SMYD3 that has been shown to impact other
components of the KRAS signaling pathway[27]. From our findings, *SUPT6H* knockout in tumor
cells enhances the efficacy of KRAS-targeting, PTM-reactive, TCR-T cells. Diminishing SPT6
activity may provide another opportunity to increase tumor immunogenicity, such as by
combining these TCR-T cells with a small molecule inhibitor of SPT6.

Our investigation into lysine methylation on $KRAS_{G12V}$ antigen revealed an intriguing role
of PTMs in changing recognition of tumor cells by cytotoxic T cells and provided a strategy to
mitigate the effects of PTMs to produce more consistently performing therapeutic TCRs. PTMs
of tumor-associated antigens have previously been implicated in altering immunogenicity of
tumor cells in several tumor types[53], and we build upon this by describing a strategy for the
development of TCRs to specifically target them. This strategy paves the way for developing
TCRs that specifically target PTMs, offering a novel avenue for personalized immunotherapy.
Moreover, we have identified TCRs that can target and kill tumor cells with mutated $KRAS_{G12V}$
and that are restricted to the most common human HLA allele, which therefore has significant
direct potential for clinical development. The widespread expression of the $KRAS_{G12V}$ mutation
across numerous different tumor types, including many solid tumors, means these TCRs may
allow development of ACT to combat these otherwise difficult-to-treat tumors.

The roles of PTMs found on tumor antigens outside of altering tumor cell immunogenicity
are likely broad. In the case of HLA-A2-presented KRAS$_{G12V}$, lysine-5 methylation in KRAS has
been speculated to affect interactions with relevant signaling proteins in the Ras family pathway
signaling[29]. Characterizing the mechanistic reasons for and consequences of PTMs on tumor
antigens should grant additional insight into possible molecular pathways that could be targeted
synergistically with, or independently of, immunotherapies targeting these antigens.
513

Materials and Methods

<u>Computational modeling of TCR:pMHC complexes.</u>
We used Alphafold to generate an initial structural model comprising a given TCR and
pMHC complex with unmodified KRAS epitope. Using the resulting model of the TCR:pMHC
complex not yet including any PTMs, we used PyRosetta[43] 2023.49 to introduce relevant PTMs
to the epitope, then reoptimize the binding energy of the TCR:pMHC complex with the PTMs
under consideration to create the final structural model. Subsequent visualizations were
generated using VMD[54]. For modeling only the pMHC complex, we used PyRosetta to take a
pMHC complex with known structure, consisting of HLA-A2 presenting a peptide with a
sequence similar to KRAS$_{G12V}$ 5-14, LLFGYPVAV (PDB: 1QSF[55]), mutated the peptide to
KRAS$_{G12V}$ 5-14 with or without PTMs, and reoptimized the binding energy of the resulting pMHC
complex.
526

<u>Generation and use of lentiviral constructs for TCRs.</u>
We created lentiviral transfer plasmids expressing our TCRs, using pRRLSIN as a basis.
We digested this plasmid with AscI and PstI restriction enzymes (New England Biolabs), with
recognition sites being introduced at appropriate locations beforehand via site-directed
mutagenesis, and gel-extracted the backbone. We then performed Gibson assembly using the
Gibson Assembly Kit (New England Biolabs) on the digested backbone and synthesized DNA
blocks (Codex DNA; Integrated DNA Technologies) encoding codon-optimized versions of our
selected TCR sequences. The Gibson assembly products were transformed into chemically
competent 5-alpha *E. coli* bacteria (New England Biolabs). The final construct thus contained a
single open reading frame for the TCR, composed of the TCRβ chain, P2A, and TCRα chain
and driven by the MSCV U3 promoter. To verify sequences of resulting clones, we used the full
plasmid sequencing service from Primordium Labs.
To generate lentivirus for TCR expression, we cultured 293T cells in RPMI 1640 media
(Thermo Fisher) with 5% FBS (Cytiva), 5% human serum (Bloodworks Northwest), 0.6% MEM

Non-Essential Amino Acids solution (Thermo Fisher), 4 mM L-glutamine (Gibco), and 100 U/ml

penicillin-streptomycin (Gibco). To generate lentivirus from the above constructs, we transfected

the 293T cells at 3 million cells/10 cm dish with Effectene transfection reagent (Qiagen)

following manufacturer guidelines, using 1.5 µg of an individual TCR construct, 0.5 µg of

pMD2.G (Addgene #12259), 1 µg of pMDLg/pRRE (Addgene #12251), and 1 µg of pRSV-Rev

(Addgene #12253). 1 d after transfection, we changed media, removing transfection reagent; 1

d following that, we filtered culture supernatant through 0.45 µm PVDF filters. The filtered

product was stored at –80ºC until used in TCR-T cell generation.

TCR-T cell generation.

We isolated CD8$^+$ T cells from primary human donor PBMCs using the EasySep Human

CD8$^+$ T cell Isolation Kit (STEMCELL Technologies) and activated them overnight using anti-

CD3/CD28 DynaBeads (Thermo Fisher). Afterward, we transduced these T cells with lentivirus

to express an engineered TCR targeting HLA-A2-presented KRAS$_{G12V}$ epitope, including

spinfection at 1,055 g for 90 minutes at 30°C. Cells were grown in "CTL media": RPMI 1640

media (Thermo Fisher) with 10% human serum (Bloodworks Northwest), 4 mM L-glutamine

(Gibco), 100 U/ml penicillin-streptomycin (Gibco), and 50 IU/ml IL-2 (Prometheus; obtained via

University of Washington Pharmacy, DOH license #DRRS.FX.60550751), with medium changes

every 2-3 d. Cells were sorted for peptide-MHC tetramer binding 6 d after transduction. These

cells were expanded via rapid expansion protocol for 8–14 d as described previously[56]. At the

end of expansion, the cells were used for functional assays.

MS of KRAS protein and ARTEMIS.

We performed ARTEMIS following previously described procedure[37] on DAN-G human

pancreatic adenocarcinoma and 293F cell lines. Briefly, we transduced these cell lines with a

construct expressing secreted, His-tagged HLA-A2 single chain dimer (SCD) with mCherry and

a construct for constitutive expression of the first 100 amino acids of human KRAS$_{G12V}$, based

on Lenti-mCherry-G12V-KRAS-IRES-Blast (Addgene #153336). Cells were sorted for SCD

expression via mCherry and further selected for the truncated KRAS$_{G12V}$ expression via

blasticidin at 10 µg/ml for 7 d. We expanded these cells and harvested culture supernatant for

two weeks. We filtered out debris from this supernatant using a 0.6 µm filter, captured HLA-A2

SCD-peptide complex with Ni-NTA agarose beads (Invitrogen) in imidazole equilibration buffer

(10mM imidazole, 300 mM NaCl, 50 mM NaH$_2$PO$_4$, pH 8), and eluted the bound peptides in a

denaturing buffer (5M guanidine-HCl, 250 mM NaCl, 50 mM NaH$_2$PO$_4$, 1 mM DTT, pH 8).

For whole KRAS protein, we performed cell lysis on HLA-A2$^+$ tumor cell lines to collect

protein as described previously[57]. From the resulting protein extract, we performed

immunoprecipitation for KRAS using MS-compatible magnetic immunoprecipitation kit (Pierce)

and anti-KRAS antibody (12063-1-AP; Thermo Fisher), following manufacturer guidelines. The

immunoprecipitated protein was run through SDS-PAGE, and a protein band at ~21 kDa size

was excised and used for liquid chromatography-MS.

Discovery of new TCRs via mutagenesis of an existing TCR targeting HLA-A2-presented

KRAS$_{G12V}$ 5-14.

To generate rationalized mutagenesis libraries of TCR$_2$ targeting the sequence within

CDR3α as selected from the structural prediction, we performed Gibson assembly to combine

synthesized saturation mutagenesis DNA blocks replacing the identified 'core' residues of

CDR3α with the remainder of the TCR$_2$ construct. We transduced these TCR libraries into

populations of Jurkat cells expressing CD8 and the Nur77-green fluorescent protein (GFP)

reporter gene, which specifically permits measurement of TCR signaling[46]. These cells were

grown in CTL media with the addition of KRAS$_{G12V}$ 5-14, with or without methylation of the

lysine-5 side chain, and presented by HLA-A2. After two weeks, we sorted these cells for

binding to tetramers of HLA-A2 with unmethylated or methylated KRAS$_{G12V}$ epitope and

amplified CDR3α sequences from sorted cells. Amplicons were sequenced by next-generation

sequencing on an Illumina MiSeq v2 flow cell with paired-end 150 bp reads.

To process sequencing results, we used custom code, trimming adapters and using

Bowtie2 to align reads to a reference sequence for the CDR3α region of TCR$_2$. We then

scanned for variations from the reference sequence and compared the frequencies of individual

sequence variants between sorted populations. The most differentially detected variants were

cloned into new lentiviral TCR constructs based on pRRLSIN, as described in the section

"Generation and use of lentiviral constructs for TCRs".

Discovery of new TCRs by stimulation of primary T cells by MHC I-presented peptides.

To enrich T cells expressing TCRs from the centrally tolerated human TCR repertoire

that could recognize HLA-A2-presented KRAS$_{G12V}$ epitope, we isolated primary CD8$^+$ T cells

from HLA-A2$^+$, healthy human donor PBMCs (**Supplementary Figure 4a**). From the remaining

PBMCs, we generated mature DCs through a differentiation cocktail consisting of GM-CSF and

IL-4 (Peprotech) for 1 d followed by TNFα, IL-1β, IL-6 (Peprotech), and prostaglandin E2 (MP

Biochemicals) for 1 d. The resulting DCs acted as APCs: they were pulsed for 4 h with synthetic

KRAS$_{G12V}$ peptides (Elim Biopharm), with or without PTMs incorporated, and stimulated the

CD8$^+$ T cells with the resulting peptide-presenting DCs. After this initial stimulation, we

continued to expand the T cells with a combination of IL-2 (Prometheus; obtained via University

of Washington Pharmacy, DOH license #DRRS.FX.60550751), IL-7, and IL-15 (Peprotech) with

medium replenishment every 2–3 d. These T cells were brought through multiple cycles of this

stimulation. In subsequent stimulations, peptide presentation was done by peptide-pulsing

PBMCs instead of mature DCs. At the end of the final round of stimulation, we stained T cells

with HLA-A2 tetramers presenting KRAS$_{G12V}$ 5-14 (with or without PTMs) and conjugated to a

fluorophore, with a different fluorophore per PTM/unmodified peptide. These cells were sorted

into separate populations based on tetramer binding patterns.

619   From the sorted T cells, we quantitated TCR clonotypes using the ImmunoSEQ assay

620   (Adaptive Biotech) and performed single cell RNA-seq using the Chromium Single Cell V(D)J

621   Enrichment Kit for T cells (10X Genomics) to identify paired TCRα/TCRβ sequences. Using

622   Cellranger 7.0.0 followed by Loupe VDJ Browser 4.0, for each population, we selected the most

623   highly enriched sequences for cloning of TCR lentiviral constructs.

625   <u>Coculture with engineered TCR-T cells and HLA-A2$^+$ tumor cells.</u>

626   HLA-A2$^+$ tumor cell lines were lentivirally transduced to express either Nuclight-Red or

Nuclight-Green nuclear-localizing fluorescent protein (Sartorius) and sorted for their expression.

For co-culture with TCR-T cells, we plated the tumor cell lines on 48-well plates at a density of

10,000 cells/well one day before adding T cells to allow for tumor cell adherence. T cells,

generated as described above, were added to these wells at a CD8$^+$ T cell:tumor cell ratio of 4:1

unless stated otherwise, with an initial dose of 12.5 IU/ml IL-2. Cocultures were imaged every

two hours at 10X magnification using an Incucyte S3 machine (Sartorius).

In vitro peptide assays.

635   To determine functional avidity of our engineered TCRs against peptides, we used either

Jurkat CD8$^+$ Nur77-GFP reporter cells or primary human HLA-A2$^+$ T cells and transduced them

with one of our TCRs. Cells were incubated with *TAP*-deficient T2 cells and synthetic peptide at

set concentrations, in the presence of GolgiStop and GolgiPlug protein transport inhibitors (BD

Biosciences) at 1:1000 dilution each for 16 h at 37°C. Cells were incubated by themselves in the

presence of the same protein transport inhibitors in addition to synthetic peptides at set

concentrations for 4 h at 37°C. At the end of incubation, live CD8$^+$ cells were measured for

expression of IFNγ (primary CD8$^+$ T cells) or Nur77-GFP (Jurkat CD8$^+$ Nur77-GFP reporter

cells) by flow cytometry. CD8+ T cells were stained with eFluor 780 Fixable Viability Dye

(Invitrogen) followed by CD8:BV421 (Biolegend), fixed and permeabilized using the BD

Cytofix/Cytoperm Kit (BD Biosciences), and stained with IFNγ:PE-Cy7 antibody (Biolegend).

Nur77-GFP Jurkat cells were stained with CD8:BV421 (Biolegend) followed by propidium iodide.

Flow cytometry analysis was performed using FlowJo v9 (BD Biosciences).

To measure the stability of peptide presentation on HLA-A2, we incubated T2 cells with

individual peptides of interest for 16 h, then measured live cells for surface expression of HLA-

A2 by flow cytometry, staining with HLA-A2:FITC antibody (Biolegend) and propidium iodide.

<u>CRISPR knockout screen and validation of genes involved in methylation processes.</u>

Genes were selected using the PANTHER database[58] which met the criteria of

containing "methyl" in the gene ontology description. The total number of genes selected was

605. A total of 1,813 sgRNA sequences for the genes of interest were collected from existing

databases[59, 60, 61]. 30 control gRNA sequences were added to the library for a total of 1,843

gRNAs (**Supplementary Table 1**). The list of sgRNAs was ordered as a custom oligonucleotide

pool from Integrated DNA Technologies.

For the creation of the CRISPR knockout library plasmid, we performed cloning following

Canver *et al.*, 2018[62]. An oligo pool was cloned into lentiCRISPRv2 (Addgene #52961) which

was digested using Esp3I-HF (New England Biolabs). (2.5 µg) was transfected into HEK293T

cells using X-tremeGENE™ 9 (Roche) with psPAX2 (1.625 µg) and pCMV-VSV-G (0.875 µg)

plasmids (Addgene #12260, #8454). Viral supernatant was collected and filtered using a 0.45

µm PVDF filter 48 hours after transfection. The plasmid-containing viral supernatant was then

transduced into the DAN-G cell line. To select for the cells which were successfully transduced,

the cells underwent puromycin selection for 7 days at 1 µg/ml concentration with medium

changes every 2-3 days. The library-containing DAN-G cells were co-cultured with the KRAS-

targeting TCR-T cells for 7 days. Following co-culture, the DAN-G cells were rinsed with PBS

and pelleted for genomic DNA extraction using QIAGEN DNeasy Blood & Tissue kit. The

sgRNA region of the genomic DNA was PCR amplified following the 'sgRNA/shRNA/ORF PCR

for Illumina Sequencing' protocol via the Broad Institute Genetic Perturbation Platform and

assessed for fragment length distribution on TapeStation D1000 (Agilent). Upon verification of

fragment length, the samples were pooled and sequenced on an Illumina MiSeq machine.

Raw NGS reads were adapter-trimmed with Cutadapt v2.9, followed by alignment to the

reference library with Bowtie v1.3. Comparisons between different conditions were performed

using the MAGeCK pipeline, using RRA and MLE algorithms for calculating statistical

significance of results. MAGeCK also predicts gene essentiality by evaluating the ranked list of

sgRNAs[63].

To verify hits from our CRISPR library screen, we transduced DAN-G cells with lentivirus

inducing single knockouts and puromycin resistance. After 7 days of puromycin selection at 1

µg/ml, the DAN-G single knockout cells was co-cultured with TCR-T cells at a 4:1 T cell:tumor

cell ratio and imaged every 4 hours at 10X magnification using an Incucyte S3 machine

(Sartorius).

To verify knockout, genomic DNA was extracted from single knockout DAN-G cells, and

the gRNA region was PCR amplified using Q5 High-Fidelity DNA Polymerase (New England

Biolabs), yielding amplicon lengths of between 500-800 base pairs. The PCR-amplified samples

were sequenced using the "Premium PCR Sequencing" service from Primordium Labs. The

resulting sequence files were aligned with the reference genome using BWA-MEM to screen for

the common mutation sites to confirm the knockout. To verify the methylation status of HLA-A2-

presented KRAS$_{G12V}$ epitope in the *SUPT6H* knockout DAN-G cells and non-knocked-out DAN-

G cells, we expanded these cells and extracted HLA-A2-presented peptide using the ARTEMIS

protocol as previously described.

To compare HLA-A2 expression of *SUPT6H* knockout vs. non-knocked-out cells, the

cells were stained with HLA-A2:PE antibody (Biolegend) and propidium iodide and analyzed via

flow cytometry. To compare KRAS expression, protein lysate was extracted from tumor cell

lines as described above and used as input for Western blots with anti-KRAS antibody (12063-

1-AP; Thermo Fisher).

Code availability

700   Code for structural modeling and TCR sequence data processing from TCR

701 mutagenesis are available at https://github.com/jihoonlee0/A2_KRAS-G12V.

Acknowledgements

We would like to thank the Fred Hutch Genomics, Bioinformatics,

Proteomics/Metabolomics, Flow Cytometry, and Cellular Imaging Shared Resources for their

respective roles in helping to run key experiments. These resources are supported by the Fred

Hutch/University of Washington/Seattle Children's Cancer Consortium Cancer Center Support

Grant (NIH P30CA015704). We also thank members of the Chapuis Lab and the Strong Lab at

Fred Hutch for their help setting up ARTEMIS for use in this study.

This study received financial support from donations to Fred Hutchinson Cancer Center
for "TCR Therapy for Patients with Pancreatic Cancers" (Fred Hutch Reference Number
SRA210802) and Affini-T Therapeutics, Inc. (Fred Hutch Reference Number 20012884). The
mass spectrometry instrumentation used in this research was funded by a grant from the NIH
Office of Research Infrastructure Programs (S10OD030225).
715

716   # References

717   1.   Schmitt TM, Aggen DH, Ishida-Tsubota K, Ochsenreither S, Kranz DM, Greenberg PD.
718   Generation of higher affinity T cell receptors by antigen-driven differentiation of
719   progenitor T cells in vitro. *Nat Biotechnol* **35**, 1188-1195 (2017).
720
721   2.   Chapuis AG*, et al.* Transferred WT1-reactive CD8+ T cells can mediate antileukemic
722   activity and persist in post-transplant patients. *Sci Transl Med* **5**, 174ra127 (2013).
723
724   3.   Chapuis AG*, et al.* T cell receptor gene therapy targeting WT1 prevents acute myeloid
725   leukemia relapse post-transplant. *Nat Med* **25**, 1064-1072 (2019).
726
727   4.   Lu D*, et al.* KRAS G12V neoantigen specific T cell receptor for adoptive T cell therapy
728   against tumors. *Nat Commun* **14**, 6389 (2023).
729
730   5.   Doran SL*, et al.* T-Cell Receptor Gene Therapy for Human Papillomavirus-Associated
731   Epithelial Cancers: A First-in-Human, Phase I/II Study. *J Clin Oncol* **37**, 2759-2768
732   (2019).
733
734   6.   Hobbs GA, Der CJ, Rossman KL. RAS isoforms and mutations in cancer at a glance. *J
735   Cell Sci* **129**, 1287-1292 (2016).
736
737   7.   Cook JH, Melloni GEM, Gulhan DC, Park PJ, Haigis KM. The origins and genetic
738   interactions of KRAS mutations are allele- and tissue-specific. *Nat Commun* **12**, 1808
739   (2021).
740
741   8.   Rive CM*, et al.* Recombinant T cell receptors specific for HLA-A*02:01-restricted
742   neoepitopes containing KRAS codon 12 hotspot mutations. *bioRxiv*,
743   2020.2006.2015.149021 (2020).
744
745   9.   Arrieta-Bolanos E, Hernandez-Zaragoza DI, Barquera R. An HLA map of the world: A
746   comparison of HLA frequencies in 200 worldwide populations reveals diverse patterns
747   for class I and class II. *Front Genet* **14**, 866407 (2023).
748

749 10. Chin L*, et al.* Essential role for oncogenic Ras in tumour maintenance. *Nature* **400**, 468-
750   472 (1999).

751
752 11. Malcolm J. W. Sim K-iH, Zachary Stotz, Zhiya Yu, Jinghua Lu, Paul Brennan, Max
753   Quastel, Geraldine M. Gillespie, Eric O. Long, James C. Yang, Peter D. Sun.
754   Identification and Structural Characterization of a mutant KRAS-G12V specific TCR
755   restricted by HLA-A3.) (2024).

756
757 12. Zhang M*, et al.* Identification and affinity enhancement of T-cell receptor targeting a
758   KRAS(G12V) cancer neoantigen. *Commun Biol* **7**, 512 (2024).

759
760 13. Leidner R*, et al.* Neoantigen T-Cell Receptor Gene Therapy in Pancreatic Cancer. *N
761   Engl J Med* **386**, 2112-2119 (2022).

762
763 14. Poole A*, et al.* Therapeutic high affinity T cell receptor targeting a KRAS(G12D) cancer
764   neoantigen. *Nat Commun* **13**, 5333 (2022).

765
766 15. Gonzalez-Galarza FF, Christmas S, Middleton D, Jones AR. Allele frequency net: a
767   database and online repository for immune gene frequencies in worldwide populations.
768   *Nucleic Acids Res* **39**, D913-919 (2011).

769
770 16. Ellis JM, Henson V, Slack R, Ng J, Hartzman RJ, Katovich Hurley C. Frequencies of
771   HLA-A2 alleles in five U.S. population groups. Predominance Of A*02011 and
772   identification of HLA-A*0231. *Hum Immunol* **61**, 334-340 (2000).

773
774 17. Bear AS*, et al.* Biochemical and functional characterization of mutant KRAS epitopes
775   validates this oncoprotein for immunological targeting. *Nat Commun* **12**, 4365 (2021).

776
777 18. Choi J*, et al.* Systematic discovery and validation of T cell targets directed against
778   oncogenic KRAS mutations. *Cell Rep Methods* **1**, 100084 (2021).

779
780 19. Mishto M*, et al.* An in silico-in vitro Pipeline Identifying an HLA-A(*)02:01(+) KRAS
781   G12V(+) Spliced Epitope Candidate for a Broad Tumor-Immune Response in Cancer
782   Patients. *Front Immunol* **10**, 2572 (2019).

783
784 20. Lahman MC*, et al.* Targeting an alternate Wilms' tumor antigen 1 peptide bypasses
785   immunoproteasome dependency. *Sci Transl Med* **14**, eabg8070 (2022).

786
787 21. Johnsen AK, Templeton DJ, Sy M, Harding CV. Deficiency of transporter for antigen
788   presentation (TAP) in tumor cells allows evasion of immune surveillance and increases
789   tumorigenesis. *J Immunol* **163**, 4224-4231 (1999).

791   22.   Maggs L, Sadagopan A, Moghaddam AS, Ferrone S. HLA class I antigen processing
792   machinery defects in antitumor immunity and immunotherapy. *Trends Cancer* **7**, 1089-
793   1101 (2021).

794
795   23.   Hardy LL, Wick DA, Webb JR. Conversion of tyrosine to the inflammation-associated
796   analog 3'-nitrotyrosine at either TCR- or MHC-contact positions can profoundly affect
797   recognition of the MHC class I-restricted epitope of lymphocytic choriomeningitis virus
798   glycoprotein 33 by CD8 T cells. *J Immunol* **180**, 5956-5962 (2008).

799
800   24.   Corbiere V*, et al.* Natural T Cell Epitope Containing Methyl Lysines on Mycobacterial
801   Heparin-Binding Hemagglutinin. *J Immunol* **204**, 1715-1723 (2020).

802
803   25.   Petersen J*, et al.* Phosphorylated self-peptides alter human leukocyte antigen class I-
804   restricted antigen presentation and generate tumor-specific epitopes. *Proc Natl Acad Sci*
805   *U S A* **106**, 2776-2781 (2009).

806
807   26.   Raposo B*, et al.* T cells specific for post-translational modifications escape intrathymic
808   tolerance induction. *Nat Commun* **9**, 353 (2018).

809
810   27.   Mazur PK*, et al.* SMYD3 links lysine methylation of MAP3K2 to Ras-driven cancer.
811   *Nature* **510**, 283-287 (2014).

812
813   28.   Chiang CY*, et al.* Methylation of KRAS by SETD7 promotes KRAS degradation in non-
814   small cell lung cancer. *Cell Rep* **42**, 113003 (2023).

815
816   29.   Yoshino H*, et al.* Identification of lysine methylation in the core GTPase domain by
817   GoMADScan. *PLoS One* **14**, e0219436 (2019).

818
819   30.   Caron E, Kowalewski DJ, Chiek Koh C, Sturm T, Schuster H, Aebersold R. Analysis of
820   Major Histocompatibility Complex (MHC) Immunopeptidomes Using Mass Spectrometry.
821   *Mol Cell Proteomics* **14**, 3105-3117 (2015).

822
823   31.   Hughes C, Ma B, Lajoie GA. De novo sequencing methods in proteomics. *Methods Mol*
824   *Biol* **604**, 105-121 (2010).

825
826   32.   Perdivara I, Deterding LJ, Przybylski M, Tomer KB. Mass spectrometric identification of
827   oxidative modifications of tryptophan residues in proteins: chemical artifact or post-
828   translational modification? *J Am Soc Mass Spectrom* **21**, 1114-1117 (2010).

829
830   33.   Peters B, Nielsen M, Sette A. T Cell Epitope Predictions. *Annu Rev Immunol* **38**, 123-
831   145 (2020).

833   34.   Ahearn I, Zhou M, Philips MR. Posttranslational Modifications of RAS Proteins. *Cold*
834   *Spring Harb Perspect Med* **8**, (2018).

835
836   35.   Miller CLW, Warner JL, Winston F. Insights into Spt6: a histone chaperone that functions
837   in transcription, DNA replication, and genome stability. *Trends Genet* **39**, 858-872
838   (2023).

839
840   36.   Reynisson B, Alvarez B, Paul S, Peters B, Nielsen M. NetMHCpan-4.1 and
841   NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by concurrent
842   motif deconvolution and integration of MS MHC eluted ligand data. *Nucleic Acids Res*
843   **48**, W449-W454 (2020).

844
845   37.   Finton KAK*, et al.* ARTEMIS: A Novel Mass-Spec Platform for HLA-Restricted Self and
846   Disease-Associated Peptide Discovery. *Front Immunol* **12**, 658372 (2021).

847
848   38.   Ooi A, Wong A, Esau L, Lemtiri-Chlieh F, Gehring C. A Guide to Transient Expression of
849   Membrane Proteins in HEK-293 Cells for Functional Characterization. *Front Physiol* **7**,
850   300 (2016).

851
852   39.   Tew BY*, et al.* Genome-wide DNA methylation analysis of KRAS mutant cell lines. *Sci*
853   *Rep* **10**, 10149 (2020).

854
855   40.   Jin H*, et al.* Epigenetic silencing of a Ca(2+)-regulated Ras GTPase-activating protein
856   RASAL defines a new mechanism of Ras activation in human cancers. *Proc Natl Acad*
857   *Sci U S A* **104**, 12353-12358 (2007).

858
859   41.   Chen M, Zhang M, Zhai L, Hu H, Liu P, Tan M. Tryptic Peptides Bearing C-Terminal
860   Dimethyllysine Need to Be Considered during the Analysis of Lysine Dimethylation in
861   Proteomic Study. *J Proteome Res* **16**, 3460-3469 (2017).

862
863   42.   Nijman HW*, et al.* Identification of peptide sequences that potentially trigger HLA-A2.1-
864   restricted cytotoxic T lymphocytes. *Eur J Immunol* **23**, 1215-1219 (1993).

865
866   43.   Chaudhury S, Lyskov S, Gray JJ. PyRosetta: a script-based interface for implementing
867   molecular modeling algorithms using Rosetta. *Bioinformatics* **26**, 689-691 (2010).

868
869   44.   Karapetyan AR*, et al.* TCR Fingerprinting and Off-Target Peptide Identification. *Front*
870   *Immunol* **10**, 2501 (2019).

871
872   45.   Jumper J*, et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
873   583-589 (2021).

875 46. Ashouri JF, Weiss A. Endogenous Nur77 Is a Specific Indicator of Antigen Receptor
876   Signaling in Human T and B Cells. *J Immunol* **198**, 657-668 (2017).

877
878 47. Hudson D, Fernandes RA, Basham M, Ogg G, Koohy H. Can we predict T cell specificity
879   with digital biology and machine learning? *Nat Rev Immunol* **23**, 511-521 (2023).

880
881 48. Bradley P. Structure-based prediction of T cell receptor:peptide-MHC interactions. *Elife*
882   **12**, (2023).

883
884 49. Moris P*, et al.* Current challenges for unseen-epitope TCR interaction prediction and a
885   new perspective derived from image classification. *Brief Bioinform* **22**, (2021).

886
887 50. Bagdonas H, Fogarty CA, Fadda E, Agirre J. The case for post-predictional
888   modifications in the AlphaFold Protein Structure Database. *Nat Struct Mol Biol* **28**, 869-
889   870 (2021).

890
891 51. Renfrew PD, Choi EJ, Bonneau R, Kuhlman B. Incorporation of noncanonical amino
892   acids into Rosetta and use in computational protein-peptide interface design. *PLoS One*
893   **7**, e32637 (2012).

894
895 52. Liu J, Wang Q, Kang Y, Xu S, Pang D. Unconventional protein post-translational
896   modifications: the helmsmen in breast cancer. *Cell Biosci* **12**, 22 (2022).

897
898 53. Srivastava AK, Guadagnin G, Cappello P, Novelli F. Post-Translational Modifications in
899   Tumor-Associated Antigens as a Platform for Novel Immuno-Oncology Therapies.
900   *Cancers (Basel)* **15**, (2022).

901
902 54. Humphrey W, Dalke A, Schulten K. VMD: visual molecular dynamics. *J Mol Graph* **14**,
903   33-38, 27-38 (1996).

904
905 55. Ding YH, Baker BM, Garboczi DN, Biddison WE, Wiley DC. Four A6-TCR/peptide/HLA-
906   A2 structures that generate very different T cell signals are nearly identical. *Immunity* **11**,
907   45-56 (1999).

908
909 56. Riddell SR, Greenberg PD. High efficiency transduction of T lymphocytes using rapid
910   expansion methods (" REM").). Google Patents (2000).

911
912 57. Adams LM, DeHart CJ, Kelleher NL. Precise Characterization of KRAS4B Proteoforms
913   by Combining Immunoprecipitation with Top-Down Mass Spectrometry. *Methods Mol*
914   *Biol* **2262**, 47-64 (2021).

916   58.   Mi H*, et al.* PANTHER version 16: a revised family classification, tree-based
917   classification tool, enhancer regions and extensive API. *Nucleic Acids Res* **49**, D394-
918   D403 (2021).

919
920   59.   Sanjana NE, Shalem O, Zhang F. Improved vectors and genome-wide libraries for
921   CRISPR screening. *Nat Methods* **11**, 783-784 (2014).

922
923   60.   Doench JG*, et al.* Optimized sgRNA design to maximize activity and minimize off-target
924   effects of CRISPR-Cas9. *Nat Biotechnol* **34**, 184-191 (2016).

925
926   61.   Concordet JP, Haeussler M. CRISPOR: intuitive guide selection for CRISPR/Cas9
927   genome editing experiments and screens. *Nucleic Acids Res* **46**, W242-W245 (2018).

928
929   62.   Canver MC*, et al.* Integrated design, execution, and analysis of arrayed and pooled
930   CRISPR genome-editing experiments. *Nat Protoc* **13**, 946-986 (2018).

931
932   63.   Li W*, et al.* MAGeCK enables robust identification of essential genes from genome-scale
933   CRISPR/Cas9 knockout screens. *Genome Biol* **15**, 554 (2014).

Figures and Tables

Figure 1. Posttranslational modification on the HLA-A2-presented $KRAS_{G12V}$ epitope alters recognition of the antigen by MHC class I-restricted TCRs and subsequent tumor cell killing. (a) $Log_{10}(EC_{50}$ (M)) comparison of recognition of different concentrations of a synthetic $KRAS_{G12V}$ 5-14 peptide (KLVVVGAVGV) presented by HLA-A2 by different MHC class I-restricted TCRs specific for the epitope. Lower values indicate higher functional avidity. (b) Killing of live CFPAC1 and DAN-G HLA-A2$^+$ $KRAS_{G12V}$$^+$ pancreatic adenocarcinoma cells cocultured with CD8$^+$ T cells expressing the TCRs shown in (a) using a 4:1 T cell:tumor cell ratio. Cocultures were recorded by Incucyte imaging. Tumor cells expressed nuclear fluorescent protein; NR: red fluorescent protein; NG: green fluorescent protein. (c) ARTEMIS MS data of peptides eluted from HLA-A2. Data is from the 293F cell line, as commonly used for proteomics assays requiring high protein expression. All cell lines were transduced with an HLA-A2 single chain secreted dimer as well as a $KRAS_{G12V}$ constitutive expression construct to increase presentation and the likelihood of detecting a presented peptide by MS, of $KRAS_{G12V}$-related epitopes. The first 100 amino acids of $KRAS_{G12V}$ are shown, with the G12V mutation highlighted in yellow. Epitopes predicted by NetMHCPan 4.1 are shown in purple; epitopes detected from ARTEMIS are shown in blue. ARTEMIS detection of $KRAS_{G12V}$ 5-14 is outlined in red. (d) Heatmap of – $log_{10}(EC_{50}$ (µg/ml)) of CD8$^+$ T cells expressing TCR_2 or TCR_{19} against candidate $KRAS_{G12V}$ epitopes, with and without methylation of the lysine-5 side chain of the epitope, presented by HLA-A2. Higher values indicate higher functional avidity. EC_{50} values were calculated from T cell exposure to peptide concentrations ranging from 1 µg/ml to 10^{-5} µg/ml. Gray squares indicate EC_{50} calculations that lacked a stable fit, e.g., due to lack of response to peptide even at high doses. (e) Rosetta structural modeling of HLA-A2 presenting $KRAS_{G12V}$ 5-14 and its methylated variants. Red circles indicate the amine group of the lysine-5 side chain; arrows indicate its movement with different methylation states. A reference point is drawn at the same position in each image to help illustrate positional changes. (f) Alanine scan of the $KRAS_{G12V}$ epitope and resulting response by CD8$^+$ T cells expressing TCR_2 and TCR_{19}. The epitope with each individual residue substituted with alanine (or, for alanine-11, substituted with either glycine or threonine) was presented by antigen-presenting cells to stimulate primary human CD8$^+$ T cells expressing a TCR, and subsequent IFNγ expression was measured. Higher IFNγ induction indicates greater tolerance for amino acid substitution at that position. X-axis numbers indicate the residue number on the KRAS protein, and letters indicate the amino acid substitution. A: alanine; G: glycine; T: threonine. Last column is unmodified $KRAS_{G12V}$ 5-14.

Figure 2. A computationally guided, rationalized mutagenesis library to develop MHC class I-
restricted TCRs that recognize methylated epitopes of HLA-A2-presented KRAS$_{G12V}$ and
achieve enhanced targeting of HLA-A2$^+$ KRAS$_{G12V}$$^+$ tumor cell lines. (a) Structural model, as
predicted by a combined Alphafold-Rosetta pipeline, of a 'core' CDR3α sequence (sequence
EDNT) of an existing TCR recognizing KRAS$_{G12V}$ 5-14 and its relation to the lysine-5 side chain
(circled in red) of the epitope as presented by HLA-A2. (b) Generation of TCR mutagenesis
libraries by mutagenesis of the core CDR3α sequence followed by transduction of the TCRs into
Nur77-GFP reporter Jurkat cells and subsequent clonal expansions. Tetramer binding patterns
of HLA-A2-KRAS$_{G12V}$ 5-14 peptide with unmodified or methylated lysine-5 side chain at the end
of cell stimulation are shown. Each circle indicates a population with a specific pattern of
unmethylated vs. methylated peptide tetramer binding that was sorted and sequenced. (c)
Examples of mutated CDR3α 'core' sequences resulting from sorted clones of the mutagenesis
library that exhibit the noted different patterns of PTM peptide recognition. (d) Heatmap of –
log$_{10}$(EC$_{50}$ (μg/ml)) of TCRs derived from the mutagenesis library and the original TCR$_2$
responding to different degrees of lysine-5 methylation of KRAS$_{G12V}$ 5-14 presented by HLA-A2,
as calculated from dose response assays. Gray squares indicate EC$_{50}$ values that lacked a
stable fit, e.g., due to limited response to peptide even at high doses. TCR$_{4UM}$ is the 4th TCR
clonotype identified from an initial sequencing dataset from sorted Jurkat cells binding to both
unmethylated ("U") and methylated ("M") tetramers; other TCRs are labeled by changes in the
'core' CDR3α sequence (e.g., EDST). (e) Killing of HLA-A2$^+$ KRAS$_{G12V}$$^+$ CFPAC1 and DAN-G
pancreatic adenocarcinoma cells by CD8$^+$ T cells expressing selected mutagenized TCRs.
Cocultures of T cells and tumor cells were recorded by Incucyte imaging. Tumor cells expressed
nuclear fluorescent protein. NR: red fluorescent protein; NG: green fluorescent protein.
993
Figure 3. Stimulation of primary human T cells with unmethylated and methylated HLA-A2-
presented KRAS$_{G12V}$ to generate MHC class I-restricted TCRs from a normal repertoire that
recognize methylated epitopes. (a) Heatmap of –log$_{10}$(EC$_{50}$ (μg/ml)) of selected TCRs
responding to the unmethylated and methylated epitopes presented by HLA-A2, as calculated
from dose response assays. Higher values indicate higher functional avidity. Gray squares
indicate EC$_{50}$ values that lacked a stable fit, e.g., due to limited response to peptide even at high
doses. (b) Killing of HLA-A2$^+$ KRAS$_{G12V}$$^+$ CFPAC1 and DAN-G pancreatic adenocarcinoma cells
by CD8$^+$ T cells expressing selected TCRs. Cocultures of fluorescently labeled tumor cells and

TCR-T cells were recorded by Incucyte imaging. Tumor cells expressed nuclear fluorescent

protein; NR: red fluorescent protein; NG: green fluorescent protein. (c) Alphafold-Rosetta

visualizations of TCR:pMHC complexes for $TCR_{A2UoM1-1}$, isolated from expansion of responding

primary human $CD8^+$ T cells from normal repertoires. Structures were generated for

unmethylated and mono/di/trimethylated versions of the epitope, presented by HLA-A2. The four

CDR3α amino acids in closest proximity to the lysine-5 side chain of $KRAS_{G12V}$ are shown with

individual molecular bonds.

1009

Figure 4. A gene knockout screen for methylation-related genes identifies SPT6 (*SUPT6H*) as a

participant in the demethylation of the HLA-A2-presented $KRAS_{G12V}$ epitope. (a) Overall

workflow: DAN-G pancreatic adenocarcinoma cells were transduced with a CRISPR knockout

library targeting methylation-related genes, and then cocultured with $CD8^+$ T cells expressing

either TCR_2 or mutagenized TCR_{EDST}, which respectively have weak or strong recognition of

methylated $KRAS_{G12V}$ epitopes. Tumor cells surviving at the end of coculture were sequenced to

identify gRNA enrichment or depletion associated with immune evasion. (b) Volcano plot

summarizing $\log_2$(fold change) of gRNA read frequencies targeting individual genes recovered

after coculture of DAN-G cells with T cells expressing TCR_{EDST} versus TCR_2. The threshold of

significance was set at $\log_{10}(p) < 1.2$ ($p < 0.05$) and is denoted by the horizontal dashed line.

Mean counts of all gRNAs per gene were used to calculate fold change values. The data point

for gRNAs targeting *SUPT6H* is specifically labeled. The knockouts depleted or enriched after

cocultures using TCR_{EDST} versus TCR_2 are colored blue and red, respectively. (c) Killing of

DAN-G cells with or without *SUPT6H* knockout in coculture with selected TCR-T cells

(untransduced or expressing TCR_2, TCR_{EDST}, or $TCR_{A2UoM1-1}$). Cocultures of T cells and tumor

cells were recorded by Incucyte imaging. Tumor cells expressed green nuclear fluorescent

protein ("NG"). (d) Kaplan-Meier survival plots for individuals who have tumors with KRAS

mutations, using public TCGA data. Left: only tumors with the $KRAS_{G12V}$ mutant, with or without

concurrent *SUPT6H* mutation. Right: tumors with any KRAS mutant, with or without a

concurrent *SUPT6H* mutation.

1030

Supplementary Figure 1. HLA-A2-presented $KRAS_{G12V}$ epitope includes a lysine residue that

can be stably presented with potential side chain methylation. (a) NetMHCPan 4.1 summary of

the top 10 peptide sequences that are most likely to be presented by HLA-A2 from the first 30

amino acids of $KRAS_{G12V}$. Peptides containing the G12V mutation (including mutations in red

text) are in black text; others are in gray. (b) $\log_{10}(EC_{50}$ (M)) comparison of recognition of

synthetic KRAS$_{G12V}$ 5-14 peptide (KLVVVGAVGV) presented by HLA-A2 by an initially

generated set of MHC class I-restricted TCRs isolated to be specific for this epitope. Lower

values indicate higher functional avidity. (c) % IFNγ$^+$ T cells, transduced with TCR$_2$, after

exposure to B-LCL cell lines expressing different Class I HLA alleles found on CFPAC1 cells

without KRAS$_{G12V}$ 5-14 peptide loaded. Cell lines are color coded by allele. (d) MS results of

RAS family peptide fragments detected after tryptic digestion of immunoprecipitated whole

KRAS protein from cell lysate of DAN-G and CFPAC1 pancreatic adenocarcinoma cell lines.

Lysine side chain trimethylation is indicated in red with an asterisk. (e) Reconstitution of properly

folded HLA-A2 surface expression on T2 cells after exogeneous loading by wild type or G12V/D

mutant KRAS 5-14 peptides with or without lysine-5 side chain methylations. (f) Additional views

of structural models of HLA-A2-presented KRAS$_{G12V}$ 5-14 peptide including possible lysine-5

side chain methylation, supplementing Figure 1e by highlighting the protrusion of the lysine-5

side chain (circled in red) out of the HLA-A2 groove. (g) Overview of structural prediction of a

TCR-pMHC complex using a combined Alphafold-Rosetta pipeline. Alphafold allows for deep

learning-based prediction of proteins and protein complexes, which we applied to predict TCR-

pMHC complex structure. However, this does not yet contain PTMs, which Alphafold cannot

incorporate reliably. Thus, we used the resulting structure as input into Rosetta, which used

physics-based energy functions to introduce PTMs and reoptimize the structure to

accommodate them.

1055

Supplementary Figure 2. TCRs developed from a mutagenesis library to recognize post-

translationally modified epitopes of HLA-A2-presented KRAS$_{G12V}$ are G12V mutation-specific.

(a) Dose response curves of CD8$^+$ T cells expressing selected TCRs generated from the HLA-

A2 KRAS$_{G12V}$ TCR mutagenesis library after exposure to wild type or G12D KRAS 5-14 peptides

presented by HLA-A2. (b) Failure of CD8$^+$ T cell killing and progressive growth of human HLA-

A2$^+$ tumor cell lines that express wild type KRAS (HeLa cervical carcinoma cell line; left) or

KRAS$_{G12D}$ (Panc1 pancreatic adenocarcinoma; right) and no KRAS$_{G12V}$. Cocultures of T cells

and tumor cells were recorded by Incucyte imaging. Tumor cells expressed nuclear fluorescent

protein. NR: red fluorescent protein; NG: green fluorescent protein. (c) Alanine scan of

KRAS$_{G12V}$ 5-14 epitope tested against TCRs selected from the mutagenesis library. Each cell

indicates the resulting expression of Nur77-GFP upon replacement of a KRAS$_{G12V}$ residue with

alanine (or in the case of alanine-11, replacement with either glycine or threonine) and with or

without lysine-5 side chain dimethylation. (d) IFNγ secretion by CD8$^+$ T cells expressing

mutagenized TCRs after exposure to synthetic peptides from the human proteome fitting the K-

x-x-V-V-x-A-x-x-x tolerance pattern identified from the alanine scan shown in (c), with a

concentration of 0.1 µg/ml for each peptide. (e) Dose response curves of CD8$^+$ T cells

expressing mutagenized TCRs after exposure to the potential human off-target peptides

demonstrated above to generate the three largest responses at 0.1 µg/ml: CFA61$_{23\text{-}32/607\text{-}616/667\text{-}}$

$_{676}$, RSLBB$_{35\text{-}44}$, and TRXR1$_{342\text{-}351}$, compared to the KRAS$_{G12V}$ epitope.

1075

Supplementary Figure 3. TCR-pMHC structural models by Alphafold-Rosetta show CDR3α

core interactions with N-terminal lysine side chain of HLA-A2-presented KRAS$_{G12V}$ epitope.

Structural predictions of unmethylated and methylated variants of KRAS$_{G12V}$ 5-14, as well as

unmethylated wild type or G12D peptide, presented by HLA-A2 to interface with (a) TCR$_2$ and

(b) mutagenized TCR$_{EDST}$ TCRs. The CDR3α 'core' sequence is shown as individual amino

acids within the TCRs. In (b), the TCR$_2$-pMHC interfaces from (a) are overlaid with transparency

to facilitate visual comparison. With TCR$_{EDST}$, residues in the core sequence of CDR3α and the

lysine-5 side chain of each variant of KRAS$_{G12V}$ epitope appear closer to each other than with

TCR$_2$.

1085

Supplementary Figure 4. TCRs identified from primary human T cells specifically target

KRAS$_{G12V}$ epitope presented by HLA-A2. (a) Experimental workflow to enrich and characterize T

cell clones with TCRs recognizing a defined peptide epitope. (b) Tetramer binding patterns of

primary human T cells stimulated with unmethylated and methylated KRAS$_{G12V}$ peptide

presented by HLA-A2. Plots also indicate gates used to sort cells with different affinities for

unmethylated and methylated epitopes. (c) Dose response curves of TCRs against wild type

and KRAS$_{G12D}$ 5-14 peptides presented by HLA-A2. (d) Tumor cell growth during coculture of

CD8$^+$ T cells expressing selected TCRs with HLA-A2$^+$, wild type KRAS HeLa and KRAS$_{G12D}$

Panc1 pancreatic adenocarcinoma cell lines. Cocultures of T cells and tumor cells were

recorded by Incucyte imaging. Tumor cells expressed nuclear fluorescent protein. NR: red

fluorescent protein; NG: green fluorescent protein. (e) Alanine scan of KRAS$_{G12V}$ 5-14 epitope

tested against TCR$_{A2UoM1\text{-}1}$. Each box indicates the resulting expression of Nur77-GFP upon

replacement of a KRAS$_{G12V}$ residue with alanine (or in the case of alanine-11, replacement with

either glycine or threonine). (f) Dose response curve of CD8$^+$ Nur77-GFP Jurkat cells

expressing TCR$_{A2UoM1\text{-}1}$ after exposure to EPIPL$_{1948\text{-}1957}$, the potential human off-target peptide

as determined from alanine scan results.

Supplementary Figure 5. Further characterization of CRISPR knockout of *SUPT6H.* (a)

Location of the gRNA binding region in the *SUPT6H* gene and summary of the genomic position

of the commonly occurring mutations detected from DAN-G cells transduced with

lentiCRISPRv2 with gRNA targeting *SUPT6H*. Red dots indicate a mutation detected from long-

read sequencing. (b) Surface expression of HLA-A2 with and without *SUPT6H* knockout in

DAN-G cells. (c) Western blot of total expressed KRAS protein with and without *SUPT6H*

knockout in DAN-G cells. KRAS expression in other pancreatic adenocarcinoma cell lines

Panc1 and CFPAC1 are shown for additional comparison. GAPDH protein expression was used

as a control.

1111

1112   **Supplementary Table 1.** List of gRNAs for CRISPR knockout screen of methylation-related

1113   genes.

Figure 1. Posttranslational modification on the HLA-A2-presented KRAS$_{G12V}$ epitope alters recognition of the antigen by MHC class I-restricted TCRs and subsequent tumor cell killing.

Supplementary Figure 1. HLA-A2-presented KRAS$_{G12V}$ epitope includes a lysine residue that can be stably presented with potential side chain methylation.

a

Peptide sequence	Residues	%ile rank of HLA-A2 binding vs. random peptides
IQLIQNHFV	21-29	1.0784
KLVVVGAVGV	5-14	2.8794
LVVVGAVGV	6-14	3.1117
QLIQNHFV	22-29	3.8599
MTEYKLVVV	1-9	5.3608
GVGKSALTI	13-21	5.6588
TIQLIQNHFV	20-29	5.9529
GVGKSALTIQL	13-23	6.9975
KSALTIQLI	16-24	7.1313
KLVVVGAV	5-12	7.6549

d

Residue

MTEYKLVVVGAGGVGKSALT

Detected in:

LVVVGAGGVGK⭐ DAN-G

LVVVGAGGVGK DAN-G, CFPAC1

⭐ Lysine side chain trimethylation

f

Unmethylated

Dimethylated

Monomethylated

Trimethylated

g

1: Initial structural prediction by Alphafold

2: Introduction of PTMs in Rosetta

Figure 2. A computationally guided, rationalized mutagenesis library to develop MHC class I-restricted TCRs that recognize methylated epitopes of HLA-A2-presented KRAS$_{G12V}$ and achieve enhanced targeting of HLA-A2$^+$ KRAS$_{G12V}^+$ tumor cell lines.

Supplementary Figure 2. TCRs developed from a mutagenesis library to recognize post-translationally modified epitopes of HLA-A2-presented KRAS$_{G12V}$ are G12V mutation-specific.

Supplementary Figure 3. TCR-pMHC structural models by Alphafold-Rosetta show CDR3α core interactions with N-terminal lysine side chain of HLA-A2-presented KRAS$_{G12V}$ epitope.

a

TCR$_2$

b

TCR$_{EDST}$ vs. TCR$_2$

Figure 3. Stimulation of primary human T cells with unmethylated and methylated HLA-A2-presented KRAS$_{G12V}$ to generate MHC class I-restricted TCRs from a normal repertoire that recognize methylated epitopes.

Supplementary Figure 4. TCRs identified from primary human T cells specifically target KRAS_{G12V} epitope presented by HLA-A2.

Figure 4. A gene knockout screen for methylation-related genes identifies SPT6 (*SUPT6H*) as a participant in the demethylation of the HLA-A2-presented KRAS$_{G12V}$ epitope.

Supplementary Figure 5. Further characterization of CRISPR knockout of *SUPT6H*.