Title: **Human macula formation involves two waves of retinoic acid signaling suppression via**
CYP26A1 **regulating cell cycle exit and cone specification**

Authors/affiliations: Philippa Harding[1*], Maja Wojtynska[1], Alexander J. Smith[1], Robin R. Ali[1], Rachael
A. Pearson[1*]

[1]Ocular Cell and Gene Therapy Group, Centre for Gene Therapy and Regenerative Medicine, King's
College London, Tower Wing, Guy's Hospital, London, SE1 9RT, UK

[*] Co-corresponding authors - rachael.pearson@kcl.ac.uk, philippa.harding@kcl.ac.uk

6673 words

Abstract

The human macula is a specialized, M/L cone-rich region of the eye, critical for high acuity vision, but
little is known about the pathways regulating its development. Transcriptional regulation by Retinoic
Acid (RA) is essential for many aspects of human eye development. Here, we report a striking biphasic
expression of the RA-catabolizing enzyme, *CYP26A1,* in early human macular development between
post-conception weeks 6-17. Early inhibition of RA signaling in human retinal organoids (hROs)
prompts early cell cycle exit, and an increase in cone photoreceptors, while late inhibition alters cone
subtype specification. Conversely, *FGF8,* which is negatively regulated by RA and vital for High Acuity
Area specification in chick, is not expressed in the nascent human macula and had no effect on hRO
photoreceptor fate.

Keywords

Macula; fovea; Retinoic Acid; FGF8; high acuity area; cone photoreceptor; development; specification,
fetal, retinal organoids

Graphical abstract:

Acronyms

CMZ – Ciliary marginal zone

CS – Carnegie Stage

CYP26 - cytochrome P450 family 26

d – Day

FGF – Fibroblast growth factor

GCL – Ganglion cell layer

HAA – High acuity area

hROs – Human stem cell derived retinal organoids

INL – Inner nuclear layer

LE – Lens epithelium

mROs – Mouse stem cell derived retinal organoids

NBL – Neuroblastic Layer

ONH – Optic Nerve Head

ONL – Outer nuclear layer

PCW – Post-conception weeks

PM – Presumptive Macula

RA – Retinoic Acid

RAR – Retinoic Acid Receptor

RARE – Retinoic Acid Response Element

RFZ – Rod-free zone

RGC – Retinal ganglion cells

RPCs – Retinal Progenitor Cells

RXR – Retinoid X receptor

 Introduction

The macula is a 5-6mm region at the centre of the retina critical for high acuity and color vision in
humans and simian primates[1,2]. Damage and degeneration in this region lead to debilitating sight loss,
significantly impacting activities such as reading, recognizing faces, and driving. The macula has a
unique architecture comprising a high density of cone photoreceptors and retinal ganglion cells (RGCs),
contrasting with the rod-dominant peripheral retina, as well as possessing a highly ordered synaptic
configuration to aid rapid phototransduction[1,3,4]. Cone density is highest (200,000 cones/mm^2) at the
very centre of the macula in the foveola, a 300µm avascular rod-free zone (RFZ) consisting exclusively
of medium (M/green) and long (L/red) wavelength-detecting cones[1,3]. The proportion of cones:rods
declines with increasing eccentricity, being ~1:1 in the fovea (1.85mm from the fovea centre), dropping
to 1:30 in the peripheral retina[1,3].

Macular formation starts early in development, with immature cones present in the presumptive
macula (PM) by post-conception week (PCW)8/Carnegie stage (CS)23[2,5]. The macula region has
accelerated differentiation compared with the rest of the retina throughout prenatal development[2,6,7],
but adult cone density is not achieved until 4-6 years[3,8–10]. Despite its importance, the molecular
mechanisms underlying human macular development remain poorly understood, owing to the lack of
equivalent structures in rodents and many other classical animal models[2,11].

In addition to humans and primates, some other species possess a broadly analogous region of the
retina adapted for high acuity vision[12]. Chickens possess an RFZ, known as an *area centralis* or High
Acuity Area (HAA)[12–14]. Recent research has shown that chick HAA formation is dependent on
regulation of retinoic acid (RA) levels[13]. RA is a morphogen that can control gene expression through
binding to retinoic acid receptors (RARs) and retinoid X receptors (RXRs), which can then engage
specific DNA sequences known as retinoic acid-response elements (RAREs)[13,15]. Upon binding to
RAREs, co-activators are recruited and lead to transcriptional activation of specific RA-regulated
genes[15,16]. RA is synthesized by retinal dehydrogenases (ALDHA1-3) and catabolized by cytochrome
P450 family 26 enzymes (CYP26s)[15,17,18]. Transcriptional regulation by RA is key to development of
multiple tissues[16], including heart organogenesis[19], limb formation[20] and hindbrain patterning[21], and is
essential for human eye development, as demonstrated by the range of congenital ocular conditions
associated with pathogenic variants of the RA signaling pathway, including microphthalmia,
anophthalmia and coloboma (MAC)[22].

Modulation of RA signaling has been implicated in the formation of HAAs in other organisms, including
high expression of *Cyp26a1* by Müller glia in the zebrafish HAA at later stages of development[23]. Strong
expression of *Cyp26a1/c1* was observed at a single spot at the presumptive HAA early in chick
development, coincident and colocalized with *Fgf8* (fibroblast growth factor 8) expression, which is
directly downregulated by RA through binding to a RARE upstream of *Fgf8* [13,16,24]. In the chick retina,
RA was found to negatively regulate *Fgf8*, with transient addition of RA leading to loss of *Fgf8*
expression, disruption of the RFZ, and a decrease in the high RGC density typical of the HAA.
Furthermore, expression of *CYP26A1* mRNA was observed in the human PM at CS22 (PCW8).
Subsequent studies have reported *CYP26A1* to be significantly upregulated in macula compared with
peripheral retina through bulk and scRNAseq in mid-gestation (PCW20) and adult retina human
retina[25]. *CYP26C1* was found to have no detectable expression in developing human retina in both
RNAseq and *in situ* experiments[13,26]. Recently, detailed analysis of *CYP26A1* and *FGF8* expression in

Rhesus monkey found that *CYP26A1* is highly expressed in the developing macular region at day 40,
after the onset of cone genesis (corresponding to human ~PCW10), while *FGF8* was not localized to
the macular region[27]. However, little is known about earlier developmental stages encompassing the
period of human macula specification. Evaluation of *CYP26A1* and *FGF8* expression has not been
performed in human fetal retina, and a conserved functional effect of RA suppression and/or
upregulated *FGF8* signaling in the initial stages of human macula specification has not been
demonstrated.
105
Understanding macula specification is important for the utilization of human stem cell-derived retinal
organoids (hROs). Stem cell-derived hROs can produce layered structures that broadly resemble the
human neural retina and closely follow the timing of human retinal development[28,29]. They have wide
therapeutic potential, serving as donor material for cell/graft transplantation[28,30–34], as well as
preclinical models for the study of development and disease mechanisms and testing new therapeutic
compounds and gene therapies. Generation of cones has been of particular interest for photoreceptor
replacement therapies and attempts to improve cone density of hROs have been plentiful, such as
through stage-specific inhibition of Notch signalling,[31] removal of RA[35] and manipulation of thyroid
hormone signaling. To date, however, hRO protocols have failed to generate macular-like cells with *in*
vivo-like spatial regionalization or the genetic signature of foveal cones[25,28,36], suggesting that more
complex signaling than simple removal of RA alone is involved, while the role of FGF8 has not been
examined. Here, we used *in situ* hybridisation in human fetal tissue and pharmacological manipulation
in hROs to investigate the spatiotemporal expression and potential roles of RA and *FGF8* in human
macular development.

Results

The RA catabolizing enzyme CYP26A1 *displays a biphasic pattern of expression in the developing human macula*

To determine expression patterns of RA catabolizing enzyme *CYP26A1* and growth factor *FGF8* in the developing human macula, RNAscope™ *in situ* hybridisation was performed on fetal retina samples between PCW6 (CS16) and PCW17 (**Figure 1**). No *CYP26A1* was visible at PCW6 (CS16/17), with weak expression first visible at PCW7 (CS18/19), localized to a single, spot-like region at the centre of the retina, approximately 1mm temporal to the optic nerve head (ONH) (**Figure 1a**), a location identified previously as the presumptive macula (PM)[6]. Within this region, *CYP26A1* was distributed throughout the neuroblastic layer (NBL), colocalizing with VSX2+ve RPCs (**Figure 1b**). At the start of PCW8 (CS20-21), much stronger expression of *CYP26A1* was observed in the VSX2+ve NBL (**Figure 1a)**, with expression reducing by the end of PCW8 (CS22-23), by which point some RPCs expressed the cone/bipolar progenitor marker, OTX2 (**Figure 2a**). These were distributed sparsely throughout the NBL of the entire retina, although OTX2 labelling was strongest in the PM region and, within this, in the more apically located cells (**Figure 2ai-ii**). The nascent GCL could also start to be defined in PCW8 samples (**Figure 2a**). No expression of the cone photoreceptor-specific markers RECOVERIN or RXRγ was visible at this timepoint (**Figure 2aii**). *CYP26A1* was also expressed in the lens epithelium (LE) and at low levels at the nasal edge of the retina and around the ONH at CS18-20 (**Supplementary Figure 1**). By PCW10, *CYP26A1* mRNA expression was visibly weaker in the PM region, indicating that expression is downregulated by this developmental stage (**Figure 1a, b**). Moreover, its expression was restricted to a band of VSX2+/OTX2+ cells located in the nascent INL. RXRγ+/RECOVERIN+/OTX2+ cone photoreceptors did not express *CYP26A1* and were located exclusively within a single cell layer (the nascent outer nuclear layer; ONL) at the apical edge of the presumptive macula; RXRγ+/RECOVERIN+ cones were not present in the peripheral retina at this stage (**Figure 2b**). At PCW12, *CYP26A1* expression was visibly increased, compared with PCW10, this time localizing to the basal edge of the VSX2+ INL; by PCW16/17, expression of *CYP26A1* was restricted to the VSX2+/SOX9+ developing Müller glia layer, and was not visible in the VSX2+/SOX9- presumptive bipolar cells, which express much higher levels of VSX2 than presumptive Müller glia[43,44] (**Figure 1b, Figure 2c**).

FGF8 *is not expressed in the developing macula, but instead is restricted to the Optic Nerve Head in the early human retina*

In the chick, *Fgf8* is expressed in a bullseye-like pattern overlapping with *Cyp26a1/c1* and is essential for forming the HAA[13]. In notable contrast, no *FGF8* mRNA labelling was observed in the human PM region between PCW6-10, during which macula cone precursors are born, indicating that *FGF8* is not required for cone specification in the human macula (**Figure 1**). *FGF8* was visible between PCW7-12, with expression highest during early PCW8; however, expression was restricted to around the presumptive optic stalk, similar to what has previously been reported for the macaque from 40 days gestation[27], as well as the developing mouse eye from E9.5-14.5[37] and chick eye from E2-E7[38], where it is understood to maintain Pax2 expression (**Figure 3**). Expression remained low during PCW12-17, and again was only seen at low levels and restricted to around the ONH. Overlapping co-expression of *CYP26A1* and *FGF8* was not observed at any time point examined.

 Presumptive macula is located more peripheral in early development relative to its central position in
the mature retina due to differential proliferation rates

In the mature adult eye, the macula is located at the centre of the retina, opposite the pupil and approximately 3.5-5.5mm temporal to the ONH[39]. At early developmental timepoints, expression of *CYP26A1* was closer to the peripheral edge of the retina than expected for the PM, relative to its more central location in the adult eye. Increased proliferation of RPCs in the peripheral, compared with central, retina has been previously demonstrated through proliferative marker staining at later timepoints in development (post-PCW8)[2]. However, due to lack of available tissue and macular markers at early stages, proliferation and how this variable growth rate affects the location of the PM has not previously been investigated. Moreover, whilst a centre-to-periphery gradient of proliferation is well known, it is less clear how this relates to the PM specifically, nor whether this region stops proliferating at a different time to the rest of the central retina.

To explore this, we examined growth and proliferation across regions of the retina throughout early development. In the first instance, we measured the distance between the centre of the region of *CYP26A1* expression and i) the peripheral edge of the retina and ii) the ONH. Distance from the centre of *CYP26A1* expression to the peripheral edge of the retina increased from 300μm at PCW7 to 9000μm at PCW17, growing in a linear manner (R^2=0.96) (**Figure 4ai-iv**). The distance from the centre of *CYP26A1* expression with respect to the ONH also grew in a linear manner (R^2=0.94), from 400μm at PCW7 to 6000μm at PCW17 (**Figure 4ai-iv**). However, comparing the rates of growth, the increase in distance between the macula and periphery was approximately two-fold higher, compared with the macula to ONH distance (a rate of 114μm/day and 62μm/day respectively, *p*>0.001). Notably, while the macula-to-periphery distance increases from 20% of the total retina length at PCW7 to 45% at PCW17, macula-to-ONH remains constant at 30% (**Figure 4av**). This shows that the distance between the macula and the peripheral edge of the retina increases at approximately twice the rate of the central region, and that the *CYP26A1*+ macula region is located more temporally early in development and shifts closer, relative to total retinal length, to the ONH as the peripheral retina grows.

We next examined regional proliferation using the proliferative marker KI67. Consistent with the expected centre-to-periphery gradient of maturation, between CS18 and CS23, the proportion of cells that were KI67+ was higher at all timepoints in the peripheral retina compared with the central retina, with twice as many KI67+ cells at CS23 (CS18: 47% ±13.6 vs 66% ±4.7, n=3; CS20: 36% ±3.4 vs 68% ±11.6, n=3; CS23: 22% ±3.7 vs 45% ±19.1, n=3) (**Figure 4bi-ii; Supplementary Figure 2a**). This difference was statistically significant at CS20, and close to statistical significance at CS18 and CS23, likely due to low sample sizes (CS18: *t*(2)=3.3, *p*=0.08; CS20: *t*(2)=6.4, *p*=0.02; CS23: *t*(2)=2.5, *p*=0.13) (**Figure 4bi-ii**).

In addition to comparing proliferation in the central and peripheral regions, we also explored potential differences between the PM and a region in the nasal retina equidistant from the periphery as the macula, i.e. at the equivalent region of the central-peripheral gradient. (**Figure 4biii; Supplementary Figure 2a-b**). Both the macular and nasal regions showed significant reductions in the proportion of proliferating cells between CS18 and CS20/CS23 (**Figure 4biii**). However, linear regression showed that the slope in the macula region was nearly twice as steep as that of the equivalent nasal region between CS18 and CS20 (-15.6%/day and -8.6%/day, respectively, *p*=0.046), indicating the rate at which

proliferation slows is twice as fast in the macula across this time window (**Figure 4biv, Supplementary**
Figure 2c).

At the earliest timepoint, CS18, the macula had one of the highest proportions of KI67+ cells of any
region, similar to the periphery (mean of 65% ±6.3 vs 66% ±4.7), while the central/nasal regions had
lower proportions of KI67 expressing cells (mean of 47% ±13.6 vs 56% ±1.6) (**Figure 4bii-iii,**
Supplementary Figure 2b). However, the number of proliferating cells in the macula halved by CS20,
resulting in a similar mean to the central retina by this timepoint (33% ±5.4 vs 36% ±3.4), which
persisted to CS23 (24% ±7.3 vs 22% ±3.7). Notably, this reduction in proliferation rate is coincident with
the timing of *CYP26A1* expression.

Human stem cell-derived retinal organoids express components of RA synthesis pathway during early
differentiation, demonstrating similarity to peripheral retina development

We next sought to examine the impact of RA signaling on hRO differentiation to better understand its
role in macula formation. We have previously reported that a 2D/3D differentiation protocol yields
hROs from pluripotent stem cells and that this method faithfully replicates the timing of human retinal
development[28] (**Supplementary Figure 3a**). This protocol involves the addition of RA from day 70 (d70)
onwards and yields retinas with ~1:4 ratio of cones:rods, a proportion similar to the outer edges of the
macula[3]. We therefore explored the expression of components of RA metabolism and FGF8 signaling
in hROs generated using this method in the first instance.

The RA synthesizing enzymes, *ALDH1A1* and *ALDH1A3,* were both upregulated early in differentiation,
at the time of hRO picking, at ~d40, compared with undifferentiated hESCs, followed by a steady
downregulation of *ALDH1A1* to d90 (**Figure 5a-b**). In contrast, qRT-PCR analysis showed limited
expression of *CYP26A1* in d40 hROs, but this was subsequently upregulated at d80, following
exogenous addition of RA to culture medium at d70, and by which time photoreceptor specification
has already occurred (**Figure 5c, Supplementary Figure 3**). CRABP2 binds to cytoplasmic RA and traffics
it to the nucleus for regulation of gene expression[40]. As *all*-trans retinoic acid concentration increases
(atRA), so *CRABP2* expression is concomitantly upregulated, making it a proxy for circulating RA
levels[40]. In hROs, *CRABP2* was upregulated at d40, compared with undifferentiated hESCs, and
continues to increase throughout differentiation, but significantly increases beyond d70, when
exogenous RA is added. This demonstrates that endogenous RA signaling occurs throughout hRO
differentiation, and the pathway is, unsurprisingly, significantly upregulated following addition of
exogenous RA (**Figure 5d**). RA signaling and lack of *CYP26A1* expression prior to cone specification in
hROs points to a more peripheral retina-like molecular developmental program in hROs produced
using this protocol.

Expression of *FGF8* mRNA was significantly increased by d40 of differentiation compared with
undifferentiated hESCs, with expression levels then steadily decreasing throughout differentiation, up
to d90 (**Figure 5e**). mRNA expression for FGF receptors *(FGFR)1-4* was detected in hESCs d0 (CT values
25-35); however, expression of *FGFR1/2/4* significantly decreased over the course of hRO
differentiation, while *FGFR3* expression increased significantly between d0 and d40, followed by
subsequent slowly decreasing expression between d40-d90 (**Figure 5f**).

Early inhibition of RA by AGN193109 leads to smaller hROs and reduced proliferation

Based on mRNA expression and immunostaining analysis showing onset of cone specification by d40 of differentiation (**Supplementary Figure 3**), hROs were dosed with pan-Retinoic Acid Receptor (RAR) antagonist AGN193109 (10µM)[41] or recombinant growth factor FGF8 (rFGF8) (100ng/ml) initially with for a single pulse (early) from d42-50 of differentiation in order to mimic the early phase of _CYP26A1_-induced RA catabolism, as seen in the human PM during PCW7-8. Dosing concentrations was determined using a dose-response viability assay and effect on gene expression (**Supplementary Figure 4**). Early dosing with 10µM AGN193109 significantly reduced expression of _CRABP2_ ($t(2)=5.77$, $p=0.03$), demonstrating reduced levels of circulating RA[40] (**Supplementary Figure 5a**).

Given the rapid reduction in proliferative cells observed in the RPCs of the macula during the first phase of _CYP26A1_ expression, relative to other regions of the retina (**Figure 4a**), we first examined growth rates in hROs dosed with RA inhibitor AGN193109. Measurements of hRO size before and after early dosing showed RA inhibition led to a very significant reduction in growth by day 60, compared with DMSO-treated control hROs ($t(32)=5.18$, $p<0.0001$) (**Figure 6a**). Quantification of immunostaining with proliferative marker KI67 showed a significantly reduced percentage of KI67+ cells in AGN193109-dosed hROs ($t(24)=2.48$, $p=0.02$), while no difference in apoptotic marker CASPASE-3 was observed ($t(29)=0.64$, $p=0.5$), indicating that the reduction in growth in RA-inhibited hROs was due to lower levels of proliferation, rather than exacerbated cell death (**Figure 6b**). These data indicate that early inhibition of RA in RPCs leads to increased cell cycle exit, which may uncover the role of the first phase of _CYP26A1_ expression in the developing human macula at PCW7-8.

Early pharmacological inhibition of RA signaling in hROs increases expression of photoreceptor markers, while addition of recombinant FGF8 has no effect, mimicking in vivo macula formation

Since application of AGN193109 led to a reduction in RPC proliferation, but no change in cell death, it is plausible to assume that RPCs are being induced to exit the cell cycle prematurely. We therefore examined whether this had any effect on the adoption of early born cell fates (predominantly cone photoreceptors and RGCs). Transient early inhibition of the RA pathway led to significant increases in the expression of the pan-photoreceptor marker _RECOVERIN_ ($t(5)=2.56$, $p=0.05$), at d60, along with cone progenitor maker _OTX2_ ($t(4)=4.58$, $p=0.01$), while expression of the cone precursor markers, _RXRγ_ and _THRβ,_ also slightly increased, although were not significant ($t(5)=1.45$, $p=0.2$); ($t(5)=1.69$, $p=0.15$, respectively) (**Figure 6ci**). No significant changes were observed in the expression of RPC marker _VSX2_ ($t(5)=1.39$, $p=0.2$) (**Figure 6c**) or RGC marker _RBPMS_ and RGC/bipolar/amacrine marker _ISLET1_ ($t(4)=1.02$, $p=0.37$; $t(4)=0.30$, $p=0.78$, respectively) (**Supplementary Figure 5b**). Quantification of immunostaining showed a significant increase in the proportion of cone progenitor cell marker OTX2+ cells in RA-inhibited hROs ($t(28)=2.4$, $p=0.02$), reflecting gene expression analysis (**Figure 6d**). Also consistent with gene expression analysis, no change in the proportion of VSX2+ RPCs ($t(29)=0.01$, $p=0.99$) or cone precursor marker RXRγ was observed ($t(26)=0.96$, $p=0.35$) (**Figure 6d, Supplementary Figure 5c**). Conversely, addition of rFGF8 between d42-50 did not lead to any changes in RPC or photoreceptor marker expression compared with PBS dosed controls (_RECOVERIN_: $t(2)=0.22$, $p=0.85$; _OTX2_: $t(2)=2.99$, $p=0.10$; _RXRγ_: $t(3)=0.23$, $p=0.83$; _THRβ_: $t(2)=1.01$, $p=0.40$; _VSX2_: $t(3)=0.39$, $p=0.72$) (**Figure 6cii**). These results indicate that early inhibition of RA signaling leads to early exit from the cell cycle and the generation of early-born OTX2+ cone progenitor cells.

 Early and late RA inhibition, mimicking biphasic CYP26A1 expression, results in increase in M/L-OPSIN,

indicating a role in cone subtype specification

To investigate the effect of the early and late phases of *CYP26A1* expression, hROs were dosed with an

early dose of AGN193109 between d42-50, alongside a second dose at d70-80 (early+mid) or a second

and third dose at d120-130 (early+mid+late). At d150, no changes in rod marker *RHODOPSIN*

expression ($t(2)=1.03$, $p=0.41$; $t(3)=0.65$, $p=0.76$; $t(3)=2.43$, $p=0.15$) or pan-cone marker *ARRESTIN-3*

($t(2)=0.82$, $p=0.50$; $t(3)=0.43$, $p=0.88$; $t(3)=1.22$, $p=0.46$) were observed between any AGN193109-

dosed organoids and DMSO controls (**Figure 7a**). Dosing with AGN193109 at the early timepoint alone

also showed no changes in expression of blue-wavelength detecting cone marker *S-OPSIN*, or

green/red-wavelength detecting cone markers *M/L-OPSIN*s at d150. However, additional pulses

(early+mid and early+mid+late) of AGN193109 both resulted in a reduction of *S-OPSIN* ($t(3)=1.67$,

$p=0.30$ and $t(3)=5.64$, $p=0.02$, respectively) and an increase in *M/L-OPSIN* mRNA expression at d150

approaching significance ($t(3)=1.70$, $p=0.29$; $t(3)=2.37$, $p=0.16$), indicative of the M/L-cone-rich fovea[42]

(**Figure 7a**). Expression of both *M-* and *L- OPSIN* showed no change with early dosing but higher

expression with both early+mid and early+mid+late dosing, although *L-OPSIN* showed a far greater

fold change (*M-OPSIN* 1.5x, *L-OPSIN* 8x), and only early+mid+late *L-OPSIN* approached significance

($t(3)=2.70$, $p=0.07$). Immunostaining of early+mid+late dosed hROs found no significant difference in

the proportion of RHODOPSIN/ARRESTIN expressing cells, or the proportion of M/L-OPSIN+/ARRESTIN-

3+ cells, but RA inhibition led to a significant reduction in S-OPSIN+/ARRESTIN-3+ expressing cells

($t(11)=3.14$, $p=0.01$) (**Figure 7b**).

In addition, all doses (early; early+mid; early+mid+late), RA-inhibited hROs showed upregulation of

the WNT inhibitor, *SFRP1,* at d150, although this was only statistically significant for early+mid+late

($t(2)=2.02$, $p=0.18$; $t(3)=2.45$, $p=0.15$; $t(3)=5.04$, $p=0.02$, respectively) (**Figure 7ci**). Early dosing also

showed a significant increase in *SFRP1* expression at d60 ($t(4)=4.50$, $p=0.01$) (**Figure 7cii**).

Taken together, this indicates that suppression of RA signaling mediated by the later phase of *CYP26A1*

expression is important for specifying green/red cones in the human fovea, and that the effects of both

early and late RA inhibition may involve inhibition of WNT signaling through SFRP1.

Discussion

In this study, we provide the first detailed spatiotemporal analysis of *CYP26A1* and *FGF8* expression in human fetal retina. We also investigated the roles of these pathways in macular development through manipulation of hROs. Our aim was to begin to unravel the precise functional roles of these pathways in producing the specialized cellular structure of the human macula.

Analysis of early-stage human macular development has been very limited to date, with small sample sizes due to scarcity of samples. The earliest previous *in situ* spatial analysis of *CYP26A1* expression in human fetal retina was CS22-23 (late PCW8), which showed expression in the presumptive macula (PM) at CS22 but switched off at CS23[13]. Here, we have defined the developmental timeline of *CYP26A1* expression, which shows a previously unreported and striking biphasic pattern of expression in the PM. The first phase of expression is initiated in the PM at PCW7 (CS18), prior to cone specification, and is maintained in RPCs until the end of PCW8. By PCW10, *CYP26A1* expression in the RPCs is markedly reduced. Closer examination of the PM region showed that during PCW7-8, *CYP26A1* is expressed by RPCs but not by cone photoreceptor precursors, which are born by PCW10 and are localized to the most apical layer of the PM. The second phase of expression begins from PCW12 and is restricted to Müller glia residing in the INL of the retina, but not expressed in the developing bipolar cells.

Our findings are supported by a very recent report of *CYP26A1* expression in rhesus macaque development[27], which presented strong *CYP26A1* expression in the macular region. However, in the macaque, *CYP26A1* expression was only assessed after cone specification (from day 40), by which time RXRγ+ cone precursor marker was expressed in the NBL of the macular region; this is equivalent to around PCW10 in the human retina (~25% through gestation)[5]. In both human and macaque retina, *CYP26A1* was only expressed in RPCs during the early macular formation, with no *CYP26A1* mRNA observed in the OTX2+/RXRγ+/RECOVERIN+ cone precursor cells of the ONL. We also observed similar *CYP26A1* expression outside the PM in LE, CMZ and around the ONH in early (PCW7-8) development as reported in macaque, although the level of expression relative to the PM, appears much lower in the human fetal samples compared with the monkey retina[27]. Our observation of *CYP26A1* expression in the Müller glial cells after PCW12 is consistent with previous descriptions for the later stages of HAA maturation in the Zebrafish[23] and the mature primate macula in the macaque[27], as well as transcriptomic data for foveal Müller cells in adult human retina[25]. Taken together, we propose a biphasic model of *CYP26A1* expression in the macula, with the first wave onset at PCW7-8 in RPCs, followed by a second phase from PCW12, specific to Müller glia.

In striking contrast to HAA formation in the chick[13], *FGF8* was not expressed in the PM at any time between PCW6 and PCW17 of development nor exhibited any spatial overlap with *CYP26A1* expression, indicating that it does not play a role in human macula formation or macula cone differentiation. *FGF8* mRNA was instead found around the ONH during the same period of development. This is also consistent with the macaque retina, where an ONH-to-periphery gradient of *FGF8* expression was observed[27]. Moreover, we found addition of rFGF8 to hRO cultures had no impact on photoreceptor marker expression. As such, this demonstrates a notable divergence in the molecular mechanisms underlying high acuity region development in different species.

We next sought to explore the potential functions of RA inhibition in human macular development by manipulating RA signaling in hROs via pharmacological inhibition of RARs[41] in hROs to mimic *CYP26A1*

expression. Increased proliferation of RPCs in the periphery, compared with the central retina, has
previously been reported for developmental timepoints after PCW8 in human tissue[2,6] and after fetal
d80 in marmoset[45]. However, due to lack of markers and access to tissue, regional variation in
proliferation, including the macula, has not been shown for earlier stages of development. We found
that RPCs of the macula had the highest proportion of proliferative cells at CS18, but that this reduced
at twice the rate of the equivalent nasal region by CS23. Indeed, the proportion of proliferative cells
was already significantly reduced by CS20, coincident with *CYP26A1* expression. This suggests that
inhibition of RA via *CYP26A1* is required to encourage cell cycle exit. Early inhibition of RA signaling in
hROs supports this notion, with a significant growth reduction compared with controls. Furthermore,
staining with KI67 showed significantly fewer proliferating cells in dosed organoids. Thus, we propose
that a reduction in RA signaling caused by *CYP26A1* during the first phase of expression encourages
cell cycle exit of RPCs within the PM. Furthermore, hROs treated early with RAR antagonist displayed
an increase the number of cone-biased progenitors at d60, consistent with early cell cycle exit of RPCs
leading to differentiation of early-born cell types, particularly cones, which dominate the macula. This
aligns with our previous data investigating the effects of exogenous RA on mouse ESC-derived retinal
organoids (mROs), which found a short pulse of RA from d14-16 (cone precursor stage) led to an
increase in S-phase cells and overall cell yield, indicating RA promotes cell proliferation in RPCs[31].

Cone subtypes are determined by two fate check points: first, between S or M/L fate, and if the latter,
between M- or L-OPSIN expression[46]. When RA signaling was inhibited early in differentiation, prior to
cone specification, no change in cone subtype at d150 was observed. However, RA inhibition during
both mid and late hRO differentiation, after cone specification, led to upregulation of *M/L-OPSIN* and
a significant downregulation of *S-OPSIN* gene expression. Whilst no significant change in the
proportion of M/L-OPSIN protein expressing cells occurred, a significant reduction in S-OPSIN
expressing cells was seen. This is notable since the central retina contains a higher density of medium
(M)/long (L) wavelength-detecting cones than short (S) wavelength cones, which dominate the
peripheral retina, and the human foveola at the centre of the macula exclusively contains M/L-cones[42].
Interestingly, a similar effect of *CYP26A1*-mediated promotion of M-cone identity has been reported
in *VAX2* knockout mice, where lack of *VAX2*-mediated restriction of *CYP26A1* expression to a stripe in
the central retina leads to the expansion of M-cones into the typically S-cone dominated ventral
retina[47]. Moreover, addition of exogenous RA to mROs after cone precursor specification led to
upregulation of S-opsin but no change in Arrestin-3 expression, indicating that alterations in RA
signaling later in development also regulates cone subtype specification in mouse development[28]. In
our hands, pulses of RA inhibition early+mid+late in hROs led to an increase *L-OPSIN* expression,
broadly aligning with a recent report by Johnston and colleagues indicating that sustained application
during the same period of differentiation (d43-d130) of RA suppresses L-cone formation, whilst
promoting M-cone fate[48], albeit that we did not see a downregulation of *M-OPSIN*.

Recent scRNAseq data of human fetal retina has shown that the WNT pathway inhibitor *SFRP1* is
expressed in early RPCs and, later in development, by Müller glia, with strongest expression in the cells
of the macula, confirmed by *in situ* hybridisation at PCW18[11]. Thus, *SFRP1* shows a remarkably similar
biphasic and spatiotemporal expression pattern to *CYP26A1* in human fetal retina. Both early and late
inhibition of RA signaling in hROs led to a significant upregulation of *SFRP1*, indicating that *CYP26A1*-
mediated RA inhibition leads to suppression of the WNT signaling pathway. Canonical WNT signaling
commonly regulates proliferation through binding of β-catenin; it is plausible, therefore, to speculate

that SFRP1-induced inhibition of the WNT-β-catenin pathway may induce cell cycle exit in early RPCs,
which leads to lineage specification of cone precursors[49]. Wnt signaling has been shown to play a role
in maintaining RPCs in a proliferative state in Drosophila[50], zebrafish[51], *Xenopus*[52] and mice[53]. Notably,
however, in chick, Wnt was not found to alter progenitor proliferation, but instead promotes
expression of markers associated with peripheral retinal cell fate, ciliary body and iris[54]. This divergence
in Wnt pathway function between human and chick may help explain the lack of *FGF8* expression in
the presumptive human macula, in contrast to the previously established co-expression with
Cyp26a1/c1 in the chick HAA[13].

Together, these data show for the first time a biphasic pattern of expression of *CYP26A1* across human
macula development. Conversely, *FGF8* expression is restricted to the region around the ONH and does
not exhibit any spatial overlap with *CYP26A1*, in contrast to the chick suggesting an important
divergence of regulatory mechanisms in high acuity specification. We propose novel roles for the two
phases of RA inhibition in macular formation: the early wave of RA inhibition (PCW7-8) stimulates RPCs
to exit the cell cycle, encouraging cone generation, while the second, later phase (from PCW12) plays
a role in specification of cone subtypes. Upregulation of *SFRP1* following RA inhibition in hROs indicates
that either, or both, of these processes may involve *SFRP1*-mediated WNT inhibition, although further
work is required to test this directly.

This work bridges a gap in fundamental knowledge of the molecular mechanisms underlying human
macular development and highlights the importance of studying human development in both animal
models and human-derived tissue. These results open two avenues of future developmental research:
1) can we resolve the molecular network downstream of *CYP26A1* to establish how these pathways
generate the unique structure of the human macula; and 2) what are the drivers of *CYP26A1*
expression in the human PM and how are they initiated? Elucidating these molecular mechanisms
could provide strategies to encourage formation of more macular-like cellular structure in *in vitro*
models, which may be used to study macular development and disease, provide a preclinical model
for drug testing to treat macular degeneration and improve efficacy of cells used for transplantation
therapies which could restore visual function in patients by cell transplants.

STAR Methods

<u>Resource Availability</u>

Corresponding Author - Further information and requests for resources and reagents should be
directed to and will be fulfilled by the corresponding author, Rachael Pearson
(rachael.pearson@kcl.ac.uk).

<u>Materials Availability</u>

This study did not generate new unique reagents.

<u>Data and Code Availability</u>

Mean data is presented as mean ±SD, as indicated in figure legends. This paper did not generate
original code. Any additional information/data is available from the corresponding contact upon
request.

<u>Human fetal retina sample collection</u>

Human fetal retina samples were collected from the Human Developmental Biology Resource
(http://www.hdbr.org/) with research ethics approval and informed consent (Project Number 200599).
Samples were stored in full compliance with the Human Tissue Act 2004.

<u>Differentiation of retinal organoids</u>

Human PSCs were maintained on Geltrex-coated six-well plates in E8. Human retinal organoids (hROs)
were generated from the H9 ESC line according to a previously published protocol[28]. In brief, hESCs
were maintained to 90% confluency, then on day 1 (d1) E6 medium was added for 2 days, followed by
Proneural Induction Medium (PIM), which was changed every 2-3 days. Cultures were maintained in
PIM for up to 40 days, during which time hROs started to appear and were individually excised and
transferred to low-binding 96-well plates, where they were maintained in 3D culture in Retinal
Differentiation Media (RDM). At d42, additional factors were added to the RDM (RDM+F) which were
maintained until d70. On d70, individual hROs were transferred into low-binding 24-well plates and
Retinoic Acid (RA) was added fresh to the RDM+F at a concentration of 1mg/ml. On d90, hROs were
transitions to Retinal Differentiation Medium (RDM90), to which fresh RA was added at a
concentration of 0.5mg/ml[28].

<u>Fetal retina/hRO sample preparation for immunohistochemistry/RNAscope™</u>

For immunohistochemistry (IHC)/RNAscope™, human fetal samples were fixed for 1 day in 10%
formalin before transferring to PBS. Samples were incubated overnight in 20% (w/v) sucrose, prior to
cryo-embedding in OCT matrix. Tissue was cut at 10μM thickness. hROs were washed with PBS and
fixed for 1 hour in 4% PFA, followed by PBS wash and incubated overnight in 20% (w/v) sucrose, prior
to cryo-embedding in OCT matrix. Tissue was cut at 10μM thickness, mounted on Superfrost glass
slides, air-dried overnight and frozen at -20°C.

RNAscope™ *in situ* hybridisation

RNAscope™ Multiplex Fluorescence Assay was performed using manufacturer's protocol for fixed
frozen tissue with manual antigen retrieval. In short, samples were baked at 60°C for 30 minutes,
postfixed with 4% PFA for 15 minutes at 4°C then dehydrated using an ethanol gradient, before 10-
minute treatment with hydrogen peroxidase. Manual antigen retrieval was performed at 80°C for 5
minutes, then sections were pretreated with protease III for 30 minutes at 40°C. Sections were
incubated with probe Hs-CYP26A1-C1 (#487741) and Hs-FGF8-C2 (#415791-C2) for 3 hours at 40°C,
before storing in 5xSSC overnight. Probes went through series of amplification steps, before developing
the signal and labelling each with OPAL cyanine 3/5 fluorophores, respectively. Slides were then
counterstained with VSX2 and Alexa-fluor 488 and DAPI according to the IHC protocol below and
mounted in DAKO mounting medium. For **Figure 1**, sample numbers were as follows: CS16/17: n=3,
CS18/19: n=2, CS20/21: n=4, CS22/23: n=4, PCW10: n=3, PCW12: n=2, PCW16/17: n=2. Regional length
measurements were taken using ImageJ and fitted and analyzed using linear regression analysis in
GraphPad Prism (v10.1.2).

Immunohistochemistry (IHC)

Cryosections were permeabilized for 1 hour at room temperature with PBS/0.1% Tween/0.5% Triton
X, then blocked for 2 hours at room temperature with PBS/0.1% Tween/5% donkey serum/1% BSA.
Primary antibodies with incubated overnight at 4°C. Alexa Fluor® secondary antibodies were incubated
for 1 hour at room temperature at 1:400 dilution, along with DAPI counterstain. Confocal images were
taken using a Leica DM5500Q and processed in ImageJ. Spot count analysis was performed using Imaris
10.0. For analysis of KI67 in human fetal tissue, the same length of region was selected at different
regions of the same retina: the macula (identified by *CYP26A1+* staining); the central region between
the macula and nerve; the region the same distance as the macula from the periphery on the nasal
side of the retina; and the periphery (Supplementary Figure 2a). For analysis of CAPSASE staining in
hROs, necrotic core was masked to ensure only cell death in the neural retina cells was included
(**Supplementary Figure 6**). Antibodies are listed in **Supplementary Table 1**. Analysis was performed
using unpaired t-tests.

qPCR gene expression analysis

For all qPCR assays, 3-5 hROs were pooled for each sample depending on collection timepoint (5 for
d40-d90, 3 for d120-150). Samples were taken from independent experiments and a minimum of n=3
independent samples were used for analysis. RNA was extracted from hRO pellets using RNeasy Micro
kit (Qiagen), according to manufacturer's instructions. Samples were eluted in 30μl and stored at -
80°C. Concentration and RNA quality analyzed using a BioDrop. cDNA was synthesized using
QuantiTect Reverse Transcription Kit, according to manufacturer's instructions. Quantitative RT-PCR
was performed using Perfecta Low-ROX MasterMix with custom primers listed in **Supplementary Table**
2 and probes from the Merck Universal Probe library or PowerTrack SYBR MasterMix with ROX as
passive reference. All samples were run in triplicate alongside 2 endogenous reference genes (*GAPDH*
and *β–actin*), along with water and no RT negative controls. Comparative CT was calculated to
determine relative gene expression as fold change from either d0 PSCs or undosed controls[55].

AlamarBlue viability assay

Optimal dosing concentrations of recombinant FGF8 and AGN193109 were determined using cell viability analysis by Alamar blue (**Supplementary Figure 4**). AlamarBlue utilizes resazurin which is reduced in presence of living cells and is non-toxic/cell permeable. Organoids were kept in separate wells of a 96-well plate and incubated for 4 hours with 10% AlamarBlue diluted in media, after which AlamarBlue was transferred to a new 96 well plate, organoids were washed 3 times with PBS and returned to media. Fluorescence of AlamarBlue at 590nm (RFU) was then measured using a plate scanner and % reduction calculated using:

$$Reduction_{AB} = \frac{RFU^{Exp} - RFU^{Neg}}{RFU^{100\%} - RFU^{Neg}} \times 100$$

In which $Reduction_{AB}$ is the percentage reduction of AlamarBlue, RFU^{Exp} is the Relative Fluorescence Units (RFU) for the experimental sample, RFU^{Neg} is the RFU for the Negative control, and $RFU^{100\%}$ is the RFU for the 100% reduced positive control.

To control for differences in hRO size, change in reduction from day 0 was calculated for each hRO:

$$\Delta Reduction = Reduction_{AB}^{Day\ X} - Reduction_{AB}^{Day\ 0}$$

Relative change in viability of treated organoids was normalized to DMSO treated controls:

$$\Delta Viability = \frac{\Delta Reduction^{Exp}}{\Delta Reduction^{Control}} \times 100$$

In which $\Delta Viability$ is the change in viability from Day 0 in the experimental sample relative to the DMSO-dosed control sample

As hRO viability was not significantly reduced upon dosing with 10μM AGN19309, but was significantly reduced upon dosing with 20μM AGN19309, 10μM AGN19309 was used for all following experiments. No reduction in viability was observed at any concentration of rFGF8.

FGF8 can self-regulate its mRNA expression[56]. Consequently, expression of *FGF8* was assessed following dosing with rFGF8. No significant difference in *FGF8* expression was observed at any concentration, however, the largest mean difference was found at 100ng/μl. Consequently, this concentration was used for all following experiments.

561   hRO growth analysis

562   Growth of hROs was analyzed by measuring organoid area with ImageJ from brightfield images taken
563   at d42 (prior to dosing) and d54 (4 days post-dosing), from which % increase in area size calculated.
564
Statistics
All means are presented as mean ±SD; N denotes number of independent experiments (i.e.,
differentiation batches) and n denotes number of images or hROs examined, where appropriate.
Statistical testing was performed in GraphPad Prism (version 10.1.2). Tests used to analyze statistical
significance (normality testing, paired t-tests, one-way ANOVAs) are specified in figure legends.

Acknowledgements
This research was funded by Fight for Sight (5139/5140), and Medical Research Council UK
(MR/T002735/2). Maja Wojtynska was funded by the Wellcome Trust as part of the Advanced
Therapies for Regenerative Medicine Wellcome Trust PhD Program (218461/Z/19/Z). This work was
made possible by the support, dedication and teamwork of the Ocular Cell and Gene Therapy, with
particular thanks to Dr E West, Dr M. Branch, Dr M. Khazim, B. Ladino, J. Kapadia, E. Lanning, M.
Margari, M. Tariq, C. Mofidi, K. Kumar, S. Guilfoyle, S. van Heerden for help with stem cell maintenance
cultures. For the purpose of open access, the authors have applied a Creative Commons Attribution
(CC BY) licence to any Author Accepted Manuscript version arising.
579

580   **Author Contributions**
581   Conceptualization – PH, RP; Data curation – PH, MW; Formal Analysis – PH; Funding acquisition – RP;
582   Investigation – PH, RP; Methodology – PH; Supervision – RP; Visualization – PH; Writing – original draft
583   – PH; Writing – review & editing – PH, AS, RA, RP
584
585   **Declaration of Interests**
586   The authors declare no competing interests.

References

1. Voigt, A. P. *et al.* Human photoreceptor cells from different macular subregions have distinct
transcriptional profiles. *Hum. Mol. Genet.* **30**, 1543–1558 (2021).

2. Hoshino, A. *et al.* Molecular Anatomy of the Developing Human Retina. *Dev. Cell* **43**, 763-
779.e4 (2017).

3. Curcio, C. A., Sloan, K. R., Kalina, R. E. & Hendrickson, A. E. Human photoreceptor topography.
J. Comp. Neurol. **292**, 497–523 (1990).

4. Provis, J. M. & Hendrickson, A. E. The Foveal Avascular Region of Developing Human Retina.
Arch. Ophthalmol. **126**, 507–511 (2008).

5. Hendrickson, A. & Zhang, C. Development of cone photoreceptors and their synapses in the
human and monkey fovea. *J. Comp. Neurol.* **527**, 38–51 (2019).

6. Hendrickson, A. Development of Retinal Layers in Prenatal Human Retina. *Am. J. Ophthalmol.*
161, 29-35.e1 (2016).

7. Hendrickson, A., Possin, D., Vajzovic, L. & Toth, C. A. Histologic Development of the Human
Fovea From Midgestation to Maturity. *Am. J. Ophthalmol.* **154**, 767-778.e2 (2012).

8. Diaz-Araya, C. & Provis, J. M. Evidence of photoreceptor migration during early foveal
development: a quantitative analysis of human fetal retinae. *Vis. Neurosci.* **8**, 505–514 (1992).

9. Springer, A. D., Troilo, D., Possin, D. & Hendrickson, A. E. Foveal cone density shows a rapid
postnatal maturation in the marmoset monkey. *Vis. Neurosci.* **28**, 473 (2011).

10. Yuodelis, C. & Hendrickson, A. A qualitative and quantitative analysis of the human fovea
during development. *Vision Res.* **26**, 847–855 (1986).

11. Lu, Y. *et al.* Single-Cell Analysis of Human Retina Identifies Evolutionarily Conserved and
Species-Specific Mechanisms Controlling Development. *Dev. Cell* **53**, 473–491 (2020).

12. Mey, J. & Thanos, S. Development of the visual system of the chick: I. Cell differentiation and
histogenesis. *Brain Res. Rev.* **32**, 343–379 (2000).

13. Silva, S. da & Cepko, C. L. Fgf8 Expression and Degradation of Retinoic Acid Are Required for
Patterning a High-Acuity Area in the Retina. *Dev. Cell* **42**, 68–81 (2017).

14. Meyer, D. & May Jr, H. C. The Topographical Distribution of Rods and Cones in the Adult
Chicken Retina. *Exp. Eye Res.* **17**, 347–355 (1973).

15. Cvekl, A. & Wang, W.-L. L. Retinoic acid signaling in mammalian eye development. *Exp. Eye
Res.* **89**, 280–291 (2009).

16. Cunningham, T. J. & Duester, G. Mechanisms of retinoic acid signalling and its roles in organ
and limb development. *Nat. Rev. Mol. Cell Biol.* **16**, 110–123 (2015).

17. Sakai, Y., Luo, T., McCaffery, P., Hamada, H. & Dräger, U. C. CYP26A1 and CYP26C1 cooperate
in degrading retinoic acid within the equatorial retina during later eye development. *Dev.
Biol.* **276**, 143–157 (2004).

18. Thatcher, J. E. & Isoherranen, N. The role of CYP26 enzymes in retinoic acid clearance. *Expert
Opin. Drug Metab. Toxicol.* **5**, 875 (2009).

19. Nakajima, Y. Retinoic acid signaling in heart development. *Genesis* **57**, e23300 (2019).

20. Stratford, T., Horton, C. & Maden, M. Retinoic acid is required for the initiation of outgrowth
in the chick limb bud. *Curr. Biol.* **6**, 1124–1133 (1996).

21. Hernandez, R. E., Putzke, A. P., Myers, J. P., Margaretha, L. & Moens, C. B. Cyp26 enzymes
generate the retinoic acid response pattern necessary for hindbrain development.
Development **134**, 177–187 (2007).

22. Thompson, B. *et al.* Genetics and functions of the retinoic acid pathway, with special
emphasis on the eye. *Hum. Genomics* **13**, 1–15 (2019).

23. Lahne, M., Yoshimatsu, T. & MacDonald, R. B. Retinoic acid signaling in the developing
zebrafish high acuity zone. *Invest. Ophthalmol. Vis. Sci.* **64**, 3630–3630 (2023).

24. Kumar, S. & Duester, G. Retinoic acid controls body axis extension by directly repressing Fgf8
transcription. *Development* **141**, 2972–2977 (2014).

25. Cowan, C. S. *et al.* Cell Types of the Human Retina and Its Organoids at Single-Cell Resolution.

Cell **182**, 1623-1640.e34 (2020).

26.   Chen, R. Single Cell Multiome Atlas of the Human Developing Retina. *Res. Sq.* (2023).

27.   Krueger, M. R., Fishman-Williams, E., Simó, S., Tarantal, A. F. & La Torre, A. Expression
patterns of CYP26A1, FGF8, CDKN1A, and NPVF in the developing rhesus monkey retina.
Differentiation **135**, 100743 (2024).

28.   Gonzalez-Cordero, A. *et al.* Recapitulation of Human Retinal Development from Human
Pluripotent Stem Cells Generates Transplantable Populations of Cone Photoreceptors. *Stem*
Cell Reports **9**, 820–837 (2017).

29.   Li, L. *et al.* Electrophysiological characterization of photoreceptor-like cells in human inducible
pluripotent stem cell-derived retinal organoids during in vitro maturation. *Stem Cells* **39**, 959–
974 (2021).

30.   Pearson, R. A. *et al.* Restoration of vision after transplantation of photoreceptors. *Nature* **485**,
99–103 (2012).

31.   Kruczek, K. *et al.* Differentiation and Transplantation of Embryonic Stem Cell-Derived Cone
Photoreceptors into a Mouse Model of End-Stage Retinal Degeneration. *Stem Cell Reports* **8**,
1659–1674 (2017).

32.   Gonzalez-Cordero, A. *et al.* Photoreceptor precursors derived from three-dimensional
embryonic stem cell cultures integrate and mature within adult degenerate retina. *Nat.*
Biotechnol. **31**, 741–747 (2013).

33.   Ribeiro, J. *et al.* Restoration of visual function in advanced disease after transplantation of
purified human pluripotent stem cell-derived cone photoreceptors. *Cell Rep.* **35**, 109022
(2021).

34.   Shirai, H. *et al.* Transplantation of human embryonic stem cell-derived retinal tissue in two
primate models of retinal degeneration. *Proc. Natl. Acad. Sci.* **113**, E81–E90 (2016).

35.   Kim, S. *et al.* Generation, transcriptome profiling, and functional validation of cone-rich
human retinal organoids. *Proc. Natl. Acad. Sci.* **116**, 10824–10833 (2019).

36.   Capowski, E. E. *et al.* Reproducibility and staging of 3D human retinal organoids across
multiple pluripotent stem cell lines. *Development* **146**, dev171686 (2019).

37.   Crossley, P. H. & Martin, G. R. The mouse Fgf8 gene encodes a family of polypeptides and is
expressed in regions that direct outgrowth and patterning in the developing embryo.
Development **121**, 439–451 (1995).

38.   Soukkarieh, C., Agius, E., Soula, C. & Cochard, P. Pax2 regulates neuronal–glial cell fate choice
in the embryonic optic nerve. *Dev. Biol.* **303**, 800–813 (2007).

39.   Bringmann, A. *et al.* The primate fovea: Structure, function and development. *Progress in*
Retinal and Eye Research vol. 66 49–84 (2018).

40.   Napoli, J. L. Functions of Intracellular Retinoid Binding-Proteins. *Subcell. Biochem.* **81**, 21
(2016).

41.   Schubert, M. & Germain, P. Retinoic Acid and Retinoid X Receptors. *Cells* **12**, 864 (2023).

42.   Bumsted, K. & Hendrickson, A. Distribution and Development of Short-Wavelength Cones
Differ Between Macaca Monkey and Human Fovea. *J. Comp. Neurol.* **516**, 502–516 (1999).

43.   Luo, Z. *et al.* Islet1 and Brn3 Expression Pattern Study in Human Retina and hiPSC-Derived
Retinal Organoid. *Stem Cells Int.* **2019**, 1–14 (2019).

44.   Wohlschlegel, J. *et al.* ASCL1 induces neurogenesis in human Müller glia. *Stem Cell Reports*
18, 2400–2417 (2023).

45.   Hendrickson, A., Possin, D., Kwan, W. C., Huang, J. & Bourne, J. A. The temporal profile of
retinal cell genesis in the marmoset monkey. *J. Comp. Neurol.* **524**, 1193–1207 (2016).

46.   Hussey, K. A., Hadyniak, S. E. & Johnston, R. J. Patterning and Development of Photoreceptors
in the Human Retina. *Front. Cell Dev. Biol.* **10**, (2022).

47.   Alfano, G. *et al.* Vax2 regulates retinoic acid distribution and cone opsin expression in the
vertebrate eye. *Development* **138**, 261–271 (2011).

48.   Hadyniak, S. E. *et al.* Retinoic acid signaling regulates spatiotemporal specification of human

689   green and red cones. *PLoS Biol.* **22**, e3002464 (2024).

690 49. Liu, J. *et al.* Wnt/β-catenin signalling: function, biological mechanisms, and therapeutic
691   opportunities. *Signal Transduct. Target. Ther. 2021 71* **7**, 1–23 (2022).

692 50. Kubo, F., Takeichi, M. & Nakagawa, S. Wnt2b controls retinal cell differentiation at the ciliary
693   marginal zone. *Development* **130**, 587–598 (2003).

694 51. Meyers, J. R. *et al.* β-catenin/Wnt signaling controls progenitor fate in the developing and
695   regenerating zebrafish retina. *Neural Dev.* **7**, (2012).

696 52. Denayer, T. *et al.* Canonical Wnt Signaling Controls Proliferation of Retinal Stem/Progenitor
697   Cells in Postembryonic Xenopus Eyes. *Stem Cells* **26**, 2063–2074 (2008).

698 53. Liu, H., Mohamed, O., Dufort, D. & Wallace, V. A. Characterization of Wnt signaling
699   components and activation of the Wnt canonical pathway in the murine retina. *Dev. Dyn.*
700   **227**, 323–334 (2003).

701 54. Cho, S.-H. H. & Cepko, C. L. Wnt2b/beta-catenin-mediated canonical Wnt signaling
702   determines the peripheral fates of the chick eye. *Development* **133**, 3167–3177 (2006).

703 55. Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative CT method.
704   *Nat. Protoc.* **3**, 1101–1108 (2008).

705 56. Crossley, P. H., Martinez, S., Ohkubo, Y. & Rubenstein, J. L. R. Coordinate expression of Fgf8,
706   Otx2, Bmp4, and Shh in the rostral prosencephalon during development of the telencephalic
707   and optic vesicles. *Neuroscience* **108**, 183–206 (2001).

F**igures**

Figure 1. ***CYP26A1* shows biphasic expression localized to the macula of PCW6-17 developing human fetal**
retina, peaking at early PCW8 and PCW16/17.
Human fetal retina samples from PCW6-17 stained *in situ* using RNAscope™ for fibroblast growth factor 8 (*FGF8,*
cyan) and RA catabolizing enzyme *CYP26A1* (*magenta*), showing *CYP26A1*+ spot-like presumptive macular (PM)
region colocalizing with RPC (retinal progenitor cell) marker, VSX2 (*green*) in the neuroblastic layer between CS18-
23, and basal to the VSX2+ layer at PCW12-17. (a) Whole retinal section (20x), (b) High magnification of *CYP26A1*+
PM regions (40x). White arrows indicate PM region. Sections were counterstained with nuclear marker, DAPI
(*greyscale*). PCW – Post-conception Weeks. Images are representative of n≥2 samples. Scale bars - 500μm.

Figure 2. Cone progenitor markers are expressed in late PCW8 presumptive macula (PM), localized to the apical
layer by PCW10 but do not co-express *CYP26A1*, while macular Müller Glia cells express *CYP26A1* in PCW17
human fetal retina. (a), At late PCW8 (CS23), OTX2 (*yellow*) is expressed in cone/bipolar-biased RPCs in the
neuroblastic layer (NBL) throughout the retina, but at highest density in the PM, while no RECOVERIN (*red*) or
RXRγ+ (*blue*) cone precursor markers are visible. (b), At PCW10, the PM region expresses a thick layer of OTX2+
cells in the NBL, as well as RXRγ+/RECOVERIN+ cone precursors localized to a single cell layer at the apical region
of the central retina, not present in the peripheral retina. *CYP26A1* expression is restricted to the neuroblastic
layer, not expressed in the RECOVERIN+/ RXRγ+/OTX2+ cone precursors. (c) At PCW17, *CYP26A1* (*magenta*)
localizes to the VSX2+ (*green*)/SOX9+ (*yellow*) developing Müller glia layer, basal to the VSX2+/SOX9- developing
bipolar cell layer. Müller glia cytoskeletal marker Vimentin (*VIM, red*) is clearly expressed, while no *FGF8* (*cyan*)
is visible in the PM. Note stains for OTX/RECOVERIN/RXRγ and VSX2/*CYP26A1* were each performed on
consecutive sections from the same respective retina for (a) and (b), and for SOX9/VIM and *CYP26A1/FGF8* in (c).
Sections were counterstained with nuclear marker, DAPI (*greyscale*). PCW – Post-conception Weeks; ONL – Outer
Nuclear Layer; INL – Inner Nuclear Layer; GCL – Ganglion Cell Layer. Images taken at 20x (ai/bi)/40x (aii-iii/bii-
iii/c) magnification. Images are representative of n≥2 samples. Scale bars – 500μm for whole retina, 200μm for
(a-b) ROIs, 100μm for (c) ROIs.

Figure 3. *CYP26A1* is highly expressed in the presumptive macula (PM) of early PCW8 (CS20) human fetal
retina, while *FGF8* is localized exclusively to the optic nerve head (ONH). Representative images of CS20 fetal
retina stained *in situ* for *CYP26A1* (*magenta*) and *FGF8 (cyan)*, showing strong expression of *CYP26A1* in the PM
colocalizing with RPC marker VSX2 (*green*), but no *FGF8*, which is only visible in RPCs around the ONH. Assay
negative control (*DapB, cyan*) and positive control (Cy5 *PPIB*, Cy3 *POLR2A, magenta*) also shown. Sections were
counterstained with nuclear marker, DAPI (*greyscale*). PCW – Post-conception Weeks; ONL – Outer Nuclear Layer;
INL – Inner Nuclear Layer; GCL – Ganglion Cell Layer. Images taken at 40x magnification Images are representative
of n=3 CS20/21 samples. Scale bars, 100μm.

Figure 4. Higher proliferation in the peripheral retina compared with the central region results in a temporally
shifted *CYP26A1*+ presumptive macula (PM) at early developmental stages (PCW7-10).
(a), (i-iii) The distance of the *CYP26A1*+ (*magenta*) PM (white arrow) to the peripheral edge of the retina (IV)
increases at a significantly higher rate (over 2-fold) between PCW7-17 compared with the central region between
the PM and the optic nerve head linear regression slope=114 and 62 respectively, *p*<0.001. (v), The distance
between the macula to periphery as a proportion of the whole retina increases over PCW7-17 (from 20%-45%),
while the distance between the macula and optic nerve remains almost constant as a proportion of the whole
retina (30%), due to the slower growth of the central compared with peripheral region (linear regression
slope=0.0033/-0.0004 respectively). n=14, linear regression with 95% confidence internals plotted. (b), (i)
Representative images of proliferative marker KI67 (*yellow*) staining in CS18-23 human fetal retinas shows: (ii)
the proportion of cells expressing KI67 was at nearly two-fold higher in the peripheral retina compared with the
central region during CS18-23 at all timepoints (CS18: *t*(2)=3.3, *p*=0.08; CS20: *t*(2)=6.4, *p*=0.02; CS23: *t*(2)=2.5,
p=0.13). (iii) The proportion of proliferating cells decreases significantly in both the macula and the equivalent

nasal region on the centre to peripheral gradient by CS20 (macula: $t(4)=6.50$, $p=0.003$; nasal: $t(4)=5.03$, $p=0.007$).
(iv) Linear regression shows that the decline in KI67+ cells in the macular region is significantly steeper than the
nasal region between CS18 and CS20, during which time *CYP26A1* expression is upregulated in the macula
(macula: -15.6, nasal: -8.7, $p=0.046$). Significance values determined by paired/unpaired t-tests: *p value<0.05,
**p value<0.01. n=3 samples/timepoint, data are represented as mean ±SD. Sections were counterstained with
nuclear marker, DAPI (*greyscale*). PCW – Post-conception Weeks. Images taken at 20x magnification. Scale bars
– 500µm.

Figure 5. Differentiating human retinal organoids express retinoic acid and FGF8 pathway components. (a)
Relative expression of RA synthesizing enzymes *ALDH1A1* and *ALDH1A3* mRNA across differentiation of hROs. (b)
immunostaining of ALDH1A1 and ALDH1A3 in d40 hROs showing expression throughout the organoid, co-stained
with neural retinal progenitor cell marker VSX2 (images are representative of n=3 hROs), counterstained with
DAPI (*greyscale*); (c-f) relative expression of (c) retinoic acid catabolizing enzyme *CYP26A1*, (d) *CRABP2,* a proxy
for circulating RA levels, (e) fibroblast growth factor 8 (*FGF8*) and (f) FGF receptors (*FGFR1/2/3/4*) mRNA. All data
shown as Log$_2$ fold change of CT values relative to d0 hESCs (5 pooled hROs per sample, n=3). Data normalized
to both *GAPDH* and *β-actin* housekeeping genes. Significance values determined by one-way repeated measures
ANOVA followed by Dunnett's test against control group, adjusting for multiple comparisons: *p value<0.05, **p
value<0.01, ***p value<0.001, data are represented as mean ±SD. Scale bars - 100µm.

Figure 6. Dosing early hROs with RA receptor inhibitor AGN193109 results in reduced growth due to reduced
proliferation and increased cone precursor marker OTX2 expression. (a), hRO area was measured at d42,
prior to dosing, and no significant difference in size was found ($t(33)=1.64$, $p=0.11$). However, following
dosing between d42-50, organoid size was measured at d54, and the % change in area was found to
be significantly lower in AGN193109 dosed hROs compared with DMSO dosed controls, demonstrating
reduced growth in RA inhibited hROs ($t(32)=5.18$, $p<0.0001$). (b), Immunostaining for (i) apoptotic
marker CASPASE-3 (*yellow*) at d60 showed no significant increase in cell death between AGN193109
dosed and control hRO ($t(29)=0.64$, $p=0.5$), (ii) while staining for proliferative marker KI67 (*green*)
showed a significant reduction in cell proliferation in RA inhibited hROs ($t(24)=2.48$, $p=0.02$). (c) (i)
Dosing with AGN193109 led to significant increases in photoreceptor marker *RECOVERIN* ($t(5)=2.56$,
p<0.05), and cone progenitor marker *OTX2* ($t(4)=4.58$, p=0.01), while early cone precursor markers
RXRγ and *THRb* show trended towards increased expression ($t(5)=1.45$, $p=0.2$)/($t(5)=1.69$, $p=0.15$), and
no increase in retinal progenitor cell (RPC) marker *VSX2* was observed ($t(5)=1.39$, $p=0.2$). (ii) However,
dosing with rFGF8 had no effect on any marker expression tested (*RECOVERIN*: $t(2)=0.22$, $p=0.85$;
OTX2: $t(2)=2.99$, $p=0.10$; *RXRγ*: $t(3)=0.23$, $p=0.83$; *THRb*: $t(2)=1.01$, $p=0.40$; *VSX2*: $t(3)=0.39$, $p=0.72$).
(d) Immunostaining of dosed and DMSO dosed control hROs at d60 showing a significant increase in
proportion of cone progenitor marker OTX2+ (blue) cells in AGN193109 dosed hROs ($t(28)=2.4$,
$p=0.02$). All qRT-PCR data shown as Log2 fold change of CT values relative to the mean of the DMSO
treated controls (5 pooled hROs per sample, n≥3, lines indicate paired samples of DMSO/AGN-dosed
samples from the same batch of differentiation to control for batch-batch variation.). All qRT-PCR data
normalized to both *GAPDH* and *b-actin* housekeeping genes. Significance values determined by
paired/unpaired t-tests: *p value<0.05, n ≥13, data are represented as mean ±SD. Images
representative of n≥13 hROs, counterstained with DAPI (*greyscale*). Scale bars - 200um.

Figure 7. Dosing late hROs with RA receptor inhibitor AGN193109 results in increased M/L-OPSIN and reduced
S-OPSIN expression, indicating shift in cone subtype specification towards fovea-like composition. Dosing with
RA receptor inhibitor AGN193109 early in differentiation (between d42-d50), mid (d70-80) and late (d120-130)
resulted in (a) no change in mRNA expression at d150 of (i) pan-cone marker *ARRESTIN-3* ($t(2)=0.82$, $p=0.5$;
$t(3)=0.43$, $p=0.88$; $t(3)=1.22$, $p=0.46$), or (ii) rod marker *RHODOPSIN* ($t(2)=1.03$, $p=0.41$; $t(3)=0.65$, $p=0.76$;

t(3)=2.43, p=0.15), while early+mid and early+mid+late dosing resulted in upregulation of mRNA for (iii)
green/red cone gene *M/L-OPSIN* approaching significance (t(3)=1.70, p=0.29/t(3)=2.37, p=0.16), and significant
reduction in (iv) blue cone gene *S-OPSIN* (t(3)=1.67, p=0.30; t(3)=5.64, p=0.02). Both (v) *M-OPSIN* (t(3)=1.27,
p=0.29/t(3)=1.53, p=0.22) and (vi) *L-OPSIN* (t(3)=1.51, p=0.23/t(3)=2.70, p=0.07) showed a trend towards
increasing for early+mid/early+mid+late dose, with early+mid+late for *L-OPSIN* approaching significance. (b) no
change in proportion of cells expressing protein (i) ARRESTIN-3 (t(11)=1.29, p=0.22); (ii) RHODOPSIN (t(11)=0.41,
p=0.69); or proportion of ARRESTIN-3+ cells expressing (iii) M/L-OPSIN (t(11)=0.14, p=0.89); but a significant
decrease in the proportion of ARRESTIN-3+ cells expressing (iv) S-OPSIN (t(11)=3.14, p=0.01). (c) Early, middle
and late dosing resulted in an increase in *SFRP1* expression at d150, although this only reached statistical
significance for triple-dosed hROs (t(2)=2.02, p=0.18; t(3)=2.45, p=0.15; t(3)=5.04, p=0.02). (c) Additionally, early
dosing with AGN193109 led to significant increase in *SFRP1* by d60 of differentiation (t(4)=4.50, p=0.01). All qRT-
PCR data shown as Log_2 fold change of CT values relative to the mean of DMSO treated controls (3-5 pooled hROs
per sample, n≥3, lines indicate paired samples of DMSO/AGN-dosed samples from the same batch of
differentiation to control for batch-batch variation.). All qRT-PCR data normalized to both *GAPDH* and *β–actin*
housekeeping genes. Significance values determined by paired/unpaired t-tests/one-way repeated measures
ANOVA followed by Dunnett's test against control group, adjusting for multiple comparisons: *p value<0.05, **p
value<0.01, data are represented as mean ±SD. Staining data representative of n>5 hROs, counterstained with
DAPI (*greyscale*). Scale bars - 500μm.

 Supplementary Tables

ALDH1A1	Rabbit	Abcam	ab23375	AB_2224009	1:200
ALDH1A3	Rabbit	Abcam	ab129815	AB_2937054	1:200
ARRESTIN-3	Goat	Novus	NBP1-37003	AB_2060085	1:100
CASPASE-3	Rabbit	Abcam	ab2302	AB_302962	1:100
CRX	Mouse	Abnova	H00001406-M02	AB_606098	1:800
KI67	Rabbit	Abcam	ab15580	AB_443209	1:100
M/L-OPSIN	Rabbit	Millipore	AB5405	AB_177456	1:100
NRL	Goat	R+D systems	AF2945	AB_2155098	1:100
ONECUT1	Mouse	Santa cruz	sc-13050	AB_2251852	1:100
OTX2	Goat	R&D Systems	AF1979	AB_2157172	1:50
RBPMS	Rabbit	PhosphoSolutions	1830	AB_2492225	1:150
RECOVERN	Rabbit	Chemicon	AB5585	AB_2253622	1:1000
RHODOPSIN	Mouse	Sigma	O4886	AB_260838	1:1000
RXRγ	Mouse	Santa Cruz	sc-365252	AB_10850062	1:200
S-OPSIN	Rabbit	Millipore	AB5407	AB_177457	1:200
SOX9	Rabbit	Sigma	AB5535	AB_2239761	1:200
VIM	Chicken	Novus	NB300	AB_922758	1:600
VSX2	Mouse	Santa Cruz	sc-365519	AB_10842442	1:200

 Supplementary Table 1 – Primary antibodies

Gene	Forward	Reverse	Universal Probe
AIPL1	GGAGAGGGAAATCGGCTCTT	TTCTCCTTGGTCTGCAGGTT	79
ALDH1A1	GTTAGCTGATGCCGACTTGG	CGCTCAACACTCCTTCGAAC	81
ALDH1A3	AAGAGGAGATTTTCGGGCCA	ACTTCAGGGCTTTGTCGAGA	66
ARR3	CACGGAGACTGTAGCTGCTA	GCTAGAGGCCAGGTTGGTAT	14
β-ACTIN	GCACCCAGCACAATGAAGAT	TCGTCATACTCCTGCTTGCT	63
CRX	GAGGCTGTGTCTTGTGCAAA	TCGAAGTCTTACCCCATCGG	12
CRABP2	CCCTGTAAGAGCCTGGTGAA	CCGTCATGGTCAGGATCAGT	93
CYP16A1	GACATGCAGGCACTAAAGCA	GGTAGAGCCCCAGGTAAGTG	5
FGF8	GAGACGGGCCTCACATCTG	TGTACCAGCCCTCGTACTTG	67
FGFR1	CGGGACAGACTGGTCTTAGG	GGTTGGGTTTGTCCTTGTCC	53
FGFR2	CAGTCATCCTGTGCCGAATG	AACTGTTACCTGTCTCCGCA	71
FGFR3	TACTCCTTCGACACCTGCAA	ACTTCTGGGAGGCCAAGTAC	68
FGFR4	GGAGTCCCGGAAGTGTATCC	GTAGTCAATGTGGTGGACGC	71
GAPDH	TTGCCCTCAACGACCACTTT	TGGTCCAGGGGTCTTACTCC	77
ISLET1	TCTCCGGATTTGGAATGGCA	CATTTGATCCCGTACAACCTGAT	-
L-OPSIN	TGCATCATCCCACTCGCTAT	TTCCTTCTCTGCCTTCTGGG	5
M-OPSIN	CTACCTCCAAGTGTGGCTGG	GAATGCCAGGACCATCACCA	13
M/L-OPSIN	CTTCCACCCTTTGATGGCTG	AGTTTCGAAACTGCCGGTTC	-
NRL	AGAGCTGTGTGCAGAAGTCT	GCAGAAGTCGTCCAATCCAC	95
ONECUT1	CAGCGTCGAACTCTACATGC	ACTCCTCCTTCTTGCGTTCA	3
OTX2	ATTGCTAGAGCAGCCCTCAC	GGAGCAGTGGAACTTACAGC	-
RBPMS	CACTGCATGCCCAGATGC	CCCACAAGACAGATTGCAGC	-
RCVRN	GCTTCGATTCCAACCTCGAC	TTACCGTCCACGTCGTAGAG	78
RHODOPSIN	TGGGCCATTAAAAGCTCAGC	GTGGAAGCTGGCAGTTTCAA	77
RXRγ	CATGAAGAGGGAAGCTGTGC	TTCTGCCTCACTCTCAGCTC	82
SFRP1	GCCCGAGATGCTTAAGTGTG	GACACACCGTTGTGCCTTG	-
S-OPSIN	GCACTGTAGCAGGTCTGGTT	CAATGGTCCAGGTAGCCAGG	72
THRβ	ATTTCGTTCTCCGGTCCTGT	TCAAGTCTTGGACCAGGGAC	9
VSX2	TGGGGATGCACAAAAAGTCG	GCTCCATCTTGTCGAGCTTG	69

829   **Supplementary Table 2 – Primers**

    Supplementary Figures

Supplementary Figure 1. Serial sections of a single human fetal retina (CS20) show expression of *CYP26A1* is
localized to a single spot-like region at the presumptive macula (PM), while *FGF8* is exclusively expressed
around the optic nerve head. (a) Representative images of serial sections through a CS20 human retina, taken
at 100μm intervals and stained for the RA catabolizing enzyme, *CYP26A1* (*magenta*), and growth factor *FGF8*
(*cyan*) mRNA using RNAscope™. Imaging shows expression of *CYP26A1* in the PM with no overlapping *FGF8*
expression, which is instead expressed around the optic nerve head. White arrows indicate PM, while white
outlined arrow heads indicate low levels of *CYP26A1* expression visible around optic nerve head (ONH) and at
nasal edge of the retina. Sections were co-stained with RPC marker, VSX2 (*green*) and counterstained with
nuclear marker, DAPI (*greyscale*). S – Superior; I – Inferior; PCW – Post-conception Weeks. Images taken at 20x
magnification. Images are representative of n=3 CS20/21 samples. Scale bars - 500μm.

Supplementary Figure 2. Proliferation in human developing retinal regions (CS18-23)
(a), Masking of macular (based on *CYP26A1+* staining), central (adjacent to the optic nerve), nasal (equivalent
distance as macula to periphery on the central-peripheral gradient on the nasal side) and peripheral regions in
Imaris 10.0. (b) Proportion of KI67+ cells in all analyzed developing retinal regions from CS18-23, showing the
macula has a similar proportion of KI67+ cells to the peripheral retina at CS18 (mean of 65/66% respectively,
$t(2)=0.33$, $p=0.78$)), and significantly different to central/nasal regions ($t(2)=2.24$, $p=0.16$; $t(2)=2.05$, $p=0.18$),
while at CS20/23, the macula had similar proportion of KI67+ cells to the central retina (CS20: means of 34/36%,
$t(2)=1.15$, $p=0.37$); CS23: means of 24/22%, $t(2)=0.38$, $p=0.74$) and significant/nearing significant difference in
proliferating cells compared with nasal/peripheral retina (CS20: $t(2)=5.26$, $p=0.03$/$t(2)=5.92$, $p=0.02$; CS23:
$t(2)=2.47$, $p=0.13$/$t(2)=2.76$, $p=0.11$ respectively. (c) Linear regression of proportion of KI67+ cells in different
retinal regions between CS18 and CS20 shows that the macular region has a steeper decline in KI67+ cells than
any other retinal region (macula: -15.62, central: -5.5, nasal: -8.7, peripheral: 0.8, $p=0.01$). n=3/timepoint, data
are represented as mean ±SD.

Supplementary Figure 3. Temporal and spatial photoreceptor development in human retinal organoids.
(a) graphical representation of hRO differentiation and "early" dosing with RA receptor inhibitor AGN193019
(10μM) and recombinant FGF8 (100ng/ml) between d42-50, followed by "mid" AGN193109 pulse at d70-80 and
"late" pulse at d120-130; (b-d) Relative mRNA expression of (b) RPC cell marker *VSX2* showing upregulation at
d40 of differentiation compared with d0, (c) photoreceptor precursor markers, *RECOVERIN*, *CRX* and *AIPL1*
showing photoreceptor specification from d40 of differentiation with expression further increasing to d90 and
(d) cone precursor markers *RXRγ*, *THRβ* and *ONECUT1* significantly upregulated by d40, with plateauing
expression from d60-70, indicating specification of cone precursors by this timepoint; (e) immunostaining of
VSX2+ retinal progenitor cells in the neuroblastic layer (NBL) from d40 of differentiation, along with retinal
ganglion cell (RGC) marker RBPMS in the inner nuclear layer (INL) from d60, OTX2+ cone-biased progenitor cells
showing increasing cell numbers from d50, and RECOVERIN+ photoreceptor precursors from d70 predominantly
localizing to the outer layers of developing hROs, and ONECUT1+ horizontal cell-biased retinal progenitors
towards the inner layers of the organoid, alongside some OTX2+/RECOVERIN+ bipolar cells/mis-localized
photoreceptor precursor cells; (f) Relative mRNA expression of mature cone markers *ARR3, OPSNSW, OPSNMW*
and *OPSNLW* and (g) mature rod markers *RHODOPSIN*, *NRL* and photoreceptor marker *RECOVERIN* showing
maturation of rods and cones by d150 of differentiation; (h) immunostaining of mature d150 retinal hROs stained
for pan-photoreceptor marker RECOVERIN, with tight multilayered NRL+ rod nuclei in the outer nuclear layer
(ONL), RHODOPSIN labelling the outer segments of rod photoreceptor cells and ARR3 labelling the cell bodies of
a single apical layer of cone photoreceptor cells showing the inner segments, with a small number of S-OPSIN+
short wavelength cones, and M/L-OPSIN+ medium/long wavelength cones, with some RECOVERIN+
photoreceptors displaced to the INL. All qRT-PCR data shown as Log_2 fold change of CT values relative to d0 hESCs
(3-5 pooled hROs per sample, n=3). qRT-PCR data normalized to both *GAPDH* and *β-actin* housekeeping genes.
Significance values determined by one-way repeated measures ANOVA followed by Dunnett's test against control

group, adjusting for multiple comparisons: *p value<0.05, **p value<0.01, ****p value<0.0001, data are
represented as mean ±SD. Staining data representative of n=3 hROs, counterstained with DAPI (*greyscale*). Scale
bars - 100μm.

Supplementary Figure 4 – Dose optimization of AGN193109 and recombinant FGF8 for hROs based on
alamarBlue cell viability assay. Fluorescence was measured at 590nm (RFU) and change in reduction at d4 and
d8, compared with d0 of dosing for each hRO, then relative change of treated compared with DMSO treated was
calculated and plotted for (a) AGN193109 and (b) recombinant FGF8 (rFGF8) (n=5). (c) Dosing with rFGF8between
d40-d50 lead to no significant change in *FGF8,* which is known to regulate its own expression Data shown as Log_2
fold change of CT values relative to PBS treated controls (5 pooled hROs per sample, n=4). Data normalized to
both *GAPDH* and *β–actin* housekeeping genes. Error bars - standard deviation. Significance values determined
by one-way repeated measures ANOVA followed by Dunnett's test against control group, adjusting for multiple
comparisons. *p value<0.05.

Supplementary Figure 5. Dosing early hROs with RA receptor inhibitor AGN193109 does not alter retinal
ganglion cell (RGC) marker or cone precursor marker expression at d60. (a-b) Dosing with RA receptor inhibitor
AGN193109 between d40-d50 leads to (a) a significant reduction in *CRABP2,* a proxy for circulating RA levels,
(t(2)=5.77, p=0.03), (b) but no significant change in RGC marker *RBPMS* (t(4)=1.02, p=0.37) or
RGC/bipolar/amacrine cell marker *ISLET1* (t(4)=0.30, p=0.78). Lines indicate paired samples of DMSO/AGN-dosed
samples from the same batch of differentiation to control for batch-batch variation. All data shown as Log_2 fold
change of CT values relative to the mean of DMSO treated controls (5 pooled hROs per sample, n≥3, lines indicate
paired samples of DMSO/AGN-dosed samples from the same batch of differentiation to control for batch-batch
variation). All qRT-PCR data normalized to both *GAPDH* and *β–actin* housekeeping genes. Significance values
determined by paired t-tests. (c) Immunostaining of AGN193109 dosed and DMSO dosed control hROs at d60
showing no significant change in the proportion of (i) RGC marker RBPMS *(yellow)* cells (t(30)=0.08, p=0.94) (ii)
cone precursor marker RXRg (*red*) (t(26)=0.96, p=0.35) or (iii) RPC marker VSX2 (staining in Figure 6) (t(29)=0.01,
p=0.99). Significance values determined by unpaired t-tests: *p value<0.05, n>13, data are represented as mean
±SD. Staining data representative of n>13 hROs, counterstained with DAPI (*greyscale*). Scale bars - 100μm.

Supplementary Figure 6 – Masking of necrotic core for CASPASE-3 staining quantification in IMARIS 10.0.

a

i Macula-Periphery

ii Macula-Nerve

iii Total Retina

iv

Regional Foetal Retina Growth

Y = 114.2*X − 4939
R² = 0.9645

Y = 61.96*X − 2176
R² = 0.9386

X axis: Days post-conception
Y axis: Distance [μm]

● Macula-Periphery
▲ Macula-Nerve

v

Regional Proportion of Total Retina

Y = 0.003303*X + 0.08772
R² = 0.9262

Y = −0.0004306*X + 0.3257
R² = 0.04464

X axis: Days post-conception
Y axis: Proportion of total retina length

● Macula-Periphery
▲ Macula-Nerve

b

i KI67/DAPI

CS18   CS20   CS23

Periphery

Centre

Macula

Nasal

ii **KI67 Expression in Central v Peripheral Developing Retina**

X axis: Carnegie Stage (18, 20, 23)
Y axis: % of KI67+ cells

● Periphery
■ Centre

iii **KI67 Expression in Macular v Nasal Developing Retina**

**
**
**
*

X axis: Carnegie Stage (18, 20, 23)
Y axis: % of KI67+ cells

● Macula
▲ Nasal

Linear regression of KI67 Expression in Macular v Nasal Developing Retina

iv

Y = −15.62*X + 346.2
R² = 0.9134

Y = −8.655*X + 212.0
R² = 0.8634

X axis: Carnegie Stage (18, 20)
Y axis: % of KI67+ cells