SENP2-based N-terminal truncation of α-synuclein in Lewy pathology propagation

Katsutoshi Taguchi[1]*, Yoshihisa Watanabe[2], and Masaki Tanaka[1]*

[1]Department of Anatomy and Neurobiology, Graduate School of Medical Science, Kyoto

Prefectural University of Medicine, Kawaramachi-Hirokoji, Kamikyo-ku, Kyoto 602-8566,

Japan

[2]Department of Basic Geriatrics, Graduate School of Medical Science, Kyoto Prefectural

University of Medicine, Kawaramachi-Hirokoji, Kamikyo-ku, Kyoto 602-8566, Japan

*Corresponding author:

Katsutoshi Taguchi, Ph.D.

Department of Anatomy and Neurobiology, Graduate School of Medical Science, Kyoto

Prefectural University of Medicine, Kawaramachi-Hirokoji, Kamikyo-ku, Kyoto 602-8566,

Japan

Tel: +81-75-251-5301; Fax: +81-75-251-5306; E-mail: ktaguchi@koto.kpu-m.ac.jp

Masaki Tanaka, M.D., Ph.D.

Department of Anatomy and Neurobiology, Graduate School of Medical Science, Kyoto

Prefectural University of Medicine, Kawaramachi-Hirokoji, Kamikyo-ku, Kyoto 602-8566,

Japan

Tel: +81-75-251-5300; Fax: +81-75-251-5306; E-mail: mtanaka@koto.kpu-m.ac.jp

Summary

α-Synuclein (αSyn) is a major component of Lewy bodies (LBs) and Lewy neurites (LNs) which are pathological features of Parkinson's disease (PD) and Dementia with Lewy bodies. In the PD brain, with disease progression, LB/LN formation is propagated from the lower brainstem to the cerebral cortex. Prion-like cell-to-cell seed-transmission has been implicated as an underlying mechanism for Lewy-pathology propagation. However, the biochemical properties and production mechanism of those pathogenic seeds are unelucidated. In this study, we ascertained that the seeds released from pathological neurons that harbour LB/LN-like aggregates have the N-terminally truncated form of αSyn. This N-terminal truncation is directly catalysed by SENP2, which is a well-known deSUMOylation enzyme. After SENP2 processing of recombinant αSyn, the SDS-resistant high-molecular oligomer formation was promoted *in vitro*. Inhibition of SENP2 activity suppressed aggregate formation and propagation in cultured neurons and mouse brains. Thus, SENP2 might be a novel therapeutic target in LB diseases.

Introduction

α-Synuclein (αSyn) is one of the major constituents of Lewy bodies (LBs) and Lewy neurites

(LNs) that constitute the well-known pathological hallmarks of synucleinopathies, including

Parkinson's disease (PD) and dementia with Lewy bodies (DLB)[1-3]. In the sporadic PD brain,

the propagation of LB/LN formation indicates a caudorostral process that is associated with

disease progression from the lower brainstem, through the basal midbrain and forebrain, into

the cerebral cortex[4,5]. Interestingly, LBs/LNs were observed even in the grafted dopaminergic

neurons derived from foetal mesencephalic tissue that has been transplanted into the striatum

of a patient with PD[6]. This suggests the presence of prion-like propagation as the underlying

mechanism in neurodegeneration[7]. In several experimental models, inoculation of preformed

fibrils (PFFs) prepared from recombinant αSyn or PD brain-derived tissue extracts into the

mouse striatum can induce LB/LN-like aggregate formation and propagation through the

anatomical connections in the synaptic contacts within the brain[8-10].

   Abnormal intracellular aggregates, such as LBs and LNs, are formed by the recruitment of

intrinsic soluble αSyn into the insoluble aggregate core (the so-called "seeds")[7,11]. Therefore,

the endogenous expression of αSyn as well as seeds, which accelerate the polymerization of

soluble αSyn, are required for the formation of aggregates. Next, the cell-to-cell transmission

of seeds is a crucial step for the propagation of Lewy pathology. However, the biochemical

properties and mechanism of production of these pathogenic seeds that are released from

pathological neurons, which harbour LBs/LNs, are unelucidated.

The evidence from recent studies implicates posttranslational modifications (PTMs) of

αSyn, such as phosphorylation, glycosylation, acetylation, ubiquitination, SUMOylation, and

truncation, in the pathophysiological propensities of αSyn[12,13]. By using histological methods

and mass spectrometric analysis, the presence of various PTMs of αSyn has been detected in

the PD brain. Phosphorylation at the Ser129 residue is a well-known PTM of

αSyn-comprising LBs/LNs and is used as a marker for the detection of these abnormal

aggregates[14]. Furthermore, ubiquitination and SUMOylation play important roles in the

regulation of αSyn's stability and aggregation ability[15,16]. The truncation of αSyn has

significant effects on the aggregation ability, cross-seeding kinetics, and fibril stability of the

resultant species[12,17,18]. Interestingly, there are some characteristic differences between the

C-terminal- and N-terminal-truncated forms of the αSyn[17,19], wherein the former shows

higher aggregation ability and the latter shows lower aggregation ability and fibril stability

with higher cross-seeding activity in cultured cells and the mouse brain. The injection of

PFFs prepared from the N-terminal 10-residue- or 30-residue-truncated αSyn, rather than

PFFs prepared from full-length αSyn, into the mouse striatum induces more abundant

aggregate formation[17], which, given the consensus on the inverse correlation of seeding

activity with fibril stability in the field of prions, possibly contributes to the lower stability of

N-terminal-truncated αSyn-derived PFFs[20]. Among the various αSyn-processing enzymes

that have been identified thus far, caspase-1 and asparagine endopeptidase are C-terminal

cleavage enzymes, whereas 20S proteasome, calpain-1, cathepsins, and matrix

metalloproteases can truncate both C-terminal and N-terminal regions, including the NAC

region of αSyn[12]. Furthermore, PFFs are processed by lysosomal cathepsin B, and this partial

cleavage enhances the seeding ability of PFFs[21].

Studies using mass spectrometric analysis have identified many cleavage sites within the

tissue extracts obtained from the PD brain[12]. For example, calpain-mediated C-terminal

truncation at 122/123 residues of αSyn has been detected in LBs/LNs[22,23]; however, the

relationship between αSyn truncation and the biochemical properties of the seeds remains

elusive.

In this study, we explored this relationship through the characterization of

seed-constituting αSyn released from PD-model neurons containing pathological αSyn

aggregates and identified novel cleavage sites in the N-terminal region.

Results

Isolation and characterization of cell-to-cell transmissible seeds

To isolate the cell-to-cell transmissible seeds released from pathological neurons that harbour

LB/LN-like abnormal aggregates, the oligomeric species of αSyn within the conditioned

medium of those pathological neurons were screened using sucrose density-gradient

ultracentrifugation and Western blotting analysis (Figure S1A–C and Methods). Accordingly,

140-kDa αSyn oligomers, separated by native-PAGE, were recovered from the interface

fraction between the concentrated conditioned medium and the 0.8 M sucrose-containing

PBS layer. Moreover, we confirmed that the LB/LN-like aggregate formation in the primary

cultured neurons was induced by the fraction-containing fresh medium (Figure S1D).

To more precisely isolate the seeds and obviate PFF contamination, isolation was achieved

with microfluidic devices (Figure 1A, B; C1 and C3 chambers), wherein seeds released from

pathological neurons were recovered from the PFF-free culture medium. Furthermore, using

this microfluidic method, 140-kDa αSyn oligomers that were separated by native-PAGE were

recovered from the interface fraction as well as the results obtained by the batch method

(Figure 1C–E). Atomic force microscopy showed that this fraction contained a uniform

population of granular molecules with a diameter of 36 ± 1.2 nm (Figure 1F, G). Next, to

further confirm that these granular molecules are composed of αSyn, these molecules were

detected by several antibodies against αSyn. For αSyn detection by immunoelectron

microscopy, the reactivity of secondary antibodies was optimized using PFFs as the positive

control (Figure S1G). These granular molecules (diameter 35 ± 2.3 nm) detected by electron

microscopy were composed of αSyn and thus labelled by certain monoclonal antibody 5G4,

but not by the O2 antibody (Figure 1H–J). 5G4 recognizes protofibrils or large oligomeric

species of αSyn[28]. To determine the molecular weight of the intact seed structure,

size-exclusion chromatography (SEC)-high-performance liquid chromatography (HPLC)

analysis was performed (Figure S1E, F). The seed structure detected with immunoelectron

microscopy was recovered in the SEC Fraction No. 8 (corresponding to approximately 1000

kDa), as determined using a high-molecular-weight standard kit. These results indicated that

seeds of approximately 1000-kDa size were produced in the pathological neurons and

subsequently released into the culture medium.

Seeds contain truncated forms of αSyn

To investigate whether seeds contained the truncated αSyn or not, the cleavage sites within

αSyn were screened using mass spectrometric analysis. As a large amount of seed protein

was required for identification of the cleavage sites, a large microfluidic device was

developed (Figure S2). Using immunoprecipitation, samples for mass spectrometric analysis

were prepared from seed fraction (Figure 2A). PFF-constituting αSyn, which was derived

from the human αSyn sequence, was prepared as a control sample for mass spectrometric

analysis because results of the artificially cleaved sites detected in PFF-constituting αSyn

were subtracted from those of the cleaved sites detected in the seed-constituting αSyn.

Protein bands, Nos. 1–5, and control bands were subjected to mass spectrometric analysis

(Figure 2B), and the results suggested the presence of 6 cleavage sites (Figure 2C). The

C-terminal cleavage site at the 119/120 residue is a 20S proteasome-processing site[24]. To our

knowledge, cleavage sites detected at the N-terminal and NAC regions have not been

reported previously. Interestingly, conserved cleavage sites (KX|KXGV), the N-terminal

sides of lysine residues located at the N-terminal repeat regions (1–4) of αSyn, were detected

(Figure 2C). Although the cleavage site at repeat 4 was detected in the results of the control

sample in the mass spectrometric analysis, this site was considered as a possible cleavage site,

because the primary amino acid sequence is completely conserved at repeat 3. These results

suggested that the N-terminus of the seed-constituting αSyn was partially processed by

unknown N-terminal lysine endopeptidase activity. This information on the conserved

processing sites is useful for identifying the processing enzyme. Therefore, we focused on

these N-terminal cleavage sites of αSyn.

αSyn is processed by SENP2

Based on the results of mass spectrometric analysis, candidate enzymes that cleave the

N-terminal side of lysine residues located at the N-terminal repeat regions of αSyn

(conserved cleavage sites in Figure 2C) were screened using the Peptidase database,

MEROPS, which showed that carboxypeptidase B, USP7, and SENP2 were candidate

enzymes that can process the N-terminal side of the lysine residues. After screening these

enzymes, each enzyme's inhibitors were selected (Table 1). To examine the inhibitory effect

of the inhibitors on aggregate formation, primary neurons were cultured and treated with

PFFs in the presence of each inhibitor. In the inhibitor assay, NSC632839 could significantly

inhibit the aggregate formation (Figure 3A–F), whereas the reagent inhibits the enzymatic

activity of not only USP7 and SENP2 but also USP2 and UCHL1[25,26]. Therefore, to examine

whether αSyn is directly processed by these enzymes, protease assays were performed using

recombinant active enzymes. Recombinant αSyn was cleaved by SENP2, and the resulting

fragment was detected clearly (Figure 3G). However, cleaved fragments of αSyn were not

detected in the presence of USP7, USP2, and UCHL1 (Figure S4A–C). SENP2 preferentially

cleaved the soluble form of αSyn, rather than PFFs. The cleaved fragment of αSyn increased,

depending on the SENP2 dose (Figure 3H, I). Interestingly, SDS-resistant

high-molecular-weight signals of αSyn were increased by high-dose SENP2 treatment. This

cleavage pattern of αSyn by SENP2 was diminished by the insertion of mutations at the

predicted cleavage sites, as represented in Figure 2D (Figure 3J).

SENP2 was ordinarily expressed in the nucleus[27]. After aggregate formation, SENP2 was

partially distributed in the aggregates (Figure 4). In cultured neurons harbouring

phosphorylated-αSyn aggregates induced by PFFs, SENP2 was partially detected in those

aggregates (Figure 4A). In the human PD brain, SENP2 was observed in the amorphous

aggregates composed of phosphorylated αSyn, but not in typical LBs, which are spheroidal

(Figure 4B and Figure S5C, D).

Inhibition of SENP2 suppresses aggregate formation and its propagation

Biochemical studies demonstrated that SENP2 cleaved the N-terminal repeat regions of αSyn

whereas its inhibitor, NSC632839, suppressed aggregate formation in cultured neurons. Next,

we studied the inhibitory effects of SENP2 through the inhibitor infusion on aggregate

formation and its propagation *in vivo*. The efficiency of the aggregate formation and

inhibitory effect of NSC632839 on the propagation in the brain was examined using

immunohistochemical methods. Injection of PFFs into the striatum (Figure 5A) showed

aggregate formation, which was propagated from the striatum to other brain regions, such as

the piriform cortex and amygdala, whereas inhibitor infusion into the lateral ventricle

decreased aggregate formation and its propagation (Figure 5B). The number of cell bodies

harbouring phosphorylated-αSyn-positive aggregates was significantly decreased in the

piriform cortex and amygdala (Figure 5C, D). In this process, endogenous expression of

soluble αSyn was required for aggregate formation, because αSyn-KO mice did not show any

spreading of phosphorylated αSyn aggregates after the PFF injection (Figure S6). These

results indicated that the inhibition of the enzymatic activity attenuated aggregate formation

and its propagation efficiency.

Main figures and figure legends

Figure 1.

Bars: 50 nm

Figure 1. Isolation and characterization of pathological seeds. (A) Schematic of a

microfluidic device for pathological seed preparation. The primary neurons were cultured in

separate chambers (C1, C2, and C3) connected through microgrooves. Preformed fibrils

(PFFs) were supplemented in C2 culture medium, with height kept constantly lower than that

of C1 and C3 media during cell culture. **(B)** Propagation of phosphorylated-αSyn

(pS-αSyn)-positive intracellular aggregates from C2 (upper panels) to C1 and C3 (lower

panels) areas. **(C)** Fractionation of concentrated culture medium (CM) recovered from C1

and C3 using sucrose density-gradient ultracentrifugation. **(D)** SDS-PAGE and

immunoblotting assay using anti-αSyn antibody (numbers indicate recovered fractions in **(C)**;

His-tagged and endogenous αSyn (arrows a and b, respectively)). **(E)** Native-PAGE and

immunoblotting assay of Fraction 4 **(C, D)**. A major band at approximately 140-kDa (arrow)

was detected using anti-αSyn antibody. **(F, G)** Atomic force microscopic imaging of the

molecules recovered in Fraction 4, containing uniform granular molecules (diameter 36 ±

1.2-nm; n=16). White square in **(F)** is magnified in **(G)**. Arrowheads in **(G)** indicate the

examples of those molecules. **(H–J)** Immunoelectron microscopic imaging of molecules in

Fraction 4; αSyn-positive granular molecules detected using anti-pan-αSyn antibody (black

arrowheads: 10-nm gold particles) in **(H–J)**. Granular molecules were recognized by

anti-aggregated αSyn antibody 5G4 (**J**; red arrowheads, 6-nm gold particles), but not by the

anti-oligomeric αSyn antibody O2 (**I**; negative for 6-nm gold particles). αSyn-positive

granular molecules (diameter 35 ± 2.3 nm; n=6). Data represent mean ± SEM. Scale bars: 20

μm and 1 μm in **B** and **F**, respectively; 100 nm in **G**, and 50 nm in **H–J**.

Figure 2.

Figure 2. Truncated αSyns are included in the pathological seeds. (A) Experimental

design to prepare samples for the detection of αSyn truncation using mass spectrometric

analysis. **(B, C)** Immunoprecipitated proteins were separated by SDS-PAGE and detected by

silver staining as in **(B)**. Gel bands indicated by the numbers 1–5 were dissected and

subjected to mass spectrometric analysis. Predicted truncation sites are shown in **(C)**.

Truncated sites detected in common amino acid sequence between human and mouse αSyn

are indicated in red, whereas sites detected in human or mouse αSyn-specific sequences are

indicated in blue or green, respectively. **(D)** Schematic representation of the predicted

cleavage sites and the corresponding insertion of each mutation to inhibit the cleavage.

247 **Figure 3.**

Figure 3. Identification of processing enzyme. (A–F) Inhibition of aggregate formation by

NSC632839. Primary neurons were treated with PFFs and cultured in the presence of DMSO

(A) or NSC632839 **(B)**. The region marked by a white square in **(A)** and **(B)** is magnified in

(C) and **(D)**, respectively. Arrowheads in **(A)** and **(C)** indicate the formation of the aggregates.

Quantitative comparison of the phosphorylated-αSyn-positive area **(E)** and DAPI-positive

area **(F)** between DMSO-treated and NSC632839-treated neurons (n=3 fluorescent images

from 2 coverslips were analysed for each treatment; **p < 0.01, two-tailed, unpaired *t*-test;

two independent cultures were performed and confirmed reproducibility). **(G)** Cleavage of

αSyn by GST-fused recombinant SENP2 for 4 h at 37°C. A cleaved fragment is indicated by

the arrowhead. Mono, αSyn monomer. **(H–J)** Dose-dependent effect of SENP2 on

αSyn-cleavage. αSyn was processed for 24 h at 37°C. The cleaved fragment (red arrowhead)

was gradually increased in the presence of a higher amount of SENP2. Black and white

arrowheads show His-tagged full-length αSyn- and GST-fused SENP2 recombinant protein,

respectively. Note that αSyn-cleavage increased high-molecular-weight SDS-resistant

immunoreactivity (asterisk). This αSyn-cleavage by SENP2 for 4 h was inhibited by the

insertion of the N-terminal mutations **(J)**. Scale bar: 20 μm in **(A–D)**.

272   **Figure 4.**

5 µm

Figure 4. Presence of endogenous SENP2 in αSyn-aggregates. (A) Presence of SENP2 in

PFF-induced αSyn-aggregates formed in primary cultured neurons. **(B)** Presence of SENP2 in

αSyn-aggregates formed in the neuron of the substantia nigra pars compacta of PD-brain.

Arrowheads indicate examples of the overlapping intense signals of SENP2 and

phosphorylated αSyn. Scale bar: 5 µm.

Figure 5.

100 μm

Figure 5. Inhibition of aggregate formation and cell-to-cell seed transmission. (A)

Schematic representation of PFF injection in the WT mouse striatum, and DMSO or

NSC632839 osmotic pump-based infusion into the lateral ventricle. C, caudal; LV, lateral

ventricle; R, rostral; Str, striatum. **(B)** Inhibitory effect of NSC632839 infusion on the

propagation of αSyn-aggregate formation in the ipsilateral sites of the striatum (injection site),

piriform cortex, and amygdala. The region marked by a white square in the amygdala is

magnified in each treatment. **(C, D)** Statistical analysis of the number of cell bodies

harbouring pathological aggregates of phosphorylated αSyn, with or without NSC632839

treatment (**C**, n=5 NSC632839-treated and n=3 DMSO-treated control mice, the ipsilateral

piriform cortex was analysed by using the two-tailed, unpaired *t*-test with Welch's correction;

D, n=3 mice in each treatment for the ipsilateral amygdala analysed by the two-tailed,

unpaired *t*-test; *$p < 0.05$). Bar: 100 μm in **(B)**.

Discussion

Recent studies indicated that cell-to-cell seed transmission is a pivotal mechanism underlying

the propagation of Lewy-pathology in the PD brain[7]; however, the properties of seeds remain

unelucidated. In this study, SEC-HPLC analysis demonstrated that seeds released from

pathological cultured neurons were recovered as αSyn oligomers (approximately 1000 kDa),

whereas native-PAGE assay showed a 140-kDa band. These results suggest that intact seeds

are composed of 140-kDa rigid oligomers as the elementary units. During electrophoresis, the

recovered seeds might become dissociated from the original high-molecular structure into

those elementary units. These seeds were preferentially recognized by monoclonal antibody

5G4, but not O2. Furthermore, LBs and LNs were recognized by these antibodies[28,29].

Although LBs/LNs have epitopes against both 5G4 and O2 on their surfaces, seeds have only

5G4 epitopes on their molecular surfaces. The different structural milieu between LBs/LNs

and the seeds might be responsible for this distinct immunoreactivity.

Biochemical and mass spectrometric analysis showed that seeds contained truncated forms

of αSyn. Interestingly, N-terminal sides of lysine residues located in the N-terminal repeat

regions of αSyn were processed by SENP2 – an enzyme that routinely removes SUMO from

various SUMOylated proteins[30]. However, in the present study, SENP2 directly cleaved

recombinant αSyn itself. After αSyn cleavage by high-dose SENP2, the formation of

SDS-resistant high-molecular oligomeric species was facilitated by this treatment (see, Figure

3H, I). Furthermore, this enzymatic activity was crucial for aggregate formation. The

inhibitory effects of NSC632839 on aggregate formation and its propagation were

significantly observable *in vitro* and *in vivo*. Therefore, we further investigated the

knockdown effects of SENP2 using AAV-assisted shRNA expression system in primary

cultured neurons. However, SENP2 knockdown severely impaired normal neuronal

development. Moreover, neuronal viability was decreased by the knockdown treatment

during the neuronal maturation. On the other hand, significant knockdown efficiency was not

obtained using matured neurons. These results suggested that SENP2 was indispensable for

neuronal development, and its protein turnover rate in immature neurons might be higher than

that in matured neurons. In the present study, NSC632839-based inhibition of SENP2 did not

induce cellular damage, and this might be related to the neuronal maturation level.

Previous studies indicated that N-terminal repeat regions of αSyn implicate lipid

membrane association with conformational changes to α-helical structure[31,32]. N-terminal

truncation causes β-sheet structure-enriched conversion within the truncated αSyn[31] and the

resultant decreased stability of PFFs[17]. Thus, the N-terminal truncation of αSyn might

promote pathogenesis, including αSyn oligomerization, and further provide an intracellular

milieu that readily accelerates seed production.

Interestingly, a recent study revealed the presence of various kinds of αSyn-PTMs, such as

acetylation, methylation, and carboxymethylation, in tissue extracts prepared from Lewy

body disease (LBD) and control brains[33]. According to this study, N-terminal lysine residues

at K12, K23, K34, and K45, which are SENP2-processing sites, are post-translationally

modified with acetylation. The seeding ability of soluble αSyn harbouring K45Q, which is

one of the acetylation mimic mutations, is significantly decreased. On the other hand, lysine

residues at K12 and K23 are modified with carboxymethylation in LBD brains, but not in

control brains. The seeding ability of αSyn might be regulated by the balance of PTMs,

including truncation, acetylation, and carboxymethylation.

SENP2 was partially distributed in aggregates of phosphorylated αSyn in cultured neurons

and PD brain. However, that was not detected in a typical LB, which might be attributable to

the maturation stages of the aggregates. In the PD brain, SENP2 was detected in amorphous

aggregates, rather than spheroidal aggregates, such as typical LBs, which are larger and more

rigid than amorphous aggregates. SENP2 cleaves soluble αSyn, and the resulting fragments

mediate aggregate formation. SENP2 may persist in immature aggregates but might be

excluded from matured aggregates.

Conclusion

Transmissible seeds contained the N-terminal truncated αSyn. SENP2 directly processed

N-terminal repeat regions of αSyn, and inhibition of its activity suppressed aggregate

formation and propagation in cultured neurons and mouse brains. The present study revealed

that SENP2 activity is essential for Lewy-pathology formation. Thus, SENP2 constitutes a

potential novel therapeutic target, and fine-tuning the PTMs of αSyn, including N-terminal

truncation, could provide an innovative therapeutic strategy to prevent the progression of LB

diseases.

Limitations of this study

The presence of the characteristic seeds and the N-terminal processing sites of αSyn in the

human brain are unconfirmed. Due to the limited amount of seeds recovered, it is currently

challenging to produce structure-specific antibodies for *in vivo* seed detection using

microfluidic devices. Further studies are required for precise seed characterization.

STAR⬛Methods

Key resources table

Antibodies

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Mouse anti-αSyn antibody, Syn-1	BD Biosciences	Cat# 610787 RRID:AB_398108
Rabbit anti-αSyn antibody, C-20,	Santa Cruz Biotechnology	Cat# sc-7011-R RRID:AB_2192953
Mouse anti-phosphorylated αSyn antibody	Wako	Cat# pSyn#64 RRID:AB_516843
Rabbit anti-phosphorylated αSyn antibody	Abcam	Cat# ab59264 RRID:AB_2270761
Mouse anti-aggregated αSyn antibody, 5G4	Merck Millipore	Cat# MABN389 RRID:AB_2716647
Mouse anti-oligomeric αSyn antibody, O2	BioLegend	Cat# 847602 RRID:AB_2632701
Rabbit anti-SENP2 antibody	GeneTex	Cat# 110504 RRID:AB_1951846
Goat Alexa488-conjugated	Thermo Fisher Scientific	Cat# A11029

anti-mouse IgG antibody		RRID:AB_2534088
Goat Alexa594-conjugated anti-mouse IgG antibody	Thermo Fisher Scientific	Cat# A11032 RRID:AB_2534091
Goat Alexa488-conjugated anti-rabbit IgG antibody	Thermo Fisher Scientific	Cat# A11034 RRID:AB_2576217
Goat Alexa594-conjugated anti-rabbit IgG antibody	Thermo Fisher Scientific	Cat# A11037 RRID:AB_2534095
Goat alkaline phosphatase-conjugated anti-mouse IgG antibody	Jackson ImmunoResearch Laboratories Inc.	Cat# 115-055-146 RRID:AB_2338538
Goat alkaline phosphatase-conjugated anti-rabbit IgG antibody	Jackson ImmunoResearch Laboratories Inc.	Cat# 111-055-144 RRID:AB_2337953
Goat biotinylated anti-rabbit IgG antibody	Vector Laboratory	Cat# BA-1000 RRID:AB_2313606
Goat 6nm colloidal gold-conjugated anti-mouse IgG antibody	Abcam	Cat# ab39614 RRID:AB_954520
Goat 10nm colloidal	BBI solutions	Cat# EM.GAR10

gold-conjugated anti-rabbit IgG antibody		RRID:AB_2715527

Recombinant proteins

human USP2 catalytic domain, CF	R&D Systems	RSD-E-504-050-50
human His6-USP7 Protein, CF	R&D Systems	E-519-025
GST-tagged human SENP2 catalytic domain	Enzo	BML-UW9765-0100
mouse active UCHL1	ProSpec	PTT-PRO-2424-10

Inhibitors

Mergepta	MuseChem	R058848
carboxypeptidase inhibitor derived from potato tuber	Sigma	C0279-5MG
NSC 632839	Selleckchem	S4922

Cell culture medium, chemicals, and other reagents

Neurobasal Medium	Gibco, Thermo Fisher Scientific	Cat# 21103049
B-27	Gibco, Thermo Fisher Scientific	Cat# 17504044
L-glutamine	Nacalai Tesque	Cat# 169-19
Penicillin-Streptomycin Solution	Nacalai Tesque	Cat# 26253-84
Polyethyleneimine	Sigma	Cat# P-3143
Papain	Nacalai Tesque	Cat# 26035-44
Sucrose	Nacalai Tesque	Cat# 30404-45
Glucose	Nacalai Tesque	Cat# 16805-35
Tris-HCl	Nacalai Tesque	Cat# 35406-62
Sodium dodecyl sulfate	Nacalai Tesque	Cat# 30400-72
Pyronin Y	Wako	Cat# 164-11581
Glycerol	Nacalai Tesque	Cat# 17018-25
2-mercaptoethanol	Nacalai Tesque	Cat# 21438-82
6-aminocaproic acid	Nacalai Tesque	Cat# 020-09
Ponceau S	Nacalai Tesque	Cat# 28322-72
Tricine	Nacalai Tesque	Cat# 34713-62

Bis-Tris/HCl	Nacalai Tesque	Cat# 05103-52
Skim milk	Nacalai Tesque	Cat# 31149-75
Tween-20	Nacalai Tesque	Cat# 35624-15
NBT	Nacalai Tesque	Cat# 24720-01
BCIP	Nacalai Tesque	Cat# 05643-11
Triton X-100	Nacalai Tesque	Cat# 12967-45
Normal goat serum	Gibco, Thermo Fisher Scientific	Cat# 16210064
FluorSave	Merck Millipore	Cat# 345789
Bovine serum albumin	Sigma	Cat# A-8022
Dithiothreitol	Nacalai Tesque	Cat# 14128-91
Dimethyl sulfoxide	Nacalai Tesque	Cat# 13408-64
DAPI	Dojindo	Cat# 340-07971
HEPES	Nacalai Tesque	Cat# 17514-44
NaCl	Nacalai Tesque	Cat# 31320-05
TCEP	Nacalai Tesque	Cat# 07277-61
EDTA	Nacalai Tesque	Cat# 15111-45
PFA	Merck Millipore	Cat# 1.04005. 1000

KCl	Nacalai Tesque	Cat# 28514-75
NaHPO$_4$ · 12H$_2$O	Nacalai Tesque	Cat# 31723-35
KH$_2$PO$_4$	Nacalai Tesque	Cat# 28721-55
Trypsin for MS analysis	Pierce, MS Grade	Cat# 90057
Dithiothreitol (DDT) for MS analysis	Sigma	Cat# 43816-10ML
Iodoacetamide for MS analysis	Wako	Cat# 097-05592
Trifluoroacetic acid for MS analysis	Wako	Cat# 206-10731
Acetonitrile for MS analysis	Wako	Cat# 018-19853
Formic acid for MS analysis	Wako	Cat# 067-04531

375 **Critical commercial assay kit**

SilverQuest™ Silver Staining Kit	Thermo Fisher Scientific	LC6070
Hematoxylin and Eosin	ScyTek Laboratories	HAE-1

Stain Kit		

Experimental models: Organisms/strains

C57BL/6N	Japan SLC, Inc.	N/A
α Syn knockout mice (B6;129X1-Sncatm1Rosl/J)	Jackson Laboratory	N/A

Software for data collection and analysis

LTQ-Orbitrap Velos Pro	Thermo Fisher Scientific	N/A
MASCOT version 2.6	Matrix Science	N/A
image browser Zen 3 blue edition for LSM900	Zeiss	N/A
LAS X Office for STELLARIS5	Leica	N/A
WinRoof 2018	Mitani Corporation	N/A
GraphPad Prism 5	GraphPad Software Inc.	N/A
Adobe Photoshop 7.0	Adobe	N/A
Microsoft PowerPoint	Microsoft	N/A

Resource availability

Lead contact

Further information and requests for resources and reagents as well as datasets and protocols

used in this study should be directed to and will be fulfilled by the lead contact, KT and MT,

as described below.

Katsutoshi Taguchi, Ph.D.

Department of Anatomy and Neurobiology, Graduate School of Medical Science, Kyoto

Prefectural University of Medicine, Kawaramachi-Hirokoji, Kamikyo-ku, Kyoto 602-8566,

Japan. Tel: +81-75-251-5301; Fax: +81-75-251-5306; E-mail: ktaguchi@koto.kpu-m.ac.jp

AND

Masaki Tanaka, M.D., Ph.D.

Department of Anatomy and Neurobiology, Graduate School of Medical Science, Kyoto

Prefectural University of Medicine, Kawaramachi-Hirokoji, Kamikyo-ku, Kyoto 602-8566,

Japan. Tel: +81-75-251-5300; Fax: +81-75-251-5306; E-mail: mtanaka@koto.kpu-m.ac.jp

Materials availability

The experimental materials and data resources used in this study are available from the

corresponding authors, MT and KT, upon reasonable request.

Data and code availability

Mass spectrometric and proteomic analyses

The raw files of the mass spectrometric analysis were used to conduct searches against the mouse and human αSyn amino acid sequences that were obtained from the Mus musculus dataset (Uniprot Proteome UP000000589 2021_03_07 downloaded), Homo sapiens dataset (Uniprot Proteome UP000005640 2022_05_25 downloaded), and the common Repository of Adventitious Proteins (cRAP; ftp://ftp.thegpm.org/fasta/cRAP). The search was conducted by using MASCOT version 2.6 (Matrix Science) via Proteome Discoverer 2.4 (Thermo Fisher Scientific).

Based on the results of mass spectrometric analysis, the candidate enzymes which process the N-terminal side of lysine residues located at the N-terminal repeat regions of αSyn were screened using the Peptidase database, MEROPS (https://www.ebi.ac.uk/merops/).

All the remaining research data are available in this article and its Supplementary Information. Furthermore, the source data have been provided with this paper.

Experimental model and study participant details

Animals

αSyn homozygous knockout (KO) mice (B6;129X1-*Snca*^{tm1Rosl}/J[34]) were purchased from the Jackson Laboratory and maintained on a C57BL/6N background. Adult male C57BL/6N

wildtype (WT) and αSyn-KO (12- to 18-week-old) mice were used in the histological

experiments. KO mice were genotyped according to the manufacturer's protocol (Jackson).

The mice were housed under a 12-h light/dark cycle in a temperature-controlled environment

with ad libitum food and water in our animal facility. All animal experimental procedures

conformed to the Guidelines for Animal Experiments, Kyoto Prefectural University of

Medicine (approval numbers: M2024-148, M2024-154, and M2024-509) and national

regulations, and the number of animals used and their suffering were minimized.

Ethics approval and consent to participate

The use of postmortem brain tissue in this study was approved by the Institutional Review

Boards of Kyoto Prefectural University of Medicine (approval no. ERB-C-1846-2). Written

informed consent for cadaver surgical training and the use of tissue samples for medical

research was obtained, both from patients themselves and their next of kin.

Consent for publication

Not applicable

Method details

Primary neuronal culture

Using a previously reported method[35], with slight modifications, primary cultured neurons

were obtained. Briefly, dissociated cells obtained from the prefrontal cortices and hippocampi

of WT mouse embryos (E15–18) were disseminated onto polyethyleneimine-coated

coverslips or into the chambers of microfluidic devices, and cultured in Neurobasal Medium

(Gibco, Thermo Fisher Scientific) supplemented with B-27 (Gibco, Thermo Fisher Scientific),

L-glutamine (Nacalai Tesque), and penicillin/streptomycin (Nacalai Tesque). Half of the

volume of the culture medium was changed every 3–4 days until use. To induce the formation

of intracellular phosphorylated αSyn aggregates, neurons were cultured in PFF-supplemented

complete Neurobasal Medium (final 20 µg/mL) for 4–5 days, unless specified otherwise.

Preparation of recombinant αSyn and PFFs

For the preparation of recombinant human αSyn from bacterial culture, plasmid vectors were

constructed as described previously[36]. Briefly, PCR fragments of human αSyn were inserted

into pTrc-His-TOPO vector (Thermo Fisher Scientific). His-tagged αSyn was purified by

cOmplete™ His-Tag Purification Column (Roche). To confirm the purity of the protein by

SDS-PAGE, the purified fraction was electrophoresed, and the protein band was stained by

Coomassie Brilliant Blue. The final protein yield was determined with Lowry's method.

PFFs were prepared according to a previously reported method, albeit with some slight

modifications[11]. Briefly, PFFs were generated by incubating purified αSyn in sterile PBS

(final concentration: 2 mg/ml) at 37°C with constant agitation for 5–6 days, then

ultracentrifuging (at 200,000g) the PFF-containing solution for 2 h, and recovering the PFF

precipitate at the bottom of the centrifuge tube. The resulting pellet was dissolved in sterile

PBS and stored at −30°C until use.

To prepare recombinant αSyn harbouring mutations at the N-terminal repeat region (Figure

2D), a DNA fragment of mouse full-length αSyn harbouring those mutations was artificially

synthesized and inserted into the pEX-A2J2 vector by Eurofins Genomics Japan. A DNA

fragment of the mutated αSyn was further cloned into pTrc-His2B vector (Thermo Fisher

Scientific). The presence of each mutation was confirmed by DNA sequencing using the ABI

PRISM3130 Genetic analyzer. The His-tagged mutated αSyn-recombinant protein was

purified using the abovementioned method described for WT αSyn.

Microfluidic devices

Transmission of pathological αSyn was reproduced *in vitro* using microfluidic devices[37]. To

isolate cell-to-cell transmissible pathological seeds, two types of microfluidic devices were

used. For small-scale preparation of the seeds, three-compartment devices (TCND500) were

purchased from Xona and cell culture was performed according to the manufacturer's

protocol. Briefly, pathological neurons harbouring intracellular phosphorylated-αSyn

aggregates were cultured in the middle chamber, C2 (Figure 1A). Seeds released from those

pathological neurons were recovered in the culture medium in the side chambers, C1 and C3,

which are connected to C2 through microgrooves. During cell culture, to prevent the

backflow of the PFF-containing medium into the C1 and C3 chambers, the medium in the C2

chamber was constantly maintained at a lower height than that of the C1 and C3 chambers.

For large-scale preparation of seeds to perform mass spectrometric analysis, a large

polydimethylsiloxane-based microfluidic device was developed (Figure S2A–C). Before the

dissemination of dissociated cells prepared from the embryonic brain, the large microfluidic

device was settled on a polyethyleneimine-coated glass dish that comprises two areas – a

central square area and a hollow square (outer) area – which are connected by microgrooves

through which the axons extend (Figure S2D, E). Pathological neurons were cultured in the

central area of this device, and the seeds released from these neuronal cells were recovered

from the medium in the outer area. During cell culture, the height of the medium in the

central area was constantly maintained lower than that in the outer area to prevent the

backflow of the PFF-containing medium.

Preparation of the seed fraction

Unless specified otherwise, the seeds were prepared using the microfluidic devices. Partly

seeds fraction was prepared using the batch method (Figure S1A–F). After the PFF treatment

on primary neurons cultured in the middle or central square chamber, half of the volume of

the seed-containing medium in the side (C1 and C3) or outer chambers was recovered every

3–4 days for 14 days. By ultrafiltration through a 3-kDa molecular-weight cutoff membrane

(Merck Millipore), this seed-containing medium was concentrated up to 10 folds. Next, a

discontinuous sucrose density gradient was prepared in a centrifuge tube purchased from

Beckman Coulter (Figure 1C and Figure S1A). Each layer comprised phosphate-buffered

saline (PBS, pH 7.4) containing the indicated concentration of sucrose. The concentrated

medium was overlaid on a 0.8 M sucrose-containing PBS layer, ultracentrifuged at 200,000g

for 12 h at 4°C, and an equal volume (250-300 μL) was aliquoted and further subjected to

biochemical analysis. Seeds were recovered on the interface between the concentrated

medium and the 0.8 M sucrose-containing PBS layer (Fraction 4 in Figure 1C).

For seed preparation using the batch method, the conditioned medium of pathological

neurons was recovered as the seed-containing medium. After 2-day culture under PFF

treatment, the PFF-containing medium was discarded, the cells were washed 4 times with

complete Neurobasal Medium and then cultured in a fresh Neurobasal Medium for 5–7 days.

This conditioned medium was collected and concentrated using ultrafiltration as described

above. The concentrated medium was fractionated using sucrose density-gradient

ultracentrifugation as well as the microfluidic method. Seeds were recovered at the interface

between the concentrated medium and 0.8 M sucrose-containing PBS and subjected to

biochemical analysis or aggregate-formation assay (Figure S1).

Immunoblotting assay

For SDS-PAGE, samples were denatured by heating with 1% 2-mercaptoethanol-containing

SDS-sample buffer (62.5 mM Tris-HCl, pH 6.8, 2% sodium dodecyl sulfate, 0.003% Pyronin

Y, and 10% glycerol) at 98°C for 3 min. Proteins were separated by 15% polyacrylamide gel

(Wako) and transferred to a polyvinylidene difluoride (PVDF) membrane (Bio-Rad

Laboratories). For native-PAGE, the samples were treated with sample buffer (15% glycerol,

0.5 M 6-aminocaproic acid, 0.001% Ponceau S, and 50 mM Bis-Tris/HCl, pH 7.0). Proteins

were separated by NativePAGE™ Bis-Tris Gel purchased from Thermo Fisher Scientific and

electrophoresed using anode (50 mM Bis-Tris/HCl, pH 7.0) and cathode (50 mM Tricine, 15

mM Bis-Tris/HCl, pH 7.0) buffers. After electrophoresis, polyacrylamide gel was immersed

in a transfer buffer containing 1% SDS, and the separated protein bands were also denatured

in this step. The denatured proteins were transferred to the PVDF membrane; the membrane

was blocked with 5% skim milk in Tris-buffered saline (TBS, pH 7.4) containing 0.1%

Tween-20 for 1 h at room temperature. After treatment with primary antibody, the membrane

was incubated with an alkaline phosphatase-conjugated secondary antibody. Protein bands

were detected using the NBT-BCIP system (Nacalai Tesque). For αSyn detection, mouse

monoclonal antibody Syn-1 (1:1000; BD Biosciences) or rabbit polyclonal antibody C-20

(1:2000; Santa Cruz Biotechnology) was used.

Immunocytochemistry

Cultured cells were fixed with 2% paraformaldehyde (PFA) in the culture medium for 10 min

at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 10 min, and blocked

with 10% normal goat serum (NGS) in PBS for 30 min. Next, cells were incubated for 1–2 h

with primary antibodies in the blocking solution, washed with PBS, and treated with

secondary antibodies. The primary antibodies were detected with Alexa488- and

Alexa594-conjugated secondary antibodies (Thermo Fisher Scientific). After labelling with

the secondary antibodies, cells were washed with PBS and then with milliQ water (Merck

Millipore), before being mounted with FluorSave (Merck Millipore).

   For αSyn detection, Syn-1 or C-20 antibodies were used. Phosphorylated αSyn was

detected by mouse monoclonal antibody pSyn#64 (1:1000; Wako) or rabbit polyclonal

antibody phospho-S129 αSyn (1:1000; Abcam). SENP2 was detected by rabbit polyclonal

antibody (1:250; GeneTex). Low-magnification images were acquired by using

immunofluorescent signals captured through a UPlanFl 10×/0.30 Ph1 objective with an

inverted fluorescent microscope IX71 (Olympus) equipped with a cooled charge-coupled

device camera (Olympus). For confocal observation, images were acquired as Z stacks (5–15

z-sections, 1-μm apart, 1024 × 1024 pixels) using a Plan-Apochromat 63×/1.40 Oil DIC

objective (Carl Zeiss) with an inverted laser-scanning confocal microscope, LSM900 (Carl

Zeiss) or using an HC PL APO CS2 63×/1.40 OIL objective (Leica) with a laser-scanning

confocal microscope, STELLARIS5 (Leica) for high-resolution imaging (Figure 4A).

Atomic force microscopy

To acquire images of the seed structure, the seed fraction was dropped onto the surface of a

freshly cleaved mica plate (NILACO) and incubated for 5 min. Images were acquired using

atomic force microscopy, SPM-9700 (SHIMADZU), operating in Dynamic Mode. The

scanning parameters were as follows: operating point, 0.146–0.166 V; drive (tapping)

frequency, 323–325 kHz; and scan rate, 1.0 Hz.

Size-exclusion chromatography

To determine the molecular weight of the seeds, size-exclusion chromatography

(SEC)–high-performance liquid chromatography-based analysis was performed using the

SMART system (Pharmacia). Briefly, 50 μL seed fraction was injected into a sample tube and

separated using the Superose 6 column comprising filtered PBS (pH 7.4) as the mobile phase.

Each sample was eluted at a 40 μL/min flow rate; each 80 μL was collected as a fraction.

Protein elution was monitored by UV absorbance at 280 nm. To obtain a standard curve for

the molecular-weight determination, gel-filtration standards (Gel Filtration Calibration Kit

HMW) purchased from GE Healthcare were separated using the same SEC column and flow

rate. To detect the presence of the seed structure recovered in these SEC fractions, each SEC

fraction was further observed using immunoelectron microscopy.

Immunoelectron microscopy

For immunolabelling of seeds or PFFs, the seed fraction or sonicated PFFs were dropped

onto the surface of formvar-coated molybdenum grids, PVF-M15 STEM Mo150P

(Okenshoji), and incubated for 10 min. After incubation, the excess solution on each grid was

discarded; the grid surface was further blocked with 1% bovine serum albumin (BSA) in PBS

for 15 min and incubated with primary antibodies for 1 h. After washing in PBS, primary

antibodies were detected with 10-nm gold (BBI Solutions) and/or 6-nm gold

(Abcam)-conjugated secondary antibodies in PBS containing 1% BSA. After labelling, the

grids were washed with PBS, stained with uranyl acetate, and observed using a JEM-1220

electron microscope (JEOL) at 80 keV.

For αSyn detection, the C-20 antibody was used as a pan-αSyn antibody. For the detection

of oligomeric species of αSyn, the O2 (1:500; BioLegend) and 5G4 (1:500; Merck Millipore)

antibodies, which recognize soluble oligomers and fibril forms of αSyn, including LBs and

LNs[28,29], were used.

Mass spectrometric analysis for screening of the αSyn-cleavage site

To detect the cleavage sites of the seed-constituting αSyn from the mass spectrometric

analysis sample, the seeds were concentrated by immunoprecipitation. Briefly, the seed

fraction was prepared using large microfluidic devices as the original fraction for

immunoprecipitation. The seeds were concentrated by immunoprecipitation after labelling by

a C-20 antibody and precipitation with protein A-conjugated magnet beads (Bio-Rad), and

then eluted by the SDS sample buffer at 56°C for 15 min. The eluted proteins were further

denatured by heating at 98°C with 1% 2-mercaptoethanol for 3 min, and separated by

SDS-PAGE. Protein bands were detected by silver staining using the SilverQuest™ Silver

Staining Kit (Thermo Fisher Scientific). The PFF-constituting αSyn was prepared as the

control for mass spectrometric analysis, because the results of the artificially cleaved sites

detected in the PFF-constituting αSyn were subtracted from those of the cleaved sites

detected in the seed-constituting αSyn. These protein bands (Nos. 1–5 and control bands,

Figure 2B) were subjected to mass spectrometric analysis.

For mass spectrometric analysis using Nano-liquid chromatography–tandem mass

spectrometry (LC–MS/MS), the proteins present in each gel slice underwent a series of

stepwise treatment as follows: reduction with 10 mM dithiothreitol (DDT) at 56°C for 1 h,

alkylation with 55 mM iodoacetamide at room temperature for 45 min in the dark, and

digestion with 10 μg/mL modified trypsin (Pierce, MS Grade) at 37°C for 16 h. The resulting

peptides were extracted using 1% trifluoroacetic acid and 50% acetonitrile, vacuum-dried,

and dissolved in a solution of 2% acetonitrile and 0.1% formic acid.

Mass spectra were acquired using an LTQ-Orbitrap Velos Pro (Thermo Fisher Scientific)

coupled with a nanoflow UHPLC system (ADVANCE UHPLC; AMR Inc.) featuring

Advanced Captive Spray SOURCE (AMR Inc.). The peptide mixtures were loaded onto a

C18 ID 0.1 mm × 20 mm, 5-μm (particle size) trap column (Acclaim PepMap 100 C18,

Thermo Fisher Scientific) and fractionated through a C18, ID 0.075 × 500 mm, particle size

3-μm column (CERI). The peptides were eluted at a flow rate of 300 nL/min with a linear

gradient of 5–35% solvent B over 60 min. Buffers A and B comprised 0.1% formic acid and

100% acetonitrile, respectively. The mass spectrometer was programmed for 11 successive

scans, including a full-scan MS over 350–2,000 m/z by orbitrap at a resolution of 60,000, and

automatic data-dependent MS/MS scans of the 10 most intense ion signals in the first

precursor scan by ion-trap detectors for the second and eleventh scans. Using a 2 m/z

isolation width and a 90-s exclusion time for molecules in the same m/z range, MS/MS

spectra were obtained with a normalized collision energy of 35% CID.

The raw files were searched against the mouse and human αSyn amino acid sequences,

including the N-terminal His-tag sequence, with Mus musculus dataset (Uniprot Proteome

UP000000589 2021_03_07 downloaded), Homo sapiens dataset (Uniprot Proteome

UP000005640 2022_05_25 downloaded), and the common Repository of Adventitious

Proteins (cRAP; ftp://ftp.thegpm.org/fasta/cRAP). The search was conducted through

MASCOT version 2.6 (Matrix Science) via Proteome Discoverer 2.4 (Thermo Fisher

Scientific). The following settings were used as the search parameters: Enzyme,

Semi-trypsin; false discovery rate (FDR), 0.01; precursor mass tolerance, 10 ppm; product

tolerance, 0.8 Da; fixed modification, carbamidomethylation of cysteine; variable

modification, oxidation of methionine, and acetylation of protein N-termini; and maximum

missed cleavages, 2.

Screening of the αSyn-processing enzyme and inhibitor assay

Based on the results of mass spectrometric analysis, the candidate enzymes which cleave the

N-terminal side of lysine residues located at the N-terminal repeat regions of αSyn

(conserved cleavage sites in Figure 2C) were screened using the Peptidase database,

MEROPS (https://www.ebi.ac.uk/merops/); next, inhibitors for each enzyme were selected

(Table 1), dissolved with dimethyl sulfoxide (DMSO), and stored at −30°C until use. To

ascertain the inhibitory effect of the inhibitors on aggregate formation, primary neurons were

cultured and treated with PFFs in the presence of each inhibitor. The following inhibitors

were purchased from the manufacturers and used at the indicated final concentration:

Mergepta (46 μM; MuseChem), carboxypeptidase inhibitor derived from potato tuber (CPI)

(1 μM; Sigma), and NSC 632839 (9 μM; Selleckchem). As a control, primary neurons were

cultured with diluted DMSO as well as each inhibitor.

Image analysis and quantification of the inhibitory effect of NSC632839 on aggregate

formation in primary cultured neurons

The phosphorylated αSyn-positive and DAPI-positive areas were measured using WinRoof

2018 software (Mitani Corporation). As shown in Figure S3, the phosphorylated-αSyn

immunoreactive area, excluding the DAPI-positive area, was quantitatively analysed because

the monoclonal antibody against phosphorylated αSyn #64 tends to react with nuclei. In

either the presence or absence of NSC632839, wherein the threshold intensity was kept

constant and set to include measurement profiles by visual inspections. Data represent the

mean ± SEM. Statistical analysis was performed using GraphPad Prism 5 with a two-tailed,

unpaired *t*-test and assessed using a significance level of $p < 0.05$ (n=3 fluorescent images

acquired from 2 coverslips were analysed for each treatment; two independent cultures were

performed, and reproducibility was confirmed).

Protease assay

To examine whether αSyn is processed by the several candidate enzymes summarized in

Table 1, a protease assay was performed using recombinant active enzymes: 7 µg soluble or

PFF forms of recombinant αSyn (substrate) was mixed with 0.5–5.2 µg of each recombinant

enzyme in the following buffer to a final volume of 100 µL: USP7 (50 mM HEPES (pH8.0),

0.1 M NaCl, and 1 mM TCEP), SENP2 (50 mM HEPES, pH 8.0; 0.15 M NaCl; and 1 mM

DTT), USP2 and UCHL1 (50 mM HEPES, pH 8.0, 0.15 M NaCl, 0.1 mM EDTA, and 1 mM

DTT). PFFs were sonicated using the Hielscher ultrasonic processor UP50H with sonotrode

MS1 (10□s on–10□s off, repeated 10 times on ice, at 100% amplitude setting) prior to use.

Each substrate was incubated with each enzyme at 37°C for the indicated duration (4–24 h).

The reaction was stopped by the addition of SDS sample buffer, and the samples were

immediately heated with SDS-sample buffer at 98°C for 3□min and subjected to

immunoblotting assay. Recombinant enzymes were purchased from the following

manufacturers: human USP2 catalytic domain, CF (R&D Systems), human His6-USP7

Protein, CF (R&D Systems), GST-tagged human SENP2 catalytic domain (Enzo), and mouse

active UCHL1 (ProSpec).

Human PD brain

The use of postmortem brain tissue in this study was approved by the Institutional Review

Boards of Kyoto Prefectural University of Medicine (approval no. ERB-C-1846-2). Written

informed consent for cadaver surgical training and the use of tissue samples for medical

research was obtained, both from patients themselves and their next of kin. The postmortem

brain specimen was obtained from a patient with PD (age: 85 years, female), fixed using

Thiel's method, with slight modifications. Briefly, a fixative containing 5% formalin was

perfused from the superior sagittal sinus. The brain was removed from the skull, immersed in

3.7% formalin for 14 days, and dissected into tissue blocks, including the substantia nigra

(SN). Tissue blocks were further fixed with 4% PFA in 0.1 M phosphate buffer (PB) for 2

days. After fixation, these tissue blocks were subjected to immunohistochemistry as described

below.

PFF injection and inhibitor infusion *in vivo*

To reproduce the propagation of LB-like pathology *in vivo*, WT mice were anesthetized using

a midazolam/medetomidine/butorphanol cocktail (4, 0.3, and 5 mg/kg, respectively), and 2

μL PFFs (7 mg/mL) was stereotaxically injected into the striatum, as reported previously[8]. A

needle inserted into the left forebrain (+0.98 mm relative to Bregma, +1.70 mm from

midline) was used to direct the inoculum into the striatum (+3.5 mm beneath the dura) and

was injected using a 10-μL Hamilton syringe at a flow rate of 0.2 μL/min (total 14 μg PFFs

per site), with the needle left in place for >20 min.

The inhibitor (NSC 632839, final concentration 73.5 μM in sterile saline) was infused into

the left lateral ventricle using an Alzet osmotic pump (Model 2006) connected to a brain

infusion kit (Type 3) in accordance with the manufacturer's instructions (Alzet) for

controlled-release of the reagent (0.15 μL/h for 42 days). Subjects were monitored regularly

following their postoperative recovery, and sacrificed at 6 weeks after the surgery. The

efficiency of the aggregate formation and inhibitory effect of NSC632839 on aggregate

formation was determined using immunohistochemistry. Data from NSC635839-infused mice

(n=5) and DMSO-infused control mice (n=3) were included in the statistical analysis as

described below.

To study the effect of endogenous αSyn expression on aggregate formation, we injected

PFF into the striatum of WT and αSyn-KO mice (Figure S6), and the PFF injections were

performed as described above.

Immunohistochemistry and Hematoxylin and Eosin (HE) staining

Unless specifically mentioned, a standard immunohistological procedure was performed on

free-floating sections prepared from mouse and human brains.

For mouse brain, immunostaining was performed as previously described[38]. Briefly, under

deep anesthesia, adult male mice were intracardially perfused with PBS followed by 4% PFA

in PBS. Brains were dissected into blocks and postfixed using the same fixative for 12 h at

4°C. Coronal sections (40 μm) were obtained using a vibratome (Dosaka-EM), permeabilized

with 0.3% Triton X-100 in PBS (PBST) for 1 h, and blocked with 10% NGS in PBST for 20

h. These sections were then incubated with the primary antibody in the blocking solution for

20 h, washed with PBST, and treated with a secondary antibody. For double immunostaining,

this staining procedure was repeated. For the detection of phosphorylated αSyn, two types of

antibodies were used: monoclonal mouse IgG antibody (#64) and polyclonal rabbit IgG

antibody (pS129 αSyn).

For the immunofluorescent method, the primary antibodies were detected with Alexa488-

and Alexa594-conjugated secondary antibodies (Thermo Fischer Scientific). Sections were

washed with PBST and then with 20 mM Tris-HCl buffer, and mounted with FluorSave

(Merck Millipore). For confocal observation, images were acquired as Z stacks (10–20

z-sections, 1-µm apart, 1024×1024 pixels) using a Plan-Apochromat 63x/1.40 Oil DIC

objective (Carl Zeiss) with an inverted laser-scanning confocal microscope, LSM900 (Carl

Zeiss).

For the immunoenzyme method, the primary antibody was detected by biotinylated

anti-rabbit IgG antibody (1:500; Vector Laboratory) for 4 h, followed by horseradish

peroxidase (HRP)-conjugated streptavidin (1:2; Nichirei Biosciences Inc.) detection for 1 h.

HRP was visualized with 3,3'-diaminobezidine tetrahydrochloride and H_2O_2 in 50 mM

Tris-buffered saline (pH 7.4). Immunosignals were acquired through the UPlanFl 60×

objective with an inverted microscope (IX71), equipped with a cooled charge-coupled device

camera (Olympus).

For the human PD brain, the dissected tissue block including the SN was cryoprotected

using 20% sucrose in 0.1 M PB (pH 7.4) and rapidly frozen in Tissue-Tek OCT compound

(Sakura Finetek Japan). Sections were obtained using a cryostat (Leica) at 30-µm thickness.

For HE staining, sections were stained on a slide glass using a Hematoxylin and Eosin Stain

Kit according to the manufacturer's standard protocol (ScyTek Laboratories). For

immunostaining with antigen retrieval, sections were immersed in formic acid for 10 min,

washed 4 times with PBS, and subjected to immunohistochemical analysis using

anti-phosphorylated αSyn (#64) and anti-SENP2 antibodies with the immunostaining

procedure described for mouse brain sections.

Quantification of the number of cell bodies harbouring pathological aggregates of

phosphorylated αSyn

The number of cell bodies harbouring aggregates of phosphorylated αSyn in the amygdala

and piriform cortex of NSC632839-treated and DMSO-treated control mouse brains was

counted. For cell-counting in each region, 3–6 sections per animal for each treatment were

subjected to the cell-counting procedure. The number of animals used in each experimental

group is indicated in the figure legends (Figure 5F, G). After the cell-counting procedure, the

mean cell numbers per animal were individually calculated in each brain region; the mean

values per experimental group were then used in the statistical analysis using GraphPad

Prism 5 by the two-tailed, unpaired *t*-test, with or without Welch's correction. Data are

expressed as the mean ± SEM. The significance levels were set at *$p < 0.05$.

Image processing

The digital images obtained on the LSM900 or IX71 were processed using Adobe Photoshop

7.0 (Adobe). Each figure set was prepared in Microsoft PowerPoint (Microsoft).

Statistical analysis and reproducibility

Statistical analyses between the two experimental groups were performed using the

two-tailed, unpaired t-test, with or without Welch's correction (GraphPad Prism 5, GraphPad

Software Inc.). The number of animals used in each experiment and the number of images

that were analysed are specified in the Methods and/or indicated in each figure legend. Data

are expressed as the mean ± standard error of the mean. The significance levels were set at p

< 0.05 (*) and $p < 0.01$ (**), as indicated. Reproducibility was confirmed by independent

experiments that were repeated at least two to three times, with similar results.

Acknowledgments

We thank Professor Masaya Ikegawa (Genomics, Proteomics and Biomedical Functions,

Department of Life and Medical Systems, Faculty of Life and Medical Sciences, Doshisha

University, Kyoto, Japan) for facilitating the pilot experiments in mass spectrometric analysis

and thank Dr. Reiko Nakagawa (Laboratory for Cell-Free Protein Synthesis, RIKEN Center

for Biosystems Dynamics Research, Kobe, Japan) for the mass spectrometric analysis and the

780 preparation of the datasheets. This work was supported by Grants-in-Aid for Scientific

781 Research from the Japan Society for the Promotion of Science (KT: 24K10514, 21K07299,

782 and 18K07371; YW: 24K10494; MT: 21K06412) and The Shimizu Foundation for

783 Immunology and Neuroscience Grant 2021 (to KT).

Author contributions

786 KT and MT conceived and designed the experiments. KT performed the experiments. YW

787 contributed to the analysis and discussion. KT and MT wrote the manuscript.

Declaration of interests

790 The authors have no potential or actual conflicts of interest.

799 **Supplemental information**

800 **Table 1. Screening of αSyn-processing enzyme**

	Candidate enzyme	Inhibitor	Inhibition of aggregate formation	Activity of αSyn-cleavage
1	Carboxypeptidase B (B2, N)	Mergepta and CPI	(-)	
2	USP2 (ubiquitin specific peptidase 2)	NSC632839	(+)	(-)
3	USP7 (ubiquitin specific peptidase 7)			(-)
4	SENP2 (Sentrin-specific protease 2)			(+)
5	UCHL1 (ubiquitin carboxy-terminal hydrolase-1)			(-)

Supplemental data figures and legends

Figure S1.

Figure S1. Isolation and characterization of pathological seeds using the batch method.

(A) Schematic diagram of fractionation by sucrose-density gradient ultracentrifugation of the

conditioned culture medium derived from pathological primary neurons harbouring Lewy

body- and Lewy neurite-like aggregates. CM, culture medium. **(B)** SDS-PAGE and an

immunoblotting assay of the fractions using anti-αSyn antibody. The numbers indicate the

recovered fractions shown in **(A)**. **(C)** Native-PAGE and immunoblotting assay of the

fractions. **(D)** The seeding ability of the pathological seeds recovered in Fraction 3. pS-αSyn,

phosphorylated αSyn. **(E)** Standard curve for HPLC (Log. molecular weight vs. elution

volume). **(F)** Immunoelectron microscopic observation of the αSyn-oligomer species

recovered in each HPLC-prepared fraction. **(G)** Immunoelectron microscopic observation of

PFFs as a control for αSyn immunolabeling. PFFs, preformed fibrils. Scale bar: 20 μm in **(D)**,

50 nm in **(F)**, and 100 nm in **(G)**.

Figure S2.

100 μm

Figure S2. Development of the microfluidic device for large-scale preparation of

pathological seeds. (A–C) Overview of microfluidic device for large-scale preparation of

seeds. **(D, E)** Primary neuronal culture using the large-scale microfluidic device for 5 days

(D) and 12 days **(E)** *in vitro*. Arrowheads show the extending axons. The region marked by a

red square in **(D)** and **(E)** is magnified in each right panel, respectively. GC, growth cone.

Scale bar: 100 µm.

Figure S3.

Figure S3. Image analysis of the inhibitory effect of NSC632839 on aggregate formation.

Phosphorylated-αSyn immunoreactive area, excluding the DAPI-positive area surrounded by

dotted lines, was quantitatively analysed because the monoclonal antibody against

phosphorylated αSyn #64 tends to react with the nuclei. Arrowheads indicate examples of

phosphorylated αSyn-positive aggregates. Scale bar: 20 μm.

Figure S4.

Figure S4. Screening assay for possible αSyn-processing enzymes. (A–C)

αSyn-processing ability of possible NSC632839-inhibited enzymes was examined by a

protease assay using recombinant proteins. As indicated by asterisks, cleaved fragments were

not detected in the presence of USP-7 **(A)**, USP-2 **(B)**, or UCHL1 **(C)**. Both PFFs and a

monomeric form of αSyn were not processed as appropriate substrates by these enzymes. The

reaction time and temperature are indicated in each figure panel.

Figure S5.

Figure S5. Presence of SENP2 in Lewy pathology. (A, B) Lewy body, Lewy neurites, and

neuromelanin observed in the substantia nigra pars compacta of the PD-brain. **(C, D)**

Presence of endogenous SENP2 in amorphous aggregates **(C)**, but not in a typical Lewy body

(D). Arrowheads indicate the presence of both SENP2 and phosphorylated αSyn. LB, Lewy

body; LN, Lewy neurites; NM, neuromelanin. Scale bar: 20 μm in **(A, B)**, and 10 μm in **(C,**

D).

Figure S6.

Figure S6. Endogenous αSyn expression is indispensable for aggregate formation. (A)

Schematic diagram of PFF injection into the striatum of the mouse brain. C, caudal; R,

rostral; Str, striatum. **(B)** Intense formation of phosphorylated-αSyn aggregates was observed

in the WT, but not αSyn-KO, mouse brain. WT, wildtype; KO, knockout. Scale bar: 50 μm.

References

873 1. Spillantini, M.G., Crowther, R.A., Jakes, R., Hasegawa, M., and Goedert, M. (1998).

874   alpha-Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease

875   and dementia with lewy bodies. Proc Natl Acad Sci U S A *95*, 6469-6473.

876   10.1073/pnas.95.11.6469.

877 2. Dickson, D.W. (2001). Alpha-synuclein and the Lewy body disorders. Curr Opin

878   Neurol *14*, 423-432. 10.1097/00019052-200108000-00001.

879 3. Stefanis, L. (2012). alpha-Synuclein in Parkinson's disease. Cold Spring Harb

880   Perspect Med *2*, a009399. 10.1101/cshperspect.a009399.

881 4. Braak, H., and Del Tredici, K. (2009). Neuroanatomy and pathology of sporadic

882   Parkinson's disease. Adv Anat Embryol Cell Biol *201*, 1-119.

883 5. Braak, H., Del Tredici, K., Rub, U., de Vos, R.A., Jansen Steur, E.N., and Braak, E.

884   (2003). Staging of brain pathology related to sporadic Parkinson's disease. Neurobiol

885   Aging *24*, 197-211. 10.1016/s0197-4580(02)00065-9.

886 6. Li, J.Y., Englund, E., Holton, J.L., Soulet, D., Hagell, P., Lees, A.J., Lashley, T., Quinn,

887   N.P., Rehncrona, S., Bjorklund, A., et al. (2008). Lewy bodies in grafted neurons in

888   subjects with Parkinson's disease suggest host-to-graft disease propagation. Nat Med

889   *14*, 501-503. 10.1038/nm1746.

890 7. Volpicelli-Daley, L., and Brundin, P. (2018). Prion-like propagation of pathology in

891   Parkinson disease. Handb Clin Neurol *153*, 321-335.

10.1016/B978-0-444-63945-5.00017-9.

8.   Luk, K.C., Kehm, V., Carroll, J., Zhang, B., O'Brien, P., Trojanowski, J.Q., and Lee,

V.M. (2012). Pathological alpha-synuclein transmission initiates Parkinson-like

neurodegeneration in nontransgenic mice. Science *338*, 949-953.

10.1126/science.1227157.

9.   Peng, C., Gathagan, R.J., Covell, D.J., Medellin, C., Stieber, A., Robinson, J.L.,

Zhang, B., Pitkin, R.M., Olufemi, M.F., Luk, K.C., et al. (2018). Cellular milieu

imparts distinct pathological alpha-synuclein strains in alpha-synucleinopathies.

Nature *557*, 558-563. 10.1038/s41586-018-0104-4.

10.   Henderson, M.X., Cornblath, E.J., Darwich, A., Zhang, B., Brown, H., Gathagan, R.J.,

Sandler, R.M., Bassett, D.S., Trojanowski, J.Q., and Lee, V.M.Y. (2019). Spread of

alpha-synuclein pathology through the brain connectome is modulated by selective

vulnerability and predicted by network analysis. Nat Neurosci *22*, 1248-1257.

10.1038/s41593-019-0457-5.

11.   Volpicelli-Daley, L.A., Luk, K.C., Patel, T.P., Tanik, S.A., Riddle, D.M., Stieber, A.,

Meaney, D.F., Trojanowski, J.Q., and Lee, V.M. (2011). Exogenous alpha-synuclein

fibrils induce Lewy body pathology leading to synaptic dysfunction and neuron death.

Neuron *72*, 57-71. 10.1016/j.neuron.2011.08.033.

12.   Suthar, S.K., and Lee, S.Y. (2023). Truncation or proteolysis of alpha-synuclein in

Parkinsonism. Ageing Res Rev *90*, 101978. 10.1016/j.arr.2023.101978.

13.   Canever, J.B., Soares, E.S., de Avelar, N.C.P., and Cimarosti, H.I. (2023). Targeting

alpha-synuclein post-translational modifications in Parkinson's disease. Behav Brain

Res *439*, 114204. 10.1016/j.bbr.2022.114204.

14.   Fujiwara, H., Hasegawa, M., Dohmae, N., Kawashima, A., Masliah, E., Goldberg,

M.S., Shen, J., Takio, K., and Iwatsubo, T. (2002). alpha-Synuclein is phosphorylated

in synucleinopathy lesions. Nat Cell Biol *4*, 160-164. 10.1038/ncb748.

15.   Rott, R., Szargel, R., Shani, V., Hamza, H., Savyon, M., Abd Elghani, F.,

Bandopadhyay, R., and Engelender, S. (2017). SUMOylation and ubiquitination

reciprocally regulate alpha-synuclein degradation and pathological aggregation. Proc

Natl Acad Sci U S A *114*, 13176-13181. 10.1073/pnas.1704351114.

16.   Savyon, M., and Engelender, S. (2020). SUMOylation in alpha-Synuclein

Homeostasis and Pathology. Front Aging Neurosci *12*, 167.

10.3389/fnagi.2020.00167.

17.   Terada, M., Suzuki, G., Nonaka, T., Kametani, F., Tamaoka, A., and Hasegawa, M.

(2018). The effect of truncation on prion-like properties of alpha-synuclein. J Biol

Chem *293*, 13910-13920. 10.1074/jbc.RA118.001862.

18.   Sorrentino, Z.A., and Giasson, B.I. (2020). The emerging role of alpha-synuclein

truncation in aggregation and disease. J Biol Chem *295*, 10224-10244.

10.1074/jbc.REV120.011743.

19.   McGlinchey, R.P., Ni, X., Shadish, J.A., Jiang, J., and Lee, J.C. (2021). The N

932   terminus of alpha-synuclein dictates fibril formation. Proc Natl Acad Sci U S A *118*.

933   10.1073/pnas.2023487118.

934   20.   Legname, G., Nguyen, H.O., Peretz, D., Cohen, F.E., DeArmond, S.J., and Prusiner,

935   S.B. (2006). Continuum of prion protein structures enciphers a multitude of prion

936   isolate-specified phenotypes. Proc Natl Acad Sci U S A *103*, 19105-19110.

937   10.1073/pnas.0608970103.

938   21.   Tsujimura, A., Taguchi, K., Watanabe, Y., Tatebe, H., Tokuda, T., Mizuno, T., and

939   Tanaka, M. (2015). Lysosomal enzyme cathepsin B enhances the aggregate forming

940   activity of exogenous alpha-synuclein fibrils. Neurobiol Dis *73*, 244-253.

941   10.1016/j.nbd.2014.10.011.

942   22.   Mishizen-Eberz, A.J., Guttmann, R.P., Giasson, B.I., Day, G.A., 3rd, Hodara, R.,

943   Ischiropoulos, H., Lee, V.M., Trojanowski, J.Q., and Lynch, D.R. (2003). Distinct

944   cleavage patterns of normal and pathologic forms of alpha-synuclein by calpain I in

945   vitro. J Neurochem *86*, 836-847. 10.1046/j.1471-4159.2003.01878.x.

946   23.   Dufty, B.M., Warner, L.R., Hou, S.T., Jiang, S.X., Gomez-Isla, T., Leenhouts, K.M.,

947   Oxford, J.T., Feany, M.B., Masliah, E., and Rohn, T.T. (2007). Calpain-cleavage of

948   alpha-synuclein: connecting proteolytic processing to disease-linked aggregation. Am

949   J Pathol *170*, 1725-1738. 10.2353/ajpath.2007.061232.

950 24. Liu, C.W., Giasson, B.I., Lewis, K.A., Lee, V.M., Demartino, G.N., and Thomas, P.J.

951   (2005). A precipitating role for truncated alpha-synuclein and the proteasome in

952   alpha-synuclein aggregation: implications for pathogenesis of Parkinson disease. J

953   Biol Chem *280*, 22670-22678. 10.1074/jbc.M501508200.

954 25. Yan, C., Huo, H., Yang, C., Zhang, T., Chu, Y., and Liu, Y. (2018). Ubiquitin

955   C-Terminal Hydrolase L1 regulates autophagy by inhibiting autophagosome

956   formation through its deubiquitinating enzyme activity. Biochem Biophys Res

957   Commun *497*, 726-733. 10.1016/j.bbrc.2018.02.140.

958 26. Nicholson, B., Leach, C.A., Goldenberg, S.J., Francis, D.M., Kodrasov, M.P., Tian, X.,

959   Shanks, J., Sterner, D.E., Bernal, A., Mattern, M.R., et al. (2008). Characterization of

960   ubiquitin and ubiquitin-like-protein isopeptidase activities. Protein Sci *17*, 1035-1043.

961   10.1110/ps.083450408.

962 27. Cubenas-Potts, C., Goeres, J.D., and Matunis, M.J. (2013). SENP1 and SENP2 affect

963   spatial and temporal control of sumoylation in mitosis. Mol Biol Cell *24*, 3483-3495.

964   10.1091/mbc.E13-05-0230.

965 28. Kovacs, G.G., Wagner, U., Dumont, B., Pikkarainen, M., Osman, A.A.,

966   Streichenberger, N., Leisser, I., Verchere, J., Baron, T., Alafuzoff, I., et al. (2012). An

967   antibody with high reactivity for disease-associated alpha-synuclein reveals extensive

brain pathology. Acta Neuropathol *124*, 37-50. 10.1007/s00401-012-0964-x.

29. Vaikath, N.N., Majbour, N.K., Paleologou, K.E., Ardah, M.T., van Dam, E., van de

Berg, W.D., Forrest, S.L., Parkkinen, L., Gai, W.P., Hattori, N., et al. (2015).

Generation and characterization of novel conformation-specific monoclonal

antibodies for alpha-synuclein pathology. Neurobiol Dis *79*, 81-99.

10.1016/j.nbd.2015.04.009.

30. Guo, C., and Henley, J.M. (2014). Wrestling with stress: roles of protein

SUMOylation and deSUMOylation in cell stress response. IUBMB Life *66*, 71-77.

10.1002/iub.1244.

31. Kessler, J.C., Rochet, J.C., and Lansbury, P.T., Jr. (2003). The N-terminal repeat

domain of alpha-synuclein inhibits beta-sheet and amyloid fibril formation.

Biochemistry *42*, 672-678. 10.1021/bi020429y.

32. Qin, Z., Hu, D., Han, S., Hong, D.P., and Fink, A.L. (2007). Role of different regions

of alpha-synuclein in the assembly of fibrils. Biochemistry *46*, 13322-13330.

10.1021/bi7014053.

33. Zhang, S., Zhu, R., Pan, B., Xu, H., Olufemi, M.F., Gathagan, R.J., Li, Y., Zhang, L.,

Zhang, J., Xiang, W., et al. (2023). Post-translational modifications of soluble

alpha-synuclein regulate the amplification of pathological alpha-synuclein. Nat

Neurosci *26*, 213-225. 10.1038/s41593-022-01239-7.

34.   Abeliovich, A., Schmitz, Y., Farinas, I., Choi-Lundberg, D., Ho, W.H., Castillo, P.E.,

988   Shinsky, N., Verdugo, J.M., Armanini, M., Ryan, A., et al. (2000). Mice lacking

989   alpha-synuclein display functional deficits in the nigrostriatal dopamine system.

990   Neuron *25*, 239-252. 10.1016/s0896-6273(00)80886-7.

991   35.   Taguchi, K., Watanabe, Y., Tsujimura, A., Tatebe, H., Miyata, S., Tokuda, T., Mizuno,

992   T., and Tanaka, M. (2014). Differential expression of alpha-synuclein in hippocampal

993   neurons. PLoS One *9*, e89327. 10.1371/journal.pone.0089327.

994   36.   Watanabe, Y., Tatebe, H., Taguchi, K., Endo, Y., Tokuda, T., Mizuno, T., Nakagawa,

995   M., and Tanaka, M. (2012). p62/SQSTM1-dependent autophagy of Lewy body-like

996   alpha-synuclein inclusions. PLoS One *7*, e52868. 10.1371/journal.pone.0052868.

997   37.   Mao, X., Ou, M.T., Karuppagounder, S.S., Kam, T.I., Yin, X., Xiong, Y., Ge, P.,

998   Umanah, G.E., Brahmachari, S., Shin, J.H., et al. (2016). Pathological alpha-synuclein

999   transmission initiated by binding lymphocyte-activation gene 3. Science *353*.

1000   10.1126/science.aah3374.

1001   38.   Taguchi, K., Watanabe, Y., Tsujimura, A., and Tanaka, M. (2016). Brain

1002   region-dependent differential expression of alpha-synuclein. J Comp Neurol *524*,

1003   1236-1258. 10.1002/cne.23901.

1004

1005