Multidirectional effect of low-intensity electrical myostimulation on gene expression and phenotype in thigh and calf muscles after one week of disuse

Running title Disuse and electromyostimulation

Anna A. Borzykh[1], Roman Y. Zhedyaev[1], Ivan I. Ponomarev[1], Tatiana F. Vepkhvadze[1],

Viktor G. Zgoda[2], Mira A. Orlova[1], Nikita E. Vavilov[2], Egor M. Lednev[1], Kristina A. Sharlo[1],

Anastasia R. Babkova[1], Pavel A. Makhnovskii[1], Boris S. Shenkman[1], Ilya V. Rukavishnikov[1],

Oleg I. Orlov[1], Elena S. Tomilovskaya[1,*], Daniil V. Popov[1,*]

[1] Institute of Biomedical Problems, Russian Academy of Sciences, Moscow, Russia;

[2] V. N. Orekhovich Research Institute of Biomedical Chemistry, Moscow, Russia

Anna A. Borzykh and Roman Y. Zhedyaev have contributed equally to this work.

Corresponding author:

Daniil V. Popov, e-mail: danil-popov@yandex.ru, https://orcid.org/0000-0002-3981-244X

Elena S. Tomilovskaya, e-mail: finegold@yandex.ru, https://orcid.org/0000-0002-9030-6876

Abstract

Aim. Low-intensity neuromuscular electrical stimulation was offered as a safe (non-traumatic) approach to prevent the loss of muscle mass, strength, and endurance in patients with severe chronic diseases and in spaceflight. However, the effects of this approach on various leg muscles are poorly investigated.

Methods. This study assessed the efficiency of low-intensity (~10% of maximal voluntary contraction) electrical stimulation in preventing the negative effects of weekly disuse (dry immersion without and with daily stimulation) on the strength and aerobic performance of the ankle plantar flexors and knee extensors, mitochondrial function in permeabilized muscle fibers, and the proteomic (quantitative mass spectrometry-based analysis) and transcriptomic (RNA-sequencing) profiles of the soleus muscle and vastus lateralis muscle.

Results. Application of electrical stimulation during dry immersion prevented a decrease in the maximal strength and a slight reduction in aerobic performance of knee extensors, as well as a decrease in maximal (intrinsic) ADP-stimulated mitochondrial respiration and changes in the expression of genes encoding mitochondrial, extracellular matrix, and membrane proteins in the vastus lateralis muscle. In contrast, for the ankle plantar flexors/soleus muscle, electrical stimulation had a positive effect only on maximal mitochondrial respiration, but accelerated the decline in the maximal strength and muscle fiber cross-sectional area, which appears to be related to the activation of inflammatory genes.

Conclusion. The data obtained open up broad prospects for the use of low-intensity electrical stimulation to prevent the negative effects of disuse for "mixed" muscles, meanwhile, the optimization of the stimulation protocol is required for "slow" muscles.

Keywords: dry immersion, skeletal muscle, electromyostimulation, transcriptome, proteome, inflammation

Practitioner Points

- Low-intensity electrical myostimulation is often used as an alternative to exercise and high-intensity electrical stimulation to prevent the loss of muscle mass and function in patients with severe chronic diseases and in spaceflight. However, its effect on muscles with different functional capacities remains uncertain.

- One week of disuse (dry immersion) lead to a comparable decrease in the maximal strength and (intrinsic) mitochondrial respiration in both the ankle plantar flexors/soleus muscle and the knee extensors/vastus lateralis muscle. Meanwhile changes in gene expression (transcriptome) were three times more pronounced in the soleus muscle than in the vastus lateralis muscle.

- Application of electrical stimulation during disuse prevented most of the negative effects of disuse in the knee extensors/vastus lateralis muscle, but accelerated the decline in the maximal strength/muscle fiber cross-sectional area in the ankle plantar flexors/soleus muscle, which may be related to the activation of genes regulating the inflammatory response.

Introduction

The initial weeks of disuse (e.g., leg immobilization, bedrest, dry immersion), as well as spaceflight, cause a decrease in leg muscle strength and mass (Narici and de Boer 2011; Hackney and Ploutz-Snyder 2012; Koryak 2020; Vikne et al. 2020; Marusic et al. 2021) at a rate of 0.4%–0.5% per day (Narici and de Boer 2011; Hackney and Ploutz-Snyder 2012). Additionally, disuse leads to a reduction in the aerobic performance of the organism, measured as the maximum oxygen consumption rate in an incremental test on a cycle ergometer or treadmill, at a rate of 0.4%–1% per day (Ade et al. 2015; Ried-Larsen et al. 2017). Whole-body

maximal aerobic performance is influenced by "central" factors (cardiac output) and "peripheral" factors (muscle O_2 diffusion and oxidative capacity). However, only a few studies have investigated the effect of disuse on aerobic performance using tests that involve small muscle mass – an approach eliminating the influence of "central" factors – such as tests focused solely on the knee extensors (Ringholm et al. 2011; Salvadego et al. 2011; Salvadego et al. 2016; Salvadego et al. 2018; Zuccarelli et al. 2021).

Low-intensity (<30% of maximal voluntary contraction) neuromuscular electrical stimulation (NMES) was offered as a safe (non-traumatic) approach and alternative to exercise and high-intensity NMES to prevent the loss of muscle mass, strength, and endurance in patients with severe chronic diseases (Quittan et al. 2001; Harris et al. 2003; Zanotti et al. 2003; Nuhr et al. 2004; Karavidas et al. 2010), as well as in spaceflight (Mayr et al. 1999; Yarmanova et al. 2015; Maffiuletti et al. 2019). However, the effects of low-intensity NMES on strength and aerobic performance in various leg muscles are poorly investigated. This study assessed, for the first time, the efficiency of low-intensity NMES in preventing the negative effects of strict disuse using a dry immersion model. Specifically, we investigated the effects of a one-week dry immersion, both with and without NMES, on the strength and aerobic performance of leg muscles (test involving small muscle mass), mitochondrial function in permeabilized muscle fibers, and the proteomic (quantitative mass spectrometry-based analysis) and transcriptomic (RNA-sequencing) profiles of skeletal muscles. Some effects of disuse (especially during dry immersion and spaceflight) are pronounced differently in leg muscles with different functional capacities. For instance, the soleus muscle – a muscle with strong postural function, which consists predominantly of type I slow muscle fibers, shows more pronounced effects than other leg muscles (LeBlanc et al. 1992; Kozlovskaya et al. 2007; Fitts et al. 2010; Tomilovskaya et al. 2019; Casuso et al. 2024). Therefore, we examined the effect of dry immersion, both with and without NMES, on the ankle plantar flexors/soleus muscle and the knee extensors/vastus lateralis muscle.

Methods

Ethical approval

The studies were approved by the Committee on Biomedical Ethics of the Institute of Biomedical Problems of the RAS (protocols № 594 of September 6, 2021, and № 620 of July 12, 2022) and complied with all the principles set forth in the Declaration of Helsinki. All volunteers gave written informed consent to participate in the study and to undergo skeletal muscle biopsy.

Study design

Ten males participated in a control 7-day dry immersion (DI, ages 25–38 years; median height 1.78 m [interquartile range 1.71–1.79 m]; body mass 70 kg [64–74 kg]; and body mass index 24 kg/m² [21–24 kg/m²]). A year later (at the same time of year), another ten males participated in a 7-day dry immersion with NMES (DI+NMES, ages 26–39 years; height 1.74 m [1.70–1.76 m]; body mass 73 kg [70–79 kg]; and body mass index 24 kg/m² [23–26 kg/m²]). Study design and test protocols were similar in both experiments.

To evaluate the effects of NMES, changes induced by DI were compared with changes induced by DI+NMES. After a familiarization session, one day before and 8–9 hours after the end of DI or DI+NMES, the maximum isometric voluntary contraction (MVC) of the ankle plantar flexors and knee extensors of the right leg and (after 10 minutes of rest) the aerobic performance of these muscles (maximum power in an incremental ramp test till exhaustion, W_{max}) were examined (Fig. 1). Muscle samples were taken using a Bergstrom needle with aspiration under local anesthesia (2 ml of 2% lidocaine) from the medial part of the vastus lateralis (VL) and soleus (Sol) muscles of the left leg 14 days before and 6 days after DI and DI+NMES (at 10:00, 3 hours after a standardized very light breakfast: 5.2 g protein, 2.7 g fat, 55 g carbohydrates, 1253 kJ). This biopsy time was chosen because various physiological tests were

performed immediately after the end of the 7-day dry immersion (not included in this study). The samples were cleared of visible fragments of connective and adipose tissue; part of the fresh tissue was used to assess mitochondrial function, another part was immediately frozen in liquid nitrogen to study the transcriptome and proteome profiles, and the last part was embedded in Tissue-Tek O.C.T. Compound (Sakura, USA) and frozen in liquid nitrogen to histological study; all frozen samples were stored at -80 ℃ until further analyses.

Fig. 1. Design of the study. Dry immersion without (DI) and with neuromuscular electrical stimulation (DI+NMES).

Neuromuscular electrical stimulation

During DI+NMES, low-intensity, low-frequency electrical stimulation of the anterior and posterior surfaces of calf and thigh muscles was performed, as described elsewhere (Poltavskaya et al. 2021), daily at 11:00 using a stimulator Stimul-01 (Biophyspribor, Russia) for 40–45 minutes. The stimulation protocol was as follows: 1 second of bipolar symmetrical rectangular pulses (1 ms, 25 Hz), followed by a 2-second pause. Every day at 17:00, low-intensity high-frequency electrical stimulation was performed, as described elsewhere (Amirova et al. 2022), on the anterior and posterior surfaces of the calf muscles, followed by the thigh muscles, using a stimulator Amplidin-EST (NTC UP RAN, Russia) for 10 minutes. This stimulation protocol consisted of 10 seconds of bipolar symmetrical pulses (2.5 kHz with a sinusoidal modulating frequency of 50 Hz), followed by a 50-second pause. Pair of electrodes (PG-901, Fiab, Italy) was placed on the proximal and distal thirds of the thigh and calf for the anterior (13 cm x 5 cm) and

posterior (13 cm x 5 cm) surfaces. For the posterior surface of the calf, one electrode was placed in the proximal part of the calf (13 cm x 5 cm), and another electrode (9 cm x 5 cm) was placed 5 cm distal to the gastrocnemius muscles. The intensity of stimulation was individually adjusted to the point of causing unpleasant sensations. The mean voltage [standard deviation] of low-frequency stimulation was 19.2 ± 4.2 V for the anterior thigh, 17.7 ± 4.6 V for the anterior calf, 17.6 ± 4.2 V for the posterior thigh, and 18.2 ± 4.3 V for the posterior calf. The mean current during high-frequency stimulation was: for both anterior and posterior surfaces of the thigh – 18.3 ± 5.2 mA and calf – 10.5 ± 2.5 mA. NMES caused visible contractions of the stimulated muscles (~10% MVC for both knee extensors and ankle plantar flexors, evaluated without stimulation of antagonist muscles).

Maximum voluntary contraction of the ankle plantar flexors and knee extensors

Isometric MVC was assessed in a sitting position (Supporting information Fig. S1). To reduce the contribution of the biarticular gastrocnemius muscles to force production (Cresswell et al. 1995; Niess et al. 2018), MVC of ankle plantar flexors was examined at a knee joint angle of 70 degrees, the upright position of the calf, and the ankle joint 90 degrees using a CalfRaise-PRO dynamometer (AntexLab, Russia). A force transducer was pressed down to the knee joint (200 N) and then reset to zero.

The MVC of the knee extensors was examined in a sitting position on a Pro System 3 dynamometer (Biodex, USA) at a knee joint angle of 110 degrees. In each test, the subject performed 5–7 attempts with a rest period of 30 seconds between attempts; the best attempt was taken into account.

Maximum aerobic performance of the ankle plantar flexors and knee extensors

Maximum aerobic power (W_{max}) of the ankle plantar flexors was assessed in an incremental ramp test to exhaustion in a sitting position, as described above. Additionally, the

distal part of the foot was placed on a 4 cm high stand (Supporting information Fig. S1); the subject performed rhythmic extensions of the metatarsophalangeal joint (raising the calf with 55% of the maximum amplitude): raising – 1.1 s, lowering – 0.3 s, rest – 0.6 s. The rhythm and amplitude of movement were set using biofeedback (analog-to-digital converter E14-440, L-CARD, Russia and PowerGraph 3 software, Interoptica-c, Russia). The incremental test was carried out using the CalfRaise-PRO ergometer: the initial load for raising the calf was 30 N, the load increment was 25 N/min, and the load during calf lowering was minimal. The test was stopped when the subject was unable to maintain the full amplitude at the given rhythm for five cycles. The power for each cycle was calculated as

$$W_i = \frac{F_i \times A_i}{T_i} \tag{1}$$

where i – cycle number, F_i – average load during muscle contraction in a cycle (N); A_i – amplitude of movement in a cycle (m); T_i – cycle duration (s). The dependence "W_i – time" was smoothed (moving average with a window of 20 s), then W_{max} was defined as the maximum value (Supporting information Fig. S1).

The maximum aerobic power of the knee extensors was assessed using a modified Ergoselect 900 cycle ergometer (Ergoline, Germany): each subject performed rhythmic knee extensions (from 80 to 160 degrees) in a reclining sitting position (hip joint 35 degrees) (Supporting information Fig. S2). The extension frequency (1 cycle/s) was set by a sound command, the initial load – 0 W, the load increment – 5 W/min, and the load during flexion was minimal. During the test, constant cadence (1 cycle/s) was simulated on the ergometer cadence transducer. The amplitude of knee extension was ~80 degrees, but the frequency of extension could be different from the specified (simulated) one (because of fatigue), therefore the power per cycle was calculated as

$$W_i = W_{Es;\, i} \times \frac{\varphi_i}{360} \times \frac{T}{T_i} \tag{2}$$

where i – cycle number, $W_{Es;\, i}$ – power per cycle set on the Ergoselect 900 ergometer (W); φ_i – change in the knee joint angle per cycle (degrees); T and T_i – specified and actual cycle duration,

respectively (s). The knee joint angle was recorded using the analog-to-digital converter E14-440 and PowerGraph 3 software. The test was stopped when the subject was unable to maintain the full amplitude in the given rhythm for 5 cycles. The dependence "W_i – time" was smoothed (moving average with a window of 20 s), and then W_{max} was defined as the maximum value (Supporting information Fig. S2).

Muscle fiber cross-sectional area

Muscle transverse sections (10 μm) were prepared using a cryotome (CM 1900, Leica Biosystems, Germany), and immunostaining was performed as previously described (Sharlo et al. 2019), using primary antibodies for slow (1:100, M8421, RRID:AB_477248, Sigma, USA) and fast muscle fibers (1:100, ACC 232, RRID:AB_3106894, Leibniz Institute DSMZ, Germany), as well as secondary antibodies AlexaFluor 488 and 546 (A32723, RRID:AB_2633275 and A11035 RRID:AB_2534093), respectively, all 1:400, Thermo Fisher Scientific, USA). Images were acquired using a fluorescent microscope Olympus IX83P2ZF3 (Olympus, Japan) and analyzed by the ImageJ software; at least 100 fibers of each type were evaluated.

Mitochondrial respiration in permeabilized muscle fibers

A piece of fresh tissue (~10 mg) was placed in ice-cold BIOPS relaxation buffer (10 mM Ca-EGTA buffer, 0.1 μM free calcium, 20 mM imidazole, 20 mM taurine, 50 mM K-MES, 0.5 mM DTT, 6.56 mM $MgCl_2$, 5.77 mM ATP, 15 mM phosphocreatinine, pH 7.1) to remove connective and adipose tissue. Then a small piece (~2 mg) was removed and the muscle fibers were separated using a pair of needles to form a bundle. The fibers were permeabilized in BIOPS buffer with saponin (50 μg/ml, 30 min) on ice with slow stirring, then the tissue was washed 2 times for 10 min in MiRO5 buffer (0.5 mM EGTA, 3 mM $MgCl_2 \cdot 6\ H_2O$, 60 mM lactobionic acid, 20 mM taurine, 10 mM KH_2PO_4, 20 mM HEPES, 110 mM D-sucrose, 1 g/l fatty acid-free

bovine serum albumin). The rate of mitochondrial respiration was assessed simultaneously on two channels on an Oxygraph polarograph (Hansatech, United Kingdom) under hyperoxic conditions (Co_2 >200 μM), sequentially adding: malate (2 mM) + pyruvate (5 mM) + glutamate (10 mM), ADP (5 mM) + $MgCl_2$ (3 mM), succinate (10 mM), cytochrome c (10 μM) and then the FCCP uncoupler was titrated (0.25 μM to 1.5 μM). The respiration rate was normalized to the mass of permeabilized muscle fibers.

RNA sequencing and data processing

A piece of frozen tissue (~15 mg) was homogenized at +4 °C in 500 μl of a lysis buffer (ExtractRNA, Evrogen, Russia) using a plastic pestle and a drill. Total RNA was isolated using a spin column (RNeasy Mini Kit, Qiagen, Germany). RNA concentration was assessed using a Qubit 4 fluorimeter (ThermoScientific, USA); RNA integrity – using capillary gel electrophoresis (TapeStation, Agilent, USA). All samples had RNA integrity (RIN) >7.8.

Strand-specific RNA libraries were prepared, as described previously (Kurochkina et al. 2024), using the NEBNext Ultra II RNA kit (New England Biolabs, USA) according to the manufacturer's recommendations. Library efficient concentration was evaluated by qPCR using 5X qPCRmix-HS SYBR kit (Evrogen, Russia); single-end sequencing was done by a NextSeq 550 analyzer (Illumina, USA), according to the manufacturer's recommendations, with a read length of 75 bp (~60 million reads/sample).

Sequencing quality was assessed using the FastQC tool (v.0.11.5, RRID:SCR_014583), and low-quality reads were removed using Trimmomatic (v.0.36, RRID:SCR_011848). High-quality reads were mapped to the human reference genome GRCh38.p13 primary assembly. The number of unique reads mapped to exons of each gene was determined using the Rsubread package (R environment) and Ensembl annotation (GRCh38.101). The DESeq2 method (paired sample analysis with Benjamini-Hochberg correction) was used to analyze differential gene expression. Differentially expressed genes were defined as protein-coding genes (as well as

polymorphic pseudogenes and translated pseudogenes) with an expression level of TPM >1 (Transcripts Per Million, kallisto tool v0.46.2), p_{adj}<0.01 and |fold change|≥1.25.

Quantitative mass spectrometry-based proteomic analysis

A piece of frozen tissue (~10 mg) was homogenized as described above in 140 µl of lysis buffer (4% sodium dodecyl sulfate, 0.1 M Tris and 0.1 M dithiothreitol, pH 7.6). The lysate was boiled (95 ºC, 5 min), transferred to an AFA microtube and sonicated (average power 20 W, 30 s × 4 using a ME220 sonicator (Covaris, USA), and centrifuged (5 min, 30,000 g). The protein concentration in the supernatant was measured by a fluorimeter Qubit 4.

The proteins (85 µg) were diluted in lysis buffer (final volume 24 µl), transferred to an S-Trap column (ProtiFi, USA), alkylated (20 mM iodoacetamide, 15 min), and hydrolyzed into the S-Trap column using trypsin and Lys-C (2 h at 47 °C, 1:15 and 1:30, respectively, all Promega, USA) in 40 µl 50 mM triethylammonium bicarbonate according to the manufacturer's recommendations. After elution, 20 µg of peptides (fluorimeter Qubit 4) were dried, resuspended in 100 mM triethylammonium bicarbonate, and labeled with TMT 10-plex or 16-plex isobaric labels (Thermo Scientific, USA) for 1.5 h according to the manufacturer's recommendations. The reaction was stopped (0.3% hydroxylamine w/v, 15 min), and samples were pooled.

The mixture of labeled peptides was separated using high pH reverse-phase LC fractionation (HPLC 1200, Agilent, USA). The peptides were concentrated on an analytical column (XBridge C18, particle size 5 µm, 4.6 x 250 mm, Waters, Ireland) in isocratic mode at a flow of 750 µl/min for 3 min in mobile phase A (15 mM ammonium acetate, pH 9,0). Twenty-four fractions were collected from 3 min to 50 min (collection time 2 min, volume 1500 µL) using a gradient elution mode with mobile phase B (15 mM ammonium acetate, pH 9.0, 80% acetonitrile, pH 9.0). Each fraction was concentrated, and then fractions (1 and 13, etc.) were combined to obtain 12 fractions.

Each fraction was analyzed three times using an HPLC Ultimate 3000 RSLC nanosystem (Thermo Scientific, USA) and a Q Exactive HF-X Hybrid Quadrupole-Orbitrap mass spectrometer (Thermo Scientific, USA) by the nanoelectrospray ion source in the positive mode of ionization (Thermo Scientific) as previously described (Kurochkina et al. 2024). The gradient (90 min) was formed by the mobile phase A (0.1% formic acid) and B (80% acetonitrile, 0.1% formic acid) at a 0.4 µL/min flow. The ionizing voltage was 2.1 kV. MS spectra were acquired at a resolution of 60,000 in the 390–1300 m/z range; fragment ions were mass scanned at a resolution of 60,000 at the range from m/z 120 to the upper m/z value as assigned by the mass to charge state of the precursor ion. All tandem MS scans were performed on ions with a charge state from z = 2+ to z = 4+. Synchronous precursor selection facilitated the simultaneous isolation of up to 40 MS2 fragment ions. The maximum ion accumulation times were set to 50 ms for precursor ions and 25 ms for fragment ions. AGC targets were set to 10^6 and 10^5 for precursor ions and fragment ions, respectively.

Search and identification of peptides and proteins were performed using a MaxQuant platform (2.1.4.0; Max Planck Institute for Biochemistry, RRID:SCR_014485) with default settings (FDR for peptides 1%, N-terminal acetylation and oxidation of methionine as variable modifications and carbamidomethylation of cysteine, as fixed modification). Further analysis was performed using a Perseus platform (1.6.5; Max Planck Institute for Biochemistry, RRID:SCR_015753). To avoid the batch effect associated with labeling with the TMT 10-plex and 16-plex kits, the average reporter ion intensity of all samples labeled with 10-plex isobaric tags was normalized to the intensity of all samples labeled with 16-plex tags, as previously described (Kurochkina et al. 2024). After filtration (potential contaminants, reverse peptides, peptides identified by site only), proteins identified by more than one peptide (unique+razor) and present in >70% of samples were taken for further analysis. Differentially expressed proteins were determined using one-way ANOVA with q-value (Benjamini-Hochberg adjusted p-value) <0.05 and Tukey's *post-hoc* test (p<0.05).

Statistical analysis

Data are presented as median and interquartile range. The nonparametric Wilcoxon test was used to evaluate the significance of changes after immersion or between different muscles. To compare changes between DI and DI+NMES, the nonparametric Mann-Whitney test was used. The significance level was 0.05.

Functional enrichment analysis for biological processes, cellular components and pathways (UNIPROT KW BP/CC, GENE ONTOLOGY BP/CC DIRECT and KEGG PATHWAY databases) was performed against all protein-coding genes (TPM >1, ~10,000 mRNA) or detected proteins using the DAVID tool v2023q4 (RRID:SCR_001881) (p_{adj}<0.05, Fisher's exact test with Benjamini correction).

Results

Strength and endurance of knee extensors and ankle plantar flexors

DI resulted in comparable reductions in knee extensor and ankle plantar flexor MVC (7% and 4%, respectively, p<0.05; Fig. 2A). NMES had different effects on different muscles: it prevented the dry immersion-induced decrease in MVC in the knee extensors (p = 0.9), but unexpectedly increased the drop in strength in the ankle plantar flexors (up to 16%, p = 0.01). Moreover, a trend toward a difference was found between the magnitude changes in these measures in DI and DI+NMES for both muscle groups (p = 0.06–0.09), indicating the multidirectional effect of NMES on various muscle groups (Fig. 2A).

DI caused a very small decrease in aerobic performance in the knee extensors only (3%, p = 0.049) (Fig. 2B). In contrast, DI+NMES led to an increase in this measure (by 5%, p = 0.039); moreover, the magnitude of change in these measures after DI differed (p = 0.01) from that after

DI+NMES, underlining the positive effect of NMES on VL. Both DI and DI+NMES had no effect on the aerobic performance of the ankle plantar flexors (Fig. 2B).

Muscle fiber cross-sectional area and mitochondrial function in the vastus lateralis and soleus muscles

After DI, a tendency to decrease in CSA was observed in type II fibers in VL (by 8.5%, $p = 0.08$); no changes were found in type I fibers in VL and in both fiber types in Sol (Fig. 2C). DI+NMES had no effect on both fiber types in VL and type I fibers in Sol, but reduced CSA in type II fibers in Sol (by 15%, $p = 0.01$; Fig. 2C, which is consistent with the loss of MVC in ankle plantar flexors; Fig. 2A). Interestingly, the magnitude of change in this measure after DI differed ($p = 0.04$) from that after DI+NMES, underlining the negative effect of NMES on Sol.

DI reduced maximum ADP-stimulated mitochondrial respiration in permeabilized VL and Sol muscle fibers by 17% to 20% ($p<0.03$). In contrast, DI+NMES increased this measure by ~50% ($p<0.04$) in both muscles (Fig. 2D). Moreover, in both muscles, the magnitude of change in these parameters after DI differed ($p<0.01$) from that after DI+NMES, underlining a strong positive effect of NMES on both muscles (Fig. 2D).

Fig. 2. Electrical stimulation prevented the dry immersion-induced decrease in maximal voluntary contraction (MVC) and a slight decrease in the endurance of the knee extensors, but increased the drop in both MVC and type II muscle fibers cross-sectional area (CSA) in the ankle plantar flexors/soleus muscle.

A and B – Changes in isometric maximum voluntary contraction (A) and aerobic performance (maximum power in the incremental ramp test, W_{max}) (B) in the knee extensors and ankle plantar flexors after dry immersion without (DI) and with neuromuscular electrical stimulation (DI+NMES); ($n = 9$–10 subjects in each group).

C and D – Changes in the cross-sectional area of type I and II muscle fibers (C, $n = 6$–10 in each group) and maximum ADP-stimulated mitochondrial respiration (Jo_2) in permeabilized fibers (D) in VL and Sol muscles; ($n = 9$ subjects in each group).

Median, interquartile range, and individual values are shown. The lower *p*-values refer to changes from the baseline (dashed line) (Wilcoxon signed-rank test), the upper ones refer to differences between the magnitude of changes in DI and DI+NMES (Mann-Whitney test).

Proteomic profile in the vastus lateralis and soleus muscles

A total 1890 proteins were detected in both muscles: mainly highly abundant myofibrillar, mitochondrial, cytoskeletal, sarcomeric, and ribosomal proteins, as well as heat shock and sarcoplasmic reticulum proteins (Supporting information Table S1). In VL, the relative content of 724 proteins was lower than in Sol (Fig. 3A); the majority of them (399 proteins) enriched functional categories related to mitochondrial proteins and regulators of fatty acid and carbohydrate metabolism (Fig. 3B, Supporting information Table S1 and S2). The relative content of 64 proteins was higher in VL than in Sol; these proteins were associated with calcium signaling (14 proteins), sarcomere (12 proteins), glycolysis (12 proteins), and glycogen metabolism (4 proteins) (Fig. 3B, Supporting information Table S1 and S2). In VL, the relative content of the "fast" myosin heavy chains 2X and 2A, encoded by the *MYH1* and *MYH2* genes, was 1.7 and 1.8 times higher ($q<0.01$) than in Sol, respectively, meanwhile the "slow" myosin heavy chain (*MYH7* gene) was 1.4 times less ($q<0.01$) (Supporting information Table S1). The marked difference in the proteomic profiles of VL and Sol is consistent with differences in protein expression between type I and type II muscle fibers showed in the human VL (Murgia et al. 2017; Murgia et al. 2021), as well as with differences in the transcriptomic profiles of these muscles and in the maximum rate of mitochondrial respiration in the baseline (Fig. 3A, C, and D and see below).

Further analysis showed that dry immersion had no effect on the protein content of detected proteins of either muscle. It should be noted that, because of the high dynamic range of the muscle proteome (Deshmukh et al. 2015), the proteomic profile in this study largely did not

include low abundant regulatory/signaling proteins. These proteins usually have a short half-life (Schwanhausser et al. 2011; Fornasiero et al. 2018) and thereby potentially could significantly change their relative content after 6 days of disuse.

Fig. 3. Pronounced differences between the vastus lateralis (VL) and soleus (Sol) muscles in the baseline (prior to dry immersion) for proteomic and transcriptomic profiles and mitochondrial respiration.

A – number up- and down-regulated mRNA/proteins in VL *vs.* Sol in the baseline ($n = 10$ subjects in each group). All detected protein-coding genes and proteins are presented in Supporting information Table S1.

B and C – Tables show the number of mRNA/proteins and the corresponding p_{adj} for the main enriched functional categories (all results of enrichment analysis are presented in Supporting information Table S2).

D – Maximum uncoupled and ADP-stimulated mitochondrial respiration (Jo_2) in permeabilized fibers in VL and Sol in the baseline. Median, interquartile range, and individual values are shown. p-values refer to differences between the magnitude of change in DI and DI+NMES (Mann-Whitney test, $n = 9$ subjects in each group).

Transcriptomic profile in the vastus lateralis and soleus muscles

After the removal of low-expressed genes, ~10,000 mRNAs were detected (Supporting information Table S1). As expected, in VL, compared with Sol, up-regulated mRNAs (1133) enriched terms related to aerobic and fatty acid metabolism enzymes, contractile apparatus, cell adhesion, and ion transporters, and down-regulated mRNAs (1326) were associated with regulators of translation, glycolytic enzymes, transmembrane transporters, and contractile proteins (Fig. 3C, Supporting information Table S1 and S2).

After both DI and DI+NMES, changes in gene expression in Sol (~1600 mRNAs) were more pronounced (2.5–4 times) than in VL (~400 mRNAs) (Fig. 4A and B, Supporting information Table S3). Interestingly, NMES had different effects on the transcriptome of muscles with different functional capacities: it slightly decreased the number of genes that changed expression in the "mixed" fiber type composition VL (607 mRNAs and 442 mRNAs in DI and DI+NMES, respectively) and increased in the "slow" Sol (1527 mRNAs and 1791 mRNAs in DI and DI+NMES, respectively, Fig. 4A and C). Moreover, NMES significantly altered the transcriptomic response to immersion, inducing the expression of multiple genes that showed no overlap with genes that responded to DI (263 and 1046 mRNAs in VL and Sol, respectively) (Fig. 4C).

The small number of up-regulated genes in VL after DI (215 mRNAs) enriched no functional categories (Fig. 4A and D, Supporting information Table S4). In contrast, after DI+NMES, up-regulated genes in VL (164 mRNAs) were associated with secreted proteins (30

mRNAs), plasma membrane proteins (61 mRNAs), and lipid transport (9 mRNAs) (Fig. 4A and D, Supporting information Table S4), which can be interpreted as a positive effect of NMES. In Sol, DI caused a large increase in gene expression (567 mRNAs) that was associated with numerous plasma membrane proteins (168 mRNAs), as well as extracellular (56 mRNAs) and immune response proteins (23 mRNAs) (Fig. 4A and D, Supporting information Table S4). Interestingly, in Sol after DI+NMES, the number of up-regulated genes slightly increased (663 mRNAs); these genes enriched functional categories associated with the extracellular region (72 mRNAs), inflammatory response (28 mRNAs), and proteasome (18 mRNAs) (Fig. 4A and D, Supporting information Table S4). In addition, several genes encoding NF-kappa B inflammatory pathway proteins, cyto/myokines, and their receptors were also up-regulated (Fig. 4D and 5, Supporting information Fig. S3).

Fig. 4. Electrical stimulation differently affected the transcriptomic response to dry immersion in the "mixed" vastus lateralis (VL) and "slow" soleus (Sol) muscles.

A – The number of mRNA that changed expression in dry immersion without (DI) and with neuromuscular electrical stimulation (DI+NMES); ($n = 10$ subjects in each group). All detected protein-coding genes are presented in Supporting information Table S3.

B – Transcriptomic response to DI (and DI+NMES) had little overlap between VL and Sol and was significantly more pronounced in Sol. The number of DEGs is presented.

C – NMES significantly altered the transcriptomic response to immersion in both muscles.

D and E – Functions of up- (D) and down-regulated (E) genes in DI and DI+NMES (all results of enrichment analysis are presented in Supporting information Table S4). The number of genes in each category is indicated, the heat map shows p_{adj}. N.S. – non significant.

Fig. 5. Application of neuromuscular electrical stimulation during dry immersion induced expression of mRNAs encoding proteins of the NF-kappa B inflammatory pathway in "slow" soleus muscle (Sol), but not in "mixed" vastus lateralis muscle (VL) (also see Supporting Information Fig S3).

DI and DI+NMES – dry immersion without and with neuromuscular electrical stimulation, respectivly; ($n = 10$ subjects in each group). The color of the circles and the heat map show the

magnitude of changes in various muscle after various interventions for each detected mRNA; *N.S.* – non-significant change.

After DI, a small number of down-regulated genes in VL (392 mRNAs) were associated with a variety of membrane (125 mRNAs), mitochondrial (67 mRNAs), and sarcomeric (15 mRNAs) proteins (Fig. 4A and E, Supporting information Table S3). NMES largely prevented these negative changes: down-regulated genes were associated only with sarcomeric proteins (12 mRNAs) (Fig. 4E, Supporting information Table S4). In contrast, in Sol, DI caused the down-regulation of many genes (960 mRNAs) associated with various mitochondrial (283 mRNAs), membrane (260 mRNAs), and sarcomeric proteins (26 mRNAs) (Fig. 4A and E, Supporting information Table S4). Unexpectedly, DI+NMES slightly increased the number of down-regulated genes in Sol (to 1128 mRNAs), especially genes encoding mitochondrial proteins (to 330 mRNAs: predominantly genes related to mitochondrial ATP synthesis coupled proton transport, tricarboxylic acid cycle, mitochondrial organization, and mitochondrial PDH complex) (Fig. 4A and E, Supporting information Table S4). Moreover, a more pronounced response (in terms of magnitude) was found for mRNAs encoding mitochondrial proteins, in particular, for the inner mitochondrial membrane (Supporting information Fig. S4). Additionally, in Sol, NMES induced a decrease in the expression of genes encoding regulators of fat metabolism (30 mRNAs) and peroxisomal proteins (21 mRNAs) (Fig. 4E).

Discussion

To assess the possibility of preventing the negative effects of disuse, the effects of a week-long dry immersion without and with low-intensity NMES on the functional capacity and gene expression in the thigh and calf muscles were compared for the first time. The generalized effects of DI and DI+NMES are presented in Fig. 6.

Parameters	Knee extensors or *VL*		Ankle flexors or *Sol*	
	DI	DI+NMES	DI	DI+NMES
Maximal voluntary contraction	↓	−	↓	↓↓
Muscle fiber cross-section area	−	−	−	↓↓
mRNA of sarcomeric proteins	−	−	−	−
Aerobic performance	↓	↑	−	−
Maximal mitochondrial respiration	↓	−/↑	↓	−/↑
mRNA of mitochondrial proteins	↓	−	↓↓	↓↓↓
mRNA of fat metabolism regulators	−	↑	−	↓
mRNA of immune response regulators	−	−	↑	−
mRNA of inflammatory response regulators	−	−	−	↑
mRNA of proteasomal related proteins	−	−	−	↑
mRNA of ECM proteins	−/↓	↑/−	↑↑/↓↓	↑↑/−
mRNA of membrane proteins	−/↓	↑/−	↑↑/↓↓	↑/−
mRNA of transmembrane proteins	↓↓	−	↑/↓↓↓	−

Fig. 6. Generalized effects of dry immersion without (DI) and with low-intensity neuromuscular electrical stimulation (DI+NMES) on the functional capacity of the knee extensors/vastus lateralis muscle (VL) and the ankle plantar flexors/soleus muscle (Sol). Arrows show the direction of change; the number of arrows reflects the severity of the change/the number of genes changed expression. Blue and red boxes indicate the positive and negative effects of NMES, respectively.

NMES prevented the decline in knee extensor strength but accelerated that in ankle plantar flexors

In this study, the decline in MVC after DI was comparable to or slightly less than in the literature (Marusic et al. 2021) and did not differ between knee extensors and ankle plantar flexors (Figs. 2 and 6). At baseline, the proteomic profiles in VL and Sol were markedly different (788 of 1890 detected proteins; Fig. 3). However, after 6 days of DI, no changes in the relative content of highly abundant muscle proteins, including sarcomeric proteins (constituting a half of the muscle protein mass), were found in either muscle. This may be explained by a similar decrease in the rate of synthesis of highly abundant muscle proteins and/or an increase in the rate of their degradation, as well as the fact that the expected decrease in muscle mass during weekly disuse is only a few percent (~2.4–3.0%) (Narici and de Boer 2011; Hackney and Ploutz-

Snyder 2012; Marusic et al. 2021); meanwhile, the half-life of many highly abundant proteins (sarcomeric and cytoskeletal proteins, extracellular matrix proteins, mitochondria and histones) is more than a week (Fornasiero et al. 2018; Rolfs et al. 2021). The absence of changes in the expression of highly abundant contractile (as well as mitochondrial) proteins is consistent with work examining the effects of 10 days of bedrest in individual VL fibers (>200) (Murgia et al. 2022). More than two hundred proteins were found to be up- or down-regulated (out of >7500 detected); down-regulated proteins predominantly regulated the interaction of the fiber with the extracellular matrix and force transmission. It should be noted that in this work for estimating relative protein content, a label-free approach with imputation of missing data was used, and the sample size was set by the number of fibers analyzed (rather than the number of samples/volunteers), which may explain the large differences in the number of detected and differentially expressed proteins with our study.

NMES had different effects on different muscle groups: it prevented the disuse-induced decrease of MVC in the knee extensors, but induced the loss of fiber cross-sectional area in Sol accelerating the decrease of MVC in ankle plantar flexors (Figs. 2 and 6). Low-intensity high- or low-frequency NMES (5%–30% MVC, 25–50 Hz, 10–60 min/day, 5–7/week, 3–8 weeks) was previously shown to increase the muscle protein synthesis rate, size, and MVC of "mixed" knee extensor muscles in various patients with decreased muscle mass (Gibson et al. 1988; Gibson et al. 1989; Quittan et al. 2001; Harris et al. 2003) and in healthy volunteers (Stefanovska and Vodovnik 1985). An increase in MVC of the ankle plantar flexors was also shown in healthy volunteers after low-intensity high-frequency NMES (20% MVC, 80 Hz, 0.5 h/day, 5/week, 2 weeks); this increase is likely due not only to the direct effect of the NMES on the muscle, but also to the neural adaptation (Bouguetoch et al. 2021). However, low-intensity low-frequency NMES of significantly longer duration (15 Hz, 6 h/day, 6/week, 4.5 weeks) reduced knee extensor MVC (by 17%) and type II muscle fiber CSA in VL (by 10%) in healthy volunteers (Shenkman et al. 2006; Shenkman et al. 2007). It can be assumed that a combination of the

effects of dry immersion and low-intensity NMES may cause similar negative effects on Sol even with a significantly shorter duration of NMES treatment. It is important to note, recruitment order during NMES is synchronous and random with respect to motor unit type, i.e. both type I and II muscle fiber were involved in contraction (Barss et al. 2018). The negative effect of a combination of NMES and dry immersion on "slow" Sol may be related to the fact that energy supply, proteomic profile, regulation of protein synthesis and breakdown (Fig. 3) (van Wessel et al. 2010; Ciciliot et al. 2013; Murgia et al. 2017; Murgia et al. 2021), and some molecular responses to disuse (e.g., activation of inflammatory response genes (see below); Fig. 5 and Supporting information Fig. S3) differ significantly between "slow" and "fast" muscle fibers/muscles. In addition, the negative effects of NMES on the ankle plantar flexors could be due to their small size, compare to the knee-extensors, resulting in the greater NMES-induced recruitment of muscle fiber (Maffiuletti 2010).

NMES prevented the decline in aerobic performance only in the knee extensors

The lack of a pronounced decrease in aerobic performance in both muscle groups (Figs. 2B and 6) and change in the relative content of more than three hundred detected mitochondrial proteins in both muscles after 6 days of DI (Supporting information Table S1 and S2), is in line with studies showing no changes in time to failure/maximum power in the one-legged knee extension test after 7-10-day bedrest (Ringholm et al. 2011; Salvadego et al. 2016), as well as activity/content of mitochondrial enzymes/mitochondrial density in VL after 5-10-day bedrest (Mikines et al. 1991; Salvadego et al. 2016; Zuccarelli et al. 2021; Marshall et al. 2022; Eggelbusch et al. 2024) and 5-11-day space flight (Edgerton et al. 1995). Taken together, these data and other studies suggest that pronounced reductions in markers of mitochondrial density (for review, see (Gram et al. 2014)) and aerobic performance at work of small muscle mass (Salvadego et al. 2011) should occur in a later stage of disuse. On the other hand, DI caused a

decrease in the maximum ADP-stimulated mitochondrial respiration rate in permeabilized fibers, which is consistent with several studies examining the effects of 6–10 days of disuse (Dirks et al. 2016; Dirks et al. 2020; Standley et al. 2020), but contradicts others (Salvadego et al. 2016; Edwards et al. 2020; Zuccarelli et al. 2021; Eggelbusch et al. 2024). The decrease in maximum mitochondrial respiration without a decrease in the relative content of mitochondrial proteins, indicates a decrease in intrinsic mitochondrial respiration related to dysregulation of the respiration (respiratory complexes). The latter corresponds to data on a sharp drop in the maximum (intrinsic) ADP-stimulated mitochondrial respiration rate after just three days of leg immobilization in VL (Miotto et al. 2019) and of dry immersion in Sol (Popov et al. 2023).

Interestingly, low-intensity NMES had a small positive effect on the aerobic performance of the knee extensors only (Figs. 2B and 6). This is consistent with studies showing that low-intensity low-frequency NMES (10–25 Hz, 5–8 weeks) increases maximal pulmonary oxygen uptake and maximal power in an incremental test with large muscle mass (meta-analyses (Sbruzzi et al. 2010; Smart et al. 2013)), as well as citrate synthase activity, a marker of mitochondrial density, in VL of patients with heart failure (Nuhr et al. 2004) and healthy volunteers (Nuhr et al. 2003; Shenkman et al. 2006; Shenkman et al. 2007). Since the NMES-induced increase in aerobic performance correlates with the total stimulation time (Smart et al. 2013), it can be assumed that the small effect of low-intensity stimulation for the thigh muscles is due to its insufficient duration, meanwhile the lack of effects for the calf/"slow" Sol muscles – with the high oxidative capacity of this muscle (and low ability to respond to a training stimulus) and/or with Sol-specific activation of inflammatory response genes induced by combined exposure to immersion and NMES (Fig. 5, Supporting information Fig. S3 and see below).

NMES has differential effects on the transcriptomic response to dry immersion in the soleus and vastus lateralis muscles

Unlike mass spectrometry-based proteomic analysis, RNA-seq makes it possible to detect all expressed protein-coding genes. In line with studies showing greater negative effect of various disuse models on phenotype of Sol than other leg muscles (LeBlanc et al. 1992; Kozlovskaya et al. 2007; Fitts et al. 2010; Tomilovskaya et al. 2019; Casuso et al. 2024), DI caused significantly greater transcriptome changes in Sol than in VL (Fig. 4). The more pronounced response in Sol and the direction of these changes are consistent with studies examining the effects of long-term (60 days) bedrest in human VL and Sol (Chopard et al. 2009) and 6 weeks of spaceflight in various leg muscles of mice (da Silveira et al. 2020). A comparison of the transcriptomic response in different muscles suggests that longer-term disuse should cause a decrease in the relative content of some highly abundant proteins (mitochondrial, sarcomeric, etc.) and in muscle function, and this decrease will be much more pronounced in Sol.

NMES significantly altered the transcriptomic response in both muscles, but the direction of these changes differed between VL and Sol. In VL, NMES reduced most of the negative transcriptomic changes caused by immersion and, moreover, increased the expression of a number of genes encoding fat metabolism regulators and extracellular matrix proteins (Figs. 4 and 6, Supporting information Table S4), which correlates with transcriptomic changes caused by regular aerobic exercise (Popov et al. 2019; Makhnovskii et al. 2021). In contrast, in Sol, DI+NMES increased the number of down-regulated genes (genes encoding mitochondrial proteins, proteasomes, and fat metabolism regulators) or the magnitude of their response (Fig. 4D and E and Supporting information Fig. S4). Moreover, in Sol, NMES induced genes encoding NF-kappa B pathway proteins, a number of cyto/myokines, and their receptors (Fig. 4D and 5, Supporting information Fig. S3) indicating an inflammatory response. Such response was previously shown in "slow" Sol on the first days of disuse (Popov et al. 2023), in contrast to VL (Abadi et al. 2009). The obtained data suggest that low-intensity NMES may amplify this effect in Sol. Activation of inflammatory signaling pathways has a negative impact on protein synthesis and breakdown, mitochondrial biogenesis, and other cellular processes (Costamagna et

al. 2015; Thoma and Lightfoot 2018; Abu Shelbayeh et al. 2023). It can be hypothesized that the activation of inflammatory genes partially explains the negative effects of NMES after a week of dry immersion: an increase in the number of down-regulated genes encoding mitochondrial proteins (and magnitude of their response) and muscle fiber cross-sectional area in Sol, a lack of preventive effect on aerobic capacity, and a negative effect on MCV in ankle plantar flexors.

Conclusions

The present work was the first to investigate *i*) the effect of a week-long dry immersion on the phenotype and gene expression in leg muscles with significantly differed functional capacities (in particular, mitochondrial respiration, proteomic and transcriptomic profile), and *ii*) the efficiency of low-intensity (~10% MVC) NMES to prevent the negative effects of dry immersion. Dry immersion-induced decreases in MVC in both muscles were comparable and occurred without a decrease in the relative content of contractile proteins, which appears to be due to disruption of neuromuscular mechanisms rather than a decrease in muscle mass. The decrease in maximum ADP-stimulated mitochondrial respiration was also comparable in both muscles and occurred without a decrease in the relative content of mitochondrial proteins/respiratory enzymes indicating dysregulation of intrinsic respiration. At the same time, it was found that changes in the transcriptome were significantly more pronounced (2.5–4 times, in terms of number of differentially expressed genes) in the "slow" soleus muscle than in the vastus lateralis muscle. This suggests that longer-term disuse will lead to a change (primarily a decrease) in the relative content of a number of highly abundant proteins in Sol and an accelerated decline in the functional capacity of the ankle plantar flexors.

Application of NMES during dry immersion prevented a decrease in MVC and a slight reduction in aerobic performance of knee extensors, as well as a decrease in maximal mitochondrial respiration in permeabilized fibers and changes in the expression of genes

encoding mitochondrial, extracellular matrix, and membrane proteins in the vastus lateralis muscle. In contrast, for the ankle plantar flexors/soleus muscle, NMES had a positive effect only on maximal mitochondrial respiration, but accelerated the decline in the MCV/muscle fiber cross-sectional area, which appears to be related to the activation of genes regulating the inflammatory response. The data obtained open up broad prospects for the use of low-intensity NMES to prevent the negative effects of disuse for "mixed" muscles, meanwhile, the optimization of the stimulation protocol is required for "slow" muscles.

Data availability statement

The data supporting these findings is available in the NCBI Gene Expression Omnibus (GEO) under accession number GSE271606 (secure token: mzaxawykznohful) and GSE271607 (secure token: enmtakquxneffmf) (RNA-seq datasets) and Supporting Information Table S1 (proteomic datasets).

Funding

The study was supported by the Ministry of Science and Higher Education of the Russian Federation under agreement № 075-15-2022-298 from 18 April 2022 about the grant in the form of subsidy from the federal budget to provide government support for the creation and development of a world-class research center, the "Pavlov Center for Integrative Physiology to Medicine, High-tech Healthcare and Stress Tolerance Technologies".

Conflict of interest disclosure

The authors declare that they have no competing interests.

Ethics approval statement

The studies were approved by the Committee on Biomedical Ethics of the Institute of Biomedical Problems of the RAS (protocols № 594 of September 6, 2021, and № 620 of July 12, 2022) and complied with all the principles set forth in the Declaration of Helsinki. All volunteers gave written informed consent to participate in the study and to undergo skeletal muscle biopsy.

Acknowledgements

The authors acknowledge the laboratory staffs at IBMP RAS for their excellent assistance in organization of dry immersion experiments, Motanova E.S. for assistance in the experiment on mitochondrial respiration, and are grateful for the opportunity to use the mass spectrometry equipment of the "Human Proteome" Core Facility (Institute of Biomedical Chemistry, Moscow).

Author contributions

OIO, IVR, EST, and DVP conception or design of the work. AAB, RYZ, IIP, TFV, VGZ, MAO, NEV, EML, KAS, ARB, PAM, BSS, IVR, EST, and DVP acquisition, analysis or interpretation of data for the work. AAB, EST, and DVP drafting the work or revising it critically for important intellectual content. All authors approved the final version of the manuscript and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All persons qualify for authorship are listed.

References

Abadi A, Glover EI, Isfort RJ, Raha S, Safdar A, Yasuda N, Kaczor JJ, Melov S, Hubbard A, Qu X et al. 2009. Limb immobilization induces a coordinate down-regulation of mitochondrial and other metabolic pathways in men and women. *PLoSOne* **4**: e6518.

Abu Shelbayeh O, Arroum T, Morris S, Busch KB. 2023. PGC-1alpha Is a Master Regulator of Mitochondrial Lifecycle and ROS Stress Response. *Antioxidants (Basel)* **12**.

Ade CJ, Broxterman RM, Barstow TJ. 2015. VO(2max) and Microgravity Exposure: Convective versus Diffusive O(2) Transport. *Med Sci Sports Exerc* **47**: 1351-1361.

Amirova L, Avdeeva M, Shishkin N, Gudkova A, Guekht A, Tomilovskaya E. 2022. Effect of modulated electromyostimulation on the motor system of elderly neurological patients. Pilot study of Russian currents also known as Kotz currents. *Front Physiol* **13**: 921434.

Barss TS, Ainsley EN, Claveria-Gonzalez FC, Luu MJ, Miller DJ, Wiest MJ, Collins DF. 2018. Utilizing Physiological Principles of Motor Unit Recruitment to Reduce Fatigability of Electrically-Evoked Contractions: A Narrative Review. *Arch Phys Med Rehabil* **99**: 779-791.

Bouguetoch A, Martin A, Grospretre S. 2021. Insights into the combination of neuromuscular electrical stimulation and motor imagery in a training-based approach. *Eur J Appl Physiol* **121**: 941-955.

Casuso RA, Huertas JR, Aragon-Vela J. 2024. The role of muscle disuse in muscular and cardiovascular fitness: A systematic review and meta-regression. *Eur J Sport Sci* **24**: 812-823.

Chopard A, Lecunff M, Danger R, Lamirault G, Bihouee A, Teusan R, Jasmin BJ, Marini JF, Leger JJ. 2009. Large-scale mRNA analysis of female skeletal muscles during 60 days of bed rest with and without exercise or dietary protein supplementation as countermeasures. *Physiol Genomics* **38**: 291-302.

Ciciliot S, Rossi AC, Dyar KA, Blaauw B, Schiaffino S. 2013. Muscle type and fiber type specificity in muscle wasting. *Int J Biochem Cell Biol* **45**: 2191-2199.

Costamagna D, Costelli P, Sampaolesi M, Penna F. 2015. Role of Inflammation in Muscle Homeostasis and Myogenesis. *Mediators Inflamm* **2015**: 805172.

Cresswell AG, Loscher WN, Thorstensson A. 1995. Influence of gastrocnemius muscle length on triceps surae torque development and electromyographic activity in man. *Exp Brain Res* **105**: 283-290.

da Silveira WA, Fazelinia H, Rosenthal SB, Laiakis EC, Kim MS, Meydan C, Kidane Y, Rathi KS, Smith SM, Stear B et al. 2020. Comprehensive Multi-omics Analysis Reveals Mitochondrial Stress as a Central Biological Hub for Spaceflight Impact. *Cell* **183**: 1185-1201 e1120.

Deshmukh AS, Murgia M, Nagaraj N, Treebak JT, Cox J, Mann M. 2015. Deep proteomics of mouse skeletal muscle enables quantitation of protein isoforms, metabolic pathways, and transcription factors. *MolCell Proteomics* **14**: 841-853.

Dirks ML, Miotto PM, Goossens GH, Senden JM, Petrick HL, van Kranenburg J, van Loon LJC, Holloway GP. 2020. Short-term bed rest-induced insulin resistance cannot be explained by increased mitochondrial H2 O2 emission. *J Physiol* **598**: 123-137.

Dirks ML, Wall BT, van de Valk B, Holloway TM, Holloway GP, Chabowski A, Goossens GH, van Loon LJ. 2016. One Week of Bed Rest Leads to Substantial Muscle Atrophy and Induces Whole-Body Insulin Resistance in the Absence of Skeletal Muscle Lipid Accumulation. *Diabetes* **65**: 2862-2875.

Edgerton VR, Zhou MY, Ohira Y, Klitgaard H, Jiang B, Bell G, Harris B, Saltin B, Gollnick PD, Roy RR et al. 1995. Human fiber size and enzymatic properties after 5 and 11 days of spaceflight. *J Appl Physiol (1985)* **78**: 1733-1739.

Edwards SJ, Smeuninx B, McKendry J, Nishimura Y, Luo D, Marshall RN, Perkins M, Ramsay J, Joanisse S, Philp A et al. 2020. High-dose leucine supplementation does not prevent muscle atrophy or strength loss over 7 days of immobilization in healthy young males. *Am J Clin Nutr* **112**: 1368-1381.

Eggelbusch M, Charlton BT, Bosutti A, Ganse B, Giakoumaki I, Grootemaat AE, Hendrickse PW, Jaspers Y, Kemp S, Kerkhoff TJ et al. 2024. The impact of bed rest on human skeletal muscle metabolism. *Cell Rep Med* **5**: 101372.

Fitts RH, Trappe SW, Costill DL, Gallagher PM, Creer AC, Colloton PA, Peters JR, Romatowski JG, Bain JL, Riley DA. 2010. Prolonged space flight-induced alterations in the structure and function of human skeletal muscle fibres. *J Physiol* **588**: 3567-3592.

Fornasiero EF, Mandad S, Wildhagen H, Alevra M, Rammner B, Keihani S, Opazo F, Urban I, Ischebeck T, Sakib MS et al. 2018. Precisely measured protein lifetimes in the mouse brain reveal differences across tissues and subcellular fractions. *Nat Commun* **9**: 4230.

Gibson JN, Morrison WL, Scrimgeour CM, Smith K, Stoward PJ, Rennie MJ. 1989. Effects of therapeutic percutaneous electrical stimulation of atrophic human quadriceps on muscle composition, protein synthesis and contractile properties. *Eur J Clin Invest* **19**: 206-212.

Gibson JN, Smith K, Rennie MJ. 1988. Prevention of disuse muscle atrophy by means of electrical stimulation: maintenance of protein synthesis. *Lancet* **2**: 767-770.

Gram M, Dahl R, Dela F. 2014. Physical inactivity and muscle oxidative capacity in humans. *Eur J Sport Sci* **14**: 376-383.

Hackney KJ, Ploutz-Snyder LL. 2012. Unilateral lower limb suspension: integrative physiological knowledge from the past 20 years (1991-2011). *Eur J Appl Physiol* **112**: 9-22.

Harris S, LeMaitre JP, Mackenzie G, Fox KA, Denvir MA. 2003. A randomised study of home-based electrical stimulation of the legs and conventional bicycle exercise training for patients with chronic heart failure. *Eur Heart J* **24**: 871-878.

Karavidas A, Arapi SM, Pyrgakis V, Adamopoulos S. 2010. Functional electrical stimulation of lower limbs in patients with chronic heart failure. *Heart Fail Rev* **15**: 563-579.

Koryak YA. 2020. Isokinetic Force and Work Capacity After Long-Duration Space Station Mir and Short-Term International Space Station Missions. *Aerosp Med Hum Perform* **91**: 422-431.

Kozlovskaya IB, Sayenko IV, Sayenko DG, Miller TF, Khusnutdinova DR, Melnik KA. 2007. Role of support afferentation in control of the tonic muscle activity. *Acta Astronautica* **60**: 285-294.

Kurochkina NS, Orlova MA, Vigovskiy MA, Zgoda VG, Vepkhvadze TF, Vavilov NE, Makhnovskii PA, Grigorieva OA, Boroday YR, Philippov VV et al. 2024. Age-related changes in human skeletal muscle transcriptome and proteome are more affected by chronic inflammation and physical inactivity than primary aging. *Aging Cell* doi:10.1111/acel.14098: e14098.

LeBlanc AD, Schneider VS, Evans HJ, Pientok C, Rowe R, Spector E. 1992. Regional changes in muscle mass following 17 weeks of bed rest. *J Appl Physiol (1985)* **73**: 2172-2178.

Maffiuletti NA. 2010. Physiological and methodological considerations for the use of neuromuscular electrical stimulation. *Eur J Appl Physiol* **110**: 223-234.

Maffiuletti NA, Green DA, Vaz MA, Dirks ML. 2019. Neuromuscular Electrical Stimulation as a Potential Countermeasure for Skeletal Muscle Atrophy and Weakness During Human Spaceflight. *Front Physiol* **10**: 1031.

Makhnovskii PA, Bokov RO, Kolpakov FA, Popov DV. 2021. Transcriptomic Signatures and Upstream Regulation in Human Skeletal Muscle Adapted to Disuse and Aerobic Exercise. *Int J Mol Sci* **22**.

Marshall RN, Smeuninx B, Seabright AP, Morgan PT, Atherton PJ, Philp A, Breen L. 2022. No effect of five days of bed rest or short-term resistance exercise prehabilitation on markers of skeletal muscle mitochondrial content and dynamics in older adults. *Physiol Rep* **10**: e15345.

Marusic U, Narici M, Simunic B, Pisot R, Ritzmann R. 2021. Nonuniform loss of muscle strength and atrophy during bed rest: a systematic review. *J Appl Physiol (1985)* **131**: 194-206.

Mayr W, Bijak M, Girsch W, Hofer C, Lanmuller H, Rafolt D, Rakos M, Sauermann S, Schmutterer C, Schnetz G et al. 1999. MYOSTIM-FES to prevent muscle atrophy in microgravity and bed rest: preliminary report. *Artif Organs* **23**: 428-431.

Mikines KJ, Richter EA, Dela F, Galbo H. 1991. Seven days of bed rest decrease insulin action on glucose uptake in leg and whole body. *J Appl Physiol (1985)* **70**: 1245-1254.

Miotto PM, McGlory C, Bahniwal R, Kamal M, Phillips SM, Holloway GP. 2019. Supplementation with dietary omega-3 mitigates immobilization-induced reductions in skeletal muscle mitochondrial respiration in young women. *FASEB J* **33**: 8232-8240.

Murgia M, Ciciliot S, Nagaraj N, Reggiani C, Schiaffino S, Franchi MV, Pisot R, Simunic B, Toniolo L, Blaauw B et al. 2022. Signatures of muscle disuse in spaceflight and bed rest revealed by single muscle fiber proteomics. *PNAS Nexus* **1**: pgac086.

Murgia M, Nogara L, Baraldo M, Reggiani C, Mann M, Schiaffino S. 2021. Protein profile of fiber types in human skeletal muscle: a single-fiber proteomics study. *Skelet Muscle* **11**: 24.

Murgia M, Toniolo L, Nagaraj N, Ciciliot S, Vindigni V, Schiaffino S, Reggiani C, Mann M. 2017. Single Muscle Fiber Proteomics Reveals Fiber-Type-Specific Features of Human Muscle Aging. *Cell Rep* **19**: 2396-2409.

Narici MV, de Boer MD. 2011. Disuse of the musculo-skeletal system in space and on earth. *Eur J Appl Physiol* **111**: 403-420.

Niess F, Fiedler GB, Schmid AI, Laistler E, Frass-Kriegl R, Wolzt M, Moser E, Meyerspeer M. 2018. Dynamic multivoxel-localized (31) P MRS during plantar flexion exercise with variable knee angle. *NMR Biomed* **31**: e3905.

Nuhr M, Crevenna R, Gohlsch B, Bittner C, Pleiner J, Wiesinger G, Fialka-Moser V, Quittan M, Pette D. 2003. Functional and biochemical properties of chronically stimulated human skeletal muscle. *Eur J Appl Physiol* **89**: 202-208.

Nuhr MJ, Pette D, Berger R, Quittan M, Crevenna R, Huelsman M, Wiesinger GF, Moser P, Fialka-Moser V, Pacher R. 2004. Beneficial effects of chronic low-frequency stimulation of thigh muscles in patients with advanced chronic heart failure. *Eur Heart J* **25**: 136-143.

Poltavskaya MG, Sviridenko VP, Brand AV, Andreev DE, Koryak YA, Veliev GO, Dikur ON, Kulikov VM, Vaisman YD, Tomilovskaya ES. 2021. The Use of "Space" Electrical Myostimulation in Clinical Cardiology on Earth. *Hum Physiology* **47**: 382-390.

Popov DV, Makhnovskii PA, Shagimardanova EI, Gazizova GR, Lysenko EA, Gusev OA, Vinogradova OL. 2019. Contractile activity-specific transcriptome response to acute endurance exercise and training in human skeletal muscle. *Am J Physiol Endocrinol Metab* **316**: E605-E614.

Popov DV, Makhnovskii PA, Zgoda VG, Gazizova GR, Vepkhvadze TF, Lednev EM, Motanova ES, Lysenko EA, Orlov OI, Tomilovskaya ES. 2023. Rapid changes in transcriptomic profile and mitochondrial function in human soleus muscle after 3-day dry immersion. *J Appl Physiol (1985)* **134**: 1256-1264.

Quittan M, Wiesinger GF, Sturm B, Puig S, Mayr W, Sochor A, Paternostro T, Resch KL, Pacher R, Fialka-Moser V. 2001. Improvement of thigh muscles by neuromuscular electrical stimulation in patients with refractory heart failure: a single-blind, randomized, controlled trial. *Am J Phys Med Rehabil* **80**: 206-214; quiz 215-206, 224.

Ried-Larsen M, Aarts HM, Joyner MJ. 2017. Effects of strict prolonged bed rest on cardiorespiratory fitness: systematic review and meta-analysis. *J Appl Physiol (1985)* **123**: 790-799.

Ringholm S, Bienso RS, Kiilerich K, Guadalupe-Grau A, Aachmann-Andersen NJ, Saltin B, Plomgaard P, Lundby C, Wojtaszewski JF, Calbet JA et al. 2011. Bed rest reduces metabolic protein content and abolishes exercise-induced mRNA responses in human skeletal muscle. *Am J Physiol Endocrinol Metab* **301**: E649-658.

Rolfs Z, Frey BL, Shi X, Kawai Y, Smith LM, Welham NV. 2021. An atlas of protein turnover rates in mouse tissues. *Nat Commun* **12**: 6778.

Salvadego D, Keramidas ME, Brocca L, Domenis R, Mavelli I, Rittweger J, Eiken O, Mekjavic IB, Grassi B. 2016. Separate and combined effects of a 10-d exposure to hypoxia and inactivity on oxidative function in vivo and mitochondrial respiration ex vivo in humans. *J Appl Physiol (1985)* **121**: 154-163.

Salvadego D, Keramidas ME, Kolegard R, Brocca L, Lazzer S, Mavelli I, Rittweger J, Eiken O, Mekjavic IB, Grassi B. 2018. PlanHab(*) : hypoxia does not worsen the impairment of skeletal muscle oxidative function induced by bed rest alone. *J Physiol* **596**: 3341-3355.

Salvadego D, Lazzer S, Marzorati M, Porcelli S, Rejc E, Simunic B, Pisot R, di Prampero PE, Grassi B. 2011. Functional impairment of skeletal muscle oxidative metabolism during knee extension exercise after bed rest. *J Appl Physiol (1985)* **111**: 1719-1726.

Sbruzzi G, Ribeiro RA, Schaan BD, Signori LU, Silva AM, Irigoyen MC, Plentz RD. 2010. Functional electrical stimulation in the treatment of patients with chronic heart failure: a meta-analysis of randomized controlled trials. *Eur J Cardiovasc Prev Rehabil* **17**: 254-260.

Schwanhausser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, Chen W, Selbach M. 2011. Global quantification of mammalian gene expression control. *Nature* **473**: 337-342.

Sharlo K, Paramonova I, Turtikova O, Tyganov S, Shenkman B. 2019. Plantar mechanical stimulation prevents calcineurin-NFATc1 inactivation and slow-to-fast fiber type shift in rat soleus muscle under hindlimb unloading. *J Appl Physiol (1985)* **126**: 1769-1781.

Shenkman BS, Lyubaeva EV, Popov DV, Netreba AI, Bravy YR, Tarakin PP, Lemesheva YS, Vinogradova OL. 2007. Chronic effects of low-frequency low-intensity electrical stimulation of stretched human muscle. *Acta Astronautica* **60**: 505-511.

Shenkman BS, Lyubaeva EV, Popov DV, Netreba AI, Tarasova OS, Vdovina AB, Tarakin PP, Lemesheva YS, Belichenko OI, Sinitsyn VE et al. 2006. Effects of chronic low-frequency electrical stimulation of human knee extensor muscles exposed to static passive stretching. *Human Physiology* **32**: 74-80.

Smart NA, Dieberg G, Giallauria F. 2013. Functional electrical stimulation for chronic heart failure: a meta-analysis. *Int J Cardiol* **167**: 80-86.

Standley RA, Distefano G, Trevino MB, Chen E, Narain NR, Greenwood B, Kondakci G, Tolstikov VV, Kiebish MA, Yu G et al. 2020. Skeletal Muscle Energetics and Mitochondrial Function Are Impaired Following 10 Days of Bed Rest in Older Adults. *J GerontolA BiolSciMedSci* **75**: 1744-1753.

Stefanovska A, Vodovnik L. 1985. Change in muscle force following electrical stimulation. Dependence on stimulation waveform and frequency. *Scand J Rehabil Med* **17**: 141-146.

Thoma A, Lightfoot AP. 2018. NF-kB and Inflammatory Cytokine Signalling: Role in Skeletal Muscle Atrophy. *Adv Exp Med Biol* **1088**: 267-279.

Tomilovskaya E, Shigueva T, Sayenko D, Rukavishnikov I, Kozlovskaya I. 2019. Dry Immersion as a Ground-Based Model of Microgravity Physiological Effects. *Front Physiol* **10**: 284.

van Wessel T, de Haan A, van der Laarse WJ, Jaspers RT. 2010. The muscle fiber type-fiber size paradox: hypertrophy or oxidative metabolism? *EurJApplPhysiol* **110**: 665-694.

Vikne H, Strom V, Pripp AH, Gjovaag T. 2020. Human skeletal muscle fiber type percentage and area after reduced muscle use: A systematic review and meta-analysis. *Scand J Med Sci Sports* **30**: 1298-1317.

Yarmanova EN, Kozlovskaya IB, Khimoroda NN, Fomina EV, Reeves JM. 2015. Evolution of russian microgravity countermeasures. *Aerospace Medicine and Human Performance* **86**: A32-A37.

Zanotti E, Felicetti G, Maini M, Fracchia C. 2003. Peripheral muscle strength training in bed-bound patients with COPD receiving mechanical ventilation: effect of electrical stimulation. *Chest* **124**: 292-296.

Zuccarelli L, Baldassarre G, Magnesa B, Degano C, Comelli M, Gasparini M, Manferdelli G, Marzorati M, Mavelli I, Pilotto A et al. 2021. Peripheral impairments of oxidative metabolism after a 10-day bed rest are upstream of mitochondrial respiration. *J Physiol* **599**: 4813-4829.

Supporting Information

Supporting Information Fig. S1. Subject position and dynamics of external load, motion amplitude, cycle duration, and calculated power during an incremental ramp test for the ankle plantar flexors.

Supporting Information Fig. S2. Subject position and dynamics of external load, knee angle amplitude, cycle duration, and calculated power during an incremental ramp test for the knee extensors.

Supporting Information Fig. S3. Expression of mRNAs encoding cyto/myokines and their receptors (A) and proteasomal proteins (B) in the "mixed" vastus lateralis (VL) and "slow" soleus (Sol) muscles after dry immersion without (DI) and with neuromuscular electrical stimulation (DI+NMES); ($n = 10$ subjects in each group). The color of the circles and the heat map show the magnitude of changes in various muscle after various intervention for each mRNA; *N.S.* – non-significant change.

Supporting Information Fig. S4. Number of differentially expressed genes (DEGs) related to some enriched functional categories from Fig. 4 and their changes in expression after 6 days of dry immersion without (DI) and with neuromuscular electrical stimulation (DI+NMES); (n = 10 subjects in each group). Common for both and unique to each intervention DEGs are shown; *p*-values relate to the difference between DI and DI+NMES (Mann-Whitney test).

Supporting Information Table S1. All detected and differentially expressed proteins (DEP) and mRNAs (DEG) in the vastus lateralis muscle (VL) *vs.* soleus muscle (S) in the baseline.

Supporting Information Table S2. Function enrichment analysis of differentially expressed proteins and mRNAs in the vastus lateralis muscle (VL) *vs.* soleus muscle (S) in the baseline. The main terms shown in Fig. 3 are depicted in red.

Supporting Information Table S3. All and differentially expressed mRNAs (DEG) in the vastus lateralis (VL) and soleus (S) muscles after 6 days of dry immersion.

Supporting Information Table S4. Function enrichment analysis of differentially expressed proteins and mRNAs in the vastus lateralis (VL) and soleus (S) muscles after 6 days of dry immersion.