Stable and dynamic gene expression patterns over diurnal and developmental timescales in *Arabidopsis thaliana*

Running title: The Circadian Clock in the Ageing Plant

Ethan J Redmond[1] (https://orcid.org/0000-0001-9105-0609), James Ronald[1] (https://orcid.org/0000-0002-8847-0378), Seth J Davis[1] (https://orcid.org/0000-0001-5928-9046), Daphne Ezer[1,*] (https://orcid.org/0000-0002-1685-6909)

[1]: Department of Biology, University of York, Wentworth Way, Heslington, York YO10 5DD, UK

[*]: Corresponding author. daphne.ezer@york.ac.uk

Summary

Developmental processes throughout the lifecycle of a plant are known to be circadian regulated. For instance, the circadian clock regulates key genes involved in the photoperiodic flowering pathway and the initiation of leaf senescence. Furthermore, signals which entrain the circadian clock, such as energy availability, are known to vary in strength over plant development. However, diurnal oscillations of the *Arabidopsis* transcriptome have typically been measured in seedlings. We collected RNA-seq data from *Arabidopsis* leaves over developmental and diurnal timescales, concurrently: every 4 hours per day, on 3 separate days after a synchronised vegetative-to-reproductive transition. Gene expression varied more over the developmental timescale than on the diurnal timescale, including genes related to a key energy sensor: the *Sucrose non-fermenting-1-related protein kinase* (SnRK1) complex. Moreover, regulatory targets of core clock genes displayed changes in rhythmicity and amplitude of expression over development. Cell-type-specific expression showed diurnal patterns that varied in amplitude, but not phase, over development. Some previously identified qRT-PCR housekeeping genes display undesirable levels of variation over both timescales. We identify which common qRT-PCR housekeeping genes are most stable across developmental and diurnal timescales. In summary, we establish the patterns of circadian transcriptional regulation over plant development, demonstrating how diurnal patterns of expression change over developmental timescales.

Introduction

Due to their sessile nature within a cyclical environment, plants have evolved an internal timekeeper called the circadian clock. It consists of a core interlocking loop of transcriptional regulators, whose components are mostly conserved across embryophytes (Petersen *et al.*,

2022; Wang *et al.*, 2024). Circadian clocks confer fitness to plants by allowing them to
coordinate responses to photoperiod, light quality, temperature, and other environmental cues
(Dodd *et al.*, 2005; Atamian *et al.*, 2016; Rubin *et al.*, 2017; Xu *et al.*, 2022). This coordination
takes place through widespread transcriptional control of the plant transcriptome, among other
methods of regulation (Nagel *et al.*, 2015; Hayama *et al.*, 2017; Ezer *et al.*, 2017; Romanowski
et al., 2020; Xiong *et al.*, 2022). Estimates of the proportion of the *Arabidopsis* and wheat
transcriptomes that are controlled by the clock range from 30% to 50% (Covington *et al.*, 2008;
Romanowski *et al.*, 2020; Rees *et al.*, 2022).

Large-scale circadian RNA-seq experiments are typically performed in seedlings or juvenile
plants, due to the ease of performing timeseries experiments under different entrainment and
free-running conditions. Yet, many of the clock genes have fundamental roles in the timing of
developmental processes (Inoue *et al.*, 2018; Wang *et al.*, 2024). A key example of this is the
evening complex (EC), a tripartite protein complex containing EARLY FLOWERING 3 and 4
(ELF3 and ELF4) and LUX ARRHYTHMO (LUX) (Nusinow *et al.*, 2011; Herrero *et al.*, 2012).
The EC, in combination with GIGANTEA (GI), acts upstream of the photoperiodic flowering
pathway in *Arabidopsis* and rice (Fowler *et al.*, 1999; Park *et al.*, 1999; Sawa & Kay, 2011;
Andrade *et al.*, 2022). Leaf senescence occurs concurrently with the vegetative-to-
reproductive transition (Redmond et al., 2024) and is also regulated by the circadian clock.
Mutations in clock genes, including *ELF3*, *PSEUDO-RESPONSE REGULATOR9* (*PRR9*) and
CIRCADIAN-CLOCK ASSOCIATED1 (*CCA1*) have all been shown to affect the onset of leaf
senescence (Sakuraba *et al.*, 2014; Song *et al.*, 2018; Kim *et al.*, 2018).

Moreover, developmentally associated processes exhibit diurnal patterns. These patterns
begin at the earliest stages of plant development and extend into all developmental stages.
Germination responds to diurnally fluctuating temperatures in many plant species (Thompson
et al., 1977). Hypocotyl development is mediated through auxin- and temperature-related
processes, via the clock-controlled *PHYTOCHROME-INTERACTING FACTORS 4 and 5*
(PIF4 and *PIF5)* (Nozue *et al.*, 2007; Seaton *et al.*, 2015). Many of the central integrators that
control flowering time are diurnally expressed, such as *FLOWERING LOCUS T* (*FT*),
SUPRESSOR OF OVEREXPRESSION OF CO1 (*SOC1*) and *LEAFY* (*LFY*) (Wendell *et al.*,
2017). Additionally, many of the transcription factors that regulate the synthesis of key plant
hormones involved in development, like auxin and ABA, are diurnally expressed (Balcerowicz
et al., 2021).

Tissue-specificity also plays a role in the link between the clock and development. For
instance, the vascular clock plays a dominant role over the epidermal clock in leaves.
Moreover, these tissue-specific clocks influence two distinct developmental processes,

flowering and hypocotyl development (Endo *et al.*, 2014). Vong *et al.,* 2024 suggested that

each cell type's transcriptional activity varied across diurnal and developmental timescales,

but it is unclear how the daily oscillations in cell-type activity vary over developmental

timescales.

There remains a large gap in our knowledge of how diurnal genes vary over development and

how developmental genes vary across the day. Here, we address this gap by measuring

diurnal gene expression over the one of the most crucial developmental transitions an annual

plant experiences: the vegetative-to-reproductive transition. We measure gene expression

over two timescales: every 24 hours (the diurnal timescale) and over approximately 2 weeks

(the developmental timescale). As individual plants experience developmental asynchrony in

their floral transition (Klingenberg, 2019; Redmond *et al.*, 2024), we use a photoperiod shift

from short days (SD; 8h light per day) to long days (LD; 16h light per day) to induce

synchronised vegetative-to-reproductive transitions. One day of LD conditions is sufficient to

induce maximum gene expression response of *FLOWERING LOCUS T* (*FT*) (Corbesier *et al.*,

1996; Krzymuski *et al.*, 2015). We therefore measure diurnal gene expression in ageing plants

after the inductive SD to LD signal.

First, we found that gene expression changed most dramatically over the developmental

timescale and that core clock genes had broadly stable phase and amplitude of expression

per day. Second, we observed the expression dynamics of targets of a key *Arabidopsis*

energy sensor vary across both scales. Third, we showed that transcriptional targets of core

clock genes exhibited differing changes of amplitude over development. Fourth, we identified

tissue-specific changes in rhythmic processes in ageing leaves. Finally, one of the most

important applications of our work is in identifying sets of genes that are stable across both

timescales, as these could serve as important controls for qRT-PCR. We suggest a filtered

set of housekeepers that we would encourage the community to use for studies which aim to

study gene expression of clock-controlled processes in ageing plants.

Materials and Methods

Plant growth conditions

Seeds from the Arabidopsis ecotype Wassilewskija (Ws-2) (Anwer *et al.*, 2020) were surface-

sterilised. They were then plated onto 1x Murashige and Skoog basal salts (Duchefa

Biochemie) supplemented with 1% w/v sucrose (ThermoFisher), 0.5% w/v MES (Melford

Bioscience) and 1.5% w/v phytoagar (Duchefa Biochemie). After four days of stratification at

4°C, these plates were then transferred to short day (SD) conditions (8h light / 16h dark; 70

µmol m^{-2} s^{-1} with vertical fluorescent lighting; Osram L 36W/830 Warm White) at 21°C and left

for 10 days. Then, 2 seedlings per pot were transferred to soil, then were thinned out to 1
seedling per pot after 3 days. 3rd and 4th true rosette leaves were tracked for each individual
plant, by placing pipette tips in soil next to these leaves and adjusting them as the leaves
moved. 12 days after transferring to soil, the conditions were changed to long day (LD; 16h
light / 8h dark) at 21°C.

Samples were collected at ZT (zeitgeiber time; hours after dawn) 0, 1, 4, 8, 12, 16, 20 on three
days. We chose to include the ZT1 timepoint due to our previous observations that gene
expression changes rapidly in the first hour after dawn (Balcerowicz *et al.*, 2021). The first day
of sample collection started 24 hours after ZT0 of the first LD (2nd day post-transition). The
second day started 5 days later (7th day post-transition). The third started another 5 days later
(12th day post-transition). 3 replicates for each time point were taken - with each replicate
consisting of the 3rd and 4th true rosette leaves of two separate plants. These were flash
frozen in liquid nitrogen and then stored at -80°C. For collections during dark conditions (i.e.
ZT16 and ZT20), this work was carried out under green-filtered light.

RNA-seq and initial data processing

RNA was extracted using the Monarch Total RNA Miniprep Kit (New England Biolabs),
according to the manufacturer's protocol. Residual genomic DNA treatment was removed
using the Invitrogen TURBO DNA-free kit (ThermoFisher), using 'Routine DNase treatment',
according to the manufacturer's protocol.

Libraries were prepared from high quality RNA by the Genomics Laboratory of the University
of York's Bioscience Technology Facility. Libraries were prepared using the NEBNext Ultra II
Directional Library prep kit for Illumina in conjunction with the NEBNext poly(A) magnetic
isolation module (New England Biolabs). Library quality was assessed using the Agilent 2100
Bioanalyzer instrument, before pooling at equimolar ratios. Pooled libraries were sent for
paired end 150 base sequencing by Novogene (UK) Ltd, on an Illumina NovaSeq 6000.

Before further processing of the raw sequencing data, FastQC v0.11.7 (Andrews, 2010) was
used to ensure sample quality passed the expected checks for RNA-seq data. Illumina
adapters and low quality bases at the 3' and 5' ends were trimmed using CutAdapt v3.4
(Martin, 2011). Reads were quantified using Salmon v1.6.0 (Patro *et al.*, 2017) against the
TAIR10 transcriptome (Berardini *et al.*, 2015).

R (version 4.2.3) was used for analysis of gene expression data (R Core Team, 2023).
Transcripts per million (TPM) was used as the measure of relative gene expression across
samples. Additionally, TPM levels per transcript isoform were combined to leave only gene-
level expression data. 13,156 genes were filtered out before further analysis if all median TPM

values (across the three biological replicates per time-of-day and sampling day) were less
than or equal to 0.5 (Supplemental Figure 1, Supplemental Table S1). Where log-transformed
gene expression is referred to in this text, the transformed expression is $(TPM + 1)$.

PCA and GO term overrepresentation

PCA was performed using 'prcomp' on log-transformed gene expression on all samples. All
GO term overrepresentation was performed using the gprofiler2 R package (version 0.2.2)
(Kolberg *et al.*, 2020). This called the g:Profiler server (version e111_eg58_p18_f463989d),
which utilises the g:SCS multiple testing correction method, and we then applied a significance
threshold of 0.05 (Kolberg *et al.*, 2023) (Supplemental Tables S2, S3, S6). Related groups of
genes were submitted as 'multi queries'.

CIBERSORTx

148　TPM tables were input into the Cell Fraction mode of the CIBERSORTx web interface
149　(https://cibersortx.stanford.edu/, 28 May 2024), using 100 permutations for the permutation
150　test to assess the p-values (Newman *et al.*, 2019). The signature matrix came from (Vong *et*
151　*al.*, 2024) and was trained on the single cell RNA-seq data from (Procko *et al.*, 2022).

Rhythmic gene prediction

To predict rhythmic genes, we applied the R package MetaCycle (version 1.2.0) to log-
transformed gene expression, only allowing the JTK_cycle method (Hughes *et al.*, 2010; Wu
et al., 2016). Samples at ZT1 were excluded since JTK_cycle required evenly spaced time
points. Genes were classified as rhythmic if and only if the 'BH.Q' value was less than 0.05.
The same method was used to classify the rhythmicity of cell types, where instead we used
unscaled cell proportions as inputs.

Data from previous publications

To identify the putative targets of SNF1 KINASE HOMOLOG10 (KIN10), we took the 'AGI
number' column from each of the 'INCREASED' and 'DECREASED' sections from
Supplementary Table S3 of ref. (Baena-González *et al.*, 2007). Putative targets of CIRCADIAN
CLOCK ASSOCIATED1 (CCA1) were collected from Dataset S2 of ref. (Nagel *et al.*, 2015).
Putative targets of EARLY FLOWERING3 (ELF3) were collected from Supplementary Table
6 of ref. (Ezer *et al.*, 2017). Housekeeping genes suggested for developmental and diurnal
time series were collected from Supplemental Table 1 of ref. (Czechowski *et al.*, 2005). Only
gene names which consistent with our processed RNA-seq were used.

Results

<u>Gene expression varies more across development than across a diurnal cycle.</u>

Since the clock is tightly linked to changes in gene expression over plant development, we
first sought to determine whether diurnal or developmental time had a greater impact on gene
expression. Principal component analysis (PCA) shows that the first principal component
primarily separates out samples based on developmental time, suggesting that variation in
gene expression across development dominates variation from diurnal oscillations (Figure
1(a)). As expected, the central day (7th day after transition) lies in between the early (2nd) and
late (12th) days. This agrees with clustering based on correlation between samples
(Supplemental Figure 2). 52.5% of variance within the filtered dataset can be explained by
the developmental-related principal component – indicating a large transcriptomic scale shift
in the 12 days after the transition (Figure 1(a), Supplemental Figure 3).

In the next two principal components (PC2 and PC3), samples clustered based on time-of-
day, not day of sampling, with samples forming a circle ordered by the time of day that samples
were collected (Figure 1(b)). These results suggest that PCA successfully isolates the
developmental and diurnal components of gene expression in our samples.

By analysing the relative contribution (i.e. loadings) of each gene to the first three principal
components, we can identify genes that may be associated with developmental (PC1) or
diurnal (PC2/3) processes (Figure 1(c), 1(d)). For instance, we can observe that the primary
circadian clock genes, as defined by (Wang *et al.*, 2024), are correctly ordered in the loadings
in Figure 1(d). This includes a precise counter-clockwise ordering of the PSEUDO-
RESPONSE REGULATOR (PRR) genes by the order of their expected phases (Webb *et al.*,
2019). The gene with the most positive loading on PC1 was PATHOGENESIS-RELATED
GENE 1 (PR1; AT2G14610), a salicylic acid responsive gene whose expression is tightly
linked to leaf senescence (Zhang *et al.*, 2013). The gene with the most negative loading on
PC1 was a proline-rich, cell wall protein AT2G10940 (Duruflé *et al.*, 2017). As expected, the
expression of PR1 increases post-transition and the expression of AT2G10940 decrease post-
transition (Figure 1(e)). We identified the 'core clock' genes, related to the vegetative-to-
reproductive transition as defined in (Wang *et al.*, 2024), with the most positive and negative
loadings on PC2. The clock gene with most positive PC2 loading was EARLY FLOWERING4
(ELF4; AT2G40080), a member of the evening complex (Nusinow *et al.*, 2011; Herrero *et al.*,
2012; Ezer *et al.*, 2017), which peaked around ZT8 in all days. The clock gene with most
negative PC2 loading was CIRCADIAN CLOCK ASSOCIATED1 (CCA1; AT2G46830), which
peaked in the early morning in all days. Interestingly, the transcript level of *CCA1* did not
appear to decrease across developmental time (i.e. as the leaves aged), contrary to previous

reports using qRT-PCR (Song *et al.*, 2018). These results suggest that we can use the PCA
loadings to identify genes that are primarily developmental or diurnally regulated.

<u>Developmental changes in gene expression are linked to metabolism, senescence, and</u>
<u>kinase-related processes.</u>

To determine the broad shifts in gene expression that occur across the developmental
timescale, we created two groups of the most early- and late-associated genes. These
consisted of genes with the most negative 10% of PC1 loadings (associated with high
expression on the 2nd day of sampling) and the most positive 10% (associated with high
expression on the 12th day of sampling). GO term overrepresented terms among early-
associated genes include 'photosynthesis' (GO:0015979) and 'generation of precursor
metabolites and energy' (GO:0006091), while those overrepresented among late-associated
genes include 'defence response' (GO:0006952) and 'plant organ senescence'
(GO:0090693). These are consistent with the vegetative-to-reproductive transition associated
GO terms identified in (Hinckley & Brusslan, 2020; Redmond *et al.*, 2024) (Supplemental Table
S2).

Kinases have been associated with regulating the floral transition, as well as diurnally
phosphorylating proteins, forming a bridge between developmental and diurnal processes
(Tsai & Gazzarrini, 2012; Uhrig *et al.*, 2021). Since 'protein serine/threonine kinase activity'
(GO:0004674) was overrepresented in the top 10% of genes by positive PC1 loading, we
visualised the distribution of all genes related to this GO term (Supplemental Table S2,
Supplemental Figure 4(a)). The PC1 (developmental time associated) loadings of these genes
were significantly different to 0 (two-sided t-test, t = 17.778, p-value < 2.2e-16) and the mean
value was 4.64e-4 (Supplemental Figure 4(a)). However, there was a much smaller deviation
from zero for the two diurnal time associated PCs (PC2 and PC3). The loadings on PC2 were
not significantly different from 0 (two-sided t-test, t = -1.8644, p-value = 0.06272; mean value
-4.47e-4) and the loadings for PC3 were significantly different (two-sided t-test, t = 7.4955, p-
value = 2.13e-13; mean value 1.83e-3), but with a much lower effect size than PC1. This
suggests that expression of kinases primarily varies over the developmental timescale.

The 4 genes from this category with the highest PC1 loading came from two kinase families.
Two of these were *WALL-ASSOCIATED KINASES* (*WAK1*, *WAK3*), which act as pectin
receptors and have roles in defence response and cell expansion (Anderson *et al.*, 2001;
Kohorn & Kohorn, 2012). A further two were *CYSTEINE-RICH KINASES* (*CRK4*, *CRK45*).
Overexpression of *CRK4* is known to enhance pattern-triggered immunity (Yeh *et al.*, 2015).
Each of these genes had the property that the expression on day 2 is consistently low
throughout the day and the expression of day 12 is consistently high. However, the expression

on day 7 spans the range of expression on day 2 and day 12, matching day 2 expression at
ZT0 and day 12 expression at ZT20. (Supplemental Figure 4(b)). This could explain the
significantly non-zero PC3 loadings of kinases, since PC3 may be incorporating some
developmental variation as well as diurnal variation.

<u>Targets of *SNF1 KINASE HOMOLOG 10 (KIN10)* Show Diurnal and Developmental</u>
<u>Expression Changes</u>

Next, we chose to analyse the diurnal and developmental variation of a well-known kinase-
containing protein complex that functions as an energy sensor in *Arabidopsis*: the *Sucrose*
non-fermenting-1-related protein kinase (SnRK1) complex (Polge & Thomas, 2007; Broeckx
et al., 2016), which can entrain the clock in response to sugars (Frank *et al.*, 2018) and can
regulate growth and development in response to sugar status (Chen *et al.*, 2017). The SnRK1
complex consists of three subunits, each of which are encoded from different homologues
(Broeckx *et al.*, 2016). *SNF1 KINASE HOMOLOG10* (*KIN10*) and *SNF1 KINASE*
HOMOLOG11 (KIN11), are protein serine/threonine kinases, which encode the α subunit of
SnRK1. Importantly, homologues of all three subunits may be expressed specifically in
different developmental stages and environmental conditions (Broeckx *et al.*, 2016). Both
KIN10 and KIN11 had increased expression over the developmental timescale (Figure 2(a)).
However, especially in the 12th day, KIN10 expression had larger diurnal variation than KIN11.
As for the β subunit, we found that KINβ1 expression had much larger variation over both the
diurnal and developmental timescales than expression of its homologues KINβ2 and KINβ3
(Supplemental Figure 5(a)). Finally, the essential gene KINβγ (KINbeta-gamma, HOMOLOG
OF YEAST SUCROSE NONFERMENTING4, SNF4) was far more stably expressed over
development than the related KINγ (KINgamma) (Supplemental Figure 5(b)) (Ramon *et al.*,
2013). This could be explained by the fact KINγ is not suspected to form a functional part of
the SnRK1 complex, as opposed to its orthologue in mammalian systems (Emanuelle *et al.*,
2015).

Since KIN10 is known to activate and repress a wide range of pathways (Baena-González *et*
al., 2007), we asked how expression of processes that act downstream of KIN10 changed
over diurnal and developmental timescales. We selected genes which increased or decreased
expression by transient *KIN10* overexpression in protoplasts and clustered them based on
expression over both timescales (Figure 2(b), 2(c)) (Baena-González *et al.*, 2007). Both sets
of targets clustered into two large subgroups – characterised by their expression patterns over
the developmental timescale. In terms of genes with increased expression in the *KIN10*
overexpressed lines*,* cluster 1 showed a large increase over age, which agrees with greater
KIN10 expression over age (Figure 2(a), 2(b)). GO terms significantly overrepresented in this
cluster (and not in cluster 2) included 'response to hypoxia' (GO:0001666), 'response to

abscisic acid' (GO:0009737), and 'carbohydrate metabolic process' (GO:0005975)
(Supplemental Table S3). These represented the range of stress responses *KIN10* is known
to affect as well as its control over metabolism (Baena-González *et al.*, 2007). Cluster 2
broadly showed a less dramatic increase in expression over age.

In terms of genes with decreased expression in the *KIN10* overexpressed lines, the two
clusters showed strikingly different expression. Cluster 1 expression broadly decreased in
expression over age, which agrees with increasing *KIN10* expression (Figure 2(a)). The
majority of these targets also peaked at ZT12 or later in the day, when starch is degraded to
provide a carbohydrate source (Figure 2(c)) (Streb & Zeeman, 2012). This cluster appeared
to represent the sink-to-source transition of ageing leaves, since both 'primary metabolic
process' (GO:0044238) and 'cellular nitrogen compound metabolic process' (GO:0034641)
were significantly overrepresented (Supplemental Table S3) (Havé *et al.*, 2017). Contrarily,
cluster 2 showed a broad increase in expression. This suggests that *KIN10* repression in
protoplasts does not always predict repression in ageing plants, at least at the transcriptional
level. Our results suggest that SnRK1 sub-components primarily changed their transcriptional
profiles developmentally, but that most genes that were downregulated in response to KIN10
expression have consistent diurnal expression patterns.

<u>Many targets of core clock genes vary their expression patterns over development</u>

Although PCs 2 and 3 (the diurnal components) accounted for only 24.6% of the variation in
our dataset and the diurnal variation of *KIN10* expression was relatively small, metabolic
signals are known to be both inputs and outputs of the circadian clock (Shin *et al.*, 2017; Frank
et al., 2018; Cervela-Cardona *et al.*, 2021). Additionally, senescence-associated changes in
period and phase have been observed in both the core clock genes and the wider circadian
transcriptome (Kim *et al.*, 2016, 2018; Buckley *et al.*, 2024). We noticed that most clock genes
had expression patterns that were consistent across development, under our fixed long-day
conditions (Supplemental Figure 6). This prompts the questions of whether clock gene *targets*
had stable expression over development and whether they were rhythmic in each day after
the SD to LD transition.

CIRCADIAN-CLOCK ASSOCIATED 1 (*CCA1*) is a core clock gene whose expression peaks
in the early morning under long days (Figure 3(a)) (Webb *et al.*, 2019; Wang *et al.*, 2024). We
collected predicted CCA1 targets from a previous ChIP-seq study, where seedlings were
grown under 12h light/ 12h dark conditions (Nagel *et al.*, 2015). We observed that the median
expression of *CCA1,* at every time of day, increased slightly over development (Figure 3(a)).
Surprisingly, we found that clusters targets of CCA1 demonstrated increasing, decreasing,
and stable mean expression per day of the experiment (Figure 3(b), 3(c), Supplemental Table

S5). Further, we found that some clusters of targets became less rhythmic over age (e.g.
clusters B and E), according to JTK_cycle, while some clusters maintained high levels (>80%)
of rhythmicity (cluster C) (Hughes *et al.*, 2010).

To show if these complicated patterns of regulation were preserved across other clock genes,
we looked at predicted targets of another clock gene, *ELF3*, from (Ezer *et al.*, 2017). *ELF3* is
a component of the EC and its expression peaks around ZT8 under long days, 8 hours later
than *CCA1* (Figure 4(a)) (Ezer *et al.*, 2017; Webb *et al.*, 2019). Intriguingly, as with *CCA1*,
many clusters showed a low proportion of rhythmic genes (<51%) on the 12th day after the SD
to LD transition. As for highly rhythmic clusters, they showed varying behaviour over
development. Clusters C and D were both highly rhythmic on the 2nd, 7th, and 12th days and
their expression broadly peaked before ZT8 on each day, suggesting that they were repressed
by the EC (Figure 4(b), 4(c); Supplemental Table S4, S5). The mean expression of cluster C
over each day slightly increased whereas this decreased for cluster D. Both clusters had an
overrepresentation of 'photosystem I' (GO:0009522) and multiple other photosynthesis related
terms (Supplemental Table S6). However, many regulatory GO terms were overrepresented
in cluster C but not in cluster D, such as 'regulation of biosynthetic process' (GO:0009889)
and 'response to red or far red light' (GO:0009639). *PHYTOCHROME-INTERACTING*
FACTOR5 (*PIF5*), which is phosphorylated and degraded in response to red-light, was in
cluster C (Shen *et al.*, 2007). This suggests that the interplay between the EC and
PHYTOCHROME B (*phyB*) (Ezer *et al.*, 2017) may be stable over development.

<u>Cell-type specific activity changes over diurnal and developmental timescales</u>

Since different clock components are enriched in various *Arabidopsis* leaf cell types (Nohales,
2021; Davis *et al.*, 2022), we wanted to know how cell-specific diurnal expression of genes
changes over age. Our previous work adapted a method, CIBERSORTx, to estimate cell type-
specific activity in bulk leaf RNA-seq samples (Newman *et al.*, 2019; Vong *et al.*, 2024). Most
cell types are predicted to have rhythmic activity on all three days of sampling (JTK_cycle,
BH.q value < 0.05; Supplemental Figure 7, Supplemental Table S8). However, some cell types
with low activity early in development are predicted not to be rhythmic, then become rhythmic
later in development, such as 'Unknown Group 3'. Vong *et al.,* 2024 suggested that these
groups may represent stressed or senescent cell types. Our results also confirm that phloem
and the unknown cell groups were expressed late in development, while the epidermal cells
were primarily transcriptionally active early in development (Figure 4(a)). We also confirm
consistent diurnal oscillations in guard cell and vascular cell transcriptional activity.

Coupling between the circadian clocks in mesophyll and vasculature has been demonstrated
(Endo *et al.*, 2014). We therefore sought to understand the behaviour of subgroups of

mesophyll. For instance, there were three groups of mesophyll cells that were each primarily
active at different times of day and different stages of development. Our results demonstrate
that the phase of their expression is consistent throughout development, but that the peak
expression increases in groups 1 and 2 over development and the minimum decreases in
group 3 (Figure 4(b)). These results highlight that any differentially expressed genes over
diurnal or developmental time may be a consequence of oscillations in which cells are
transcriptionally active, rather than a homogenous change of expression across all cell types.

<u>Known housekeeping genes display variation over both developmental and diurnal timescales</u>

Our previous analysis found many examples of genes that change their expression diurnally
and / or over development. However, it is also important to find sets of genes that are stably
expressed under both conditions. These are important controls in experiments like reverse
transcription followed by quantitative polymerase chain reaction (qRT-PCR), where gene
expression must be normalised to a housekeeper gene (Czechowski *et al.*, 2005; Souček *et*
al., 2017). Researchers often refer to well-known lists of housekeeping genes, which are
suggested to be stable under relevant experimental conditions. We calculated the coefficient
of variation (CV; standard deviation divided by mean) from our dataset of the suggested
housekeeping genes for developmental and diurnal experiments from ref. (Czechowski *et al.*,
2005). A high CV value indicates that a gene shows high biological variability gene expression
is not comparable across different times of day or developmental stages. Although the mean
CV value of the housekeeping genes was lower than the mean CV value across all genes,
many housekeeping genes still showed undesirable levels of variation (Figure 6(a)). For
example, two commonly used housekeeping genes, *PROTEIN PHOSPHATASE 2A* (*PP2A)*
SUBUNIT3 and *YELLOW-LEAF-SPECIFIC GENE8 (YLS8),* had large expression changes
over development (Figure 6(b)). *YLS8* approximately doubled in expression from the 2^{nd} day
of sampling to the 12^{th} day. However, some housekeeping genes did show promisingly low
CV scores. Genes *ARABIDOPSIS RNA POLYMERASE B13.6* (*ATRPB13.6;* AT3G52090*),*
AT3G03070*,* and AT3G12260 had the lowest CV values across all housekeeping genes
(Figure 6(a)). Despite the low CV values, we observed that these genes still displayed
variation across the diurnal timescale, when visualised on a relative expression scale (Figure
6(c)). We suggest that combinations of these genes may be useful as controls for qRT-PCR.

Discussion

We measured gene expression via RNA-seq in ageing plants, across both diurnal and
developmental timescales. Using PCA, we observed that most of the variation in samples can
be explained by a developmental PC, with a smaller contribution from two diurnal PCs (Figure

1). These PCA loadings help us establish the extent to which any individual gene varies its
expression across diurnal and developmental timescales.

Although kinases are thought to associate diurnal and developmental programmes, we found
that they primarily change their expression across developmental scales. However, many
genes whose expression changes are perturbed by KIN10 have diurnal expression patterns,
which is consistent with previous suggestions that there is crosstalk between the SnRK1
complex and circadian processes (Shin *et al.*, 2017; Chen *et al.*, 2017; Frank *et al.*, 2018).

We did not observe changes in the phase or period of core clock genes (Supplemental Figure
6). This is unsurprising given that our ageing plants were exposed to LD (entrainment)
conditions on the days of sampling, rather than free-running conditions, and it highlights the
stability of the central oscillator. However, it has previously been observed that the clock can
have different free-running periods in younger versus older leaves (Kim *et al.*, 2016). This is
consistent with our and previous reported observations that metabolic signalling inputs to the
clock, such as KIN10, have different levels of expression across plant development (Baena-
González *et al.*, 2007; Frank *et al.*, 2018).

Despite the relatively stable expression of CCA1 and ELF3 over development, their predicted
targets cluster into groups that either decrease or increase in mean expression (Figure 3(b),
Figure 4(b)). This could be because of changing patterns of core clock gene binding to
regulatory regions of the genome. However, these patterns may instead result from
downstream processing of circadian signals. For example, the clock influences senescence
through both transcriptional and *miR164*-mediated post-translation control of ORESARA1
(ORE1) (Kim *et al.*, 2018). The extent of these post-translations mechanisms could be
assessed by measuring the changes in the diurnal proteome and phosphoproteome in ageing
plants (Uhrig *et al.*, 2021). Our work suggests that there is value in spanning both
developmental and diurnal timescales and suggests that other complementary datasets, such
as ChIP-seq, proteomics, and phosphoproteomics, should be performed at more than one
timepoint per day and multiple stages of leaf development (Nagel *et al.*, 2015; Ezer *et al.*,
2017; Li *et al.*, 2023).

Diurnal rhythms of gene expression have previously been observed in different cell types,
through luciferase assays, qRT-PCR, and single-nucleus RNA-seq (Endo *et al.*, 2014;
Greenwood *et al.*, 2022; Qin *et al.*, 2023). Here, we utilised CIBERSORTx to deconvolute
gene expression from bulk RNA-seq data into contributions from leaf-specific cell types,
allowing us to see how diurnal patterns change over development (Newman *et al.*, 2019). We
identified some cell types that became rhythmic only at the latest stages of development. We
also showed that subgroups of mesophyll cells exhibit different timings of expression across

both timescales. It remains to be seen whether the reported coordination of the clock across
cell types is maintained across development (Endo *et al.*, 2014; Gould *et al.*, 2018; Schmal *et*
al., 2019).

We provide a valuable resource for the community to search for appropriate housekeeping
genes when researching the clock in ageing plants. Although existing housekeeping genes
show lower variability than background gene expression, we recommend that researchers use
at least one of our three suggested genes. Our suggested genes showed little variation across
the developmental timescale. Using multiple housekeeping genes to normalise qRT-PCR data
may counterbalance the unavoidable diurnal variation of housekeeping genes (Vandesompele
et al., 2002).

Acknowledgements
This project was undertaken on the Viking Cluster, which is a high-performance compute
facility provided by the University of York. The authors are grateful for computational support
from the University of York High Performance Computing service, Viking and the Research
Computing team. They also acknowledge support from the Horticulture Facility and the
Genomics Laboratory in the Technology Facility at the Department of Biology, University of
York.

Funding
This work was supported by funding from the Biotechnology and Biological Sciences Research
Council (BBSRC)—D.E. and S.J.D.: BB/V006665/1; D.E.: BB/S506795/1; and S.J.D.:
BB/N018540/1. The authors also acknowledge BBSRC White Rose DTP studentships
(BB/M011151/1 and BB/T007222/1) to J.R. (Ref.: 1792522) and E.J.R. (Ref.: 2444228). D.E.
was also supported by the Royal Society (RGS\R2\212345).

Competing interests
No competing interests are declared.

Author contributions
E.J.R.: conceptualization, methodology, software, formal analysis, investigation, writing—
original draft, and visualization. J.R.: conceptualization, investigation, and writing—review and
editing. S.J.D.: writing—review and editing, supervision, and funding acquisition. D.E.:
conceptualization, methodology, writing—original draft, writing—review and editing,
supervision, and funding acquisition.

Data availability

Raw and processed sequencing data have been deposited in the NCBI Gene Expression Omnibus (GEO) database under accession number GSE242964 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE242964).

Scripts used to produce the figures have been deposited in a GitHub repository https://github.com/stressedplants/AgeingCircadian

References

Anderson CM, **Wagner TA**, **Perret M**, **He ZH**, **He D**, **Kohorn BD**. **2001**. WAKs: cell wall-associated kinases linking the cytoplasm to the extracellular matrix. *Plant molecular biology* **47**: 197–206.

Andrade L, **Lu Y**, **Cordeiro A**, **Costa JMF**, **Wigge PA**, **Saibo NJM**, **Jaeger KE**. **2022**. The evening complex integrates photoperiod signals to control flowering in rice. *Proceedings of the National Academy of Sciences of the United States of America* **119**: e2122582119.

Andrews S. **2010**. FastQC: a quality control tool for high throughput sequence data.

Anwer MU, **Davis A**, **Davis SJ**, **Quint M**. **2020**. Photoperiod sensing of the circadian clock is controlled by EARLY FLOWERING 3 and GIGANTEA. *The Plant journal: for cell and molecular biology* **101**: 1397–1410.

Atamian HS, **Creux NM**, **Brown EA**, **Garner AG**, **Blackman BK**, **Harmer SL**. **2016**. Circadian regulation of sunflower heliotropism, floral orientation, and pollinator visits. *Science* **353**: 587–590.

Baena-González E, **Rolland F**, **Thevelein JM**, **Sheen J**. **2007**. A central integrator of transcription networks in plant stress and energy signalling. *Nature* **448**: 938–942.

Balcerowicz M, **Mahjoub M**, **Nguyen D**, **Lan H**, **Stoeckle D**, **Conde S**, **Jaeger KE**, **Wigge PA**, **Ezer D**. **2021**. An early-morning gene network controlled by phytochromes and cryptochromes regulates photomorphogenesis pathways in Arabidopsis. *Molecular plant* **14**: 983–996.

Berardini TZ, **Reiser L**, **Li D**, **Mezheritsky Y**, **Muller R**, **Strait E**, **Huala E**. **2015**. The Arabidopsis information resource: Making and mining the "gold standard" annotated reference plant genome. *Genesis* **53**: 474–485.

Broeckx T, **Hulsmans S**, **Rolland F**. **2016**. The plant energy sensor: evolutionary conservation and divergence of SnRK1 structure, regulation, and function. *Journal of experimental botany* **67**: 6215–6252.

Buckley CR, **Boyte JM**, **Albiston RL**, **Hyles J**, **Beasley JT**, **Johnson AAT**, **Trevaskis B**, **Fournier-Level A**, **Haydon MJ**. **2024**. A circadian transcriptional sub-network and EARLY FLOWERING 3 control timing of senescence and grain nutrition in bread wheat. *bioRxiv*: 2024.02.19.580927.

Cervela-Cardona L, **Yoshida T**, **Zhang Y**, **Okada M**, **Fernie A**, **Mas P**. **2021**. Circadian Control of Metabolism by the Clock Component TOC1. *Frontiers in plant science* **12**: 683516.

Chen L, **Su Z-Z**, **Huang L**, **Xia F-N**, **Qi H**, **Xie L-J**, **Xiao S**, **Chen Q-F**. **2017**. The AMP-
activated protein kinase KIN10 is involved in the regulation of autophagy in Arabidopsis.
Frontiers in plant science **8**: 1201.

Corbesier L, **Gadisseur I**, **Silvestre G**, **Jacqmard A**, **Bernier G**. **1996**. Design in Arabidopsis
thaliana of a synchronous system of floral induction by one long day. *The Plant journal: for cell*
and molecular biology **9**: 947–952.

Covington MF, **Maloof JN**, **Straume M**, **Kay SA**, **Harmer SL**. **2008**. Global transcriptome
analysis reveals circadian regulation of key pathways in plant growth and development.
Genome biology **9**: R130.

Czechowski T, **Stitt M**, **Altmann T**, **Udvardi MK**, **Scheible W-R**. **2005**. Genome-wide
identification and testing of superior reference genes for transcript normalization in
Arabidopsis. *Plant physiology* **139**: 5–17.

Davis W, **Endo M**, **Locke JCW**. **2022**. Spatially specific mechanisms and functions of the
plant circadian clock. *Plant physiology* **190**: 938–951.

Dodd AN, **Salathia N**, **Hall A**, **Kévei E**, **Tóth R**, **Nagy F**, **Hibberd JM**, **Millar AJ**, **Webb AAR**.
2005. Plant circadian clocks increase photosynthesis, growth, survival, and competitive
advantage. *Science* **309**: 630–633.

Duruflé H, **Hervé V**, **Balliau T**, **Zivy M**, **Dunand C**, **Jamet E**. **2017**. Proline Hydroxylation in
Cell Wall Proteins: Is It Yet Possible to Define Rules? *Frontiers in plant science* **8**: 1802.

Emanuelle S, **Hossain MI**, **Moller IE**, **Pedersen HL**, **van de Meene AML**, **Doblin MS**, **Koay**
A, **Oakhill JS**, **Scott JW**, **Willats WGT**, *et al.* **2015**. SnRK1 from Arabidopsis thaliana is an
atypical AMPK. *The Plant journal: for cell and molecular biology* **82**: 183–192.

Endo M, **Shimizu H**, **Nohales MA**, **Araki T**, **Kay SA**. **2014**. Tissue-specific clocks in
Arabidopsis show asymmetric coupling. *Nature* **515**: 419–422.

Ezer D, **Jung J-H**, **Lan H**, **Biswas S**, **Gregoire L**, **Box MS**, **Charoensawan V**, **Cortijo S**, **Lai**
X, **Stöckle D**, *et al.* **2017**. The evening complex coordinates environmental and endogenous
signals in Arabidopsis. *Nature plants* **3**: 17087.

Fowler S, **Lee K**, **Onouchi H**, **Samach A**, **Richardson K**, **Morris B**, **Coupland G**, **Putterill**
J. **1999**. GIGANTEA: a circadian clock-controlled gene that regulates photoperiodic flowering
in Arabidopsis and encodes a protein with several possible membrane-spanning domains.
The EMBO journal **18**: 4679–4688.

Frank A, **Matiolli CC**, **Viana AJC**, **Hearn TJ**, **Kusakina J**, **Belbin FE**, **Wells Newman D**,
Yochikawa A, **Cano-Ramirez DL**, **Chembath A**, *et al.* **2018**. Circadian Entrainment in
Arabidopsis by the Sugar-Responsive Transcription Factor bZIP63. *Current biology: CB* **28**:
2597-2606.e6.

Gould PD, **Domijan M**, **Greenwood M**, **Tokuda IT**, **Rees H**, **Kozma-Bognar L**, **Hall AJ**,
Locke JC. **2018**. Coordination of robust single cell rhythms in the Arabidopsis circadian clock
via spatial waves of gene expression. *eLife* **7**.

Greenwood M, **Hall AJW**, **Locke JCW**. **2022**. High Spatial Resolution Luciferase Imaging of
the Arabidopsis thaliana Circadian Clock. *Methods in molecular biology* **2398**: 47–55.

Havé M, **Marmagne A**, **Chardon F**, **Masclaux-Daubresse C**. **2017**. Nitrogen remobilization
during leaf senescence: lessons from Arabidopsis to crops. *Journal of experimental botany*
68: 2513–2529.

Hayama R, **Sarid-Krebs L**, **Richter R**, **Fernández V**, **Jang S**, **Coupland G**. **2017**. PSEUDO
RESPONSE REGULATORs stabilize CONSTANS protein to promote flowering in response
to day length. *The EMBO journal* **36**: 904–918.

Herrero E, **Kolmos E**, **Bujdoso N**, **Yuan Y**, **Wang M**, **Berns MC**, **Uhlworm H**, **Coupland G**,
Saini R, **Jaskolski M**, *et al.* **2012**. EARLY FLOWERING4 recruitment of EARLY
FLOWERING3 in the nucleus sustains the Arabidopsis circadian clock. *The Plant cell* **24**: 428–
443.

Hinckley WE, **Brusslan JA**. **2020**. Gene expression changes occurring at bolting time are
associated with leaf senescence in Arabidopsis. *Plant direct* **4**: e00279.

Hughes ME, **Hogenesch JB**, **Kornacker K**. **2010**. JTK_CYCLE: an efficient nonparametric
algorithm for detecting rhythmic components in genome-scale data sets. *Journal of biological*
rhythms **25**: 372–380.

Inoue K, **Araki T**, **Endo M**. **2018**. Circadian clock during plant development. *Journal of plant*
research **131**: 59–66.

Kim H, **Kim HJ**, **Vu QT**, **Jung S**, **McClung CR**, **Hong S**, **Nam HG**. **2018**. Circadian control of
ORE1 by PRR9 positively regulates leaf senescence in Arabidopsis. *Proceedings of the*
National Academy of Sciences of the United States of America **115**: 8448–8453.

Kim H, **Kim Y**, **Yeom M**, **Lim J**, **Nam HG**. **2016**. Age-associated circadian period changes in
Arabidopsis leaves. *Journal of experimental botany* **67**: 2665–2673.

Klingenberg CP. **2019**. Phenotypic Plasticity, Developmental Instability, and Robustness:
The Concepts and How They Are Connected. *Frontiers in Ecology and Evolution* **7**.

Kohorn BD, **Kohorn SL**. **2012**. The cell wall-associated kinases, WAKs, as pectin receptors.
Frontiers in plant science **3**: 88.

Kolberg L, **Raudvere U**, **Kuzmin I**, **Adler P**, **Vilo J**, **Peterson H**. **2023**. g:Profiler-
interoperable web service for functional enrichment analysis and gene identifier mapping
(2023 update). *Nucleic acids research* **51**: W207–W212.

Kolberg L, **Raudvere U**, **Kuzmin I**, **Vilo J**, **Peterson H**. **2020**. gprofiler2 -- an R package for
gene list functional enrichment analysis and namespace conversion toolset g:Profiler.
F1000Research **9**.

Krzymuski M, **Andrés F**, **Cagnola JI**, **Jang S**, **Yanovsky MJ**, **Coupland G**, **Casal JJ**. **2015**.
The dynamics of FLOWERING LOCUS T expression encodes long-day information. *The Plant*
journal: for cell and molecular biology **83**: 952–961.

Li M, **Yao T**, **Lin W**, **Hinckley WE**, **Galli M**, **Muchero W**, **Gallavotti A**, **Chen J-G**, **Huang S-**
SC. **2023**. Double DAP-seq uncovered synergistic DNA binding of interacting bZIP
transcription factors. *Nature communications* **14**: 2600.

Martin M. **2011**. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* **17**: 10–12.

Nagel DH, **Doherty CJ**, **Pruneda-Paz JL**, **Schmitz RJ**, **Ecker JR**, **Kay SA**. **2015**. Genome-
wide identification of CCA1 targets uncovers an expanded clock network in Arabidopsis.

Proceedings of the National Academy of Sciences of the United States of America **112**: E4802-10.

Newman AM, **Steen CB**, **Liu CL**, **Gentles AJ**, **Chaudhuri AA**, **Scherer F**, **Khodadoust MS**, **Esfahani MS**, **Luca BA**, **Steiner D**, *et al.* **2019**. Determining cell type abundance and expression from bulk tissues with digital cytometry. *Nature biotechnology* **37**: 773–782.

Nohales MA. **2021**. Spatial Organization and Coordination of the Plant Circadian System. *Genes* **12**.

Nozue K, **Covington MF**, **Duek PD**, **Lorrain S**, **Fankhauser C**, **Harmer SL**, **Maloof JN**. **2007**. Rhythmic growth explained by coincidence between internal and external cues. *Nature* **448**: 358–361.

Nusinow DA, **Helfer A**, **Hamilton EE**, **King JJ**, **Imaizumi T**, **Schultz TF**, **Farré EM**, **Kay SA**. **2011**. The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* **475**: 398–402.

O'Malley RC, **Huang S-SC**, **Song L**, **Lewsey MG**, **Bartlett A**, **Nery JR**, **Galli M**, **Gallavotti A**, **Ecker JR**. **2016**. Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. *Cell* **165**: 1280–1292.

Park DH, **Somers DE**, **Kim YS**, **Choy YH**, **Lim HK**, **Soh MS**, **Kim HJ**, **Kay SA**, **Nam HG**. **1999**. Control of circadian rhythms and photoperiodic flowering by the Arabidopsis GIGANTEA gene. *Science (New York, N.Y.)* **285**: 1579–1582.

Patro R, **Duggal G**, **Love MI**, **Irizarry RA**, **Kingsford C**. **2017**. Salmon provides fast and bias-aware quantification of transcript expression. *Nature methods* **14**: 417–419.

Petersen J, **Rredhi A**, **Szyttenholm J**, **Mittag M**. **2022**. Evolution of circadian clocks along the green lineage. *Plant physiology* **190**: 924–937.

Polge C, **Thomas M**. **2007**. SNF1/AMPK/SnRK1 kinases, global regulators at the heart of energy control? *Trends in plant science* **12**: 20–28.

Procko C, **Lee T**, **Borsuk A**, **Bargmann BOR**, **Dabi T**, **Nery JR**, **Estelle M**, **Baird L**, **O'Connor C**, **Brodersen C**, *et al.* **2022**. Leaf cell-specific and single-cell transcriptional profiling reveals a role for the palisade layer in UV light protection. *The Plant cell* **34**: 3261–3279.

Qin Y, **Liu Z**, **Gao S**, **Long Y**, **Zhu X**, **Liu B**, **Gao Y**, **Xie Q**, **Nohales MA**, **Xu X**, *et al.* **2023**. *A day in the life of Arabidopsis:* 24-hour time-lapse single-nucleus transcriptomics reveal cell-type specific circadian rhythms. *bioRxiv*: 2023.12.09.570919.

R Core Team. **2023**. R: A language and environment for statistical computing.

Ramon M, **Ruelens P**, **Li Y**, **Sheen J**, **Geuten K**, **Rolland F**. **2013**. The hybrid four-CBS-domain KINβγ subunit functions as the canonical γ subunit of the plant energy sensor SnRK1. *The Plant journal: for cell and molecular biology* **75**: 11–25.

Redmond EJ, **Ronald J**, **Davis SJ**, **Ezer D**. **2024**. Single-plant-omics reveals the cascade of transcriptional changes during the vegetative-to-reproductive transition. *The plant cell*: 2023.09.11.557157.

Rees H, **Rusholme-Pilcher R**, **Bailey P**, **Colmer J**, **White B**, **Reynolds C**, **Ward SJ**, **Coombes B**, **Graham CA**, **de Barros Dantas LL**, *et al.* **2022**. Circadian regulation of the transcriptome in a complex polyploid crop. *PLoS biology* **20**: e3001802.

Romanowski A, **Schlaen RG**, **Perez-Santangelo S**, **Mancini E**, **Yanovsky MJ**. **2020**. Global transcriptome analysis reveals circadian control of splicing events in Arabidopsis thaliana. *The Plant journal: for cell and molecular biology* **103**: 889–902.

Rubin MJ, **Brock MT**, **Davis AM**, **German ZM**, **Knapp M**, **Welch SM**, **Harmer SL**, **Maloof JN**, **Davis SJ**, **Weinig C**. **2017**. Circadian rhythms vary over the growing season and correlate with fitness components. *Molecular ecology* **26**: 5528–5540.

Sakuraba Y, **Jeong J**, **Kang M-Y**, **Kim J**, **Paek N-C**, **Choi G**. **2014**. Phytochrome-interacting transcription factors PIF4 and PIF5 induce leaf senescence in Arabidopsis. *Nature communications* **5**: 4636.

Sawa M, **Kay SA**. **2011**. GIGANTEA directly activates Flowering Locus T in Arabidopsis thaliana. *Proceedings of the National Academy of Sciences of the United States of America* **108**: 11698–11703.

Schmal C, **Ono D**, **Myung J**, **Pett JP**, **Honma S**, **Honma K-I**, **Herzel H**, **Tokuda IT**. **2019**. Weak coupling between intracellular feedback loops explains dissociation of clock gene dynamics. *PLoS computational biology* **15**: e1007330.

Seaton DD, **Smith RW**, **Song YH**, **MacGregor DR**, **Stewart K**, **Steel G**, **Foreman J**, **Penfield S**, **Imaizumi T**, **Millar AJ**, *et al.* **2015**. Linked circadian outputs control elongation growth and flowering in response to photoperiod and temperature. *Molecular systems biology* **11**: 776.

Shen Y, **Khanna R**, **Carle CM**, **Quail PH**. **2007**. Phytochrome induces rapid PIF5 phosphorylation and degradation in response to red-light activation. *Plant physiology* **145**: 1043–1051.

Shin J, **Sánchez-Villarreal A**, **Davis AM**, **Du S-X**, **Berendzen KW**, **Koncz C**, **Ding Z**, **Li C**, **Davis SJ**. **2017**. The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner. *Plant, cell & environment* **40**: 997–1008.

Song Y, **Jiang Y**, **Kuai B**, **Li L**. **2018**. CIRCADIAN CLOCK-ASSOCIATED 1 Inhibits Leaf Senescence in Arabidopsis. *Frontiers in plant science* **9**: 280.

Souček P, **Pavlů J**, **Medveďová Z**, **Reinöhl V**, **Brzobohatý B**. **2017**. Stability of housekeeping gene expression in Arabidopsis thaliana seedlings under differing macronutrient and hormonal conditions. *Journal of plant biochemistry and biotechnology* **26**: 415–424.

Streb S, **Zeeman SC**. **2012**. Starch metabolism in Arabidopsis. *The Arabidopsis book* **10**: e0160.

Thompson K, **Grime JP**, **Mason G**. **1977**. Seed germination in response to diurnal fluctuations of temperature. *Nature* **267**: 147–149.

Tsai AY-L, **Gazzarrini S**. **2012**. AKIN10 and FUSCA3 interact to control lateral organ development and phase transitions in Arabidopsis: Interaction between FUSCA3 and AKIN10. *The Plant journal: for cell and molecular biology* **69**: 809–821.

Uhrig RG, **Echevarría-Zomeño S**, **Schlapfer P**, **Grossmann J**, **Roschitzki B**, **Koerber N**, **Fiorani F**, **Gruissem W**. **2021**. Diurnal dynamics of the Arabidopsis rosette proteome and phosphoproteome. *Plant, cell & environment* **44**: 821–841.

Vandesompele J, **De Preter K**, **Pattyn F**, **Poppe B**, **Van Roy N**, **De Paepe A**, **Speleman F**.
2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging
of multiple internal control genes. *Genome biology* **3**: RESEARCH0034.

Vong GYW, **McCarthy K**, **Claydon W**, **Davis SJ**, **Redmond EJ**, **Ezer D. 2024**. AraLeTA: An
Arabidopsis leaf expression atlas across diurnal and developmental scales. *Plant physiology*
195: 1941–1953.

Wang F, **Han T**, **Jeffrey Chen Z**. **2024**. Circadian and photoperiodic regulation of the
vegetative to reproductive transition in plants. *Communications biology* **7**: 579.

Webb AAR, **Seki M**, **Satake A**, **Caldana C**. **2019**. Continuous dynamic adjustment of the
plant circadian oscillator. *Nature communications* **10**: 550.

Wendell M, **Ripel L**, **Lee Y**, **Rognli OA**, **Torre S**, **Olsen JE**. **2017**. Thermoperiodic control of
floral induction involves modulation of the diurnal FLOWERING LOCUS T expression pattern.
Plant & cell physiology **58**: 466–477.

Wu G, **Anafi RC**, **Hughes ME**, **Kornacker K**, **Hogenesch JB**. **2016**. MetaCycle: an integrated
R package to evaluate periodicity in large scale data. *Bioinformatics* **32**: 3351–3353.

Xiong L, **Zhou W**, **Mas P**. **2022**. Illuminating the Arabidopsis circadian epigenome: Dynamics
of histone acetylation and deacetylation. *Current opinion in plant biology* **69**: 102268.

Xu X, **Yuan L**, **Yang X**, **Zhang X**, **Wang L**, **Xie Q**. **2022**. Circadian clock in plants: Linking
timing to fitness. *Journal of integrative plant biology* **64**: 792–811.

Yeh Y-H, **Chang Y-H**, **Huang P-Y**, **Huang J-B**, **Zimmerli L**. **2015**. Enhanced Arabidopsis
pattern-triggered immunity by overexpression of cysteine-rich receptor-like kinases. *Frontiers*
in plant science **6**: 322.

Zhang K, **Halitschke R**, **Yin C**, **Liu C-J**, **Gan S-S**. **2013**. Salicylic acid 3-hydroxylase
regulates *Arabidopsis* leaf longevity by mediating salicylic acid catabolism. *Proceedings of the*
National Academy of Sciences of the United States of America **110**: 14807–14812.

Figure 1: (a) Every individual replicate plotted on the first two principal components (PCs), using log-transformed gene expression as inputs. The labelled numbers represent the time of day relative to dawn (zeitgeber time; ZT). Colour and shape represent the day of sample collection, according to the legend at the bottom. **(b)** The same as (a) but for the second and third PCs. **(c)** The distribution of genes according to the loadings on PC1 and PC2. Dashed vertical lines represent the 10% and 90% percentiles on PC1. Genes highlighted in red are plotted in (e). **(d)** The distribution of genes according to the loadings on PC2 and PC3. Core clock genes (as defined by Wang et al. 2024) are highlighted in red. **(e)** Expression of the genes with most positive and negative loadings on PC1 and two core clock genes (CCA1 and ELF4). The x-axis represents ZT on each day. Points represent individual biological replicates. Lines are drawn through the median value of the biological replicates.

Figure 2: (a) Expression of the two genes which encode the *Arabidopsis* orthologue of the α subunit of SnRK1, on the same y-axis so they can be compared. **(b)** Clustering of genes which showed increased expression, in KIN10 overexpressed *Arabidopsis* protoplasts, based on normalised gene expression over both timescales. The method 'ward.D2' was used to create the hierarchical clustering of the rows. **(c)** Same as (b) based on genes which showed decreased expression.

Figure 3: (a) Expression of CCA1. **(b)** Heatmap of targets from CCA1, from ChIP-seq data in (Nagel et al., 2015). Values represent log-transformed then z-scored gene expression, so expression can be compared across different genes. **(c)** Timeseries of targets, grouped by cluster. These follow the same vertical order as in (b). Dashed lines represent mean expression over all genes plotted and all timepoints. The percentages refer to the proportion of genes which were significantly rhythmic in each day, as predicted by JTK_cycle.

Figure 4: (a) Expression of ELF3. **(b)** Heatmap of targets from ELF3, from ChIP-seq data in (Ezer et al., 2017). Values represent log-transformed then z-scored gene expression, so expression can be compared across different genes. **(c)** Timeseries of targets, grouped by cluster, with the same labels as in Figure 3(c).

Figure 5: (a) Heatmap of cibersort outputs. Values represent the predicted proportion of gene expression from each sample that came from each cell type. Values show represent the median value of the three biological replicates, which have been scaled and then values higher than 3 or lower than -3 have been truncated. Rows have been clustered with hierarchical clustering using the "complete" method. **(b)** The proportion of three different groups of mesophyll cell types. Points represent the values for the individual replicates and lines are drawn through median values.

Figure 6: (a) Coefficient of variation (CV; calculated by standard deviation / mean) versus mean expression of genes. Both values are calculated including all 63 samples (all replicates at all timepoints). Only genes with a TPM > 10 are shown. Samples which were suggested by (Czechowski et al., 2005) as good housekeeping qPCR genes for diurnal experiments or developmental experiments (or both) are highlighted. **(b)** Expression of 5 housekeeping genes selected from (a). PP2A subunit 3 and YLS8 have relatively high CV values, and the other genes have the lowest 3 CV values. The y axes are fixed so the developmental variation can be compared across genes. **(c)** Same as (b), except the y axes vary to show diurnal variation.