Accept-reject decision-making revealed via a quantitative and ethological study of *C. elegans* foraging

Authors: Jessica A. Haley[1,2], Tianyi Chen[2], Mikio Aoi[3,4]*, and Sreekanth H. Chalasani[2]**[Lead Contact]

Affiliations:

[1] Neurosciences Graduate Program, University of California San Diego, La Jolla, CA, USA.

[2] Molecular Neurobiology Laboratory, Salk Institute for Biological Studies, La Jolla, CA, USA.

[3] Halıcıoğlu Data Science Institute, University of California San Diego, La Jolla, CA, USA.

[4] Department of Neurobiology, University of California San Diego, La Jolla, CA, USA.

*Correspondence: maoi@ucsd.edu

**Correspondence: schalasani@salk.edu

ORCID: 0000-0001-6282-7124 (J.A.H), 0009-0003-7572-2155 (T.C.), 0000-0002-7052-880X (M.A.), 0000-0003-2522-8338 (S.H.C.)

Abstract

Decision-making is a ubiquitous component of animal behavior that is often studied in the context of foraging. Foragers make a series of decisions while locating food (food search), choosing between food types (diet or patch choice), and allocating time spent within patches of food (patch-leaving). Here, we introduce a framework for investigating foraging decisions using detailed analysis of individual behavior and quantitative modeling in the nematode *Caenorhabditis elegans*. We demonstrate that *C. elegans* make *accept-reject* patch choice decisions upon encounter with food. Specifically, we show that when foraging amongst small, dispersed, and dilute patches of bacteria, *C. elegans* initially *reject* several bacterial patches, opting to prioritize exploration of the environment, before switching to a more exploitatory foraging strategy during subsequent encounters. Observed across a range of bacterial patch densities, sizes, and distributions, we use a quantitative model to show that this decision to *explore* or *exploit* is guided by available sensory information, internal satiety signals, and learned environmental statistics related to the bacterial density of recently encountered and *exploited* patches. We behaviorally validated model predictions on animals that had been food-

deprived, animals foraging in environments with multiple patch densities, and null mutants with

defective chemosensation. Broadly, we present a framework to study ecologically relevant foraging

decisions that could guide future investigations into the cellular and molecular mechanisms

underlying decision-making.

Keywords

decision-making, foraging, exploration, exploitation, dietary choice, *C. elegans*, quantitative modeling

Introduction

Decision-making is frequently defined as the selection of a course of action among several alternatives. The ubiquity and importance of decision-making has led to its use in describing a broad range of behaviors from taxes of unicellular organisms to economics and politics in human society [1, 2]. In decision neuroscience, laboratory experiments have investigated the behavioral choices of animals in well-controlled environments or tasks [3]. While these experiments have led to key insights into mechanisms underlying decision-making and related cognitive processes [4], a comprehensive understanding of decision-making remains elusive. Many researchers have advocated for a more neuroethological approach – suggesting that experiments designed to understand problems the brain evolved to solve offer a more rigorous framework [3, 5, 6]. Thus, one approach to enrich our understanding of decision-making is to look at the decisions made by animals foraging in naturalistic environments.

Foraging animals make a hierarchy of decisions to locate food (food search), choose between different food types (diet or patch choice), and allocate time spent within patches of food items (patch-leaving) [5, 7-10]. These foraging decisions often require that an animal *exploit* an environment for known resources or *explore* it for potentially better opportunities elsewhere [9, 11]. This exploration-exploitation trade-off requires cognitive computations such as learning the spatiotemporal distribution of food, route planning, estimation of food availability, and decision-making [12, 13]. According to optimal foraging theory, foragers may seek to maximize their rate of net energy gained over time [8, 9, 14] by using internal and external information to guide decision-making, especially in environments where resources are sparsely distributed or fluctuating. Although many studies have provided insight into the motivations, behavioral implementations, and genes associated with foraging decisions [9, 10], a more rigorous framework for exploring the neuronal mechanisms underlying foraging decisions is essential to establish a comprehensive understanding of decision-making.

The microscopic nematode *Caenorhabditis elegans* is well suited for investigating the cellular and molecular basis of foraging decisions [15]. A myriad of genetic tools, behavioral assays, and neuronal imaging techniques have been developed to take advantage of the species' quick reproductive cycle, isogeneity, optical transparency, and ease of maintenance [16-18]. While *C. elegans* typically feed upon a diversity of bacterial types in the wild [19], they are commonly maintained in the laboratory on agar plates containing large patches of the bacteria *Escherichia coli* as a food source

[16]. However, even in these simplified laboratory conditions and despite having a numerically

simple nervous system of only 302 neurons [16, 20], *C. elegans* display complex and robust species-

typical behaviors [21] involving learning and memory [22-25], and decision-making [26-31]. Recent

studies have demonstrated that ecologically focused environmental enrichment permits identification

of novel behaviors and gene functions in *C. elegans* and other animals [32-35].

Foraging animals often must make one of two types of decisions: *stay-switch* or *accept-reject*.

Stay-switch decisions refer to scenarios where an individual experiences diminishing returns with the

current action and must decide when to switch to a new action (e.g., patch-leaving and area-

restricted search) [5, 7-10]. In contrast, *accept-reject* decisions refer to situations where an individual

must decide between engaging with an option or ignoring it in search of a better one (e.g., diet or

patch choice) [5, 8, 9]. *Stay-switch* decisions have been well described in *C. elegans* [25, 26, 36-41] and

enable them to successfully explore an environment and exploit the bacteria within [28, 42-44].

However, to our knowledge, *accept-reject* decisions have not yet been demonstrated in *C. elegans*. While

C. elegans have been shown to alter food preferences in a diet choice assay, this behavior has only

been described by a set of *stay-switch* decisions [31, 36, 44]. The ability to make *accept-reject* decisions is

likely advantageous for foraging in fluctuating or variable environments as rejecting an encountered

food item of low quality creates the opportunity for a subsequent encounter with preferred food.

Thus, while *C. elegans* can successfully forage in patchily distributed environments, it is not known

whether *C. elegans* make decisions to *exploit* a patch of food upon encounter and how this decision-

making process changes over a series of patch encounters.

In this study, we aimed to identify if *C. elegans* make *accept-reject* decisions upon encounter

with bacterial patches. To answer this question, we performed a detailed analysis of the behavior of

individual animals foraging in an ecologically relevant environment where bacterial patches were

dilute and dispersed. In conditions where the bacterial density was much lower than that of

previously visited patches, animals first explored the environment before *accepting* patches for

exploitation. This explore-then-exploit strategy persists even when only one bacterial patch is

present in the environment. Using a theoretical framework, we found that this initial exploration

reflects a series of *accept-reject* decisions guided by signals related to the bacterial density of current

and recently *explored* and *exploited* patches as well as the animal's state of satiety.

Results

103   ***C. elegans* forage in patchy environments with an explore-then-exploit strategy**

104   To investigate whether *C. elegans* make *accept-reject* decisions upon encounter with bacterial

105   patches, we observed the behavior of animals foraging on an agar surface containing an isometric

106   grid of small, low density bacterial patches (Figure 1A, Video 1-2). Animals were confined to an

107   arena where their behavior was recorded and tracked for 60 minutes (Figure 1A, Figure 1 –

108   supplement 1, Video 3). During this time, animals were observed to move about the arena (Figure

109   1A-C, Figure 1 – supplement 2, Video 4) with each animal encountering, on average, ~8.4 total

110   patches and ~5.3 unique patches during the one-hour recording (Figure 1D). These patch

111   encounters ranged in duration from seconds to tens of minutes and could be classified as either

112   short or long (2+ minutes) using a Gaussian mixture model (Figure 1E, Figure 1 – supplement 3).

113   Notably, animals were significantly more likely to stay on patch for longer durations at later time

114   points (Figure 1F) which corresponded to an overall increase in their probability of residing on

115   patch (Figure 1G, Figure 1 – supplement 4). Specifically, we found that while animals initially resided

116   on bacterial patches at levels close to those predicted by chance, an animal's probability of residing

117   on a bacterial patch increased significantly after ~10 minutes. These findings, that patch duration

118   and residence increased over time, suggest that animals are more likely to exploit an encountered

119   patch as time goes on.

120   Given that the bacterial density of the experimental patches was approximately 20 times

121   more dilute than that of conditions animals experienced during development and immediately

122   preceding the assay, it was possible that the initial delay in exploitation could be due to animals'

123   inability to detect the presence of bacteria in these more dilute patches. We therefore set out to

124   determine if animals detected patches at all time points. Previous studies have shown that *C. elegans*

125   display an immediate and marked slowdown upon encounter with the edge of a food patch and

126   sustain these slower speeds on food [28, 45]. Therefore, velocity can be used as a proxy for an

127   animal's ability to sense a bacterial patch. Following this precedent, we quantified the instantaneous

128   velocity of individuals in our assay (Figure 1H). We observed significant deceleration of an animal's

129   instantaneous velocity upon encounter with the patch edge (Figure 1I), with no obvious difference

130   between short duration, early time point (Figure 1J) and long duration, late time point (Figure 1K)

131   encounters. Animals consistently displayed marked slowdown when approaching the patch edge

132   (Figure 1L), achieving an average deceleration of -19.5 $\mu m/s^2$ upon encounter with the bacterial

133   patch (Figure 1M, Figure 1 – supplement 5). This slowdown was statistically significant for both long

134   and short duration encounters (Figure 1M). Further, consistent with *C. elegans* behavior on larger and

135 more densely seeded patches in other studies [46], animals in our assay maintained significantly lower

136 average velocities on patch (~52 μm/s) compared to off patch (~198 μm/s) (Figure 1N,O). These

137 results suggest that, despite detecting the availability of food during early patch encounters, *C. elegans*

138 opt to initially prioritize exploration of the environment before switching to a more exploitatory

139 foraging strategy during subsequent patch encounters.

The timing of the switch from exploration to exploitation varies with bacterial density

To determine whether this explore-then-exploit foraging strategy is dependent upon food-

related characteristics of the environment, we varied the density of the bacterial patches. By diluting

bacterial stocks and controlling growth time, we created 12 bacterial density conditions with relative

density ranging from 0 to ~200 (Figure 2A, Figure 2 – supplement 1-3). At the low end of this range

(densities less than ~0.5), animals do not appear to detect bacterial patches as indicated by on-patch

velocities matching those of animals foraging on bacteria-free (density 0) patches (Figure 2B,C). At

the high end of the range (density ~200), bacterial density is comparable to the environments

animals experienced during development and immediately prior to the assay (Figure 2 – supplement

2-3).

Previous studies have shown that in diet choice assays where animals are given the option

between patches of varying quality or density, animals spend more time on preferred patches [31, 36,

44]. We hypothesized that even when all patches in the environment are of equal density, an efficient

forager should still modulate the amount of time spent on the bacterial patches in a density-

dependent manner. To test this, behavior was recorded and tracked for one hour as animals foraged

in patchy environments matching one of the bacterial density conditions (Figure 2D, Figure 2 –

supplement 4). Consistent with our hypothesis, we found that the total time animals spent on

bacterial patches increased monotonically with increasing bacterial density following a sigmoidal

trend (Figure 2E). Further, the probability of an animal residing on patch as a function of time was

highly dependent on the bacterial density (Figure 2F, Figure 2 – supplement 5). At the highest

densities (50 and 200), animals spent nearly 100% of time in the assay on bacterial patches, while at

the lowest densities (0 to 0.3), animals spent chance levels of time on patch for the duration of the

experiment. At intermediate densities (0.5 to 10) animals initially resided on patches at rates

predicted by chance, later transitioning to more time spent on patch. Notably, this delayed increase

in patch residence is density dependent with animals switching earlier in environments with patches

of greater density.

Consistent with our previous observation that animals reside on patches for either short or

long durations (Figure 1E), we observed two distinct patch encounter types: 1) short (less than 2

minute) patch visits and 2) long (often tens of minutes) patch visits (Figure 2G). Using a two-

component Gaussian mixture model (GMM), we classified all encounters based on the duration of

these patch visits and animals' average velocity during the encounter (Figure 2H, Figure 2 –

supplement 6). We identified two behaviorally distinct clusters: 1) short duration and fast velocity

(*explore*), and 2) long duration and slow velocity (*exploit*) encounters. The *explore* encounters suggest

that an animal has *rejected* a patch, opting to continue exploration of the environment. In contrast,

the *exploit* encounters suggest that an animal has *accepted* a patch and is consuming the bacteria

within.

While we previously found that animals consistently slowed down upon encounter with the

patch edge when foraging in relative density ~10 (Figure 1J-M), we observed that a large portion of

short duration, exploratory encounters were not accompanied by this slow down (Figure 2 –

supplement 7A-B), especially for the lowest bacterial densities tested. Therefore, to determine which

exploratory encounters displayed evidence that the patch was detected by the animal, we further

classified encounters by an animal's minimum velocity on patch, deceleration upon patch entry, and

maximum change in the animal's velocity upon encounter (i.e., the difference between the peak

velocity immediately before the patch encounter and the minimum velocity achieved during the

patch encounter) (Figure 2I, Figure 2 – supplement 7, Video 5). We again observed two behaviorally

distinct clusters: 1) large slow-down and slow on-patch velocity, and 2) small slow-down and fast on-

patch velocity. We classified these encounters as *sensing* and *non-sensing*, respectively, using a semi-

supervised quadratic discriminant analysis (QDA) approach. The presence of a slow-down and the

subsequent continuation of slower on-patch velocity suggest that an animal sensed the bacteria,

while the absence of a slowdown suggests that the animal likely did not perceive the encounter. For

simplicity, we refer to 1) *non-sensing* encounters as *searching* (i.e., on-patch behavior that matches off-

patch behavior where an animal appears to be searching for food); 2) *sensing*, short duration

encounters as *sampling* (i.e., evaluating a patch's suitability as a food source, but ultimately *rejecting* the

patch and continuing to explore the environment); and 3) *sensing*, long duration encounters as

exploiting (i.e., *accepting* a patch and consuming its bacteria) (Figure 2J). Consistent with our

interpretation of *searching* encounters, the average velocities that animals achieved during *search* on

(~200 µm/s) and off (~207 µm/s) patch were similar, especially when compared to markedly slower

on-patch velocities during *sample* (~114 µm/s) and *exploit* (~57 µm/s) encounters (Figure 2K). In

summary, we have identified three distinct patch encounter types: encounters that were not detected

(*search*) and encounters that were detected with the animal subsequently deciding to *reject* (*sample*) or

accept (*exploit*) the patch.

We next characterized how the frequency of encounter types varied as a function of time

and patch density. The behavior of 443 animals (20-50 worms per condition) was tracked and

classified at each time point (Figure 2L, Video 6). In low density conditions (0 to 0.3), animals spent

nearly all of their time *searching* on or off patch. With increasing bacterial patch density, animals

spent less time *searching* and more time *sampling* and *exploiting*. Animals foraging in medium density

patches (0.5 to 10) initially spent most of their time *sampling* patches upon encounter. However,

animals eventually switched to mostly *exploiting* patches. Animals foraging on high density patches

(50 and 200) almost exclusively *exploited*. Classification of on-patch behavior across all densities

tested revealed a consistent switch from *exploratory* behaviors (*searching* and *sampling*) to *exploitation*

(Figure 2M) with the timing of the switch occurring earlier for animals foraging in environments

with higher bacterial patch density (Figure 2N). The delay in exploitation corresponded to a

comparatively large number of initial exploratory encounters with animals on medium density

conditions (0.5 to 10) *exploiting* after an average of 4.8 to 18.7 exploratory encounters (Figure 2O).

To summarize, for densities similar to those experienced immediately preceding the experiment (50

and 200), animals immediately *exploit* available bacteria. On the other hand, for bacterial densities

that appear to be below an animal's sensory threshold (0 to 0.3), animals never *exploit* patches and

spend the entire 60-minute assay *searching* for food. When foraging in medium density patches (0.5 to

10), animals initially explore the environment via a combination of *searching* and *sampling* encounters

and eventually switch to *exploiting*. Notably, the timing of this switch is density-dependent with

animals switching earlier when foraging on higher density patches.

Animals explore before exploiting even when only one patch is available

Given that bacterial patches within an environment were relatively invariable and that

bacterial growth during the experiment was negligible (Figure 2 – supplement 3), it was surprising

that animals were often willing to *reject* 10+ patches before *exploiting* a patch of the same density as

those previously *rejected* (Figure 2O). This *sampling* behavior may be beneficial when food is patchily

distributed in the environment as exploiting a dilute patch may result in a lost opportunity for future

encounter with a higher quality patch. Thus, we hypothesized that an animal's willingness to *accept* a

patch whose density matched that of a previously *rejected* patch could be a specific feature of a

patchily distributed environment where numerous observations of individual patches could contribute to learning the features of a changing environment. To address this, we observed whether animals were willing to *reject* a patch numerous times before *accepting* it even when only one patch is available in the environment. We created environments with single large (~8.3 mm) or small (~1.8 mm) diameter patches with relative density ranging from 0 to ~400 (Figure 3A-C). Behavior was tracked for 60 minutes as animals foraged in these environments (Figure 3D-E, Figure 3 – supplement 1-2). We found that even in these single patch environments, animals initially *explore* (*search* and *sample*) before *exploiting* (Figure 3F-G) in a density-dependent manner. Just as when foraging in a patchily distributed environment (Figure 2N), animals *explored* (*search* and *sample*) the small and large single patches numerous times prior to *exploiting* with the timing of this switch occurring significantly earlier with increasing bacterial density (Figure 3H-I). These results suggest that the explore-then-exploit strategy employed by *C. elegans* is not specific to a patchily distributed environment. Rather, animals may *sample* the same patch numerous times before deciding to *exploit*, especially when foraging on dilute patches. The decision of when to stop *exploring* and start *exploiting* is highly dependent upon the density of bacteria within a patch with the switch occurring earlier when animals encounter patches of higher density.

External and internal sensory information guide the decision to explore or exploit

While our behavioral analysis demonstrated that *C. elegans* modulate their decision to *explore* or *exploit* in a density-dependent manner, it remains unclear whether these animals use more complex cognitive processes such as learning and memory to guide their foraging decision. It is plausible that animals are making decisions using simple heuristics given only available sensory information about the current patch. However, previous studies have shown that food-related behaviors in *C. elegans* can be driven by internal states [31] as well as memories of the environment [25, 36, 42]. Thus, to better understand the factors driving the exploitation decision, we implemented a generalized linear model (GLM) to test how foraging decisions are influenced by the density of the current patch as well as additional food-related factors such as an animal's level of satiety and prior experience (Figure 4A).

We considered that the probability of exploiting a patch upon any given encounter can be described by a logistic function:

$$p(y_k = 1 | \boldsymbol{\beta} \cdot \boldsymbol{x}_k) \; = \; \frac{1}{1 + e^{-\boldsymbol{\beta} \cdot \boldsymbol{x}_k}}$$

where $p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)$ represents the conditional probability that an animal exploited during

patch encounter k; $\boldsymbol{x}_k \in R^n$ is a vector of covariates; and $\boldsymbol{\beta} \in R^n$ is a vector of weights that

describes how much each covariate influences the animal's choice. In models compared here, $\boldsymbol{x}_k$

includes a combination of a constant element and covariates that may empirically influence the

decision to *exploit*. We considered the simplest model where every encounter is independent with

fixed probability of *exploiting* (i.e., $p(y_k = 1|\beta_0)$) where β_0 represents the average animal's

propensity to *exploit*. We compared this null model against a set of nested models containing

covariates that vary from encounter to encounter and relate to: 1) the relative density of the

encountered patch k (ρ_k), 2) the duration of time spent off food since departing the last *exploited*

patch (τ_s), 3) the relative density of the patch encountered immediately before encounter k (ρ_h),

and 4) the relative density of the last *exploited* patch (ρ_e). Using this model design (i.e., $\boldsymbol{\beta} \cdot \boldsymbol{x}_k =$

$\beta_0 + \beta_k \rho_k + \beta_s \tau_s + \beta_h \rho_h + \beta_e \rho_e$), we assessed how each of these covariates influences the

probability of *exploitation* at each patch encounter k (Figure 4A).

Given that our aim is to understand what factors influence an animal's decision to *explore* or

exploit, we rationalized that encounters where the animal did not detect the presence of bacteria were

unlikely to contribute to decision-making. Therefore, we excluded *searching* encounters from our

analysis where the bacterial patch was not *sensed*. To account for the uncertainty in our classification

of *sensing*, we probabilistically included patch encounters with frequency equal to the probability that

the bacterial patch was *sensed* (Figure 4B) as previously estimated by semi-supervised QDA (Figure

2I, Figure 2 – supplement 7). In doing so, we simulated sets of observations of *sensed* (*sample* and

exploit) encounters. Further, to account for the uncertainty in our classification of patch encounters

as *exploration* or *exploitation*, we fit our GLM to the probability of *exploiting* $p(y_k = 1|\boldsymbol{z}_k)$ as

previously estimated by GMM (Figure 2H, Figure 2 – supplement 6) rather than fitting to direct

observations of exploitation y_k (see Models of exploitation probability).

When selecting covariates for our model, we calculated the Bayesian information criterion

(BIC) – a model selection test to avoid overfitting by penalizing increases in the number of

parameters – and observed improved model performance upon addition of each covariate (Figure 4

– supplement 1A). Further, we found that all covariates (i.e., ρ_k, τ_s, ρ_h, and ρ_e) significantly

contribute to the decision to *exploit* $p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)$ as indicated by coefficient values significantly

greater than or less than zero (Figure 4C, Figure 4 – supplement 1B). Specifically, we found that β_k

and β_s are significantly greater than zero which suggests that animals are more likely to *exploit*

patches with increasing patch density ρ_k and duration of food-deprivation τ_s. On the other hand,

β_h and β_e are significantly negative, which suggests that the greater the density of a recently

encountered or *exploited* patch, ρ_h and ρ_e, the less likely an animal should be willing to *exploit* the

current patch.

In order to better understand how each of the covariates in our model affect the *exploitation*

decision, we examined the probability of observing *exploitation* for the first time as a function of the

cumulative number of patches an animal encountered. Exploitation events were simulated from a

Bernoulli distribution with probability estimated by the model (i.e.,

$y_k | \boldsymbol{x}_k \sim \text{Bern}\big(p(y_k = 1 | \boldsymbol{\beta} \cdot \boldsymbol{x}_k)\big)$) or observed from our classification of *exploitation* (i.e.,

$y_k | \boldsymbol{z}_k \sim \text{Bern}\big(p(y_k = 1 | \boldsymbol{z}_k)\big)$). Although the probability of *exploiting* for the first time was not

expressly fit by our model, this derived metric highlights the model's ability to predict the delay in

exploitation observed for animals foraging on low to medium density bacterial patches in our

experiments (Figure 4D, Figure 4 – supplement 2). For the simplest model where every encounter is

independent with fixed probability of *exploiting* $p(y_k = 1 | \beta_0)$, we find that the probability of

exploiting for the first time is greatest for the first encounter and decreases for every subsequent

encounter (Figure 4D, Figure 4 – supplement 2), fitting a geometric distribution.

We next considered whether the observed delay in *exploitation* could be explained by a simple

density-dependent model (i.e., $p(y_k = 1 | \beta_0 + \beta_k \rho_k)$). This model predicts that animals are more

likely to *exploit* bacterial patches of greater density with probability proportional to the relative

density of the current patch ρ_k. While this density-dependent model accurately predicted that the

first *exploitation* occurs earlier on average for higher density patches, it also predicted that, regardless

of patch density, the first *exploitation* is most frequently observed on the first encounter which did

not match the observed distribution (Figure 4D, Figure 4 – supplement 2). We subsequently

considered that the onset of food-deprivation during *exploration* of these low-density environments

could drive the observed time-dependent delay in *exploitation*. We assume that animals become

satiated during *exploitation* and use the duration of time an animal spent searching off-food since the

last *exploitation* event (i.e., τ_s) as a proxy for the animal's likely decrease in satiety. With the addition

of this satiety-related signal (i.e., $p(y_k = 1 | \beta_0 + \beta_k \rho_k + \beta_s \tau_s)$), we observed a slight delay in when

animals are predicted to first *exploit* for the lowest density conditions tested (Figure 4D, Figure 4 –

supplement 2). To validate our finding that animals use a satiety-related signal to inform decision

making, we applied our model to observations of animals foraging in patchily distributed

environments following 3 hours of food-deprivation (Figure 4 – supplement 3A-B). We found that

only with the inclusion of a satiety-related covariate in our model, were we able to reliably predict

the immediate *exploitation* that food-deprived animals exhibit (Figure 4 – supplement 3C). Altogether,

these results suggest that animals use available external and internal sensory information to guide the

decision to *exploit*.

Prior experience guides the decision to *explore* or *exploit*

Although the current patch density and satiety covariates significantly influence the decision

to *exploit*, this model (i.e., $p(y_k = 1|\beta_0 + \beta_k\rho_k + \beta_s\tau_s)$) does not accurately predict the extent that

exploitation was delayed. We hypothesized that because animals had experienced very densely seeded

bacterial patches immediately prior to the assay, animals may have built an expectation that high

density bacterial patches are available in the new environment. We would therefore expect an initial

suppression in the probability of *exploiting* while animals updated their expectation using information

learned from *sampling* patches in the current environment. To test whether the decision to *exploit*

could be influenced by prior experience, we added a history-dependent covariate to our model (i.e.,

$p(y_k = 1|\beta_0 + \beta_k\rho_k + \beta_s\tau_s + \beta_h\rho_h)$ where ρ_h is the relative density of the patch encountered

immediately before encounter k). We find that inclusion of the density of the most recently

encountered patch significantly improves our prediction of *exploitation* (Figure 4D, Figure 4 –

supplement 2). We then considered whether the animal's expectation of bacterial density in the

environment might be a longer lasting memory. Specifically, we considered if the relative density of

the most recently *exploited* patch ρ_e could have influenced the animal's decision to *exploit*. The

addition of this second history-dependent term to our model (i.e.,

$p(y_k = 1|\beta_0 + \beta_k\rho_k + \beta_s\tau_s + \beta_h\rho_h + \beta_e\rho_e)$) further improved the fit of our model (Figure 4D,

Figure 4 – supplement 2) suggesting that *C. elegans* use recent experiences of encountered and

exploited patches to guide their decision to *exploit*. Animals may use this learned information to create

and update an expectation for the relative density of bacterial patches available in their environment

and compare that estimate against the density of a subsequently encountered patch. Altogether,

these results suggest that animals use prior experience to keep track of available resources and

modulate their decision to *explore* or *exploit*, a strategy that is highly beneficial for behavioral

adaptation in a changing environment.

To validate the role of prior experience in guiding foraging decisions, we observed animals

foraging in environments with multiple patch densities (Figure 4E, Figure 4 – supplement 4A-C).

Our model predicts that the lower the density of recently encountered and/or *exploited* patches, the

more likely an animal should be to *exploit* a subsequent patch. Without the history-dependent terms

ρ_h and ρ_e, our model predicts that no relationship should exist between prior experience and the

probability of *exploiting* the current patch. To test which model best fit the animals' behavior, we

calculated the conditional probabilities of *exploiting* $p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)$ for this new multi-density,

multi-patch data set with more nuanced combinations of current (ρ_k) and recent patch (ρ_h and ρ_e)

density using the coefficients $\boldsymbol{\beta}$ learned previously (Figure 4C, Figure 4 – supplement 4D). We

observed significantly better predictions of *exploitation* probability for the model including prior

experience as compared to the model without the terms ρ_h and ρ_e (Figure 4F) as confirmed by a

likelihood-ratio test. We observed an augmentation in *exploitation* when animals move from low to

high density (e.g., 1 to 4 or 1 to 7) as predicted by the model with history-dependence, but not the

model without. It is important to note, that this improvement in prediction was achieved without

fitting the model to this new multi-density data set. In summary, our modeling results suggest that

animals evaluate bacterial density, monitor internal signals of satiety, and leverage prior experience to

drive their decision to *exploit* a bacterial patch upon encounter.

371 Chemosensation is required for the evaluation of patch density

The quantitative modeling applied in this study provides a useful framework for testing the

influence of behaviorally relevant features on decision-making. To validate this approach and further

elucidate how animals evaluate current and recent patch density, we assessed the foraging behavior

of mutants deficient in a sensory modality (Figure 4 – supplement 5). We found that animals with

reduced function of non-ciliated mechanoreceptor neurons due to a null mutation in the *mec-4* gene

behave similarly to wild-type N2 animals, displaying increasingly delayed *exploitation* as patch density

decreases (Figure 4G, Figure 4 – supplement 5). Contrastingly, animals with reduced function of

ciliated chemoreceptor neurons due to a null mutation in the *osm-6* gene displayed immediate

exploitation even while foraging on low density patches (Figure 4G, Figure 4 – supplement 5). These

results indicate altered decision-making in the chemosensory mutants. Specifically, it appears that the

influence of patch density on the *exploitation* decision is reduced in *osm-6* mutants and that these

animals are more likely to *accept* a patch by default.

To further elaborate on these findings, we fit our GLM to each of the three data sets (i.e.,

N2, *mec-4*, *osm-6*) and assessed the difference in the estimated distributions of coefficients $\boldsymbol{\beta}$ between

mutant and wild-type models (Figure 4H). We hypothesized that the density-related covariates (ρ_k,

ρ_h, and ρ_e) would have smaller magnitude (i.e., less influence on decision-making) for animals with

reduced sensation of bacterial density. Consistent with this hypothesis, we found a significant

reduction in the magnitude of the influence of current patch density (i.e., β_k) on the decision to

exploit in *osm-6* mutants. Further, we found a significant increase in the intercept β_0 in *osm-6* mutants

which corresponds to the overall increase in *exploitation* observed in these animals (Figure 4G, Figure

4 – supplement 5). These results suggest that chemosensation plays an important role in evaluating

the density of the current patch and that when this sensory modality is impaired, animals modify

their default behavioral strategy to prioritize *exploitation*. We also observed a marginal increase in our

estimate of the influence of recently *exploited* patches (i.e., β_e) in *osm-6* mutants which suggests that

reduced efficacy in the evaluation of current patch density also affects learning and memory of

patch density. No significant changes were detected in the *mec-4* mutants. Although we cannot rule

out a role for mechanosensation in contributing to the assessment of patch density, this modality

likely provides less salient information than chemosensation. Altogether, these results suggest that

chemosensation is necessary for the evaluation of patch density and that when this modality is

impaired, animals prioritize *exploitation*. Further, these results demonstrate that this quantitative

modeling approach can be used to probe mechanisms underlying decision-making.

Discussion

Like all foraging animals, *C. elegans* must choose between *exploiting* an environment for known

resources and *exploring* it for potentially better opportunities elsewhere [5, 7-11]. Previous studies

have shown that *C. elegans* use *stay-switch* decisions to efficiently explore a patchily distributed

environment and exploit the bacteria within [28, 42-44]. In this study, we show that *C. elegans* also

make *accept-reject* decisions while foraging. Specifically, we observe that animals often initially *reject* low

density bacterial patches, opting to prioritize exploration of the environment, before switching to a

more exploitatory foraging strategy during subsequent encounters. This explore-then-exploit

strategy was consistent across a range of bacterial patch densities, sizes, and distributions. In order

to better understand this phenomenon and identify the factors that contribute to the decision to

exploit, we leveraged a quantitative modeling approach to hypothesis testing. We found that animals

do not employ a default explore-then-exploit strategy; rather, this behavior reflects a series of *accept-*

reject decisions guided by multimodal information related to an animal's environment and internal

state. Specifically, we show that each decision to *explore* or *exploit* is informed by available sensory

information, internal satiety signals, and learned environmental statistics related to the bacterial

density of recently encountered and *exploited* patches.

C. elegans make _accept-reject_ foraging decisions

While *C. elegans* have been shown to alter food preferences in a diet choice assay, this

behavior has only been described by a set of *stay-switch* decisions [31, 36, 44]. Specifically, *C. elegans*

were found to modulate their patch-leaving frequency to *stay* on high quality bacterial patches and

switch with declining food quality and quantity [26, 36, 40, 41]. The overall effect of leaving low

quality patches more frequently is that animals are more likely to reside on high quality patches.

Thus, *stay-switch* decision-making enables *C. elegans* to efficiently allocate time spent between

exploring or exploiting in a manner consistent with optimal foraging theory [5, 7-11]. However,

optimal foraging theory also predicts that an animal should never exploit (i.e., always *reject*) a low

density patch if higher density patches are sufficiently abundant as inclusion of the low density

patch type in the animals' diet decreases the overall rate of energy gained [5, 9]. Thus, while leaving

low density patches more frequently drives an overall increase in the rate of energy gained, it can still

result in sub-optimal foraging behavior.

Here, we demonstrate that *C. elegans* are capable of making *accept-reject* decisions upon

encounter with bacterial patches. We employed unbiased and quantitative methods – GMM and

semi-supervised QDA – to estimate whether animals *sensed* a patch and, if they did, whether they

accepted or *rejected* it. We found that animals initially *rejected* low density patches, opting to continue

exploration of the environment, and that this *reject* decision did not reflect an animal's inability to

detect the presence of bacteria in these dilute patches. Further, we found that the decision to *explore*

or *exploit* is density-dependent and robust across a range of bacterial densities, distributions, and

sizes. The initial preference for *exploration* during foraging on low density patches suggests that

animals may be *searching* for denser bacterial patches such as those experienced immediately

preceding the assay and throughout development. This explore-then-exploit strategy is advantageous

for animals when bacterial patch density is low or variable. On the other hand, we observed

immediate exploitation of high density patches. In these environments, an explore-then-exploit

strategy is disadvantageous, as delaying exploitation when patch density is already high results in

spending more energy on food search with no expected increase in energy gained during *exploitation*

of a subsequently encountered patch. Thus, consistent with an energy maximizing optimal forager,

C. elegans explores first when bacterial patch density is low, but immediately *exploits* when bacterial

patch density is sufficiently high. Altogether, our results demonstrate that *C. elegans* make a distinct

decision to either *explore* or *exploit* a patch upon encounter and that the density-dependence of this

decision is consistent with predictions from optimal foraging theory.

Our finding that *C. elegans* are willing to *reject* numerous encountered patches is somewhat

surprising, especially given that *C. elegans* likely possess limited spatial memory [25] and do not

appear to detect the presence of bacterial patches prior to encounter as slow-down only occurs

within one body-length of the patch edge (Figure 2B). Further, *C. elegans* are likely unaware of how

many patches are in the environment as evidenced by their willingness to *reject* a patch even when

only one patch is available. Taken together, this suggests that animals are "knowingly" passing up an

opportunity to *exploit* an immediately available food source for the potential opportunity to *exploit*

preferred food later. This type of behavior has been described min humans and other animals in the

context of delayed gratification [47, 48], intertemporal choice [47, 49, 50], self-control [51, 52], and

optimal stopping [53, 54]. Notably, few animals display the capacity for delayed gratification when

given the option between eating an immediately available but less preferred food item and waiting

for a preferred food item [55-58]. Even fewer animals are willing to wait if the delayed reward is not

visible or increases in quantity rather than quality [59, 60]. Our observation that *C. elegans* can exhibit

delayed gratification suggests that this and similar processes require less complex cognition than

previously considered or that we may have underestimated the cognitive capacity of *C. elegans*.

Prior experience and satiety guide foraging decision-making

Stay-switch decisions in *C. elegans* are strongly modulated by a variety of internal and external

sensory cues over a range of behavioral timescales. For example, patch-leaving frequency can be

modulated by an animal's prior experience [25, 42], metabolic status, arousal state [31], as well as the

presence of pathogens [61, 62], chemorepellents [63], predators [64, 65], and varying levels of

environmental O_2 and CO_2 [40, 66].

Here, we used a quantitative framework to show that prior experience (i.e., the density of

current and recently encountered and *exploited* bacterial patches) guides *accept-reject* decision-making.

According to early models in optimal foraging theory, foragers behave as if they have *complete*

information about the environment (e.g., animals are assumed to know the average density of bacterial

patches) [9]. Later theoretical and empirical work expanded models of patch choice to accommodate

situations where animals have *incomplete information* (e.g., environments with fluctuating resources)

and, therefore, must both acquire information and forage [8, 10, 67]. This work suggests that

animals may continuously sample areas in their environment to keep track of resource availability.

Here, we found that animals initially choose to *sample* patches and that the duration of this period of

sampling is correlated with the degree of mismatch between current and recently experienced and

exploited patches (i.e., animals spend more time *sampling* as the magnitude of difference between the

patch density experienced immediately before and during the experiment increases). When there is

no difference between current and prior experience (i.e., animals foraging on patches of relative

density 200), animals rarely *sample*. These results suggest that *C. elegans* use *sampling* as a way of re-

evaluating the overall quality of patches available in the environment when a change is detected.

Further, the results of our model suggest that animals likely use the information learned during

sampling encounters to guide their decision to *explore* or *exploit* on subsequent patch encounters. We

behaviorally validated this hypothesis by applying our model predictions to animals foraging in

environments with multiple patch densities. We predicted and observed that the lower the density of

recently encountered or *exploited* patches and the higher the density of the current patch, the more

likely an animal would be to *exploit*. These results suggest that *C. elegans* can learn and remember

features of recent experiences and use that learned information to guide efficient decision-making.

Incorporating prior experience into the decision to *explore* or *exploit* currently available food is highly

beneficial when foraging in a fluctuating environment. *C. elegans* thus may have evolved this ability to

maximize foraging in the wild where animals experience a boom-and-bust environment [68].

In addition to prior experience, internal state signals serve a critical role in modifying

decision-making in *C. elegans* [31, 69]. For example, food-deprived animals are more likely to cross an

aversive barrier (e.g., copper, primaquine, glycerol) in search of food as compared to well-fed

controls [69, 70]. Here, we find that a satiety-related signal likely drives the decision to *exploit*.

Specifically, our model suggests that as animals' satiety decreases, their willingness to accept a lower

quality patch increases. We confirmed this prediction by analyzing the behavior of food-deprived

animals, which immediately *exploit* even when bacterial patches are very dilute. Combined with the

effect of prior experience, these results suggest that the decision to *explore* or *exploit* likely requires

integration of food-related internal and external cues. Future studies should probe if additional

factors related to the animal's biotic and abiotic environment [35, 71-74] as well as alternative

motivations (e.g., reproduction and predatory avoidance) [75-78] further modify these foraging

decisions.

Quantitative and ethological approach to investigating decision-making

Naturalistic observations of *C. elegans* behavior are necessary to achieve a more complete cellular, molecular, and genetic understanding of how the nervous system has evolved to drive animal behavior. For example, large patches of densely seeded bacteria have primarily been used in experiments assessing *C. elegans* foraging [20, 21, 79-81]. However, wild nematodes experience a boom-and-bust environment [68] that may be more consistent with sparse, patchily distributed bacteria [28]. Thus, just as quiescence is only observed on high quality bacterial patches [82], additional behavioral states may be discovered when assessing foraging in dilute patchy environments. By conducting detailed analyses of ecologically inspired foraging behaviors in individual animals, we discovered that *C. elegans* make *accept-reject* decisions upon encounter with bacterial patches. By studying foraging in these environments containing small, dilute bacterial patches, we increased the frequency of opportunities for *accept-reject* decision-making (i.e., patch encounters and patch-leaving events are more frequent) and decreased the condition-specific bias towards *accept* decisions (i.e., animals are less likely to *accept* patches when moved between environments of differing bacterial density). We expect that further investigation into naturalistic behaviors could reveal additional insights into more complex behaviors and decision-making in *C. elegans* and other animals.

In this study, we fine-tuned an assay that enables observation of ecologically relevant decision-making in freely moving animals. We posit that future studies can leverage this assay to investigate the neuronal mechanisms underlying decision-making. The frequency of decision-making opportunities (i.e., numerous *accept-reject* decisions could be observed in the span of minutes) and consistency of patch choice decisions across hundreds of animals and a range of environmental conditions tested supports this claim. Further, we anticipate that the *accept-reject* decision occurs during a narrow time window – possibly at the time of encounter with the patch edge or shortly thereafter. Therefore, we expect that this assay could be adapted for experiments utilizing calcium imaging in freely-moving animals [17, 30, 83-87] to monitor neuronal activity as animals make these decisions.

Finally, we used simple quantitative models to dissect and validate the components underlying decision-making. Specifically, we employed a logistic regression GLM which we adapted to accommodate our uncertainty in classification of *sensing* and *exploitation* for each encounter. The GLM enabled us to conduct a sequence of hypothesis tests probing the influence of food-related

signals on the decision to *exploit*. Further, the GLM enabled us to identify the simplest model that
could explain the observed delay in *exploitation*. While even the null model predicted that, on average,
numerous *exploratory* encounters were likely to occur before the first *exploitation*, models including
satiety and prior experience significantly better predicted the extent of this delay. We demonstrated
that we could validate these model predictions in food-deprived animals, animals exploring multiple
patch densities, and animals with sensory deficiencies. We propose that this approach to investigating
behavior and decision-making is incredibly powerful for refining hypotheses and enables subsequent
investigation leveraging the diversity of cellular, molecular, and genetic tools available in *C. elegans*.
553

Abbreviations

555 Bayesian information criterion (BIC), Gaussian mixture model (GMM), generalized linear model
556 (GLM), kernel density estimate (KDE), Kullback-Leibler (KL) divergence, lysogeny broth (LB),
557 nematode growth medium (NGM), optical density at 600 nm (OD_{600}), quadratic discriminant
558 analysis (QDA)
559

Acknowledgements

561 We thank members of the Chalasani lab for comments on the manuscript and James Fitzgerald for
562 advice on modeling. This research was funded by grants from the National Science Foundation
563 (J.A.H.), UCSD Undergraduate Summer Research Award (T.C.), NIH RO1 MH096881, Dorsett
564 Brown Foundation, and Salk Innovation Grant (S.H.C.).
565

Author Contributions

Jessica Haley: Conceptualization, Methodology, Software, Validation, Formal analysis,
Investigation, Resources, Data Curation, Writing – Original Draft, Writing – Reviewing & Editing,
Visualization, Supervision, Project administration, Funding acquisition. **Tianyi Chen:** Investigation,
Data Curation, Writing – Reviewing & Editing. **Mikio Aoi:** Methodology, Formal analysis, Writing –
Reviewing & Editing, Visualization, Supervision. **Sreekanth Chalasani:** Conceptualization,
Resources, Writing – Reviewing & Editing, Supervision, Funding acquisition.
573

Declaration of Interests

575 The authors declare no competing interests.

Materials and Methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifier
bacterial strain (*Escherichia coli*)	OP50	Caenorhabditis Genetics Center	WBStrain00041969
bacterial strain (*Escherichia coli*)	OP50-GFP	Caenorhabditis Genetics Center	WBStrain00041972
organism/strain (*Caenorhabditis elegans*)	N2: wild isolate	Caenorhabditis Genetics Center	WBStrain00000001
organism/strain (*Caenorhabditis elegans*)	PR811: [osm-6(p811) V]	Caenorhabditis Genetics Center	WBStrain00030796
organism/strain (*Caenorhabditis elegans*)	TU253: [mec-4(u253) X]	Caenorhabditis Genetics Center	WBStrain00035037
software, algorithm	StreamPix 8	NorPix	RRID:SCR_015773
software, algorithm	WormLab	MBF Bioscience	RRID:SCR_017669
software, algorithm	MATLAB 2024a	Mathworks	RRID:SCR_001622
software, algorithm	Photoshop 2024	Adobe	RRID:SCR_014199
software, algorithm	Illustrator 2024	Adobe	RRID:SCR_010279
software, algorithm	Code for analysis	https://github.com/ shreklab/Haley-et-al-2024	

Data and Code Availability

All data reported in this paper will be shared by the lead contact upon request. All original code has been uploaded to GitHub (https://github.com/shreklab/Haley-et-al-2024) and are publicly available as of the date of publication. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request. Source data files contain the summarized data for all plots.

Bacterial cultures

Stock liquid cultures of the OP50 strain of *Escherichia coli* were prepared via inoculation of a single colony in sterile lysogeny broth (LB) grown overnight at room temperature. Stock liquid cultures were subsequently stored at 4°C for up to 6 weeks.

Solutions of OP50 for each experiment were prepared via a series of dilutions in LB (Figure 1 – supplement 1A). 50 mL of OP50 stock solution was centrifuged at 3000 rpm for 5 minutes. After removal of supernatant, approximately 500 to 1000 μL of saturated liquid culture remained.

The bacterial density of this saturated solution was estimated by measuring the optical density at 600

nm (OD_{600}) of a 1:50 dilution of the homogenized solution on a spectrophotometer (Molecular

Devices SpectraMax Plus 384). The saturated solution was subsequently diluted to achieve an OD_{600}

of ~10 ($\mu = 10.22$, $\sigma = 0.31$). Measurements of the number of colony forming units in the "10"

solution estimated 13.1×10^9 cells per mL on average. Additional densities (OD_{600} = {0.05, 0.1, 0.5, 1,

2, 3, 4, 5}) were prepared via dilution of the "10" solution with LB and kept on ice to prevent

bacterial growth. A "0" density solution was prepared with just LB.

For all experiments, these density solutions were seeded onto cold, low-moisture nematode

growth medium (NGM) plates (3% agar) to facilitate pipetting of small, circular, quick-drying

patches. Unless otherwise noted, seeded plates were immediately returned to 4°C after patches dried

to prevent bacterial growth. Plates were stored at 4°C for an average of 6 days before

experimentation.

For experiments using an OP50 strain expressing green fluorescent protein (OP50-GFP)

[88], cultures were prepared as above, but with the addition of 100 μg/mL of carbenicillin to the

liquid LB. All bacterial strains used in this study are listed in the Key Resources table.

Nematode cultures

Caenorhabditis elegans strains were maintained under standard conditions at 20°C on NGM

plates (1.7% agar) seeded with stock liquid culture of OP50 [16]. All experiments were performed

on young adult hermaphrodites, picked as L4 larvae the day before the experiment (μ= 24.6 hours, σ

= 3.4). Unless otherwise indicated, well-fed animals of the standard *C. elegans* strain N2 Bristol were

used for experiments. Transgenic strains were always compared to matched controls tested in parallel

on the same days. All *C. elegans* strains used in this study are listed in the Key Resources table.

Assay preparation and recording

Single-density, multi-patch assay

Several days before the experiment, condition and acclimation plates were prepared (Figure 1

– supplement 1A). For the condition plates, a large circular arena (30 mm in diameter) made of clear

transparency film (6 mil, PET) was cut using a computer-controlled cutting machine (Cricut Maker

3) and placed in the center of a 10 cm petri dish filled with 25 mL of NGM (3% agar). These arenas

function to corral animals with a low probability of escape [89]. A pipetting template made of the

same material was designed and cut with an isometric grid consisting of 19 circular holes spaced 6

mm apart from center-to-center. The pipetting template was overlaid on top of the arena and 0.5 μL

of OP50 solution (OD_{600} = {0, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10}) was pipetted into each hole. After

drying, the template was immediately removed, and the petri dish lid replaced to prevent

contamination and dehydration. For two conditions ("1 (12H)" and "1 (48H)"), plates were seeded

with OD_{600} = 1 and then left at room temperature for 12 or 24 hours, respectively, prior to storage at

4°C. For all other conditions, plates were immediately stored at 4°C after drying. Acclimation plates

were prepared by seeding NGM (3% agar) plates with one large patch of 200 μL of OD_{600} = 1

grown for 24 hours at room temperature.

Approximately 24 hours before the experiment, acclimation plates and "1 (48H)" condition

plates were transferred from 4°C to room temperature. After plates warmed to room temperature

(~1 hour), 30-60 L4 animals were picked onto the acclimation plates. Plates were then stored at

20°C until the experiment for a combined bacterial growth time of approximately 48 hours (24

hours from freshly seeded to 4°C; 24 hours from 4°C to experiment). The combined bacterial

growth time for all condition plates was approximately 1 hour (μ= 1.16 hours, σ = 0.23) unless

otherwise noted as "12" (μ= 13.63 hours, σ = 0.50) or "48" hours (μ= 49.28 hours, σ = 2.94).

On the day of the experiment, each set of condition plates was transferred from 4°C to

room temperature. After one hour, 4 young adult animals were gently transferred to an empty NGM

plate to limit the spread of bacteria from acclimation to condition plates. Animals were then quickly

transferred to a condition plate. Animals were transferred using cylindrical plugs excised from clean

3% agar [64, 89].

Condition plates were subsequently placed face down on a piece of glass suspended above

an edge-lit backlight (Advanced Illumination). Recordings were acquired using PixeLink cameras

(PL-B741F) combined with Navitar lenses (1-60135 and 1-6044) and Streampix 8 software. Behavior

was recorded for one hour at 1024x1024 pixels and 3 frames per second (fps) with spatial resolution

of ~33 pixels per mm (μ= 32.68 pixels/mm, σ = 1.70).

The assay was repeated over numerous days with every condition assayed on each day when

possible. The order of seeding assay plates and recording behavior was randomized for each day to

compensate for the potential effects of time on animal age and bacterial density. Further, the average

temperature (μ= 21.99°C, σ = 0.95) and humidity (μ= 51.39%, σ = 11.67) during each experiment

were recorded. As a result of randomization, animal age, bacterial growth time, temperature, and

humidity did not significantly vary between density conditions.

Large, single-patch assay

Plates were prepared as described for the single-density, multi-patch assay with the exception

that only one large patch was created by pipetting 20 μL of OP50 solution (OD_{600} = {0, 0.05, 0.1,

0.5, 1, 2, 3, 4, 5, 10}) into the center of the 30 mm diameter arena.

Small, single-patch assay

Plates were prepared as described for the single-density, multi-patch assay with the exception

that only one small patch was created by pipetting 0.5 μL of OP50 solution (OD_{600} = {0, 0.5, 1, 5,

10}) into the center of a 9 mm diameter arena. Given the smaller arena size, only one young adult

animal was transferred into the arena. Behavior was recorded for one hour at higher spatial (μ =

104.76 pixels/mm, σ = 17.6) and temporal resolution (8 fps). The acquisition set-up was otherwise

unaltered except for the removal of a 0.25x Navitar lens (1-6044) from the light path.

Multi-density, multi-patch assay

Plates were prepared as described for the single-density, multi-patch assay with the exception

that varying combinations of OD_{600} = {1, 5, 10} (i.e., OD_{600} = {1, 5, 10, 1+5, 1+10, 1+5+10}) were

pipetted onto each assay plate in a patterned isometric grid of 18 0.5 μL droplets. Behavior was

recorded for two hours. To compensate for the longer duration recordings, NGM agar plates were

poured without Bacto™ peptone (BD 211677) to prevent bacterial growth. The result of this

change was that the relative density of bacterial patches was ~30% lower as compared to

comparable 0.5 μL patches in the single-density, multi-patch assay and the small, single-patch assay

(Figure 2 – supplement 2-3). Further, to prevent censorship of our data at the beginning of the

recording, we formed a small droplet of S-Complete solution [90] in the middle of the arena where

no bacterial patch was placed. Animals were transferred to this droplet using an eyelash pick

immediately prior to the assay plate being placed on the imaging set-up. The recording was started

once the droplet evaporated and animals were free to crawl about the arena. This process ensured

that recordings began prior to an animal's first patch encounter.

Bacterial patch location detection

Most bacterial densities tested in these experiments were too dilute to be visible under

normal imaging conditions. Therefore, several strategies were employed to accurately detect the

location of bacterial patches. First, a small reference dot was cut into the arena and pipetting

templates enabling consistent and traceable patch placement. Further, prior to each behavior

recording, a "contrast" video was acquired wherein a piece of dark cardstock was passed between

the light source and the condition plate (Figure 1 – supplement 1B-D, Video 2). This produced an

effect where diffraction of light through the patches provided enough contrast to view the patches.

Binary masks of the circular arena and patches were extracted from these videos in MATLAB using

the Image Processing Toolbox and further refined manually in Photoshop (Adobe, 2024). Arena

masks were used to calculate the scale (pixels/mm) of each image. In early experiments, the

"contrast" video was acquired prior to worms being added to the condition plate, which resulted in

displacement of the arena within the camera's field-of-view. Image registration using MATLAB's

Computer Vision Toolbox was performed to accurately map the patches detected in the "contrast"

video onto the behavioral recording.

Bacterial patch density estimation

As described above (see Assay preparation and recording), we varied the relative density of

bacterial solutions by diluting OP50 in LB and growing patches for different lengths of time. As a

result of these procedures, the relative bacterial density throughout the assay was not known and

needed to be estimated. To determine the relative density of these bacterial patches, we seeded

plates with fluorescently labeled OP50-GFP under identical conditions as in our assays (Figure 2 –

supplement 1A). We imaged these bacterial patches for several hours using a Zeiss Axio Zoom.V16.

Images were analyzed using the Image Processing Toolbox in MATLAB. After correcting for

inconsistent illumination across the field-of-view and normalizing images using matched controls

(Figure 2 – supplement 1B-C), we detected the location of each bacterial patch and extracted a

fluorescence intensity profile along each patch's radius (Figure 2 – supplement 1D). Consistent with

previous studies, we found that even in extremely dilute conditions, bacterial density was greater at

the patch edge where actively proliferating bacteria are concentrated [43, 91]. Therefore, we detected

the patch edge and computed the peak amplitude of each bacterial patch at all time points assayed

(Figure 2 – supplement 1E). We then performed linear regression on the values for each condition

to create models of peak amplitude as a function of time. For low density conditions (i.e., $OD_{600} =$

$\{0.05, 0.1\}$) of small patches, fluorescent signals could not be detected. We therefore estimated these

functions from a multinomial regression of values taken from $OD_{600} = \{0.5, 1, 2\}$. Through this

process, we used the total time that bacteria were allowed to grow at room temperature for each

condition plate to estimate the peak amplitude at each time point (Figure 2 – supplement 3). Finally,

to minimize confusion about density conditions between experiments, we computed values for

relative density by linearly scaling estimated peak amplitude values so that 0.5 μL patches seeded

with $OD_{600} = 10$ and grown for 1 hour at room temperature would have relative density equal to 10.

We labeled each condition using the relative density rounded to one significant digit.

Behavioral tracking

WormLab® (MBF Bioscience) was used for tracking animal behavior (Figure 1 – supplement

1E-G, Video 3). For each video, we adjusted thresholds for pixel intensity and worm dimensions to

enable automatic tracking. WormLab then fit the worm's body at every frame and stitched together a

track of each worm across frames. Subsequent manual corrections were required for instances

where the software created new worm tracks. This occurred most frequently when animals: 1)

encountered the arena border, 2) collided with each other, or 3) overlapped with bubbles or dust

particles embedded in the agar. To correct for these discontinuities, we manually "stitched" worm

tracks together enabling us to keep track of each animal's location throughout the duration of the

experiment. We exported the location of each animal's midpoint for further analysis in MATLAB

(Mathworks, 2024a). For small, single-patch assays where higher spatial resolution enabled reliable

distinction between the head and tail of the animal, we manually confirmed the head-tail and

exported 25 points along the animal's midline (Figure 1 – supplement 2A-C).

Animals were excluded from analyses when 1) the animal's location could only be tracked for

less than 75% of the video due to the animal escaping the arena and when 2) the location of

bacterial patches could not be reliably assessed due to missing or low-quality "contrast" videos (see

Bacterial patch location detection).

Patch encounter detection

As the head position of animals in our experiments was difficult to track due to low spatial

resolution and limited automated detection of head position using WormLab, we instead tracked the

midpoint position of the worm's body and estimate patch encounters using a set of measured

criteria. To do this, we tracked the behavior of worms in higher resolution (~105 pixels/mm

compared to ~33 pixels/mm and 8 fps compared to 3 fps), single patch assays (Figure 1 –

supplement 2A-C). We found that when the head of the animal was in contact (i.e., within 1 pixel)

with the patch edge (see Bacterial patch location detection), the animal's midpoint was on average

0.46024 mm away (Figure 1 – supplement 2D). We therefore defined a patch encounter as the time

when an animal's midpoint came within 0.46024 mm of the patch edge. However, this definition led to two types of errors: 1) putative entry events where the animal merely passed by the patch and 2) putative leaving events where the animal did not fully leave the patch.

To remove putative entry events where the animal nearly passed by the patch, we defined an additional distance threshold based on the distance between the animal's midpoint and the patch edge for all time points when the head was within the patch (Figure 1 – supplement 2E). The midpoint was almost always (99% of time) within 0.28758 mm of the patch edge. Therefore, we removed patch encounters where the midpoint never got closer than 0.28758 mm from the patch edge. Although this criterion removes most "near miss" events, some events remain where the animal approached the patch, but its head never entered. Further decreasing this threshold (e.g., midpoint must be on patch) would result in real events being excluded due to the animal taking on specific postures where the midpoint remains outside the patch while feeding within the patch. Therefore, we chose to be more liberal (i.e., more false positive patch encounters than false negatives) in our detection of patch encounters. Subsequent analysis of these putative patch encounters removed the majority of remaining false positives (see Patch encounter classification as sensing or non-sensing).

To address putative leaving events and avoid splitting a single patch encounter into multiple encounters, we considered that a lawn leaving event is most frequently defined as an event where all body parts have left the patch [31]. When tracking only the midpoint, it is not possible to ensure that the worm's entire body has exited the patch. Using the criterion that an animal exits a patch when it's midpoint exceeds 0.46024 mm from the patch edge accurately predicts exit events most of the time, but under certain scenarios (e.g., animals maintain an outstretched feeding posture [64]) false leaving events were detected. To exclude these leaving events, we analyzed the variability in the distance from the midpoint to the patch edge during on- and off-patch events (Figure 1 – supplement 2F). We found that on-patch events had low distance variability while off-patch events displayed a bimodal distribution, with some events having high variability as expected, and some events having low variability. We fit a two-component Gaussian mixture model to the off-patch distance variability and found that low variability leaving events could be reliably excluded when the standard deviation of off patch distances was less than 0.13259. We subsequently combined putative patch encounters where variability was below this threshold, resulting in fewer overall encounters.

Patch encounter classification as exploration or exploitation

To classify patch encounters as *exploration* or *exploitation* events, patch encounters were

detected (see Patch encounter detection) and two features were computed: 1) duration of patch

encounter and 2) average on-patch velocity. Both the duration and velocity features displayed a

bimodal distribution with two visible clusters: one with short duration encounters and high on-patch

velocity and one with long duration encounters and low on-patch velocity (Figure 2 – supplement

6A). To classify patch encounters into one of these clusters, we used a two-component Gaussian

mixture model (GMM) to estimate $p(y_k = 1|\mathbf{z}_k)$ where for each encounter k, $\mathbf{z}_k$ are the log

transforms of the duration and average velocity of the patch encounter, $y_k = 1$ indicates an

exploitation, and $y_k = 0$ indicates an *exploration*. We optimized the GMM across cross-validated

replicates with respect to the regularization value α to minimize the posterior variance given by

$$\sum_k p(y_k = 1|\mathbf{z}_k)\, p(y_k = 0|\mathbf{z}_k)$$

We found that $\alpha = 0.025$ best separated the *exploration* and *exploitation* clusters (Figure 2 –

supplement 6B). The posterior probability of *exploitation* $p(y_k = 1|\mathbf{z}_k)$ was subsequently estimated

for all encounters (Figure 2 – supplement 6C-D).

Patch encounter classification as sensing or non-sensing

To classify patch encounters as *sensing* or *non-sensing* events, we computed animals' 1)

deceleration upon encounter with the patch edge, 2) minimum velocity during the patch encounter,

and 3) maximum change in velocity between the peak velocity immediately prior to the start of the

patch encounter and the minimum velocity during the encounter (Figure 2 – supplement 7A).

Deceleration was defined as the slope of the line fit on an animal's velocity between -1.5 seconds

before and 6.5 seconds after the start of an encounter. Minimum velocity was defined as the

absolute minimum velocity observed during the duration of the patch encounter. The maximum

change in velocity was computed by subtracting the minimum velocity from the peak velocity within

10 seconds of the patch encounter. The combination of these three metrics reveals two non-

Gaussian clusters representing encounters where animals sensed the patch (i.e., slow minimum

velocity, large changes in velocity) or did not appear to detect it (i.e., maintained fast minimum

velocity, with little to no change). We estimated the probability of *sensing* $p(v_k = 1|\mathbf{w}_k)$ where for

each encounter k, $\mathbf{w}_k = (s_k, t_k, u_k)$ are the three velocity-related features (i.e., minimum velocity,

deceleration, and maximum change in velocity, respectively), $v_k = 1$ indicates an encounter that was

sensed, and $v_k = 0$ indicates an encounter which did not appear to be *sensed*. Using a semi-supervised

approach to quadratic discriminant analysis (QDA), we labeled a subset of all encounters and

iteratively estimated labels on the remaining unlabeled data (Figure 2 – supplement 7C-E, Video 4).

All encounters with bacteria-free patches (i.e., LB only, relative density 0) were labeled as true

negatives $v_k = 0$. Across 1000 replicates, we probabilistically included *exploitation* encounters $y_k = 1$

estimated from the distribution $y_k | z_k \sim \text{Bern}\big(p(y_k = 1 | z_k)\big)$ as true positives $v_k = 1$. Using this

approach, the conditional probability of *sensing* $p(v_k = 1 | w_k)$ was iteratively estimated for all

encounters using QDA and averaged across replicates.

Although we found that the two clusters were successfully discriminated by this semi-

supervised QDA approach, a small subset of encounters (252 of 20109 encounters) could not be

reliably classified in this manner as the onset of the patch encounter was not observed. This type of

data censoring occurred when the behavioral recording was started after the animal had already

entered the patch. Although we could compute deceleration-related metrics for these encounters

with the assumption that the animal entered the patch at the first frame, these measurements are

censored with a bias towards lower magnitudes of deceleration. Therefore, to better predict the

conditional probabilities of *sensing* for these 252 encounters, we assumed that the minimum velocity

on-patch s_k was a reliable metric and marginalized the conditional probabilities over the other two

metrics (t_k and u_k) as defined by

$$p(v_k = 1 | s_k) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p(v_k = 1 \mid s_k, t_k, u_k)\, p(t_k, u_k \mid s_k)\ dt\ du$$

We numerically integrated using an adaptive quadrature method over the product of the QDA

estimated conditional probability distribution and the kernel density estimate of the joint probability

distribution. This procedure resulted in a slight increase ($\mu = +0.0615$, $\sigma = 0.1832$) in our estimation

of the probability of *sensing* (Figure 2 – supplement 7H).

Altogether, this classification approach resulted in low false positive and false negative rates.

Specifically, 3.35% of the time encounters with bacteria-free patches were incorrectly identified as

sensing while 2.92% of the time *exploitatory* encounters were incorrectly identified as *non-sensing*.

Further, this classification removed many of the remaining near miss patch encounters in which

animals came close to but did not truly enter the patch (see Patch encounter detection). For all

analyses, we excluded near miss encounters where the animal's midpoint never entered the patch and

the probability of *sensing* was less than 5% (i.e., $p(v_k = 1 | w_k) < 0.05$).

Models of exploitation probability

846 We consider that the probability of *exploiting* a patch upon any given encounter can be

847 described by a logistic function:

$$p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k) = \frac{1}{1 + e^{-\boldsymbol{\beta} \cdot \boldsymbol{x}_k}}$$

where $p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)$ represents the conditional probability that an animal exploited during

patch encounter k; $\boldsymbol{x}_k \in R^n$ is a vector of covariates; and $\boldsymbol{\beta} \in R^n$ is a vector of weights that

describes how much each covariate influences the animal's choice. In models compared here, $\boldsymbol{x}_k$

includes a combination of a constant element as well as covariates that may empirically influence the

decision to exploit: 1) the log-transformed relative density of the encountered patch k (ρ_k), 2) the

duration of time spent off food since departing the last *exploited* patch (τ_s), 3) the log-transformed

relative density of the patch encountered immediately before encounter k (ρ_h), and 4) the log-

transformed relative density of the last *exploited* patch (ρ_e).

The observations of *exploitation* and *sensation* for our data are derived predictions from

classification models (see Patch encounter classification as exploration or exploitation and Patch

encounter classification as sensing or non-sensing). As such, the labels for both whether a patch

encounter was *sensed* by the animal $p(v_k = 1|\boldsymbol{w}_k)$ and whether the animal *exploited* the patch

$p(y_k = 1|\boldsymbol{z}_k)$, are both reported as probabilities rather than direct measurements.

To account for uncertainty in *sensation*, we produced 100 sets of "encounter sampled"

observations wherein we probabilistically included *sensed* encounters $v_k = 1$ estimated from the

distribution $v_k|\boldsymbol{w}_k \sim \text{Bern}(p(v_k = 1|\boldsymbol{w}_k))$. In doing so, we assume that only patch encounters

recognized by the animal guide the decision to *exploit* (Figure 4B). After removing *non-sensing*

encounters where $v_k = 0$, we defined observations of all covariates (i.e., ρ_k, τ_s, ρ_h, and ρ_e) using

only *sensed* encounters.

To account for uncertainty in *exploitation* we modified the standard maximum likelihood. To

understand our approach, consider that standard logistic regression models binary observations as

realizations of the Bernoulli distribution $y_k|\boldsymbol{x}_k \sim \text{Bern}(p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k))$ and that the likelihood

for this model is given by

$$\prod_k p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)^{y_k} \left(1 - p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)\right)^{1-y_k}$$

However, we do not have direct observations of y_k and cannot evaluate the logistic likelihood

directly. Rather, we have posterior probabilities of *exploitation* determined by a classifier, denoted by

$p(y_k = 1|\mathbf{z}_k)$, where $\mathbf{z}_k$ are a set of velocity-related features that are distinct from $\mathbf{x}_k$ (see Patch

encounter classification as exploration or exploitation). Thus, rather than maximizing the Bernoulli

likelihood, we accommodate our uncertainty about y_k by learning the regression parameters that

minimize the Kullback-Leibler (KL) divergence between $p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)$ and the reference probability

distribution $p(y_k|\mathbf{z}_k)$ as given by

$$
\begin{aligned}
\mathcal{D}_{KL}[p(y_k|\mathbf{z}_k)||p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)] &\equiv \sum_{y_k} p(y_k|\mathbf{z}_k) \log\left[\frac{p(y_k|\mathbf{z}_k)}{p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)}\right] \\
&= C_{\boldsymbol{\beta}} - \sum_{y_k} p(y_k|\mathbf{z}_k) \log[p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)] \\
&= C_{\boldsymbol{\beta}} - \sum_{y_k} p(y_k|\mathbf{z}_k) \log\left[p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)^{y_k} \left(1 - p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)\right)^{1-y_k}\right] \\
&= C_{\boldsymbol{\beta}} - p(y_k = 1|\mathbf{z}_k) \log[p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)] \\
&\quad - (1 - p(y_k = 1|\mathbf{z}_k)) \log[1 - p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)] \\
&= C_{\boldsymbol{\beta}} \\
&\quad - \log\left[p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)^{p(y_k=1|\mathbf{z}_k)} \left(1 - p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)\right)^{1-p(y_k=1|\mathbf{z}_k)}\right]
\end{aligned}
$$

where $C_{\boldsymbol{\beta}}$ is a term that is constant with respect to $\boldsymbol{\beta}$, and the second term is equivalent to the

logarithm of the logistic likelihood with $p(y_k = 1|\mathbf{z}_k)$ replacing y_k. Put plainly, minimizing the sum

of the KL divergence between our classifier probabilities and logistic regression probabilities over all

encounters k is equivalent to maximizing the logistic likelihood with our classifier-based *exploitation*

probabilities $p(y_k = 1|\mathbf{z}_k)$ as observations instead of direct measurements of *exploitation* y_k as

given by

$$
\sum_k \log\left[p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)^{p(y_k=1|\mathbf{z}_k)} \left(1 - p(y_k|\boldsymbol{\beta} \cdot \mathbf{x}_k)\right)^{1-p(y_k=1|\mathbf{z}_k)}\right]
$$

To train our model, we therefore fit a standard logistic regression GLM where we provided sets of

observations of covariates $\mathbf{x}_k$ and response variables $p(y_k = 1|\mathbf{z}_k)$ for *sensed* encounters.

Exploitation of single-density, multi-patch environments

As described above, we used an "encounter sampling" protocol to generate 100 replicate

observations of only *sensed* encounters by removing all encounters that were not *sensed* (i.e., $v_k = 0$)

as randomly estimated from the distribution $v_k | \boldsymbol{w}_k \sim \text{Bern}\big(p(v_k = 1 | \boldsymbol{w}_k)\big)$ (Figure 4B). For

example, we can consider an animal foraging in this single-density, multi-patch environment.

Although we detected 10 total encounters $K = k_1, k_2, \ldots, k_{10}$, several of these encounters had low

probabilities of sensing $p(v_k = 1 | \boldsymbol{w}_k)$ (Figure 4B). We use our "encounter sampling" protocol to

simulate *sensation* of the patch during each of these 10 encounters and find that $v_k = 0$ for

k_1, k_3, k_6 and k_8. By removing these 4 non-sensed encounters, we generate a new observation

comprised of the 6 *sensed* encounters $K_1^* = k_2, k_4, k_5, k_7, k_9, k_{10} = k_{1*}, k_{2*}, \ldots, k_{6*}$. Using only

the data related to this new sequence of encounters K_1^*, we generated a vector of covariates (i.e., ρ_k,

τ_s, ρ_h, and ρ_e). Importantly, this procedure considers time spent during the *non-sensed* encounters

k_1, k_3, k_6 and k_7 the same as time spent searching off patch. We repeated this process 100 times to

generate new observations of encounter sequences (i.e., $K_1^*, K_2^*, \ldots, K_{100}^*$). In doing so, we reduced

the number of encounters from 6,560 total encounters (as defined in Patch encounter detection) to

2,604-2,698 ($\mu = 2659.8$, $\sigma = 15.9$) *sensed* encounters for each of the 100 replicates.

Additionally, we used a "worm sampling" protocol to generate 500 hierarchically

bootstrapped replicates [92]. Specifically, we took our vector of 443 animals $A = a_1, a_2, \ldots, a_{443}$

and resampled with replacement (e.g., $A_1^* = a_{317}, a_{88}, \ldots, a_{130} = a_{1*}, a_{2*}, \ldots, a_{443*}$) to generate

500 new samples $A_1^*, A_2^*, \ldots, A_{500}^*$. When combined, the "encounter sampling" and "worm

sampling" protocols created 50,000 unique replicates of observations of the covariates $\boldsymbol{x}_k$ and

response variables $p(y_k = 1 | \boldsymbol{z}_k)$ which were used to estimate the coefficients $\boldsymbol{\beta}$ (Figure 4C). A null

distribution of coefficients $\boldsymbol{\beta}^*$ was estimated by shuffling the response variable vector (Figure 4 –

supplement 1B). A two-tailed, one-sample bootstrap hypothesis test was used to assess whether our

covariate estimates were significantly greater than or less than 0 (i.e., $\boldsymbol{p} = 2 \times \min[P(\boldsymbol{\beta} \leq$

$\boldsymbol{0}), P(\boldsymbol{\beta} \geq \boldsymbol{0})]$).

Exploitation of food-deprived animals in single-density, multi-patch environments

To validate model predictions related to the satiety signal covariate τ_s, a novel data set of

well-fed and 3-hour food-deprived animals foraging in single-density, multi-patch environments of

relative density 5 was generated (Figure 4 – supplement 3A-B). The "encounter sampling" protocol

described above was used to generate 100 replicates of observations of the covariates x_k and

response variables $p(y_k = 1|z_k)$ for this data set. As a result of "encounter sampling", we reduced

the number of encounters from 42 total encounters to 30-37 ($\mu = 34.6$, $\sigma = 1.3$) *sensed* encounters

for food-deprived animals and 235 total encounters to 94-107 ($\mu = 99.8$, $\sigma = 2.1$) *sensed* encounters

for well-fed animals. Coefficient values $\boldsymbol{\beta}$ were re-estimated in the absence of the satiety signal τ_s

(Figure 4 – supplement 3C) using the original single-density, multi-patch data set. Using the mean

coefficient values $\boldsymbol{\beta}$ for both models (with and without satiety), we predicted the conditional

probabilities of exploiting $p(y_k = 1|\boldsymbol{\beta} \cdot x_k)$ for the new data set for well-fed and food-deprived

animals. We subsequently simulated a series of *exploitation* events from a Bernoulli distribution using

the estimated (i.e., $y_k|x_k \sim \text{Bern}(p(y_k = 1|\boldsymbol{\beta} \cdot x_k))$) and observed (i.e.,

$y_k|z_k \sim \text{Bern}(p(y_k = 1|z_k))$) *exploitation* probabilities. These simulated *exploitations* were used to

generate distributions of the probability of an *exploitation* occurring for the first time as a function

of the number of encounters (Figure 4 – supplement 3D).

Exploitation of multi-density, multi-patch environments

To validate model predictions related to the prior experience covariates ρ_h and ρ_e, a novel

data set of animals foraging on multi-density, multi-patch environments was generated (Figure 4E,

Figure 4 - supplement 4A-C). The "encounter sampling" protocol described above was used to

generate 100 replicates of observations of the covariates x_k and response variables $p(y_k = 1|z_k)$

for this data set. As a result of "encounter sampling", we reduced the number of encounters from

4,296 total encounters (350-984 total encounters per condition) to 1,702-1,745 ($\mu = 1,724.6$, $\sigma = 9.9$)

sensed encounters. Coefficient values $\boldsymbol{\beta}$ were re-estimated in the absence of the recently encountered

and *exploited* patch density terms ρ_h and ρ_e (Figure 4 – supplement 4D) using the original single-

density, multi-patch data set. Using the mean coefficient values $\boldsymbol{\beta}$ for both models (with and without

prior experience), we predicted the probabilities of *exploiting* $p(y_k = 1|\boldsymbol{\beta} \cdot x_k)$ for the new data set.

To generate the heat maps of predicted behavior (Figure 4F), we varied ρ_k and ρ_h as well as ρ_k and

ρ_e across a range of values and set the remaining covariate terms to the average values observed in

the data set. To generate the heat maps of observed behavior (Figure 4F), we averaged the observed

probabilities of *exploiting* $p(y_k = 1|z_k)$ for each pairing of ρ_k and ρ_h as well as ρ_k and ρ_e and

linearly interpolated values between these 9 points.

Exploitation of sensory mutants in single-density, multi-patch environments

To test the utility of our model in identifying covariate-specific phenotypes in animals with

varied genotypes, a novel data set of wild-type animals and sensory mutants foraging on single-

density, multi-patch environments was generated (Figure 4 - supplement 5A-B). The "encounter

sampling" and "worm sampling" protocols described above were used to generate 50,000 replicates

of observations of the covariates $\boldsymbol{x}_k$ and response variables $p(y_k = 1|\boldsymbol{z}_k)$ for this data set. As a

result of "encounter sampling", we reduced the number of encounters from 1,352 total encounters

(27-193 total encounters per condition) to 862-894 ($\mu = 879.4$, $\sigma = 7.4$) *sensed* encounters. As the

number of encounters were significantly less in these data sets as compared to the original single-

density, multi-patch data set, we used ridge regression to regularize the magnitude of the covariates.

We optimized each model across cross-validated replicates by varying the regularization value λ. We

found that $\lambda_{\text{N2}} = 0.0175$, $\lambda_{mec-4} = 0.0223$, and $\lambda_{osm-6} = 0.0614$ maximized the mean log-

likelihood of each model (Figure 4 – supplement 5E). Coefficient values $\boldsymbol{\beta}$ were estimated for each

of the three strains tested using these values (Figure 4 – supplement 5D). To assess whether the

magnitude of coefficients significantly differed between wild-type and mutant models, we conducted

a two-sample test of the mean of differences as given by

$$Z = \frac{\mu_{\text{mutant}} - \mu_{\text{N2}}}{\sqrt{\sigma_{\text{mutant}}^2 + \sigma_{\text{N2}}^2}}$$

966   where μ and σ are the mean and standard deviation, respectively, of the distribution of a covariate

967   across replicates. Statistical significance was calculated as a left-tailed test $\phi(Z)$ for the density

968   coefficients (β_k, β_h, and β_e) and a two-tailed test $\phi(-|Z|)$ for the other coefficients (β_0 and β_s)

969   where ϕ is the standard normal cumulative distribution function.

References

1. Lee D (2013). Decision making: From neuroscience to psychiatry. *Neuron* **78** (2): 233-248. https://doi.org/10.1016/j.neuron.2013.04.008.

2. Budaev S, Jørgensen C, Mangel M, Sigrunn E, and Giske J (2019). Decision-making from the animal perspective: Bridging ecology and subjective cognition. *Frontiers in Ecology and Evolution* **7**, 164. https://doi.org/10.3389/fevo.2019.00164.

3. Mobbs D, Trimmer PC, Blumstein DT, and Dayan P (2018). Foraging for foundations in decision neuroscience: Insights from ethology. *Nature Reviews Neuroscience* **19** (7): 419-427. https://doi.org/10.1038/s41583-018-0010-7.

4. Gold JI, and Shadlen MN (2007). The neural basis of decision making. *Annual Review of Neuroscience* **30**: 535-574. https://doi.org/10.1146/annurev.neuro.29.051605.113038.

5. Stephens DW (2008). Decision ecology: Foraging and the ecology of animal decision making. *Cognitive, Affective, & Behavioral Neuroscience* **8** (4): 475-484. https://doi.org/10.3758/CABN.8.4.475.

6. Hall-McMaster S, and Luyckx F (2019). Revisiting foraging approaches in neuroscience. *Cognitive, Affective, & Behavioral Neuroscience* **19**: 225-230. https://doi.org/10.3758/s13415-018-00682-z.

7. Schoener TW (1971). Theory of feeding strategies. *Annual Review of Ecology and Systematics* **2**: 369-404. https://doi.org/10.1146/annurev.es.02.110171.002101.

8. Pyke GH, Pulliam HR, and Charnov EL (1977). Optimal foraging: A selective review of theory and tests. *The Quarterly Review of Biology* **52** (2): 137-154. https://doi.org/10.1086/409852.

9. Stephens DW, and Krebs JR. (1986). Foraging Theory (Princeton University Press).

10. Stephens DW, Brown JS, and Ydenberg RC. (2007). Foraging: Behavior and Ecology (University of Chicago Press). 10.7208/9780226772653.

11. Nonacs P (2000). State dependent behavior and the Marginal Value Theorem. *Behavioral Ecology* **12** (1): 71-83. https://doi.org/10.1093/oxfordjournals.beheco.a000381.

12. Hills TT (2010). Animal foraging and the evolution of goal-directed cognition. *Cognitive Science* **30** (1): 3-41. https://doi.org/10.1207/s15516709cog0000_50.

13. Calhoun AJ, and Hayden BY (2015). The foraging brain. *Current Opinion in Behavioral Sciences* **5**: 24-31. https://doi.org/10.1016/j.cobeha.2015.07.003.

14. Charnov EL (1976). Optimal foraging, the marginal value theorem. *Theoretical Population Biology* **9** (2): 129-136. https://doi.org/10.1016/0040-5809(76)90040-x.

15. Haley JA, and Chalasani SH (2024). *C. elegans* foraging as a model for understanding the neuronal
basis of decision-making. *Cellular and Molecular Life Sciences* **81**, 252.
https://doi.org/10.1007/s00018-024-05223-1.

16. Brenner S (1974). The genetics of *Caenorhabditis elegans*. *Genetics* **77** (1): 71-94.
https://doi.org/10.1093/genetics/77.1.71.

17. Faumont S, and Lockery SR (2006). The awake behaving worm: Simultaneous imaging of
neuronal activity and behavior in intact animals at millimeter scale. *Journal of Neurophysiolovy*
95: 1976-1981. https://doi.org/10.1152/jn.01050.2005.

18. Boulin T, and Hobert O (2012). From genes to function: The *C. elegans* genetic toolbox. *Wiley*
Interdisciplinary Review Developmental Biology **1** (1): 114-137. https://doi.org/10.1002/wdev.1.

19. Samuel BS, Rowedder H, Braendle C, Félix M-A, and Ruvkun G (2016). *Caenorhabditis elegans*
responses to bacteria from its natural habitats. *PNAS* **113** (27): E3941-E3949.
https://doi.org/10.1073/pnas.1607183113.

20. White JG, Southgate E, Thomson JN, and Brenner S (1986). The structure of the nervous
system of the nematode *Caenorhabditis elegans*. *Philisophical Transactions of the Royal Society B* **314**
(1165): 1-340. https://doi.org/10.1098/rstb.1986.0056.

21. de Bono M, and Maricq AV (2005). Neuronal substrates of complex behaviors in *C. elegans*.
Annual Review of Neuroscience **28**: 451-501.
https://doi.org/10.1146/annurev.neuro.27.070203.144259.

22. Colbert HA, and Bargmann CI (1995). Odorant-specific adaptation pathways generate olfactory
plasticity in *C. elegans*. *Neuron* **14** (4): 803-812. https://doi.org/10.1016/0896-6273(95)90224-
4.

23. Zhang Y, Lu H, and Bargmann CI (2005). Pathogenic bacteria induce aversive olfactory learning
in *Caenorhabditis elegans*. *Nature* **438**: 179-184. https://doi.org/10.1038/nature04216.

24. Ardiel EL, and Rankin CH (2010). An elegant mind: Learning and memory in *Caenorhabditis*
elegans. *Learning & Memory* **17**: 191-201. https://doi.org/10.1101/lm.960510.

25. Calhoun AJ, Tong A, Sharpee TO, and Chalasani SH (2015). Neural mechanisms for evaluating
environmental variability in *Caenorhabditis elegans*. *Neuron* **86**: 428-441.
https://doi.org/10.1016/j.neuron.2015.03.026.

26. Bendesky A, Tsunozaki M, Rockman MV, Kruglyak L, and Bargmann CI (2011). Catecholamine
receptor polymorphisms affect decision-making in *C. elegans*. *Nature* **472** (7343): 313-318.
https://doi.org/10.1038/nature09821.

27. Faumont S, Lindsay TH, and Lockery SR (2012). Neuronal microcircuits for decision making in
C. elegans. *Current Opinion in Neurobiology* **22** (4): 580-591.
https://doi.org/10.1016/j.conb.2012.05.005.

28. Iwanir S, Brown AS, Nagy S, Najjar D, Kazakov A, Lee KS, Zaslaver A, Levine E, and Biron D
(2016). Serotonin promotes exploitation in complex environments by accelerating decision-
making. *BMC Biology* **14** (9). https://doi.org/10.1186/s12915-016-0232-y.

29. Tanimoto Y, Yamazoe-Umemoto A, Fujita K, Kawazoe Y, Miyanishi Y, Yamazaki SJ, Fei X,
Busch KE, Gengyo-Ando K, Nakai J, Iino Y, Iwasaki Y, Hashimoto K, and Kimura KD
(2017). Calcium dynamics regulating the timing of decision-making in *C. elegans*. *eLife* **6**:
e21629. https://doi.org/10.7554/eLife.21629.

30. Ji N, Madan GK, Fabre GI, Dayan A, Baker CM, Kramer TS, Nwabudike I, and Flavell SW
(2021). A neural circuit for flexible control of persistent behavioral states. *eLife* **10**: e62889.
https://doi.org/10.7554/eLife.62889.

31. Scheer E, and Bargmann CI (2023). Sensory neurons couple arousal and foraging decisions in *C.
elegans*. *eLife* **12**: RP88657. https://doi.org/10.7554/eLife.88657.1.

32. Petersen C, Dirksen P, and Schulenburg H (2015). Why we need more ecology for genetic
models such as *C. elegans*. *Trends in Genetics* **31** (3): 120-127.
https://doi.org/10.1016/j.tig.2014.12.001.

33. Volgin AD, Yakovlev OV, Demin KA, de Abreu MS, Rosemberg DB, Meshalkina DA,
Alekseeva PA, Friend AJ, Amstislavskaya TG, and Kalueff AV (2018). Understanding the
role of environmental enrichment in zebrafish neurobehavioral models. *Zebrafish* **15** (5): 425-
432. https://doi.org/10.1089/zeb.2018.1592.

34. Kempermann G (2019). Environmental enrichment, new neurons and the neurobiology of
individuality. *Nature Reviews Neuroscience* **20** (4): 235-245. https://doi.org/10.1038/s41583-
019-0120-x.

35. Guisnet A, Maitra M, Pradhan S, and Hendricks M (2021). A three-dimensional habitat for *C.
elegans* environmental enrichment. *PLoS One* **16** (1): e0245139.
https://doi.org/10.1371/journal.pone.0245139.

36. Shtonda BB, and Avery L (2006). Dietary choice behavior in *Caenorhabditis elegans*. *Journal of
Experimental Biology* **209** (1): 89-102. https://doi.org/10.1242/jeb.01955.

37. Hills T, Brockie PJ, and Maricq AV (2004). Dopamine and glutamate control area-restricted
search behavior in *Caenorhabditis elegans*. *Journal of Neuroscience* **24** (5): 1217-1225.
https://doi.org/10.1523/JNEUROSCI.1569-03.2004.

38. Calhoun AJ, Chalasani SH, and Sharpee TO (2014). Maximally informative foraging by
Caenorhabditis elegans. *eLife* **3**: e04220. https://doi.org/10.7554/eLife.04220.001.

39. Gray JM, Hill JJ, and Bargmann CI (2005). A circuit for navigation in *Caenorhabditis elegans*. *PNAS*
102 (9): 3184-3191. https://doi.org/10.1073/pnas.0409009101.

40. Milward K, Busch KE, Murphy RJ, de Bono M, and Olofsson B (2011). Neuronal and molecular
substrates for optimal foraging in *Caenorhabditis elegans*. *PNAS* **108** (51): 20672-20677.
https://doi.org/10.1073/pnas.1106134109.

41. Olofsson B (2014). The olfactory neuron AWC promotes avoidance of normally palatable food
following chronic dietary restriction. *Journal of Experimental Biology* **217** (10): 1790-1798.
https://doi.org/10.1242/jeb.099929.

42. Pradhan S, Quilez S, Homer K, and Hendricks M (2019). Environmental programming of adult
foraging behavior in *C. elegans. Current Biology* **29**: 2867-2879.
https://doi.org/10.1016/j.cub.2019.07.045.

43. Gloria-Soria A, and Azevedo RBR (2008). *npr-1* regulates foraging and dispersal strategies in
Caenorhabditis elegans. Current Biology **18** (21): 1694-1699.
https://doi.org/10.1016/j.cub.2008.09.043.

44. Madirolas G, Al-Asmar A, Gaouar L, Marie-Louise L, Garza-Enríquez A, Rodríguez-Rada V,
Khona M, Dal Bello M, Ratzke C, Gore J, and Pérez-Escudero A (2023). *Caenorhabditis elegans*
foraging patterns follow a simple rule of thumb. *Communications Biology* **6**, 841.
https://doi.org/10.1038/s42003-023-05220-3.

45. Fujiwara M, Sengupta P, and McIntire SL (2002). Regulation of body size and behavioral state of
C. elegans by sensory perception and the EGL-4 cGMP-dependent protein kinase. *Neuron* **36**
(6): 1091-1102. https://doi.org/10.1016/s0896-6273(02)01093-0.

46. Sawin ER, Ranganathan R, and Horvitz HR (2000). *C. elegans* locomotory rate is modulated by
the environment through a dopaminergic pathway and by experience through a serotonergic
pathway. *Neuron* **26** (3): 619-631. https://doi.org/10.1016/S0896-6273(00)81199-X.

47. Fawcett TW, McNamara JM, and Houston AI (2012). When is it adaptive to be patient? A
general framework for evaluating delayed rewards. *Behavioural Processes* **89** (2): 128-136.
https://doi.org/10.1016/j.beproc.2011.08.015.

48. Susini I, Safryghin A, HIllemann F, and Wascher CAF (2021). Delay of gratification in non-
human animals: A review of inter-specific variation in performance. *bioRxiv*.
https://doi.org/10.1101/2020.05.05.078659.

49. Constantino SM, and Daw ND (2015). Learning the opportunity cost of time in a patch-foraging
task. *Cognitive, Affective, & Behavioral Neuroscience* **15**: 837-853.
https://doi.org/10.3758/s13415-015-0350-y.

50. Carter EC, Pedersen EJ, and McCullough ME (2015). Reassessing intertemporal choice: Human
decision-making is more optimal in a foraging task than in a self-control task. *Frontiers in
Psychology* **6**, 95. https://doi.org/10.3389/fpsyg.2015.00095.

51. Hayden BY (2018). Why has evolution not selected for perfect self-control? *Philisophical
Transactions of the Royal Society B* **374** (1766). https://doi.org/10.1098/rstb.2018.0139.

52. Stephens DW, and Anderson D (2001). The adaptive value of preference for immediacy: When
shortsighted rules have farsighted consequences. *Behavioral Ecology* **12** (3): 330-339.
https://doi.org/10.1093/beheco/12.3.330.

53. Ferguson TS (1967). Sequential Decision Problems. In Mathematical Statistics: A Decision
Theoretic Approach, Z.W. Birnbaum, and E. Lukacs, eds. (Academic Press).

54. Robbins H (1970). Optimal stopping. *The American Mathematical Monthly* **77** (4): 333-343.
https://doi.org/10.2307/2316139.

55. Mischel W, and Ebbesen EB (1970). Attention in delay of gratification. *Journal of Personality and*
Social Psychology **16** (2): 329-337. https://doi.org/10.1037/h0029815.

56. Rosati AG, Stevens JR, Hare B, and Hauser MD (2007). The evolutionary origins of human
patience: Temporal preferences in chimpanzees, bonobos, and human adults. *Current Biology*
17 (19): 1663-1668. https://doi.org/10.1016/j.cub.2007.08.033.

57. Schnell AK, Boeckle M, Rivera M, Clayton NS, and Hanlon RT (2021). Cuttlefish exert self-
control in a delay of gratification task. *Proceedings of the Royal Society B: Biological Sciences* **288**,
1946. https://doi.org/10.1098/rspb.2020.3161.

58. Range F, Brucks D, and Virányi Z (2020). Dogs wait longer for better rewards than wolves in a
delay of gratification task: But why? *Animal Cognition* **23** (3): 443-453.
https://doi.org/10.1007/s10071-020-01346-7.

59. Hillemann F, Bugnyar T, Kotrschal K, and Wascher CAF (2014). Waiting for better, not for
more: Corvids respond to quality in two delay maintenance tasks. *Animal Behaviour* **90**: 1-10.
https://doi.org/10.1016/j.anbehav.2014.01.007.

60. Miller R, Frohnwieser A, Schiestl M, McCoy DE, Gray RD, Taylor AH, and Clayton NS (2020).
Delayed gratification in New Caledonian crows and young children: Influence of reward type
and visibility. *Animal Cognition* **23** (1): 71-85. https://doi.org/10.1007/s10071-019-01317-7.

61. Pujol N, Link EM, Liu LX, Kurz CL, Alloing G, Tan M-W, Ray KP, Solari R, Johnson CD, and
Ewbank JJ (2001). A reverse genetic analysis of components of the Toll signaling pathway in
Caenorhabditis elegans. *Current Biology* **11** (11): 809-821. https://doi.org/10.1016/S0960-
9822(01)00241-X.

62. Melo JA, and Ruvkun G (2012). Inactivation of conserved *C. elegans* genes engages pathogen-
and xenobiotic-associated defenses. *Cell* **149** (2): 452-466.
https://doi.org/10.1016/j.cell.2012.02.050.

63. Pradel E, Zhang Y, Pujol N, Matsuyama T, Bargmann CI, and Ewbank JJ (2007). Detection and
avoidance of a natural product from the pathogenic bacterium *Serratia marcescens* by
Caenorhabditis elegans. *PNAS* **104** (7): 2295-2300. https://doi.org/10.1073/pnas.0610281104.

64. Quach KT, and Chalasani SH (2022). Flexible reprogramming of *Pristionchus pacificus* motivation
for attacking *Caenorhabditis elegans* in predator-prey competition. *Current Biology* **32**: 1-14.
https://doi.org/10.1016/j.cub.2022.02.033.

65. Pribadi A, Rieger MA, Rosales K, Reddy KC, and Chalasani SH (2023). Dopamine signaling
regulates predator-driven changes in *Caenorhabditis elegans'* egg laying behavior. *eLife* **12**:
e83957. https://doi.org/10.7554/eLife.83957.

66. Busch KE, and Olofsson B (2012). Should I stay or should I go? *Worm* **1** (3): 182-186.
https://doi.org/10.4161/worm.20464.

67. Krebs JR, and Inman AJ (1992). Learning and foraging: Individuals, groups, and populations. *The*
American Naturalist **140**: S63-S84. https://doi.org/10.1086/285397.

68. Frézal L, and Félix M-A (2015). *C. elegans* outside the Petri dish. *eLife* **4**: e05849.
https://doi.org/10.7554/eLife.05849.

69. Matty MA, Lau HE, Haley JA, Singh A, Chakraborty A, Kono K, Reddy KC, Hansen M, and
Chalasani SH (2022). Intestine-to-neuronal signaling alters risk-taking behaviors in food-
deprived *Caenorhabditis elegans*. *PLoS Genetics* **18** (5): e1010178.
https://doi.org/10.1371/journal.pgen.1010178.

70. Ezcurra M, Tanizawa Y, Swoboda P, and Schafer WR (2011). Food sensitizes *C. elegans*
avoidance behaviors through acute dopamine signaling. *EMBO Journal* **30** (6): 1110-1122.
https://doi.org/10.1038/emboj.2011.22.

71. Anderson JL, Albergotti L, Ellebracht B, Huey RB, and Phillips PC (2011). Does
thermoregulatory behavior maximize reproductive fitness of natural isolates of *Caenorhabditis*
elegans. *BMC Evolutionary Biology* **11**, 157. https://doi.org/10.1186/1471-2148-11-157.

72. Petersen C, Dirksen P, Prahl S, Strathmann EA, and Schulenburg H (2014). The prevalence of
Caenorhabditis elegans across 1.5 years in selected North German locations: The importance of
substrate type, abiotic parameters, and *Caenorhabditis* competitors. *BMC Ecology* **6** (14), 4.
https://doi.org/10.1186/1472-6785-14-4.

73. Fraune S, and Bosch TCG (2010). Why bacteria matter in animal development and evolution.
BioEssays **32** (7): 571-580. https://doi.org/10.1002/bies.200900192.

74. Dirksen P, Assié A, Zimmerman J, Zhang F, Tietje A-M, Marsh SA, Félix M-A, Shapira M,
Kaleta C, Schulenburg H, and Samuel BS (2020). CeMbio - The *Caenorhabditis elegans*
microbiome resource. *G3 Genes|Genomes|Genetics* **10** (9): 3025-3039.
https://doi.org/10.1534/g3.120.401309.

75. Ding SS, Muhle LS, Brown AEX, Schumacher LJ, and Endres RG (2020). Comparison of
solitary and collective foraging strategies of *Caenorhabditis elegans* in patchy food distributions.
Philosophical Transactions of the Royal Society B **375**, 37520190382.
https://doi.org/10.1098/rstb.2019.0382.

76. Lipton J, Kleemann G, Ghosh R, Lints R, and Emmons SW (2004). Mate searching in
Caenorhabditis elegans: A genetic model for sex drive in a simple invertebrate. *Journal of*
Neuroscience **24** (34): 7427-7434. https://doi.org/10.1523/JNEUROSCI.1746-04.2004.

77. Barrios A, Nurrish S, and Emmons SW (2008). Sensory regulation of *C. elegans* male mate-
searching behavior. *Current Biology* **18** (23): 1865-1871.
https://doi.org/10.1016/j.cub.2008.10.050.

78. Matsuura T, Sato T, and Shingai R (2005). Interactions between *Caenorhabditis elegans* individuals
during chemotactic responses. *Zoological Science* **22** (10): 1095-1103.
https://doi.org/10.2108/zsj.22.1095.

79. Gray J, and Lissmann HW (1964). The locomotion of nematodes. *Journal of Experimental Biology*
41: 135-154. https://doi.org/10.1242/jeb.41.1.135.

80. Croll NA (1975). Behavioural analysis of nematode movement. *Advances in Parasitology* **13**: 71-122.
https://doi.org/10.1016/s0065-308x(08)60319-x.

81. Croll NA (1975). Components and patterns in the behavior of the nematode *Caenorhabditis*
elegans. *Journal of Zoology* **176** (2): 159-176. https://doi.org/10.1111/j.1469-
7998.1975.tb03191.x.

82. You Y-j, Kim J, Raizen DM, and Avery L (2008). Insulin, cGMP, TGF-beta signals regulate food
intake and quiescence in *C. elegans*: A model for satiety. *Cell Metabolism* **7** (3): 249-257.
https://doi.org/10.1016/j.cmet.2008.01.005.

83. Prevedel R, Yoon Y-G, Hoffmann M, Pak N, Wetzstein G, Kato S, Schrödel T, Raskar R,
Zimmer M, Boyden ES, and Vaziri A (2014). Simultaneous whole-animal 3D imaging of
neuronal activity using light-field microscopy. *Nature Methods* **11**: 727=730.
https://doi.org/10.1038/nmeth.2964.

84. Nguyen JP, Shipley FB, Linder AN, Plummer GS, Liu M, Setru SU, Shaevitz JW, and Leifer AM
(2015). Whole-brain calcium imaging with cellular resolution in freely behaving *Caenorhabditis*
elegans. *PNAS* **113** (8): E1074-E1081. https://doi.org/10.1073/pnas.1507110112.

85. Venkatachalam V, Ji N, Wang X, Clark C, Mitchell JK, Klein M, Tabone CJ, Florman J, Ji H,
Greenwood J, Chisholm AD, Srinivasan J, Alkema M, Zhen M, and Samuel ADT (2015).
Pan-neuronal imaging in roaming *Caenorhabditis elegans*. *PNAS* **113** (8): E1082-E1088.
https://doi.org/10.1073/pnas.1507109113.

86. Voleti V, Patel KB, Li W, Perez Campos C, Bharadwaj S, Yu H, Ford C, Casper MJ, Yan RW,
Liang W, Wen C, Kimura KD, Targoff KL, and Hillman EMC (2019). Real-time volumetric
microscopy of *in vivo* dynamics and large-scale samples with SCAPE 2.0. *Nature Methods* **16**
(10): 1054-1062. https://doi.org/10.1038/s41592-019-0579-4.

87. Hallinen KM, Dempsey R, Scholz M, Yu X, Linder AN, Randi F, Sharma AK, Shaevitz JW, and
Leifer AM (2021). Decoding locomotion from population neural activity in moving *C.*
elegans. *eLife* **10**: e66135. https://doi.org/10.7554/eLife.66135.

88. Labrousse A, Chauvet S, Couillault C, Kurz CL, and Ewbank JJ (2000). *Caenorhabditis elegans* is a
model host for *Salmonella typhimurium*. *Current Biology* **10** (23): 1543-1545.
https://doi.org/10.1016/S0960-9822(00)00833-2.

89. Quach KT, Hughes GA, and Chalasani SH (2024). Interdependence between SEB-3 and NLP-
49 peptides shifts across predator-induced defensive behavioral modes in *Caenorhabditis*
elegans. *eLife* **13**: RP98262. https://doi.org/10.7554/eLife.98262.1.

90. Sulston JE, and Brenner S (1974). The DNA of *Caenorhabditis elegans*. *Genetics* **77** (1): 95-104.
1221   https://doi.org/10.1093/genetics/77.1.95.

1222   91. Hallatschek O, Hersen P, Ramanathan S, and Nelson DR (2007). Genetic drift at expanding
1223   frontiers promotes gene segregation. *PNAS* **104** (50): 19926-19930.
1224   https://doi.org/10.1073/pnas.0710150104.

1225   92. Saravanan V, Berman GJ, and Sober SJ (2020). Application of the hierarchical bootstrap to
1226   multi-level data in neuroscience. *Neurons, Behavior, Data analysis, and Theory* **3** (5).
1227

Figures

Figure 1: *C. elegans* forage in a patchy environment with an explore-then-exploit strategy.

(A) An example animal's midbody location (colored to represent time in the experiment) as it forages in an environment bounded by a large (30 mm diameter) arena containing 19 small (~1.8 mm diameter) bacterial patches (gray). Each patch was made by pipetting ~0.5 μL of OP50 *E. coli* diluted to OD_{600} ~10 and grown at room temperature for ~1 hour. (B) Distance between the example animal's midbody position and the nearest patch edge (positive indicates inside patch) is plotted (black) for every time point. Putative encounters with a bacterial patch are indicated (gray). (C) Patch encounters (colored to represent the duration of the encounter) for 50 individuals foraging in these environments are plotted. (D) The number of total and unique patch encounters for each animal are shown. (E) Duration for each patch encounter was computed and classified as either short (0-2 minutes) or long (2-60 minutes) using a Gaussian mixture model. The distributions of all observed short and long encounters are plotted with duration binned logarithmically. (F) The observed start time of each patch encounter is shown for all short and long duration encounters. Long duration encounters occur significantly later (one-tailed Mann-Whitney U-Test). (G) The probability of residing on patch was computed for all worms across time (black) and compared to the probability of residing on patch if patch locations were semi-randomly permuted (pink). Smoothed median values are plotted with bootstrap-derived 2.5% and 97.5% quantiles shown in shaded regions. Time points where observed probabilities of residing on patch significantly exceed permuted probabilities are indicated by a black line (one-tailed Fisher's exact test with Benjamini-Hochberg correction). (H) The track of the example animal in (A) is replotted with color used to represent the animal's instantaneous velocity at each time point. (I) Velocity of the example animal over time is plotted (black) alongside patch encounters (gray) as previously identified in (B). Example encounters – one early, short duration (green) and one late, long duration (blue) – are indicated. (J,K) A 60-second time window surrounding the start of these example encounters is enlarged. (L) Velocity trajectories were aligned to patch entry for every encounter. Mean encounter-aligned (black/gray) and randomly-aligned (pink) trajectories are plotted for each animal (light) and across all animals (dark). (M) Deceleration upon encounter with the patch edge is plotted for every encounter and grouped by duration type. Deceleration was significantly lower for encounter-aligned as compared to randomly-aligned trajectories (pink) for all duration types (one-tailed Mann-Whitney U-Tests with Bonferroni Correction). (N) Mean velocity as a function of the distance from the edge of bacterial patches (computed for 50 μm bins) is shown for every animal (gray) and across all animals (black). (O) Each animal's mean velocity during time spent on and off (midpoint at least -0.46 mm from patch edge) patch is shown. Velocity on patch was significantly slower than off patch (one-tailed paired-sample *t*-test). Sample data for one animal (worm #1) are shown in (A-C,H-K). Summary data for all animals (N = 50 worms) and encounters (N = 419 total encounters) are shown in (D-G,L-O). Violin plots in (D,F,M,O) give the KDE and quartiles for each measure. Asterisks denote statistical significance (***p<0.001). See also Figure 1 – supplement 1-5 and Video 1-4.

Figure 2: The timing of the switch from explore-to-exploit is density-dependent.

(A) The relative density (as estimated by fluorescently-labeled OP50-GFP) is shown for small (~1.8 mm diameter) bacterial patches made by pipetting ~0.5 μL droplets of OP50 *E. coli* diluted in LB to a range of optical densities (OD_{600} = {0, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10} and controlling growth time at room temperature (hours = {1, 12, 48}). For (A-G,L-O), gray-scale color saturation is proportional to the relative density of each condition and corresponds to labels in (A). (B) The mean velocities of animals foraging in environments containing patches matching one of the 12 bacterial densities are plotted as a function of the distance from the edge of bacterial patches (computed for 50 μm bins). (C) Animals' average on-patch velocity is plotted as a function of the relative density of bacteria. Compared to animals foraging amongst bacteria-free patches containing only LB (relative density 0), animals foraging on bacterial patches with relative density of 0.5 or greater display significantly slower on-patch velocities (one-tailed Mann-Whitney U-Tests with Bonferroni Correction). (D) The midbody location (colored to represent time in the experiment) of example animals foraging in environments containing patches (gray) of relative density 0, 1, 5, 10, and 200 are shown. (E) The total time each animal spent on patch is plotted as a function of the relative density of bacteria. Time on patch increased monotonically with increasing bacterial density (Kendall's τ correlation, p<0.001) following a sigmoidal trend. (F) Smoothed median values of the probability of worms residing on patch over time for each density condition are plotted. Time points where observed probabilities of residing on patch either match (pink) or significantly exceed (gray) the probability of residing on patch if patch locations were semi-randomly permuted are indicated (one-tailed Fisher's exact test with Benjamini-Hochberg correction). (G) A KDE of the distribution of encounter durations is plotted for each density condition. (H) For each encounter, the average velocity of the animal during the encounter and the duration of that encounter are plotted on a double-logarithmic plot with color representing the probabilities of clustering classification as *search* (orange), *sample* (green), or *exploit* (blue). Contours showing the first, second, and third standard deviation of the GMM used to classify *explore* and *exploit* encounters are shown as shaded ellipses with saturation corresponding to standard deviation. KDEs for distributions of average on-patch velocity and encounter duration are plotted for each encounter type. (I) For each encounter, the minimum on-patch velocity and maximum change in velocity are plotted. Contours showing the separation of *sensing* and *non-sensing* encounters as estimated by semi-supervised QDA are indicated. A KDE for the distribution of the maximum change in velocity is plotted for each encounter type. (J) Features used to classify encounters as *search*, *sample*, or *exploit* are summarized. (K) KDEs of the distributions of animals' velocities are shown for all timepoints during search off and on patch as well as during sample and exploit encounters. (L) Ethograms of patch encounters (colored to represent the probability of classification as *search*, *sample*, and *exploit*) are shown for 443 individuals. (M) The average proportion of each encounter type over time is plotted. (N) Time elapsed and (O) number of encounters occurring prior to the first *exploitation* event are plotted for every animal (blue). In the event that no *exploitation* event occurred, the maximum observed time and encounters are plotted (red-orange). Both time and encounter number before *exploitation* decrease monotonically with increasing patch density (Kendall's τ correlation, p<0.001) following a sigmoidal trend. Summary data for all animals (N = 443 total worms; N = 20-50 worms per condition) and encounters (N = 6,560 total encounters; N = 46-876 encounters per condition) are shown in (A-C,E-I,K-O). Asterisks denote statistical significance (***p<0.001). See also Figure 2 – supplement 1-7 and Video 5-6.

Figure 3: *C. elegans* explore first even when only one patch is available.

(A) Relative density of large (~8.3 mm diameter) and (B) small (~1.8 mm diameter) bacterial patches was varied by pipetting 20 μL and 0.5 μL, respectively, droplets of OP50 *E. coli* diluted in LB to a range of optical densities (OD_{600} = {0, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10} and controlling growth time at room temperature (hours = {1, 48}). (C) Large patches were formed in the center of 30 mm arenas while small patches were formed in the center of 9 mm arenas. (D-E) The midbody location (colored to represent time in the experiment) of example animals foraging in environments containing a single (D) large or (E) small patch (gray) are shown. (F-G) Ethograms of patch encounters (colored to represent the probability of classification as *search*, *sample*, and *exploit*) for (F) 144 individuals (8-15 per condition) foraging on a single large patch and (G) 191 individuals (27-38 per condition) foraging on a single small patch are shown. (H-I) Time elapsed prior to the first *exploitation* event (blue) for animals foraging on (H) large and (I) small patches is plotted for every animal. When *exploitation* was not observed, time elapsed in the experiment is plotted (red-orange). Animals in large and small patch environments *exploited* higher density patches significantly earlier (Kendall's τ correlation, p < 0.001) following a sigmoidal trend. See also Figure 3 – supplement 1-2.

Figure 4: Exploitation decisions are driven by available sensory information, satiety, and prior experience.

(A) Schematic of the covariates x_k used in our logistic regression model to predict the probability of *exploiting* a patch upon any given encounter $p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)$. $\boldsymbol{x}_k$ includes the relative density of the encountered patch k (ρ_k), the duration of time spent off food since departing the last *exploited* patch (τ_s), the relative density of the patch encountered immediately before encounter k (ρ_h), and the relative density of the last *exploited* patch (ρ_e). (B) To account for uncertainty in our classification of *sensation*, we produced 100 sets of observations wherein we probabilistically included *sensed* encounters ($v_k = 1$) estimated from the distribution $v_k|\boldsymbol{w}_k \sim \text{Bern}(p(v_k = 1|\boldsymbol{w}_k))$ where $p(v_k = 1|\boldsymbol{w}_k)$ is the probability that the patch was *sensed* as estimated in Figure 2I. To account for uncertainty in our classification of *exploitation*, we substitute the response variable y_k with our estimate of the probability that the patch was *exploited* $p(y_k = 1|\boldsymbol{z}_k)$ as estimated in Figure 2H, a procedure analogous to including *exploitations* ($y_k = 1$) estimated from the distribution $y_k|\boldsymbol{z}_k \sim \text{Bern}(p(y_k = 1|\boldsymbol{z}_k))$. A schematic of these procedures is shown (see Models of exploitation probability). (C) Coefficient values for each covariate in the GLM were estimated across 50000 replicates (500 replicates of hierarchically bootstrapped animals in combination with 100 sets of probabilistically *sensed* encounters). All coefficients are significantly greater than or less than 0 (two-tailed, one-sample bootstrap hypothesis tests with Bonferroni correction). (D) Using observed $p(y_k = 1|\boldsymbol{z}_k)$ and estimated $p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)$ probabilities of *exploitation*, *exploitation* events were simulated from a Bernoulli distribution. Distributions of the probability of first *exploiting* as a function of the number of encounters as estimated by the model with covariates added one-by-one and as observed are shown for animals foraging in single-density, multi-patch environments of relative density 1 or 10. (E) A schematic of multi-density, multi-patch assays where animals foraged in environments containing small (~1.8 mm diameter) bacterial patches (gray) of varying combinations of OP50 *E. coli* with relative densities 1, 4, and 7 is shown. (F) The probability of exploitation as estimated with ($p(y_k = 1|\beta_0 + \beta_k\rho_k + \beta_s\tau_s + \beta_h\rho_h + \beta_e\rho_e)$) and without ($p(y_k = 1|\beta_0 + \beta_k\rho_k + \beta_s\tau_s)$) the history-dependent terms and as observed $p(y_k = 1|\boldsymbol{z}_k)$ is shown for every pairing of current patch density (ρ_k) combined with the density of recently encountered (ρ_h) or *exploited* (ρ_e) patches. Heat maps for observed values were interpolated between the 9 patch density pairings tested. (G) *Exploitation* events were simulated from the distribution $y_k|\boldsymbol{z}_k \sim \text{Bern}(p(y_k = 1|\boldsymbol{z}_k))$ and used to calculate the probability of observing *exploitation* for the first time as a function of the number of encounters for N2, *mec-4*, and *osm-6* animals foraging in single-density, multi-patch environments of relative density 1 or 10. (H) Coefficient values for each covariate were estimated using ridge regression models for each strain across 50000 replicates. A subset of these coefficients significantly varied between wild-type and mutant strains (mean of differences tests with Benjamini-Hochberg correction). Summary data for all animals (N = 443 total worms; N = 20-50 worms per condition) and encounters ($\mu = 2659.8$, $\sigma = 15.9$) in the single-density, multi-patch assay are shown in (C,D). Summary data for all animals (N = 198 total worms; N = 20-40 worms per condition) and *sensed* encounters ($\mu = 1{,}724.6$, $\sigma = 9.9$) in the multi-density, multi-patch assay are shown in (F). Summary data for all wild-type and mutant animals (N = 221 total worms; N = 14-44 worms per condition) and encounters (N = 1,352 total encounters) in a single-density, multi-patch assay are shown in (G,H). Violin plots in (C,H) show the KDE and quartiles for each measure. Asterisks denote statistical significance († $p_{unadjusted}<0.05$; *$p<0.05$; **$p<0.01$; ***$p<0.001$). See also Figure 4 – supplement 1-5.

Figure Supplements

Figure 1 – supplement 1: Assay preparation, arena and patch detection, and behavioral tracking.

(A) Graphic describing the steps to prepare assay plates. Created with BioRender. (B) Example frames from a "contrast video" showing how a piece of dark cardstock passed between the light source and the assay plate enables visualization of dilute bacterial patches. (C) An image of the bacterial patches is generated from the "contrast video". (D) Masks of the arena (black) and bacterial patches (gray) were constructed using image processing techniques and, when applicable, subsequent manual corrections (see Bacterial patch location detection). Arena masks were used to calculate the image scale in pixels/mm. (E) Example frame from a behavior video containing four adult *C. elegans* is shown. (F) WormLab was used to identify the midbody location of animals in each frame (see Behavioral tracking). An inset shows detection of one of the four worms. (G) An example animal's tracked location as it forages in this environment is plotted with color used to represent time (dark blue = 0 mins; dark red = 60 mins).

Figure 1 – supplement 2: Defining a patch encounter using high resolution behavioral recordings.

(A) Example frame from a behavior video containing one adult *C. elegans* is shown. (B) WormLab was used to track the body of the animal in each frame (see Behavioral tracking). (C) An example image showing the center spline of the example animal as well as the location of the bacterial patch (gray) and arena (black). The inset shows the animal's center spline and bacterial patch superimposed on top of the raw image. (D) Example body postures/orientations (black) of an animal during frames when its head is on the patch edge. The distance (blue lines) between the head (black dot) and midpoint (blue dot) of the animal was calculated. A histogram of this head-to-midpoint distance is plotted, and the median value (blue) is indicated (sample size: 220 animals; 74839 frames). The median head-to-midpoint distance was subsequently used as a threshold for detection of patch encounter and leaving events. For one example animal, the distance between the head (black) or midbody (blue) position and the nearest patch edge (positive = inside patch; negative = outside patch) is plotted for every time point. The edge of the patch (solid black line) and threshold for encounter detection (dashed -.- black line) are indicated. Putative patch encounter (green) and leaving (red) events are shown. (E) Example body postures/orientations of an animal when its head is on the patch. The distance between the midpoint of the animal and the patch edge (blue lines) was calculated. A histogram of this midpoint-to-patch distance is plotted, and the 1st percentile (blue) is indicated (sample size: 220 animals; 1571163 frames). This value is used a threshold for excluding putative encounters where the animal's midpoint never comes within 0.28768 mm of the patch edge. A 1.5 minute time window of the example animal's position in (D) is shown. The edge of the patch (solid line), threshold for encounter detection (dashed -.- line), and threshold for encounter exclusion (dashed – line) are indicated. Example putative encounters are shown. The first two sets of enter and exit events were excluded. The third set shown was included. The included encounter is shaded in gray. (F) The standard deviation of the distance between the location of the patch edge and the midpoint of the animal was calculated for every putative on-patch encounter and off-patch event (sample size: 220 animals; 2945 putative on-patch events; 2725 putative off-patch events). A subset of putative off-patch events had low standard deviations matching those of on-patch events. A one-dimensional Gaussian mixture model was fit to the off-patch standard deviation values and a threshold was set where the posterior probability of events falling in the high or low variance cluster was 0.5. A 1.5 minute time window of the example animal's position in (D) is shown. Examples of two excluded off-patch events and one included event are shown.

behavior video (1024 x 1024 px)　　　　WormLab tracking　　　　worm spline and patch detection

D　① Identify putative patch entries and exits by detecting when the midpoint is within 0.46024 mm of the patch edge

head on patch edge　　head-to-midpoint distance (mm)

E　② Exclude encounters where the midpoint never comes within 0.28768 mm of the patch edge

head on patch　　midpoint-to-patch distance (mm)

F　③ Exclude patch leaving events where the standard deviation of the midpoint-to-patch distance while off patch is below 0.22221

std(distance from patch edge)

1426

1427 **Figure 1 – supplement 3: Classifying encounters based on duration using a Gaussian**
1428 **mixture model.**
1429
(A) As in Figure 1E, duration for each patch encounter (N = 419 events; 50 worms) was computed
and classified as either short or long using a Gaussian mixture model (GMM). The distributions of
all observed short and long encounters are plotted with duration binned logarithmically. The
probability density functions for the two component Gaussians are shown. (B) The posterior
probabilities for classification as short or long are sorted for all encounters, highlighting the low
posterior variance of the GMM. (C) Given that some patch encounters were not fully observed (i.e.,
~22% of animals were already on patch at the start of the recording and ~96% of animals were on
patch when the recording ended), we identified that ~14% of observed patch durations were left-
censored (i.e., having actual duration ≥ observed duration). Thus, to confirm findings in Figure 1E-
F, we re-classified only non-censored observations where both the entry and exit of a patch
encounter were observed. As in Figure 1C, patch encounters for 50 individuals foraging in these
patchy environments are plotted across time. Encounters that were uncensored (pink) and censored
(gray) are indicated. (D) The distribution of the duration of only fully observed (uncensored)
encounters (N = 360 events; 50 worms) is plotted. The component Gaussians for a GMM fitting
this subset of data are shown. (E) Sorted posterior probabilities for the uncensored model are
shown. (F) The observed start time of each uncensored patch encounter is shown for all short and
long duration encounters. Long duration encounters occur significantly later (Mann-Whitney U-Test,
***p<0.001). Violin plots show the KDE and quartiles for each duration type. (G) Posterior
probabilities for classification in the long duration cluster are shown for the two models. The
uncensored data model results in a slight increase in posterior probabilities of long duration
classification, but ultimately affected classification of only one of the 419 encounters.
1451

1452

1453 **Figure 1 – supplement 4: Permuting patches to test for significance of the observed time-**
1454 **dependent increase in patch residence.**

1455

(A) Locations of bacterial patches (dark gray) are shown with the threshold for putative encounters
(light gray) set at 0.46024 mm as described in Figure 1 – supplement 2 and Patch encounter
detection. (B) Patches were permuted via randomized rotation and translation with the restriction
that patches could not overlap with other patches nor with the arena. (C,D) Midbody location of the
animal over time was overlaid on the original and semi-randomly permuted patches. (E,F) On patch
events were subsequently identified. This permutation procedure was repeated for 1000 replicates
for every worm tested. Smoothed median values of the probability of worms residing on patch over
time was compared for every time point in Figure 1G, 2F and Figure 2 – supplement 5.

1464

1465

1466 **Figure 1 – supplement 5: Analyzing deceleration upon encounter with a patch.**

1467

(A) To quantify the magnitude of slow-down upon encounter with the patch edge, we averaged the
velocity trajectories across encounters and found the velocity reached a maximum at 1.5 seconds
before (green) and a minimum at 6 seconds after (red) the start of the encounter (gray). (B,C)
Subsequently, for every encounter, a line (blue) was fit to the animal's velocity (black) for that 8
second time window surrounding the start of the encounter. Deceleration was defined as the slope
of this line. Examples matching (B) the short duration encounter in Figure 1J and (C) the long
duration encounter in Figure 1K are shown.

1475

1476

1477 **Figure 2 – supplement 1: Obtaining intensity profiles for fluorescent bacterial patches.**
1478
(A) Fluorescently labeled OP50-GFP bacterial patches were seeded onto agar plates under
conditions matching those in the experimental assay. Small patches (0.5 μL) were pipetted in an
isometric grid (6 mm center-to-center spacing) on a 16 x 25 mm rectangular template comparable to
the one used in behavioral assays. Large patches (20 or 200 μL) were seeded directly onto agar plates.
Brightfield and fluorescence images were acquired for every condition at multiple time points of
bacterial growth. The fluorescence intensity profile of these images was obtained. (B) Due to
uneven illumination within the field-of-view of our imaging system, matched fluorescence images of
an empty agar plate were acquired at each time point. These "background" images were smoothed
using a two-dimensional averaging filter to remove noise. (C) Fluorescence images of the bacterial
patches were normalized to these background images. (D) Locations of bacterial patches within the
template were automatically detected using image processing techniques. Each patch was then
radially segmented in bins of equal area. The mean pixel intensity value for each bin was computed
and used to create an intensity profile of the patch. (E) The patch border and peak were detected
using signal processing techniques. Specifically, the patch border was identified by finding a peak in
the curvature κ of the edge profile. Border amplitude was defined as the magnitude of the
difference between the patch border and peak. Relative border amplitude was defined as the border
amplitude divided by the exposure time. A line was fit to the values of relative border amplitude
across numerous time points and experiment days (5 replicates shown as varying color points).

A ① Brightfield and fluorescence images were acquired of OP50-GFP bacterial patches seeded uniformly using a template.

brightfield (2208 x 2752 px)   fluorescence (3000 ms exposure)   fluorescence intensity profile

B ② As the diffusion of light in our imaging system resulted in uneven illumination of the field-of-view, matched fluorescence images of an empty agar plate were acquired for each time point. This "background" image was smoothed using a 2D averaging filter.

fluorescence background   fluorescence intensity profile   smoothed fluorescence background

C ③ Fluorescence images of the OP50-GFP bacterial patches were normalized to the smoothed background, removing the majority of the illumination-dependent spatial variance in signal intensity.

fluorescence normalized   normalized intensity profile

D ④ Arena (16 x 25 mm) and bacterial patch locations were automatically detected via image processing techniques. Each patch was radially segmented in bins of equal area to compute an intensity profile of the patch.

E ⑤ Using the profiles curvature and maximum value, the patch border and peak were detected. Border amplitude was defined as the difference in magnitude between the peak and border pixel intensities. Border amplitude was divided by exposure time to compute a relative border amplitude that could be compared across time points and patch densities.

1497

1498 **Figure 2 – supplement 2: Example fluorescence profiles of bacterial patches under varied**
1499 **growth conditions.**
1500
Fluorescently-labeled OP50-GFP bacterial patches were imaged as described in Figure 2 –
supplement 1 and Bacterial patch density estimation. Inverted images (darker saturation = more
bacteria) are shown here to highlight the range of bacterial densities tested in these experiments. All
patches are shown on the same spatial scale and using one of the two linear saturation scales: 1)
regular – normalized to show variance in high density patches; 2) augmented (~22x) – normalized to
show variance in low density patches). Each patch shown is labeled with the optical density (OD_{600})
at time of seeding and the amount of time bacteria was grown for at room temperature (i.e., 1, 12,
or 48 hours). (A) An example large (200 μL) bacterial patch of the type that animals experienced
immediately prior to the assays in this paper. (B) Example small (0.5 μL) bacterial patches of greatest
density. (C) Example small (0.5 μL) bacterial patches of lower density. The saturation of each image
has been uniformly augmented to ~22x darker to the extent that the OD_{600} 10 (1H) bacterial patch is
the same patch as the one in B. (D) Example small (0.5 μL) bacterial patches of lower density seeded
on NGM agar plates lacking peptone. (E) Example medium (20 μL) bacterial patches of greater
density. (F) Example medium (20 μL) bacterial patches of lowest density. The OD_{600} 1 (1H) bacterial
patch is the same patch as the one shown in E, but ~22x darker.

A   200 µL - Ø 21.3 mm

OD$_{600}$ = 1 (48H)

5 mm

B   0.5 µL - Ø 1.7 mm

OD$_{600}$ = 1 (48H)   OD$_{600}$ = 1 (12H)   OD$_{600}$ = 10 (1H)

C   0.5 µL - Ø 1.7 mm (augmented saturation)

OD$_{600}$ = 10 (1H)   OD$_{600}$ = 5 (1H)   OD$_{600}$ = 4 (1H)   OD$_{600}$ = 3 (1H)

OD$_{600}$ = 2 (1H)   OD$_{600}$ = 1 (1H)   OD$_{600}$ = 0.5 (1H)

D   0.5 µL - Ø 1.7 mm - no peptone (augmented saturation)

OD$_{600}$ = 10 (1H)   OD$_{600}$ = 5 (1H)   OD$_{600}$ = 4 (1H)   OD$_{600}$ = 3 (1H)

OD$_{600}$ = 2 (1H)   OD$_{600}$ = 1 (1H)   OD$_{600}$ = 0.5 (1H)

E   20 µL - Ø 8.3 mm

OD$_{600}$ = 1 (48H)   OD$_{600}$ = 10 (1H)   OD$_{600}$ = 5 (1H)   OD$_{600}$ = 4 (1H)   OD$_{600}$ = 3 (1H)

OD$_{600}$ = 2 (1H)   OD$_{600}$ = 1 (1H)

F   20 µL - Ø 8.3 mm (augmented saturation)

OD$_{600}$ = 1 (1H)   OD$_{600}$ = 0.5 (1H)

1516

1517 **Figure 2 – supplement 3: Quantifying the relative density of bacterial patches across**
1518 **conditions and time points.**
1519
(A) Relative density of OP50-GFP bacterial patches matched to all assay conditions in this paper are
shown. Relative density represents the relative border amplitude (as defined in Figure 2 –
supplement 1 and Bacterial patch density estimation) normalized to the mean density of patches
seeded with 0.5 μL of OD_{600} = 10 grown for 1 hour on plates with peptone (blue asterisk). In this
manner, the bacterial patches experienced by animals in Figure 1 have been set to an average relative
density of 10. Violin plots show the KDE and quartiles for each condition. (B) Linear fits for the
relative density of bacterial patches in each condition as a function of time are plotted. Saturation of
each line corresponds to the relative density as depicted in (A). For each experimental plate, the
relative density of bacterial patches was estimated by applying the coefficients from these linear
regressions to the total amount of time the bacteria was grown at room temperature.
1530

1531

 Figure 2 – supplement 4: Example traces of animals foraging in environments with varying
bacterial density.

Traces showing the midbody location of 12 example animals as they forage within a 30 mm arena
are plotted with color used to represent time (dark blue = 0 mins; dark red = 60 mins). Locations of
bacterial patches are indicated in gray. The relative density of bacterial patches in each environment
is noted. Animals shown foraging on relative density 0, 1, 5, 10, and 200 are the same as those in
Figure 2D.

1541

1542 **Figure 2 – supplement 5: Time- and density-dependent increase in patch residence.**
1543

Probability of residing on patch was computed for all worms across time (black) and compared to
the probability of residing on patch for semi-randomly permuted patch locations (pink) as described
in Figure 1 – supplement 4. Smoothed median values are plotted with bootstrap-derived 2.5% and
97.5% quantiles shown in shaded regions. Time points where observed probabilities of residing on
patch significantly exceed permuted probabilities are indicated by a black line (one-tailed Fisher's
Exact Tests with Benjamini-Hochberg correction, ***p < 0.001). The smoothed median values for
observed data are shown in Figure 2F. Data for relative density 10 match those in Figure 1G.
1551

1552

1553 **Figure 2 – supplement 6: Classifying encounters as exploration or exploitation.**
1554

(A) The average velocity of the animal during an encounter and the duration of that encounter are
plotted for each encounter as visualized on a double-logarithmic plot. Contours showing the first,
second, and third standard deviation of the two-dimensional Gaussian mixture model (GMM) are
shown as shaded ellipses with saturation corresponding to standard deviation. Encounters for all
data in the paper are individually plotted as in Figure 2H but with color representing the posterior
probability of clustering classification as *explore* (green) or *exploit* (blue). (B) Posterior variance of the
data as defined in Patch encounter classification as *exploration* or *exploitation* was calculated for 1000
replicates of GMMs with varied regularization value α. The classifier with minimum posterior
variance (i.e., $\alpha = 0.025$) was used. (C) The posterior probabilities for classification as *explore*
(green) or *exploit* (blue) are sorted for all encounters, highlighting the low posterior variance of the
GMM. (D) Posterior probabilities for only the data in Figure 2 are shown.
1566

1567

Figure 2 – supplement 7: Classifying encounters as sensing or non-sensing.

(A) Minimum velocity (min. velocity on patch), maximum change in velocity (max. Δ velocity), and deceleration (Δ velocity) were calculated as described in Patch encounter classification as sensing or non-sensing. (B) Maximum change in velocity for every density condition is shown. Violin plots show the KDE and quartiles for each condition. (C) Deceleration upon encounter and minimum on-patch velocity during the encounter are plotted for each encounter. A three-dimensional parabolic boundary between *sensing* and *non-sensing* clusters was fit using semi-supervised QDA and a subset of labeled data (blue circles = sensing, dark orange circles = non-sensing). Regions identified by the classifier as *non-sensing* or *sensing* are indicated by colored contours (orange = *non-sensing*, green = *sensing*). Encounters for all data in the paper are individually plotted as in Figure 2I but with color representing the conditional probability of clustering classification as *non-sensing* (orange) or *sensing* (green). (D) Maximum change in velocity and minimum on-patch velocity during the encounter are plotted for each encounter in the paper as in (C). (E) Minimum velocity on patch, maximum change in velocity, and deceleration are shown for every encounter in the paper. See Video 5 for three-dimensional rotation of this plot. (F) The probabilities for classification as *sense* (green) or *non-sense* (orange) are sorted for all encounters. (G) Probabilities for only the data in Figure 2 are shown. (H) For a subset of encounters that were censored (i.e., recording did not observe animals entering the patch), probabilities of *sensing* were estimated using the semi-supervised QDA approach as well as via marginalization over the conditional probabilities using only the observed minimum velocity on patch. Marginalized probabilities were used for these values as described in Patch encounter classification as sensing or non-sensing.

Emit

Figure 3 – supplement 1: Example traces of animals foraging in environments with one large bacterial patch.

Example traces showing the midbody location of 11 animals as they forage within a 30 mm arena are plotted with color used to represent time (dark blue = 0 mins; dark red = 60 mins). Location of bacterial patches are indicated in gray. The relative density of bacterial patches in each environment is noted. The animal shown foraging on relative density 10 is the same as that in Figure 3D.

1599 **Figure 3 – supplement 2: Example traces of animals foraging in environments with one**
1600 **small bacterial patch.**

1601

1602 Example traces showing the midbody location of 6 animals as they forage within a 9 mm arena are
1603 plotted with color used to represent time (dark blue = 0 mins; dark red = 60 mins). Location of
1604 bacterial patches are indicated in gray. The relative density of bacterial patches in each environment
1605 is noted. The animal shown foraging on relative density 10 is the same as that in Figure 2E.

1606

1607

Figure 4 – supplement 1: Model selection for nested GLMs.

(A) Values of log-likehood, Aikake Information Criterion (AIC), and Bayesian Information Criterion (BIC) were computed for 50,000 replicates of the linear regression model described in Models of exploitation probability. Each of these metrics is plotted here across 100 replicates of "encounter samples" (i.e., we removed encounters where the animal likely did not *sense* the encounter $v_k = 0$ as estimated from the distribution $v_k | w_k \sim \text{Bern}(p(v_k = 1 | w_k))$) and 500 replicates of hierarchically bootstrapped "worm samples" (i.e., we resampled animals with replacement and included all encounters of the resampled animals). Values for each metric at each replicate are colored to match their respective color bar. Metrics were computed for the addition of each covariate to the model. Lower AIC and BIC values for the full model ((i.e., $\boldsymbol{\beta} \cdot \boldsymbol{x}_k = \beta_0 + \beta_k \rho_k + \beta_s \tau_s + \beta_h \rho_h + \beta_e \rho_e$), represent better model performance even with penalization for increasing the number of parameters. (B) Coefficients $\boldsymbol{\beta}^*$ for the null distribution (i.e., observations of the response variable $p(y_k = 1 | \boldsymbol{z}_k)$ shuffled relative to the covariates $\boldsymbol{x}_k$) are plotted. Only β_0 was significantly greater than or less than 0 (two-tailed, one-sample bootstrap hypothesis tests with Bonferroni correction). Violin plots show the KDE and quartiles for each covariate across the 50,000 replicates. Asterisks denote statistical significance (*p<0.05; **p<0.01; ***p<0.001). These $\boldsymbol{\beta}^*$ represent the null distribution for the $\boldsymbol{\beta}$ in Figure 4C.

1628 **Figure 4 – supplement 2: Probability of first exploitation as a function of the number of**
1629 **encounters.**
1630
Histograms of the probability of first *exploiting* as a function of the number of encounters as
estimated by the models and as observed are shown for animals foraging in single-density, multi-
patch environments of all 12 density conditions. *Exploitation* events were simulated from a Bernoulli
distribution with probability estimated by the model $y_k \,|\boldsymbol{x}_k \sim \text{Bern}\big(p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)\big)$ where
covariates were added one at time as well as from the observed probabilities estimated by the
Gaussian mixture model classification $y_k|\boldsymbol{z}_k \sim \text{Bern}\big(p(y_k = 1|\boldsymbol{z}_k)\big)$ in Figure 2H. Histograms for
relative density 1 and 10 are the same as those in Figure 4D.

Figure 4 – supplement 3: Exploitation decision of food-deprived animals is well-predicted
by the model.
1641

(A) Schematic of experiments used to test satiety's influence on the decision to *exploit*. As in all
experiments, animals were acclimated to high density (relative density ~200) patches for ~24 hours.
Well-fed animals were then removed from these acclimation plates, cleaned of bacteria, and then
immediately transferred to the assay plate with relative density 5. Food-deprived animals moved
about a bacteria-free arena for 3 hours prior to the assay. Examples of animal behavior while
foraging in acclimation-like, bacteria-free, and single-density, multi-patch environments of relative
density 5 are shown. (B) Patch encounters for 28 food-deprived and 28 well-fed individuals are
plotted across time. Each encounter is colored to match its probability of classification as *search*
(orange), *sample* (green), and *exploit* (blue). (C) Coefficient values were re-estimated across 50000
replicates as in Figure 4C for a model with the satiety term removed (i.e.,
$p(y_k = 1|\beta_0 + \beta_k\rho_k + \beta_h\rho_h + \beta_e\rho_e)$). All coefficients are significantly greater than or less than 0
(two-tailed, one-sample bootstrap hypothesis tests with Bonferroni correction; ***p<0.001). Violin
plots show the KDE and quartiles for each measure. (D) Histograms of the probability of first
exploiting as a function of the number of encounters are shown for food-deprived and well-fed
animals. Model predictions of *exploitation* events were simulated from a Bernoulli distribution
$y_k|\boldsymbol{x}_k \sim \text{Bern}\big(p(y_k = 1|\boldsymbol{\beta} \cdot \boldsymbol{x}_k)\big)$ where the coefficient values $\boldsymbol{\beta}$ corresponded to those previously
estimated for our linear regression model with (Figure 4C) and without (Figure 4 – supplement 3C)
the satiety term using the data set shown in Figure 2 (i.e., the 443 well-fed animals foraging in one of
12 density conditions). Observed *exploitations* were simulated from a Bernoulli distribution using the
probability of *exploitation* estimated by our GMM classifier $y_k|\boldsymbol{z}_k \sim \text{Bern}\big(p(y_k = 1|\boldsymbol{z}_k)\big)$ in Figure
2H. Summary data for all animals (N = 56 total worms; N = 28 worms per condition) and
encounters (N = 277 total encounters; N = 42-235 encounters per condition) are shown in (B,D).
1664

1665

1666 **Figure 4 – supplement 4: Example traces and classified behavior of animals foraging in**
1667 **environments with multiple patch densities.**
1668
(A) Example traces showing the midbody location of 7 animals as they forage within a 30 mm arena
are plotted with color used to represent time (dark blue = 0 mins; dark red = 60 mins). Locations of
bacterial patches are indicated in gray with saturation indicating the relative density. Environments
contained combinations of patches with relative density 1, 4, and 7 as noted. (B) Patch encounters
for 200 individuals (20-40 per condition) are plotted across time. Each encounter is colored to match
its relative density. (C) Patch encounters are colored to match their probability of classification as
search (orange), *sample* (green), and *exploit* (blue). (D) Coefficient values were re-estimated across
50000 replicates as in Figure 4C for a model with the history-dependent terms removed (i.e.,
$p(y_k = 1|\beta_0 + \beta_k\rho_k + \beta_s\tau_s)$). All coefficients are significantly greater than or less than 0 (two-
tailed, one-sample bootstrap hypothesis tests with Bonferroni correction; **p<0.01; ***p<0.001).
Violin plots show the KDE and quartiles for each measure. Summary data for all animals (N = 198
total worms; N = 20-40 worms per condition) and encounters (N = 3,493 total encounters; N =
282-795 encounters per condition) are shown in (B,C).

Figure 4 – supplement 5: Example traces and classified behavior of animals with chemosensory or mechanosensory deficiencies.

(A) Example traces showing the midbody location of 9 animals as they forage within a 30 mm arena are plotted with color used to represent time (dark blue = 0 mins; dark red = 60 mins Location of bacterial patches are indicated in gray. The relative density of bacterial patches in each environment is noted. 3 strains of *C. elegans* were tested (grey = N2, pink = *mec-4*, orange = *osm-6*) across 3 conditions of relative density (1, 5, and 10). (B) Patch encounters for 76 individuals (16-44 per density condition) of each strain are plotted across time. Each encounter is colored to match its probability of classification as *search* (orange), *sample* (green), and *exploit* (blue). (C) Histograms of the probability of first *exploiting* as a function of the number of encounters are shown for each strain and density condition. Observed *exploitations* were simulated from a Bernoulli distribution using the posterior probability of *exploitation* defined by our GMM classifier $y_k|z_k \sim Bern\big(p(y_k = 1|z_k)\big)$ in Figure 2H. Histograms of each strain for relative density conditions 1 and 10 are the same as those in Figure 4G. (D) Coefficients $\boldsymbol{\beta}^*$ for the null distribution (i.e., observations of the response variable $p(y_k = 1|z_k)$ were shuffled relative to the covariates $\boldsymbol{x}_k$) are plotted. Violin plots show the KDE and quartiles for each covariate across the 50,000 replicates. These $\boldsymbol{\beta}^*$ represent the null distribution for the $\boldsymbol{\beta}$ in Figure 4H. (E) The ridge regression parameter λ was optimized for each strain to maximize the mean log-likelihood of the model. Summary data for all animals (N = 221 total worms; N = 14-44 worms per condition) and encounters (N = 1,352 total encounters; N = 27-193 encounters per condition) are shown in (B,C).

Video Legends

Video 1. *C. elegans* foraging in a patchily distributed environment.
Four adult *C. elegans* forage for one hour in a 30 mm arena containing an isometric grid of small, low density bacterial patches.

Video 2. Visualizing dilute bacterial patches using diffraction of light.
A piece of dark cardstock passed between the light source and the assay plate enables visualization of dilute bacterial patches.

Video 3. Tracking the midbody location of an animal foraging in a patchily distributed environment.
The midbody location of one of the four animals shown in Video 1 is shown as the animal forages for one hour in a 30 mm arena containing an isometric grid of small, low density bacterial patches. The animal shown corresponds to the example animal observed in Figure 1A-B,H-K.

Video 4. Tracking the entire body of a foraging animal.
The position of the entire body of an animal is shown as the animal forages for one hour in a 9 mm arena containing one small, low density bacterial patch. The head is indicated by a small dot. The animal shown corresponds to the example animal observed in Figure 3D.

Video 5. Classifying encounters as *sensing* based on quantification of slow-down.
Three metrics quantifying the magnitude of slow-down upon patch encounter were computed. A semi-self-supervised quadratic discriminant analysis was trained on a subset of labeled data and used to classify all encounters as *sensing* (green) or *non-sensing* (orange). Data correspond with Figure 2I and Figure 2 – supplement 7C-E.

Video 6. Foraging behavior of an animal with encounters labeled as *search*, *sample*, or *exploit*.
The midbody location of the animal in Video 3 is shown and labeled by the probability of classification as *searching* (i.e., off patch *exploration* and patch encounters that were not *sensed*), *sampling* (i.e., patch encounters that were *sensed*, but not *exploited*), and *exploiting* at every time point. The animal shown corresponds to the example animal observed in Figure 1A-B,H-K.