Ocular hypertension impairs axonal transport in the optic nerve head leading to neurodegeneration in a novel Cre-inducible mouse model of myocilin glaucoma.

Balasankara Reddy Kaipa[1], Ramesh Kasetti[2], Linya Li[1], J. Cameron Millar[2], William Cho[1], Dorota Skowronska-Krawczyk[1], Prabhavathi Maddineni[3], Yogapriya Sundaresan[1], and Gulab S. Zode[1]*

[1]Gavin Herbert Eye Institute-Center for Translational Vision Research, Department of Ophthalmology, Department of Physiology and Biophysics, University of California Irvine School of Medicine, Irvine, CA 92697, USA
[2]Department of Pharmacology and Neuroscience, North Texas Eye Research Institute, University of North Texas Health Science Center at Fort Worth, TX 76107, USA
[3]Department of Ophthalmology, School of Medicine, University of Missouri, Columbia, MO

*Corresponding author: Gulab S. Zode, PhD
829 Health Sciences Rd, Irvine, CA 92617
Telephone: 949-824-4366
(gzode@hs.uci.edu)

Running title: IOP elevation impairs axonal transport leading to neurodegeneration.
Keywords: Glaucoma, trabecular meshwork, retinal ganglion cells, neuronal loss, ocular hypertension, myocilin, protein misfolding, mouse model of glaucoma, axonal transport deficits.

Disclosure Statement: Authors declare no conflict of interest.

Abstract

Background: Degeneration of optic nerve (ON) axons and loss of retinal ganglion cells (RGCs) are the pathological hallmarks of Primary Open Angle Glaucoma (POAG). Elevation of intraocular pressure (IOP) due to dysfunction of trabecular meshwork (TM) is known to induce neurodegeneration. However, the early pathological events of glaucomatous neurodegeneration are poorly understood due to lack of robust and faithful mouse model that replicates all features of human POAG. Here, we report the generation and characterization of a novel Cre-inducible transgenic mouse model of myocilin (MYOC), the leading known genetic cause of POAG. Using this model, we further explore early pathological events of glaucomatous neurodegeneration due to chronic IOP elevation.

Methods: We generated a Cre-inducible transgenic mouse model expressing DsRed-tagged Y437H mutant of human myocilin (*Tg.CreMYOC*^{Y437H}). A single intravitreal injection of helper adenovirus (HAd) 5 expressing empty cassette or Cre was performed in adult *Tg.CreMYOC*^{Y437H} mice, and glaucoma phenotypes including IOP, outflow facility, structural and functional loss of RGCs, ON degeneration, gliosis, and axonal transport deficits were examined at various stages of IOP elevation.

Results: An intravitreal injection of HAd5-Cre led to selective *MYOC* expression in the TM at the level similar to endogenous *Myoc*. Expression of mutant MYOC induced

biochemical and ultrastructural changes in TM leading to reduced outflow facility and significant IOP elevation. Notably, sustained IOP elevation led to significant functional and structural loss of RGCs and progressive ON degeneration. Glaucomatous neurodegeneration was associated with activation of astrocytes and neurodegenerative changes in the optic nerve head (ONH) region. Remarkably, chronic IOP elevation blocked anterograde axonal transport at the ONH prior to axonal degeneration and RGC loss. Interestingly, impaired axonal transport was associated with loss of cytoskeleton proteins including microtubules and neurofilaments resulting into accumulation of mitochondria in the ONH and neuronal dysfunction.

Conclusions: Our studies indicate that Cre-inducible *Tg.CreMYOC*[Y437H] mice recapitulates all glaucoma phenotypes observed in POAG patients. Importantly, sustained IOP elevation impairs axonal transport at ONH leading to glaucomatous neurodegeneration.

Introduction

Glaucoma is a group of multifactorial neurodegenerative diseases characterized by progressive optic neuropathy. It is the second leading cause of irreversible vision loss with more than 70 million people affected worldwide [1, 2], and this number is estimated to increase to 111.6 million by the year 2040 [3]. Primary open angle glaucoma (POAG) is the most common form of glaucoma, accounting for approximately 70% of all the cases [1]. POAG is characterized by progressive loss of retinal ganglion cell (RGC) axons and irreversible loss of vision [1, 2]. Elevated intraocular pressure (IOP) is a major risk factor, and the only treatable risk factor for POAG [4]. The trabecular meshwork (TM), a molecular sieve-like structure, regulates IOP by constantly adjusting the resistance to aqueous humor (AH) outflow. In POAG, increased resistance to AH outflow elevates IOP leading to neurodegeneration [5-7]. This increase in outflow resistance is associated with dysfunction of the TM[8-12]. Most of the current POAG treatment strategies do not target the underlying pathology of TM and RGCs, and vision loss continues to progress in some POAG patients[13]. To understand these pathological mechanisms of TM dysfunction/IOP elevation and glaucomatous neurodegeneration, there is an unmet need to develop simple and reliable animal model that can faithfully replicate all features of POAG.

Currently, several mouse models of ocular hypertension (OHT) are utilized to study pathophysiology of glaucomatous neurodegeneration[14, 15]. These include DBA/2J mice and inducible mouse models that physically block AH outflow through the TM leading to OHT and neuronal loss [16-21]. These models have provided important mechanistic understanding of glaucomatous neurodegeneration. However, these models do not allow study of TM pathology since these models elevate IOP by physically blocking outflow pathway [18-20]. Also, these models do not truly represent human POAG phenotype in which the AH outflow pathway is open. Importantly, these mouse model destroy the TM, which can induce acute and sudden IOP-induced glaucomatous neurodegeneration. These models are often technically challenging to replicate in the laboratory settings and exhibit variable phenotypes as well as exhibit other confounding features such as ocular inflammation. Developing a mouse model of known genetic cause of POAG represents an ideal strategy to understand the pathophysiology of POAG.

Mutations in myocilin (*MYOC*) are the most common genetic cause of POAG [6, 22, 23]. *MYOC* mutations are responsible for most cases of Juvenile-onset OAG (JOAG)[24]. *MYOC*-associated JOAG presents clinically with high IOP in young children's and progresses rapidly to vision loss[6, 23]. Mouse models with these genetic alterations in *MYOC* gene mimicking human POAG have proved invaluable tools for understanding the pathogenesis of and designing treatment strategies for POAG[25-27]. Previously, we have developed a transgenic mouse of myocilin POAG by random insertion of human mutant myocilin (*Tg-MYOC*[Y437H]) and demonstrated that *Tg-MYOC*[Y437H] mice develop glaucoma phenotypes closely resembling those seen in patients with POAG caused by the Y437H *MYOC* mutation[27]. However, *Tg-MYOC*[Y437H] mice presented few drawbacks including 1) possible unknown gene mutations and multiple copies of transgene due to random integration, 2) mild phenotype on pure C57BL/6J background limiting the study of glaucomatous neurodegeneration, 3) possibility of transgene silencing with subsequent breeding and 4) lack of specific antibodies to detect myocilin in mouse TM. To overcome these limitations, we utilized a TARGATT site-specific knock in strategy[28] to generate transgenic mice expressing human mutant *MYOC*. This technology utilizes serine integrase, PhiC31 (ΦC31) to insert any gene of interest (a single copy) into a preselected intergenic and transcriptionally active genomic locus (H11), which has been engineered with a docking site. This allows stable and site-specific transgene integration. Since mutant MYOC is toxic to TM cells[27, 29], we exploited Cre-lox system to develop inducible mouse model in which mutant MYOC is expressed in tissue of interest upon Cre expression. Here, we report the development and characterization of a Cre-inducible transgenic mice expressing DsRed-tagged Y437H mutant of human myocilin (*Tg.CreMYOC*[Y437H]). We further utilized this model to investigate early events of glaucomatous neurodegeneration in response to chronic IOP elevation. Importantly, we demonstrate that sustained IOP elevation induced by mutant MYOC impairs axonal transport in optic nerve head (ONH) leading to glaucomatous neurodegeneration.

Methods
Animal husbandry: Animals were housed and bred in a standard 12-hour light/12-hour dark conditions. Animals were fed standard chow *ad libitum* and housed in cages with dry bedding. Two to three months old C57BL/6J mice were obtained from the Jackson Laboratory (Bar Harbor, ME, USA). mT/mG mice were obtained from Dr. *Krzysztof Palczewski* lab and bred as described previously[30]. *Tg.CreMYOC*[Y437H] transgenic mice on pure C57BL/6J background were generated in the lab as described below. Animal studies were performed in agreement with the guidelines of the Association for Research in Vision and Ophthalmology (ARVO) Statement for the Use of Animals in Ophthalmic and Vision Research. Experimental protocols were approved by appropriate IACUC and Biosafety Committees at the University of California, Irvine (UCI). Mice were euthanized via inhalation of carbon dioxide followed by cervical dislocation as described previously [31].

Generation of *Tg.CreMYOC*[Y437H] mice: *Tg.CreMYOC*[Y437H] mice were generated using the TARGATT site-specific transgenic technology (Applied Stem Cell) according to the manufacturer's protocol[28]. Briefly, the cDNA for MYOC with Y437H mutation was subcloned into a TARGATT plasmid (AST-3050, Applied Stem Cell) that contains the attB

recombination site. The expression cassette contained a STOP signal flanked with loxP site in the front of Y437H MYOC cDNA under the control of CAG promoter. A mixture of donor plasmid (containing the attB sites and the transgene) and the ΦC31 mRNA was injected into the pronuclei of H11P3-C57BL/6 mouse embryos. Genomic DNA of a founder was analyzed using primer pairs for the transgene and H11 locus to verify site-specific insertion. The primer set; AST 2005 (Applied Stem Cell, Inc) was utilized to detect H11 site specific integration. Transgene positive mice were bred with C57BL/6J mice and heterozygous mice were utilized for experiments. The following primers were utilized to detect the transgene during regular genotyping. Forward primer: CTCAGCAGATGCTACCGTCA and reverse primer: GCACCTTGAAGCGCATGAA.

Cre recombinase induction by viral vectors: Ad5-Cre or HAd5-Cre under the control of CMV promoter was utilized to selectively induce Cre activity in the TM. Ad5 and HAd5 expressing empty cassette was utilized as a control. These vectors were purchased from Viral Core Facility at the University of Iowa. Mice were anaesthetized using 2.5% isoflurane plus 100% oxygen. A single intravitreal injection of Ad5 or HAd5 expressing Cre or empty cassette (2×10^7 pfu/eye) was performed as described previously[29, 31-33]. Viral injections were performed in 3 to 6 months old male and female *Tg.CreMYOC^{Y437H}* mice in all experiments except for axonal transportation in which 15-month-old *Tg.CreMYOC^{Y437H}* mice were utilized.

Antibodies and reagents: The following antibodies were used in the current study. Myocilin monoclonal antibody (catalog # 60357-1-Ig, Proteintech), Dsred antibody (catalog # 600-401-379, Rockland), KDEL (catalog # Ab12223, Abcam), ATF4 (catalog # SC-200, Santa Cruz Biotechnology)[34], CHOP (catalog # 13172, Novus Biologicals), GRP78 (catalog # ab21685, Abcam), GRP94 (catalog # 11402, Santa Cruz Biotechnology), RBPMS (catalog # 118619, Gene Tex), GAPDH (catalog # 3683, Cell Signaling Technology), TOM20 (catalog # 11802-1-Ap, Proteinteck), TUJ1(catalog # GTX130245, Genetex), GFAP (catalog #, ab4674-1001, Abcam), Fibroneclin (catalog # AB2413-1001), Neurofilament H (catalog # 80170l, Biolegend), Phalloidin stain for actin (catalog #12956, cell signaling), and ant-alpha SMA (catalog # ab5694, Abcam).

Real-time PCR: The anterior segment of Cre^- and Cre^+ *Tg.CreMYOC^{Y437H}* mice was submerged in RNAprotect Cell Reagent (catalog #76526, QIAGEN). Subsequently, the anterior segment was pelleted by centrifugation for 5 minutes at 600g and then subjected to total RNA extraction using a RNeasy Mini Kit (QIAGEN). Following RNA extraction, real-time PCR was carried out using the SYBR® Green Quantitative RT-qPCR Kit (catalog # QR0100, Sigma Aldrich) with the following primers: Forward primer, 5'-TGACTTGGCTGTGGATGAAG-3'; Reverse primer: 5'-TTGTCTCCCAGGTTTGTTCG-3'.

Western blot analysis: Ocular tissues were carefully dissected and lysed in RIPA buffer (Thermo Fisher) as described previously[27, 34]. Approximately 20–30 µg of total protein was separated on denaturing 4–12% gradient polyacrylamide ready-made gels (NuPAGE Bis–Tris gels, Life Technologies, Grand Island, NY, USA) and subsequently transferred onto PVDF membranes. The blots were blocked with 10% non-fat dried milk for 1 hour

and then incubated overnight with specific primary antibody at 4 °C on a rotating shaker. Membranes were washed three washes with phosphate-buffered saline (PBS)/Tween buffer (PBST) and were incubated with the corresponding HRP-conjugated secondary antibody for 90 minutes. Proteins were visualized on the LI-COR Odyssey Fc image system using ECL detection reagents (Super Signal West Femto Maximum Sensitivity Substrate; Life Technologies)[29, 35]. The same blot was subsequently incubated with a GAPDH antibody (Cell Signaling Technology Inc) to ensure the equal protein loading.

Anterior segment flat mount: Whole anterior segment flat mount was performed to image DsRed-MYOC levels in the entire TM region. Cre[-] and Cre[+] *Tg.CreMYOC[Y437H]* mice were enucleated, fixed with 4% paraformaldehyde (PFA), and dissected carefully to separate anterior and posterior segment. Anterior segments were then transferred onto a slide and cut into 4 quadrants to place tissue flat on the slide. Anterior segments were mounted with DAPI-mounting solution and images were captured using a Keyence microscope (Itasca, IL, USA).

RNA scope: In situ hybridization was performed using the RNAscope® Multiplex Fluorescent Assay v2 (ACD Diagnostics) following modifications as described before [36]. Briefly, fresh frozen histologic sections of mouse eyes were pretreated according to the manusfactors protocol using hydrogen peroxide and target retrieval reagents including protease IV. Probes were then hybridized according to the protocol and then detected with TSA Plus® Fluorophores fluorescein, cyanine 3, and cyanine 5. Sections were mounted with Prolong Gold Antifade (Thermo Fisher) with coverslip for imaging and imaged using confocal microscope (SP8, Leica). Probes specific for *Myoc* (Cat # 506401-C3) and *MYOC* (Cat # 506541) transcripts were designed by the manufacturer.

Immunostaining: Enucleated eyes were fixed with 4% PFA for 3 hours. The eyes were embedded in paraffin or OCT compound (Tissue-Tek). Sections of 5 or 10 microns were cut and utilized for immunostaining as described previously[27, 35]. For paraffin-embedded samples, the slides were deparaffinized and rehydrated, followed by antigen retrieval with citrate buffer (pH = 6). For OCT-embedded sections, slides were washed with PBS. The slides were then incubated with blocking buffer (10% goat serum and 0.5% triton-X-100 in PBS) for 2 hours. The slides were incubated with the primary antibody in blocking buffer overnight. After three washes with PBS, the slides were incubated with the appropriate Alexa Fluor secondary antibodies (Life Technologies, Grand Island, NY, USA) for 2 hours. Following three final washes in PBS, the sections were mounted with DAPI-mounting solution, and images were captured using a Keyence microscope (Itasca, IL, USA) as described previously [35, 37].

Mouse slit-lamp examination: To evaluate ocular abnormalities and inflammation in HAd5-Empty and HAd5-Cre-injected eyes, slit-lamp microscopy (SL-D7; Topcon) was performed as described previously[27].

IOP measurements: IOP measurements were conducted using the TonoLab® rebound tonometer (Colonial Medical Supply, Franconia, NH, USA) under anesthetic conditions (isoflurane (2.5%); oxygen (0.8 L/min)) as described previously [34] All IOP assessments

were performed between 10 AM and 2 PM in a masked manner. In addition, IOP measurements were replicated by another technician in a masked manner. The final IOP value was obtained by averaging six individual IOP measurements. Conscious IOPs were measured as described previously[38].

Aqueous humor outflow facility: The outflow facility in Cre⁻ and Cre⁺ *Tg.CreMYOC^Y437H* mice was assessed using the constant-flow infusion method as previously described[38] Briefly, mice were anesthetized with a ketamine/xylazine solution (100/10 mg/kg), and body temperature was maintained at a physiological level using a 37°C electric heating pad throughout the procedure. Topical ocular proparacaine HCl (0.5%) was administered to induce corneal anesthesia. Anterior chambers were cannulated with 32-gauge ½" steel needles (Steriject, Keeler, USA) through the cornea without contacting the iris, anterior lens capsule, or corneal endothelium. The needles were connected to tubing running to a pressure transducer. The opposite end of the transducer was connected by more tubing to a 100 µL microsyringe (Hamilton Co., Reno, NV, USA) loaded into a microdialysis infusion pump, which was filled with sterile PBS. Eyes were infused at a flow rate ranging from 0.1 µL/min to 0.5 µL/min (in 0.1 µL/min increments), and three stabilized pressures at 15-minute intervals were recorded for each flow rate. The aqueous humor outflow facility was then calculated as the reciprocal of the slope (determined by simple linear regression) of the linear part of a plot of mean stabilized pressure as the ordinate and flow rate as the abscissa. In a small proportion of cases (13%), data points obtained at flow rates of 0.5 µL/min deviated from linearity. Such points were excluded from the analysis.

Pattern electroretinography (PERG): The RGC function was assessed using binocular snout pattern electroretinography (PERG) system (JORVEC Corp., Miami, FL, USA) as described previously[35]. Briefly, mice were anesthetized with an intraperitoneal injection of ketamine/xylazine mixture (100, 10 mg/kg, respectively). Anesthetized mice were positioned on a temperature-controllable metal base at a fixed distance (10 cm) from LED monitors and maintained at a constant body temperature (37°C) using a rectal probe. To prevent corneal dryness during recording, a small amount of hypromellose eye drops was applied topically[31, 35]. The PERG was simultaneously derived from each eye using subcutaneous electrodes placed at the snout (active), back of the head (reference), and tail (ground) in response to contrast reversal of gratings generated from two LED screens operating at slightly different frequencies. Two consecutive readings were averaged, and amplitudes (P1–N2) and latencies indicating RGC soma function were shown graphically.

Whole-mount retina staining with RBPMS: Total number of RGCs was analyzed using whole-mount retina staining with RBPMS antibody as described previously[35]. The enucleated eyes were fixed with 4% PFA for 12 hours at 4 °C. After rinsing the eyes with PBS, the anterior segment was removed, and the retinas were carefully separated from the posterior cup. The isolated retinas were incubated with blocking buffer (PBS containing 10% goat serum and 0.2% Triton X-100) for 12 hours at 4 °C. Subsequently, the retinas were incubated with the RBPMS antibody for 3 days at 4 °C followed by a 2-hour wash in PBS. After washing, the retinas were incubated with a corresponding secondary antibody (goat anti-rabbit 568, 1:500; Invitrogen) for 2 hours at room

temperature, washed with PBS three times, and then mounted with a mounting medium containing DAPI nuclear stain. For RGC counting, a minimum of 16 non-overlapping images from the entire retina were captured at 200x magnification using a Keyence fluorescence microscope (Itasca, IL, USA), and RBPMS-positive cells were counted using ImageJ software[29, 31, 35].

Assessment of optic nerve degeneration: Optic nerve axonal degeneration was performed using a paraphenylenediamine (PPD) staining as described previously[31, 35] In brief, optic nerves were fixed overnight in PBS containing 3% glutaraldehyde/paraformaldehyde at 4°C. The optic nerves were rinsed twice for 10 minutes with 0.1M phosphate buffer and once with 0.1M sodium acetate buffer. Optic nerves were then dehydrated in graded ethanol concentrations. After embedding in resin, transverse semithin sections (1 µm) were cut and stained with 1% PPD for 10 minutes. Ten images without overlap were captured using confocal Leica microscope and counted manually using Image J software. The sum of total surviving axons counted approximately equaled to 10% of the total optic nerve cross-sectional area.

Transmission electron microscopy (TEM): To perform TEM imaging of the outflow pathway, 1 µm sections were processed for TEM analysis as described previously[35] The animals were perfusion-fixed with 4% PFA, and eyes were enucleated. The anterior chamber and optic nerve were dissected carefully and further fixed with 1.5% paraformaldehyde and 1.5% glutaraldehyde in 100 mM sodium cacodylate buffer (pH 7.2) for at least 24 hours. After aldehyde fixation, the eyes were incubated with 1% osmium tetroxide in 100 mM sodium cacodylate buffer, followed by ethanol dehydration and embedding of the samples in Eponate 812 resin. Sections (1 µm) were cut using an ultramicrotome (EM UC6; Leica) and stained with uranyl acetate and Reynolds lead citrate. Images were taken with a transmission electron microscope (JEM-1230; JEOL) equipped with a 2K × 2K CCD camera (USC1000; Gatan Inc.) at the University of Iowa Central Microscopy Research Facility.

Visual electrode potential: Mice were dark adapted for eight hours in a light-proof box. The dark-adopted animals were anaesthetized with an intraperitoneal injection of ketamine/xylazine mixture (100, 10 mg/kg, respectively). Pupils were dilated using a single drop of 0.5% tropicamide for five minutes. Visual evoked potentials (VEPs) were recorded using the Celeris D430 rodent ERG system (Diagnosys LLC, MA, USA) under maintained dark conditions as described previously[39]. Mice were positioned on a heated platform to regulate body temperature, and corneal moisture was maintained with 1-2% hydroxypropyl methylcellulose. Corneal electrodes integrated with stimulators were applied to lubricated corneas. Ground and reference needle electrodes were subcutaneously inserted into the tail, cheek, and subdermal region of the head, approximating the visual cortex. The midpoint between the ears served as the reference point. Although recording locations were not histologically verified, they were assumed to be consistent based on an impedance of 1.5-10 kΩ. Each eye was subjected to 100 flashes of 1 Hz, 0.05 cd s/m² white light delivered through corneal stimulators. Recordings were captured for 300 ms at a sampling rate of 2000 Hz. Two trials consisting of 50 sweeps were performed for each mouse. VEP data was filtered with

low and high frequency cutoffs of 1.25 Hz and 100 Hz, respectively. The amplitude of N1 was calculated as the distance from baseline to the most pronounced negative peak, while P1 amplitude was determined from the deepest negative peak to the subsequent positive peak. Latency was defined as the interval between stimulus onset and the initial response peak[40].

Axonal transport: To assess axonal anterograde transport mechanism, fluorescently labeled cholera toxin B (CTB) was injected intravitreally and its transport across ON to superior colliculus (SC) was tracked using fluorescence microscope as described previously[35]. We utilized 15-month old $Tg.CreMYOC^{Y437H}$ mice for CTB injections based on a previous study that indicated that axonal transportation impairment is more evident in old mice[41]. 7-weeks empty or Cre-injected $Tg.CreMYOC^{Y437H}$ mice were anesthetized using isoflurane and an intravitreal injection of 3 µL of 0.1% CTB (reconstituted in PBS) conjugated with either Alexa Fluor 555 or Alexa Fluor 480 (Invitrogen, Life Technologies, Grand Island, NY, USA) was performed. 48-hours later, mice were euthanized, and eyes and whole brains were fixed in 4% PFA at 4°C for 12 hours. Subsequently, tissues were washed with PBS, cryoprotected in a sucrose gradient (10-30%), embedded in OCT compound, and cryo-sectioned at a thickness of 10 µm. Axonal anterograde transport of CTB to ON and SC was directly visualized using fluorescence microscopy. Tissue sections were washed with PBS, mounted with a DAPI-containing mounting medium, and imaged using a Keyence fluorescence microscope[35].

Statistics: Prism 9.0 software (GraphPad, San Diego, CA, USA) was used for statical analyses. Data was represented as mean ± SEM. $P < 0.05$ will be considered statistically significant. The student's t-test was used to compare 2 groups. For comparison of different treatments, 2-way ANOVA was used followed by a Bonferroni post-hoc correction.

Results

Generation of an inducible mouse model of myocilin POAG: Using TARGATT site-specific knock-in strategy, we developed a Cre-inducible transgenic mice that expresses DsRed-tagged Y437H mutant of human *MYOC* (Referred as $Tg.CreMYOC^{Y437H}$). In normal conditions, mutant human *MYOC* is not expressed. Expression of Cre recombinase leads to removal of Stop cassette and DsRed-fused mutant *MYOC* is expressed (Figure 1A). A single copy of transgene is inserted into a preselected intergenic and transcriptionally active genomic locus (H11) that has been engineered with a docking site for stable and site-specific transgene integration. To confirm a site-specific knock-in of transgene at H11 site, we first performed PCR using primers specific to integration site (SI. 1A), which demonstrated a stable integration of transgene in one founder line. This founder was further bred with C57BL/6J mice and offspring were utilized for subsequent studies. For routine genotyping, primers specific to DsRed were utilized, which confirmed the presence of the transgene (SI. 1B).

Helper Ad5-Cre induces mutant myocilin selectively in the TM: We have previously shown that intravitreal injection of Ad5 exhibits specific tropism to mouse TM[31, 34, 42, 43,]

we therefore utilized Ad5 expressing Cre recombinase to selectively induce mutant myocilin in the TM of *Tg.CreMYOC[Y437H]* mice (SI. 2A-B). A single intravitreal injection of Ad5-empty or Ad5-Cre (2x 10^7 pfu/eye) was performed in 3-months old *Tg.CreMYOC[Y437H]* mice. Anterior segment cross sections demonstrated robust DsRed expression in mouse TM (SI. 2A) of 5-weeks Ad5-Cre-injected *Tg.CreMYOC[Y437H]* mice. No DsRed was detected in other ocular tissues including retina. Western blot analysis of iridocorneal angle, retina, and sclera clearly demonstrated that mutant myocilin is selectively induced in the TM tissue of 5-weeks Ad5-Cre-injected *Tg.CreMYOC[Y437H]* mice (SI. 2B). Since Ad5 tends to induce ocular inflammation[31, 34], we performed slit lamp imaging, which revealed moderate ocular inflammation after Ad5-Cre injections (SI. 2C). To reduce ocular inflammation, we next utilized helper Ad5 (HAd5), which lacks most of viral sequences except cis-acting elements essential for viral replication and packaging. HAd5 is known to exhibit minimal immunogenicity while maintaining robust tropism for tissue of interest[44]. A single intravitreal injection of HAd5 expressing Cre or empty cassette was performed in 3-months old *Tg.CreMYOC[Y437H]* mice. Slit-lamp imaging demonstrated no visible signs of ocular inflammation in HAd5-empty or Cre injected *Tg.CreMYOC[Y437H]* mice (Figure 1B). Next, we examined whether HAd5 expressing Cre exhibits functional activity in mouse TM using fluorescence reporter mTmG mice. These mice express tdTomato in all tissue; expression of Cre induces conversion of tdTomato to GFP [30]. We performed an intravitreal injection of HAd5-empty or Cre, and GFP/tdTomato expression was examined using confocal imaging of anterior segment cross-sections (Figure 1C). Compared to empty-injected mTmG mice, which only expressed tdTomato, Cre-injected mice exhibited GFP expression selectively in TM cells, and efficiency of conversion was nearly 100%. These data indicate that a single intravitreal injection of HAd5-cre is highly efficient in transducing the mouse TM.

Next, we investigated whether HAd5-cre induces mutant myocilin in the TM. A single intravitreal injection of HAd5-Cre or HAd5-empty (2x 10^7 pfu/eye) was performed in 2-months old *Tg.CreMYOC[Y437H]* mice and DsRed expression in various ocular tissues was evaluated (Figure 1D-F). Whole mount anterior segment imaging demonstrated mutant myocilin expression throughout the TM of Cre-injected *Tg.CreMYOC[Y437H]* mice (Figure 1D). Western blot analysis of iridocorneal angle, retina, choroid, and sclera demonstrated the presence of MYOC protein only in the iridocorneal angle tissues of HAd5-Cre induced *Tg.CreMYOC[Y437H]* mice (Figure 1E). MYOC protein was detected at ~75 KDa, which represents DsRed-tagged mutant MYOC (Figure 1E). Although endogenous myocilin was detected at 50KDa in all ocular tissues, mutant MYOC was only detected in iridocorneal angle tissues of *Tg.CreMYOC[Y437H]* mice and no mutant MYOC protein was detected in retina and sclera. Interestingly, we observed that expression of human mutant MYOC increased the level of endogenous myocilin protein in Cre-injected iridocorneal angle tissue lysates compared to control. QPCR analysis of anterior segment tissues using primers specific to mutant MYOC confirmed the presence of MYOC mRNA in Cre-induced *Tg.CreMYOC[Y437H]* mice (SI. 3). Confocal imaging of anterior segment cross-section revealed a robust and selective induction of mutant MYOC in the TM of *Tg.CreMYOC[Y437H]* mice (Figure 1F).

Since mutant *MYOC* is driven by CAG promoter, which can lead to overexpression of mutant *MYOC*, we next compared mRNA transcript of mutant *MYOC* with endogenous *Myoc* using RNA scope (Figure 1G). As expected, no mRNA transcript for mutant *MYOC* was detected in Cre⁻*Tg.CreMYOC^{Y437H}* mice while abundant endogenous *Myoc* was detected in the TM region. Importantly, Cre-induced *Tg.CreMYOC^{Y437H}* mice demonstrated the presence of mutant *MYOC* transcript selectively in the TM region. Interestingly, we observed upregulation of endogenous *Myoc* expression in Cre⁺*Tg.CreMYOC^{Y437H}* mice compared to Cre⁻*Tg.CreMYOC^{Y437H}* mice. The expression level of human mutant *MYOC* in Cre-injected eyes was similar to endogenous *Myoc* observed in control *Tg.CreMYOC^{Y437H}* mice. Together, these data established that HAd5-Cre selectively induces mutant myocilin in the TM of *Tg.CreMYOC^{Y437H}* mice at the level similar to endogenous *Myoc* without causing ocular inflammation.

Expression of mutant MYOC reduces TM outflow and elevates IOP significantly in HAd5-Cre-injected *Tg.CreMYOC^{Y437H}* mice. Three-month-old *Tg.CreMYOC^{Y437H}* mice were injected intravitreally with HAd5-empty or Cre, and IOPs were monitored weekly. Starting from 2-weeks of injection, Cre-treated mice *Tg.CreMYOC^{Y437H}* exhibited significantly higher and sustained IOP compared to empty-injected *Tg.CreMYOC^{Y437H}* mice (Figure 2A). An independent conscious (without anesthesia) IOP measurements confirmed a pronounced and significant IOP elevation in HAd5-Cre injected *Tg.CreMYOC^{Y437H}* mice (SI. 4). Measurement of outflow facility using the constant flow infusion method displayed significantly reduced outflow facility in 5-weeks HAd5-Cre injected *Tg.CreMYOC^{Y437H}* mice compared to control group (11.75 nL/min/mmHg in Cre⁺ vs. 22.32 nL/min/mmHg in Cre⁻ *Tg.CreMYOC^{Y437H}* mice) (Figure 2B). Together, these findings demonstrate that HAd5-Cre induces significant and sustained IOP elevation due to reduced aqueous humor outflow facility from the TM in *Tg.CreMYOC^{Y437H}* mice.

Mutant myocilin induces ultrastructural and biochemical changes in the TM. In POAG, increased outflow resistance is associated with ultrastructural and biochemical changes in the TM including increased extracellular matrix (ECM) deposition, actin cytoskeleton changes and induction of ER stress[8, 12, 37, 45]. We next investigated whether mutant myocilin leads to morphological changes in the TM of *Tg.CreMYOC^{Y437H}* mice. *H & E* staining of anterior segment from 5-weeks Cre-injected *Tg.CreMYOC^{Y437H}* mice demonstrated open angle and no obvious morphological changes in the anterior chamber structures (SI 5). We next performed transmission electron microscopy (TEM) to examine ultrastructural changes in the TM. Low magnification TEM analysis demonstrated that iridocorneal angle is open in both empty and Cre-injected *Tg.CreMYOC^{Y437H}* mice (Figure 3A). Higher magnification TEM images revealed loosely bound collagen fibers, ECM deposition, and disrupted TM integrity in Cre⁺ *Tg.CreMYOC^{Y437H}* mice compared to Cre⁻ *Tg.CreMYOC^{Y437H}* mice (Figure 3B). We further confirmed these findings using immunostaining. Anterior segments were immunostained with antibodies for fibronectin (FN) and actin (Figure 3C). Both FN and actin were increased and co-localized with MYOC in the TM of Cre-injected *Tg.CreMYOC^{Y437H}* mice. Alpha smooth muscle actin (SMA) was also increased and strongly co-localized with mutant MYOC in Cre-injected *Tg.CreMYOC^{Y437H}* mice (Figure 3D). Previous studies have shown that mutant myocilin expression in TM is associated with ER stress[27, 46-49]. We therefore examined whether

mutant myocilin expression induces ER stress markers in the TM of Cre-induced *Tg.CreMYOC*^{Y437H} mice (Figure 3E). Immunostaining (SI 6) and Western blot analysis (Figure 3E) demonstrated increased ER chaperones (GRP78), ATF4 and CHOP selectively in the TM of Cre-injected *Tg.CreMYOC*^{Y437H} mice. Together, these findings indicate that expression of mutant myocilin induces ultrastructural and biochemical changes in the TM leading to its dysfunction and IOP elevation in *Tg.CreMYOC*^{Y437H} mice.

Mutant MYOC-induced sustained IOP elevation leads to functional and structural loss of RGCs: We next evaluated whether sustained IOP elevation induced by mutant MYOC is sufficient to cause functional and structural loss of RGCs. To evaluate functional loss of RGCs, we performed PERG at 5, 10, and 15 weeks of injection (Figure 4A-C). Representative PERG graphs and its analysis demonstrated no significant effect on PERG at 5-weeks, but significantly reduced PERG amplitudes and increased latencies were observed at 10 and 15 weeks Cre-injected *Tg.CreMYOC*^{Y437H} mice (Figure 4A-C). To determine the structural loss of RGCs, we next performed whole mount retina staining with RBPMS antibody (Figure 4D-E). As shown in representative RBPMS images, 15-weeks Cre-injected *Tg.CreMYOC*^{Y437H} mice exhibited moderate loss of RGCs in the periphery of retina (Figure 4E). RGC counting further confirmed a significantly reduced RGCs in the periphery of Cre-injected *Tg.CreMYOC*^{Y437H} mice compared to controls at 15 weeks of injection (Figure 4E). Overall, there was 33% loss of RGCs in Cre-injected *Tg.CreMYOC*^{Y437H} mice compared to controls. We did not observe RGC loss at 10-weeks of Cre-injection in *Tg.CreMYOC*^{Y437H} mice (SI. 7). These data indicate that sustained IOP elevation induced by mutant myocilin leads to RGC functional and structural loss of RGCs in *Tg.CreMYOC*^{Y437H} mice.

Mutant MYOC-induced sustained IOP elevation leads to optic nerve degeneration in *Tg.CreMYOC*^{Y437H} mice. We investigated whether sustained IOP elevation induced by mutant myocilin leads to optic nerve degeneration using PPD staining (Figure 5A). Representative images of PPD stained ON from Cre-injected *Tg.CreMYOC*^{Y437H} mice demonstrated optic nerve degeneration as evident from darkly stained axons, active gliosis, and glial scar formation. Approximately 20% and 45% axonal loss was observed in 10 and 15-weeks of Cre-injected mice respectively compared to empty-injected *Tg.CreMYOC*^{Y437H} mice (Figure 5B). To further confirm these findings, we next performed immunostaining for GFAP on retinal cross-section of Cre-injected *Tg.CreMYOC*^{Y437H} mice (Figure 5C). Immunostaining for GFAP revealed a prominent increase in GFAP reactivity in ONH region suggesting axonal injury. Axonal degeneration was associated with decreased neuronal marker, Tuj1 and increased mitochondrial accumulation (increased TOM20) in the ONH region (Figure 5D). These data establish that sustained IOP elevation induces ON degeneration and gliosis in the ONH of Cre-injected *Tg.CreMYOC*^{Y437H} mice.

Impaired axonal transport at the ONH precedes neuronal loss in *Tg.CreMYOC*^{Y437H} mice. Since *Tg.CreMYOC*^{Y437H} mice exhibit well-defined timelines for RGC axon loss, we further sought to understand the early pathogenic events preceding neuronal loss. RGC loss and ON degeneration studies suggested that axonal changes at ONH may be the first site of damage in ocular hypertensive *Tg.CreMYOC*^{Y437H} mice. We therefore

hypothesize that IOP-induced neurodegenerative changes in the ONH precede neuronal loss. Since we did not observe RGC structural loss until 15-weeks of Cre-injection, we examined whether neurodegenerative changes occur in ONH the prior to RGC loss at 7-weeks of Cre-injection. To test this, 15-month *Tg.CreMYOC[Y437H]* mice were injected with HAd5-empty or Cre and IOPs were monitored. IOP measurements confirmed a significant IOP elevation in 6-weeks Cre-injected *Tg.CreMYOC[Y437H]* mice (Figure 6A). PERG, which measures the function of RGC soma revealed no significant functional loss of RGC soma in 5-weeks Cre-injected *Tg.CreMYOC[Y437H]* mice (Figure 4A-C). Whole mount RBPMS staining revealed no significant loss of RGCs in 10-weeks Cre-injected *Tg.CreMYOC[Y437H]* mice (SI 7). Interestingly, VEP, which measures post-retinal function of visual system including ON and visual centers of the brain demonstrated significant loss of VEP amplitudes at 7-weeks suggesting axonal dysfunction in *Tg.CreMYOC[Y437H]* mice (Figure 6B). To explore whether axonal dysfunction occurs at early stages of neuronal loss due to sustained IOP, we examined anterograde transport of fluorescently labeled CTB as described previously[35]. 7-weeks empty or Cre-injected *Tg.CreMYOC[Y437H]* mice were intravitreally injected with green fluorescently tagged CTB dye (Figure 6C). 48-hours after injections, anterograde transport of CTB through optic nerve and superior colliculus (SC) was monitored via fluorescence microscope. As expected, CTB was transported to SC in HAd5-empty injected *Tg.CreMYOC[Y437H]* mice. However, CTB transportation was completely blocked at the ONH and no CTB was observed in SC of Cre-injected *Tg.CreMYOC[Y437H]* mice. CTB fluorescence measurement in SC demonstrated a significant (~72%) loss of CTB in SC of Cre-injected *Tg.CreMYOC[Y437H]* mice. Altogether, these data indicate that mutant MYOC-induced ocular hypertension impairs axonal transport blockage at the ONH prior to RGC loss in *Tg.CreMYOC[Y437H]* mice.

Loss of microtubules and neurofilaments are associated with impaired axonal transport at the ONH of *Tg.CreMYOC[Y437H]* mice: To further understand the molecular insights into impaired axonal transport, we analyzed CTB transport in cross-section of retina along with ON. CTB was blocked in the proximal region of ON in Cre-injected *Tg.CreMYOC[Y437H]* mice (Figure 7A). TEM analysis of ON at 8-week of Cre-injection demonstrated ultrastructural changes including loss of neurofilament and microtubule structures (Figure 7B-C). The presence of myelin around axons in Cre-injected mice indicated intact axons but accumulation of intracellular materials and loss of microtubules and neurofilament suggest that these ultrastructural changes in ON axons precede axonal degeneration. To confirm, TEM findings, we performed immunostaining on retinal cross-section obtained from 7-weeks injected *Tg.CreMYOC[Y437H]* mice (Figure 7D-G). Cre-injected *Tg.CreMYOC[Y437H]* mice exhibited dramatic increase in GFAP reactivity and decreased microtubules (Tuj1) and neurofilament (NF) loss in the ONH region of Cre-injected *Tg.CreMYOC[Y437H]* mice compared to control *Tg.CreMYOC[Y437H]* mice. Importantly, we observed increased mitochondrial accumulation in RGC soma (Figure 7G) and axons at ONH region in 12-weeks Cre-injected *Tg.CreMYOC[Y437H]* mice (Figure 5D). Together, our studies suggest that loss of microtubules and neurofilament, which are required for axonal transport is associated with impaired axonal transport leading to accumulation of defective organelles and causing axonal dysfunction.

Discussion

In the present study, we describe the development of novel Cre-inducible mouse model of MYOC-associated POAG. We further report that HAd5 drives the TM-specific expression of mutant MYOC leading to TM dysfunction and IOP elevation. Importantly, a single intravitreal injection of HAd5-Cre is sufficient to induce sustained IOP elevation, which leads to glaucomatous neurodegeneration in a highly predictable manner. Induction of mutant MYOC via Cre expression in adult mice allowed us to study early events of glaucomatous neurodegeneration. Specifically, we observed that sustained IOP elevation impairs axonal transport at the ONH prior to axonal degeneration and RGC soma loss. Interestingly, our studies suggest that loss of key neuronal cytoskeleton proteins, microtubules and neurofilaments are likely to be responsible for impaired axonal transport resulting in an increased mitochondrial accumulation at the ONH and inducing nueornal dysfunction. These mechanistic insights into early events of axonal degeneration provide valuable targets to develop neuroprotective therapies for glaucoma.

Previously, we have developed a transgenic mouse model (*Tg-MYOC^Y437H*) by random insertion of mutant MYOC[27]. This transgenic mouse model exhibited robust glaucoma phenotypes on mixed background, and we discovered the pathological role of chronic ER stress in the pathophysiology of glaucomatous TM damage using this model[26, 27, 31]. However, these mice exhibited mild glaucoma phenotypes with subsequent breeding onto pure C57BL/6J background [34, 50, 51]. We reasoned this to silencing of transgene likely due to the toxic nature of mutant MYOC. In addition, our previous transgenic mouse model presented several limitations including lack of robust glaucomatous neurodegeneration on pure C57 background, unknown copies of mutant MYOC and lack of specific antibodies to distinguish human versus mouse myocilin. Development of *Tg.CreMYOC^Y437H* mice has overcome these key limitations of previously reported transgenic mouse model of MYOC-glaucoma. These include targeted insertion of mutant MYOC in the preselected intergenic and transcriptionally active genomic locus (H11). This allows stable and site-specific transgene integration of single copy of mutant MYOC. Also, *Tg.CreMYOC^Y437H* mice are developed on pure C57BL/6J background. DsRed is fused with mutant *MYOC*, which allowed us to distinguish mutant *MYOC* from endogenous *Myoc*. Indeed, Western blot and RNA scope analysis demonstrated the expression levels of mutant MYOC, which is similar to endogenous *Myoc*. We utilized RNA scope to probe both endogenous *Myoc* and mutant MYOC using specific probes, and our results clearly demonstrate that *Tg.CreMYOC^Y437H* mice express mutant *MYOC* at a similar level to endogenous *Myoc*. This was further supported by Western blot analysis, which demonstrated protein levels of human mutant MYOC fused with DsRed is less than endogenous mouse myocilin in Cre-treated *Tg.CreMYOC^Y437H* mice and it is comparable to endogenous MYOC in untreated *Tg.CreMYOC^Y437H* mice. These data indicates that glaucoma phenotypes in *Tg.CreMYOC^Y437H* mice are not a result of overexpression of mutant MYOC. Both RNA scope and Western blot clearly demonstrated that expression of mutant *MYOC* induced endogenous *Myoc*. This is consistent with previous findings that TM cells induce MYOC in response to various insults including glucocorticoid treatment [24, 52, 53]. Although the exact function of endogenous myocilin is unclear, previous studies have suggested its role as a stress-response pathway [54].

The Cre-*LoxP* recombinase system is an effective and widely used experimental tool to investigate gene of interest in tissue/cell and/or time-specific manner[55]. Cre-inducible mice are often crossed with tissue-specific Cre mouse lines to induce Cre-expression. In our study, we employed viral vector to induce Cre expression in TM. Crossing *Tg.CreMYOC^Y437H* mice with TM-specific Cre mouse line would be more ideal to induce mutant MYOC in TM. However, there are no TM-specific Cre mouse lines available. The global Cre mouse lines would express mutant MYOC in other cell types including in ciliary body and retina, which can directly result in cell death of ciliary epithelium or RGCs. Previous studies have shown that mutant MYOC is toxic to cells [27] and degenerations of the non-pigmented epithelium in the ciliary body was observed in older *Tg-MYOC^Y437H* mice [51]. We also observed that induction of mutant MYOC selectively in RGCs via an intravitreal injection of AAV2-Cre leads to severe RGC loss within 3 weeks of injection in *Tg.CreMYOC^Y437H* mice (Data not shown).

We and others have shown that Ad5 exhibits optimum tropism to TM without affecting other ocular cells but it does produce ocular inflammation, which can itself produce IOP elevation [32, 43]. Since HAd5 produces less inflammation, we utilized HAd5-Cre to express Cre selectively in the TM. DsRed imaging and Western blot data clearly demonstrated that a single intravitreal injection of HAd5-Cre leads to robust Cre activity and induces mutant MYOC in the TM without causing any discernable ocular inflammation. HAd5-Cre induced mutant MYOC expression throughout the TM. Importantly, no Cre activity was observed in retina including RGCs. We choose intravitreal injection based on our recent studies demonstrating that intravitreal injections provide a depot for optimum transduction of viral vectors to the TM [34]. It is likely that intracameral injection of HAd5-Cre will produce similar Cre activity.

Our studies show that *Tg.CreMYOC^Y437H* mice exhibit all features of human POAG including IOP elevation and reduced outflow facility due to TM dysfunction. Like POAG, we observed ultrastructural and biochemical changes associated with TM dysfunction. Importantly, sustained IOP elevation leads to progressive axonal degeneration, which is associated with RGC loss. Contrary to other existing glaucoma models, most of *Tg.CreMYOC^Y437H* mice develop ocular hypertension and exhibit neuronal loss. Nearly all eyes treated with Cre developed IOP elevation and glaucomatous neurodegeneration. Glaucoma phenotypes in Cre-inducible *Tg.CreMYOC^Y437H* mice are highly predictable, and the exact timeline of each phenotype can be easily tracked. Mutant MYOC is induced within a week, elevating IOP starting from 2-weeks of Cre injection, which is followed by axonal transport deficits at 7 weeks. Axonal loss is observed at 10-weeks and RGC soma loss is observed at 15-weeks (Figure 8). These features make *Tg.CreMYOC^Y437H* mouse model highly attractive to study both TM dysfunction and neuronal loss in glaucoma. Although we did not investigate long-term effects of mutant MYOC on TM dysfunction and neuronal loss, it is likely that IOP elevation will sustain throughout the life of mice resulting into severe neuronal loss as observed in POAG.

Our study further demonstrates that sustained IOP elevation impairs axonal transport in the ONH prior to axonal degeneration and RGC structural loss. Axonal transport deficits are considered an important pathological feature of glaucomatous neurodegeneration[56,

[57]. Previous studies have shown that impaired axonal transport is associated with glaucomatous neurodegeneration [41, 58, 59]. We have recently demonstrated that axonal transport persists during initial stages of axonal degeneration in mouse model of glucocorticoid-induced glaucoma[35]. In addition, several other studies have shown that impaired axonal transport in ONH region occurs in an inducible model of ocular hypertension[35, 59-62]. Altogether, these studies support our findings that initial axonal transport block occurs at the ONH in glaucoma. It is poorly understood how ocular hypertension impairs axonal transport at the ONH. In 7-weeks of Cre-injected *Tg.CreMYOC^{Y437H}* mice, which demonstrated complete blockage of axonal transport, we observed no significant loss of axon as evident from intact myelin. However, TEM analysis demonstrated intracellular ultrastructural changes including swollen axons, increased mitochondrial accumulation and gliosis indicating axonal dysfunction. Interestingly, we observed a significant loss of VEP, which measures post-retinal visual pathways (ON and visual center of the brain)[63] while RGC function was not significantly affected until 10-weeks of injection. Since axonal transport is essential for RGC soma survival including transport of organelles, synaptic components, vesicles, and neurotrophic factors, it is possible that impaired axonal transport leads to RGC axonal degeneration and soma loss. Consistent with this, we observed increased mitochondrial accumulation in RGC soma and ON axons in 12-weeks Cre-injected *Tg.CreMYOC^{Y437H}* mice. Although it is not clear how sustained IOP elevation impairs axonal transport, TEM and immunostaining demonstrated dramatic loss of cytoskeleton proteins including microtubules and neurofilament associated with impaired axonal transport.

Both microtubules and neurofilaments are key cytoskeleton proteins required for axonal transport and loss of these proteins is associated with glaucomatous neurodegeneration[61, 64, 65]. Here, we propose that loss of microtubule and neurofilament at the ONH due to chronic IOP elevation is likely to responsible for impaired axonal transport. This is the first study showing that IOP elevation induces loss of microtubules impairing axonal transport in the ONH prior to neuronal loss. Since both microtubules and neurofilament assembly is regulated by autophagy [66], it is possible that chronic IOP elevation leads to compromised autophagy, which results into loss of neurofilament and microtubule assembly accumulating defective organelles in the ONH region. Consistent with this, several studies have suggested that chronic IOP elevation leads to compromised autophagy in the RGC axons[67, 68]. Alternatively, loss of microtubules, which facilitate formation and transport of autophagic vesicles carrying defective organelles may contribute to accumulation of autophagic vesicles in the ONH region including defective mitochondria as evident in our studies. Further understanding of these early events of axonal degeneration provides a therapeutic window for regeneration strategies to support dying axons.

In summary, we developed a Cre-inducible mouse model of POAG that faithfully replicates all features of human POAG including IOP elevation due to TM dysfunction and IOP-dependent glaucomatous neurodegeneration. Strikingly, our studies demonstrate that sustained IOP impairs axonal transport likely due to loss of cytoskeleton proteins in the ONH prior to axonal degeneration and RGC soma loss (Figure 8). These early pathological events lead to accumulation of defective organelles including mitochondria in ONH region and axonal degeneration. Targeting these early neuronal changes may provide a novel neuroprotective strategy. We expect that *Tg.CreMYOC^{Y437H}* mouse model will be a valuable tool to study the pathophysiology of glaucomatous damage to TM and RGC axons and provide new targets for the treatment.

Figure 8: Ocular hypertension disrupts axonal transport in the optic nerve head resulting in neurodegeneration.

Authors Contributions: GSZ, and BK designed research studies. BK, RK, and LL, performed experiments and analyzed data. JCM, YS, PM, WC and DSK assisted in some experiments. BK and GSZ wrote the manuscript. All authors discussed the results and implications and commented on the manuscript at all stages.

Acknowledgments. These studies were supported by the National Institutes of Health (EY034333 and EY026177). The authors acknowledge support from NIH grant EY034238 and from an unrestricted grant from Research to Prevent Blindness to the Gavin Herbert Eye Institute at the University of California, Irvine.

Data Availability Statement: All the datasets used and/or analyzed in the present study is available from the corresponding author on reasonable request.

Figures:

Figure 1. HAd5-Cre recombinase induces mutant MYOC in mouse TM. (**A**) Design Strategy: *Tg.CreMYOC*^{Y437H} mice were engineered using a TARGATT gene knock in strategy in which DsRed-tagged human Y437H mutant *MYOC* was inserted in a transcriptionally active genomic locus (H11). A stop cassette prevents the expression of mutant MYOC-DsRed fusion protein until Cre recombinase is introduced. Following Cre expression, the stop cassette is excised allowing the expression of the mutant *MYOC* fused with DsRed only in targeted cells. (**B**) Representative slit lamp images showing that no obvious ocular inflammation is associated with either HAd5-empty or Cre-injected *Tg.CreMYOC*^{Y437H} mice (n=6). (**C**) HAd5-Cre was injected intravitreally (2 x10^7 pfu/eye) in mT/mG fluorescence-based reporter mice and the conversion from tdTomato to GFP was examined 1-week post-injection using confocal microscopy. (n=4). (**D-F**) *Tg.CreMYOC*^{Y437H} mice were injected intravitreally with HAd5-empty or Cre and MYOC induction was examined in the TM using (**D**) Confocal imaging of DsRed protein in whole mount anterior segment, (**E**) Western blot analysis of various ocular tissues using MYOC antibody showing the presence of human mutant myocilin in anterior segment tissues of Cre-injected eyes; AS-anterior segment; M-empty lane; CS-choroid and sclera, and (**F**) Confocal imaging of DsRed protein in anterior segment cross-section in Cre- and Cre+ *Tg.CreMYOC*^{Y437H} mice (n=4). (**G**) RNAscope analysis of *MYOC* (green) and *Myoc* (pink) transcripts in the anterior segment cross-section of Cre- and Cre+ *Tg.CreMYOC*^{Y437H} mice (n=3). Last panel show DsRed protein expression in the same slide. Note that DsRed florescence may be less compared other images due to fluorescence quenching during sample processing (TM, trabecular meshwork; CB, ciliary body; SC, Schlemm's canal). Arrow shows TM.

Figure 2: Intravitreal administration of HAd5-Cre elevates IOP and reduces outflow facility in *Tg.CreMYOC^{Y437H}* mice. 3-months old *Tg.CreMYOC^{Y437H}* mice received a single intravitreal injection of either HAd5-Empty or Cre in both eyes. (**A**) weekly IOP measurements demonstrated significant and sustained IOP elevation in Cre-injected *Tg.CreMYOC^{Y437H}* mice compared to HAd5-Empty injected mice. (n = 14 in Empty and n=18 in Cre-injected group, analyzed by 2-WAY ANOVA with multiple comparisons, ****P <0.0001). (**B**) Outflow facility measurements showed a significant reduction in outflow facility at 5-weeks Cre-injected *Tg.CreMYOC^{Y437H}* mice (n=8) compared to HAd5-Empty-injected mice (n=9) (unpaired t-test, two-tailed, mean ± SEM **P<0.0072).

Figure 3. Mutant MYOC-induced ocular hypertension is associated with ultrastructural and biochemical changes in the TM. Representative low (**A**) and high (**B**) magnification TEM images of 8-weeks HAd5-Empty and Cre-injected *Tg.CreMYOC*^{Y437H} mice showing the presence of loosely bound collagen fibers, ECM deposition and loss of TM integrity in juxtacanalicular connective tissue (JCT) region of Cre-injected *Tg.CreMYOC*^{Y437H} mice (n=4 in each group). (TM, trabecular meshwork; CB, ciliary body; SC, Schlemm's canal; CL, Collagen fibers; ECM, Extra cellular matrix). (**C-D**) Representative images of immunostaining for DsRed-MYOC with fibronectin and actin (**C**) or with alpha smooth muscle actin (SMA) (**D**) in HAd5-Empty (top panel) and ocular hypertensive HAd5-Cre (bottom panel) injected *Tg.CreMYOC*^{Y437H} mice. Mutant MYOC induced fibronectin, actin and SMA protein levels in mouse TM of Cre-injected *Tg.CreMYOC*^{Y437H} mice. (n=3). (**E**) Western blot analysis of iridocorneal angle tissue lysates showing that mutant MYOC induces ER stress in Cre-injected *Tg.CreMYOC*^{Y437H} mice. (n=3). E-HAd5-empty; AS-anterior segment.

Figure 4: Sustained IOP elevation leads to functional and structural loss of RGCs in Cre-injected *Tg.CreMYOC^Y437H* mice: 3 to 6-months old *Tg.CreMYOC^Y437H* mice were injected intravitreally with HAd5-Empty or HAd5-Cre in both eyes, and IOP was monitored weekly to ensure IOP elevation. PERG was performed at 5, 10 and 15 weeks to assess the function of RGCs. A representative PERG graph (**A**) and its analysis (**B-C**) demonstrated significantly reduced PERG amplitude (**B**) and increased latency (**C**) starting from 10-weeks of Cre injections indicating functional loss of RGCs in Cre-injected *Tg.CreMYOC^Y437H* mice**.** (n = 7 in HAd5-Empty and n=6 in HAd5-Cre), 2-WAY ANOVA with multiple comparisons (****P <0.0001). (**D-E**) RGCs loss was further analyzed using whole mount retina staining with RBPMS antibody. Representative image of RBPMS staining of different regions of retina (**D**) and its analysis (**E**) revealed a significant loss (33%) of RGCs in 15-weeks Cre-injected *Tg.CreMYOC^Y437H* mice compared to HAd5-Empty injected mice. (n = 7 HAd5-Empty and 6 in HAd5-Cre), 2-WAY ANOVA with multiple comparisons (****P<0.00001).

Figure 5: Sustained IOP elevation leads to optic nerve degeneration in Cre-injected *Tg.CreMYOC^{Y437H}* mice. Optic nerves were subjected to PPD staining to assess optic nerve degeneration. (**A**) Representative images of PPD-stained optic nerves showing mild axonal degeneration as evident from darkly stained axons (yellow arrow) and the presence of glial scar formation (blue arrow) in Cre-injected *Tg.CreMYOC^{Y437H}* mice and (**B**) the mean axonal counts showing a significant loss of ON axons (20% at 10-weeks and 45% at 15-weeks) in Cre-induced *Tg.CreMYOC^{Y437H}* mice. (n = 5 in Empty and n = 7 at 10 weeks and n=6 in Empty and n=6 at 15 weeks, unpaired t-test, two-tailed, ****$P <$ 0.0001). (**C**) Immunostaining for GFAP demonstrated gliosis in the ONH region of Cre-injected *Tg.CreMYOC^{Y437H}* mice (n=3). (**D**) Immunostaining for TOM20 and microtubule demonstrated increased mitochondrial accumulation in RGC axons in ON of Cre-injected *Tg.CreMYOC^{Y437H}* mice (n=3).

Figure 6: Anterograde transport deficits precede RGC degeneration in _Tg.CreMYOC^{Y437H}_ mice: 15-months old _Tg.CreMYOC^{Y437H}_ mice were injected intravitreally with HAd5-Empty or Cre and glaucoma phenotypes and anterograde axonal transport mechanisms were investigated. **(A)** IOP measurement revealed sustained and significant IOP elevation at 6-weeks of injection (n=6; unpaired t-test; P=0.0058). **(B)** VEP measurements demonstrated a significant loss of post-retinal visual pathway function in 7-weeks Cre-injected _Tg.CreMYOC^{Y437H}_ mice (n=5; unpaired t-test; P=0.0374). (**C-D**) Mutant MYOC-induced ocular hypertension leads to axonal transport deficits in Cre-injected _Tg.CreMYOC^{Y437H}_ mice. Representative images of CTB fluorescence (**C**) and its analysis (**D**) in 7-weeks HAd5-Empty or Cre-injected _Tg.CreMYOC^{Y437H}_ mice. Empty-injected _Tg.CreMYOC^{Y437H}_ mice exhibited an uninterrupted transport of CTB along the entire length of optic nerve to the SC. However, CTB transport was blocked significantly at the ONH region and no CTB was detected in SC of 7-weeks Cre-injected _Tg.CreMYOC^{Y437H}_ mice (n = 6 in each group).

Figure 7: Reduced microtubules and neurofilament are associated with axonal transport deficits in the ONH region. (**A**) Whole eyes along with ON from above CTB-injected *Tg.CreMYOC^Y437H* mice were sectioned to image CTB transport along ON. Retinal cross-sections demonstrated the blockage of CTB transport in proximal region of ON in Cre-injected *Tg.CreMYOC^Y437H* mice. (n=3). (**B**) Low and (**C**) high magnification TEM analysis of optic nerves Empty or Cre-injected *Tg.CreMYOC^Y437H* mice, which demonstrated cytoskeleton degeneration, organelle accumulation and the presence of glial scar prior to RGC degeneration in 7-weeks Cre-injected *Tg.CreMYOC^Y437H* mice (n=4). (**D-F**) Immunohistochemical analysis of cytoskeleton marker, neurofilament (**D**),

microtubules using Tuj1 antibody (**E**) and gliosis using GFAP antibody (**F**) demonstrated reduced microtubules and neurofilament and increased GFAP in the ONH region of Cre-injected *Tg.CreMYOC^{Y437H}* mice. (**G**) Immunostaining for TOM20 demonstrated increased mitochondrial accumulation in the RGCs of 12-weeks Cre-injected *Tg.CreMYOC^{Y437H}* mice (n=3). (# organelle accumulation; + glial scar). (**Neurofilament; ## microtubule).

References:

1.   Quigley HA, Broman AT. The number of people with glaucoma worldwide in 2010 and 2020. Br J Ophthalmol. 2006;90(3):262-7. doi: 10.1136/bjo.2005.081224. PubMed PMID: 16488940; PMCID: PMC1856963.
2.   Quigley HA. Neuronal death in glaucoma. Prog Retin Eye Res. 1999;18(1):39-57. doi: 10.1016/s1350-9462(98)00014-7. PubMed PMID: 9920498.
3.   Tham YC, Li X, Wong TY, Quigley HA, Aung T, Cheng CY. Global prevalence of glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-analysis. Ophthalmology. 2014;121(11):2081-90. Epub 20140626. doi: 10.1016/j.ophtha.2014.05.013. PubMed PMID: 24974815.
4.   Rosenthal J, Leske MC. Open-angle glaucoma risk factors applied to clinical area. J Am Optom Assoc. 1980;51(11):1017-24. PubMed PMID: 7440868.
5.   Tamm ER, Braunger BM, Fuchshofer R. Intraocular Pressure and the Mechanisms Involved in Resistance of the Aqueous Humor Flow in the Trabecular Meshwork Outflow Pathways. Prog Mol Biol Transl Sci. 2015;134:301-14. Epub 20150709. doi: 10.1016/bs.pmbts.2015.06.007. PubMed PMID: 26310162.
6.   Kwon YH, Fingert JH, Kuehn MH, Alward WL. Primary open-angle glaucoma. N Engl J Med. 2009;360(11):1113-24. doi: 10.1056/NEJMra0804630. PubMed PMID: 19279343; PMCID: PMC3700399.
7.   Weinreb RN, Leung CK, Crowston JG, Medeiros FA, Friedman DS, Wiggs JL, Martin KR. Primary open-angle glaucoma. Nat Rev Dis Primers. 2016;2:16067. Epub 20160922. doi: 10.1038/nrdp.2016.67. PubMed PMID: 27654570.
8.   Hoare MJ, Grierson I, Brotchie D, Pollock N, Cracknell K, Clark AF. Cross-linked actin networks (CLANs) in the trabecular meshwork of the normal and glaucomatous human eye in situ. Invest Ophthalmol Vis Sci. 2009;50(3):1255-63. Epub 2008/10/28. doi: 10.1167/iovs.08-2706. PubMed PMID: 18952927.
9.   Kasetti RB, Maddineni P, Millar JC, Clark AF, Zode GS. Increased synthesis and deposition of extracellular matrix proteins leads to endoplasmic reticulum stress in the trabecular meshwork. Sci Rep. 2017;7(1):14951. Epub 20171102. doi: 10.1038/s41598-017-14938-0. PubMed PMID: 29097767; PMCID: PMC5668243.
10.   Vranka JA, Kelley MJ, Acott TS, Keller KE. Extracellular matrix in the trabecular meshwork: intraocular pressure regulation and dysregulation in glaucoma. Exp Eye Res. 2015;133:112-25. doi: 10.1016/j.exer.2014.07.014. PubMed PMID: 25819459; PMCID: PMC4379427.
11.   Raghunathan VK, Benoit J, Kasetti R, Zode G, Salemi M, Phinney BS, Keller KE, Staverosky JA, Murphy CJ, Acott T, Vranka J. Glaucomatous cell derived matrices differentially modulate non-glaucomatous trabecular meshwork cellular behavior. Acta Biomater. 2018;71:444-59. Epub 2018/03/11. doi: 10.1016/j.actbio.2018.02.037. PubMed PMID: 29524673; PMCID: PMC5899949.
12.   Keller KE, Peters DM. Pathogenesis of glaucoma: Extracellular matrix dysfunction in the trabecular meshwork-A review. Clin Exp Ophthalmol. 2022;50(2):163-82. Epub 20220117. doi: 10.1111/ceo.14027. PubMed PMID: 35037377; PMCID: PMC9199435.
13.   Chang EE, Goldberg JL. Glaucoma 2.0: neuroprotection, neuroregeneration, neuroenhancement. Ophthalmology. 2012;119(5):979-86. Epub 20120218. doi: 10.1016/j.ophtha.2011.11.003. PubMed PMID: 22349567; PMCID: PMC3343191.
14.   Johnson TV, Tomarev SI. Rodent models of glaucoma. Brain Res Bull. 2010;81(2-3):349-58. Epub 20090418. doi: 10.1016/j.brainresbull.2009.04.004. PubMed PMID: 19379796; PMCID: PMC2830899.
15.   Pang IH, Clark AF. Inducible rodent models of glaucoma. Prog Retin Eye Res. 2020;75:100799. Epub 20190923. doi: 10.1016/j.preteyeres.2019.100799. PubMed PMID: 31557521; PMCID: PMC7085984.

16.   Aihara M, Lindsey JD, Weinreb RN. Experimental mouse ocular hypertension: establishment of the model. Invest Ophthalmol Vis Sci. 2003;44(10):4314-20. doi: 10.1167/iovs.03-0137. PubMed PMID: 14507875.

17.   Mabuchi F, Aihara M, Mackey MR, Lindsey JD, Weinreb RN. Optic nerve damage in experimental mouse ocular hypertension. Invest Ophthalmol Vis Sci. 2003;44(10):4321-30. doi: 10.1167/iovs.03-0138. PubMed PMID: 14507876.

18.   Sappington RM, Carlson BJ, Crish SD, Calkins DJ. The microbead occlusion model: a paradigm for induced ocular hypertension in rats and mice. Invest Ophthalmol Vis Sci. 2010;51(1):207-16. Epub 20091022. doi: 10.1167/iovs.09-3947. PubMed PMID: 19850836; PMCID: PMC2869054.

19.   Zhang J, Li L, Huang H, Fang F, Webber HC, Zhuang P, Liu L, Dalal R, Tang PH, Mahajan VB, Sun Y, Li S, Zhang M, Goldberg JL, Hu Y. Silicone oil-induced ocular hypertension and glaucomatous neurodegeneration in mouse. Elife. 2019;8. Epub 20190515. doi: 10.7554/eLife.45881. PubMed PMID: 31090540; PMCID: PMC6533060.

20.   John SW, Smith RS, Savinova OV, Hawes NL, Chang B, Turnbull D, Davisson M, Roderick TH, Heckenlively JR. Essential iris atrophy, pigment dispersion, and glaucoma in DBA/2J mice. Invest Ophthalmol Vis Sci. 1998;39(6):951-62. PubMed PMID: 9579474.

21.   Anderson MG, Smith RS, Hawes NL, Zabaleta A, Chang B, Wiggs JL, John SW. Mutations in genes encoding melanosomal proteins cause pigmentary glaucoma in DBA/2J mice. Nat Genet. 2002;30(1):81-5. Epub 20011217. doi: 10.1038/ng794. PubMed PMID: 11743578.

22.   Stone EM, Fingert JH, Alward WL, Nguyen TD, Polansky JR, Sunden SL, Nishimura D, Clark AF, Nystuen A, Nichols BE, Mackey DA, Ritch R, Kalenak JW, Craven ER, Sheffield VC. Identification of a gene that causes primary open angle glaucoma. Science. 1997;275(5300):668-70. doi: 10.1126/science.275.5300.668. PubMed PMID: 9005853.

23.   Libby RT, Gould DB, Anderson MG, John SW. Complex genetics of glaucoma susceptibility. Annu Rev Genomics Hum Genet. 2005;6:15-44. doi: 10.1146/annurev.genom.6.080604.162209. PubMed PMID: 16124852.

24.   Fingert JH, Stone EM, Sheffield VC, Alward WL. Myocilin glaucoma. Surv Ophthalmol. 2002;47(6):547-61. doi: 10.1016/s0039-6257(02)00353-3. PubMed PMID: 12504739.

25.   Kasetti RB, Maddineni P, Kodati B, Nagarajan B, Yacoub S. Astragaloside IV Attenuates Ocular Hypertension in a Mouse Model of TGFβ2 Induced Primary Open Angle Glaucoma. Int J Mol Sci. 2021;22(22). Epub 20211119. doi: 10.3390/ijms222212508. PubMed PMID: 34830390; PMCID: PMC8619727.

26.   Zode GS, Bugge KE, Mohan K, Grozdanic SD, Peters JC, Koehn DR, Anderson MG, Kardon RH, Stone EM, Sheffield VC. Topical ocular sodium 4-phenylbutyrate rescues glaucoma in a myocilin mouse model of primary open-angle glaucoma. Invest Ophthalmol Vis Sci. 2012;53(3):1557-65. Epub 20120321. doi: 10.1167/iovs.11-8837. PubMed PMID: 22328638; PMCID: PMC3339917.

27.   Zode GS, Kuehn MH, Nishimura DY, Searby CC, Mohan K, Grozdanic SD, Bugge K, Anderson MG, Clark AF, Stone EM, Sheffield VC. Reduction of ER stress via a chemical chaperone prevents disease phenotypes in a mouse model of primary open angle glaucoma. J Clin Invest. 2011;121(9):3542-53. Epub 20110808. doi: 10.1172/JCI58183. PubMed PMID: 21821918; PMCID: PMC3163970.

28.   Tasic B, Hippenmeyer S, Wang C, Gamboa M, Zong H, Chen-Tsai Y, Luo L. Site-specific integrase-mediated transgenesis in mice via pronuclear injection. Proc Natl Acad Sci U S A. 2011;108(19):7902-7. Epub 20110404. doi: 10.1073/pnas.1019507108. PubMed PMID: 21464299; PMCID: PMC3093482.

29.   Kasetti RB, Maddineni P, Kiehlbauch C, Patil S, Searby CC, Levine B, Sheffield VC, Zode GS. Autophagy stimulation reduces ocular hypertension in a murine glaucoma model via autophagic degradation of mutant myocilin. JCI Insight. 2021;6(5). Epub 20210308. doi: 10.1172/jci.insight.143359. PubMed PMID: 33539326; PMCID: PMC8021112.

30.   Muzumdar MD, Tasic B, Miyamichi K, Li L, Luo L. A global double-fluorescent Cre reporter mouse. Genesis. 2007;45(9):593-605. doi: 10.1002/dvg.20335. PubMed PMID: 17868096.

31.   Kasetti RB, Patel PD, Maddineni P, Patil S, Kiehlbauch C, Millar JC, Searby CC, Raghunathan V, Sheffield VC, Zode GS. ATF4 leads to glaucoma by promoting protein synthesis and ER client protein load. Nat Commun. 2020;11(1):5594. Epub 20201105. doi: 10.1038/s41467-020-19352-1. PubMed PMID: 33154371; PMCID: PMC7644693.

32.   Patil SV, Kaipa BR, Ranshing S, Sundaresan Y, Millar JC, Nagarajan B, Kiehlbauch C, Zhang Q, Jain A, Searby CC, Scheetz TE, Clark AF, Sheffield VC, Zode GS. Lentiviral mediated delivery of

CRISPR/Cas9 reduces intraocular pressure in a mouse model of myocilin glaucoma. Res Sq. 2023. Epub 20231219. doi: 10.21203/rs.3.rs-3740880/v1. PubMed PMID: 38196579; PMCID: PMC10775399.

33.   Zode GS, Sethi A, Brun-Zinkernagel AM, Chang IF, Clark AF, Wordinger RJ. Transforming growth factor-β2 increases extracellular matrix proteins in optic nerve head cells via activation of the Smad signaling pathway. Mol Vis. 2011;17:1745-58. Epub 20110628. PubMed PMID: 21738403; PMCID: PMC3130727.

34.   Patil SV, Kaipa BR, Ranshing S, Sundaresan Y, Millar JC, Nagarajan B, Kiehlbauch C, Zhang Q, Jain A, Searby CC, Scheetz TE, Clark AF, Sheffield VC, Zode GS. Lentiviral mediated delivery of CRISPR/Cas9 reduces intraocular pressure in a mouse model of myocilin glaucoma. Scientific Reports. 2024;14(1):6958. doi: 10.1038/s41598-024-57286-6.

35.   Maddineni P, Kasetti RB, Patel PD, Millar JC, Kiehlbauch C, Clark AF, Zode GS. CNS axonal degeneration and transport deficits at the optic nerve head precede structural and functional loss of retinal ganglion cells in a mouse model of glaucoma. Mol Neurodegener. 2020;15(1):48. Epub 20200827. doi: 10.1186/s13024-020-00400-9. PubMed PMID: 32854767; PMCID: PMC7457267.

36.   Xu Q, Rydz C, Nguyen Huu VA, Rocha L, Palomino La Torre C, Lee I, Cho W, Jabari M, Donello J, Lyon DC, Brooke RT, Horvath S, Weinreb RN, Ju WK, Foik A, Skowronska-Krawczyk D. Stress induced aging in mouse eye. Aging Cell. 2022;21(12):e13737. Epub 20221117. doi: 10.1111/acel.13737. PubMed PMID: 36397653; PMCID: PMC9741506.

37.   Peters JC, Bhattacharya S, Clark AF, Zode GS. Increased Endoplasmic Reticulum Stress in Human Glaucomatous Trabecular Meshwork Cells and Tissues. Invest Ophthalmol Vis Sci. 2015;56(6):3860-8. doi: 10.1167/iovs.14-16220. PubMed PMID: 26066753; PMCID: 4468426.

38.   Millar JC, Clark AF, Pang IH. Assessment of aqueous humor dynamics in the mouse by a novel method of constant-flow infusion. Invest Ophthalmol Vis Sci. 2011;52(2):685-94. Epub 20110203. doi: 10.1167/iovs.10-6069. PubMed PMID: 20861483.

39.   Du SW, Komirisetty R, Lewandowski D, Choi EH, Panas D, Suh S, Tabaka M, Radu RA, Palczewski K. Conditional deletion of miR-204 and miR-211 in murine retinal pigment epithelium results in retinal degeneration. J Biol Chem. 2024;300(6):107344. Epub 20240504. doi: 10.1016/j.jbc.2024.107344. PubMed PMID: 38705389; PMCID: PMC11140208.

40.   Liu S, Xiang K, Lei Q, Qiu S, Xiang M, Jin K. An optimized procedure to record visual evoked potential in mice. Exp Eye Res. 2022;218:109011. Epub 20220301. doi: 10.1016/j.exer.2022.109011. PubMed PMID: 35245512.

41.   Crish SD, Sappington RM, Inman DM, Horner PJ, Calkins DJ. Distal axonopathy with structural persistence in glaucomatous neurodegeneration. Proc Natl Acad Sci U S A. 2010;107(11):5196-201. Epub 20100301. doi: 10.1073/pnas.0913141107. PubMed PMID: 20194762; PMCID: PMC2841892.

42.   McDowell CM, Luan T, Zhang Z, Putliwala T, Wordinger RJ, Millar JC, John SW, Pang IH, Clark AF. Mutant human myocilin induces strain specific differences in ocular hypertension and optic nerve damage in mice. Exp Eye Res. 2012;100:65-72. Epub 20120503. doi: 10.1016/j.exer.2012.04.016. PubMed PMID: 22575566; PMCID: PMC3612883.

43.   Millar JC, Pang IH, Wang WH, Wang Y, Clark AF. Effect of immunomodulation with anti-CD40L antibody on adenoviral-mediated transgene expression in mouse anterior segment. Mol Vis. 2008;14:10-9. Epub 20080109. PubMed PMID: 18246028; PMCID: PMC2267727.

44.   Lee D, Liu J, Junn HJ, Lee EJ, Jeong KS, Seol DW. No more helper adenovirus: production of gutless adenovirus (GLAd) free of adenovirus and replication-competent adenovirus (RCA) contaminants. Exp Mol Med. 2019;51(10):1-18. Epub 20191028. doi: 10.1038/s12276-019-0334-z. PubMed PMID: 31659156; PMCID: PMC6817846.

45.   Rohen JW, Lütjen-Drecoll E, Flügel C, Meyer M, Grierson I. Ultrastructure of the trabecular meshwork in untreated cases of primary open-angle glaucoma (POAG). Exp Eye Res. 1993;56(6):683-92. doi: 10.1006/exer.1993.1085. PubMed PMID: 8595810.

46.   Jacobson N, Andrews M, Shepard AR, Nishimura D, Searby C, Fingert JH, Hageman G, Mullins R, Davidson BL, Kwon YH, Alward WL, Stone EM, Clark AF, Sheffield VC. Non-secretion of mutant proteins of the glaucoma gene myocilin in cultured trabecular meshwork cells and in aqueous humor. Hum Mol Genet. 2001;10(2):117-25. doi: 10.1093/hmg/10.2.117. PubMed PMID: 11152659.

47.   Joe MK, Sohn S, Hur W, Moon Y, Choi YR, Kee C. Accumulation of mutant myocilins in ER leads to ER stress and potential cytotoxicity in human trabecular meshwork cells. Biochem Biophys Res Commun. 2003;312(3):592-600. doi: 10.1016/j.bbrc.2003.10.162. PubMed PMID: 14680806.

48.   Liu Y, Vollrath D. Reversal of mutant myocilin non-secretion and cell killing: implications for glaucoma. Hum Mol Genet. 2004;13(11):1193-204. Epub 20040406. doi: 10.1093/hmg/ddh128. PubMed PMID: 15069026.

49.   Yam GH, Gaplovska-Kysela K, Zuber C, Roth J. Aggregated myocilin induces russell bodies and causes apoptosis: implications for the pathogenesis of myocilin-caused primary open-angle glaucoma. Am J Pathol. 2007;170(1):100-9. doi: 10.2353/ajpath.2007.060806. PubMed PMID: 17200186; PMCID: PMC1762699.

50.   Zhu W, Gramlich OW, Laboissonniere L, Jain A, Sheffield VC, Trimarchi JM, Tucker BA, Kuehn MH. Transplantation of iPSC-derived TM cells rescues glaucoma phenotypes in vivo. Proc Natl Acad Sci U S A. 2016;113(25):E3492-500. Epub 20160606. doi: 10.1073/pnas.1604153113. PubMed PMID: 27274060; PMCID: PMC4922164.

51.   Zhang X, Xi G, Feng P, Li C, Kuehn MH, Zhu W. Intraocular pressure across the lifespan of Tg-MYOC(Y437H) mice. Exp Eye Res. 2024;241:109855. Epub 20240305. doi: 10.1016/j.exer.2024.109855. PubMed PMID: 38453040.

52.   Tamm ER. Myocilin and glaucoma: facts and ideas. Prog Retin Eye Res. 2002;21(4):395-428. doi: 10.1016/s1350-9462(02)00010-1. PubMed PMID: 12150989.

53.   Clark AF, Steely HT, Dickerson JE, Jr., English-Wright S, Stropki K, McCartney MD, Jacobson N, Shepard AR, Clark JI, Matsushima H, Peskind ER, Leverenz JB, Wilkinson CW, Swiderski RE, Fingert JH, Sheffield VC, Stone EM. Glucocorticoid induction of the glaucoma gene MYOC in human and monkey trabecular meshwork cells and tissues. Invest Ophthalmol Vis Sci. 2001;42(8):1769-80. PubMed PMID: 11431441.

54.   Anderssohn AM, Cox K, O'Malley K, Dees S, Hosseini M, Boren L, Wagner A, Bradley JM, Kelley MJ, Acott TS. Molecular chaperone function for myocilin. Invest Ophthalmol Vis Sci. 2011;52(10):7548-55. doi: 10.1167/iovs.11-7723. PubMed PMID: 21873671; PMCID: PMC3183979.

55.   Kim H, Kim M, Im SK, Fang S. Mouse Cre-LoxP system: general principles to determine tissue-specific roles of target genes. Lab Anim Res. 2018;34(4):147-59. Epub 20181231. doi: 10.5625/lar.2018.34.4.147. PubMed PMID: 30671100; PMCID: PMC6333611.

56.   Fahy ET, Chrysostomou V, Crowston JG. Mini-Review: Impaired Axonal Transport and Glaucoma. Curr Eye Res. 2016;41(3):273-83. Epub 20150630. doi: 10.3109/02713683.2015.1037924. PubMed PMID: 26125320.

57.   Dias MS, Luo X, Ribas VT, Petrs-Silva H, Koch JC. The Role of Axonal Transport in Glaucoma. Int J Mol Sci. 2022;23(7). Epub 20220401. doi: 10.3390/ijms23073935. PubMed PMID: 35409291; PMCID: PMC8999615.

58.   Sharoukhov D, Bucinca-Cupallari F, Lim H. Microtubule Imaging Reveals Cytoskeletal Deficit Predisposing the Retinal Ganglion Cell Axons to Atrophy in DBA/2J. Invest Ophthalmol Vis Sci. 2018;59(13):5292-300. doi: 10.1167/iovs.18-24150. PubMed PMID: 30383181; PMCID: PMC6218151.

59.   Chidlow G, Ebneter A, Wood JP, Casson RJ. The optic nerve head is the site of axonal transport disruption, axonal cytoskeleton damage and putative axonal regeneration failure in a rat model of glaucoma. Acta Neuropathol. 2011;121(6):737-51. Epub 20110211. doi: 10.1007/s00401-011-0807-1. PubMed PMID: 21311901; PMCID: PMC3098991.

60.   Anderson DR, Hendrickson A. Effect of intraocular pressure on rapid axoplasmic transport in monkey optic nerve. Invest Ophthalmol. 1974;13(10):771-83. PubMed PMID: 4137635.

61.   Balaratnasingam C, Morgan WH, Bass L, Cringle SJ, Yu DY. Time-dependent effects of elevated intraocular pressure on optic nerve head axonal transport and cytoskeleton proteins. Invest Ophthalmol Vis Sci. 2008;49(3):986-99. doi: 10.1167/iovs.07-1090. PubMed PMID: 18326722.

62.   Dias MS, Luo X, Ribas VT, Petrs-Silva H, Koch JC. The Role of Axonal Transport in Glaucoma. International Journal of Molecular Sciences. 2022;23(7):3935. PubMed PMID: doi:10.3390/ijms23073935.

63.   Ridder WH, 3rd, Nusinowitz S. The visual evoked potential in the mouse--origins and response characteristics. Vision Res. 2006;46(6-7):902-13. Epub 20051020. doi: 10.1016/j.visres.2005.09.006. PubMed PMID: 16242750.

64.   Taniguchi T, Shimazawa M, Hara H. Alterations in neurofilament light in optic nerve in rat kainate and monkey ocular hypertension models. Brain Res. 2004;1013(2):241-8. doi: 10.1016/j.brainres.2004.04.023. PubMed PMID: 15193534.

65.   Wilson GN, Smith MA, Inman DM, Dengler-Crish CM, Crish SD. Early Cytoskeletal Protein Modifications Precede Overt Structural Degeneration in the DBA/2J Mouse Model of Glaucoma. Front

Neurosci. 2016;10:494. Epub 20161103. doi: 10.3389/fnins.2016.00494. PubMed PMID: 27857681; PMCID: PMC5093131.

66.   Rao MV, Darji S, Stavrides PH, Goulbourne CN, Kumar A, Yang DS, Yoo L, Peddy J, Lee JH, Yuan A, Nixon RA. Autophagy is a novel pathway for neurofilament protein degradation in vivo. Autophagy. 2023;19(4):1277-92. Epub 20220921. doi: 10.1080/15548627.2022.2124500. PubMed PMID: 36131358; PMCID: PMC10012948.

67.   Nettesheim A, Dixon A, Shim MS, Coyne A, Walsh M, Liton PB. Autophagy in the Aging and Experimental Ocular Hypertensive Mouse Model. Invest Ophthalmol Vis Sci. 2020;61(10):31. doi: 10.1167/iovs.61.10.31. PubMed PMID: 32797200; PMCID: PMC7441338.

68.   Dixon A, Shim MS, Nettesheim A, Coyne A, Su CC, Gong H, Liton PB. Autophagy deficiency protects against ocular hypertension and neurodegeneration in experimental and spontanous glaucoma mouse models. Cell Death Dis. 2023;14(8):554. Epub 20230824. doi: 10.1038/s41419-023-06086-3. PubMed PMID: 37620383; PMCID: PMC10449899.