Placozoan secretory cell types implicated in feeding, innate immunity and regulation of behavior

Tatiana D. Mayorova[1,#], Thomas Lund Koch[2], Bechara Kachar[3], Jae Hoon Jung[1], Thomas S. Reese[1], Carolyn L. Smith[4]*

[1]Laboratory of Neurobiology, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, United States of America

[2]Department of Biochemistry, University of Utah, Salt Lake City, Utah, United States of America

[3]Section on Structural Cell Biology, Laboratory of Cell Structure and Dynamics, National Institute on Deafness and Other Communication Disorders, National Institutes of Health, Bethesda, MD, United States of America

[4]Light Imaging Facility, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, United States of America

[#] Current address: Department of Biology, University of Toronto Mississauga, Mississauga, Ontario, Canada

*Corresponding author

E-mail: smithca@ninds.nih.gov (CLS)

Short title: Placozoan secretory cell types

Acknowledgements

We thank the Electron Microscopy Facility of the National Institute of Neurological

Disorders and Stroke, the Central Microscopy Facility of the Marine Biological

Laboratory, Christine A. Winters (NINDS) and Katherine Hammar (MBL) for sample

preparations and technical support. We thank Professor Adriano Senatore, University

of Toronto, for valuable comments on the manuscript. This work was supported by the

Intramural Research Program of the National Institute of Neurological Disorders and

Stroke, National Institutes of Health, National Institutes of Health, Bethesda, MD, USA.

Abstract

Placozoa are millimeter-sized, flat, irregularly shaped ciliated animals that crawl on surfaces in warm oceans feeding on biofilms, which they digest externally. They stand out from other animals due to their simple body plans. They lack organs, body cavities, muscles and a nervous system and have only seven broadly defined morphological cell types, each with a unique distribution. Analyses of single cell transcriptomes of four species of placozoans revealed greater diversity of secretory cell types than evident from morphological studies, but the locations of many of these new cell types were unknown and it was unclear which morphological cell types they represent. Furthermore, there were contradictions between the conclusions of previous studies and the single cell RNAseq studies. To address these issues, we used mRNA probes for genes encoding secretory products expressed in different metacells in *Trichoplax adhaerens* to localize cells in whole mounts and in dissociated cell cultures, where their morphological features could be visualized and identified. The nature and functions of their secretory granules were further investigated with electron microscopic techniques and by imaging secretion in live animals during feeding episodes. We found that two cell types participate in disintegrating prey, one resembling a lytic cell type in mammals and another combining features of zymogen gland cells and enterocytes. We identified secretory epithelial cells expressing glycoproteins or short peptides implicated in defense. We located seven peptidergic cell types and two types of mucocytes. Our findings reveal mechanisms that placozoans use to feed and protect themselves from pathogens and clues about neuropeptidergic signaling. We compare placozoan secretory cell types with cell types in other animal phyla to gain insight about general

65 evolutionary trends in cell type diversification, as well as pathways leading to the

66 emergence of synapomorphies.

Introduction

Placozoans are millimeter-sized, flat, irregularly shaped animals that crawl on surfaces

in shallow zones of tropical and subtropical oceans [1–3]. They have no nervous

system, muscles or internal digestive cavity. They feed on microalgae and

cyanobacteria, which they digest externally in the space between their lower epithelium

and the substrate ([4], Fig 1A). They are of interest in an evolutionary context due to

their phylogenetic position as sister to the clade that includes Cnidaria and Bilateria[5–7]

(Fig 1B) and their simple body plans (Fig 1C) and lifestyles. [8–12] Several dozen

distinct placozoan haplotypes have been identified based on sequencing the

mitochondrial large ribosomal subunits of specimens collected in different parts of the

globe. Based on analyses of nuclear gene sequences of 26 haplotypes, Placozoa is

proposed to include 2 classes, 4 orders, 5 families, 8 genera and 26 species/haplotypes

[13]. Placozoans primarily propagate by binary fission or budding [14], processes that

generate genetically identical clones though sexual reproduction might occur[15] .

Fig 1. *Trichoplax adhaerens (TH1)*. (A) Merged brightfield and fluorescence image of

TH1 pausing to feed on *Rhodomonas salina* microalgae (red particles). The image was

captured with a confocal microscope (reproduced from [4]). Cells in the ventral

epithelium secreted substances that lysed algae, releasing fluorescent phycoerythrin

(red clouds). (B) Phylogenetic tree for Metazoa [11] and phylogenetic relationships

among four placozoan species here studied. (C) A drawing of main cell types in *TH1* as

observed by transmitted electron microscopy (reproduced with permission from [16]).

DEC – dorsal epithelial cell; VEC – ventral epithelial cell.

Although placozoans are genetically diverse, they look alike except for variations in size

and shape. Electron microscopic studies of representatives of two proposed orders,

Trichoplax adhaerens (*TH1), Trichoplax sp. (TH2*; order Trichoplacea), and *Hoilungia*

hongkongensis (HH13; order Hoilungea) showed that they have morphologically

identical cell types and body plans [16–20]. The most prevalent cells in the lower

epithelium are columnar ventral epithelial cells (VEC), which have an apical cilium

surrounded by a collar of microvilli. The cilia are motile and adhere to the substrate

during their effective strokes, thereby allowing the animal to crawl on the substrate and

to change shape [4,21,22]. Interspersed among the ciliated VEC are lipophil cells (LC),

so called because they contain multiple large lipophilic secretory granules. When *T.*

adhaerens pause to feed on algae, LC secrete a large apical granule whose contents

lyses nearby algae [4]. The lower epithelium also contains mucocytes that secrete

mucus [18], which the animal requires to crawl on the substrate. The upper epithelium

is composed primarily of monociliated cells with broad apical endings that comprise the

upper the surface. Their cell bodies are narrower and protrude below. Both dorsal and

ventral epithelia contain several morphologically distinct types of gland cells that

possess a cilium and may be sensory [18,22,23]. The space between the upper and

lower epithelial layers is occupied by a layer of fiber cells, which have long branching

processes that contact each other as well as the other cell types. Fiber cells

phagocytose bacteria and cellular debris and engage in wound healing [24], functions

associated with macrophage-like cells in other animals. Inside the rim of the animal,

where dorsal and ventral epithelia meet, are regularly spaced crystal cells that appear to

be functional statocysts [25].

Bioinformatic analyses of *TH1* [5,6,26–29] reveal a rich repertoire of secretory proteins

including digestive enzymes, proteins implicated in innate immunity in other animals,

and precursors of peptides, some of which elicit changes in the behavior of the animals

[22,23]. Single cell RNA sequencing (RNA-Seq) analyses of *TH1* and three other

species of Placozoa [6,30] revealed that many of these secretory products are

differentially expressed. Cells expressing several digestive enzymes have been

localized by fluorescence *in situ* hybridization (FISH) [6,18] and some peptidergic cell

types have been localized by immunolabeling [18,22,23]. However, the cells that

express many secretory products remained unidentified.

In the present study, we used FISH probes for genes encoding secretory products that

are specifically expressed in *TH1* metacells classified as lipophil, gland, or epithelial and

a subset of the peptidergic metacells to localize the cells in whole mounts and in

dissociated cell cultures and by staining them with fluorescent dyes or immunolabels.

The compositions of the secretory granules in LC, VEC and dorsal epithelial cells were

further investigated with electron microscopic techniques. We used fluorescent dyes

and light microscopy to visualize secretory processes in live animals feeding on algae.

Our findings yield a more detailed picture of the placozoan body plan than was apparent

from previous morphological studies and provide insight into the mechanisms that the

animals use to feed and maintain homeostasis. Comparative analyses of the

secretomes of metacells in *TH1,TH2*, *HH13,* and *Cladtertia collaboinventa* (*CH23)*

revealed numerous similarities and several intriguing differences suggesting important

differences across placozoan species.

Results

Main secretory cell types in the ventral epithelium

Trichoplax adhaerens H1 lipophil cell (LC) metacell clusters highly express multiple

genes that are not expressed or expressed at much lower levels in other metacells (S1

Fig, S3 Fig) identified by single cell RNA-Seq studies [6,30] We obtained fluorescence

in situ hybridization (FISH) probes for three of them: *Ta 58643* (*Ta Tetraspanin*); *Ta*

29105 (*Ta GABA transporter*); *Ta 63996* (*TH1* secretory protein). We used probes for

*TH1*orthologs of the precursors of *trypsin (Ta 63128)*, *chymotrypsin* (Ta 63088) and

secretory phospholipase A2 (*sPLA2; Ta 57870*) to label the cell types classified as

"digestive gland cells" in the RNA-Seq studies[6,31].

Cells (presumably LC) co-labelled with probes for *Ta tetraspanin*, *Ta GABA transporter*

and *Ta 63996* were present throughout the central region of the ventral epithelium but

absent in a zone <10 µm from the rim (Fig2A and S2A Fig). Cells labeled with probes

for digestive enzymes (*Ta trypsin*, *Ta chymotrypsin*) were interspersed with LC in the

central part of the animal but were absent <60 μm from the rim (Fig 2B and S2B Fig).

The label in these cells was closer to the ventral surface (Fig 2B, bottom and S2B Fig)

than the label in LC (Fig 2A, B, and S2A, B Fig). Dissociated cells expressing LC

specific markers were larger (~8 μm) than cells expressing digestive enzymes (<5 μm;

Fig 2C, D, and S2C, D Figs) and contained large granules, a hallmark of LC. Unlabeled

ventral epithelial cells, identified by their small sizes and cylindrical shapes, also were

present (Fig 2D). We previously reported that LC express digestive enzymes based on

sequential labeling with vital dyes in live dissociated cultures and FISH probes for

precursors of digestive enzymes after fixation of the cells [18]; in this study, we found

that the expression level of digestive enzymes by LC was very low, as reported [6].

Fig 2. Fluorescence *in situ* hybridization (FISH) localization of expression of

lipophil and digestive gland cells markers in *T. adhaerens* H1 wholemounts (A, B)

and dissociated cell preparations (C, D). Images of wholemounts are horizontal (xy)

and vertical (xz, from boxed region on xy) maximum intensity projections encompassing

~1/4 of the diameter of the animal (edge of the animal is outlined white). Mucocytes are

labeled with WGA. (A) Lipophil cell markers (*Ta Tetraspanin*, *Ta* 63996, and *Ta GABA*

transporter) are co-expressed in scattered clusters ~8 μm in diameter throughout the

central region of the animal, starting ~ 10 μm from the rim. (B) Digestive cell markers

(*Ta Trypsin* and *Ta Chymotrypsin*) are highly expressed in a region starting ~60 μm from

the rim. Cells expressing a lipophil specific marker (*Ta GABA transporter*) are

interspersed among the digestive gland cells in this region. (C and D) Lipophil cell

markers (C, *Ta GABA transporter*, and D, *Ta Tetraspanin*) and digestive gland cell

markers (C, *Ta PLA2*, and D, *Ta Chymotrypsin*) are expressed in different populations of

cells. Nuclei are labelled with DAPI (maximum intensity projections merged with DIC).

Color separated images corresponding to A-D are shown in S2 Fig. fc – fiber cells; ec –

epithelial cells; lc – lipophil cells. Scale bars 10 µm.

Lipophil cells in thin section from animals prepared by high pressure freezing and freeze

substitution contained granules of varying sizes and appearances (Fig 3A). Granules in

the basal part of the cell, the location of the Golgi complex where the granules originate

[4], ranged in size from a few hundred nm to 2 µm and were electron lucent (the same

as background) (Fig 3A,B). Some granules contained membrane bound profiles with

content that resembled cytoplasm (Fig 3B). Analysis of serial thin sections (Fig 3B),

freeze fracture replicas (Fig 3C), and tomograms (Fig 3D, E) of these granules showed

that the profiles represented cytoplasmic protrusions into the granule. Granules closer

to the apical pole of the cell were up to 5 µm in diameter and contained variable

amounts of electron dense material as well as areas that were electron lucent (Fig 3A,

B). The largest granule resided close to the apical surface of the cell. These apical

granules often had a ring of electron dense material under their membrane (Fig 3A, F),

but some were completely devoid of electron dense material. Small (<100 nm)

membrane bound inclusions typically were present near the portion of the granule

membrane that was closest to the apical surface of the cell (Fig 3A, F). Examination of

these small inclusions in serial sections confirmed that they resided in the interior of the

granule. We saw a single example of a LC granule with an open fusion pore and the

content of the granule, including small membrane bound inclusions, exposed to the

exterior (Fig 3F). The continuity of the granule and plasma membranes at the edge of

the pore leads us to believe that this profile represents a stage in exocytosis of a LC

granule.

Fig 3. Ultrastructural features of lipophil cells and the structure of a highly

expressed lipophil cell secretory protein. (A) Transmission electron microscopy

(TEM) image of a thin section from a lipophil cell (LC) with osmiophilic material in its

granules. The inset shows a TEM image of a section from an animal fixed with osmium,

which better preserved the lipidic content of the granules and their cores. (B) SEM

image of a section taken in a backscatter mode shows that granules in basal portions of

LC have a dense core (artificially colored red) surrounded by electron lucent content.

Inset is an enlarged view of the boxed region. Arrowhead indicates an apical LC

granule. (C) Freeze fracture replica showing a basal part of a LC with multiple granules.

Note dual component nature of the granules. A fracture through the interior of a core

reveals heterogeneous content resembling cytoplasm. Arrowhead marks a protrusion

emanating inward from the granule wall. (D) TEM image of 100 nm thick section

showing basal region of a LC as inferred from its proximity to the dorsal side of the

animal. Red box depicts a region with a protrusion inside a granule, further studied by

EM tomography in (E). (E) Upper panel is an EM projection showing direct connection

between the protrusion and the inner surface of the granule. Lower panel is a 1.75 nm

thick virtual slice through a reconstructed volume of the tomogram. Note that ER

(arrowhead) penetrates the protrusion. (F) TEM image of an apical part of an LC

showing exocytosis (arrowhead) of a large granule. Note osmiophilic material and

membranous particles located near the site of exocytosis. (G) Freeze fracture replica

showing an apical part of a LC and its apical granule. (H) A protein present exclusively

in LC secretome is largely composed of alpha helices (AlphaFold per-residue

confidence score, pLDDT, color-coded: dark blue > 90 very high confidence; light blue

90□>□pLDDT > 70 confident; yellow 70□>□pLDDT > 50 low confidence; and orange <

50 very low confidence) and is positively charged (electrostatic map created with

ChimeraX). dec – dorsal epithelial cell; e – e-face; fc – fiber cell; lc – lipophil cell; p – p-

face. Scale bars 5 µm (B, D), 800 nm (A, C, F, G), and 50 nm (E).

The variable appearance and patchy distribution of the content of LC granules

suggested that the content was partially extracted by the procedure used for freeze

substitution and fixation. The apical granule in LC in thin sections from animals fixed in

glutaraldehyde and osmium simultaneously in aqueous solution at room temperature

was a large electron dense sphere with one or two electron lucent inclusions inside

(clearer than background plastic; Fig 3A, inset). The electron dense material may

represent unsaturated lipids, which are osmiophilic [31] and might be extracted by the

organic solvents used for freeze-substitution of frozen specimens. The clear inclusions

contain substances that do not bind the uranyl acetate and lead citrate grid stains

suggesting that they are not proteinaceous. Analysis of freeze fracture replicas from

rapidly frozen animals provided further evidence of the heterogeneous nature of the

content of LC granules (Fig 3C, G). A fracture face through the outer content of the

granule was smooth in appearance while the fracture face of the granule core was

rough.

The uncharacterized protein *Ta 63996* that served as one of our markers for LC is the

second most highly expressed genes in *TH1* LC metacells (S1 and S3 Figs) and the

identical gene in *TH2* (*006628*) is the most highly expressed gene in *TH2* LC metacells

(S4 Fig). The gene was annotated as laminin subunit *beta,* but it lacks a laminin

domain and laminin-type EGF domains. It has only weak homology with a region

located downstream of the critical laminin domains in laminin beta subunit orthologs in

other animals and is much shorter (346 aa versus ~1700 aa) than in other species. The

gene has a signal peptide and no transmembrane domains. Only LC express this

putative secretory protein in *TH1* although some cells classified as "upper-epithelial-like"

express it in *TH2*. The protein structure predicted by AlphaFold includes extended

helical regions with abundant basic amino acids (Fig. 3H). Many lysines are grouped in

pairs, composing a KK motif, a well-known cleavage site in neuropeptide prohormones.

However, this motif is not conserved beyond Trichoplacidae, and LC do not express any

known prohormone convertases or cathepsins. We found no other highly expressed

secretory proteins that lack transmembrane domains in *Trichoplax* LC metacells.

Two LC metacell clusters, LC1 and LC2, were identified in *HH13* (S5 Fig) and *CH23* (S6

Fig). The *HH13* LC2 metacell cluster expressed a gene whose sequence was 34%

identical and 52% positive to the uncharacterized secretory protein *Ta 63996*. In *CH23*,

both LC1 and LC2 metacells expressed an ortholog (e^{-45}) of *Ta 63996*. All LC metacells

in *TH1, TH2*, *HH13* and *CH23* contained a gene identical to the Tetraspanin (*Ta 58643*)

we used as a second marker for LC. The third marker we used for LC, *Ta 20105,* was

annotated as a solute carrier (*GABA transporter*). It was expressed in all metacells

classified as LC in *TH1* and *TH2* but only in LC1 metacells in *HH13* and *CH23*. All LC

metacell clusters in *TH1, TH2, HH13* and *CH23* highly expressed several fatty-acid

binding proteins, V-type ATPases and solute carriers, and the lysosomal membrane

glycoprotein, LAMP2 (S3-6 Figs).

We further investigated the morphology of cells that express digestive enzymes by co-

labeling dissociated cell preparations with FISH probes for the precursors of trypsin and

sPLA2 and an antibody against acetylated tubulin, a component of cilia. Cells co-

expressing *Ta trypsin* and *Ta sPLA2* often occurred in clusters (Fig 4A). The cells were

small and cylindrical in shape and many of them possessed a cilium. Labeled cells

lacking a cilium may have lost their cilium during dissociation since cilia that were not

attached to a cell also were present. Ciliated ventral epithelial cells (VEC) are the most

prevalent cell type in the ventral epithelium. Given the high density of cells expressing

digestive enzymes in the central part of the ventral epithelium and their possession of a

cilium, we identify "digestive gland cells" as a subtype of VEC (S1 and S3-6 Figs).

Fig 4. Two monociliated ventral epithelial cell (VEC) types: digestive gland cells

and cells expressing putative antimicrobial peptides (AMPs). (A) Cells secreting

digestive enzymes bear a cilium (combination of FISH for *Ta Trypsin* and *Ta PLA2* and

immunolabelling for tubulin; left panel shows fluorescence merged with DIC). (B, C)

TEM shows that both peripheral (B) and central (C) ciliated epithelial cells have ~500

nm diameter granules located close to the apical surface. Boxed regions are magnified

in insets. (D) The granules in peripheral cells are more electron dense and abundant

than those in cells in more central regions. ∗ p<0.05. (E) Trypsin is highly expressed in

the central region of the ventral epithelium, while *AMP1* expression is restricted to the

peripheral region. (F) In dissociated cell preparations, two distinct subpopulations of

cells expressing either *AMP1* or *Ta Trypsin* are apparent. Most *AMP1* expressing cells

strongly express *AMP2* whereas *Ta Trypsin* expressing cells show weak *AMP2*

expression (not visible in E). A dissociated trypsin+ cell (F, green) contains one red

AMP2 grain. (G) Venn diagrams based on cell counts show that three quarters of VEC

express either *AMP1* or *Ta Trypsin* along with *AMP2*. Some mucocytes and DEC

express *AMP1* with or without either *AMP2* or *Ta Trypsin*. Only a few lipophil and fiber

cells express *AMP1* or *Ta Trypsin*. Color coding for Venn diagrams is the same as the

fluorescence colors on E and F; double co-expression is indicated as strips of

respective colors, triple co-expression is white, and the absence of expression is dark

gray. Numbers of counted cells are shown. Scale bars: 10 µm (A, E, F) and 1 µm (B,

C).

As VEC in the peripheral part of the ventral epithelium (pVEC; within ~60 µm of the rim)

do not express digestive enzymes we speculated that their ultrastructure might differ

from that of the centrally located VEC (cVEC) that express genes encoding precursors

of digestive enzymes. Comparison of pVEC and cVEC within the same thin sections

revealed differences in their secretory granules (Fig 4B-D). Peripheral VEC, identified

based on their proximity to the rim, had numerous electron dense secretory granules

near their apical surfaces (Fig 4B, D). Ventral epithelial cells located in the central

region, in the vicinity of LC, had fewer granules (p=1.96E-05) and the granules were

less electron dense (p=1.26E-17) (Fig 4C, D).

Metacells identified as "epithelial" in *TH1* [30] or "lower epithelial" [6] expressed multiple

genes that were predicted precursors of secretory proteins/peptides (InterPro). We

noticed that the sequences of five of them resembled those of arminins, a type of

antimicrobial peptide (AMP) expressed in *Hydra* [32], in that they were ~150- 250 aa in

length, had a signal peptide, a highly acidic N-terminal region and alkali C-terminal

region that includes aliphatic amino acids and DE cleavage motifs (S7 Fig) common in

placozoan and cnidarian prepropeptides [33,34]. The N-terminal domain of one of them,

Ta 55945, was annotated "alpha defensin" (Interpro). The signal peptide and acidic N-

terminal regions of arminin prepropeptides are removed to produce active arminin

peptides. Arminin peptides have a C-terminal glycine and are thought to be amidated.

The sequences of the *Trichoplax* arminin-like peptides have a C-terminal glycine, as is

required for amidation by peptidylglycine α-amidating monooxygenase (PAM). An AMP

analysis tool (Antimicrobial Peptide Data, Univ. Nebraska Medical Center) identified

hydrophobic and basic regions in the C-terminal part of the Ta AMP sequences that

might interact with membranes.

We obtained FISH probes for two of the arminin-like genes, *Ta 55945* and *Ta 56030*,

and will refer to them as *AMP1* and *AMP2*, respectively. Both probes strongly labeled

cells in the peripheral part of the ventral epithelium (Fig 4E). The AMP expressing cells

were densely packed and showed little overlap with cells expressing digestive enzymes.

Scattered cells in the dorsal epithelium showed moderate expression of *AMP1* or both

AMP1 and *AMP2*. Most of the AMP-expressing cells in dissociated cell preparations had

morphological features typical of VEC: they were small, cylindrical in shape and

possessed a cilium (Fig 4F). They often occurred in clusters with other AMP-expressing

cells. Most *AMP1* expressing VEC strongly expressed *AMP2*. *Trypsin*-expressing VEC

weakly expressed *AMP2*, containing fewer and smaller fluorescent grains than cells co-

expressing *AMP1* and *AMP2*. Some WGA-positive mucocytes and DEC expressed

AMP1 with or without *AMP2*. A small fraction of LC and fiber cells contained one or a

few *AMP1* or *Ta trypsin* positive grains (Fig 4G) and may represent cells

transdifferentiating from cVEC to lipophil or fiber cells, respectively [6].

The genes encoding *AMP1*, *AMP2*, and a third arminin-like prepropeptide (*Ta 60631;*

AMP3) are the three most highly expressed genes in *Trichoplax* H1 classified as "lower

epithelial" metacells [6] . Our findings demonstrate that these metacells correspond to

peripheral VEC (pVEC; S1 and S3 Figs). The *TH2* metacells classified as "lower

epithelial" contain genes that are nearly identical to *AMP1*, *AMP2* and *AMP3* (S4 Fig).

The *AMP1* and *AMP3* genes are the most highly expressed genes these metacells.

Hoilungia H13 and *CH23* metacells classified as "lower epithelial" contain a gene that is

more than 55% identical and 70% similar to *AMP3* and a gene that is more than 40%

identical and 60% similar to *AMP2* (S5 and S6 Figs). No *AMP1* ortholog was found in

HH13 or *CH23*.

Main secretory cell types in the dorsal epithelium

The most prevalent cells in the dorsal epithelium are monociliated dorsal epithelial cells

(DEC) [16]. Their broad (~10 µm) polygonal-shaped apices pave the dorsal surface.

Their cell bodies are narrower and extend into the interior, where they are surrounded

by processes of fiber cells. Freeze fracture replicas of the apices of DEC revealed

exoplasmic (e) faces and protoplasmic (p) faces of numerous, large (500 nm) secretory

granules (Fig 5A). Electron micrographs of transverse thin sections through the apices

of DEC showed dark secretory granules near their dorsal surfaces (Fig 5B). The dark

granules bound the lectin wheat-germ agglutinin (WGA) conjugated to nanogold (Fig

5B, inset). WGA also labeled the outer surfaces of DEC. At the rim, cells with

morphological characteristics of DEC were adjacent to cells with morphological

characteristics of VEC (transition zone). Secretory granules in pVEC were grey or dark

like those in DEC but bound much less WGA label (Fig 5C, insets).

Fig 5. Main secretory cells in the dorsal epithelium. (A) Freeze fracture replica at

the apex of a dorsal epithelial cell (DEC) imaged in TEM reveals e- or p-faces of

numerous ~500 nm diameter secretory granules. (B, C) TEM of ultrathin sections

labelled with nanogold conjugated WGA. Transverse section in the dorsal epithelium (B)

shows multiple WGA-stained granules in ciliated DECs; insets show enlarged view of

boxed regions. Transverse section at the transition region between dorsal and ventral

epithelia (C, the border is demarcated by dotted line) shows that DEC granules bind

more WGA than do morphologically similar granules in VEC. (D-H) Confocal images of

wholemounts (D, F) and dissociated cell preparations (E, G, H). Many DEC express *Ta*

Intelectin 60661 (D, whole mount; E, dissociated cells) as evident from co-labeling with

WGA. Mucocytes (m) label intensely with WGA, but do not express Ta *Intelectin 60661*.

Other cell types (VEC, lipophil (lc) and fiber (fc) cells) are not labeled. (F) *Ta ELPE* is

expressed in nearly all DEC and a few VEC (see xz inset in F and color separated

images in S9A Fig); (G, I) About half of *Ta ELPE*+ co-expresses *Ta Intelectin 60661*. (H)

Some VEC, identified based on their small sizes and cylindrical shapes express *Ta*

ELPE but not *Ta Intelectin 60661*. (J) Cells co-expressing *Ta ELPE* and *Ta Intelectin*

60661 are larger than those expressing only *Ta ELPE*. e – e face; fc – fiber cells; lc –

lipophil cells; m – mucocytes; p – p face.

Many DEC in wholemounts and dissociated cell preparations, identified based on their

content of granules stained with WGA conjugated to a fluorescent dye, were labeled by

a FISH probe for *Ta 60661* (Fig 5D, E), a gene that is highly expressed in *TH1* metacells

classified as "epithelial" or "upper epithelial" [6,30]. Labeled cells were present

throughout the dorsal epithelium except in a region within ~15 µm of the rim (Fig 5 D,

E). Their distribution was patchy, suggesting that only a subset of DEC expressed this

gene.

Trichoplax H1 gene *Ta 60661* is identical to a gene discovered in a search for orthologs

of genes implicated in innate immunity in *TH2* [35]. The gene was identified as

belonging to a class of secreted glycoproteins called "intelectins" based on its content of

a fibrinogen-related domain (FReD). Intelectins have been implicated in defense against

bacteria in both vertebrates and invertebrates [36]. Ten of thirty-one intelectin-like

genes identified in *TH2* are uniquely expressed in *TH1* and *TH2* metacells classified as

"upper epithelial" (S1 Fig, S3, 4 Figs). These metacells also contain a gene annotated

"membrane-associated mucin 4-like glycoprotein" based on the presence of von

Willebrand factor type D (VWD), Scavenger Receptor Cysteine-Rich (SRCR) and

multiple EGF-like domains. *Hoilungia H13* metacells classified as "upper epithelial"

include genes encoding five of the intelectins and the mucin-4 gene, but not intelectins

Ta 60661 and *Ta 61411* (S5 Fig). Instead, these genes are expressed in a metacell

classified as "gland". *Cladtertia H23* metacells classified as "upper epithelial" express

four of the intelectins expressed in upper epithelial metacells in *TH1*, *TH2* and *HH13*

(S6 Fig).

The metacells representing DEC also express a prepropeptide identified in *TH1*

genomes and transcriptomes that is predicted to produce peptides with C-terminal

amino acids Glu-Leu-Pro-Glu, or ELPE [6,23,33] . We will refer to this gene as *Ta ELPE*.

The names, sequences and predicted mature peptides produced by this and other

prepropeptide genes investigated in this study are listed in S8 Text.

A FISH probe for *Ta ELPE* prepropeptide labeled cells throughout the dorsal epithelium

including those <15 μm of the rim (Figs 5F and 6A1 and S9A Fig). Approximately half of

the *Ta ELPE*-positive DEC co-expressed *Ta Intelectin 60661* (Fig 5G-I). A few VEC,

identified based on their small sizes and cylindrical shapes, expressed *Ta ELPE* but not

intelectin *Ta Intelectin 60661* (Fig 5F, H). The mean area of cells expressing both *Ta*

Intelectin 60661 and *Ta ELPE* was greater (p=0.01) than that of cells expressing *Ta*

Intelectin 60661 without *Ta ELPE* (Fig 5G, H, J) because the latter group included the

Ta ELPE-positive VEC, which were smaller than DEC.

Scattered cells in the dorsal epithelium and a few cells in the ventral epithelium located

10 to 40 µm from the rim were labeled with a probe for a gene encoding a secretory

protein with no known functional domains, *Ta 63786* (Fig 6A1). The labeled cells did not

express *Ta ELPE* (Fig 6A1 inset and A2). The *Ta 63786+* cells in the ventral epithelium,

but not those in the dorsal epithelium, co-expressed a gene *Ta 64402* encoding a

secretory peptide that bears the same *Hydra* arminin-like features as *AMPs 1*, *2* and *3*,

including a signal for C-terminal amidation, which we call *AMP4* (Fig 7E, F).

Dissociated *Ta 63786+* cells had a cilium but were smaller and narrower (p=0.004) than

Ta ELPE+ and *Ta Intelectin 60661+* DEC cells (Fig 6A2, A3). The metacells that

express *Ta 63786* and *AMP4* were classified as "peptidergic" or "epithelial unknown"

[6,30]. These metacells also contain a gene encoding an intelectin (*Ta 64393*) that is not

expressed in any other metacell (S1 and S3 Figs). *Trichoplax* H2 metacells classified

as "gland", or "epithelial lower" [6], express genes identical to *Ta 63786* and *Ta 64393*,

and a gene that is nearly identical to *AMP4* (S4 Fig). The DEC labeled by the probe for

Ta 63786, but not the probe for *AMP4* likely correspond to *TH1* metacells C102-107 (S3

Fig) and *TH2* metacells C133 and 134 (S4 Fig), which highly express *Ta 63786* but do

not express *AMP4* or the intelectin. Metacells *TH1* C103 and *TH2* C133 and 134 were

reported to co-express *Ta 63786* and *Ta ELPE* [6] but we did not observe cells co-

expressing these genes in our microscopy studies. *Hoilungia H13* metacells C177-183

express orthologs of *Ta 63786* and *Ta 64393* intelectin (S5 Fig), but no *AMP4* ortholog

was found in this species. *Cladtertia H23* lacks orthologs of *Ta 63786*, *Ta 64393*

intelectin and *AMP4*. No ortholog of *Ta ELPE* was found in *HH13* or *CH23*.

Fig 6. Localization and characterization of peptidergic secretory cells. *Trichoplax*

wholemounts and dissociated cell preparations were labeled with FISH probes for

different prepropeptides and other cell-type specific proteins. Mucocytes were labeled

with fluorescent-conjugated WGA in wholemounts. Cilia were immunolabeled with an

antibody against acetylated tubulin and nuclei were labeled with DAPI in dissociated cell

preparations. Circle plots and box plots use the same color coding as fluorescence

images; co-expression is indicated as stripes of respective colors. (A). Prepropeptide *Ta*

ELPE (A1) was expressed by DEC throughout the animal and by some VEC near the

rim. Interspersed among the *Ta ELPE+* DEC were scattered cells that expressed an

uncharacterized protein (*Ta 63786*). Probes for *Ta ELPE* and *Ta 63786* labeled distinct

populations of ciliated cells in dissociated cell preparations (A1, circle plot; A2). *Ta*

ELPE+ cells were larger in area than *Ta 63786+* cells (A3). (B). *Ta LF* prepropeptide

(B1) was strongly expressed in a row of cells in the dorsal epithelium 5 - 10 μm from the

rim and more weakly expressed in scattered cells in the ventral epithelium in a region

starting 40 μm from the rim. The yellow channel in the color-merged xy and xz images

were dislayed with gamma= 0.54 to enhance the visibility of the weakly labeled cells.

The yellow channel is displayed with gamma=1.0 (linear) in the upper xz image. The *Ta*

LF+ ventral epithelial cells co-expressed an astacin-like metalloendopeptidase (*Ta*

26557; B1, B2), while dorsal *Ta LF+* cells did not (B1, B3, B4). Both *Ta LF+/Ta 26557+*

cells (B2) and *LF+/Ta 26557-* cells (B3) were small ciliated cells (B5). Circle plot B6

shows that *Ta LF+* cells (bright yellow) were distinct from *Ta ELPE+* cells (light yellow)

and *Ta Intelectin 60661+* cells (red stripes). (C). Mucocytes were labeled with

fluorescent WGA and a probe for a secreted oligosaccharide binding protein (*Ta*

OligoBP; 63702; C1). The central population of mucocytes co-expressed *Ta RWa*

prepropeptide (C1-3). Both *Ta RWa-* and *Ta RWa+* mucocytes lacked a cilium (C2,

tubulin label absent). The rectangular insets show *Ta RWa-* (top) and *Ta RWa+*

mucocytes (bottom) without the magenta (*Ta OligoBP*) fluorescence channel. (D). *Ta*

ELP prepropeptide and an astacin-like metalloendopeptidase (*Ta 54934*; D1) were co-

expressed in a row of cells in the ventral epithelium 15 to 30 µm from the rim. The

labeled cells were ciliated (D2, D3), co-expressed both genes (D1-4) and had an area of

about 9 mm^2 (D5). (E). Probes for *Ta SIFGa* prepropeptide and for a second

uncharacterized secretory protein (*Ta 60437*) labeled separate populations of cells: *Ta*

SIFGa+ cells were prevalent in the dorsal epithelium of the rim (E1) and more sparsely

distributed in the dorsal and ventral epithelium further in the interior; *Ta 60437+* cells

were in a row 20 to 30 µm from the rim (E1). Both the *Ta SIFGa+* and *Ta 60437+* cells

were ciliated (E2, E3). The *Ta 60437+* cells were larger than the *Ta SIFGa+* cells (E4,)

and larger than *Ta ELP+/Ta 54934+* cells (E4, red), which were in the same area (D1).

(F). A subset of *Ta SIFGa+* cells in the ventral epithelium >10 µm from the rim

expressed the *Ta FFNP* prepropeptide (F1, F2). The yellow channel was displayed with

gamma=0.75 to enhance the visibility of weakly stained cells. Color separated images

displayed with gamma=1.0 (linear) are shown in S9B Fig. Both *Ta SIFGa+/Ta FFNP-*

cells (F3) and *Ta SIFGa+/Ta FFNP+* cells (F4, F5) bear a cilium. The *Ta SIFGa+/Ta*

FFNP+ cells were larger than the *Ta SIFGa+/Ta FFNP-* cells (F6). * p<0.05. Scale bars

20 μm (A1, B1, F1), 10 μm (C1, D1, E1) and 5 μm in all dissociated cells images.

Fig 7. Localization and characterization of peptidergic cells. (A) *Trichoplax*

wholemount (top panel, horizontal maximum intensity projection; bottom panels, vertical

projections of boxed region). Mucocytes were labeled with fluorescent WGA. Cells co-

expressing *Ta LF* and astacin *Ta 26557* are interspersed among mucocytes and *Ta*

PLA2+ cells in the central part of the ventral epithelium while cells that express *Ta LF*

prepropeptide without astacin are most prevalent close to the rim. (B) Dissociated cell

preparation. Nuclei were labeled with DAPI. Color separated and merged views show

multiple cells that co-express *Ta LF* and astacin *Ta 26557* (yellow arrowheads), a cell

that expresses astacin without *Ta LF* (magenta arrowhead) and cells that express *Ta*

PLA2 (white arrowheads). (C) Many cells co-express *Ta LF* and astacin *Ta 26557,* but

few cells co-express *Ta LF+* or *Ta 26557 and Ta PLA2*. (D) All labeled cells are similar in

size with *Ta PLA2+/Ta 26557+* cells slightly larger. (E) Horizontal and vertical maximum

intensity projections of a wholemount labeled with probes for *Ta LF* prepropeptide,

secretory protein *Ta 63786*, putative antimicrobial peptide *AMP4* and fluorescent WGA.

The images span nearly the entire width of the animal – note the mucocyte and the *Ta*

LF+ cell at the upper right. Cells in a region of the ventral epithelium within 10 to 40 μm

of the rim co-express *Ta 63786* and *AMP4*; cells in the peripheral part of the dorsal

epithelium express Ta LF or Ta 63786 but do not express *AMP4*. (F) Enlarged view of

partially dispersed cells co-expressing *Ta 63786* and *AMP4*. (G) Horizontal and vertical

maximum intensity projections of an animal labeled with probes for *Ta LF*, *Ta WPPF*

prepropeptides, *AMP2*, and the lectin WGA. The images encompass ~1/3 the diameter

of the animal. Cells expressing *Ta WPPF* prepropeptide are distributed in the dorsal

epithelium in a region starting ~40 µm from the rim and do not express *Ta LF* or *AMP2*.

Scale bars 20 µm in whole mount images and 5 µm in dissociated cells images.

Morphology and locations of peptidergic secretory cells

A probe for the precursor of Leu-Phe (*Ta LF*) peptides labeled a row of cells in the

dorsal epithelium 5-10 µm from the rim and more weakly labeled scattered cells in the

ventral epithelium >40 µm from the rim (Fig 6B1). A probe for a secreted astacin-like

metalloendopeptidase, *Ta 26557*, intensely labeled the *Ta LF*+ cells in the ventral

epithelium but not the *Ta LF*+ cells in the dorsal epithelium (Figs 6B1-4 and 7A). Both

Ta LF+/*Ta astacin*+ and *Ta LF*+/*Ta astacin*- cells were cylindrical in shape, possessed a

cilium (Figs 6B2, B3, and 7B), and were similar in size (Figs 6B5 and 7C, D). The

positions of *Ta LF*+/*Ta astacin*+ cells are like those of cells classified as ventral Type 3

gland cells in ultrastructural studies of *Trichoplax* [18]. Type 3 gland cells contain small

granules with textured content. The *Ta LF*+/*Ta astacin*+ cells were intermingled with

cells expressing *Ta sPLA2*, but only a few cells co-expressed *Ta astacin* and *Ta sPLA2*

(Fig 7A-D). The *Ta LF*+/*Ta astacin*- cells located near the rim of the dorsal epithelium

did not express *Ta ELPE* or the DEC-specific intelectin *Ta 60661* (Fig 6B6). Based on

the high expression of *Ta LF* and lack of expression of astacin *Ta 26557*, these cells

represent *TH1* metacell C41 (S1 Fig) and metacells *TH1* C188, *TH2* C210, and *HH13*

C244 (S3-5 Figs). The cells that co-express *Ta LF* and astacin *Ta 26557* represent *TH1*

metacells C42 (S1 Fig) and C185 and C186 (S3 Fig), *TH2* C215 (S4 Fig), and *HH13*

C251 and C252 (S5 Fig). No ortholog of LF prepropeptide was found in *CH23* but

metacell C233 contained an ortholog of astacin *Ta 26557* (S6 Fig). This metacell

contained prepropeptide genes that bore no resemblance to *Ta LF* prepropeptide.

Mucocytes identified by labeling with WGA conjugated to a fluorescent dye were in the

ventral epithelium in a zone 10 and 30 μm from the rim and in the central region >50 μm

from the rim (Fig 6C1), as reported [18]. A probe for an oligosaccharide binding protein,

Ta 63702, labeled both peripheral and central mucocytes (Fig 6C1). A probe for the

Arg-Trp-amide (*Ta RWa*) prepropeptide labeled the central mucocytes but not the

peripheral mucocytes (Fig 6C1-3). Dissociated mucocytes were ovoid cells ~8 μm in

length and lacked a cilium, as evident from the absence of staining for acetylated tubulin

(Fig 6C2). Based on co-expression of *Ta 63702* and the *Ta RWa* prepropeptide, the

central population of mucocytes corresponds to *TH1* metacell C36 (S1 Fig), *TH2*

metacells C195 and C196 (S4 Fig), *HH13* metacells C213, C216, and C277 (S5 Fig)

and *CH23* metacell C209 (S6 Fig). These metacells express PAM, the enzyme that

converts a glycine at the C terminal of a peptide to an amide, a secreted glycoprotein

with VWD, TIL, C8 and PTS domains characteristic of gel-forming mucins[37], and a

secretory protein with no known functional domains, *Ta 62229*, that is not expressed in

other metacells. *Trichoplax* H2 metacell C196, *HH13* metacell C218, and *CH23* metacell

C210 contain orthologs of *Ta 63702* and the gel-forming mucin but lack the RWa

prepropeptide gene, express very little PAM and express only low levels of *Ta 62229*.

These metacells likely represent the RWa negative mucocytes that are in the peripheral

part of ventral epithelium. Expression of *Ta RWa* prepropeptide was not reported in *TH1*

in the more recent scRNAseq study [6]. However, *TH1* metacells C165 and C166

contain orthologs of Ta 63702 and the gel-forming mucin (S3 Fig) suggesting that these

metacells represent mucocytes. Metacell C165 contains PAM and *Ta 62229* but C166

does not, suggesting C165 represents central mucocytes and C166 peripheral

mucocytes. The metacells representing mucocytes in *TH2*, *HH13* and *CH23* contain

orthologs of a homeobox DBX1-B-like transcription factor that is not present in any other

metacells. The GenBank sequence for the *TH1* ortholog of DBX1 is incomplete and

consequently the gene was not included in the scRNAseq data for *TH1*.

Cells labeled by probes for the Ta-<u>e</u>ndomorphin-<u>l</u>ike <u>p</u>eptide (*TaELP*) prepropeptide [22]

(S8 Text) and a secreted astacin-like metalloendopeptidase, *Ta 54934*, were in a narrow

zone in the ventral epithelium 10-20 µm from the rim (Fig 6D1). The positions of these

cells correspond with those classified as Type 1 gland cells in ultrastructural studies of

Trichoplax [18]. Type 1 gland cells contain larger and more electron dense granules

than Type 3 gland cells. Dissociated *TaELP* -expressing cells had a cilium (Fig 6D2,

D3), nearly always co-expressed both genes (Fig 6D4), and were relatively small (<5

µm in diameter, Fig 6D5). The *TH1* metacell that expresses *TaELP* prepropeptide and

the astacin *Ta 54934* (C32, Fig 1) highly expresses an additional astacin-like gene, *Ta*

54935, that is not expressed in any other metacells. *TaELP* expression was not reported

in *TH1*[6], but a metacell expressing *Ta 54934* and *Ta 54935* was found: C183 (S3 Fig).

A metacell expressing *TaELP* ortholog was reported in *TH2*: C210. However, H2 C210

did not express astacin orthologs *Ta 54934* and *Ta 54935*. Instead, these astacins were

expressed in H2 C214 (S4 Fig). Moreover, H2 C210 highly expresses LF prepropeptide,

which we found was expressed in distinct sets of cells (Fig 5B1). We suspect that H2

metacell C214 may represent *TaELP* expressing cells. *Hoilungia H13* metacells C248

and C249 expressed orthologs of the *TaELP* prepropeptide and both astacin-like

metalloendopeptidases (S5 Fig). No ortholog of *TaELP* prepropeptide was found in

CH23. However, *CH23* metacell C232 expresses genes identical to astacins *Ta 54934*

and *Ta 54935*. Interestingly, metacells that express these astacins in all four species

also express one or two G-protein coupled receptors (GPCR) with a C-type lectin

domain (CTLD) [35].

A probe for the Ser-Ile-Phe-Gly-amide (*Ta SIFGa*) prepropeptide labeled cells in the

dorsal epithelium near the rim and scatted cells in the dorsal and ventral epithelium

further in the interior (Fig 6E1). The metacell expressing high levels of *Ta SIFGa*

prepropeptide (C46; Fig 1) was reported to express a secreted protein with no known

functional domains, *Ta 60437*, that was not expressed in any other metacell. Our probe

for *Ta 60437* did not label *Ta SIFGa*-expressing cells but instead labeled a row of cells

in the ventral epithelium 20-30 µm from the rim (Fig 6E1). The positions of these cells

correspond with those of Type 1 gland cells. Both cell types possessed a cilium (Fig

6E2, E3). *Ta 60437*+ were larger than cells expressing *Ta SIFGa* (p = 3.32E-4,

Bonferroni corrected p value) and cells co-expressing *Ta ELP* and astacin-like

metalloendopeptidase *Ta 54934* (p = 0.02, Bonferroni corrected p value), which were in

the same area (Fig 6E4). This fact, along with the location of *Ta 60437*+ cells, makes it

likely that these are Type 1 gland cells, which are larger than other ciliated cells due to

their possession of big secretory granules [18]. The *Ta 60437* gene is highly expressed

in metacells C167 in *TH1 (S3 Fig)*, C197 in *TH2 (S4 Fig)*, C197 and C219 -C222 in

HH13 (S5 Fig) and C111-112 in *CH23* (S6 Fig).

A probe for the Phe-Phe-Asn-Pro-amide (*Ta FFNPa*) labeled the *Ta SIFGa*+ cells in the

central part of the ventral epithelium, but not those in the peripheral part of the ventral

epithelium or in the dorsal epithelium (Fig 6F1, F2). The positions of the cells that co-

express these genes correspond with those classified as ventral Type 3 gland cells [18].

Both *FFNPa+ and FFNPa-* cells possessed a cilium (Fig 6F3-F5), but the *FFNPa+* cells

were larger than *the FFNPa-* cells (p=7.86E-06; Fig5F6). Cells expressing *Ta SIFGa*

and *Ta FFNPa* correspond to metacell C38 (S1 Fig) and C168 (S3 Fig). No expression

of SIFGa was reported in the metacell expressing FFNPa in *TH2* (S4 Fig), but *HH13*

metacell C223 and *CH23* metacell C213 express both SIFGa prepropeptide and very

high levels of FFNPa prepropeptide (S5 and S6 Figs). The SIFGa+ cells in the dorsal

epithelium and in the rim of the ventral epithelium correspond to *TH1* metacells C46 (S1

Fig) and C189 (S3 Fig), *TH2* C217 (S4 Fig), *HH13* C253-254 (S5 Fig), and *CH23* C221

(S6 Fig) based on high expression of SIFGa prepropeptide and absence of FFNPa

prepropeptide.

A probe for the prepropeptide precursor of the peptide WPPF labeled small (diameter

~5 µm) cells distributed throughout the central part of the dorsal epithelium starting ~40

µm from the rim (Fig 7G). Cells expressing *Ta WPPF* represent metacell *TH1* C181 (S3

Fig), *TH2* C213 (S4 Fig), *HH13* C246 (S5 Fig), and *CH23* C230 (S6 Fig).

Roles of lipophil cells and digestive gland cells in feeding

Both main secretory cell types in the ventral epithelium, LC and ciliated VEC, have been

implicated in feeding. Lipophil cells secrete granules whose content lyses algae [4].

VEC constitutively pinocytose extracellular macromolecules such as ferritin [38], HRP,

dextran [39] and mucus [18] and, likely, the contents of lysed algae. The expression of

high levels of *Ta Trypsin*, *Ta Chymotrypsin* and *Ta sPLA2* in centrally located VEC led

us to investigate whether they secrete digestive enzymes during feeding.

We used a fluorescent indicator for trypsin activity, BZiPAR [40], to test whether

digestive gland cells secrete trypsin during feeding episodes. To visualize LC granule

secretion, we supplemented seawater with a lipophilic dye, LipidTOX, which stains

intact granules, and the membrane dye FM1-43, which stains the contents of secreted

granules [4]. Animals encountering *Rhodomonas salina* algae on the substrate

manifested behaviors typical of *Trichoplax* during feeding episodes [4]: they ceased

gliding and changing shape, and their margins spread and became more closely

attached to the substrate (Fig 8A and S10 Fig; S11 and S12 Movies). Some, although

not all, LC in the vicinity of algae released their large apical secretory granule, the

contents of which became stained with FM1-43 (Fig 8B, F and S10 Fig). The stained

content initially was a diffuse cloud but then became more concentrated. When a LC

released its granule, the surrounding area of the ventral epithelium moved farther away

from the substrate. These movements, together with the slow scan speeds (~1

frame/sec) used for these experiments, made it difficult to visualize secretion of

LipidTOX-stained granules, which happens rapidly, so we used the appearance of FM1-

43-stained spots to monitor granule secretion. Some animals contracted during lipid

granule secretion and their rims detached from the substrate, but then they flattened

and reattached (not illustrated). Thereafter, the rim remained close to the substrate, but

the central portion of the animal moved further away, forming a "feeding pocket"

between the lower surface and the substrate. The secretion of a granule often was

followed within <2 sec by lysis of algae in the vicinity (Fig 8C, F). As algae lysed, they

became intensely stained with LipidTOX and FM1-43. Following lysis of algae, cells in

the central region of the animal began to display "churning" movements, slow swirling

movements of large groups of cells and, sometimes, faster oscillations of small groups

of cells (S11 and S12 Movies) [4]. Trypsin activity monitored with BZiPAR became

detectable ~ 1 minute after LC granule secretion and continued to increase over the

course of 3 to 10 minutes (Fig 8D, E, G; S10 Fig; S11 and S12 Movies). Algae that had

been lysed by the content of LC granules gradually disappeared but algae that were not

lysed remained intact (Fig 8G). Lysed algae that became trapped under the rim and

therefore were protected from enzymes in the feeding pocket did not disappear,

implicating digestive enzymes in the disappearance of the lysed algae. The partially

digested content of the lysed algae likely was pinocytosed by VEC and transferred to

lysosomes for intracellular digestion [41]. When the animal resumed gliding, the

remaining BZiPAR fluorescence diffused from the feeding pocket into the surrounding

seawater. Some animals pausing over algae did not secrete LC granules and,

consequently, no algae were lysed. Although these animals displayed movements

typical of animals feeding on algae, no trypsin activity was detected at their lower

surfaces (not illustrated).

Fig 8. Secretory behaviors of *Trichoplax* associated with external digestion of algae. (A-E) Confocal images of a *Trichoplax* feeding on *R. salina* algae; (F, G) normalized fluorescence intensity measured in selected regions. A starved animal was transferred to a cover glass chamber containing algae in seawater with a lipophilic dye (LipidTOX, red) that stains lipophil cell granules; FM1-43 (cyan), a membrane dye used here to visualize the secreted contents of lipophil cell granules; and a fluorescent indicator for trypsin activity (BZiPAR, green). The insets numbered with lowercase letters are enlarged view of rectangular region on a respective image numbered with an uppercase letter. (A, a) At t=-1.9 sec, the animal had ceased moving in a region containing algae (blue, phycoerythrin autofluorescence, arrows) and debris (cyan) representing algae remnants from an earlier feeding episode. The animal body is outlined white, and the feeding pocket is outlined red. (B, b) At t=0 sec, lipophil cell granule secretion was evident in the feeding pocket due to the sudden appearance of small (<5 µm) diffuse clouds and bright particles (arrowheads on b) of FM1-43-stained material. (C, c) At 5.7 sec, some algae near sites of lipophil granule secretion swelled, lysed, and became intensely stained with LipidTOX and FM1-43 (pink in merged, arrows in c). Other algae (blue) remained intact and were not stained with LipidTOX or FM1-43. (D, d) By 195.4 sec, diffuse BZiPAR fluorescence (green) filled the feeding pocket. The lysed algae no longer were visible, but the intact algae remained (arrows in d). (E) Intensity encoded BZiPAR fluorescence images show increasing trypsin activity between 70.6 and 233.5 sec. (F) Details of first events in feeding: Lipophil granules (cyan; the fluorescence profile obtained for the area outlined yellow in b) were secreted

approximately synchronously at 0 sec and this was followed by a rapid increase of

fluorescence in nearby algae (four fluorescence profiles obtained for four pink algal cells

in c; different shades of magenta). (G) Evidence of digestion: algae affected by lipophil

granules (different shades of magenta) burst and released their content. Secretion of

trypsin, as indicated by BZiPAR fluorescence (green; fluorescence profile obtained for

the region outlined red in B), began 40-50 sec after lipophil discharge and was

associated with a decline of fluorescence intensity of the lysed algal cells. Those algae

not affected by lipophil granules remained constant in intensity throughout the feeding

episode (two fluorescence profiles obtained for two individual algal cells; different

shades of blue). Scale bars 100 µm.

718 Discussion

We used FISH probes for genes specifically expressed in metacells identified by

analysis of single cell transcriptomes of *Trichoplax adhaerens* H1 (*TH1)* [6,30] to map

the distributions of different cell types in *TH1* whole mounts and to identify the cells in

dissociated cell cultures where their morphology could be visualized by differential

interference contrast microscopy and fluorescent labels used to identify hallmark

features. We found that two cell types in the central part of the ventral epithelium,

lipophil cells (LC) and a subset of the monociliated ventral epithelial cells (VEC),

participate in lysis and digestion of microalgae in animals pausing to feed. The dorsal

epithelium and the peripheral part of the ventral epithelium contain secretory cells that

express genes with sequences resembling precursors of glycoproteins and/or secretory

peptides implicated in defense in other animals. Electron microscopic studies of LC,

dorsal epithelial cells (DEC) and VEC revealed ultrastructural features of their secretory

granules that, together with the transcriptomes of these cell types, provide clues about

the compositions and biogenesis of their secretory granules. We identified metacells

representing mucocytes and found that mucocytes in the central part of the ventral

epithelium differ from those in the periphery. We mapped the distributions and identified

morphological features of seven peptidergic cell types. Our findings provide a more

detailed picture of the placozoan body plan (Fig 9) than was apparent from previous

microscopic studies and reveal cellular mechanisms placozoans use to obtain nutrients

and clues about their defenses against pathogens. We compare the sequences of the

main secreted proteins and expression profiles of single cell transcriptomes of four

placozoan species and confirm a close relationship between *Trichoplax adhaerens* H1

and H2, partial divergence of *Hoilungia hongkongensis H13* and greater divergence of

Cladtertia collaboinventa H23.

Fig 9. Summary diagrams of distributions of *Trichoplax adhaerens* secretory cell

types. (A) Secretory proteins, expressed specifically in DEC, pVEC, cVEC , mucocytes

, and lipophil cells (peptidergic cells represented in panel B are omitted in panel

A). Many central DEC co-express *Ta Intelectin* and the precursor of ELPE peptide but

peripheral DEC do not express *Ta Intelectin*. Peripheral VEC highly express precursors

of putative antimicrobial peptides (AMP's), while cVEC highly express digestive

enzymes. Lipophil cells and mucocytes differentially express their own specific genes

(see legend). (B) Secretory products of peptidergic cell types. Localization of peptidergic

cell types is based on measurements from animals labeled with *FISH* probes for peptide

precursors (Figs 6, 7). Peptidergic cells located in the central zone of the animal co-

express additional secretory products unlike the cells located at the periphery. Only

mucocytes in the central zone co-express *TaRWa*. Diagrams represent vertical cross

sections of a region encompassing ~20% of the diameter of an animal ~500 µm in

diameter. Cell dimensions are based on measurements from animals prepared for

transmission electron microscopy by high pressure freezing and freeze substitution

[16].

A subset of monociliated ventral epithelial cells secrete digestive enzymes

Probes for *Ta trypsin*, *Ta chymotrypsin* and *Ta sPLA2*, which are highly expressed in

metacells classified as "digestive gland cells" [6,30], labeled cells in the central part of

the ventral epithelium with morphological features typical of ventral epithelial cells

(VEC): they possessed an apical cilium and microvilli and had electron dense secretory

granules near their apical surface. Although genes associated with cilia were not

reported in these metacells, we found that they express genes implicated in cilia

structure and function.

We used fluorescent dyes that label the content of lipophil cell granules and a

fluorescent indicator for trypsin activity to monitor secretion by lipophil cells (LC) and

VEC in *Trichoplax* while they were consuming algae. After the animal crawled onto a

patch of algae and ceased moving, LC in the vicinity of algae secreted a large granule

whose content lysed nearby algae within less than 2 seconds[4]. Activation of the

fluorescent trypsin indicator became detectable after about one minute and the

fluorescence increased over the duration of the feeding episode. Detection of trypsin

activity provides indirect evidence that central VEC (cVEC) secrete digestive enzymes

during feeding episodes. Lysis of algae apparently is required to allow the digestive

enzymes to penetrate algae because intact algae were not decomposed even when

exposed to digestive enzymes in the feeding pocket.

Trichoplax VEC are thought to take up nutrients and further digest them intracellularly

because markers such as ferritin and fluorescent dextran were observed inside

pinocytotic and endosomal vesicles in VEC after addition to the ambient seawater

[38,39]. The gastrodermis in many animals with internal digestive systems contains

absorptive cells, called enterocytes, that take up nutrients released from food partially

digested by gland cells enzymes [41]. In *Trichoplax*, cVEC appear to combine functions

of digestive gland cells and absorptive cells; this could be the reason why little similarity

was found at the transcriptomic levels between placozoan digestive gland cells and

digestive gland cells in other species [6].

793   **Lipophil cells – a placozoan synapomorphy?**

The observation that the content of granules secreted by LC during feeding episodes in

Trichoplax rapidly lyses algae led us to search LC metacells for genes that might

encode proteins with sequences resembling those of pore-forming proteins [43–46]. We

found identical genes in *TH1* and *TH2* LC metacells that encode a protein with a signal

peptide and no transmembrane domain; this protein has no orthologs in the NCBI

database of non-redundant protein sequences and bears no known domains. The gene

is the most highly expressed gene in *TH2* LC metacells, and the second most highly

expressed gene in *TH1* LC metacells. The predicted protein is mainly composed of

helical regions with positively charged areas that might interact with membranes, like

the alpha-helical class of pore-forming proteins [47], nucleoporins [48], or ninjurin [49]. A

protein with high sequence similarity to the *TH1*/H2 genes is expressed in one of the

two metacell clusters classified as LC in *HH13* (LC2) and an ortholog (e^{-45}) is expressed

in both LC1 and LC2 metacells in *CH23*. It is likely that this secretory protein is

packaged in LC granules and participates in permeabilizing prey organisms. A thin

section from a LC revealed an apical granule with an open fusion pore, suggesting that

LC granule secretion occurs by conventional exocytosis.

The outer part of the LC apical granule binds osmium, as evident by electron

microscopy in thin sections from samples fixed with osmium, indicating the presence of

unsaturated lipids [31]. In thin sections of LC from frozen and freeze substituted

samples, the apical granules contained variable amounts of electron dense material as

well as small vesicles whose content resembled cytoplasm. Granules deeper in the cell

bodies of LC contained progressively less electron dense material and larger membrane

enclosed vesicles as well as finger-shaped protrusions from the membrane surrounding

the granule. The ultrastructure of LC apical granules bears some resemblance to that

of lytic granules in rat natural killer cells (NKC), which likewise contain electron dense

cores and small membrane-enclosed vesicles [50–52]. Natural killer cell granules are

considered secretory lysosomes or lysosome related organelle (LRO). The affinity of

acidophilic Lysotracker dyes for LC granules demonstrates that their content is acidic

[4,16], like the content of NKC granules.

The presence of lipids in LC granules is intriguing and not a feature shared with NKC

granules, although LRO in other types of cells sometimes contain lipids [53,54].

Lipophil metacells highly express several fatty-acid binding proteins [6] that might serve

as chaperones to deliver lipids to their granules. *Trichoplax* can lyse and consume

cyanobacteria in addition to microalgae [4], indicating that the content of LC granules

can penetrate both bacterial and eukaryotic membranes. The identities of the lytic

components and their molecular targets in microalgae and cyanobacteria are important

questions that remain to be addressed.

The gene expression profiles of LC appear unlike those of cells in the digestive systems

of other animals [6]. However, many animal lineages have evolved unique types of

secretory cells that they use to capture, incapacitate, or digest prey and to defend

against predators. Well documented examples include cnidarian cnidocytes [55–58],

ctenophore colloblasts [57,59,60], and venom secreting cells in arthropods, for example

[61,62].

Cells in the peripheral part of the ventral epithelium express

putative antimicrobial peptides

Metacells classified as "epithelial" or "lower epithelial" in scRNAseq studies of *TH1*

[6,30] based on expression of genes associated with the structure and function of cilia

highly express genes with structural similarities to precursors of arminins, antimicrobial

peptides found in *Hydra* endodermal cells [32,63,64]. Probes for the mRNA encoding

two of these arminin-like genes (referred to here as *AMP1* and *AMP2*) labeled

monociliated cells in the peripheral part of ventral epithelium (pVEC). These cells had

ultrastructural features like digestive cVEC but possessed darker and more numerous

secretory granules. All *TH1* metacells classified as "lower epithelial" express both *AMP1*

and *AMP2* and a third arminin-like gene, *Ta 60631 (AMP3)*. Metacells classified as

"lower epithelial" in *TH2* express genes that are nearly identical to *AMP1* and *AMP3* and

a subset of them express an ortholog of *AMP2*, but *HH13* and *CH23* "lower epithelial"

metacells express only an ortholog of *AMP2*.

A probe for a different arminin-like *Trichoplax* gene, *AMP4*, labeled cells in a narrow

zone in the ventral epithelium ~15 to 30 μm from the rim. Their positions correspond to

those of cells classified as Type 1 gland cells in ultrastructural studies of *Trichoplax* [18].

The metacells that contain *AMP4* also express genes encoding an intelectin and an

uncharacterized secretory protein, *Ta 63786* that were present only in this metacell

cluster.

The genes encoding *AMP1*, *AMP2*, and *AMP3* are the three most highly expressed

genes in *Trichoplax* H1 "lower epithelial" metacells and the *TH2* orthologs of *AMP1* and

AMP3 are the most highly expressed genes in *TH2* "lower epithelial" metacells.

Similarly, arminins are among the most highly expressed genes in most *Hydra* species

that have been studied [63,64]. Labeling with mRNA probes for nine arminin paralogs in

Hydra wholemounts showed that eight were expressed in the gastrodermis (endoderm)

and one in the epidermis. *Hydra vulgaris* metacells containing arminins were classified

as "endodermal epithelial" [65] and likely correspond to absorptive gastrodermal cells

(enterocytes). Likewise, AMP expression in *Trichoplax* is mostly located to the ventral

epithelium, an anatomical structure involved in digestion and presumably corresponding

to cnidarian gastroderm[66].

Two types of mucocytes arrayed in distinct patterns in the ventral epithelium

We identified metacells representing mucocytes in whole mounts and dissociated cell

cultures of *TH1* based on labeling with fluorescent WGA and a probe for an

oligosaccharide binding protein, *Ta 63702*, that was highly expressed in a single *TH1*

metacell (C36) [30]. Mucocytes were abundant in the peripheral part of the ventral

epithelium and more sparsely distributed further in the interior, as reported [18].

Mucocytes in the central region of the ventral epithelium, but not those in the periphery,

co-expressed a gene for prepropeptide that is predicted to produce RWamide peptides.

We identified metacells in *TH2*, *HH13* and *CH23* that express a gene identical to *Ta*

63702 and a subset of these metacells contain a gene encoding the RWamide

prepropeptide. Although no *Ta RWa* gene was found in the second RNAseq analysis of

TH1 [6], we identified metacells that likely represent central and peripheral mucocytes in

this dataset. All metacells representing mucocytes in the more recent RNAseq datasets

contain a gene encoding an intelectin (Ta 62229) and a protein with VWD, TIL, C8 and

PTS domains characteristic of gel-forming mucins [37]. The RWamide peptides and

intelectin likely are packaged and secreted along with mucus since mucocytes contain

only one type of secretory granule [18]. Animal mucosa often contain lectins and

antimicrobial peptides [67]. However, the predicted products of the *Ta RWa* do not have

electrostatic properties typical of antimicrobial peptides. It is possible that the peptides

are used for intercellular signaling, although synthetic RWa peptides had no apparent

effect on the behavior of *Trichoplax* when added to their culture dishes [23]

Genes encoding gel-forming mucins with VWD, TIL, C8 and PTS domains were found in

single cell transcriptomes of the cnidarian *Hydra vectensis* and the ctenophore *M. leidyi*

[30,68]. Mucus-like substances have been observed on epithelia of adult demosponges

[69] and cells resembling mucocytes were found by electron microscopy in

metamorphosing larvae of several classes of sponges[70–72]. The distribution of

mucus-secreting cells across animal phyla and presence of mucus-like genes in

Choanoflagellates [73] suggests that the common ancestor of Metazoa may have

possessed cells that secreted a mucus-like substance.

907 Dorsal epithelial cells secrete lectins implicated in defense

Cells with morphological characteristics of DEC were labeled with a probe for mRNA

encoding the ELPE prepropeptide and many of them were labeled with a probe for an

intelectin, *Ta 60661*. The *TH1* metacells expressing these genes were classified as

"epithelial" or "upper epithelial" [6,30] and include genes encoding ten intelectins that

are not expressed in any other metacells. *Trichoplax* H2 "upper epithelial" metacells

contain identical intelectin genes. *Hoilungia* H13 "upper epithelial" metacells contain

nearly identical orthologs of five of them and *CH23* "upper epithelial" metacells contain

four of them. These intelectins likely are packed inside the electron dense granules in

DEC, like intelectins in goblet cells in mammalian intestine [74]. These metacells also

express a gene encoding a membrane associated mucin 4-like glycoprotein that may be

a component of the glycocalyx observed on the apical surfaces of DEC by electron

microscopy.

Intelectins in chordate epithelia are thought to protect against invasion by pathogens by

binding glycans on their surfaces and aggregating the cells[36]. Less is known about

the prevalence and functions of intelectins in non-chordates, although the available

evidence suggests that the primary function is defense [36,75].

Peptidergic cell types

We mapped the positions and studied the morphology of cells expressing seven of the

fourteen prepropeptides identified by scRNAseq in *TH1* [6]. Peptides that are the

predicted/possible products of the six of these prepropeptides (WPPF, ELPE, SIFGa,

FFNPa, LF, TaELP) elicited changes in the behavior of *Trichoplax* when added to their

culture dishes[22,23]. We anticipated that the positions of the peptidergic cells and the

responses elicited by peptides they secrete might provide clues about their functions.

Cells that expressed the *Ta SIFGa* prepropeptide were most prevalent in a narrow zone

near the rim of the dorsal and ventral epithelium but were also present in more central

regions of the epithelia. The *Ta SIFG+* cells in the central part of the ventral epithelium

co-expressed the *Ta FFNPa* peptide precursor. Addition of low concentrations (<500

nM) of SIFGa to dishes containing *Trichoplax* rapidly (<1 min) elicited contraction of the

entire animal. Higher concentrations of the peptide caused the animal to fold up and

detach from the substrate[22,23]. The peptide FFNPamide elicited an expansion of the

area of the animal and an increase in the frequency of spontaneous pauses in

movement. The latencies of the responses to FFNPa were longer (> 7 minutes) than

those of the contractions elicited by SIFGamide, suggesting that simultaneous secretion

of SIFGa and FFNPa by cells that co-express these peptides might elicit contraction

followed by relaxation. Detachment from the substrate elicited by secretion of

endogenous SIFGa may allow the animal to move between the substrate and air/water

interface, as animals maintained in culture often do. The location of the cells that co-

expressed *Ta SIFGa* and *Ta FFNPa* suggests that they may participate in orchestrating

behaviors that o*cc*ur during feeding.

Cells expressing the gene for LF prepropeptide were in a narrow zone near the rim of

the dorsal epithelium and distributed throughout the central part of the ventral

epithelium. The *Ta LF*+ cells in the ventral epithelium co-expressed a secreted astacin-

like metalloendopeptidase. Applying LF to *Trichoplax* elicited a large expansion in the

animals' area. In addition, animals that were translocating stopped and rotated in place

[23], indicating a change in the collective behavior of the monociliated VEC that propel

crawling. Secretion of endogenous LF by cells located at the rim could promote

adhesion of the rim to the substrate by relaxing the contractile cytoskeletons of cells in

the rim.   The location of the cells that co-expressed *Ta LF* and astacin *Ta 26557*

suggests that they may have roles in feeding.

Cells co-expressing the *TaELP* prepropeptide and an astacin-like metalloendopeptidase

were in the ventral epithelium near the rim. Their locations did not correspond to those

of cells labeled with antiserum against *TaELP* prepropeptide or against the peptide

YPFFamide (human endomorphin 2), which instead labeled mucocytes[18,22]. We

previously reported that adding YPFFamide or QDYPFFamide to a dish containing

moving *Trichoplax* consistently caused them to stop moving and arrested ciliary beating,

but the predicted products of the *TaELP* prepropeptide (QDYPFFGN or pQDYPFFGN)

arrested movement of fewer than half of the animals tested [22]. The observation that

antiserum against YPFFamide did not label the cells that express *TaELP* prepropeptide

suggests that these cells do not produce the peptide QDYPFFamide and casts doubt on

the idea that the function of these cells is to detect food and arrest movement during

feeding episodes[22]

The *Ta ELPE* prepropeptide was expressed in a large fraction of DEC and by cells near

the rim of the ventral epithelium. Cells expressing the *Ta WPPF* prepropeptide were

interspersed among DEC in the central part of the dorsal epithelium. Bath application of

ELPE or WPPF were reported to elicit behaviors reminiscent of behaviors observed in

animals feeding on biofilms: periodic cessation of gliding a*cc*ompanied by churning

movements of cells in central regions of the ventral epithelium[23]. The ELPE peptide

elicited a small expansion in the animals' area, comparable to that observed in animals

pausing to feed, whereas the WPPF peptide elicited a much larger expansion. The

positions of the cells that express *Ta ELPE* or *Ta WPPF* make it unlikely that their

primary function is to initiate feeding behaviors.

We identified orthologs for four of the seven studied prepropeptides across all four

placozoan species. Additionally, the TaELPE precursor was detected only in TH1 and

TH2. This pattern of phylogenetic distribution and sequence divergence reflects what is

observed across cnidarian classes and bilaterian phyla [33,34]. These findings further

emphasize the biological divergence among placozoan species, despite their

morphological similarities. The morphology and expression profiles of placozoan

peptidergic cells closely resemble those of sensory-secretory cells found in Cnidaria

and Bilateria [6, 16, 18]. Further studies on peptide receptors, along with their

associated signaling pathways and cellular interactions, are necessary to better

understand their biological roles and functions.

Conclusions

Placozoa provide an opportunity to explore the biology of animals with body plans and

lifestyles reminiscent of ancestors of present day metazoa [76–81]. The availability of

single cell transcriptomes of four placozoan species [6,30] provides a rich resource for

investigating the functions and evolutionary histories of their cell types. Building on this

foundation, we linked single-cell transcriptomic data with previously described

morphological cell types and explore the roles of their secretory products.

1004

Placozoan secretory cell types share morphological features with those found in other

metazoans, with several expressing homologs of functionally characterized proteins.

The ventral epithelial cells (VECs) in the central part of the animal secrete digestive

enzymes, similar to zymogen gland cells in cnidarians and bilaterians. In contrast,

peripheral VECs express putative antimicrobial peptides (AMPs), akin to gastrodermal

cells in various animals. Both central and peripheral VECs possess a cilium and

microvilli and engage in pinocytosis to absorb nutrients released from partially digested

food, resembling enterocytes in cnidarians and bilaterians. Defensive epithelial cells

(DECs) in placozoans express glycoproteins associated with pathogen defense across

metazoans, while mucocytes produce orthologs of gel-forming mucins, a key

component of mucus. Additionally, placozoans exhibit a variety of peptidergic cell types,

each characterized by distinct distributions and gene expression profiles. Lipophil cells

secrete lytic substances like certain secretory cells in other animals. However, the

composition of their secretory granules appears unique to placozoans. Collectively, our

work highlights the diversity of secretory cells and their products in Placozoa, a phylum

with a unique phylogenetic position within the metazoan tree of life. These findings

provide valuable insights for comparative studies across animals, advancing our

understanding of the evolution of secretory cells.

1023

Methods

Animals

Trichoplax adhaerens (Schultze, 1883) of the Grell (1971) strain, a gift from Leo Buss

(Yale University), were kept in Petri dishes with artificial seawater (ASW; Instant Ocean,

Blacksburg, VA, USA) supplemented with 1% Micro Algae Grow (Florida Aqua Farms,

Dade City, FL, USA) and red algae (*Rhodomonas salina*, Provasoli-Guillard National

Center for Culture of Marine Plankton, East Boothbay, ME, USA), as described

previously [16]. Water was partially changed once a week.

1032

1033   Cell dissociation

1034   To prepare dissociated cells, a group of animals was rinsed in calcium and magnesium

1035   free ASW (calcium-free ASW)[82] and then incubated in 0.25% trypsin in calcium-free

1036   ASW for 2 h. The animals were transferred to normal ASW and triturated with a glass

1037   Pasteur pipette until the suspension was homogeneous.

1038

1039   Fluorescence *in situ* hybridization

In situ hybridization was performed with probes and reagents from Advanced Cell

Diagnostics, Inc. (ACD, Hayward, CA, USA) using protocols developed to optimize

staining in *Trichoplax* [18,24]. The RNA sequences were retrieved from PubMed or from

a *T. adhaerens* transcriptome database, access to which we were granted by Adriano

Senatore (University of Toronto). RNAscope probes for multiplex fluorescence *in situ*

hybridization were designed by ACD. Catalog names and numbers, gene names, and

annotations, and mRNA accession numbers are listed in Table 1. Further details are

available in the ACD product catalog.

1048

1049   **Table 1. RNAscope probes for multiplex fluorescence *in situ* hybridization.**

1050

RNAscope probe	Catalog #	Gene	Annotation	Accession number
Ta-57870-C2	561141-C2	Ta-57870	PLA2	XM_002113901.1
Ta-63128	561031	Ta-63128	Trypsin	XM_002115568.1
Ta-Chymotrypsin-C2	572831-C2	Ta-63088	Chymotrypsin	XM_002109248.1
Ta-PhospholipaseA2	572841	Ta-63140	PLA2	XM_002116279.1
Ta-57870-C2	561141-C2	Ta-57870	PLA2	XM_002113901.1
Ta-56030-C2	56030-C2	Ta-56030	AMP2	XM_002111672.1
Ta-55945-C3	55945-C3	Ta-55945	AMP1	XM_002111637.1
Ta-64402-C2	1313051-C2	Ta- 64402	AMP3	XM_002118041.1
Ta-58643-C1	1138211-C1	Ta-58643	Tetraspanin	XM_002114763.1
Ta-63786	1209181-C1	Ta-63786	Ta Sec Prot	XM_002110591.1
Ta-64037-C2	1222681-C2	Ta-64037	Ta Sec Prot	XM_002113441.1
Ta-63996-C2	1138221-C2	Ta-63996	Ta Sec Prot	XM_002113590.1
Ta-29105-C3	1138231-C3	Ta-29105	Tansporter	XM_002115362.1

Ta-63702-C3	823661-C3	Ta-63702	Oligosac BP SIFGamide	XM_002110164.1
Ta-60185-C2	487491-C2	Ta-60185	Pre	XM_002116138.1
None	856541-C2	EST	RWamide Pre FFNPamide	GR951923.1
Ta-63942-C2	823731-C2	Ta-63942	pre	XM_002112788.1
Ta-51275	561121	Ta-51275	LF Pre	XM_002108379.1
Ta-51275-C3	561121-C3	Ta-51275	LF Pre	XM_002108379.1
Ta-YPFFamide-O1-C2	1228931-C2	AQX36197.1	TaELP Pre	KY675296.1
Ta-56359	561051	Ta- 56359	WPPF Pre	XM_002112355.1
Ta-60185	487491	Ta-60185	SIFGa Pre	XM_002116138.1
Ta-64280-C3	8236681-C3	Ta-64280	ELPE Pre	XM_002116364.1
Ta-26557-C3	561111-C3	Ta-26557	Astacin	Tricho_evg1529192
Ta-54934	560991	Ta-54934	Astacin	XM_002110640.1
Ta-60661-C3	1241961-C3	Ta-60661	Intelectin	XM_002116619.1
Ta-60661-C2	1217761-C2	Ta-60661	Intelectin	XM_002116619.1
3-Plex Neg. Control	320871			

1051

1052   Abbreviations: Ta – Triaddraft-; PLA2 – PhopholipaseA2; AMP– Putative antimicrobial

1053   peptide; Ta Sec Prot – Predicted *T. adhaerens* secretory protein; Oligosac BP –

1054   Oligosac*ch*aride binding protein; Pre – Prepropeptide

1055

Animals and dissociated cells were transferred to silanized cover slips or Superfrost

Plus Gold glass slides (Thermo Fisher Scientific, Pittsburgh, PA, USA) with a drop of

ASW mixed 1:1 with 0.97 M mannitol (in water). Samples were left on cover slips/slides

to adhere for about 2 h. Then the liquid was blotted, and the cover slips/slides were

plunged into prechilled tetrahydrofuran on dry ice and kept overnight. Wholemount

samples were transferred to 3% acetic acid in methanol at −20 °C for 30 min followed

by a mixture of formalin and methanol (1:10), initially at −20 °C and then at room

temperature (RT) for 30 min. The wholemount samples were rinsed twice in methanol,

dried for 5 min and then treated with Protease IV for 30 min. The dissociated cell

samples were transferred directly to a mixture of formalin and methanol (1:10), initially

at −20°C and then at RT for 30 min. The cell samples were then rinsed once with

methanol, twice with ethanol followed by descending concentrations of ethanol in PBS

(70% and 50%) and PBS. The cell samples were treated with Protease III diluted 1:15 in

PBS for 15 min. Hybridization was performed with RNAscope Fluorescent Multiplex

Reagent Kit (# 320850) a*cc*ording to supplier's instructions. The negative control 3-Plex

Negative Control Probe gave no labelling as did the sense probes for two of the genes

studied (TaYPFFamide, Cat. #488721; and Ta 26557, Cat #1222671-C3). Samples

were counterstained with 1:200 wheat germ agglutinin (WGA) conjugated to Alexa 647

(# <u>W32466</u>, Thermo Fisher Scientific, Waltham, MA) or CF405M (# 29028, Biotium,

Hayward, CA, USA) and/or DAPI, and mounted in ProLong™ Gold antifade reagent

(# <u>P36934</u>, Invitrogen, Eugene, OR, USA). Dissociated cell preparations were also

subjected to immunolabelling for tubulin after RNAscope steps to visualize cilia. For

that, samples rinsed with wash buffer after the last (AMP4) hybridization step were

further washed twice with PBS (pH7.4) and once with blocking buffer (BB, 3% goat

serum, 2% horse serum, 1% BSA in PBS). Then, primary anti-acetylated tubulin mouse

antibody (#T7451; Sigma) diluted 1:500 in BB was applied. After overnight incubation

with the primary antibody, samples were washed with PBS and incubated in secondary

Atto-655 goat anti-mouse antibody (#50283; Sigma) or Alexa 555 goat anti-mouse

(#A21422; ThermoFisher) diluted 1:200 in blocking buffer. Finally, samples were rinsed

from secondary antibody with PBS, briefly incubated in DAPI, and mounted in

ProLong™ Gold antifade reagent (ThermoFisher). The samples were examined with a

Plan-Apochromat 40X NA 1.3 or 63X NA 1.4 Plan-Apochromat objective on a LSM 800

or LSM 880 AIry laser scanning confocal microscope with DIC and fluorescence optics

(Carl Zeiss Microscopy, LLC). The areas of dissociated cells visualized with DIC optics

were estimated by measuring the length of the long axis of the cell body and the

perpendicular axis and calculated using the equation for an ellipse. Cells were

considered labeled if they possessed at least two fluorescent grains. To demonstrate

expression patterns in wholemounts, we created xy maximum intensity projections in

Zen software (Carl Zeiss Microscopy, LLC). In order to show the distribution of a signal

across the animal thickness, we made xz projections for selected strip regions,

indicated on the xy images with a narrow box. Labelling with each probe was done at

least twice; the results of independently repeated experiments were similar.

1098

Time-Lapse Microscopy of Living Animals

1099

1100

To monitor secretion of lipophil cell granules and trypsin during feeding episodes,

animals first were transferred to a 35 mm petri dish containing ASW and kept for one to

three hours and then incubated for 10 minutes in ASW containing FM1-43 (N-(3-

Triethylammoniumpropyl)-4-(4-(Dibutylamino) Styryl) Pyridinium Dibromide; Thermo

Fisher; #T3163; 1µg/ml) and LipidTOX (HCS LipidTOX Deep Red neutral lipid stain;

Thermo Fisher; #34477; 3 µl/ml). The animals were transferred to a RC-40LP chamber

containing ASW with FM1-43, LipidTOX, BZiPAR (Rhodamine 110, bis-(N-CBZ-L-

isoleucyl-L-prolyl-L-arginine amide), dihydrochloride; Biotium; #10208; 8 µM) and

Rhodomonas salina algae. LipidTOX stains intact lipophil cell granules and FM1-43

stains the contents of secreted granules, as described previously [4]. BZiPAR is a

fluorescence indicator for trypsin activity [40] The behavior of the animals was

monitored with a 32-channel spectral detector and a 10X NA 0.45 Plan-Apochromat

objective on a LSM880 confocal microscope with 488 nm, 561 nm and 647 nm

illumination. Images (512 X 512 bits) were captured at ~1.7 frames/second. Reference

spectra used for linear unmixing were collected from the following samples: (1) FM1-43,

ventral epithelium of an animal labeled with FM1-43; (2) LipidTOX, lipophil cell granule

in an animal labeled with LipidTOX; (3) algae; autofluorescence; (4) BZiPAR; seawater

containing BZiPAR and 0.25% trypsin. Bicubic interpolation was used to enlarge the

images and reduce pixilation. Still images used for illustration in Fig 8 were created by

averaging two su*cce*ssive frames to improve signal. Fiji software was used to build

fluorescence profiles shown in Fig 8.

1122

Electron microscopy
1123

1124   For transmission electron microscopy (TEM), scanning electron microscopy (SEM), and

1125   electron microscope tomography animals were high pressure frozen, freeze-substituted

1126   and embedded as described previously [16]. WGA-nanogold (EY Laboratories, San

1127   Mateo, CA, USA) labelled thin sections were prepared as described[18]

1128

1129   For transmission electron microscopy, the ultrathin sections were observed in a JEOL

1130   200-CX (JEOL, Japan) at 120–200 kV and were imaged with an AMT camera mounted

1131   below the microscope column. Image processing and color intensity measurement was

1132   done with Fiji.

1133

1134   The protocol for serial sectioning followed by SEM imaging in backscatter mode was

1135   published elsewhere [18]. Series of these images were used to study the nature of

1136   lipophil granules protrusions and to count granules in the pVEC and cVEC.

1137

For EM tomography, we used ~100 nm sections and a transmission electron

microscope, JEOL 1400 (JEOL, Japan) at an a*cce*lerating voltage of 120kV. Images

were taken at ~1° tilt intervals from -69° to 69° at a single tilt axis at the magnification

x5000. The tilt images were aligned using an electron tomography software package,

EM3D (em3d.org), upgraded by Jae Hoon Jung and Eun Jin Choi in Dr. Reese's

laboratory, and reconstructed to generate a reconstructed volume or tomogram by

Simultaneous Iterative Reconstruction Technique (SIRT) with 10 iterations using the

algorithm implemented in a tomography program, called VEM (Volume Electron

Microscopy), developed by Jae Hoon Jung and Eun Jin Choi. The examination of the

virtual slices was carried out using both EM3D and VEM. The virtual slices through the

tomogram were one-voxel thick, which is 1.8 nm.

1149

To prepare freeze fractured samples, *T. adhaerens* were placed on the surface of a

gelatin cushion and directly frozen by contact with a sapphire disk glued to the surface

of a copper block cooled to $-186\,°C$ using a LifeCell CF-100 slam-freezing machine.

Samples were transferred to a Balzer freeze-fracture apparatus. Replicas were

prepared by freeze-fracturing the specimen at $-110\,°C$ and at a vacuum of 10^{-} Torr,

rotary shadowing with platinum and carbon, and backing the replica with carbon. The

replicas were cleaned with sodium hypochlorite, mounted on grids, and examined using

a JEOL 200-CX at 120 kV.

1158

Protein/peptide predictions

Deep TMHMM [83] was used to predict signal peptides and transmembrane alpha

helices. InterPro online tool [84]was used to predict protein topology and conserved

domains. AlphaFold protein database [85] was used to predict secondary structures of

the *Trichoplax* proteins. ChimeraX-1.5 [86,87]was used to render a secondary structure

of the proteins and color protein surface by electrostatic potential.

1165

Regions of prepropeptides were annotated and classified into signal peptide region

using SignalP6.0 [https://pubmed.ncbi.nlm.nih.gov/34980915/], prohormone convertase

cleavage sites (dibasic residues), mature peptide (based on characteristics of cnidarian

neuropeptide processing [88]. C-terminal glycine residues were predicted to be

processed into amide-groups, and N-terminal glutamates were predicted to be

processed into pyroglutamates.

1172

1173 Quantitative image processing and Statistics

We used Paleontological Statistics (PaSt), freely available software [89], to do statistical

comparisons and build boxplots. Mann-Whitney test was applied, since many variables

did not follow the Gaussian distribution. For multiple samples, we used an ANOVA test

followed by Mann-Whitney post hoc pairwise test with the Bonferroni correction. In

boxplots, the box represents 25-75 quartiles, and the *median* is shown with a horizontal

line inside the box. "Whiskers" show the minimal and maximal values excluding

outliers. Each value is plotted as a dot on top of a boxplot. A*c*curate area-proportional

Venn diagrams were drawn using eulerAPE program [90] or PaSt.

1182

1183 Gene expression profiles

The data for S1Fig was downloaded from [30]. The fold change and percentage

expression were extracted for the selected genes across all the metacells and

visualized using the tidyverse package in R. Gene expression heatmaps were

generated from [https://sebelab.crg.eu/placozoa_cell_atlas/] using orthologous genes

from TH1, TH2, HH13, and CH23. The genes and meta cells were manually annotated

using InkScape.

1190

Acknowledgements

We thank the Electron Microscopy Facility of the National Institute of Neurological

Disorders and Stroke, the Central Microscopy Facility of the Marine Biological

Laboratory, Christine A. Winters (NINDS) and Katherine Hammar (MBL) for sample

preparations and technical support. We thank Professor Adriano Senatore for valuable

comments on the manuscript. This work was supported by the Intramural Research

Program of the National Institute of Neurological Disorders and Stroke, National

Institutes of Health, Bethesda, MD, United States of America.

1199

References

1.   Grell KG, Ruthmann A. Placozoa. In: Harrison FW, Westfall JA, editors.
Microscopic Anatomy of Invertebrates. New York: Wiley-Liss; 1991. pp. 13–27.
Available:
http://download.springer.com/static/pdf/813/art%3A10.1007%2Fs00427-004-
0390-
8.pdf?auth66=1363711205_d5c9623bde6301d7ab57239eaefe9a93&ext=.pdf
2.   Pearse VB, Voigt O. Field biology of placozoans (Trichoplax): distribution,
diversity, biotic interactions. Integr Comp Biol. 2007;47: 677–92.
doi:10.1093/icb/icm015
3.   Eitel M, Schierwater B. The phylogeography of the Placozoa suggests a taxon-
rich phylum in tropical and subtropical waters. Mol Ecol. 2010;19: 2315–27.
doi:10.1111/j.1365-294X.2010.04617.x
4.   Smith CL, Pivovarova N, Reese TS. Coordinated feeding behavior in trichoplax,
an animal without synapses. Steele RE, editor. PLoS One. 2015;10: e0136098.
doi:10.1371/journal.pone.0136098

5.   Srivastava M, Begovic E, Chapman J, Putnam NH, Hellsten U, Kawashima T, et
al. The Trichoplax genome and the nature of placozoans. Nature. 2008;454: 955–
60. doi:10.1038/nature07191
6.   Najle SR, Grau-Bové X, Elek A, Navarrete C, Cianferoni D, Chiva C, et al.
Stepwise emergence of the neuronal gene expression program in early animal
evolution. Cell. 2023;186: 4676-4693.e29. doi:10.1016/j.cell.2023.08.027
7.   Schierwater B, DeSalle R. Placozoa. Current Biology. 2018;28: R97–R98.
doi:10.1016/J.CUB.2017.11.042
8.   Moroz LL, Kocot KM, Citarella MR, Dosung S, Norekian TP, Povolotskaya IS, et
al. The ctenophore genome and the evolutionary origins of neural systems.
Nature. 2014;510: 109–114. doi:10.1038/nature13400
9.   Pisani D, Pett W, Dohrmann M, Feuda R, Rota-Stabelli O, Philippe H, et al.
Genomic data do not support comb jellies as the sister group to all other animals.
Proceedings of the National Academy of Sciences. 2015;112: 201518127.
doi:10.1073/pnas.1518127112
10.   Ryan JF, Pang K, Schnitzler CE, Nguyen A-D, Moreland RT, Simmons DK, et al.
The genome of the ctenophore Mnemiopsis leidyi and its implications for cell type
evolution. Science. 2013;342: 1242592. doi:10.1126/science.1242592
11.   Schultz DT, Haddock SHD, Bredeson J V., Green RE, Simakov O, Rokhsar DS.
Ancient gene linkages support ctenophores as sister to other animals. Nature.
2023;618: 110–117. doi:10.1038/s41586-023-05936-6
12.   Simion P, Philippe H, Baurain D, Jager M, Richter DJ, Di Franco A, et al. A Large
and Consistent Phylogenomic Dataset Supports Sponges as the Sister Group to
All Other Animals. Current Biology. 2017;27: 958–967.
doi:10.1016/j.cub.2017.02.031
13.   Tessler M, Neumann JS, Kamm K, Osigus HJ, Eshel G, Narechania A, et al.
Phylogenomics and the first higher taxonomy of Placozoa, an ancient and
enigmatic animal phylum. Front Ecol Evol. 2022;10: 1016357.
doi:10.3389/FEVO.2022.1016357/BIBTEX
14.   Thiemann M, Ruthmann A. Alternative modes of asexual reproduction in
Trichoplax adhaerens (Placozoa). Zoomorphology. 1991;110: 165–174.
doi:10.1007/BF01632872
15.   Signorovitch AY, Dellaporta SL, Buss LW. Molecular signatures for sex in the
Placozoa. Proc Natl Acad Sci U S A. 2005;102: 15518–22.
doi:10.1073/pnas.0504031102
16.   Smith CL, Varoqueaux F, Kittelmann M, Azzam RN, Cooper B, Winters CA, et al.
Novel cell types, neurosecretory cells, and body plan of the early-diverging
metazoan Trichoplax adhaerens. Current Biology. 2014;24: 1565–1572.
doi:10.1016/j.cub.2014.05.046
17.   Gruber-Vodicka HR, Leisch N, Kleiner M, Hinzke T, Liebeke M, McFall-Ngai M, et
al. Two intracellular and cell type-specific bacterial symbionts in the placozoan
Trichoplax H2. Nature Microbiology 2019 4:9. 2019;4: 1465–1474.
doi:10.1038/s41564-019-0475-9
18.   Mayorova TD, Hammar K, Winters CA, Reese TS, Smith CL. The ventral
epithelium of trichoplax adhaerens deploys in distinct patterns cells that secrete

digestive enzymes, mucus or diverse neuropeptides. Biol Open. 2019;8.
doi:10.1242/bio.045674

19. Romanova DY, Varoqueaux F, Daraspe J, Nikitin MA, Eitel M, Fasshauer D, et al.
Hidden cell diversity in Placozoa: ultrastructural insights from Hoilungia
hongkongensis. Cell Tissue Res. 2021;385: 623–637. doi:10.1007/S00441-021-
03459-Y

20. Eitel M, Francis WR, Varoqueaux F, Daraspe J, Osigus H-J, Krebs S, et al.
Comparative genomics and the nature of placozoan species. Tyler-Smith C,
editor. PLoS Biol. 2018;16: e2005359. doi:10.1371/journal.pbio.2005359

21. Bull MS, Prakash VN, Prakash M. Ciliary flocking and emergent instabilities
enable collective agility in a non-neuromuscular animal. 2021 [cited 20 May 2024].
Available: https://arxiv.org/abs/2107.02934v1

22. Senatore A, Reese TS, Smith CL. Neuropeptidergic integration of behavior in
Trichoplax adhaerens, an animal without synapses. Journal of Experimental
Biology. 2017;220: 3381–3390. doi:10.1242/JEB.162396

23. Varoqueaux F, Williams EA, Grandemange S, Schierwater B, Jeke, Fasshauer D.
High Cell Diversity and Complex Peptidergic Signaling Underlie Placozoan
Behavior. Current Biology. 2018;28: 3495–3501. doi:10.1016/j.cub.2018.08.067

24. Mayorova TD, Hammar K, Jung JH, Aronova MA, Zhang G, Winters CA, et al.
Placozoan fiber cells: mediators of innate immunity and participants in wound
healing. Sci Rep. 2021;11: 23343. doi:10.1038/S41598-021-02735-9

25. Mayorova TD, Smith CL, Hammar K, Winters CA, Pivovarova NB, Aronova MA, et
al. Cells containing aragonite crystals mediate responses to gravity in Trichoplax
adhaerens (Placozoa), an animal lacking neurons and synapses. Steele RE,
editor. PLoS One. 2018;13: e0190905. doi:10.1371/journal.pone.0190905

26. Wong YY, Le P, Elkhatib W, Piekut T, Senatore A. Transcriptome profiling of
Trichoplax adhaerens highlights its digestive epithelium and a rich set of genes for
fast electrogenic and slow neuromodulatory cellular signaling. 2019.
doi:10.21203/RS.2.14504/V1

27. Ringrose JH, van den Toorn HWPP, Eitel M, Post H, Neerincx P, Schierwater B, et
al. Deep proteome profiling of Trichoplax adhaerens reveals remarkable features
at the origin of metazoan multicellularity. Nat Commun. 2013;4: 1408.
doi:10.1038/ncomms2424

28. Jékely G. Global view of the evolution and diversity of metazoan neuropeptide
signaling. Proc Natl Acad Sci U S A. 2013;110: 8702–7.
doi:10.1073/pnas.1221833110

29. Nikitin M. Bioinformatic prediction of Trichoplax adhaerens regulatory peptides.
Gen Comp Endocrinol. 2014;212: 145–155. doi:10.1016/j.ygcen.2014.03.049

30. Sebé-Pedrós A, Chomsky E, Pang K, Lara-Astiaso D, Gaiti F, Mukamel Z, et al.
Early metazoan cell type diversity and the evolution of multicellular gene
regulation. Nat Ecol Evol. 2018;2: 1176–1188. doi:10.1038/s41559-018-0575-6

31. Belazi D, Solé-Domènech S, Johansson B, Schalling M, Sjövall P. Chemical
analysis of osmium tetroxide staining in adipose tissue using imaging ToF-SIMS.
Histochem Cell Biol. 2009;132: 105–115. doi:10.1007/S00418-009-0587-
Z/FIGURES/8

32.   Augustin R, Anton-Erxleben F, Jungnickel S, Hemmrich G, Spudy B, Podschun R,
1307   et al. Activity of the novel peptide arminin against multiresistant human pathogens
1308   shows the considerable potential of phylogenetically ancient organisms as drug
1309   sources. Antimicrob Agents Chemother. 2009;53: 5245–5250.
1310   doi:10.1128/AAC.00826-09/ASSET/DDE78552-AE28-472D-8CBE-
1311   1EE103DCACEB/ASSETS/GRAPHIC/ZAC0120985750004.JPEG
1312   33.   Koch, TL. Molecular Evolution of Peptide Signaling in Placozoa and Cnidaria. Ph.
1313   D., University of Copenhagen. 2020.
1314   34.   Hauser F, Koch TL, Grimmelikhuijzen CJP. Review: The evolution of peptidergic
1315   signaling in Cnidaria and Placozoa, including a comparison with Bilateria. Front
1316   Endocrinol (Lausanne). 2022;13: 973862.
1317   doi:10.3389/FENDO.2022.973862/BIBTEX
1318   35.   Kamm K, Schierwater B, DeSalle R. Innate immunity in the simplest animals -
1319   placozoans. BMC Genomics. 2019;20: 5. doi:10.1186/s12864-018-5377-3
1320   36.   Doolittle RF. Structural and Functional Diversity of Fibrinogen-Related Domains.
1321   The Evolution of the Immune System: Conservation and Diversification. 2016;
1322   275–294. doi:10.1016/B978-0-12-801975-7.00011-6
1323   37.   Lang T, Hansson GC, Samuelsson T. Gel-forming mucins appeared early in
1324   metazoan evolution. Proc Natl Acad Sci U S A. 2007;104: 16209–14.
1325   doi:10.1073/pnas.0705984104
1326   38.   Ruthmann A, Behrendt G, Wahl R. The ventral epithelium of Trichoplax adhaerens
1327   (Placozoa): Cytoskeletal structures, cell contacts and endocytosis.
1328   Zoomorphology. 1986;106: 115–122. doi:10.1007/BF00312113
1329   39.   Smith CL, Reese TS. Adherens junctions modulate diffusion between epithelial
1330   cells in Trichoplax adhaerens. Biological Bulletin. 2016;231. doi:10.1086/691069
1331   40.   Leytus SP, Melhado LL, Mangel WF. Rhodamine-based compounds as
1332   fluorogenic substrates for serine proteinases. Biochemical Journal. 1983;209:
1333   299–307. doi:10.1042/BJ2090299
1334   41.   Hartenstein V, Martinez P. Phagocytosis in cellular defense and nutrition: a food-
1335   centered approach to the evolution of macrophages. Cell and Tissue Research
1336   2019 377:3. 2019;377: 527–547. doi:10.1007/S00441-019-03096-6
1337   42.   Hartenstein V, Martinez & P. Phagocytosis in cellular defense and nutrition: a
1338   food-centered approach to the evolution of macrophages. [cited 21 May 2024].
1339   doi:10.1007/s00441-019-03096-6
1340   43.   Parker MW, Feil SC. Pore-forming protein toxins: from structure to function. Prog
1341   Biophys Mol Biol. 2005;88: 91–142. doi:10.1016/J.PBIOMOLBIO.2004.01.009
1342   44.   Bischofberger M, Iacovache I, Gisou Van Der Goot F. Pathogenic Pore-Forming
1343   Proteins: Function and Host Response. Cell Host Microbe. 2012;12: 266–275.
1344   doi:10.1016/J.CHOM.2012.08.005
1345   45.   Iacovache I, Bischofberger M, van der Goot FG. Structure and assembly of pore-
1346   forming proteins. Curr Opin Struct Biol. 2010;20: 241–246.
1347   doi:10.1016/J.SBI.2010.01.013
1348   46.   Liu X, Lieberman J. Knocking 'em Dead: Pore-Forming Proteins in Immune
1349   Defense. Annu Rev Immunol. 2020;38: 455–485. doi:10.1146/ANNUREV-
1350   IMMUNOL-111319-023800/CITE/REFWORKS

47. Cosentino K, Ros U, García-Sáez AJ. Assembling the puzzle: Oligomerization of
α-pore forming proteins in membranes. Biochimica et Biophysica Acta (BBA) -
Biomembranes. 2016;1858: 457–466. doi:10.1016/J.BBAMEM.2015.09.013

48. Jeudy S, Schwartz TU. Crystal Structure of Nucleoporin Nic96 Reveals a Novel,
Intricate Helical Domain Architecture. Journal of Biological Chemistry. 2007;282:
34904–34912. doi:10.1074/JBC.M705479200

49. David L, Borges JP, Hollingsworth LR, Volchuk A, Jansen I, Garlick E, et al. NINJ1
mediates plasma membrane rupture by cutting and releasing membrane disks.
Cell. 2024;187: 2224-2235.e16. doi:10.1016/J.CELL.2024.03.008

50. Blott EJ, Griffiths GM. Secretory lysosomes. Nat Rev Mol Cell Biol. 2002;3: 122–
31. doi:10.1038/nrm732

51. Griffiths GM, Argon Y. Structure and Biogenesis of Lytic Granules. Curr Top
Microbiol Immunol. 1995;198: 39–58. doi:10.1007/978-3-642-79414-8_3

52. Neighbour PA, Huberman HS, Kress Y. Human large granular lymphocytes and
natural killing: Ultrastructural studies of strontium-induced degranulation. Eur J
Immunol. 1982;12: 588–595. doi:10.1002/EJI.1830120711

53. Weaver TE, Na CL, Stahlman M. Biogenesis of lamellar bodies, lysosome-related
organelles involved in storage and secretion of pulmonary surfactant. Semin Cell
Dev Biol. 2002;13: 263–270. doi:10.1016/S1084952102000551

54. Schroeder LK, Kremer S, Kramer MJ, Currie E, Kwan E, Watts JL, et al. Function
of the Caenorhabditis elegans ABC transporter PGP-2 in the biogenesis of a
lysosome-related fat storage organelle. Mol Biol Cell. 2007;18: 995–1008.
doi:10.1091/MBC.E06-08-
0685/ASSET/IMAGES/LARGE/ZMK0030779830009.JPEG

55. Fautin DG, Mariscal RN. Cnidaria: anthozoa. In: Harrison FW, Ruppert EE,
editors. Microscopic Anatomy of Invertebrates Volume 2: Placozoa, Porifera,
Cnidaria, and Ctenophora. Siley-Lis; 1991. pp. 267–358.

56. Steger J, Cole AG, Denner A, Lebedeva T, Genikhovich G, Ries A, et al. Single-
cell transcriptomics identifies conserved regulators of neuroglandular lineages.
Cell Rep. 2022;40. doi:10.1016/J.CELREP.2022.111370

57. Babonis LS, Martindale MQ. Old Cell, New Trick? Cnidocytes as a Model for the
Evolution of Novelty. Integr Comp Biol. 2014;54: 714–722.
doi:10.1093/ICB/ICU027

58. Fautin DG. Structural diversity, systematics, and evolution of cnidae. Toxicon.
2009;54: 1054–1064. doi:10.1016/J.TOXICON.2009.02.024

59. Hernandez-Nicaise M-L. Ctenophora. In: Harrison F, Westfall JA, editors.
Microscopic Anatomy of Invertebrates Volume 2. New York: Wiley-Liss; 1991. pp.
359–418.

60. Babonis LS, DeBiasse MB, Francis WR, Christianson LM, Moss AG, Haddock
SHD, et al. Integrating Embryonic Development and Evolutionary History to
Characterize Tentacle-Specific Cell Types in a Ctenophore. Mol Biol Evol.
2018;35: 2940–2956. doi:10.1093/MOLBEV/MSY171

61. Schendel V, Müller CHG, Kenning M, Maxwell M, Jenner RA, Undheim EAB, et al.
The venom and telopodal defence systems of the centipede Lithobius forficatus
are functionally convergent serial homologues. BMC Biology 2024 22:1. 2024;22:
1–22. doi:10.1186/S12915-024-01925-X

62. von Reumont BM, Undheim EAB, Jauss RT, Jenner RA. Venomics of Remipede
Crustaceans Reveals Novel Peptide Diversity and Illuminates the Venom's
Biological Role. Toxins 2017, Vol 9, Page 234. 2017;9: 234.
doi:10.3390/TOXINS9080234
63. Klimovich A, Bosch TCG. Novel technologies uncover novel 'anti'-microbial
peptides in Hydra shaping the species-specific microbiome. Philos Trans R Soc
Lond B Biol Sci. 2024;379. doi:10.1098/RSTB.2023.0058
64. Franzenburg S, Walter J, Künzel S, Wang J, Baines JF, Bosch TCG, et al. Distinct
antimicrobial peptide expression determines host species-specific bacterial
associations. Proc Natl Acad Sci U S A. 2013;110: E3730–E3738.
doi:10.1073/PNAS.1304960110/SUPPL_FILE/PNAS.201304960SI.PDF
65. Siebert S, Farrell JA, Cazet JF, Abeykoon Y, Primack AS, Schnitzler CE, et al.
Stem cell differentiation trajectories in Hydra resolved at single-cell resolution.
Science (1979). 2019;365.
doi:10.1126/SCIENCE.AAV9314/SUPPL_FILE/AAV9314_SIEBERT_SM.PDF
66. Schierwater B, de Jong D, DeSalle R. Placozoa and the evolution of Metazoa and
intrasomatic cell differentiation. Int J Biochem Cell Biol. 2009;41: 370–379.
doi:10.1016/J.BIOCEL.2008.09.023
67. Bakshani CR, Morales-Garcia AL, Althaus M, Wilcox MD, Pearson JP, Bythell JC,
et al. Evolutionary conservation of the antimicrobial function of mucus: a first
defence against infection. NPJ Biofilms Microbiomes. 2018;4: 14.
doi:10.1038/s41522-018-0057-2
68. Sebé-Pedrós A, Saudemont B, Chomsky E, Plessier F, Mailhé M-P, Renno J, et
al. Cnidarian Cell Type Diversity and Regulation Revealed by Whole-Organism
Single-Cell RNA-Seq. Cell. 2018;173: 1520-1534.e20.
doi:10.1016/J.CELL.2018.05.019
69. Kornder NA, Esser Y, Stoupin D, Leys SP, Mueller B, Vermeij MJA, et al. Sponges
sneeze mucus to shed particulate waste from their seawater inlet pores. Current
Biology. 2022;32: 3855-3861.e3. doi:10.1016/J.CUB.2022.07.017
70. Leys SP, Degnan BM. Cytological basis of photoresponsive behavior in a sponge
larva. Biological Bulletin. 2001;201: 323–338.
doi:10.2307/1543611/ASSET/IMAGES/LARGE/1N0610531011.JPEG
71. Nakanishi N, Stoupin D, Degnan SM, Degnan BM. Sensory Flask Cells in Sponge
Larvae Regulate Metamorphosis via Calcium Signaling. Integr Comp Biol.
2015;55: 1018–1027. doi:10.1093/icb/icv014
72. Ereskovsky A V., Tokina DB, Bézac C, Boury-Esnault N. Metamorphosis of
cinctoblastula larvae (Homoscleromorpha, porifera). J Morphol. 2007;268: 518–
528. doi:10.1002/JMOR.10506
73. Brunet T, King N. The Origin of Animal Multicellularity and Cell Differentiation.
Developmental Cell. Elsevier; 2017. pp. 124–140.
doi:10.1016/j.devcel.2017.09.016
74. Wrackmeyer U, Hansen GH, Seya T, Danielsen EM. Intelectin: A novel lipid raft-
associated protein in the enterocyte brush border. Biochemistry. 2006;45: 9188–
9197. doi:10.1021/BI060570X/ASSET/IMAGES/LARGE/BI060570XF00009.JPEG

75. Hanington PC, Zhang SM. The Primary Role of Fibrinogen-Related Proteins in
Invertebrates Is Defense, Not Coagulation. J Innate Immun. 2010;3: 17–27.
doi:10.1159/000321882
76. Arendt D, Benito-Gutierrez E, Brunet T, Marlow H. Gastric pouches and the
mucociliary sole: setting the stage for nervous system evolution. Phil Trans R Soc
Lond B. 2015;370: 20150286. doi:10.1098/rstb.2015.0286
77. Brunet T, King N. The origin of animal multicellularity and cell differentiation. Dev
Cell. 2017;43: 124–140. doi:10.1016/j.devcel.2017.09.016
78. Cavalier-Smith T. Origin of animal multicellularity: precursors, causes,
consequences-the choanoflagellate/sponge transition, neurogenesis and the
Cambrian explosion. Phil Trans R Soc Lond B. 2017;372: 20150476.
doi:10.1098/rstb.2015.0476
79. Nielsen C. Six major steps in animal evolution: are we derived sponge larvae?
Evol Dev. 2008;10: 241–257. doi:10.1111/j.1525-142x.2008.00231.x
80. Smith CL, Mayorova TD. Insights into the evolution of digestive systems from
studies of Trichoplax adhaerens. Cell Tissue Res. 2019;377: 353–367.
doi:10.1007/s00441-019-03057-z
81. Syed T, Schierwater B. Trichoplax adhaerens: Discovered as a missing link,
forgotten as a hydrozoan, re-discovered as a key to Metozoan evolution. Vie et
Milieu. 2002;52: 177–187.
82. Calcium- and magnesium-free artificial seawater (CMF-ASW). Cold Spring Harb
Protoc. 2009;2009: pdb.rec12053. doi:10.1101/PDB.REC12053
83. Hallgren J, Tsirigos KD, Damgaard Pedersen M, Juan J, Armenteros A, Marcatili
P, et al. DeepTMHMM predicts alpha and beta transmembrane proteins using
deep neural networks. bioRxiv. 2022; 2022.04.08.487609.
doi:10.1101/2022.04.08.487609
84. Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, et al.
InterPro in 2022. Nucleic Acids Res. 2023;51: D418–D427.
doi:10.1093/NAR/GKAC993
85. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al.
AlphaFold Protein Structure Database: massively expanding the structural
coverage of protein-sequence space with high-accuracy models. Nucleic Acids
Res. 2022;50: D439–D444. doi:10.1093/NAR/GKAB1061
86. Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, et al.
UCSF ChimeraX: Meeting modern challenges in visualization and analysis.
Protein Sci. 2018;27: 14–25. doi:10.1002/PRO.3235
87. Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, et al.
UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. Protein Sci. 2021;30: 70–82. doi:10.1002/PRO.3943
88. Grimmelikhuijzen CJP, Leviev I, Carstensen K. Peptides in the Nervous Systems
of Cnidarians: Structure, Function, and Biosynthesis. Int Rev Cytol. 1996;167: 37–
89. doi:10.1016/S0074-7696(08)61345-5
89. Hammer DAT, Ryan PD, Hammer Ø, Harper DAT. Past: Paleontological Statistics
Software Package for Education and Data Analysis. Palaeontologia Electronica.
2001;4: 178. Available: http://palaeo-electronica.orghttp://palaeo-
electronica.org/2001_1/past/issue1_01.htm.

90. Micallef L, Rodgers P. eulerAPE: Drawing Area-Proportional 3-Venn Diagrams
Using Ellipses. PLoS One. 2014;9: 101717. doi:10.1371/journal.pone
1489

Supporting information captions

1491

S1 Fig. Expression of selected genes across *T. adhaerens* metacells (columns) identified by single cell RNA sequencing. Data are from [30]. Cell types are identified based on data from the present study and [6,18,24,30]. Dot color represents fold change of gene expression and dot size represents percentage of the total UMI in the given metacell.

1497

S2 Fig. Color separated *FISH i*mages of the micrographs shown in Fig.2. (A, B) Separated channels corresponding to horizontal (xy) and vertical (xz) projections of color-merged *FISH* images of TH1 wholemounts in Fig 2A and B. (C, D) Separated DIC and fluorescence channels of the merged images of dissociated cells in Fig 2C and D.

1502

S3-6 Fig. Normalized expression of selected genes across metacells for *TH1, TH2*, HH13, and CH23. Cell types are identified based on data from the present study and [6,18,30]. Expression data from: https://sebelab.crg.eu/placozoa_cell_atlas/

1506

S7 Fig. Secondary structure predictions and electrostatic potential maps for putative *Trichoplax* AMPs (AMP1 (A), AMP2 (B), and AMP3 (C) and *Hydra vulgaris* arminin (D). Secondary structures are predicted with Alpha Fold and peptide surfaces are colored by electrostatic potential with ChimeraX.

1511

1512 **S8 Text. Placozoan prepropeptides, predicted cleavage sites and processing**

1513 **motifs**. Underlined regions represent predicted signal peptides. Highlighted yellow

1514 represents the mature peptide region, green represents prohormone convertase

1515 cleavage sites, cyan represents pyroglutamate, and magenta represents C-terminal

1516 amide.

1517

1518 **S9 Fig. Color separated *FISH i*mages of TH1 wholemounts corresponding to**

1519 **horizontal (xy) and vertical (xz) of color-merged *FISH* images in Fig 5F (A) and Fig**

1520 **6F1 (B)**.

1521

S10 Fig. Secretory behaviors of *Trichoplax* feeding on *R. salina* algae. Lipophil cell

granules were labeled with LipidTOX (red). The fluorescent membrane dye FM1-43

(cyan) was added to the seawater to label cell membranes and the contents of LC

granules. BZiPAR, a fluorescent indicator of trypsin activity (green), was added to detect

secretion of trypsin. Algae are visible by autofluorescence and FM1-43 staining (pink in

merged images). At the beginning of the sequence, the peripheral part of the animal

was closely attached to the substrate, while the central part was invaginated, forming a

feeding pocket enclosing algae. At t=0 sec, lipophil granule secretion was evident due to

the appearance of small FM1-43-stained spots (cyan/white; arrowheads) in the feeding

pocket. At t=12 sec, several algae (pink) in the feeding pocket were lysed and released

material stained by FM1-43 (cyan). By 150 – 287 sec, trypsin activity was evident in the

feeding pocket and the lysed algae were decomposing. Bottom panels show intensity-

coded images of BZiPAR trypsin activity indicator at sequential timepoints for the boxed

region.

1536

S11 Movie. *Trichoplax* feeding on *R. salina* algae (same animal as illustrated in Fig

8). Lipophil cell granules were labeled with LipidTOX (red). The fluorescent membrane

dye FM1-43 (cyan) was added to the seawater to label cell membranes and the

contents of LC granules. BZiPAR, a fluorescent indicator of trypsin activity, was added

to detect secretion of trypsin. Algae are visible by autofluorescence (magenta). Time

stamp is at upper left. Lipophil cell granule secretion begins at ~50 sec as evident from

the appearance of small FM1-43-stained puncta in the vicinity of algae. Green

fluorescence generated by proteolysis of BZiPAR becomes visible at ~110 sec and

reaches a peak at ~270 sec. Content of lysed algae is stained by FM1-43, but staining

declines as the algae decompose. The rim of the animal remains closely attached to the

substrate forming a feeding pocket that confines the contents of the lysed algae and the

BZiPAR fluorescence.

1549

1550 **S12 Movie. *Trichoplax* feeding on *R. salina* algae (same animal as illustrated in**

1551 **S10 Fig)**. Staining is the same as in S11 Movie. Lipophil granule secretion begins at

1552 ~14 sec as evident from the appearance of small clouds of FM1-43-stained material in

1553 the vicinity of algae. The clouds coalesce into puncta within seconds. Algae begin to

1554 lyse at ~20 sec. Trypsin activity becomes visible at ~45 sec.

1555

1556

1557

1558

1559

A

B

METAZOA
PARAHOXOZOA
PLANULOZOA

Ctenophora

Porifera

Cnidaria

Bilateria

Placozoa

Trichoplax adhaerens H1

Trichoplax sp. H2

Cladtertia collaboinventa H23

Hoilungia hongkongensis H13

C

Dorsal cilium

Fiber cell

DEC

Crystal cell

Gland cell

VEC

Lipophil cell

Ventral cilium

A

e p

p

p

e e

e

500 nm

B

1 μm

500 nm

C

1 μm

500 nm

D

TaIntelectin
WGA

10 μm

E

TaIntelectin
WGA
DAPI

dec

fc

lc

lc

lc

vec

m

m

10 μm

F

TaIntelectin
TaELPE
WGA

10 μm

I

34   35

J   μm²

*

G

TaIntelectin
TaELPE
DAPI
Tub

H

5 μm

A1
Ta63786
TaELPE
WGA

15
1
15

20 μm

A2
DAPI
Tub

5 μm

A3
μm²
*

B1
Ta26557
TaLF
WGA

B2

B3
DAPI
Tub

B4
1
13
13

B5
μm²
ns

B6
24
2 1
25
19

C1
TaOligoBP
TaRWa
WGA

10 μm

C2

DAPI
Tub

C3
78 105
1

5 μm

D1
Ta54934
TaELP
WGA

D2
DAPI
Tub

D3

D4
12

D5
μm²

E1
Ta60437
TaSIFGa
WGA

E2
DAPI
Tub

E3

E4
μm²
* *
ns

F1
Ta54934
TaELP
WGA
TaFFNPa
TaSIFGa
WGA

F2
38 50

F6
μm²
*

F3
DAPI
Tub

F4

F5

A

DEC

Muc

pVEC

cVEC

LC

TaTetraspanin;
Ta63996;
TaGABAtransporter

TaTrypsin;
TaChimotrypsin;
TaPLA2

● AMP2   ● TaIntelectin   ● AMP1   ● TaELPE   ● TaOligoBP

B

● TaRWa   ● TaFFNPa   ● TaAstacin (Ta26557)   ● TaELP; TaAstacin (Ta54934)   ● TaWPPF   ● TaLF   ● TaSIFGa   ● TaOligoBP