1
2
3
4 **Phosphorylation-induced SUMOylation promotes Ulk4 condensation at ciliary tip**

5 **to transduce Hedgehog signal**

6
7
8
9

10 Mengmeng Zhou[1], Yuhong Han[1], and Jin Jiang[1,2,*]
16 [1]Department of Molecular Biology, [2]Department of Pharmacology, University of Texas

17 Southwestern Medical Center, Dallas, TX 75390, USA

21 *Corresponding authors: jin.jiang@utsouthwestern.edu
22
23

Abstract

Hedgehog (Hh) signaling controls embryonic development and adult tissue homeostasis through the Gli family of transcription factors. In vertebrates, Hh signal transduction depends on the primary cilium where Gli is thought to be activated at the ciliary tip, but the underlying mechanism has remained poorly understood. Our previous study showed that two Unc-51-like kinase (Ulk) family members Ulk4 and Stk36 colocalize with Gli2 at ciliary tip to promote Gli2 phosphorylation upon Shh stimulation, and that ciliary tip localization of Ulk4/Stk36 depends on phosphorylation of Ulk4 by Stk36 (Zhou et al. 2023). Here we show that phosphorylation of Ulk4 drives its phase separation to form biomolecular condensates that recruit Stk36 and Gli2. Mechanistically, Stk36-mediated phosphorylation of Ulk4 promotes its SUMOylation in response to Shh, and subsequent interaction between SUMO and a SUMO-Interacting-Motif (SIM) in the C-terminal region of Ulk4 drives Ulk4 self-assembly to form biomolecular condensates. Phospho-mimetic mutation of Ulk4 sufficed to drive Ulk4/Stk36/Gli2 condensation at ciliary tip, resulting in constitutive Shh pathway activation. Taken together, our results suggest that phosphorylation-dependent SUMOylation of Ulk4 promotes kinase-substrate condensation at ciliary tip to transduce the Hh signal. We speculate that phase separation driven by phosphorylation-dependent SUMOylation may represent a general mechanism for kinase-mediated signaling.

Introduction

The Hh family glycoproteins play essential roles in embryonic development and adult

tissue homeostasis (Jiang and Hui 2008; Briscoe and Therond 2013). Aberrant Hh

signaling contributes to a wide range of human diseases including birth defects and

cancer (Villavicencio et al. 2000; Nieuwenhuis and Hui 2005; Jiang 2022). Hh signaling

in vertebrates critically depends on primary cilia (Goetz and Anderson 2010). In

response to Hh stimulation, Hh pathway components including the GPCR family protein

and Hh signal transducer Smoothened (Smo) and pathway transcriptional factors Gli

proteins accumulate in primary cilia (Corbit et al. 2005; Haycraft et al. 2005; Rohatgi et

al. 2007; Chen et al. 2009; Tukachinsky et al. 2010; Chen et al. 2011). Gli proteins are

thought to be activated in primary cilia because Gli2 variants that fail to accumulate in

the cilia cannot be activated both in cultured cells and in mouse embryos (Santos and

Reiter 2014; Liu et al. 2015; Han et al. 2017); however, the underlying mechanism has

remained poorly understood.

Our previous studies showed that two Ulk family kinases, Ulk3 and Stk36, act in

parallel to convert full-length Gli2 ($Gli2^F$) into its activator form ($Gli2^A$) by directly

phosphorylating Gli2 at multiple sites in mammalian cells (Han et al. 2019; Zhou et al.

2022; Zhou and Jiang 2022). More recently, we found that another Ulk family member

Ulk4, a pseudokinase, acts in conjunction with Stk36 to promote Gli2 phosphorylation

and Hh pathway activation by forming a complex with Stk36 and regulating Stk36 ciliary

accumulation (Zhou et al. 2023). Interestingly, we found that Stk36 and Ulk4 depend on

each other for their ciliary localization and that ciliary tip localization of Ulk4 depends on

Stk36-mediated phosphorylation of Ulk4 on T1021/T1023 (Zhou et al. 2023).

Here we provide evidence that phosphorylation of Ulk4 by Stk36 drives its phase

separation to form biomolecular condensates that recruit Stk36, Gli2 and Sufu. We find

that Stk36-mediated phosphorylation of Ulk4 at T1021/T1023 promotes its SUMOylation

on an adjacent site (K1030) in response to Shh, and that interaction between SUMO

and a SUMO-Interacting-Motif (SIM) in the C-terminal region of Ulk4 drives Ulk4 self-

association to form biomolecular condensates. Remarkably, a phospho-mimetic Ulk4

variant can form condensates that recruit Stk36, Gli2 and Sufu in the absence of Shh,

leading to constitutive Gli2 phosphorylation and pathway activation.

78
79
80
81 **Results**
82
83
84 **Phosphorylation of Ulk4 is sufficient to drive its ciliary tip accumulation**
85
Our previous study revealed that phosphorylation of Ulk4 by Stk36 is necessary for its

ciliary tip accumulation (Zhou et al. 2023). To determine whether Ulk4 phosphorylation

is sufficient to drive its ciliary tip accumulation, we generated a phospho-mimetic form of

human Ulk4 (hUlk4) by converting two previously identified Stk36 sites (T1021 and

T1023) to E (Ulk4-EE). C-terminally HA-tagged wild type Ulk4 (Ulk4-WT) and Ulk4-EE

were expressed in NIH3T3 cells via lentiviral infection, followed by Shh stimulation. As

shown in Figure 1A, Ulk4-WT-HA accumulated at ciliary tip only after Shh stimulation

whereas Ulk4-EE-HA accumulated at ciliary tip regardless of Shh treatment.

Quantification of ciliary tip Ulk4 signals revealed that Shh dramatically increased Ulk4-

WT-HA levels but had little if any effect on Ulk4-EE-HA levels at ciliary tips (Figure 1B),

suggesting that Stk36-mediated phosphorylation of Ulk4 is sufficient to promote its

ciliary tip accumulation.

We next asked whether Ulk4-EE could bring Stk36 to the ciliary tip in the

absence of Shh. NIH3T3 cells were infected with lentivirus expressing Myc-tagged

Stk36 (Myc-Stk36) and Ulk4-WT-HA or Ulk4-EE-HA, followed by immunostaining with

antibodies against the epitope tags to monitor ciliary accumulation of Myc-Stk36 and

Ulk4-WT-HA/Ulk4-EE-HA. Cells co-expressing Myc-Stk36 and Ulk4-WT-HA showed no

ciliary tip accumulation of either Ulk4-WT-HA or Myc-Stk36 (Figure1C and D). By

contrast, cells co-expressing Myc-Stk36 and Ulk4-EE-HA showed ciliary tip

accumulation of both Ulk4-EE-HA and Myc-Stk36 (Figure1C and D), suggesting that

Ulk4-EE could bring Stk36 to the ciliary tip even in the absence of Shh.

107
108
109 **Gli2 ciliary tip accumulation is promoted by Stk36/Ulk4**
110
In response to Shh, Gli proteins and their binding partner and Hh pathway inhibitor Sufu

accumulate at ciliary tip while Smo accumulates along the cilium axoneme (Corbit et al.

2005; Haycraft et al. 2005; Rohatgi et al. 2007; Chen et al. 2009; Tukachinsky et al.

2010; Chen et al. 2011). The mechanisms that control ciliary tip localization of Gli

proteins are poorly understood although Kif7/Gli interaction has been implicated to play

an important role (Haque et al. 2022). The observations that Stk36/Ulk4 colocalized with

Gli2 at ciliary tip and that Ulk family kinases including Ulk3 and Stk36 acted upstream of

Gli2 to promote Shh pathway activation let us to speculate that Gli ciliary tip localization

could be regulated by these kinases. Indeed, as shown in Figure 2A and B, Shh-

induced ciliary accumulation of Gli2 and Sufu was diminished in Stk36 and Ulk3 double

knockout (DKO) NIH3T3 cells (NIH3T3^{DKO}) (Han et al. 2019). Expression of wild type

Stk36 (Stk36-WT) but not its kinase inactive form (Stk36-AA; Zhou et al. 2023) restored

ciliary tip accumulation of both Gli2 and Sufu in response to Shh (Figure 2A and B).

Furthermore, expression of Ulk4-EE but not Ulk4-WT in NIH3T3^{DKO} cells restored ciliary

tip localization of both Gli2 and Sufu, which correlated with ciliary localization of Ulk4-EE

(Figure 2C-F). These results suggest ciliary tip localization of GIli2 and Sufu is promoted

by Stk36 and Ulk4.

128
129
130 **Phosphorylation drives Ulk4 condensation**

131
The ciliary tip accumulation of Hh signaling components resembles biomolecular

condensate formation that depends on multivalent interactions and protein

concentrations (Banani et al. 2017; Shin and Brangwynne 2017). Indeed, Stk36, Ulk4,

Gli and Sufu physically interact with one another (Han et al. 2015; Preuss et al. 2020;

Zhou et al. 2023). Upon Hh stimulation, Smo is accumulated and activated in primary

cilia, which may result in local activation of Stk36 kinase and Stk36-mediated

phosphorylation of Ulk4, leading to their condensation that also recruits Gli2/Sufu. To

test this hypothesis, we asked whether Stk36-mediated phosphorylation of Ulk4 can

drive Stk36/Ulk4 condensation in the cytoplasm when both proteins were

overexpressed in HEK293T cells. Constitutively active Stk36 (Stk36-EE) or its kinase

inactive form (Stk36-AA) was transfected into HEK293T cells together with Ulk4-WT. As

shown in Figure 3A, coexpression of Stk36-EE with Ulk4-WT led to the formation of

Ulk4 puncta that also contained Stk36. However, coexpression of Stk36-AA with Ulk4-

WT failed to induce the formation of Ulk4/Stk36 puncta. The formation of Ulk4/Stk36

puncta depends on phosphorylation of Ulk4 by Stk36 because HEK293T cells co-

expressing a phospho-deficient form of Ulk4 (Ulk4-2TA) and Stk36-EE failed to form

Ulk4/Stk36 puncta (Figure 3B). Strikingly, HEK293T cells expressing Ulk4-EE formed

Ulk4 condensates that also recruited endogenous Gli2 and Sufu into the condensates

(Figure 3C). These results suggest that phosphorylation of Ulk4 by Stk36 promotes the

formation of Ulk4 condensates that can recruit its interaction proteins including Stk36,

Gli2 and Sufu.

Most membraneless biomolecular condensates form through Liquid-Liquid phase

separation (LLPS) that can be disrupted by 1, 6-hexanediol (Sabari et al. 2018;

McKnight 2024). Indeed, the condensates of GFP-tagged YAP (GFP-YAP), which is

formed via LLPS (Cai et al. 2019), were disrupted by treating cells with 5% 1, 6-

hexanediol for 5 min (Figure 3D). To determine the biomaterial property of the

condensates organized by Ulk4, we treated HEK293T cells that contained CFP-

Stk36/Ulk4-EE condensates with 5% 1, 6-hexanediol for 5 to 30 minutes. We found that

CFP-Stk36/Ulk4-EE condensates were resistant to 1, 6-hexanediol even after treatment

for 30 min (Figure 3D; Figure 3- figure supplement 1), suggesting that Stk36/Ulk4

condensates are more rigid than LLPS and may resemble gel-like or solid-state

condensates (Woodruff et al. 2017; Putnam et al. 2019).

164
165
166   **Shh induces SUMOylation of Ulk4 depending on Stk36-mediated phosphorylation**
167
We next asked how Ulk4 phosphorylation promotes its condensation. Phosphorylation

can regulate other post-translational modifications (PTMs) such as SUMOylation, and

SUMOylation has been implicated in the formation of biomolecular condensates (Shen

et al. 2006; Yang and Gregoire 2006; Keiten-Schmitz et al. 2021). Interestingly, like

Ulk4-mediated condensation, condensates formed by a synthetic construct containing

multiple SUMO and SIM domains, RFP-SUMO6-SIM10 (Banani et al. 2016), were also

resistant to 1, 6-hexanediol treatment (Figure 3D; Figure 3- figure supplement 1).

Therefore, we asked whether phosphorylation induces Ulk4 condensation through

promoting its SUMOylation. Coexpression of Ulk4-HA with Fg-Stk36-EE but not with Fg-

Stk36-AA in HEK293T cells resulted in a mobility shift of Ulk4-HA indicative of SUMO

conjugation (Figure 4A; top panel). Probing with anti-SUMO1 antibody confirmed that

higher molecular weight bands contained SUMOylated Ulk4 (Figure 4A; bottom panel).

Mutating the Stk36 sites on Ulk4 (Ulk4-2TA) abolished Ulk4 SUMOylation stimulated by

Stk36-EE (Figure 4B). On the other hand, Ulk4-EE was SUMOylated without

coexpression of Stk36-EE, and SUMOylation of Ulk4-EE was abolished by a

SUMOylation inhibitor TAK981 (Fig. 4C)(Langston et al. 2021). These results

demonstrated that Stk36-mediated phosphorylation at T1021/T1023 can induce Ulk4

SUMOylation. Of note, SUMOylated Ulk4 tends to form high molecular weight species

on the SDS-PAGE, a phenomenon also observed for SUMOylated nuclear hormone

receptors (Poukka et al. 2000; Tian et al. 2002).

To determine whether Shh induces SUMOylation of Ulk4 depending on its

phosphorylation by Stk36, Ulk4-WT or Ulk4-2TA was expressed in NIH3T3 cells via

lentiviral infection, followed by Shh stimulation. As shown in Figure 4D, Shh induced

SUMOylation of Ulk4-WT but not Ulk4-2TA.

SUMOylation sites have consensus motifs: ψ$\underline{K}$xE/D or E/DxKψ, where ψ is a

large hydrophobic amino acid and x could be any amino acid (Matic et al. 2010).

SUMOylation of many proteins was promoted by phosphorylation on sites adjacent to

the SUMOylation sites (Yang and Gregoire 2006). Inspection of Ulk4 C-terminal region

identified a SUMOylation consensus site (ESK$_{1030}$L) adjacent to the Stk36

phosphorylation sites (Figure 4E). To determine whether phosphorylation of Ulk4 at

T1021/T1023 could promote SUMOylation at K1030, we carried out *in vitro*

SUMOylation experiments. GST fusion proteins containing a Ulk4 fragment that

encompasses both the Stk36 sites and the putative SUMOylation site (GST-Ulk4C;

Figure 4E) were incubated with immunopurified Stk36-EE and recombinant CK1 for *in*

vitro phosphorylation, followed by incubation with SUMO1 and SUMO enzymes (E1 and

E2) for *in vitro* SUMOylation. As shown in Figure 4F, GST-Ulk4C was SUMOylated after

kinase phosphorylation. Of note, CK1 was included to boost Stk36 kinase activity in the

in vitro phosphorylation assay. Mutation of either K1030 to R (KR) or T1021/T1023 to

A/A (2TA) abolished *in vitro* SUMOylation of GST-Ulk4C (Figure 4F). Furthermore, the

K1030R mutation abolished Stk36-induced SUMOylation of Ulk4 and the constitutive

SUMOylation of Ulk4-EE (Ulk4-EE-KR) in HEK293T cells, as well as Shh-induced Ulk4

SUMOylation in NIH3T3 cells (Figure 4G-I). Taken together, these results demonstrate

that Shh induces SUMOylation of Ulk4 on K1030 depending on its phosphorylation at

T1021/T1023.

SUMOylation promotes Ulk4 self-association via SUMO/SIM interaction

SUMOylation mediates protein-protein interaction by binding to SIM, which consists of a

stretch of hydrophobic amino acids adjacent to acidic residues (Song et al. 2004).

Interestingly, Ulk4 contains a SIM motif ($_{1145}$EDLLLL$_{1151}$DLE) located C-terminally to

phosphorylation-mediated SUMOylation site (Figure. 4E). To determine whether

SUMOylation of Ulk4 promotes its self-association through interacting with SIM, we

examined the interaction between Flag-tagged Ulk4-EE (Ulk4-EE-Fg), which is

constitutively SUMOylated, or Ulk4-EE-KR-Fg, which is no longer SUMOylated (Figure

4H), with HA-tagged wild type (Ulk4-WT-HA) or SIM-deficient Ulk4 (Ulk4-ΔSIM-HA;

Figure. 4E). These constructs were transfected into HEK293T cells, followed by co-

immunoprecipitation (Co-IP) and western blot analysis. As shown in Figure 4J, Ulk4-EE-

Fg formed a complex with Ulk4-WT-HA, which was diminished when SIM was mutated

in Ulk4-WT-HA (Ulk4-ΔSIM-HA) or the SUMOylation site was mutated in Ulk4-EE-Fg

(Ulk4-EE-KR-Fg). We also generated a C-terminal fragment of Ulk4-EE from aa1004-

aa1275 of (C272-EE-Fg). C272-EE-Fg formed a complex with Ulk4-WT-HA albeit at

lower affinity compared with Ulk4-EE-Fg; however, C272-EE-Fg failed to form a

complex with Ulk4-ΔSIM-HA (Figure 4K). These observations suggest that 1) SUMO-

SIM interaction facilitates Ulk4 self-association through its C-terminal region and 2) the

N-terminal region of Ulk4 can also mediate Ulk4 self-association independent of SUMO-

SIM interaction.

SUMOylation and SIM are essential for Ulk4 condensation and Ulk4-mediated Shh

pathway activation

To determine whether SUMOylation and SIM mediates Ulk4 condensation as is the

case for PLM bodies (Shen et al. 2006), we coexpressed Stk36-EE with wild type (Ulk4-

WT), SUMOylation deficient (Ulk4-KR), or SIM mutated Ulk4 (Ulk4-ΔSIM) in HEK293T

cells. As shown in Figure 5A, while Ulk4-WT formed condensates that contained Stk36-

EE, neither Ulk4-KR nor Ulk4-ΔSIM formed condensates but rather exhibited diffused

distribution in HEK293T cells.

Next, we expressed human Ulk4-WT, Ulk4-KR, or Ulk4-ΔSIM via lentiviral

infection in NIH3T3 cells with endogenous Ulk4 depleted by shRNA and treated the

cells with or without Shh. In contrast to Ulk4-WT, which was accumulated at the ciliary

tip after Shh treatment, neither Ulk4-KR nor Ulk4-ΔSIM accumulated at ciliary tip in

response to Shh stimulation (Figure 5B and C). While Ulk4-WT rescued Gli2

phosphorylation, *Ptch1* and *Gli1* expression induced by Shh in Ulk4 depleted cells,

neither Ulk4-KR nor Ulk4-ΔSIM rescued Gli2 phosphorylation and Shh target gene

expression (Figure 5D-F). These results suggest that Shh-induced Ulk4 SUMOylation

and subsequent SUMO-SIM interaction is essential for Ulk4 condensation at ciliary tip

and its role in mediating Shh pathway activation.

Ulk4-EE exhibits constitutive pathway activity depending on its SUMOylation

Ulk4-EE formed condensates in the cytoplasm when expressed in HEK293T cells and

accumulated at the ciliary tips in NIH3T3 cells in the absence of Shh. Treating cells with

the SUMOylation inhibitor TAK-981 abolished Ulk4 condensation and ciliary tip

accumulation (Figure 6A-C), suggesting that Ulk4-EE condensation and constitutive

ciliary tip localization may depend on its SUMOylation. To determine whether Ulk4-EE

condensation and constitutive ciliary tip localization depend on its SUMOylation at

K1030, we examined Ulk4-EE-KR-Fg in which K1030 was mutated to R. In contrast to

Ulk4-EE-Fg, Ulk4-EE-KR-Fg neither formed condensates in HEK293T cells nor

accumulated at ciliary tip when expressed in NIH3T3 cells (Figure 6D-F).

Consistent with its constitutive ciliary tip accumulation and its ability to recruit

Stk36 and Gli2/Sufu to the ciliary tip (Figure 1A, C and Figure 2C-F), Ulk4-EE also

exhibited constitutive pathway activity as indicated by Gli2 phosphorylation and

transcriptional activation of *Ptch1* and *Gli1* in the absence of Shh (Figure 6G-L).

Blocking the SUMOylation of Ulk4-EE with either TAK-981 treatment or the K1030R

mutation inhibited its constitutive pathway activity (Figure 6G-L). Knockdown of Stk36

attenuated Shh-independent activation of *Ptch1* and *Gli1* induced by Ulk4-EE (Figure 7-

figure supplement 1A and B), suggesting that the constitutive pathway activity of Ulk4-

EE depends on Stk36. Indeed, Ulk4-EE could activate Stk36 as indicated by the

phosphorylation of the activation loop in the Stk36 kinase domain (p158T/pS1590;

Figure 7-figure supplement 1C) even though Stk36 is normally activated by Shh

independent of Ulk4 (Zhou et al. 2023). Taken together, these results suggest that

phosphorylation-mediated SUMOylation and subsequent SUMO-SIM interaction drives

Ulk4 condensation that recruits Stk36 and Gli2 at ciliary tip, which is essential for Ulk4's

role in mediating Shh signal transduction.

278
279
280 **Discussion**
281
282
How Hh converts the latent Gli transcription factor into its activator form at ciliary tip has

remained an enigma in the field. Our previous study revealed that Stk36/Ulk4 acts in

parallel with Ulk3 to promote Gli2 phosphorylation and activation in a manner depending

on ciliary tip localization of Stk36/Ulk4, and that Shh-induced ciliary tip accumulation of

this kinase/pseudokinase pair depends on phosphorylation of Ulk4 by Stk36 (Zhou et al.

2023). Here we demonstrate that phosphorylation of Ulk4 induces its SUMOylation and

that subsequent SUMO-SIM mediated protein-protein interaction promotes ciliary tip

condensation of Ulk4 that also recruits Stk36 and Gli2-Sufu, leading to Gli2

phosphorylation and activation (Figure 7). We further showed that Shh-induced

SUMOylation of Ulk4 depended on Ulk4 phosphorylation at T1021/T1023 and that a

phospho-mimetic Ulk4 variant (Ulk4-EE) formed condensates that recruited Stk36 and

Gli2 in the absence of Shh, leading to constitutive pathway activation.

Biomolecular condensates have been implicated in many cellular processes in

which they function to facilitate biochemical reactions by simultaneously concentrating

substrates and enzymes in the same subcellular compartments (Banani et al. 2017;

Shin and Brangwynne 2017; Su et al. 2021). The formation of biomolecular

condensates through phase separation is often constitutive or regulated in response to

stress signals. Whether phase separation can be induced by extracellular signals such

as morphogens has been largely underexplored. Here we showed that Shh induces the

formation of Ulk4 condensates that recruit Stk36 and Gli2 at ciliary tip to facilitate Gli2

phosphorylation and activation, providing a mechanistic insight into how Shh activates

Gli proteins at ciliary tip. The formation of Stk36-Ulk4-Gli2 condensates at ciliary tip in

response to Shh stimulation is likely due to local activation of Stk36 kinase by activated

Smo accumulated at primary cilia and relatively high concentration of Stk36, Ulk4, Gli2

and Sufu at ciliary tips. Indeed, overexpression of Ulk4-EE or Stk36-EE/Ulk4 in

HEK293T cells led to the formation of Stk36/Ulk4 condensates in the cytoplasm that

could also recruit endogenous Gli2 and Sufu.

In most cases, the formation of membraneless biomolecular condensates is

driven by liquid-liquid phase separation (LLPS) mediated through intrinsically disordered

regions (IDRs)/low-complexity domains (LCDs) (Banani et al. 2017; Shin and

Brangwynne 2017). Here we showed that phosphorylation-regulated SUMOylation of

Ulk4 and subsequent SUMO-SIM interaction is responsible for Shh-induced

Stk36/Ulk4/Gli2 condensation at ciliary tip, which provides a mechanism that links

kinase-mediated signaling and biomolecular condensation. Because phosphorylation-

induced SUMOylation motifs are present in many regulatory proteins (Yang and

Gregoire 2006), and SUMOylation has been implicated as a mechanism for condensate

formation (Keiten-Schmitz et al. 2021), we speculate that phase separation driven by

phosphorylation-dependent SUMOylation could be generally utilized in other settings

and may represent a more general mechanism for kinase-mediated signaling.

Unlike liquid-like condensates whose formation is mediated by IDRs/LCDs and

can be disrupted by 1,6 Hexanediol (Sabari et al. 2018; McKnight 2024),

Ulk4/Stk36/Gli2 condensates are resistant to 1,6 Hexanediol treatment (Figure 3D;

Figure 3-figure supplement 1), thus may resemble gel-like or solid-state condensates

(Woodruff et al. 2017; Putnam et al. 2019). Interestingly, SUMO/SIM-mediated

condensation of a synthetic construct (RFP-SUMO6-SIM10) also formed gel-like/solid-

state condensates because they were resistant to 1,6 Hexanediol treatment (Figure 3D;

Figure 3-figure supplement 1). It would be interesting to determine whether other

condensates mediated by SUMO-SIM interaction such as PLM also exhibit gel-

like/solid-state property.

Although Shh-induced SUMOylation of Ulk4 and SUMO-SIM-mediated

interaction are essential for the formation of Ulk4/Stk36/Gli2/Sufu condensates, other

protein-protein interaction events could also be important for the ciliary tip accumulation

of these intracellular Hh signaling components. Indeed, a recent study revealed that

Kif7-Gli2 interaction is essential for their ciliary tip accumulation (Haque et al. 2022). In

the future, it would be interesting to explore the functional relationship between Kif7 and

the Ulk family kinases in the regulation of ciliary accumulation of Gli proteins.

339
340
341
342

Materials and methods

DNA constructs

Myc-Gli2 was described previously (Han et al. 2019). Myc-Stk36, Fg-Stk36-WT, Fg-Stk36-EE, Fg-Stk36-AA, Ulk4-WT-HA, Ulk4-AA-HA, GST-Ulk4C-WT, GST-Ulk4C-2TA were generated previously (Zhou et al. 2023). C-terminally 3XHA-tagged human Ulk4-EE, Ulk4-KR, Ulk4-ΔSIM and C-terminally 3XFg-tagged Ulk4-EE, Ulk4-EE-KR, Ulk4-C272-EE were subcloned into the *FUGW* vector between EcoRI and BamHI sites. All GST fusion proteins were subcloned into the *pGEX 4T-1* vector between EcoRI and XhoI sites. Ulk4C-KR, Ulk4-C272-EE, Ulk4-KR, Ulk4-ΔSIM, Ulk4-EE, Ulk4-EE-KR plasmids were constructed by PCR-mediated site-directed mutagenesis.

Cell culture, transfection, and Shh treatment

NIH3T3 cells (ATCC, CRL-1658) were cultured in Dulbecco's modified Eagle's medium (DMEM, Sigma) with10% bovine calf serum (BCS, Gibco) and 1% penicillin-streptomycin (Sigma). GenJet Plus *in vitro* DNA transfection kit was used for transfection experiments according to the manufacturer's instruction (SignaGen). HEK293T cells (ATCC, CRL-11268) were cultured in DMEM containing 10% fetal bovine serum (FBS, Gibco) and 1% penicillin-streptomycin. Transfection was performed using the PolyJet *in vitro* DNA transfection kit (SignaGen) according to the instruction. For Shh treatment, the NIH3T3 cells were cultured at 50%-confluent densities in plate or chamber slides for one day and starved in serum-free medium (0.5% BCS) for 12 hours to allow ciliation. The recombinant human Shh N-terminal fragment (1 ng/ml; R&D

Systems, #8908-SH-005) was then added to the same serum-free medium for 12 hours

or overnight.

Immunostaining, immunoprecipitation, and western blotting analysis

Immunostaining and quantification were carried out as previously described (Han et al.

2019; Zhou et al. 2023). Images were captured using Zeiss LSM700 laser scanning

microscope. Fluorescence signal intensities were measured in the respective channel

by ImageJ. 20 cells were randomly picked for each experimental group to measure

ciliary tip fluorescence signal intensities. For Co-IP experiments were carried out as

previously described (Zhou et al. 2023). Western blot was carried out using standard

protocol as previously described (Zhou et al. 2023). Antibodies used in this study were

as follows: Mouse anti-Myc (Santa Cruz Biotechnology; 9E10), Mouse anti-Flag (Sigma;

M2), Mouse anti-HA (Santa Cruz Biotechnology; F7), Mouse anti-acetylated Tubulin

(Sigma; T7451; staining:1:1000), Rabbit anti-Myc (Abcam; ab9106; staining 1:2000),

Rabbit anti-HA (AbCam; ab9110; staining 1:500), Rabbit anti-SUMO1 (4930, Cell

Signaling), Rabbit anti-Sufu (Abcam;ab28083; staining 1:500). Goat anti-HA (NOVUS;

NB600-362; 1:400), Goat anti-Gli2 (R&D system; AF3635; staining 1:1000), Rabbit anti-

pGli2 (pS230/232) (Han et al. 2019), and Rabbit anti-pStk36 (pT158/pS159) (Shi et al.

2011; Zhou et al. 2023)

SUMOylation Assay

Cell-based SUMOylation assay was carried out as previously described (Ma et al. 2016;

Han and Jiang 2022). Briefly, cell pellets were lysed at 4 °C for 10 min with lysis buffer

containing 20 mM Tris HCl pH 8.0, 150 mM NaCl, 5 mM EDTA, 1% IGPAL CA-630, 10%

glycerol, 1 mM DTT, 20 mM NEM, 1X protease inhibitors cocktail, 1X phosSTOP

phosphatase inhibitors. SDS was added into the lysates to a final concentration of 1%

and the mixture was boiled 5 min. After 5-fold dilution with cool lysis buffer, the mixture

was subjected to immunoprecipitation and western blot analysis.

In vitro SUMOylation experiments were performed using the SUMOylation Assay

Kit (Abcam; ab139470) following manufacturer's instruction. Flag-tagged Stk36-EE was

transiently expressed in HEK293T cells, purified with anti-Fg agarose (Sigma; A2220)

and eluted with Flag peptide (Sigma; F4799). The GST and GST fusion proteins were

purified with glutathione beads (GE; 17-0756-01) and the resulting agarose were

subjective to phosphorylation by Stk36-EE/CK1 as previously described (Zhou et al.

2022). The products of *in vitro* kinase assays were used as the substrates for *in vitro*

SUMOylation assays.

1,6 hexanediol treatment

Cultured HEK293T cells were transfected with GFP-Stk36 and Ulk4-EE-HA, GFP-YAP

or RFP-SUMO6SIM10 DNA constructs. After 48 hours, cells were treated with 5% 1,6

hexanediol for 5-30 minutes. Images were taken before and after treatment.

Lentivirus production and infection

To package the lentivirus, Ulk4 shRNA vector or *FUGW* lentiviral constructs were co-

transfected with packaging plasmids (PSPAX2 and PMDG2) into HEK293T cells using

PolyJet. After 48 hours to 72 hours of incubation, the supernatant was transferred to a

15 mL centrifuge tube and centrifuged at 2500 RPM for 15 minutes. After filtered

through a 0.45 mM filter, the supernatant was collected into a new sterile tube

(Beckman; 344059) and centrifuged at 20,000 g for 2 hours. The sediment was re-

suspended, aliquoted in small volume of 10% FBS culture medium and stored at -80^{o}C

for future use. For cell infection, lentiviruses were added to 70% confluent wild type

NIH3T3 cells together with Polybrene (Sigma, Cat. No. H9268) for overnight to get

stable NIH3T3 cell lines.

RNA interference

The mouse Ulk4 shRNA lentiviral vector was purchased from Sigma (Sigma;

TRCN0000328268). The Non target shRNA Plasmid DNA (Sigma; SHC016) was used

as a control. The target sequence of mouse Ulk4 shRNA is:

CTGCGAAGATTATCGAGAATG. The sequences for Stk36 siRNA were: 5'-

GGUAUACUGGCUUCAGAAA-3' (Cat# SASI_ Mm02 _00345637) and 5'-

GCCUUAUGUGCUUUGCUGU-3' (Cat# SASI_Mm01_00167751). The sequences for

negative control were: 5'-UUCUCCGAACGUGUCACG U-3'. The knockdown efficiency

was validated via real-time PCR.

Quantitative RT-qPCR

Total RNA was extracted from 1×10^{6} cells using the RNeasy Plus Mini Kit (Qiagen;

74134), and cDNA was synthesized with High-Capacity cDNA Reverse Transcription Kit

(Applied Biosystems; 01279158) and qPCR was performed using Fast SYBR Green

Master Mix (Applied Biosystems; 2608088) and a Bio-Rad CFX96 real-time PCR

system (BIO-RAD; HSP9601). The RT-qPCR was performed in triplicate for each of

three independent biological replicates. Quantification of mRNA levels was calculated

using the comparative CT method. Primers: Gli1, 5'-GTGCACGTTTGAAGGCTGTC-3'

and 5'-GAGTGGGTCCGATTCTGGTG-3'; Ptch1, 5'-GAAGCCACAGAAAACCCTGTC-3'

and 5'-GCCGCAAGCCTTCTCTACG-3'; Ulk4, 5'- ATGCAGAGTGTGATTGCGTTG-3'

and 5'- GAGTGGCAGTTTCTGTGAACA-3'; Stk36, 5'-

CGCATCCTACACCGAGATATGA-3' and 5'- AAATCCAAAGTCACAGAGCTTGA-3';

GAPDH, 5'-GTGGTGAAGCAGGCATCTGA-3' and 5'-GCCATGTAGGCCATGAGGTC-

3'.

Acknowledgements

We thank Bing Wang and Yong Suk Cho for helps during performing the experiments.

J. Jiang is a Eugene McDermott Endowed Scholar in Biomedical Science at the

University of Texas Southwestern.

449
450
451   **Declaration of interests**

452
453   The authors declare no competing interests.

454
455

Reference

Banani SF, Lee HO, Hyman AA, Rosen MK. 2017. Biomolecular condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Biol* **18**: 285-298.

Banani SF, Rice AM, Peeples WB, Lin Y, Jain S, Parker R, Rosen MK. 2016. Compositional Control of Phase-Separated Cellular Bodies. *Cell* **166**: 651-663.

Briscoe J, Therond PP. 2013. The mechanisms of Hedgehog signalling and its roles in development and disease. *Nat Rev Mol Cell Biol* **14**: 418-431.

Cai D, Feliciano D, Dong P, Flores E, Gruebele M, Porat-Shliom N, Sukenik S, Liu Z, Lippincott-Schwartz J. 2019. Phase separation of YAP reorganizes genome topology for long-term YAP target gene expression. *Nat Cell Biol* **21**: 1578-1589.

Chen MH, Wilson CW, Li YJ, Law KK, Lu CS, Gacayan R, Zhang X, Hui CC, Chuang PT. 2009. Cilium-independent regulation of Gli protein function by Sufu in Hedgehog signaling is evolutionarily conserved. *Genes Dev* **23**: 1910-1928.

Chen Y, Sasai N, Ma G, Yue T, Jia J, Briscoe J, Jiang J. 2011. Sonic Hedgehog dependent phosphorylation by CK1alpha and GRK2 is required for ciliary accumulation and activation of smoothened. *PLoS biology* **9**: e1001083.

Corbit KC, Aanstad P, Singla V, Norman AR, Stainier DY, Reiter JF. 2005. Vertebrate Smoothened functions at the primary cilium. *Nature* **437**: 1018-1021.

Goetz SC, Anderson KV. 2010. The primary cilium: a signalling centre during vertebrate development. *Nat Rev Genet* **11**: 331-344.

Han Y, Jiang J. 2022. Cell-Based Assays for Smoothened Ubiquitination and Sumoylation. *Methods Mol Biol* **2374**: 139-147.

Han Y, Shi Q, Jiang J. 2015. Multisite interaction with Sufu regulates Ci/Gli activity

through distinct mechanisms in Hh signal transduction. *Proc Natl Acad Sci U S A*

112: 6383-6388.

Han Y, Wang B, Cho YS, Zhu J, Wu J, Chen Y, Jiang J. 2019. Phosphorylation of Ci/Gli

by Fused Family Kinases Promotes Hedgehog Signaling. *Dev Cell* **50**: 610-626

e614.

Han Y, Xiong Y, Shi X, Wu J, Zhao Y, Jiang J. 2017. Regulation of Gli ciliary localization

and Hedgehog signaling by the PY-NLS/karyopherin-beta2 nuclear import

system. *PLoS biology* **15**: e2002063.

Haque F, Freniere C, Ye Q, Mani N, Wilson-Kubalek EM, Ku PI, Milligan RA,

Subramanian R. 2022. Cytoskeletal regulation of a transcription factor by DNA

mimicry via coiled-coil interactions. *Nat Cell Biol* **24**: 1088-1098.

Haycraft CJ, Banizs B, Aydin-Son Y, Zhang Q, Michaud EJ, Yoder BK. 2005. Gli2 and

gli3 localize to cilia and require the intraflagellar transport protein polaris for

processing and function. *PLoS Genet* **1**: e53.

Jiang J. 2022. Hedgehog signaling mechanism and role in cancer. *Semin Cancer Biol*

85: 107-122.

Jiang J, Hui CC. 2008. Hedgehog signaling in development and cancer. *Dev Cell* **15**:

801-812.

Keiten-Schmitz J, Roder L, Hornstein E, Muller-McNicoll M, Muller S. 2021. SUMO:

Glue or Solvent for Phase-Separated Ribonucleoprotein Complexes and

Molecular Condensates? *Front Mol Biosci* **8**: 673038.

Langston SP, Grossman S, England D, Afroze R, Bence N, Bowman D, Bump N, Chau

R, Chuang BC, Claiborne C et al. 2021. Discovery of TAK-981, a First-in-Class

Inhibitor of SUMO-Activating Enzyme for the Treatment of Cancer. *J Med Chem*

64: 2501-2520.

Liu J, Zeng H, Liu A. 2015. The loss of Hh responsiveness by a non-ciliary Gli2 variant.

Development **142**: 1651-1660.

Ma G, Li S, Han Y, Li S, Yue T, Wang B, Jiang J. 2016. Regulation of Smoothened

Trafficking and Hedgehog Signaling by the SUMO Pathway. *Dev Cell* **39**: 438-

451.

Matic I, Schimmel J, Hendriks IA, van Santen MA, van de Rijke F, van Dam H, Gnad F,

Mann M, Vertegaal AC. 2010. Site-specific identification of SUMO-2 targets in

cells reveals an inverted SUMOylation motif and a hydrophobic cluster

SUMOylation motif. *Mol Cell* **39**: 641-652.

McKnight SL. 2024. Protein domains of low sequence complexity-dark matter of the

proteome. *Genes Dev* **38**: 205-212.

Nieuwenhuis E, Hui CC. 2005. Hedgehog signaling and congenital malformations. *Clin

Genet* **67**: 193-208.

Poukka H, Karvonen U, Janne OA, Palvimo JJ. 2000. Covalent modification of the

androgen receptor by small ubiquitin-like modifier 1 (SUMO-1). *Proc Natl Acad

Sci U S A* **97**: 14145-14150.

Preuss F, Chatterjee D, Mathea S, Shrestha S, St-Germain J, Saha M, Kannan N,

Raught B, Rottapel R, Knapp S. 2020. Nucleotide Binding, Evolutionary Insights,

and Interaction Partners of the Pseudokinase Unc-51-like Kinase 4. *Structure* **28**: 1184-1196 e1186.

Putnam A, Cassani M, Smith J, Seydoux G. 2019. A gel phase promotes condensation of liquid P granules in Caenorhabditis elegans embryos. *Nat Struct Mol Biol* **26**: 220-226.

Rohatgi R, Milenkovic L, Scott MP. 2007. Patched1 regulates hedgehog signaling at the primary cilium. *Science* **317**: 372-376.

Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, Abraham BJ, Hannett NM, Zamudio AV, Manteiga JC et al. 2018. Coactivator condensation at super-enhancers links phase separation and gene control. *Science* **361**.

Santos N, Reiter JF. 2014. A central region of Gli2 regulates its localization to the primary cilium and transcriptional activity. *J Cell Sci* **127**: 1500-1510.

Shen TH, Lin HK, Scaglioni PP, Yung TM, Pandolfi PP. 2006. The mechanisms of PML-nuclear body formation. *Mol Cell* **24**: 331-339.

Shi Q, Li S, Jia J, Jiang J. 2011. The Hedgehog-induced Smoothened conformational switch assembles a signaling complex that activates Fused by promoting its dimerization and phosphorylation. *Development* **138**: 4219-4231.

Shin Y, Brangwynne CP. 2017. Liquid phase condensation in cell physiology and disease. *Science* **357**.

Song J, Durrin LK, Wilkinson TA, Krontiris TG, Chen Y. 2004. Identification of a SUMO-binding motif that recognizes SUMO-modified proteins. *Proceedings of the National Academy of Sciences of the United States of America* **101**: 14373-14378.

Su Q, Mehta S, Zhang J. 2021. Liquid-liquid phase separation: Orchestrating cell

signaling through time and space. *Mol Cell* **81**: 4137-4146.

Tian S, Poukka H, Palvimo JJ, Janne OA. 2002. Small ubiquitin-related modifier-1

(SUMO-1) modification of the glucocorticoid receptor. *Biochem J* **367**: 907-911.

Tukachinsky H, Lopez LV, Salic A. 2010. A mechanism for vertebrate Hedgehog

signaling: recruitment to cilia and dissociation of SuFu-Gli protein complexes.

Journal Of Cell Biology **191**: 415-428.

Villavicencio EH, Walterhouse DO, Iannaccone PM. 2000. The sonic hedgehog-

patched-gli pathway in human development and disease. *Am J Hum Genet* **67**:

1047-1054.

Woodruff JB, Ferreira Gomes B, Widlund PO, Mahamid J, Honigmann A, Hyman AA.

2017. The Centrosome Is a Selective Condensate that Nucleates Microtubules

by Concentrating Tubulin. *Cell* **169**: 1066-1077 e1010.

Yang XJ, Gregoire S. 2006. A recurrent phospho-sumoyl switch in transcriptional

repression and beyond. *Mol Cell* **23**: 779-786.

Zhou M, Han Y, Jiang J. 2023. Ulk4 promotes Shh signaling by regulating Stk36 ciliary

localization and Gli2 phosphorylation. *eLife* **12**.

Zhou M, Han Y, Wang B, Cho YS, Jiang J. 2022. Dose-dependent phosphorylation and

activation of Hh pathway transcription factors. *Life Sci Alliance* **5**.

Zhou M, Jiang J. 2022. Gli Phosphorylation Code in Hedgehog Signal Transduction.

Front Cell Dev Biol **10**: 846927.

568
569
570 **Figure 1. Stk36-mediated phosphorylation of Ulk4 suffices to drive its ciliary tip**

571 **accumulation**

572
(A-B) Representative images of immunostaining (A) and quantification (B) of ciliary tip

localized Ulk4-HA (green in A) in NIH3T3 cells infected with Ulk4 shRNA and lentivirus

expressing human Ulk4-WT-HA or Ulk4-EE-HA in the presence or absence of Shh-N.

(C-D) Representative images of immunostaining (C) and quantification (D) of ciliary tip

localized Ulk4-HA (green in C) or Myc-Stk36 (blue in C) in NIH3T3 cells infected with

Ulk4-WT-HA or Ulk4-EE-HA lentivirus. Primary cilia were marked by acetylated tubulin

(Ac-tub) staining (red in A and C). The intensity of ciliary-localized Ulk4-HA or Myc-

Stk36 was measured by Image J. 20 cells were randomly selected from each

experimental group for quantification. Data are mean ± SD from three independent

experiments. ***$p < 0.001$ (Student's t test). Scale bars are 5 μm.

583
584
585 **Figure 2. Stk36/Ulk4 promotes ciliary tip accumulation of Gli2 and Sufu**
586
(A-B) Representative images of immunostaining (A) and quantification (B) of ciliary tip

localized endogenous Sufu (green in A) or Gli2 (blue in A) in control or *Ulk3^{KO} Stk36^{KO}*

(DKO) NIH3T3 cells infected with Stk36-WT or Stk36-AA lentivirus and treated with or

without Shh-N.

(C-F) Representative images of immunostaining (C, E) and quantification (D, F) of

ciliary tip localized Ulk4-HA (green in C, E), Gli2 (blue in C) or Sufu (blue in E) in DKO

cells infected with Ulk4-WT-HA or Ulk4-EE-HA lentivirus.

Primary cilia were marked by acetylated tubulin (Ac-tub) staining (red in A, C and E).

The intensity of ciliary-localized Sufu, Gli2 or Ulk4-HA was measured by Image J. 20

cells were randomly selected from each experimental group for quantification. Data are

mean ± SD from three independent experiments. ***$p < 0.001$ (Student's t test). Scale

bars are 5 μm.

599
600
Figure 3. Overexpression of phosphorylated Ulk4 drives its condensation outside

primary cilia

(A-B) Representative images of immunostaining in HEK293T cells co-transfected with

the indicated Ulk4-HA and Fg-Stk36 expression constructs. Arrowheads indicate Ulk4

(green in A and B) and Stk36 (red in A and B) puncta.

(C) Representative images of immunostaining in HEK293T cells transfected with Ulk4-

EE-HA or Ulk4-AA-HA expression construct. Arrowheads indicate Ulk4-HA (green in C),

Gli2 (red in C) or Sufu (blue in C) puncta.

(D) Representative images of HEK293T cells expressing the indicated constructs before

and after treatment with 5% 1, 6-hexanediol for 5 minutes.

Scale bars are 6 μm.

612
613
614 **Figure 4. Stk36-mediated phosphorylation promotes Ulk4 SUMOylation**
615
(A-B) Western blot analysis of Ulk4 SUMOylation in HEK293T cells co-transfected with

the indicated Fg-Stk36 and Ulk4-HA constructs. Cells were lysed in the presence of the

deSUMOylation inhibitor NEM. Ulk4-HA was immunoprecipitated with anti-HA antibody,

followed by western blot analysis with HA and SUMO1 antibodies.

(C) Western blot analysis of Ulk4 SUMOylation in HEK293T cells transfected with Ulk4-

WT-HA or Ulk4-EE-HA construct and treated with or without TAK-981.

(D) Western blot analysis of Ulk4 SUMOylation in NIH3T3 cells infected with the

indicated Ulk4 constructs and treated with or without Shh-N.

(E) Schematic diagram of human Ulk4 with wild type and mutated sequences of Stk36

sites, SUMOylation site and SIM motif shown.

(F) *in vitro* phosphorylation and SUMOylation of Ulk4. GST-fusion proteins containing

wild type or mutant Ulk4C fragments were incubated with Fg-Stk36-EE purified from

HEK293T cells and recombinant CK1, followed by *in vitro* SUMOylation by incubating

with recombinant SUMO1, E1 and E2.

(G, H) Western blot analysis of Ulk4 SUMOylation in HEK293T cells co-transfected with

the indicated Fg-Stk36 and Ulk4-HA constructs.

(I) Western blot analysis of Ulk4 SUMOylation in NIH3T3 cells infected with Ulk4-WT-

HA or Ulk4-KR-HA construct and treated with or without Shh-N.

(J, K) Co-IP experiments to show Ulk4 self-association promoted by SUMO/SIM

interaction. HEK293T cells were transfected with the indicated Ulk4 constructs, followed

by immunoprecipitation and western blot analysis with the indicated antibodies.

Figure 5. SUMOylation promotes Ulk4 condensation, ciliary tip accumulation, and

pathway activation depending on SIM

(A) Representative images of immunostaining of HEK293T cells co-transfected with Fg-

Stk36-EE and Ulk4-WT-HA, Ulk4-KR-HA or Ulk4-ΔSIM-HA constructs. Arrowheads

indicate Ulk4 (green in A), Stk36 (red in A), or Gli2 (blue in A) puncta.

(B-C) Representative images of immunostaining (B) and quantification (C) of ciliary tip

localized Ulk4-HA (green in B) in NIH3T3 cells infected with Ulk4 shRNA and the

indicated human Ulk4 lentivirus in the presence or absence of Shh-N. Primary cilia were

marked by acetylated tubulin (Ac-tub) staining (red in A) and nuclei by DAPI (Blue in A).

(D) Western blot analysis of Myc-Gli2 phosphorylation on S230/S232 in NIH3T3 cells

expressing the control (Ctrl) or Ulk4 shRNA and the indicated Ulk4 lentiviral constructs

and treated with or without Shh-N.

(E-F) Relative *Gli1* (E) and *Ptch1* (F) mRNA levels in NIH3T3 cells expressing the

control (Ctrl) or Ulk4 shRNA and the indicated Ulk4 lentiviral constructs and treated with

or without Shh-N.

The intensity of ciliary localized wild type and mutant Ulk4-HA was measured by Image

J. 20 cells were randomly selected from each experimental group for quantification.

Data are mean ± SD from three independent experiments. ***p < 0.001 (Student's t

test).

Scale bars in A and B are 6 μm and 5 μm, respectively.

659
Figure 6. Ulk4-EE exhibits constitutive Hh pathway activity depending on its

SUMOylation

(A) Representative images of immunostaining of HEK293T cells transfected with the

indicated Ulk4 expression constructs and treated with or without the SUMOylation

inhibitor TAK981. Arrowheads indicate Ulk4 puncta. Scale bars are 6 μm.

(B, C) Representative images of immunostaining (B) and quantification (C) of ciliary tip

localized Ulk4-WT-HA or Ulk4-EE-HA (green) and endogenous Gli2 (blue) in NIH3T3

cells treated with or without the SUMOylation inhibitor TAK981. Primary cilia were

marked by acetylated tubulin (Ac-tub) staining (red). Scale bars are 5 μm.

(D) Representative images of immunostaining of HEK293T cells transfected with Ulk4-

EE-Fg or Ulk4-EE-KR-Fg expression construct. Arrowheads indicate Ulk4 puncta. Scale

bars are 6 μm.

(E, F) Representative images of immunostaining (E) and quantification (F) of ciliary tip

localized Ulk4-Fg (green) and endogenous Gli2 (blue) in NIH3T3 cells infected with

Ulk4-EE-Fg or Ulk4-EE-KR-Fg lentivirus. Primary cilia were marked by acetylated

tubulin (Ac-tub) staining (red). Scale bars are 5 μm.

(G, J) Western blot analysis of Myc-Gli2 phosphorylation on S230/S232 in NIH3T3 cells

co-infected with Myc-Gli2 and the indicated Ulk4 lentiviral constructs and treated with or

without the SUMOylation inhibitor TAK981.

(H, I, K, L) Relative *Ptch1* (H, K) and *Gli1* (I, L) mRNA levels in NIH3T3 cells infected

with the indicated Ulk4 lentiviral constructs and treated with or without the SUMOylation

inhibitor TAK981.

The intensity of ciliary-localized wild type and mutant Ulk4-HA was measured by Image

J. 20 cells were randomly selected from each experimental group for quantification.

Data are mean ± SD from three independent experiments. ***$p < 0.001$ (Student's t

test).

Figure 7. Phosphorylation-induced SUMOylation promotes kinase/substrate

condensation at ciliary tip

A working model in which Shh induces Ulk4/Stk36/Gli2/Sufu condensation at primary

cilia. In response to Shh, Stk36 phosphorylates Ulk4 on T1021/T1023, which induces

SUMOylation of Ulk4 on K1030. Subsequently, SUMO/SIM interaction drives the

formation of Ulk4 condensates at ciliary tip that recruit Stk36, Gli2, and Sufu.

Figure 3-figure supplement 1. Stk36/Ulk4 condensates are resistant to 1, 6-

hexanediol

Representative images of HEK293T cells expressing the indicated constructs before

and after treatment with 5% 1, 6-hexanediol for 10, 20, and 30 minutes.

Scale bars are 5 μm.

Figure 7-figure supplement 1. The constitutive pathway activity of Ulk4-EE

depends on Stk36

(A, B) Relative *Ptch1* (A) and *Gli1* (B) mRNA levels in NIH3T3 cells infected with the

indicated Ulk4 lentiviral constructs and treated with control (Ctrl) or Stk36 siRNA.

(C) Western blot analysis of Stk36 phosphorylation on pT158/pS159 in NIH3T3 cells co-

infected with the indicated Ulk4 and Stk36 lentiviral constructs.