A unified rodent atlas reveals the cellular complexity and evolutionary divergence of the

dorsal vagal complex

Cecilia Hes[1,2], Abigail J. Tomlinson[3], Lieke Michielsen[4], Hunter J. Murdoch[1,5], Fatemeh

Soltani[1,2], Maia Kokoeva[1,5,6], Paul V. Sabatini[1,2,6]*****

*Corresponding author: paul.sabatini@mcgill.ca

Author affiliations:

1. McGill University Health Centre Research Institute, Montreal QC, Canada
2. Division of Experimental Medicine, Department of Medicine, McGill University,
Montreal QC, Canada
3. Department of Internal Medicine, University of Michigan, Ann Arbor MI, USA
4. Department of Human Genetics, Leiden University Medical Center, Leiden, The
Netherlands
5. Integrated Program in Neuroscience, McGill University, Montreal QC, Canada
6. Division of Endocrinology, Department of Medicine, McGill University, Montreal QC,
Canada
19

Keywords:

DVC, area postrema, nucleus of the solitary tract, cell atlas, GLP1, PDGFRA, CCK, TH

Abstract
The dorsal vagal complex (DVC) is a region in the brainstem comprised of an intricate network
of specialized cells responsible for sensing and propagating many appetite-related cues.
Understanding the dynamics controlling appetite requires deeply exploring the cell types and
transitory states harbored in this brain site. We generated a multi-species DVC cell atlas using
single nuclei RNAseq (sn-RNAseq), thorough curation and harmonization of mouse and rat
data which includes >180,000 cells and 123 cell identities at 5 granularities of cellular
resolution. We report unique DVC features such as Kcnj3 expression in Ca$^+$-permeable
astrocytes as well as new cell populations like neurons co-expressing *Th* and *Cck*, and a leptin
receptor-expressing neuron population in the rat area postrema which is marked by expression
of the progenitor marker, *Pdgfra*. In summary, our findings suggest there are distinct cellular
populations specific to the DVC compared to other brain sites and our comprehensive atlas is
a valuable tool for the study of this metabolic center.
35

Introduction
Regulating appetite in response to changing environmental conditions to maintain energy
balance requires the intricate interplay of multiple organs. Within the central nervous system,
an incredible diversity of discrete brain regions and cell types control appetite([1]). More
specifically, the dorsal vagal complex (DVC) of the brainstem, comprising the area postrema
(AP), nucleus of the solitary tract (NTS) and dorsal motor nucleus of the vagus (DMV), acts as
a key nexus point for metabolic signals, evidenced by the dense expression of many relevant
hormone receptors([2]). Separately, the DVC also serves as the primary site of gut-based signals
via vagal afferents([3-5]). Functionally, the DVC was long considered a site promoting meal
termination([6]) but growing evidence suggests the DVC also regulates long-term energy
balance([7, 8]).
47
Much of the interest in DVC neurons' role in appetite and energy balance stems from their role
as therapeutic targets for obesity and anorexia in cancer cachexia([9, 10]). For many years,
most pharmacotherapies failed to produce efficacious and long-lasting weight loss in obesity
and increase appetite in cancer cachexia([11]). More recently however, the glucagon-like
peptide 1 receptor (GLP1R) agonists are eclipsing the weight loss achieved by previous
generations of therapeutics([12]). While expressed across the central nervous system, GLP1R
expression is particularly enriched in DVC neurons([13]) and it is the DVC that mediates many
of the effects of GLP1R agonists([14]). Separately, an emerging strategy for treating cancer
cachexia is inhibiting growth differentiation factor 15 (GDF15) signaling in the DVC by blocking
the action of GDNF Family Receptor Alpha Like (GFRAL)([15]), the receptor for GDF15([16]).
Given the clinical significance of the DVC for this spectrum of conditions, understanding the

molecular heterogeneity and function of this region is essential to understand appetite
regulation without major side effects.
61
Traditionally, identification and ensuing study of DVC cell types was limited to known
neurotransmitter and receptor-expressing cell types. For instance, the cholecystokinin and
monoamine-expressing cells, demarcated by tyrosine hydroxylase (*Th*) expression, were
previously shown to be non-overlapping cell populations with unique roles in appetite
suppression(*17*). More recently, the advent of single-cell mRNA sequencing approaches made
unbiased molecular censuses of heterogenous populations possible. Such studies focusing
on the mouse have revealed the complexity of the mouse DVC, comprised of dozens of
transcriptionally distinct cell types(*2, 18, 19*). Moving forward, unifying information from multiple
single-nuclei RNA sequencing (snRNA-seq) datasets of the DVC into a reference atlas is
needed to more fully understand the cells within the DVC and generate a common and scalable
classification for cell classes. Such a reference atlas is also needed to contrast the cell types
found within the mouse to other research models, including the rat, which are prominent
models for DVC biology(*2, 20*).
75
To address this, we generated a comprehensive snRNA-seq based atlas of the mouse and rat
DVC. This atlas features mouse and rat labelled cells with increasing granularity and an
accompanying computational toolbox including the mouse and rodent atlas label transfer
environment using treeArches. The mouse DVC atlas reveals a higher degree of cellular
complexity than previously appreciated. The cellular census of the rat DVC exhibits many
similarities to the mouse but also rat-specific cells including a novel population of leptin
receptor and *Pdgfra*-expressing neurons localized to the AP. We then used this unified atlas
to uncover the cell types that transcriptionally respond to meal consumption which may hold
the potential of suppressing appetite without gastrointestinal side effects.
85

Results
The murine dorsal vagal complex cell classes
To generate a snRNA-seq atlas of the mouse DVC, we isolated the DVC from 30 adult
C57BL/6J mice and subjected them to 10X Genomics-based snRNA barcoding and
sequencing (**Fig.1A**). From the CellRanger pre-processing pipeline(*21*) we obtained 110,167
cells of which 99,740 were retained after processing. Cell identities were initially mapped to 5
databases from different brain regions and then curated manually using 473 cell identity
markers (**Tables S1-S2**). We distinguished clusters of neurons and glial, vascular and
connective tissue cells, a first layer of cell identity (**Fig.1B**; **Table S3**). We further describe a
second layer of cellular resolution, which are subgroups of the first layer that contain the cell
classes (**Fig.1C**; **Table S3**).

Among the ten neuronal cell classes identified at layer 2 resolution, we found two major groups

which we called 'magnaclass1' and 'magnaclass2' (**Fig.S1**). Magnaclass1 neurons

predominately express genes required for inhibitory neurotransmission (e.g. *Gad1*, *Gad2*) and

share transcriptomic markers with the Sall3-class of neurons, whereas magnaclass2 neurons

express mostly excitatory transcripts (e.g. the glutamate transporter VGLUT2) also found in the

Lhx2-class and monoamine-class (**Fig.1D-E**; **Fig.S1**). However, as described in other brain

areas([22, 23]), evidence of co-transmission/co-release of multiple primary neurotransmitters

was found in at least 55% of neurons regardless of their cell class (**Fig.1D-E**; **Table S4**). We

also identified 8 transcriptionally distinct neural classes in addition to the magnaclasses. The

Golli-class expresses myelin-associated genes such as *Mag* and *Mog*([24-26]) (**Fig.1D**; **Table**

S5). Moreover, we describe the corelautus-class (Latin: 'washed nucleus'), distinguished by a

large number of downregulated genes in the nuclei compared to other neuronal classes, and

a class with genes normally upregulated in astrocytes (e.g. *Slc6a11*) and related to clearance

and uptake of glutamate, gamma-aminobutyric acid (GABA), Ca^+ and bicarbonate ([27, 28]),

which we named 'pH-related class' (**Table S5**). When interrogating snRNA-seq data from the

hypothalamus([29]) (**Fig.S2**), another brain site important for energy balance, we could not find

clusters sharing all markers with our neuronal classes, indicating DVC-specific neuronal

programs (**Fig.S2C**).

<u>The murine DVC cells at their highest resolution</u>

To better define the heterogeneity of the mouse DVC, we further assigned a clustering-based

third layer of cellular resolution resulting in fifty cell identities, of which 15 are non-neuronal

(**Fig.2A**). At this resolution we differentiated between two major groups of astrocytes.

Differential expression in synapse-related factors revealed a group of highly excitable

astrocytes with high expression of the inwardly rectifying GIRK1 (*Kcnj3*) channel([30]) and lack

of expression of the AMPAr subunit GluA2 (*Gria2*), additionally rendering them K^+/Na^+/Ca^+-

permeable([31, 32]) (**Fig.2B**). We speculate these astrocytes are specific to the DVC as we could

not find Kcnj3+; Gria2- astrocytes in the hypothalamus, cortex or hippocampus([29, 33])

(**Fig.S2D**; **Fig.S3**). In addition, we distinguished a spectrum of DVC mature oligodendrocytes,

including a group resembling the intermediate oligodendrocytes described in the other brain

sites that decrease in number with age([34]) (**Fig.2C**). This gradient of expression of markers

including *Fyn*, *Opalin* and *Anln* in oligodendrocytes (**Fig.2C**), do not respond to a

developmental trajectory (**Fig.S4**). Although 'myelinating-intermediate' and 'myelinating'

oligodendrocytes include cells with overlapping gene expression programs, there is no

evidence that each oligodendrocyte will follow a single trajectory (**Fig.S4**). Furthermore, to

properly separate some cell identities, we performed sub-clustering in three DVC clusters (i.e.

clusters 23, 26 and 27) (**Fig.S5A**). This permitted us to label pre-myelinating oligodendrocytes,

endothelial and mural cells, and distinguish two fibroblast sub-types (**Fig.S5**). One of these
subtypes expresses high *Stk39* previously described in fibroblast stress-response(*35*) and so
was termed 's-fibroblasts' for stress-associated fibroblasts (**Fig.S5B**), which we failed to detect
in the hypothalamus (**Fig.S2E**). Additionally, there was no evidence of reactive microglia as we
mapped minimal levels of the activation genes *Cxcl10*, *Cd5*, *Cxcl9* and *Zbp1*(*36*), therefore we
posit that our DVC microglial cells are in a basal state (**Fig.S6**). We also obtained the DVC gene
expression markers of the main cell types in the central nervous system and neurons (**Fig.2D**).
142
To more thoroughly classify DVC neural types, we subset and re-clustered the 10 neural
classes from layer 2 which resulted in 35 neuronal identities as a third layer of granularity
(**Fig.3A**; **Table S5**). These identities include 'preganglionic' or *Chat*-expressing neurons and a
large number of neurons with unspecific markers which we called 'mixed neurons' (**Table S5**).
A mixed neurons-like group of cells was previously described in the murine hypothalamus(*37*).
Additionally, we split the monoamine-class into four layer-3 cell identities (i.e. M0, M1, M2 and
M3) with specific cell markers (**Fig.3B**; **Table S5**). M0 and M1 neurons express genes for
enzymes synthesizing norepinephrine and serotonin, but M0 expresses minimal levels of the
transporter VMAT2 (*Slc18a2*) (**Fig.3C**; **Table S5**). Due to the low proportion of cells expressing
Th and *Tph2* in M2 and M3 neurons, we consider them monoamine-modulators as they seem
to synthesize trace amines (**Fig.3C**; **Table S5**). Of note, the receptor for GDF15, GFRAL, is
expressed heavily by a subset of M0 neurons, some of which co-express the calcium sensing
receptor (*Casr*) gene (**Fig.3C**).
156
157   We named 'GLP1' a neuronal cluster with high expression of the pre-proglucagon gene (*Gcg*)
158   (**Fig.3D**; **Table S5**), which also expresses high levels of both the prolactin and leptin
159   receptors(*38*) (**Fig.3E**). Additionally, these cells have the highest proportion of DVC neuronal
160   co-expression of glutamate and GABA related genes (**Fig.3F**).
161
162   In our analysis we found a subset of monoaminergic neurons that express the cholecystokinin
163   (*Cck*) gene (**Fig.4A**). While the *Cck* and *Th*-expressing DVC neurons are considered to be non-
164   overlapping cell types with distinct physiologies(*39*), we confirmed co-expression of *Th* and
165   *Cck* results by *in-situ* hybridization (**Fig.4B**). The majority of this overlap occurs in cells
166   belonging to the COE-Atp10a and monoamine class (i.e. M0, M3) cell identities (**Fig.4A**).
167

<u>The murine DVC cell atlas</u>
To generate a comprehensive murine DVC database from multiple datasets, we utilized
treeArches(*40*) to harmonize our snRNA-seq DVC labels with that from a publication by Ludwig
and collaborators (i.e. the 'Ludwig dataset')(*2*) (**Fig.5A-B**). After building our initial cell hierarchy
(**Fig.S7**), we integrated our dataset with the Ludwig dataset, giving a total of 171,868 cells

(Fig.5B). Next, the labeled cell identities from the Ludwig dataset(*2*) were incorporated into our
tree based on the similarity between the transcriptomic programs of our labeled identities and
the Ludwig-labeled identities using progressive learning through scHPL(*41*) (Fig.5A). Some of
the Ludwig dataset identities failed to be incorporated into our tree, like the oligodendrocyte
precursor cells (OPCs), which contain a mix of OPCs and pre-myelinating oligodendrocytes
(Fig.S8A). The mix of cell programs makes it impossible for treeArches to correctly add one
branch of the Ludwig OPCs to the tree as ideally it would be added as a subpopulation of both
the OPCs and pre-myelinating oligodendrocytes, which is not biologically possible. The
resultant murine hierarchy has a fourth layer of cellular resolution as some of Ludwig identities,
mainly neurons, are subgroups of our cell identities (Fig.S8B).
183
We confirmed the validity of our cell hierarchy using three methods. First, by using the learned
model to predict the labels of the cells we used to develop them (i.e. this study and the Ludwig
dataset), for which we obtained highly concordant cell labeling (Fig.5C). Secondly, we used
the learned model to predict the labels of another publicly available dataset from Dowsett and
collaborators (i.e. the 'Dowsett dataset')(*18*) (Fig.5D). Using this approach, we were able to
successfully label 95% of these cells at high resolution (i.e. layer 3 or 4 of cell identity
granularity); further validating our combined DVC labels. Finally, we manually mapped the
marker genes from Ludwig(*2*) and Dowsett(*18*) neuronal clusters in our dataset and confirmed
the sub-groups of neurons among our identities pointed out by treeArches (Fig.S8B).
193
An integrated rodent DVC cell hierarchy
To identify rat DVC cell clusters and subclusters and contrast these with the mouse DVC atlas,
we generated a *de novo* ad libitum-fed rat dataset (cells n=12,167) (Fig.6A) and labeled it by
mapping the top 5 gene expression markers of each of our murine layer-3 cell identities (Table
S6). After manual curation at three granularities (Fig.6B-C; Table S7), we obtained 52 DVC rat
identities at high cellular resolution (Fig.6C) including a small group of 'unspecific' cells (n=35)
that we could not confidently label because its top gene expression markers have unknown
functions and none of our known identities' genes clearly marked this population (Fig.6B-C;
Table S6).
203
We established 10, layer-3 neuronal identities not previously found in the murine data (Fig.6C;
Fig.S9A; Table S7). Notably, two of those did not belong to any of the existing murine neuronal
classes (i.e. layer-2 of cellular granularity), therefore we created two new layer-2 classes
(Fig.6B). One of these, the immunity-related neurons, have high cystatin C, *Inpp5d* and *Csf1r*
expression, genes commonly associated with immunity-related processes in microglia and
with immunomodulation in cancer(*42-44*) (Fig.7; Table S7). The second neural class identified
in rat but not mouse DVC were the 'Pdgfra neurons' (Fig.7). These neurons express *Pdgfra*,

Arhgap31 and *Itga9*, known markers for progenitor cells for which we named this cell class
'ortus-akin' (Latin: 'origin') because they are similar to cells giving origin to other cells(*45*)
(**Fig.7A**; **Table S7**). Interestingly, this neuronal class not only expressed *Pdgfra*, but also other
OPC signature genes (**Fig.7A**). Mapping the expression levels of metabolic-associated
receptors revealed that both immunity-akin and ortus-akin express GFRAL, prolactin receptor
and leptin receptor (**Fig.7B**), although they seem to have a varied neurotransmitter profile
(**Fig.S9B**). To confirm the presence of *Pdgfra*-expressing neurons in the rat DVC, we performed
immunostaining in the mouse and rat brainstem. Within the mouse, we found PDGFRA-
immunoreactivity(IR) in the two morphologically distinct cells; matching OPC and blood vessel-
associated fibroblast morphologies(**Fig.7C**). However, within the rat, we identified PDGFRA-IR
in cells with three morphologies, one of which resembled neurons with an enlarged cell body
and protruding cellular processes (**Fig.7C**). To determine if these PDGFRA-IR cells were
neurons, we co-stained with the neural marker, HuC/D and noted several PDGFRA and HuC/D
copositive cells (**Fig.7D**); confirming our sequencing data that a PDGFRA-expressing neural
class exists in the rat DVC and localizing these cells to the area postrema.
226
To construct an integrated rodent DVC cell atlas, we incorporated the labeled rat dataset into
our hierarchy using treeArches(*40*) (**Fig.8**). The rodent data was projected onto the mouse
datasets (**Fig.8A**) and the 42 high resolution rat cell identities were appended to the tree
(**Fig.8B**; **Fig.S10**). The final hierarchy has 123 labels of which 99 are high resolution according
to our cellular granularity (i.e. layers 3 to 5) (**Fig.8B**). In addition it has 20 layer-2 identities and
7 cell identities that are novel from the rat dataset (**Fig.8B**).
233
<u>*Hcrt* and *Agrp* expression in the DVC</u>
Since *Agrp* has been described to be expressed in the DVC(*46*), we mapped this and other
neuropeptide genes in DVC neurons. Surprisingly, we found *Hcrt* and *Agrp* mRNA in neurons
in our dataset (**Fig.S11A**). We then performed *in-situ* hybridizations for these transcripts in both
the mouse hypothalamus and DVC. While both *Hcrt* and *Agrp* were readily detectable in the
hypothalamus, we failed to observe signal within the DVC (**Fig.S11B**). Since we also found
expression of *Hcrt* and *Agrp* in neurons of the Dowsett and Ludwig datasets(*2, 18*) (**Fig.S11C**),
we then mapped the sequenced reads from these transcripts and confirmed that most of them
mapped to intronic and non-coding regions of these genes (**Fig.S11D**). Furthermore, our rat
DVC dataset was pre-processed excluding introns from the reads mapping, and the *Hcrt* and
Agrp expression levels observed is minimal (**Fig.S11E**), which is in agreement with the non-
coding location of the reads. These data explain absence of signal from our *in-situ*
hybridizations and suggests DVC cells do not produce HCRT or AGRP neuropeptides.
247

Meal-related transcriptional programs in the murine DVC

As the DVC acts as a primary node for meal-related signals(*47*), we wanted to define which of

our cell identities responded to meal consumption and thus performed differential gene

expression analysis between DVC cell types from mice that were euthanized under fasting

conditions, with ad libitum access to food or two hours post-refeeding following an overnight

fast (**Fig.1A**; **Fig.9**). This analysis revealed widespread transcriptional changes across nearly

every neural cell identity, the exceptions being corelautus, Golli class and pH classes (**Fig.9A**).

Additionally, the size of the transcriptional changes were relatively small, with approximately

90% of the differentially expressed genes having increases or decreases between 0.5 and 1.0

log2 fold-change, with only ~10% of the gene changes being ≥1 log2 fold-change between

conditions (**Fig.9A**). Of the responding neurons, magnaclasses 1 and 2, which represent large

populations of predominately inhibitory and excitatory neurons (**Fig.S1**), showed the greatest

number of differentially expressed genes following refeeding for both comparisons refed

versus fasting, and refed versus ad libitum food access (**Fig.9A**). To understand how refeeding

transcriptionally alters inhibitory compared with excitatory DVC neurons, we compared

differentially expressed genes between these two classes across conditions. Intriguingly,

despite being in transcriptionally distinct classes, we found meal consumption altered many of

the same genes (**Fig.9B-D**), suggesting these cells may be receiving similar postprandial

signals which are propagated via these cells in unique manners. This is supported by the

similar receptor expression profile of magnaclass 1 and 2 (**Fig.S12**). One notable exception is

Cckbr which is upregulated in magnaclass 2 and downregulated in magnaclass 1 in refeeding

when compared with ad libitum fed mice, while in refed versus fasted mice (**Table S9**). We also

note that oligodendrocytes contained the highest number of differentially expressed genes in

refeeding (**Fig.9A**), however the roles of oligodendrocytes in refeeding responses within the

DVC have yet to be determined and the biologic significance of these changes are presently

unclear.

Discussion

The recent success of DVC-based therapies(*12, 48*) have moved DVC biology to the forefront

of weight control therapies and there is now clear clinical impetus to fully define DVC cell types

and their functions. Previous studies detailing DVC cell types describe a high degree of

heterogeneity(*2, 18, 19*), however, these studies lacked an in-depth analysis of

neurotransmitter profiles of neurons, any cross-species evaluation of DVC cell types, or an

analysis of how meal consumption transcriptionally regulates these cells. We therefore

developed *de novo* mouse and rat DVC cell atlases to address these points.

To more fully describe the cellular complexity within the DVC, we combined a pipeline of cell
label transfer using brain existing published datasets and thorough curation of cell identities,
which resulted in a 4-layer mouse cell hierarchy and a 5-layer rodent cell hierarchy of the DVC.
Within this hierarchy we found several unique features of DVC glial cells. First, we found highly
excitable Ca$^+$-permeable astrocytes in the DVC in which *Gfap* is almost exclusively expressed
and they coexpress *Kcnj3* (GIRK1 potassium channel). These cells largely lack *Gria2*,
consistent with findings in Bergmann glial cells in which colocalization of *Gria2* and GFAP was
found to alter neuron-glial interactions([49]). Comparing Ca$^+$-permeable DVC astrocytes to
hypothalamic astrocytes([29]), we found the same expression pattern with minimal overlap
between *Gria2* and *Gfap.* However, DVC astrocytes seem to have some unique properties
when compared with hypothalamic, cortical and hippocampal astrocytes. Principally DVC
astrocytes express the GIRK1 channel, likely rendering them more excitable. Additionally,
while we identified hypothalamic fibroblasts which express high *Aldh1a1* similar to the DVC p-
fibroblasts, we failed to find a group equivalent to s-fibroblasts. Other cell populations seem to
be consistent to what has been found in other brain sites, like the intermediate
oligodendrocytes that are found enriched in the hypothalamus and the corpus callosum in
mice([34]). In summary, these findings show that some cell populations may be specific to the
DVC compared to other brain sites and we were able to capture them in our atlas, further
supporting its utility for studying this brain region.
303
Within the mouse neural classes, we found overlap between *Cck* and *Th*-expressing cells. As
these cells have previously been described as non-overlapping, this was unexpected([17]). Our
in-situ hybridization confirmed the presence of *Cck/Th* positive cells within the NTS and AP.
We believe the discrepancy may be due to detection methods as these cells were initially
defined as non-overlapping using immunofluorescent detection of *Th* protein and a *Cck*
genetic reporter([17]). Within the NTS, *Cck* neurons are strongly aversive whereas *Th*-labelled
neurons are non-aversive([50]). The role of dual Cck/Th positive neurons remains to be
determined. While no functional characterization of *Cck* and *Th* neurons of the AP has been
performed, we posit that as some *Th*-expressing AP neurons also express *Gfral* and *Casr*, both
having potently aversive ligands (e.g. GDF15, deoxynivalenol)([16, 51]); they are likely to
mediate aversive anorexia.
315
Our snRNA-Seq data also resulted in detected transcripts for both *Agrp* and *Hcrt*. While both
neuropeptides have long been considered to have hypothalamic-exclusive expression, a
recent report describes *Agrp*-expressing cells within the mouse DVC([46]). To confirm our
sequencing data, we performed *in-situ* hybridizations for both *Agrp* and *Hcrt* but failed to
detect any transcripts within the DVC, in agreement with this finding is that the mapped reads

within the *Agrp* and *Hcrt* gene overwhelmingly mapped to non-coding regions, suggesting to

us that the AGRP and HCRT neuropeptides are not produced by DVC cells in rodents.

Beside the classical dichotomy of inhibitory or GABAergic and excitatory glutamatergic,

neurons, subpopulations of neurons have been shown to be capable of co-releasing GABA

and glutamate(*52*) and other neurotransmitter combinations(*53*). By co-expression analysis of

genes implicated in synthesis or neurotransmitter vesicle packing, we confirmed that over 50%

the DVC neurons co-transmit/co-release more than one primary neurotransmitter. This complex

functionality of the DVC is further exemplified by the presence of neurons with seemingly mixed

programs which we called 'mixed neurons'. Similar groups have been described in the

hypothalamus(*37*) suggesting that some neurons may have hybrid-functionalities. As these

mixed neurons share transcriptomic features with many other cell identities, we foresee them

to be difficult to target or to be affected by treatments not directed to them as an off-target

effect. Finally, we were not able to clearly associate our neuronal classes with the cell classes

that had been previously identified in hypothalamic snRNA-seq data(*29*) which is in line with

the complementarity of different metabolic centers in the brain suggesting that the

hypothalamic and brain stem neuronal populations involved in feeding control and energy

balance, share only some functionality, possibly reflective of their complementary roles in

metabolic regulation.

To perform a cross-species analysis, we generated a rat-based DVC atlas and compared it

with the mouse DVC. Our rat dataset was smaller than the mouse data, it is possible some of

the of the new neuronal cell identities (e.g. M_sema, M4) may contain cells belonging to

multiple cell types, as treeArches did not appended these identities to the tree. Despite this,

we corroborated two new cell types in the rat DVC that are not present in mice the *Pdgfa*-

expressing (ortus-akin class) and the immunity-related neurons. *Pdgfra* is a well described

marker gene for OPCs and fibroblasts within the central nervous system(*45*) but, to our

knowledge, has not been associated with neurons. Notably, although the PDGFRA immuno-

signal in HuC/D+ neurons is clearly discernible, it is less intense than in OPCs and fibroblasts.

Since it was previously demonstrated that OPCs can give rise to neurons in the adult brain(*54*),

it is possible that these PDGFRA+/HuCD+ cells in the AP represent a transient stage of

PDGFRA+ OPCs differentiating into neurons.

Regardless of their maturity, we speculate that the Pdgfra neurons (ortus-akin class) have roles

in appetite regulation as they express leptin receptor and *Gfral*. Interestingly, while multiple

studies have failed to detect *Lepr* via Cre-based reporters(*38, 55*) or leptin-induced

phosphorylated STAT3 in the AP of mice(*56*), *Lepr* and leptin-induced phosphorylated STAT3

are detectable within the rat AP(*57-59*). We believe one explanation for this species
discrepancy in *Lepr* expression is the absence of the ortus class neurons in mice.

DVC-targeting obesity therapeutics, including amylin and GLP1 mimetics, produce clinically
relevant weight loss but are associated with multiple gastrointestinal adverse events and there
is a need to develop new obesity therapies that can reduce appetite without unwanted
gastrointestinal side effects(*60, 61*). Indeed, the DVC cells that regulate appetite suppression
and nausea are separable(*50*), making targeting such cells an attractive target for
pharmacotherapy. As meal consumption is generally not associated with gastrointestinal
distress, we sought to identify refeeding responsive cells within the mouse DVC. From this
analysis, we found nearly all neural cell identities are transcriptionally altered by meal
consumption. Therefore, identifying neurons that promote non-aversive anorexia via the
transcriptional response to refeeding as the basis of pharmacotherapy is unlikely to yield
specific targetable pathways.

In general, the limitations of our study are that it is based on nuclear RNA which may reflect a
fraction of biological changes, and that cell viability and isolating nuclei from cells are
challenging. In addition, for meal consumption, we were unable to analyze responses in
individual mice but due the number of cells recovered from each mouse DVC, pooling was the
most amenable option in our pipeline.

As the amount of snRNA-Seq data performed on the DVC continues to grow, there is a need
for a common, scalable label set that can be applied to new studies and integrate new
sequencing data. To develop such a label set, we started with a *de novo* DVC atlas, we
generated three layers of cellular granularity for both neural and non-neuronal cell types. Then,
using the treeArches pipeline(*40*) we harmonized our snRNA-Seq labels with the Ludwig
dataset(*2*) and increased our cellular granularity from three to four layers of resolution. This
approach of manual curation and harmonization enhanced the reach of our cell atlas to capture
the complexity of the DVC neurons since not all groups seem to contain easily delineable
subgroups of cells. For example, eleven of the labels incorporated into our cell hierarchy in this
pipeline belong to the monoaminergic neuronal class, which seem to encompass multiple
subgroups of specialized cells. We further constructed the rodent DVC cell atlas which resulted
in a fifth layer of cell identity to the monoaminergic M3 neurons. Although the number of
identities increases as layers are more specific, this gives this unified atlas greater adaptability
to different research needs. Additionally, our labels can be accurately harmonized with new
databases regardless of size and new DVC snRNA-Seq data can be added to the tree to
generate new branches. Thus, our unified rodent DVC atlas is highly amenable to future
research elucidating new biologic and therapeutic insights into DVC cell types.

Materials and methods

Experimental design

To generate a snRNA-Seq based atlas of the mouse dorsal vagal complex (DVC), we isolated the DVC from 30 adult C57BL/6J mice and subjected them to 10x Genomics-based single-nuclei RNA barcoding and sequencing. Mice were either fasted overnight (n=5), fasted overnight and then refed 2 hours prior to euthanasia (n=10), fed ad libitum (n=5), injected with LiCl (n=5) or injected with vehicle (n=5). Conserved cell clusters across these groups were then identified.

Animals

All animals were bred in the Unit for Laboratory Animal Medicine at the University of Michigan and the Research Institute of the McGill University Health Centre. Procedures performed were approved by the University of Michigan Committee on the Use and Care of Animals, in accordance with Association for the Assessment and Approval of Laboratory Animal Care and National Institutes of Health guidelines. Alternatively, procedures were approved by the animal care committee of McGill University.

Mice for sn-RNA sequencing assay

Wild-type C57BL/6J mice (Jackson laboratories) were given ad libitum access to food (Purina Lab Diet 5001) and water in temperature-controlled (22°C) rooms on a 12-hour light-dark cycle with daily health checks. Mice were euthanized by isoflurane anesthesia and decapitated, then the brain was removed from the skull and aligned in a brain matrix. The cerebellar cortex was removed and the DVC was microdissected using AP-centric targeting and flash frozen in liquid nitrogen. Frozen tissue was stored at -80°C until use.

For assessment of feeding-related gene expression, mice were either ad libitum fed or food was removed at the onset of the dark cycle and the animals were fasted overnight. Animals were then euthanized the next morning under ad libitum fed or fasting conditions. A subset of fasted animals (n=5) were provided food and allowed to refeed for 2 hours prior to euthanasia.

Mice for *in-situ* hybridization assays

Mice were provided with ad libitum access to food (Purina Lab Diet 5001) and water in temperature-controlled (22°C) rooms on a 12 h light-dark cycle with daily health status checks. C57BL/6J mice (Jackson laboratories) were euthanized by isoflurane anesthesia followed by CO_2 asphyxiation. Mice were then trans-cardially perfused with phosphate buffered saline for 3 minutes followed by 5 minutes perfusion with 10% formalin. Brain tissue was then collected

and postfixed for 24 hours in 10% formalin before transfer into 30% sucrose for a minimum of

24 hours. Brains were then sectioned as 30 μm thick, free floating sections.

<u>Rats</u>

Wild-type Sprague Dawley rats (Charles River) were provided with given access to food (Purina

Lab Diet 5001) and water in temperature-controlled (22°C) rooms on a 12-hour light-dark cycle

with daily health checks. Rats were euthanized by intraperitoneal injections of pentobarbital

and decapitated, then the brain was removed from the skull and aligned in a brain matrix. The

cerebellar cortex was removed and the DVC was dissected out in an AP-centric manner then

flash frozen in liquid nitrogen. Frozen tissue was stored at -80°C until use.

<u>Isolation of nuclei from DVC tissue</u>

Frozen tissue (from 2 to 3 mixed sex mice of the same treatment or 1 rat per sample) was

homogenized in Lysis Buffer (EZ Prep Nuclei Kit, Sigma-Aldrich) with Protector RNAase

Inhibitor (Sigma-Aldrich) and filtered through a 30mm MACS strainer (Miltenyi). Filtered

samples were centrifuged at 500 rcf for 5 minutes at 4°C, and pelleted nuclei were

resuspended in fresh-made wash buffer (10 mM Tris Buffer, pH 8.0, 5 mM KCl, 12.5 mM MgCl2,

1% BSA with RNAse inhibitor) before undergoing a second filtration and centrifugation. The

pelleted nuclei were resuspended in wash buffer with propidium iodide (Sigma-Aldrich) and

underwent FACS sorting on a MoFlo Astrios Cell Sorter to remove debris. PI+ nuclei were

collected and centrifuged at 100 rcf for 5 minutes at 4°C and resuspended in wash buffer to

obtain a concentration of 750-1,200 nuclei/μL in preparation for sequencing.

<u>Single-nuclei RNA-sequencing</u>

Library preparation was handled by the Advanced Genomics Core at the University of

Michigan. RT mix was added to target approximately 10,000 nuclei recovered per sample and

loaded onto the 10× Chromium Controller chip. The Chromium Single Cell 3′ Library and Gel

Bead Kit v3, Chromium Chip B Single Cell kit, and Chromium i7 Multiplex Kit were used for

subsequent RT, cDNA amplification, and library preparation, as instructed by the

manufacturer. Libraries were sequenced on an Illumina NovaSeq 6000 (pair ended with read

lengths of 150 nt).

<u>Processing of the snRNA-seq files</u>

From the snRNA-seq, we obtained four FASTQ files (two per sequencing lane, containing

forward and reverse sequences respectively) per mouse treatment and two files with the

forward and reverse sequences for the rat samples. We pre-processed the murine FASTQ files

using either CellRanger 5.0.1 with inclusion of introns or CellRanger 7.0.1 with default

parameters (which include intronic sequences) to align sequencing reads to the murine
genome(*21*). We aligned to the *Mus musculus* refdata-gex-mm10-2020-A reference genome.
For the rat files, we used CellRanger 3.0.1 with no introns inclusion and aligned using the *Rattus*
norvegicus genome assembly Rnor_6.0(*21*). We obtained 110,167 and 13,360 cells from the
murine and rat experiments, respectively. In addition to these two datasets, we also used the
following steps to process the murine Dowsett dataset(*18*) publicly available as a raw gene
expression matrix set of files. We processed the gene expression matrices in R(*62*) with Seurat
v5 and SeuratObject v5 to remove low quality cells(*63*). For processing and analysis, R
v.4.3.1(*62*) was used. We filtered cells with ≥500 number of genes mapped and >1.22 RNA
unique molecular identifiers (UMIs) counts/genes mapped ratio (based on the first 2
percentiles per sample) (**Table S10**). Subsequently, we merged the data per treatment,
calculated the median RNA content of clusters at resolution 1.0 and those below the first
quartile of the median distribution were removed (**Table S11**). Additionally, we used
DoubletFinder v.2.0.3 based on the expected doublets rates by 10X Genomics after adjusting
for homotypic doublets modeled as the sum of squared annotation frequencies(*64*) (**Table**
S12). On each iteration we processed the data with libraries scaled to 10,000 UMIs per cell
and log-normalized. We identified the most variable genes computing a bin Z-score for
dispersion based on 20 bins average expression. We regressed UMI counts and used
principal component (PC) analysis for dimensionality reduction on to the top 2,000 most
variable genes. We used the first 30 PCs for *k*-nearest neighbors clustering and for uniform
manifold approximation and projection (UMAP) projections using Seurat v5(*63*) default
parameters. Murine samples integration in a single matrix was done with Harmony(*65*), after
which clustering and UMAP projections were performed using the harmony embeddings
instead of PCs.
494

Mapping of our murine data to existing brain databases
We mapped the resulting murine integrated high quality singlets to the celldex v1.12.0 built-in
mouse RNA-seq reference and other four mouse databases(*37, 66-68*) (**Table S1**) using
SingleR v2.4.1(*69*) and scRNAseq v2.16.0(*70*) packages in R(*62*). Since each database has
its own cell type nomenclature, in our murine dataset we established a 'likely-cell type' to be
unanimous among all databases, for example if all databases labeled a cell as 'neuron'. We
initially labeled the cells with conflicting or unassigned label by one or more databases as
'unknown'. In order to confirm the cell type of all the cells labeled and to identify the unknown
cell identities, we mapped 473 markers from the literature. For this, the average expression per
marker gene was obtained on scaled data for each cluster at resolution 1.0 (i.e. 48 clusters)
(**Table S2**). We also manually visualized UMAP projections from all these markers in our
dataset. Some clusters (e.g. cluster 27) showed in the previous step to contain cells belonging
to different cell types, so we subset and reprocessed these clusters using the Harmony

embeddings to assign their cell identity based on marker expression at the smallest resolution

that allowed to separate them (**Fig.S5**).

<u>Assigning high resolution clustering-based cell identity in the murine dataset</u>

In the murine dataset, for all non-neuronal clusters at resolution 1.0 gene expression markers

were obtained using the FindConservedMarkers function from Seurat v5(*63*) package in R(*62*).

The function was run using the MAST algorithm(*71*) on scaled data with a log fold-change (FC)

threshold of 0.25. Only upregulated marker genes detected on 40% of all cells in that cluster,

were retained. Since the FindConservedMarkers gives a log2FC output per sample, we

calculated the average log2FC across samples and the proportion of samples for which a gene

had an adjusted p-value ≤0.05. A marker gene was considered as such if it had an average

log2FC>1 across samples and was significant (i.e. had an adjusted p-value ≤0.05) in more

than 80% of the samples. Furthermore, neurons were subset, reprocessed and subjected to

new clustering through Seurat v5(*63*). We obtained the gene markers per cluster at resolution

1.0 as described. We named the clusters based on the known functions of the

upregulated/downregulated genes (e.g. myelinating oligodendrocytes), peculiarities of the cell

groups (e.g. Ca$^+$-permeable astrocytes) or their upregulated gene expression markers if the

previous methods were unsuitable (e.g. Ano2 neurons) (**Table S5**). All visualizations, unless

otherwise stated, were done using ggplot2 v3.5.0 and ComplexHeatmap v2.18.0 packages in

R(*62*).

<u>Oligodendrocytes trajectory inference</u>

We subset the oligodendrocytes (i.e. pre-myelinating, myelinating intermediate and

myelinating) in our murine dataset and placed each cell on an inferred cellular trajectory using

their transcriptomic data in R(*62*). We used the SCORPIUS v1.0.9 package(*72*) with default

parameters to perform dimensionality reduction by PCA on log-normalized counts and to

obtain the trajectory image, with a random seed of 1,000. We additionally performed the

analysis on the Harmony embeddings from the original data integration.

<u>Gene expression files format interconversions</u>

To convert Seurat ('RDS') DVC datasets to 'h5ad' format and vice versa, we used reticulate

v1.35.0(*73*) and anndata 0.7.5.6(*74*) packages in R(*62*) with import of scipy, scanpy, numpy

and anndata modules from Python v3.9(*75*). To convert .csv and .tsv datasets (i.e. the Ludwig

dataset(*2*) and the cortex/hippocampus database(*33*)), we converted those to 'h5ad' format in

Python v3.9(*75*).

 Murine DVC hierarchy construction

We converted our murine labeled count matrix to h5ad. We used treeArches(*40*) in Python
v3.9(*75*) to create a manual tree using the three layers of cellular granularity in our database.
We reprocessed the samples using the treeArches pipeline(*40*) to normalize the count data
(counts normalized per cell and then log-normalized), identify the top 2,500 most variable
genes and integrate the samples using scVI(*76*). This reference latent space obtained after
integration was used to generate the UMAP embeddings. We trained our manual tree based
on the cell identity layer-3 labels using default parameters.

We subset the Ludwig dataset to match the initial 2,500 most variable genes from our datasets
and performed surgery to incorporate labels from the Ludwig dataset(*2*). The two layers of
granularity established by Ludwig(*2*) were combined in the variable "identity_layer3" to contain
the highest granularity of cell identity for neurons and non-neuronal cell types. Next, we
normalized the count data as specified for our murine dataset, and mapped the Ludwig dataset
on the reference latent space using scArches. We then used the learn_tree function with default
parameters for hierarchical progressive learning from scHPL v1.0.5(*41*) which yielded a
hierarchy of harmonized cell labels from both datasets. We printed all trees using matplotlib
v3.8.2(*77*) and scHPL v1.0.5(*41*). All steps performed and scripts used in this phase are
available in the GitHub repository: https://github.com/LabSabatini/DVC_cell_atlas.

Corroboration of the validity of our murine cell hierarchy using treeArches prediction modality
Using treeArches(*40*) and its dependencies in Python v3.9(*75*) we compared the original cell
labels in our and Ludwig datasets with the resulting prediction of the label for each cell using
our learned hierarchy. We used the predict_labels function from scHPL v1.0.5(*41*) with default
parameters, and the latent representation obtained when we constructed our murine hierarchy.
We compared the original and the predicted labels through a heatmap visualization. Next, we
predicted the labels of new data, a murine dataset by Dowsett which comprises the NTS, a
part of the DVC(*18*). We processed the Dowsett data similarly to the Ludwig dataset and
obtained the query latent representation. We then used the predict_labels function from scHPL
v1.0.5(*41*) with default parameters to transfer the labels in our hierarchy to the Dowsett dataset
based on gene expression similarity among the 2,500 initial most variable genes. We then
compared the resulting labels from this prediction to the manual mapping of our cell identities
based on maker expression (**Table S3**) using UMAP embeddings in R(*62*).

Labeling of our rat DVC dataset
We processed our rat dataset as previously described to obtain high-quality singlets. Based
on the three layers of cell identity that we created to label the murine data from our experiments,
we calculated the average expression of the top 5 marker genes for each of the layer-3

identities (except duplicated genes and mouse-specific genes) on every rat cluster to identify
those that were equivalent and could be labeled as the murine data (**Tables S5** and **S6**). Since
this dataset is smaller than the mouse data, we decided to use clusters at resolution of 2.0 to
capture more heterogeneity and delineate better the non-neuronal cell populations. Clusters
27 and 29 were further subset and reclustered since we found them to contain a mix of
endothelial cells, mural cells, tanycytes and fibroblasts based on marker expression. The
smallest resolution that allowed to separate these populations was used to label the cells.
Neurons were also subset and we mapped the murine neuronal classes markers to resolution
2.0 markers which were manually curated to delineate better the different populations within
them. For all cell identities, and because we only applied one treatment to the rats (i.e. fed ad
libitum), we used the FindMarkers function from Seurat v5(*63*) package in R(*62*) to obtain gene
expression markers of each rat cell population. The function was run using the MAST
algorithm(*71*) with same parameters used for the murine data (**Table S7**).
595

<u>Construction of the rodent DVC hierarchy</u>
We then integrated our rat labeled dataset to the murine samples in Python v3.9(*75*) using the
initial top 2,500 most variable genes initially defined in our mouse data. Incorporation of rat
labels was done as previously described for the Ludwig dataset using the learn_tree function
for hierarchical progressive learning from scHPL v1.0.5(*41*). The cell populations that were not
rejected but placed at the bottom of the tree were moved manually to their belonging parent
branches (**Fig.8A**; **Fig.S10**). For example, the 'smooth_muscle_rat' label was moved to be a
children node of the mural cells node (**Fig.8A**). If the parent branch did not exist, we created
it. For example, the 'immunity_related_rat' label belonged to a new rat neuronal class, therefore,
we incorporated this parent node within the neurons node and then we moved the
'immunity_related_rat' label to be a children node (**Fig.8A**). After manual curation, we re-trained
the tree. We printed all trees using matplotlib v3.8.2(*77*) and scHPL v1.0.5(*41*). All steps
performed and scripts used in this phase are available at the GitHub repository
https://github.com/LabSabatini/DVC_cell_atlas for reproducibility.
610

<u>Mapping gene expression in other brain-site snRNA-seq datasets</u>
We downloaded labeled datasets from two brain regions different from the DVC: the
HypoMap(*29*) and the cortex/hippocampus databases(*33*). For the HypoMap data, we subset
the database by randomly selecting cells belonging to 32 included hypothalamic samples. We
processed the datasets in Seurat v5(*63*) as previously described and obtained Harmony(*65*)
embeddings to obtain UMAP projections. The labels used in the visualization are the original
curated labels provided on each dataset. Neurons were further subset from the processed
datasets and reprocessed as previously described. The cortex/hippocampus dataset was

processed in the same manner, except that since it mainly contained astrocytes, these were
subset.

Differential gene expression analysis in murine treatments
We subset the murine dataset neurons and glial cells at layer 2 granularity of identity and
obtained differential gene expression for pair-wise comparison of treatments (i.e. refed versus
ad libitum fed, refed versus fasted and LiCl versus vehicle injected mice). We used the
FindMarkers function from Seurat v5(*63*) package in R(*62*) using the MAST algorithm(*71*) on
log-normalized counts with a minimal log fold-change threshold of 0.1 and default parameters.
The number of cells per treatment can be found in **Table S13**.

Immunofluorescent staining
Free-floating brain sections from mouse and rat were washed with PBS, three times for five
minutes. The sections were then blocked for one hour in PBS containing 0.1% Triton X-100 and
3% normal donkey serum (Fisher Scientific). The sections were incubated overnight at room
temperature with Goat anti-PDGFRA (Invitrogen, CAT#, 1:200), HuC/HuD (Invitrogen, A-21271,
1:30). The following day, sections were washed and incubated for two hours with Alexflour488
and Cy3 conjugated secondary antibody (Jackson immunoresearch, 1:500). Tissues were then
washed three times in PBS before being mounted onto glass slides, covered in mounting media
(Fluoromount-G, Southern Biotechnology) and coverslipped. Imaging was performed using an
Olympus BX61 microscope and a Zeiss LSM780-NLO laser scanning confocal microscope
equipped with IR-OPO lasers at the Molecular Imaging Platform, Research Institute of the
McGill University Health Centre (RI-MUHC), Montreal, CA.

In-situ hybridization and imaging
Brain sections containing the DVC from four wild-type C57BL/6J mice obtained as previously
described, were fixed on glass slides and stored at -20°C for no more than 36 hours. We
followed the manufacturer's ACD 323100 user manual for the RNAscope® Multiplex
Fluorescent Reagent Kit v2 Assay for fixed frozen tissue samples. Briefly, we rinsed the slides
with 1X PBS and incubated them at room temperature (RT) for 10 minutes after adding ~1-2
drops of H_2O_2 per section. We removed the H_2O_2 from slides and rinsed them twice with distilled
water (diH_2O). Next, we submerged the slides in 200 mL of hot diH_2O for 10 seconds followed
by 200mL of RNAscope® 1X Target Retrieval Reagent for 5 minutes, using a steamer with lid.
After briefly transferring the slides to RT diH_2O, we submerged them in 100% ethanol for 3
minutes and allowed them to air dry. We applied approximately 1-2 drops of RNAscope®
Protease III to each section and incubated them at 40°C for 30 minutes in a HybEZ™ oven with

diH$_2$O wet paper in the tray. After rinsing the slides with diH$_2$O, we hybridized the probes at
40°C for 2 hours.

From this point on, we performed all steps by incubating the slides at 40°C in the HybEZ™
oven with humid tray followed by rinse using RNAscope® 1X Wash Buffer. We applied
approximately 1-2 drops of RNAscope® reagents AMP1, AMP2, AMP3 and then intercalated
HRP channel (15 minutes), fluorophore (30 minutes) and HRP blocker (15 minutes) for each
channel present in the probe mix. The mix of probes used for the *Th*/*Cck* co-expression assay
was RNAscope® Mm-Th-C2 (317621-C2) and Mm-Cck-C3 (402271-C3) diluted in probe
diluent as specified in the user manual. For *Hcrt* we used RNAscope® Mm-Hcrt-C2 (490461-
C2) and for *Agrp*, Mm-Agrp (400711), in a dilution as specified in the user manual. The used
fluorophores were Cy3 for *Cck* and *Hcrt* (1:1,500), and Cy5 for *Th* and *Agrp* (1:1,000) diluted
in RNAscope® TSA buffer. DAPI dye (1:1,000) was applied at the end of the assay, and the
samples were stored for ~48 hours at -4°C, protected from light, before imaging. We imaged
the sections using a Zeiss LSM780-NLO laser scanning confocal microscope with IR-OPO
lasers at the Molecular Imaging Platform at the RI-MUHC, Montreal, CA.

References
1.   L. Gautron, J. K. Elmquist, K. W. Williams, Neural control of energy balance: translating
circuits to therapies. *Cell* **161**, 133-145 (2015).
2.   M. Q. Ludwig, W. Cheng, D. Gordian, J. Lee, S. J. Paulsen, S. N. Hansen, K. L.
Egerod, P. Barkholt, C. J. Rhodes, A. Secher, L. B. Knudsen, C. Pyke, M. G. Myers,
Jr., T. H. Pers, A genetic map of the mouse dorsal vagal complex and its role in
obesity. *Nat Metab* **3**, 530-545 (2021).
3.   L. Bai, S. Mesgarzadeh, K. S. Ramesh, E. L. Huey, Y. Liu, L. A. Gray, T. J. Aitken, Y.
Chen, L. R. Beutler, J. S. Ahn, L. Madisen, H. Zeng, M. A. Krasnow, Z. A. Knight,
Genetic Identification of Vagal Sensory Neurons That Control Feeding. *Cell* **179**, 1129-
1143 e1123 (2019).
4.   W. Han, L. A. Tellez, M. H. Perkins, I. O. Perez, T. Qu, J. Ferreira, T. L. Ferreira, D.
Quinn, Z. W. Liu, X. B. Gao, M. M. Kaelberer, D. V. Bohorquez, S. J. Shammah-
Lagnado, G. de Lartigue, I. E. de Araujo, A Neural Circuit for Gut-Induced Reward.
Cell **175**, 665-678 e623 (2018).
5.   E. K. Williams, R. B. Chang, D. E. Strochlic, B. D. Umans, B. B. Lowell, S. D. Liberles,
Sensory Neurons that Detect Stretch and Nutrients in the Digestive System. *Cell* **166**,
209-221 (2016).
6.   M. W. Schwartz, Central nervous system regulation of food intake. *Obesity (Silver*
Spring) **14 Suppl 1**, 1S-8S (2006).
7.   W. Cheng, E. Ndoka, J. N. Maung, W. Pan, A. C. Rupp, C. J. Rhodes, D. P. Olson, M.
G. Myers, Jr., NTS Prlh overcomes orexigenic stimuli and ameliorates dietary and
genetic forms of obesity. *Nat Commun* **12**, 5175 (2021).
8.   W. Qiu, C. R. Hutch, Y. Wang, J. Wloszek, R. A. Rucker, M. G. Myers, D. Sandoval,
Multiple NTS neuron populations cumulatively suppress food intake. *Elife* **12**, (2023).
9.   T. Borner, C. G. Liberini, T. A. Lutz, T. Riediger, Brainstem GLP-1 signalling
contributes to cancer anorexia-cachexia syndrome in the rat. *Neuropharmacology*
131, 282-290 (2018).

10. C. W. le Roux, O. Steen, K. J. Lucas, E. Startseva, A. Unseld, A. M. Hennige, Glucagon and GLP-1 receptor dual agonist survodutide for obesity: a randomised, double-blind, placebo-controlled, dose-finding phase 2 trial. *Lancet Diabetes Endocrinol* **12**, 162-173 (2024).

11. S. M. Advani, P. G. Advani, H. M. VonVille, S. H. Jafri, Pharmacological management of cachexia in adult cancer patients: a systematic review of clinical trials. *BMC Cancer* **18**, 1174 (2018).

12. M. M. Veniant, S. C. Lu, L. Atangan, R. Komorowski, S. Stanislaus, Y. Cheng, B. Wu, J. R. Falsey, T. Hager, V. A. Thomas, M. Ambhaikar, L. Sharpsten, Y. Zhu, V. Kurra, R. Jeswani, R. K. Oberoi, J. R. Parnes, N. Honarpour, J. Neutel, J. L. Strande, A GIPR antagonist conjugated to GLP-1 analogues promotes weight loss with improved metabolic parameters in preclinical and phase 1 settings. *Nat Metab* **6**, 290-303 (2024).

13. J. M. Adams, H. Pei, D. A. Sandoval, R. J. Seeley, R. B. Chang, S. D. Liberles, D. P. Olson, Liraglutide Modulates Appetite and Body Weight Through Glucagon-Like Peptide 1 Receptor-Expressing Glutamatergic Neurons. *Diabetes* **67**, 1538-1548 (2018).

14. K. P. Huang, A. A. Acosta, M. Y. Ghidewon, A. D. McKnight, M. S. Almeida, N. T. Nyema, N. D. Hanchak, N. Patel, Y. S. K. Gbenou, A. E. Adriaenssens, K. A. Bolding, A. L. Alhadeff, Dissociable hindbrain GLP1R circuits for satiety and aversion. *Nature* **632**, 585-593 (2024).

15. I. Medic, S. Gulati, H.-J. Lenz, D. Mahalingam, J. Thomas, J. Luo, J. Zha, A. DePaoli, C. Tran-Muchowski, C. Tseng, H. Lieu, Initial results of a cohort of advanced prostate cancer patients in a phase Ia study of NGM120, a first-in-class anti- GDNF family receptor alpha like (GFRAL) antibody. *Annals of Oncology* **33**, S1186-S1186 (2022).

16. L. Yang, C. C. Chang, Z. Sun, D. Madsen, H. Zhu, S. B. Padkjaer, X. Wu, T. Huang, K. Hultman, S. J. Paulsen, J. Wang, A. Bugge, J. B. Frantzen, P. Norgaard, J. F. Jeppesen, Z. Yang, A. Secher, H. Chen, X. Li, L. M. John, B. Shan, Z. He, X. Gao, J. Su, K. T. Hansen, W. Yang, S. B. Jorgensen, GFRAL is the receptor for GDF15 and is required for the anti-obesity effects of the ligand. *Nat Med* **23**, 1158-1166 (2017).

17. C. W. Roman, S. R. Sloat, R. D. Palmiter, A tale of two circuits: CCK(NTS) neuron stimulation controls appetite and induces opposing motivational states by projections to distinct brain regions. *Neuroscience* **358**, 316-324 (2017).

18. G. K. C. Dowsett, B. Y. H. Lam, J. A. Tadross, I. Cimino, D. Rimmington, A. P. Coll, J. Polex-Wolf, L. B. Knudsen, C. Pyke, G. S. H. Yeo, A survey of the mouse hindbrain in the fed and fasted states using single-nucleus RNA sequencing. *Mol Metab* **53**, 101240 (2021).

19. C. Zhang, J. A. Kaye, Z. Cai, Y. Wang, S. L. Prescott, S. D. Liberles, Area Postrema Cell Types that Mediate Nausea-Associated Behaviors. *Neuron* **109**, 461-472 e465 (2021).

20. S. M. Fortin, R. K. Lipsky, R. Lhamo, J. Chen, E. Kim, T. Borner, H. D. Schmidt, M. R. Hayes, GABA neurons in the nucleus tractus solitarius express GLP-1 receptors and mediate anorectic effects of liraglutide in rats. *Sci Transl Med* **12**, (2020).

21. G. X. Zheng, J. M. Terry, P. Belgrader, P. Ryvkin, Z. W. Bent, R. Wilson, S. B. Ziraldo, T. D. Wheeler, G. P. McDermott, J. Zhu, M. T. Gregory, J. Shuga, L. Montesclaros, J. G. Underwood, D. A. Masquelier, S. Y. Nishimura, M. Schnall-Levin, P. W. Wyatt, C. M. Hindson, R. Bharadwaj, A. Wong, K. D. Ness, L. W. Beppu, H. J. Deeg, C. McFarland, K. R. Loeb, W. J. Valente, N. G. Ericson, E. A. Stevens, J. P. Radich, T. S. Mikkelsen, B. J. Hindson, J. H. Bielas, Massively parallel digital transcriptional profiling of single cells. *Nat Commun* **8**, 14049 (2017).

22. C. E. Vaaga, M. Borisovska, G. L. Westbrook, Dual-transmitter neurons: functional implications of co-release and co-transmission. *Curr Opin Neurobiol* **29**, 25-32 (2014).

23. N. X. Tritsch, A. J. Granger, B. L. Sabatini, Mechanisms and functions of GABA co-release. *Nat Rev Neurosci* **17**, 139-145 (2016).

24. C. F. Landry, J. A. Ellison, T. M. Pribyl, C. Campagnoni, K. Kampf, A. T. Campagnoni, Myelin basic protein gene expression in neurons: developmental and regional

changes in protein targeting within neuronal nuclei, cell bodies, and processes. *J.*
Neurosci. **16**, 2452-2462 (1996).
25.   D. Lutz, G. Loers, R. Kleene, I. Oezen, H. Kataria, N. Katagihallimath, I. Braren, G.
Harauz, M. Schachner, Myelin basic protein cleaves cell adhesion molecule L1 and
promotes neuritogenesis and cell survival. *J Biol Chem* **289**, 13503-13518 (2014).
26.   V. T. Cheli, D. A. Santiago Gonzalez, V. Spreuer, V. Handley, A. T. Campagnoni, P. M.
Paez, Golli Myelin Basic Proteins Modulate Voltage-Operated Ca(++) Influx and
Development in Cortical and Hippocampal Neurons. *Mol Neurobiol* **53**, 5749-5771
(2016).
27.   M. Melone, S. Ciappelloni, F. Conti, Plasma membrane transporters GAT-1 and GAT-3
contribute to heterogeneity of GABAergic synapses in neocortex. *Front Neuroanat* **8**,
72 (2014).
28.   S. M. Theparambil, P. S. Hosford, I. Ruminot, O. Kopach, J. R. Reynolds, P. Y.
Sandoval, D. A. Rusakov, L. F. Barros, A. V. Gourine, Astrocytes regulate brain
extracellular pH via a neuronal activity-dependent bicarbonate shuttle. *Nat Commun*
11, 5073 (2020).
29.   L. Steuernagel, B. Y. H. Lam, P. Klemm, G. K. C. Dowsett, C. A. Bauder, J. A. Tadross,
T. S. Hitschfeld, A. Del Rio Martin, W. Chen, A. J. de Solis, H. Fenselau, P. Davidsen, I.
Cimino, S. N. Kohnke, D. Rimmington, A. P. Coll, A. Beyer, G. S. H. Yeo, J. C. Bruning,
HypoMap-a unified single-cell gene expression atlas of the murine hypothalamus. *Nat*
Metab **4**, 1402-1419 (2022).
30.   Y. Kubo, T. J. Baldwin, Y. N. Jan, L. Y. Jan, Primary structure and functional
expression of a mouse inward rectifier potassium channel. *Nature* **362**, 127–133
(1993).
31.   N. Burnashev, A. Khodorova, P. Jonas, P. J. Helm, W. Wisden, H. Monyer, P. H.
Seeburg, B. Sakmann, Calcium-Permeable AMPA-Kainate Receptors in Fusiform
Cerebellar Glial Cells. *Science* **256**, 1566-1570 (1992).
32.   J. R. P. Geiger, T. Melcher, D. S. Koh, B. Sakmann, S. P. H., P. Jonas, H. Monyer,
Relative Abundance of Subunit mRNAs Determines Gating and Ca*+ Permeability of
AMPA Receptors in Principal Neurons and Interneurons in Rat CNS. *Neuron* **15**, 193-
204 (1995).
33.   M. Y. Batiuk, A. Martirosyan, J. Wahis, F. de Vin, C. Marneffe, C. Kusserow, J.
Koeppen, J. F. Viana, J. F. Oliveira, T. Voet, C. P. Ponting, T. G. Belgard, M. G. Holt,
Identification of region-specific astrocyte subtypes at single cell resolution. *Nat*
Commun **11**, 1220 (2020).
34.   S. Marques, A. Zeisel, S. Codeluppi, D. van Bruggen, A. Mendanha Falcao, L. Xiao, H.
Li, M. Haring, H. Hochgerner, R. A. Romanov, D. Gyllborg, A. Munoz Manchado, G. La
Manno, P. Lonnerberg, E. M. Floriddia, F. Rezayee, P. Ernfors, E. Arenas, J. Hjerling-
Leffler, T. Harkany, W. D. Richardson, S. Linnarsson, G. Castelo-Branco,
Oligodendrocyte heterogeneity in the mouse juvenile and adult central nervous
system. *Science* **352**, 1326-1329 (2016).
35.   M. Kasai, Y. Omae, Y. Kawai, A. Shibata, A. Hoshino, M. Mizuguchi, K. Tokunaga,
GWAS identifies candidate susceptibility loci and microRNA biomarkers for acute
encephalopathy with biphasic seizures and late reduced diffusion. *Sci Rep* **12**, 1332
(2022).
36.   A. B. DePaula-Silva, C. Gorbea, D. J. Doty, J. E. Libbey, J. M. S. Sanchez, T. J. Hanak,
D. Cazalla, R. S. Fujinami, Differential transcriptional profiles identify microglial- and
macrophage-specific gene markers expressed during virus-induced
neuroinflammation. *J Neuroinflammation* **16**, 152 (2019).
37.   J. N. Campbell, E. Z. Macosko, H. Fenselau, T. H. Pers, A. Lyubetskaya, D. Tenen, M.
Goldman, A. M. Verstegen, J. M. Resch, S. A. McCarroll, E. D. Rosen, B. B. Lowell, L.
T. Tsai, A molecular census of arcuate hypothalamus and median eminence cell
types. *Nat Neurosci* **20**, 484-496 (2017).
38.   W. Cheng, E. Ndoka, C. Hutch, K. Roelofs, A. MacKinnon, B. Khoury, J. Magrisso, K.
S. Kim, C. J. Rhodes, D. P. Olson, R. J. Seeley, D. Sandoval, M. G. Myers, Jr., Leptin
receptor-expressing nucleus tractus solitarius neurons suppress food intake
independently of GLP1 in mice. *JCI Insight* **5**, (2020).

39.   A. A. Worth, R. Shoop, K. Tye, C. H. Feetham, G. D'Agostino, G. T. Dodd, F. Reimann,
F. M. Gribble, E. C. Beebe, J. D. Dunbar, J. T. Alexander-Chacko, D. K. Sindelar, T.
Coskun, P. J. Emmerson, S. M. Luckman, The cytokine GDF15 signals through a
population of brainstem cholecystokinin neurons to mediate anorectic signalling. *Elife*
9,   (2020).
40.   L. Michielsen, M. Lotfollahi, D. Strobl, L. Sikkema, M. J. T. Reinders, F. J. Theis, A.
Mahfouz, Single-cell reference mapping to construct and extend cell-type hierarchies.
NAR Genom Bioinform **5**, lqad070 (2023).
41.   L. Michielsen, M. J. T. Reinders, A. Mahfouz, Hierarchical progressive learning of cell
identities in single-cell data. *Nat Commun* **12**, 2799 (2021).
42.   V. Chou, R. V. Pearse, 2nd, A. J. Aylward, N. Ashour, M. Taga, G. Terzioglu, M. Fujita,
S. B. Fancher, A. Sigalov, C. R. Benoit, H. Lee, M. Lam, N. T. Seyfried, D. A. Bennett,
P. L. De Jager, V. Menon, T. L. Young-Pearse, INPP5D regulates inflammasome
activation in human microglia. *Nat Commun* **14**, 7552 (2023).
43.   S. O. Kleeman, T. M. Thakir, B. Demestichas, N. Mourikis, D. Loiero, M. Ferrer, S.
Bankier, Y. Riazat-Kesh, H. Lee, D. Chantzichristos, C. Regan, J. Preall, S. Sinha, N.
Rosin, B. Yipp, L. G. N. de Almeida, J. Biernaskie, A. Dufour, P. Tober-Lau, A.
Ruusalepp, J. L. M. Bjorkegren, M. Ralser, F. Kurth, V. Demichev, T. Heywood, Q.
Gao, G. Johannsson, V. H. Koelzer, B. R. Walker, H. V. Meyer, T. Janowitz, Cystatin C
is glucocorticoid responsive, directs recruitment of Trem2+ macrophages, and
predicts failure of cancer immunotherapy. *Cell Genom* **3**, 100347 (2023).
44.   N. Hagan, J. L. Kane, D. Grover, L. Woodworth, C. Madore, J. Saleh, J. Sancho, J. Liu,
Y. Li, J. Proto, M. Zelic, A. Mahan, M. Kothe, A. A. Scholte, M. Fitzgerald, B. Gisevius,
A. Haghikia, O. Butovsky, D. Ofengeim, CSF1R signaling is a regulator of
pathogenesis in progressive MS. *Cell Death Dis* **11**, 904 (2020).
45.   S. H. Kang, M. Fukaya, J. K. Yang, J. D. Rothstein, D. E. Bergles, NG2+ CNS glial
progenitors remain committed to the oligodendrocyte lineage in postnatal life and
following neurodegeneration. *Neuron* **68**, 668-681 (2010).
46.   T. P. Bachor, E. Hwang, E. Yulyaningsih, K. Attal, F. Mifsud, V. Pham, E. Vagena, R.
Huarcaya, M. Valdearcos, C. Vaisse, K. W. Williams, P. J. Emmerson, A. W. Xu,
Identification of AgRP cells in the murine hindbrain that drive feeding. *Mol Metab* **80**,
101886 (2024).
47.   M. K. Holt, The ins and outs of the caudal nucleus of the solitary tract: An overview of
cellular populations and anatomical connections. *J Neuroendocrinol* **34**, e13132
(2022).
48.   D. M. Breen, H. Kim, D. Bennett, R. A. Calle, S. Collins, R. M. Esquejo, T. He, S.
Joaquim, A. Joyce, M. Lambert, L. Lin, B. Pettersen, S. Qiao, M. Rossulek, G. Weber,
Z. Wu, B. B. Zhang, M. J. Birnbaum, GDF-15 Neutralization Alleviates Platinum-Based
Chemotherapy-Induced Emesis, Anorexia, and Weight Loss in Mice and Nonhuman
Primates. *Cell Metab* **32**, 938-950 e936 (2020).
49.   K. Tsuzuki, S. Ishiuchi, GluR2 expressed by glial fibrillary acidic protein promoter
decreases the number of neurons. *Front. Biosci. (Landmark Ed)* **13**, 2784–2796
(2008).
50.   W. Cheng, I. Gonzalez, W. Pan, A. H. Tsang, J. Adams, E. Ndoka, D. Gordian, B.
Khoury, K. Roelofs, S. S. Evers, A. MacKinnon, S. Wu, H. Frikke-Schmidt, J. N. Flak, J.
L. Trevaskis, C. J. Rhodes, S. I. Fukada, R. J. Seeley, D. A. Sandoval, D. P. Olson, C.
Blouet, M. G. Myers, Jr., Calcitonin Receptor Neurons in the Mouse Nucleus Tractus
Solitarius Control Energy Balance via the Non-aversive Suppression of Feeding. *Cell*
Metab **31**, 301-312 e305 (2020).
51.   A. R. Patel, H. Frikke-Schmidt, P. V. Sabatini, A. C. Rupp, D. A. Sandoval, M. G. Myers,
Jr., R. J. Seeley, Neither GLP-1 receptors nor GFRAL neurons are required for aversive
or anorectic response to DON (vomitoxin). *Am J Physiol Regul Integr Comp Physiol*
324, R635-644 (2023).
52.   D. H. Root, C. A. Mejias-Aponte, S. Zhang, H. L. Wang, A. F. Hoffman, C. R. Lupica, M.
Morales, Single rodent mesohabenular axons release glutamate and GABA. *Nat*
Neurosci **17**, 1543-1551 (2014).

53. A. Saunders, A. J. Granger, B. L. Sabatini, Corelease of acetylcholine and GABA from
cholinergic forebrain neurons. *Elife* **4**, (2015).
54. S. C. Robins, E. Trudel, O. Rotondi, X. Liu, T. Djogo, D. Kryzskaya, C. W. Bourque, M.
V. Kokoeva, Evidence for NG2-glia derived, adult-born functional neurons in the
hypothalamus. *PLoS One* **8**, e78236 (2013).
55. N. Wada, S. Hirako, F. Takenoya, H. Kageyama, M. Okabe, S. Shioda, Leptin and its
receptors. *J Chem Neuroanat* **61-62**, 191-199 (2014).
56. H. Munzberg, J. S. Flier, C. Bjorbaek, Region-specific leptin resistance within the
hypothalamus of diet-induced obese mice. *Endocrinology* **145**, 4880-4889 (2004).
57. P. M. Smith, P. Brzezinska, F. Hubert, A. Mimee, D. H. Maurice, A. V. Ferguson, Leptin
influences the excitability of area postrema neurons. *Am J Physiol Regul Integr Comp*
Physiol **310**, R440-448 (2016).
58. C. G. Liberini, C. N. Boyle, C. Cifani, M. Venniro, B. T. Hope, T. A. Lutz, Amylin
receptor components and the leptin receptor are co-expressed in single rat area
postrema neurons. *Eur J Neurosci* **43**, 653-661 (2016).
59. L. Huo, L. Maeng, C. Bjorbaek, H. J. Grill, Leptin and the control of food intake:
neurons in the nucleus of the solitary tract are activated by both gastric distension and
leptin. *Endocrinology* **148**, 2189-2197 (2007).
60. S. R. Smith, L. J. Aronne, C. M. Burns, N. C. Kesty, A. E. Halseth, C. Weyer, Sustained
weight loss following 12-month pramlintide treatment as an adjunct to lifestyle
intervention in obesity. *Diabetes Care* **31**, 1816-1823 (2008).
61. J. P. H. Wilding, R. L. Batterham, S. Calanna, M. Davies, L. F. Van Gaal, I. Lingvay, B.
M. McGowan, J. Rosenstock, M. T. D. Tran, T. A. Wadden, S. Wharton, K. Yokote, N.
Zeuthen, R. F. Kushner, S. S. Group, Once-Weekly Semaglutide in Adults with
Overweight or Obesity. *N Engl J Med* **384**, 989-1002 (2021).
62. R. C. Team. (R Foundation for Statistical Computing, Vienna, Austria, 2023).
63. Y. Hao, T. Stuart, M. H. Kowalski, S. Choudhary, P. Hoffman, A. Hartman, A.
Srivastava, G. Molla, S. Madad, C. Fernandez-Granda, R. Satija, Dictionary learning for
integrative, multimodal and scalable single-cell analysis. *Nat Biotechnol* **42**, 293-304
(2024).
64. C. S. McGinnis, L. M. Murrow, Z. J. Gartner, DoubletFinder: Doublet Detection in
Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-
337 e324 (2019).
65. I. Korsunsky, N. Millard, J. Fan, K. Slowikowski, F. Zhang, K. Wei, Y. Baglaenko, M.
Brenner, P. R. Loh, S. Raychaudhuri, Fast, sensitive and accurate integration of single-
cell data with Harmony. *Nat Methods* **16**, 1289-1296 (2019).
66. R. Chen, X. Wu, L. Jiang, Y. Zhang, Single-Cell RNA-Seq Reveals Hypothalamic Cell
Diversity. *Cell Rep* **18**, 3227-3241 (2017).
67. R. A. Romanov, A. Zeisel, J. Bakker, F. Girach, A. Hellysaz, R. Tomer, A. Alpar, J.
Mulder, F. Clotman, E. Keimpema, B. Hsueh, A. K. Crow, H. Martens, C. Schwindling,
D. Calvigioni, J. S. Bains, Z. Mate, G. Szabo, Y. Yanagawa, M. D. Zhang, A. Rendeiro,
M. Farlik, M. Uhlen, P. Wulff, C. Bock, C. Broberger, K. Deisseroth, T. Hokfelt, S.
Linnarsson, T. L. Horvath, T. Harkany, Molecular interrogation of hypothalamic
organization reveals distinct dopamine neuronal subtypes. *Nat Neurosci* **20**, 176-188
(2017).
68. B. Tasic, V. Menon, T. N. Nguyen, T. K. Kim, T. Jarsky, Z. Yao, B. Levi, L. T. Gray, S. A.
Sorensen, T. Dolbeare, D. Bertagnolli, J. Goldy, N. Shapovalova, S. Parry, C. Lee, K.
Smith, A. Bernard, L. Madisen, S. M. Sunkin, M. Hawrylycz, C. Koch, H. Zeng, Adult
mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat Neurosci* **19**,
335-346 (2016).
69. D. Aran, A. P. Looney, L. Liu, E. Wu, V. Fong, A. Hsu, S. Chak, R. P. Naikawadi, P. J.
Wolters, A. R. Abate, A. J. Butte, M. Bhattacharya, Reference-based analysis of lung
single-cell sequencing reveals a transitional profibrotic macrophage. *Nat Immunol* **20**,
163-172 (2019).
70. D. Risso, M. Cole. (2023).
71. G. Finak, A. McDavid, M. Yajima, J. Deng, V. Gersuk, A. K. Shalek, C. K. Slichter, H.
W. Miller, M. J. McElrath, M. Prlic, P. S. Linsley, R. Gottardo, MAST: a flexible statistical

framework for assessing transcriptional changes and characterizing heterogeneity in
single-cell RNA sequencing data. *Genome Biol* **16**, 278 (2015).
72.   R. Cannoodt, W. Saelens, D. Sichien, S. Tavernier, S. Janssens, M. Guilliams, B.
Lambrecht, K. De Preter, Y. Saeys, SCORPIUS improves trajectory inference and
identifies novel modules in dendritic cell development. *Preprint at bioRxiv*, (2016).
73.   K. Ushey, J. Allaire, Y. Tang. (2024).
74.   R. Cannoodt. (2023).
75.   G. Van Rossum, F. L. Drake. (CreateSpace, Scotts Valley, CA, 2009).
76.   A. Gayoso, R. Lopez, G. Xing, P. Boyeau, V. Valiollah Pour Amiri, J. Hong, K. Wu, M.
Jayasuriya, E. Mehlman, M. Langevin, Y. Liu, J. Samaran, G. Misrachi, A. Nazaret, O.
Clivio, C. Xu, T. Ashuach, M. Gabitto, M. Lotfollahi, V. Svensson, E. da Veiga Beltrame,
V. Kleshchevnikov, C. Talavera-Lopez, L. Pachter, F. J. Theis, A. Streets, M. I. Jordan,
J. Regier, N. Yosef, A Python library for probabilistic analysis of single-cell omics data.
Nat Biotechnol **40**, 163-166 (2022).
77.   J. D. Hunter, Matplotlib: A 2D graphics environment. *Computing in Science &*
Engineering **3**, 90-95 (2007).
78.   H. Thorvaldsdottir, J. T. Robinson, J. P. Mesirov, Integrative Genomics Viewer (IGV):
high-performance genomics data visualization and exploration. *Brief Bioinform* **14**,
178-192 (2013).
79.   M. J. Park, K. Chung, Endogenous lectin (RL-29) expression of the autonomic
preganglionic neurons in the rat spinal cord. The Anatomical Record 254, 54-60
(1999).
80.   C. O. Miranda, K. Hegedus, H. Wildner, H. U. Zeilhofer, M. Antal, Morphological and
neurochemical characterization of glycinergic neurons in laminae I-IV of the mouse
spinal dorsal horn. J Comp Neurol 530, 607-626 (2022).
81.   F. Guo, Y. Wang, TCF7l2, a nuclear marker that labels premyelinating
oligodendrocytes and promotes oligodendroglial lineage progression. Glia 71, 143-
154 (2023).
82.   E. Weihe, C. Depboylu, B. Schutz, M. K. Schafer, L. E. Eiden, Three types of tyrosine
hydroxylase-positive CNS neurons distinguished by dopa decarboxylase and VMAT2
co-expression. Cell Mol Neurobiol 26, 659-678 (2006).
83.   J. Sun, Y. Song, Z. Chen, J. Qiu, S. Zhu, L. Wu, L. Xing, Heterogeneity and Molecular
Markers for CNS Glial Cells Revealed by Single-Cell Transcriptomics. Cell Mol
Neurobiol 42, 2629-2642 (2022).
84.   M. Dallaporta, M. S. Bonnet, K. Horner, J. Trouslard, A. Jean, J. D. Troadec, Glial cells
of the nucleus tractus solitarius as partners of the dorsal hindbrain regulation of
energy balance: a proposal for a working hypothesis. Brain Res 1350, 35-42 (2010).
85.   J. Kalucka, L. de Rooij, J. Goveia, K. Rohlenova, S. J. Dumas, E. Meta, N. V.
Conchinha, F. Taverna, L. A. Teuwen, K. Veys, M. Garcia-Caballero, S. Khan, V.
Geldhof, L. Sokol, R. Chen, L. Treps, M. Borri, P. de Zeeuw, C. Dubois, T. K.
Karakach, K. D. Falkenberg, M. Parys, X. Yin, S. Vinckier, Y. Du, R. A. Fenton, L.
Schoonjans, M. Dewerchin, G. Eelen, B. Thienpont, L. Lin, L. Bolund, X. Li, Y. Luo, P.
Carmeliet, Single-Cell Transcriptome Atlas of Murine Endothelial Cells. Cell 180, 764-
779 e720 (2020).
86.   N. Dani, R. H. Herbst, C. McCabe, G. S. Green, K. Kaiser, J. P. Head, J. Cui, F. B.
Shipley, A. Jang, D. Dionne, L. Nguyen, C. Rodman, S. J. Riesenfeld, J. Prochazka, M.
Prochazkova, R. Sedlacek, F. Zhang, V. Bryja, O. Rozenblatt-Rosen, N. Habib, A.
Regev, M. K. Lehtinen, A cellular and spatial map of the choroid plexus across brain
ventricles and ages. Cell 184, 3056-3074 e3021 (2021).
87.   B. Ray, J. A. Bailey, S. Sarkar, D. K. Lahiri, Molecular and immunocytochemical
characterization of primary neuronal cultures from adult rat brain: Differential
expression of neuronal and glial protein markers. J Neurosci Methods 184, 294-302
(2009).
88.   S. E. Lauri, M. Ryazantseva, E. Orav, A. Vesikansa, T. Taira, Kainate receptors in the
developing neuronal networks. Neuropharmacology 195, 108585 (2021).
89.   Y. H. Song, J. Yoon, S. H. Lee, The role of neuropeptide somatostatin in the brain and
its application in treating neurological disorders. Exp Mol Med 53, 328-338 (2021).

90.   H. W. Song, K. L. Foreman, B. D. Gastfriend, J. S. Kuo, S. P. Palecek, E. V. Shusta,
Transcriptomic comparison of human and mouse brain microvessels. Sci Rep 10,
12358 (2020).
91.   L. Xu, Y. Yao, Central Nervous System Fibroblast-Like Cells in Stroke and Other
Neurological Disorders. Stroke 52, 2456-2464 (2021).
92.   U. Ernsberger, H. Rohrer, Sympathetic tales: subdivisons of the autonomic nervous
system and the impact of developmental studies. Neural Dev 13, 20 (2018).
93.   C. R. Gajera, H. Emich, O. Lioubinski, A. Christ, R. Beckervordersandforth-Bonk, K.
Yoshikawa, S. Bachmann, E. I. Christensen, M. Gotz, G. Kempermann, A. S. Peterson,
T. E. Willnow, A. Hammes, LRP2 in ependymal cells regulates BMP signaling in the
adult neurogenic niche. J Cell Sci 123, 1922-1930 (2010).
94.   A. MacDonald, B. Lu, M. Caron, N. Caporicci-Dinucci, D. Hatrock, K. Petrecca, G.
Bourque, J. A. Stratton, Single Cell Transcriptomics of Ependymal Cells Across Age,
Region and Species Reveals Cilia-Related and Metal Ion Regulatory Roles as Major
Conserved Ependymal Cell Functions. Front Cell Neurosci 15, 703951 (2021).
95.   H. W. Jeong, R. Dieguez-Hurtado, H. Arf, J. Song, H. Park, K. Kruse, L. Sorokin, R. H.
Adams, Single-cell transcriptomics reveals functionally specialized vascular
endothelium in brain. Elife 11, (2022).
96.   K. Huizer, D. A. M. Mustafa, J. C. Spelt, J. M. Kros, A. Sacchetti, Improving the
characterization of endothelial progenitor cell subsets by an optimized FACS protocol.
PLoS One 12, e0184895 (2017).
97.   E. Gerrits, Y. Heng, E. Boddeke, B. J. L. Eggen, Transcriptional profiling of microglia;
current state of the art and future perspectives. Glia 68, 740-755 (2020).
98.   P. T. Shah, J. A. Stratton, M. G. Stykel, S. Abbasi, S. Sharma, K. A. Mayr, K. Koblinger,
P. J. Whelan, J. Biernaskie, Single-Cell Transcriptomics and Fate Mapping of
Ependymal Cells Reveals an Absence of Neural Stem Cell Function. Cell 173, 1045-
1057 e1049 (2018).
99.   C. G. Causing, A. Gloster, R. Aloyz, S. X. Bamji, E. Chang, J. Fawcett, G. Kuchel, F. D.
Miller, Synaptic Innervation Density Is Regulated by Neuron-Derived BDNF. Neuron
18, 257–267 (1997).
100. Y. Zhang, S. A. Sloan, L. E. Clarke, C. Caneda, C. A. Plaza, P. D. Blumenthal, H.
Vogel, G. K. Steinberg, M. S. Edwards, G. Li, J. A. Duncan, 3rd, S. H. Cheshier, L. M.
Shuer, E. F. Chang, G. A. Grant, M. G. Gephart, B. A. Barres, Purification and
Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional
and Functional Differences with Mouse. Neuron 89, 37-53 (2016).
101. F. Langlet, Tanycyte Gene Expression Dynamics in the Regulation of Energy
Homeostasis. Front Endocrinol (Lausanne) 10, 286 (2019).
102. O. Uriarte Huarte, D. Kyriakis, T. Heurtaux, Y. Pires-Afonso, K. Grzyb, R. Halder, M.
Buttini, A. Skupin, M. Mittelbronn, A. Michelucci, Single-Cell Transcriptomics and In
Situ Morphological Analyses Reveal Microglia Heterogeneity Across the Nigrostriatal
Pathway. Front Immunol 12, 639613 (2021).
103. R. E. Pepper, K. A. Pitman, C. L. Cullen, K. M. Young, How Do Cells of the
Oligodendrocyte Lineage Affect Neuronal Circuits to Influence Motor Function,
Memory and Mood? Front Cell Neurosci 12, 399 (2018).
104. A. Maegawa, K. Murata, K. Kuroda, S. Fujieda, Y. Fukazawa, Cellular Profiles of
Prodynorphin and Preproenkephalin mRNA-Expressing Neurons in the Anterior
Olfactory Tubercle of Mice. Front Neural Circuits 16, 908964 (2022).
105. S. T. Hentges, V. Otero-Corchon, R. L. Pennock, C. M. King, M. J. Low,
Proopiomelanocortin expression in both GABA and glutamate neurons. J Neurosci 29,
13684-13690 (2009).
106. W. Sun, A. Cornwell, J. Li, S. Peng, M. J. Osorio, N. Aalling, S. Wang, A. Benraiss, N.
Lou, S. A. Goldman, M. Nedergaard, SOX9 Is an Astrocyte-Specific Nuclear Marker in
the Adult Brain Outside the Neurogenic Regions. J Neurosci 37, 4493-4507 (2017).
107. A. T. Lee, D. Vogt, J. L. Rubenstein, V. S. Sohal, A class of GABAergic neurons in the
prefrontal cortex sends long-range projections to the nucleus accumbens and elicits
acute avoidance behavior. J Neurosci 34, 11519-11525 (2014).

108. M. Vanlandewijck, L. He, M. A. Mae, J. Andrae, K. Ando, F. Del Gaudio, K. Nahar, T.
Lebouvier, B. Lavina, L. Gouveia, Y. Sun, E. Raschperger, M. Rasanen, Y. Zarb, N.
Mochizuki, A. Keller, U. Lendahl, C. Betsholtz, A molecular atlas of cell types and
zonation in the brain vasculature. Nature 554, 475-480 (2018).
109. M. Carlen, K. Meletis, C. Goritz, V. Darsalia, E. Evergren, K. Tanigaki, M. Amendola, F.
Barnabe-Heider, M. S. Yeung, L. Naldini, T. Honjo, Z. Kokaia, O. Shupliakov, R. M.
Cassidy, O. Lindvall, J. Frisen, Forebrain ependymal cells are Notch-dependent and
generate neuroblasts and astrocytes after stroke. Nat Neurosci 12, 259-267 (2009).
110. J. Garcia-Revilla, A. Boza-Serrano, A. M. Espinosa-Oliva, M. S. Soto, T. Deierborg, R.
Ruiz, R. M. de Pablos, M. A. Burguillos, J. L. Venero, Galectin-3, a rising star in
modulating microglia activation under conditions of neurodegeneration. Cell Death Dis
13, 628 (2022).
111. A. C. Yang, R. T. Vest, F. Kern, D. P. Lee, M. Agam, C. A. Maat, P. M. Losada, M. B.
Chen, N. Schaum, N. Khoury, A. Toland, K. Calcuttawala, H. Shin, R. Palovics, A. Shin,
E. Y. Wang, J. Luo, D. Gate, W. J. Schulz-Schaeffer, P. Chu, J. A. Siegenthaler, M. W.
McNerney, A. Keller, T. Wyss-Coray, A human brain vascular atlas reveals diverse
mediators of Alzheimer's risk. Nature 603, 885-892 (2022).
112. S. Chasseigneaux, Y. Moraca, V. Cochois-Guegan, A. C. Boulay, A. Gilbert, S. Le
Crom, C. Blugeon, C. Firmo, S. Cisternino, J. L. Laplanche, E. Curis, X. Decleves, B.
Saubamea, Isolation and differential transcriptome of vascular smooth muscle cells
and mid-capillary pericytes from the rat brain. Sci Rep 8, 12272 (2018).
113. Z. Gao, K. Ure, J. L. Ables, D. C. Lagace, K. A. Nave, S. Goebbels, A. J. Eisch, J.
Hsieh, Neurod1 is essential for the survival and maturation of adult-born neurons. Nat
Neurosci 12, 1090-1092 (2009).
114. T. Kodama, S. Guerrero, M. Shin, S. Moghadam, M. Faulstich, S. du Lac, Neuronal
classification and marker gene identification via single-cell expression profiling of
brainstem vestibular neurons subserving cerebellar learning. J Neurosci 32, 7819-
7831 (2012).
115. J. Manjarrez-Marmolejo, J. Franco-Pérez, Gap Junction Blockers- An Overview of their
Effects on Induced Seizures in Animal Models. Current Neuropharmacology 14, 759-
771 (2016).
116. T. Paretkar, E. Dimitrov, Activation of enkephalinergic (Enk) interneurons in the central
amygdala (CeA) buffers the behavioral effects of persistent pain. Neurobiol Dis 124,
364-372 (2019).
117. Z. Zhang, Z. Ma, W. Zou, H. Guo, M. Liu, Y. Ma, L. Zhang, The Appropriate Marker for
Astrocytes: Comparing the Distribution and Expression of Three Astrocytic Markers in
Different Mouse Cerebral Regions. Biomed Res Int 2019, 9605265 (2019).
118. A. M. Jurga, M. Paleczna, J. Kadluczka, K. Z. Kuter, Beyond the GFAP-Astrocyte
Protein Markers in the Brain. Biomolecules 11, (2021).
119. V. Dyachuk, A. Furlan, M. K. Shahidi, M. Giovenco, N. Kaukua, C. Konstantinidou, V.
Pachnis, F. Memic, U. Marklund, T. Müller, C. Birchmeier, K. Fried, P. Ernfors, I.
Adameyko, Parasympathetic neurons originate from nerve-associated peripheral glial
progenitors. Science 345, 82-87 (2014).
120. M. S. Erwig, J. Patzig, A. M. Steyer, P. Dibaj, M. Heilmann, I. Heilmann, R. B. Jung, K.
Kusch, W. Möbius, O. Jahn, K.-A. Nave, H. B. Werner, Anillin facilitates septin
assembly to prevent pathological outfoldings of central nervous system myelin. eLife
8, (2019).
121. J. Spaas, W. M. A. Franssen, C. Keytsman, L. Blancquaert, T. Vanmierlo, J. Bogie, B.
Broux, N. Hellings, J. van Horssen, D. K. Posa, D. Hoetker, S. P. Baba, W. Derave, B.
O. Eijnde, Carnosine quenches the reactive carbonyl acrolein in the central nervous
system and attenuates autoimmune neuroinflammation. J Neuroinflammation 18, 255
(2021).
122. J. DeSisto, R. O'Rourke, H. E. Jones, B. Pawlikowski, A. D. Malek, S. Bonney, F.
Guimiot, K. L. Jones, J. A. Siegenthaler, Single-Cell Transcriptomic Analyses of the
Developing Meninges Reveal Meningeal Fibroblast Diversity and Function. Dev Cell
54, 43-59 e44 (2020).

123. D. S. Liu, T. L. Xu, Cell-Type Identification in the Autonomic Nervous System. Neurosci
Bull 35, 145-155 (2019).
124. B. J. Kang, D. A. Chang, D. D. Mackay, G. H. West, T. S. Moreira, A. C. Takakura, J.
M. Gwilt, P. G. Guyenet, R. L. Stornetta, Central nervous system distribution of the
transcription factor Phox2b in the adult rat. J Comp Neurol. 503, 627-641 (2007).
125. M. Adler, N. Moriel, A. Goeva, I. Avraham-Davidi, S. Mages, T. S. Adams, N. Kaminski,
E. Z. Macosko, A. Regev, R. Medzhitov, M. Nitzan, Emergence of division of labor in
tissues through cell interactions and spatial cues. Cell Rep 42, 112412 (2023).
126. A. Steimel, J. Suh, A. Hussainkhel, S. Deheshi, J. M. Grants, R. Zapf, D. G. Moerman,
S. Taubert, H. Hutter, The C. elegans CDK8 Mediator module regulates axon guidance
decisions in the ventral nerve cord and during dorsal axon navigation. Dev Biol 377,
385-398 (2013).
127. T. Daiber, C. J. VanderZwan-Butler, G. J. Bashaw, T. A. Evans, Conserved and
divergent aspects of Robo receptor signaling and regulation between Drosophila
Robo1 and C. elegans SAX-3. Genetics 217, (2021).
128. S. Schmetsdorf, U. Gärtner, T. Arendt, Constitutive Expression of Functionally Active
Cyclin-Dependent Kinases and Their Binding Partners Suggests Noncanonical
Functions of Cell Cycle Regulators in Differentiated Neurons. Cerebral Cortex 17,
1821–1829 (2007).
129. K. K. Kummer, M. Zeidler, T. Kalpachidou, M. Kress, Role of IL-6 in the regulation of
neuronal development, survival and function. Cytokine 144, 155582 (2021).
130. P. März, J. G. Cheng, R. A. Gadient, P. H. Patterson, T. Stoyan, U. Otten, S. Rose-
John, Sympathetic neurons can produce and respond to interleukin 6. Proc Natl Acad
Sci 95, 3251-3256 (1998).
131. K. M. McGrail, J. M. Phillips, K. J. Sweadner, Immunofluorescent localization of three
Na,K-ATPase isozymes in the rat central nervous system: both neurons and glia can
express more than one Na,K-ATPase. J Neurosci 11, 381-391 (1991).
132. H. Poulsen, H. Khandelia, J. P. Morth, M. Bublitz, O. G. Mouritsen, J. Egebjerg, P.
Nissen, Neurological disease mutations compromise a C-terminal ion pathway in the
Na(+)/K(+)-ATPase. Nature 467, 99-102 (2010).
133. A. Abousaab, J. Warsi, B. Elvira, F. Lang, Caveolin-1 Sensitivity of Excitatory Amino
Acid Transporters EAAT1, EAAT2, EAAT3, and EAAT4. J Membr Biol 249, 239-249
(2016).
134. E. R. Bongarzone, C. W. Campagnoni, K. Kampf, E. C. Jacobs, V. W. Handley, V.
Schonmann, A. T. Campagnoni, Identification of a new exon in the myelin proteolipid
protein gene encoding novel protein isoforms that are restricted to the somata of
oligodendrocytes and neurons. J Neurosci 19, 8349-8357 (1999).
135. L. F. Nunnelly, M. Campbell, D. I. Lee, P. Dummer, G. Gu, V. Menon, E. Au, St18
specifies globus pallidus projection neuron identity in MGE lineage. Nat Commun 13,
7735 (2022).
136. C. Plachez, C. Lindwall, N. Sunn, M. Piper, R. X. Moldrich, C. E. Campbell, J. M.
Osinski, R. M. Gronostajski, L. J. Richards, Nuclear factor I gene expression in the
developing forebrain. J Comp Neurol 508, 385-401 (2008).
137. J. Fraser, A. Essebier, A. S. Brown, R. A. Davila, D. Harkins, O. Zalucki, L. P. Shapiro,
P. Penzes, B. J. Wainwright, M. P. Scott, R. M. Gronostajski, M. Boden, M. Piper, T. J.
Harvey, Common Regulatory Targets of NFIA, NFIX and NFIB during Postnatal
Cerebellar Development. Cerebellum 19, 89-101 (2020).
138. L. Dubois , A. Vincent, The COE--Collier/Olf1/EBF--transcription factors: structural
conservation and diversity of developmental functions. 108 3-12, (2001).
139. Y. Liu, T. Lear, O. Iannone, S. Shiva, C. Corey, S. Rajbhandari, J. Jerome, B. B. Chen,
R. K. Mallampalli, The Proapoptotic F-box Protein Fbxl7 Regulates Mitochondrial
Function by Mediating the Ubiquitylation and Proteasomal Degradation of Survivin. J
Biol Chem 290, 11843-11852 (2015).
140. H. Kim, N. C. Berens, N. E. Ochandarena, B. D. Philpot, Region and Cell Type
Distribution of TCF4 in the Postnatal Mouse Brain. Front Neuroanat 14, 42 (2020).
141. A. Sirp, A. Shubina, J. Tuvikene, L. Tamberg, C. S. Kiir, L. Kranich, T. Timmusk,
Expression of alternative transcription factor 4 mRNAs and protein isoforms in the

developing and adult rodent and human tissues. Front Mol Neurosci 15, 1033224 (2022).

142. R. D. Hodge, T. E. Bakken, J. A. Miller, K. A. Smith, E. R. Barkan, L. T. Graybuck, J. L. Close, B. Long, N. Johansen, O. Penn, Z. Yao, J. Eggermont, T. Hollt, B. P. Levi, S. I. Shehata, B. Aevermann, A. Beller, D. Bertagnolli, K. Brouner, T. Casper, C. Cobbs, R. Dalley, N. Dee, S. L. Ding, R. G. Ellenbogen, O. Fong, E. Garren, J. Goldy, R. P. Gwinn, D. Hirschstein, C. D. Keene, M. Keshk, A. L. Ko, K. Lathia, A. Mahfouz, Z. Maltzer, M. McGraw, T. N. Nguyen, J. Nyhus, J. G. Ojemann, A. Oldre, S. Parry, S. Reynolds, C. Rimorin, N. V. Shapovalova, S. Somasundaram, A. Szafer, E. R. Thomsen, M. Tieu, G. Quon, R. H. Scheuermann, R. Yuste, S. M. Sunkin, B. Lelieveldt, D. Feng, L. Ng, A. Bernard, M. Hawrylycz, J. W. Phillips, B. Tasic, H. Zeng, A. R. Jones, C. Koch, E. S. Lein, Conserved cell types with divergent features in human versus mouse cortex. Nature 573, 61-68 (2019).

143. T. Kuroda, S. Yasuda, S. Tachi, S. Matsuyama, S. Kusakawa, K. Tano, T. Miura, A. Matsuyama, Y. Sato, SALL3 expression balance underlies lineage biases in human induced pluripotent stem cell differentiation. Nat Commun 10, 2175 (2019).

144. F. Morello, D. Borshagovski, M. Survila, L. Tikker, S. Sadik-Ogli, A. Kirjavainen, N. Estartus, L. Knaapi, L. Lahti, P. Toronen, L. Mazutis, A. Delogu, M. Salminen, K. Achim, J. Partanen, Molecular Fingerprint and Developmental Regulation of the Tegmental GABAergic and Glutamatergic Neurons Derived from the Anterior Hindbrain. Cell Rep 33, 108268 (2020).

145. L. Wolf, C. S. Gao, K. Gueta, Q. Xie, T. Chevallier, N. R. Podduturi, J. Sun, I. Conte, P. S. Zelenka, R. Ashery-Padan, J. Zavadil, A. Cvekl, Identification and characterization of FGF2-dependent mRNA: microRNA networks during lens fiber cell differentiation. G3 (Bethesda) 3, 2239-2255 (2013).

146. H. Liu, M. Aramaki, Y. Fu, D. Forrest, Retinoid-Related Orphan Receptor beta and Transcriptional Control of Neuronal Differentiation. Curr Top Dev Biol 125, 227-255 (2017).

147. M. G. Del Barrio, S. Bourane, K. Grossmann, R. Schule, S. Britsch, D. D. O'Leary, M. Goulding, A transcription factor code defines nine sensory interneuron subtypes in the mechanosensory area of the spinal cord. PLoS One 8, e77928 (2013).

148. Y. Li, A. Osuma, E. Correa, M. A. Okebalama, P. Dao, O. Gaylord, J. Aburas, P. Islam, A. E. Brown, P. Kratsios, Establishment and maintenance of motor neuron identity via temporal modularity in terminal selector function. Elife 9, (2020).

149. D. Moruzzo, L. Nobbio, B. Sterlini, G. G. Consalez, F. Benfenati, A. Schenone, A. Corradi, The Transcription Factors EBF1 and EBF2 Are Positive Regulators of Myelination in Schwann Cells. Mol Neurobiol 54, 8117-8127 (2017).

150. M. A. Jimenez, P. Akerblad, M. Sigvardsson, E. D. Rosen, Critical role for Ebf1 and Ebf2 in the adipogenic transcriptional cascade. Mol Cell Biol 27, 743-757 (2007).

151. H. Yu, M. Rubinstein, M. J. Low, Developmental single-cell transcriptomics of hypothalamic POMC neurons reveal the genetic trajectories of multiple neuropeptidergic phenotypes. Elife 11, (2022).

152. I. Singh, L. Wang, B. Xia, J. Liu, A. Tahiri, A. El Ouaamari, M. B. Wheeler, Z. P. Pang, Activation of arcuate nucleus glucagon-like peptide-1 receptor-expressing neurons suppresses food intake. Cell Biosci 12, 178 (2022).

153. L. Xiao, D. P. Merullo, T. M. I. Koch, M. Cao, M. Co, A. Kulkarni, G. Konopka, T. F. Roberts, Expression of FoxP2 in the basal ganglia regulates vocal motor sequences in the adult songbird. Nat Commun 12, 2617 (2021).

154. T. T. Du, J. B. Dewey, E. L. Wagner, R. Cui, J. Heo, J. J. Park, S. P. Francis, E. Perez-Reyes, S. J. Guillot, N. E. Sherman, W. Xu, J. S. Oghalai, B. Kachar, J. B. Shin, LMO7 deficiency reveals the significance of the cuticular plate for hearing function. Nat Commun 10, 1117 (2019).

155. M. A. Wozniak, B. M. Baker, C. S. Chen, K. L. Wilson, The emerin-binding transcription factor Lmo7 is regulated by association with p130Cas at focal adhesions. PeerJ 1, e134 (2013).

156. T. Kawaue, A. Shitamukai, A. Nagasaka, Y. Tsunekawa, T. Shinoda, K. Saito, R. Terada, M. Bilgic, T. Miyata, F. Matsuzaki, A. Kawaguchi, Lzts1 controls both neuronal delamination and outer radial glial-like cell generation during mammalian cerebral development. Nat Commun 10, 2780 (2019).

157. O. Randlett, L. Poggi, F. R. Zolessi, W. A. Harris, The oriented emergence of axons from retinal ganglion cells is directed by laminin contact in vivo. Neuron 70, 266-280 (2011).

158. N. Ichikawa-Tomikawa, J. Ogawa, V. Douet, Z. Xu, Y. Kamikubo, T. Sakurai, S. Kohsaka, H. Chiba, N. Hattori, Y. Yamada, E. Arikawa-Hirasawa, Laminin alpha1 is essential for mouse cerebellar development. Matrix Biol 31, 17-28 (2012).

159. E. S. Liau, S. Jin, Y. C. Chen, W. S. Liu, M. Calon, S. Nedelec, Q. Nie, J. A. Chen, Single-cell transcriptomic analysis reveals diversity within mammalian spinal motor neurons. Nat Commun 14, 46 (2023).

Acknowledgements

We would like to thank authors of the Ludwig and the Dowsett publications for making publicly available their datasets. We thank M. Myers Jr. for the valuable help provided; S. Hebert and S. Bailey for their insights and suggestions in regards to the bioinformatic analysis; members of the Kokoeva lab for their insights on the *in-situ* hybridization and immunohistochemistry pipelines; S. Feng and M. Fu for technical assistance for images acquisition at the Molecular Imaging Platform at the RI-MUHC, and A. Duensing for assistance with relevant file transfer.

Funding

This research was also supported by grants from Canadian Institutes for Health Research (PJT180590 for PVS and 202010PJT for MK) the Natural Sciences and Engineering Research Council of Canada (RGPIN-2022-03390). HJM was funded in part by the Canada First Research Excellence Fund and Fonds de recherche du Quebec, awarded to the Healthy Brains, Healthy Lives initiative.

Author contributions

Conceptualization: CH and PS

Methodology: CH, AB, LM, MK and PS

Sample collection: AB and PS

Software: CH, LM and HJM

Data analysis: CH, LM, MK and PS

Data curation: CH and HJM

Validation of findings: CH, FS, HJM and PS

Imaging: CH and PS

Visualization: CH, MK and PS

Supervision: CH and PS

Writing—original draft: CH and PS —review and editing: CH, AB, LM, FS, HJM, MK and PS

Funding acquisition: MK and PS

1250

1251   <u>Competing interests</u>

1252   The authors declare that they have no competing interests.

1253

<u>Data and materials availability</u>

All data needed to evaluate the conclusions in the paper are present in the paper and/or the

Supplementary Materials. FASTQ files have been deposited in the NCBI Sequence Read

Archive (Accession: PRJNA1161292). In addition, our murine and rat datasets have been

made available for visualization and analysis through the Broad Institute Single Cell Portal

(https://singlecell.broadinstitute.org/single_cell/study/SCP2773). The labels from our cell atlas

can be transferred to new datasets through treeArches using the available hosted Jupyter

Notebook in Google Colab (https://colab.research.google.com/drive/1v983gcQxIwBN-

4lGx18JXPweZoKqMbwX?usp=sharing). Original code used for constructing the murine and

rodent cell hierarchies and predicting labels using treeArches are available in the GitHub

repository https://github.com/LabSabatini/DVC_cell_atlas.

 Figure legends

Figure 1: Two layers of cell identity for the murine DVC cells from snRNA-seq

data

A. Adult C57BL/6J mice were subjected to 10X Genomics-based snRNA barcoding

and sequencing after dissection of their DVC (n=30). **B.** Labeled UMAP plot of the

mouse DVC (cells n=99,740) at the lowest resolution (layer 1) including 4 cell

identities, and **C.** the cell class (layer 2) including 18. **D.** Heatmap of the main marker

genes (average log2FC expression >80 percentiles of upregulated genes) for each

neuronal class of the mouse DVC (MAST algorithm; neurons n= 48,131; adj. p<0.05).

Each column is one cell and each row is one marker gene. **E.** Stacked barplot of log-

normalized counts of genes for 10 transcripts (i.e. *Gad2*, *Gad1*, *Slc6a5*, *Slc17a6*,

Slc18a2, *Addc*, *Sst*, *Npy*, *Cck* and *Slc5a7*) related to transmission/release of 8

neurotransmitters and neuropeptides, by neuronal cell class. Each bar represents one

neuron (n=48,131).

snRNA-seq=single-nuclei RNA-sequencing; DVC=dorsal vagal complex; FACs=fluorescence-

activated cell sorting; UMAP=uniform manifold approximation and projection; FC=fold-change;

adj.=adjusted; GABA=gamma-aminobutyric acid

1282

Figure 2: Third layer of non-neuronal murine DVC cell identities

A. Labeled UMAP plot of the mouse DVC with the 15 non-neuronal cell identities shown

(total cells n=99,740; non-neuronal cells n=51,609). **B.** UMAP plot showing scaled

expression of *Gfap*, *Gria2* and *Kcnj3* genes in Ca^+-permeable astrocytes (circled; total

cells n=99,740; Ca^+-permeable astrocytes n=1,719). **C.** Barplot of log-normalized

expression of six genes in oligodendrocytes (premyelinating n=267; myelinating

intermediate n=25,692; myelinating n= 6,170). Only premyelinating oligodendrocytes

lack expression of *Mobp*, involved in myelin formation. Reduction of *Synpr* and *Opalin*

expression is observed as *Anln* and *Aff3* increase in myelinating oligodendrocytes. **D.**

Balloon plot of the main cell types of non-neuronal cells and neurons showing average

log-normalized counts of their marker genes (MAST algorithm; adj. p<0.05). Markers

shown are upregulated genes in >80% of cells per group with average log2FC>4, or

upregulated in >70% of cells with average log2FC>8.

DVC=dorsal vagal complex; UMAP=uniform manifold approximation and projection; APNTS=Area

postrema and nucleus of the solitary tract; OL=oligodendrocyte; OPC=oligodendrocyte precursor cell;

Smc=smooth muscle cells

1299

1300 **Figure 3: Neuronal populations of the murine DVC**

1301 **A.** Labeled UMAP plot with the neuronal layer 3 cell identities (neurons n=48,131). **B.**

1302 Monoamine class balloon plot showing average log-normalized expression of marker

genes for each cell identity (MAST algorithm; M0 cells n=770, M1 cells n=551, M2 cells n=666 and M3 cells n=2,593; adj. p<0.05), and **C.** heatmap of expression of monoamine related genes. In the heatmap, each column represents one cell and each row, one gene. **D.** Barplot of *Gcg* average log-normalized expression by neuronal cell identity. **E.** UMAP plot showing the expression of leptin receptor and prolactin receptor genes in neurons. The GLP1 cluster is highlighted (GLP1 cells n=153). **F.** Stacked barplot of the proportion of cells expressing one or more GABA-related genes (i.e. *Slc32a1*, *Gad1*, *Gad2*) and glutamate-related genes (i.e. *Slc17a6*, *Slc17a7*). The proportion of cells per cell group co-expressing GABA and glutamate associated genes are shown in purple. Only cells with log-normalized counts >0 were considered to express the genes.

DVC=dorsal vagal complex; UMAP=uniform manifold approximation and projection; Res.=resolution; GABA=gamma-aminobutyric acid

Figure 4: *Th* and *Cck* co-expression in the DVC

A. UMAP plot of neuronal *Th* and *Cck* scaled expression and cells with co-expression of both genes (neurons n=48,131; *Th*-expressing neurons n=764; *Cck*-expressing neurons n=3,821; *Th/Cck* co-expressing neurons n=80). On the right plot, cells highlighted in magenta are the co-expressing neurons. The three neuronal cell identities in which the majority of nuclear co-expression of *Th* and *Cck* mRNA is found, as well as the percentage of total co-expressing neurons, are shown. **B.** *In-situ* hybridization of *Cck* and *Th* mRNA in coronal mouse DVC sections corresponding to -7.2mm and -7.56mm relative to bregma showing overlap of *Th/Cck*. Some of the cells with overlapping signal are highlighted with an arrow in the enhanced merged image.

DVC=dorsal vagal complex; UMAP=uniform manifold approximation and projection

Figure 5: The murine DVC cell hierarchy

A. Dendrogram of the harmonized hierarchy which incorporates cells from Ludwig and our datasets. Orange cell identities are a fourth layer of cell identity resolution obtained as some of the Ludwig dataset identities are subgroups of our original cell identities at their highest resolution. Magenta labels represent layer 3 of cellular granularity. These two layers are considered high-resolution. **B.** UMAP plot of the integration between our and Ludwig datasets using treeArches (cells n=171,868). **C.** Pairwise heatmap showing the proportion of cells originally labeled in this study and in Ludwig dataset (y-axis), predicted to belong to each identity group (x-axis) by treeArches using our learned harmonized hierarchy. In blue are the labels considered non-specific for a high-resolution cell identity (n=2,324; 1.3%). In pink are the cell labels for rejected cells, therefore not assigned any identity by treeArches (n=3,108; 1.8%).

D. UMAP plot of the Dowsett dataset labeled through treeArches using the learned

harmonized hierarchy representation from our murine DVC atlas. The 'unlabeled' cells

include those rejected by the algorithm and thus without an assigned identity, and

those with an unspecific layer-3/4 label. For example, some were labeled 'neuron' or

'monoamine-class' but without a high-resolution cell identity.

UMAP=uniform manifold approximation and projection; APNTS=Area postrema and nucleus of the

solitary tract; Lnc=long non-coding; OL=oligodendrocyte; OPC=oligodendrocyte precursor cell;

Neur=neuronal; COE=collier/Olf1/EBF transcription factor

1349

Figure 6: The snRNA-seq derived cell identities for the rat DVC

A. Schematic of the pipeline for snRNA-seq of the rat DVC. **B.** Labeled UMAP plots of

the low resolution (i.e. layers 1 and 2 of granularity) and **C.** high resolution (i.e. layer

3) cell identities in our rat DVC dataset (cells n=12,167). We labeled four layer-1,

nineteen layer-2 and fifty-two layer-3 cell identities. Those labels novel to this dataset

and not present in the murine data are highlighted with a pink ♦ symbol. We found a

small cluster (cells n=35) that we could not corroborate its identity at any layer of

cellular granularity that we named 'unspecific'.

snRNA-seq=single-nuclei RNA-sequencing; DVC=dorsal vagal complex; FACs=fluorescence-

activated cell sorting; UMAP=uniform manifold approximation and projection; APNTS=Area postrema

and nucleus of the solitary tract; Lnc=long non-coding; OL=oligodendrocyte; OPC=cursor cell;

Neur=neuronal; COE=collier/Olf1/EBF transcription factors

1362

Figure 7: Two novel neuronal classes specific to the rat DVC

A. Balloon plots comparing the expression of the marker genes of the two novel

neuronal classes found in rats (cells n=12,167): immunity-akin and the ortus-akin

classes (framed in gray). Expression is shown across the layer-2 rat cell identities.

Based on the expression of two DVC neuronal markers *Mtus2* and *Syt1*, we

corroborated those cell classes contain neurons, and not microglial or

vascular/endothelial cells (In orange and red, respectively). The immunity-akin class

(n=48 neurons) shares high expression of *Cst3*, *Inpp5d*, *Csf1r*, *Slco2b1* and *Lyn* with

microglial cells. Meanwhile, the ortus-akin class (n=34 neurons) shows higher overlap

of gene expression markers (i.e. *Arhgap31*, *Pdgfra*, *Bcas1*, *Ppfibp1* and *Itga9*) with

OPCs (highlighted with a ♦ symbol). Mouse cell identities (cells n=99,740) do not

show these expression patterns (framed in black). Average expression was calculated

using log-normalized counts. **B.** Violin plots of scaled expression of 10 genes (i.e.

Glp1r, Gfral, Calcr, Ramp3, Gipr, Lepr, Cckar, Cckbr ,Mc4r and *Prlr*) coding for

metabolism-associated receptors in the Pdgfra and immunity-related rat neurons

(Pdgfra neurons n=34; immunity-related neurons n=48). Pdgfra neurons are the only

layer-3 identity within the ortus-akin class. An overlapping dotplot shows one dot per

cell. **C.** Immunofluorescent detection of PDGFRA in mouse and rat DVC. Pink

arrowheads point to cells with OPC morphology, white arrowheads point to cells with

fibroblast morphology and the yellow arrowhead points to cells with neural

morphology.   **D.** Co-staining for PDGFRA and HuC/D in rat DVC with high

magnification (right images) of area postrema PDGFRA-expressing cells . Yellow

arrowheads indicate colocalization of HuC/D and PDGFRA.

DVC=dorsal vagal complex; OPC=oligodendrocyte precursor cell; AP=area postrema; NTS=nucleus

of the solitary tract; PDGFRA= Platelet-derived growth factor receptor A; CC=central canal

1388

Figure 8: The rodent DVC cell hierarchy

A. UMAP plot of the integration between the murine hierarchy and our rat dataset using

treeArches (cells n=184,035). **B.** Dendrogram of the harmonized hierarchy of our

labeled rat dataset and the murine DVC hierarchy. We highlight the incorporated rat

cell identities (blue and magenta). Those in magenta are the novel rat identities

established by us. The immunity-akin and the ortus-akin classes were not found in the

murine datasets. The M3 and M5 rat identities are subgroups of a Ludwig layer 4 cell

identity, therefore yields a fifth layer within the monoamine neuronal class.

UMAP=uniform manifold approximation and projection; DVC=dorsal vagal complex; APNTS=Area

postrema and nucleus of the solitary tract; Lnc=long non-coding; OL=oligodendrocyte;

OPC=oligodendrocyte precursor cell; Neur=neuronal; COE=collier/Olf1/EBF transcription factors

1400

Figure 9: Meal-related transcriptional changes in the mouse DVC

A. Horizontal barplots of the number of differential genes between refed and ad libitum

fed, and refed and fasted mice in glial cells and neurons per layer-2 cell identity (MAST

algorithm on log-normalized counts; adj. p<0.05; refed neurons n=14,396; refed glial

cells n=18,019; ad libitum fed neurons n= 9,885; ad libitum fed glial cells n=4,610;

fasted neurons n=8,829; fasted glial cells n=12,496). **B.** Venn Diagrams of the

differentially expressed genes in magnaclass 1 and magnaclass 2 neurons between

refed and ad libitum fed, and between refed and fasted treatments (MAST algorithm

on log-normalized counts). The number of overlapping log2FC≥1 upregulated genes

between the two magnaclasses per treatment are highlighted. The treatment-induced

changes in only one cell class are shown as non-overlapping. The percentage is

based on the total differential genes surveyed per comparison. **C.** Volcano plots of the

differential genes in neurons belonging to the magnaclass 1 and the magnaclass 2

between refed and ad libitum fed, and **D.** refed and fasted treatments (MAST algorithm

on log-normalized counts; adj. p<0.05). Each point represents one gene. Only genes

upregulated or downregulated with a log2FC≥2 are labeled. Since we considered a

minimum log fold-change of 0.1 between treatments per cell group, those genes with
very low variance (i.e. log fold-change<0.1) were excluded from the differential
expression analysis, therefore a variable number of genes are shown per comparison
per identity, in the volcano plots.
DVC=dorsal vagal complex; vr=versus; OPC=oligodendrocyte precursor cell;   FC=fold-change;
adj.=adjusted; NS=non-significant
1423

 Supplementary Files

 Figure S1: Mapping of *Gad2* and VGLUT2 in neurons

UMAP plot of the murine neurons (n=48,131) showing scaled log-normalized expression of **A.**
Gad2, a gene related to GABA synthesis and transmission, **B.** and VGLUT2, a glutamate
transporter. Cells with higher expression of *Gad2* belong to the neuronal magnaclass1, while
cells with higher expression of VGLUT2 (*Slc17a6*) to the magnaclass2 (both highlighted in
yellow in adjacent plots). However, the expression is not mutually exclusive among both
classes, nor every neuron belonging to each class express the respective transcript.
UMAP=uniform manifold approximation and projection; GABA=gamma-aminobutyric acid; VGLUT2=vesicular
glutamate transporter-2

1434

Figure S2: Mapping of DVC populations in a murine hypothalamic dataset
UMAP plots of a randomly selected subset of the HypoMap database (cells n=76,260; samples
n=32) labeled **A.** by sample and **B.** by original curated cell class. **C.** Balloon plot showing
average expression of the top 5 upregulated gene markers of each neuronal DVC class (MAST
algorithm; adj. p<0.05) in the neurons of the HypoMap subset (n=45,679). We selected the
neuronal curated class cells from our initial HypoMap subset and reprocessed them, including
clustering through *k*-nearest neighbors algorithm using the first 30 PCs and Harmony reduction
embeddings. Resulting clusters at resolution 0.5 are shown. Average expression in the
HypoMap neurons was calculated using log-normalized counts. **D.** Set of UMAP plots showing
scaled expression of three genes (i.e. *Gfap*, *Gria2* and *Kcnj3*) which have a specific pattern of
expression in murine DVC Ca$^+$-permeable astrocytes. The astrocytic population is circled in
purple (astrocytes n=7,629) and while *Gfap* and *Gria2* expression is in agreement to what we
observe in Ca$^+$-permeable astrocytes, in the hypothalamic astrocytes *Kcnj3* is not expressed.
E. Set of UMAP plots of expression of *Aldh1a1* and *Stk39* which are markers of p-fibroblasts
and s-fibroblasts in the DVC. The hypothalamic fibroblasts are circled in magenta (fibroblasts
n=300).
DVC=dorsal vagal complex; UMAP=uniform manifold approximation and projection; AMPAr=α-amino-3-hydroxy-
5-methyl-4-isoxazolepropionic acid receptor; GIRK1=G protein-coupled inwardly rectifying potassium channel 1;
OPC=Oligodendrocyte precursor cell; NG=*Ng2* glial cell

1454

Figure S3: Mapping of DVC astrocytes in a murine cortical/hippocampal dataset
UMAP plots of a cortical/hippocampal astrocytes dataset by **A.** sample, **B.** brain area and **C.**
original astrocyte type (cell n=1,811). **D.** Set of UMAP plots showing scaled expression of three
genes (i.e. *Gfap, Gria2* and *Kcnj3*) which have a specific pattern of expression in murine DVC
Ca$^+$-permeable astrocytes. There is some overlap in *Gfap* and *Gria2* expression and almost no
expression of *Kcnj3*, different from what we observe in DVC Ca$^+$-permeable astrocytes.

1461

Figure S4: Trajectory inference of the oligodendrocyte lineage

Dotplot of the two principal components obtained through dimensionality reduction using the
main 50 principal components analysis on log-normalized gene expression data in
oligodendrocytes (cells n=32,129) **A.** merged and **B.** by sample, and on Harmony embeddings
in oligodendrocytes **C.** merged and **D.** by sample. Cellular trajectory was inferred using
SCORPIUS also on normalized gene expression data (**A.** and **B.**) and on the Harmony
embeddings (**C.** and **D.**), and drawn across the data points (black line). Each data point
represents one cell.

OL=oligodendrocyte

1471

Figure S5: Murine DVC clusters

A. UMAP plot of the mouse DVC clusters at resolution 1 (cells n=99,740). Cells from mice
exposed to 5 food and aversion-related treatments were integrated using the Harmony
algorithm in R and the resulting embeddings were used to calculate the clusters and UMAP
projections. We subset and re-clustered three clusters containing oligodendrocytes, vascular
cells and fibroblasts, respectively (highlighted). **B.** Violin plots of the two types of fibroblasts in
cluster 26, distinguished by *Skt39* (stress-response 's-fibroblasts') and *Aldh1a1* (perivascular
'p-fibroblasts') expression. Each data point represents one cell (s-fibroblasts n=244; p-
fibroblasts n=559). **C.** UMAP plot of subclusters from clusters 23 (oligodendrocytes) and **D.** 27
(vascular cells) using the Harmony embeddings. Gene expression markers for cell populations
were mapped (**Table S2**) and the smallest resolution that allowed to separate them was further
used to label the cells. Cells in subclusters 1 and 2 from cluster 23 were labeled
'premyelinating_OL' whilst subclusters 0 and 3 were labeled 'myelinating_intermediate_OL'. In
cluster 27, subcluster 7 was labeled 'pericyte', subcluster 4 was labeled 'smooth_muscle' and
the rest of the clusters were considered to be endothelial cells.

DVC=dorsal vagal complex; Res.=resolution; UMAP=uniform manifold approximation and projection;
OL=oligodendrocyte; smc=smooth muscle cell.

1489

1490 Figure S6: Mapping of markers for reactive microglia

1491 UMAP plot of the microglia from the mouse DVC showing scaled log-normalized expression of
1492 *Cd5*, *Zbp1*, *Cxcl9* and *Cxcl10*. Microglia was subset and re-processed using the harmony
1493 integration embeddings to obtain the UMAP coordinates (n=2,787).

1494 UMAP=uniform manifold approximation and projection; DVC=dorsal vagal complex

1495

Figure S7: Initial murine cell hierarchy using our samples

A. UMAP plot of the integration of our samples (based on the five mouse treatments
administered to mice prior to euthanasia; see **Methods**) using treeArches. There were two
cohorts of refed mice. **B.** UMAP plot of the fifty layer-3 cell identities obtained using the

treeArches pipeline. **C.** Dendrogram showing the initial hierarchy of the three layers of cell
identity established by us in this study.

1505

<u>Figure S8: Incorporation of the Ludwig dataset to our initial hierarchy using treeArches</u>
A. Barplot of log-normalized expression of OPC markers (i.e. *Pdgfra*, *Cspg4*, *C1ql1*) and
premyelinating OL markers (i.e. *Bmp4*, *Fyn*) in the OPCs-labeled cells from the Ludwig dataset
(total cells n=1,770; cells expressing >1 marker gene n=1,104). Cells labeled as OPCs in the
Ludwig dataset are a mix of OPC and premyelinating OL. Some cells lack expression of the
five marker genes used in this plot, they are represented on the left side with an empty bar and
those expressing *Fyn*/*Bmp4* but not expressing the OPC markers are likely premyelinating OLs.
Cells were ordered using hierarchical clustering based on these five gene expression data
through the Ward's method in R. We only indicate the 'likely cell identity' for cells with
expression of the five marker genes. **B.** UMAP showing the manual mapping of Ludwig and
Dowsett neuronal clusters and identities in neurons from this study. We used the gene
expression markers for all clusters from Ludwig and Dowsett to identify which of our neurons
shared identity with those described in both publications. Only groups and clusters
successfully mapped manually are shown. The Ludwig identities are in black, Dowsett
identities are in blue and our identities are in purple. Most of neuronal clusters in both publicly
available datasets, are subgroups of our cell identities.

1525

<u>Figure S9: Rat DVC neurons</u>
A. Labeled UMAP plot with the neuronal layer 3 cell identities in our rat dataset (neurons
n=4,721). Neurons were subset and re-processed to obtain the UMAP coordinates. **B.** Violin
plots of scaled expression of 12 genes (i.e. *Gad2, Gad1, Slc6a5, Slc17a7, Slc17a6, Th, Ddc,*
Nos1, Sst, Npy, Cck and *Slc5a7*) related to transmission/release of GABA, glutamate,
monoamines, nitric oxide, somatostatin, neuropeptide-Y, cholecystokinin and acetylcholine in
the Pdgfra and immunity-related rat neurons (Pdgfra neurons n=34; immunity-related neurons
n=48). An overlapping dotplot shows one data point representing each cell.

1536

1537 <u>Figure S10: Dendrogram after incorporation of the rat dataset to the murine hierarchy</u>

1538 Dendrogram (i.e. tree) resulting from incorporating our rat dataset (cells n=12,167) to the
1539 existing murine DVC cell hierarchy using treeArches. The rat cell populations in red that did
1540 not show similarity with the previously existing murine identities are appended at the bottom
1541 by treeArches (shown in red).

1542

<u>Figure S11: Expression of *Agrp* and *Hcrt* in DVC neurons</u>
A. UMAP plots of the neurons (n=48,131) from our murine DVC dataset showing scaled
expression of *Agrp* and *Hcrt*. **B.** *In-situ* hybridization images of *Agrp* and *Hcrt* in the DVC show
no exonic mRNA. We included tissue sections of the arcuate nucleus of the hypothalamus and
in the lateral hypothalamus as controls, and *Agrp* and *Hcrt* were present in these sites,
respectively. For each reaction, tissue from the same mouse was used through two different
sections, one including the DVC and one including the hypothalamic region. **C.** UMAP plots of
the Dowsett (n=8,288) and the Ludwig datasets neurons (n=49,392) showing scaled
expression of *Agrp* and *Hcrt*. **D.** Genome visualization of reads coverage in the *Agrp* and *Hcrt*
regions of the *Mus musculus* refdata-gex-mm10-2020-A reference from two of our mouse
sample sequencing replicates. Resulting BAM files from the original murine DVC FASTQ files
were obtained after processing. We visualized the reads from these BAM files that mapped to
the *Agrp* and *Hcrt* genes using IGV v2.16.1(*78*). The coverage of these genes (*Agrp* region:
chr8:105,566,180–105,582,268; *Hcrt* region: chr11:100,761,693–100,762,931) are shown in
gray. The majority of the reads mapped to the intronic regions of *Agrp*, and to the last part of
the 3'-end of exon 2 and the 3'-untranslated region of *Hcrt*. **E.** UMAP plot of rat DVC neurons
(n=4,721) showing scaled expression of *Agrp* and *Hcrt*. For all UMAP projections, neurons
from each dataset were subset based on our labeling for our murine and rat datasets, the
treeArches resulting labeling for the Dowsett dataset, and the original labeling from Ludwig
and collaborators. Then the neurons were processed using Seurat(*63*) and integrated using
Harmony(*65*), to yield the final UMAP embeddings.
DVC=dorsal vagal complex; UMAP=uniform manifold approximation and projection; AP=area postrema;
NTS=nucleus of the solitary tract; ARH=arcuate nucleus of the hypothalamus; 3v=third ventricle; ME=median
eminence; LH=lateral hypothalamus. BAM=binary alignment map

1567

<u>Figure S12: Expression of metabolism-associated receptors in the DVC magnaclasses</u>
Violin plots of scaled expression of 10 genes (i.e. *Glp1r, Gfral, Calcr, Ramp3, Gipr, Lepr, Cckar,*
Cckbr ,Mc4r and *Prlr*) coding for metabolism-associated receptors in the magnaclass1 and
magnaclass2 neurons of the mouse DVC in fasted (magnaclass1 n=2,756; magnaclass2
n=2,789) and refed (magnaclass1 n=4,586; magnaclass2 n=4,423).
DVC=dorsal vagal complex
1574

1575   <u>Table S1: Murine brain databases used to map our unlabeled mouse dataset</u>

1576   We used four databases included in the scRNAseq v2.16.0(*34, 37, 66, 67, 70*) R package and

1577   the celldex v1.12.0 built-in mouse RNA-seq reference (which we called 'SingleR' database)

1578   using SingleR (*69*) function to transfer labels to our dataset in R(*62*). The name, brain area and

1579   reference in which each database was made public are specified.

1580

<u>Table S2: Literature-derived markers per central nervous system cell identity</u>

We thoroughly searched in the literature for established markers at the mRNA and/or protein

level. Some markers were found to be described for more than one cell population which are

specified in columns 'notes' and 'notes_2'. After initial labeling of our murine dataset with the

databases as described, we proceeded to obtain the average expression (on log-normalized

counts) of each gene coding for the markers for the labeled cells (i.e. astrocyte, endothelial,

microglia, neuron and oligodendrocyte) and unlabeled cells (i.e. unknown). We also calculated

the average expression for each cluster in our dataset, established during our processing

pipeline, at resolution 1.0.

1590

<u>Table S3: Literature-derived markers per central nervous system cell identity</u>

Breakdown of the cell identities (except neurons at layer 3 of cellular resolution) we established

and corroborating using a combination of markers surveyed in **Table S2** and additional markers

established using the FindConservedMarkers function from Seurat v5(*63*) package, indicating

a cell state derived from the literature. For example, 's_fibroblasts' were named after finding

high expression of *Stk39*, a stress-related marker(*35*).

1597

<u>Table S4: Expression of neurotransmitter-related genes in our murine neurons</u>

Expression of genes coding for neurotransmitter synthesis enzymes TH, DBH, TPH, GABA,

NOS, prohormone SST and primary neurotransmitter transporters VMAT2, VGLUT1, VGLUT2,

GLYT2 per neuron in our dataset. Expression and glutamate and GABA release are specified

as binary variables (i.e. 0=no expression; 1=expression) in the green and blue columns,

respectively. The percentile of the log-normalized expression is shown. ~55% of the neurons

display expression of genes related to the synthesis/release of >1 primary neurotransmitter.

1605

<u>Table S5: Description of our DVC layer-3 neuronal cell identities established in mice</u>

Neurons were subset and reprocessed before labeling at layer-3 granularity. The MAST

algorithm(*71*) was used through Seurat v5(*63*)on scaled data with a log fold-change threshold

of 0.25 and detection threshold on ≥40% of cells on each neuronal cluster at resolution 1.0. We

searched the resulting gene expression markers in the literature and compiled them. We then

based the names for our cell identities on these genes.

1649

1650   <u>Table S13: Labeled cells in our murine dataset</u>

1651   Number of cell per treatment per identity layer 2.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9