Morphological profiling reveals neuroprotection via mitochondrial uncoupling in human dopaminergic neurons

Short title: Neuroprotection via Mitochondrial Uncoupling

Vyron Gorgogietas[1,#], Amélie Weiss[2,#], Loïc Cousin[2], David Hoffmann[2], Karen Schmitt[2], Arnaud Ogier[2], Peter A. Barbuti[1,3,4], Bruno F.R. Santos[1,4], Ibrahim Boussaad[1], Annika Wittich[5], Andrea Zaliani[5], Ole Pless[5], Rejko Krüger[1,6,7,8], Peter Sommer[2], Johannes H. Wilbertz[2,*]

[1] Luxembourg Centre for Systems Biomedicine (LCSB), Esch-sur-Alzette, Luxembourg

[2] Ksilink, Strasbourg, France

[3] Department of Neurology, Columbia University Irving Medical Center, New York, NY 10032, USA

[4] Disease Modelling and Screening Platform, Esch-sur-Alzette, Luxembourg

[5] Fraunhofer Institute for Translational Medicine and Pharmacology ITMP, ScreeningPort, Hamburg, Germany

[6] Centre Hospitalier de Luxembourg (CHL), Luxembourg, Luxembourg

[7] Transversal Translational Medicine, Luxembourg Institute of Health (LIH), Strassen, Luxembourg

[8] Parkinson's Research Clinic, Centre Hospitalier de Luxembourg (CHL), Luxembourg, Luxembourg

[#] shared first author

* correspondence to: Johannes Wilbertz, Ksilink, 16 Rue d'Ankara, 67000 Strasbourg, France, Telephone: +33(0) 7 66 50 27 99, Email: johannes.wilbertz@ksilink.com

Abstract

Parkinson's disease (PD) involves multiple pathological processes in midbrain dopaminergic (mDA) neurons, including protein degradation defects, vesicular trafficking disruption, endolysosomal dysfunction, mitochondrial issues, and oxidative stress. Current PD models often lack complexity and focus on single phenotypes. We used patient-derived SNCA triplication (SNCA-4x) and isogenic control (SNCA-corr) mDA neurons, applying high-content imaging-based morphological profiling to identify and rescue multiple phenotypes. Screening 1,020 compounds, we identified top-scoring compounds that restored healthy profiles in SNCA-4x neurons, increasing Tyrosine hydroxylase (TH) and decreasing α-synuclein (αSyn) levels. Several hits were linked to mitochondrial biology. Tyrphostin A9, a mitochondrial uncoupler, and several of its structural analogues decreased ROS levels, normalized mitochondrial membrane potential, and increased respiration. Western blotting confirmed that Tyrphostin A9 reduces αSyn levels. Our study highlights the neuroprotective potential of mild mitochondrial uncoupling in mDA neurons.

Introduction

Parkinson's disease (PD), with its diverse phenotypic, neuropathological, and genotypic manifestations, is increasingly acknowledged as a complex disorder rather than a singular disease entity [1–3]. While traditional models have been instrumental in devising symptomatic therapies to address PD's motor symptoms, they have been less successful in developing neuroprotective strategies [4]. Genetic stratification of patients into subgroups has emerged as a crucial approach [5,6]. Over the past 25 years, significant progress has been made in identifying genes associated with PD, including those responsible for rare monogenic forms of the disease [2]. These monogenic forms have proven invaluable for PD research, as patient-based cell models exhibit disease-specific cellular phenotypes that mirror those found in postmortem brain tissue [7–11]. This approach suggests that targets for pharmaceutical interventions might be identifiable based on rare but potent molecular signatures [12–14].

Point mutations, as well as duplications and triplications of the wild-type alpha-synuclein gene (*SNCA*),

are responsible for causing autosomal dominantly inherited PD [15–18]. Patients with *SNCA* triplication

(SNCA-4x) express α-synuclein protein (αSyn) from four alleles resulting in a roughly 2-fold increase of

αSyn, and typically exhibit an early disease onset, accelerated disease progression, and significant

dementia. In contrast, patients with SNCA duplications often display a more conventional late-onset PD

phenotype. The severity of clinical trajectories in SNCA multiplication appears to correlate with gene

dosage. Patient-derived induced pluripotent stem cells (iPSC) carrying the *SNCA* triplication have been

successfully differentiated into midbrain dopaminergic (mDA) neurons and organoids [19–22]. Despite their

young age, iPSC mDA neurons exhibit PD-relevant phenotypes. *SNCA* gene triplication leads to an

increase in αSyn levels resulting in oligomeric αSyn pathology and elevated αSyn extracellular release

[19,21,23,24]. Multiple pathophysiological defects correlate with increased αSyn protein expression, such as

altered lysosomal and mitochondrial functions, dysregulated autophagy increased endoplasmic reticulum

stress, and heightened oxidative stress levels [20,22,24–28]. Due to the interconnection of many of the

identified phenotypes, it is currently an open question whether a rescue of one key or multiple cellular PD

hallmarks is necessary to achieve a significant neuroprotective effect. Imaging-based profiling is a

strategy that transforms the rich data in biological images into multidimensional profiles of extracted

features, which can be analyzed to reveal biological signatures, aiding in disease understanding and drug

discovery, with advancements in machine learning and computational technologies enhancing its potential

[29]. By capturing the observable characteristics of cells under different conditions, morphological profiling

can provide valuable insights into the complex cellular mechanisms underlying PD [8,9,11].

In this study, we find that *SNCA* gene triplication results in a multidimensional alteration of the

morphological profile in mDA neurons. We further identify multiple small molecular compounds

correcting this altered profile to align with the phenotype of isogenic controls (SNCA-corr) mDA

neurons. One of the identified compounds, Tyrphostin A9, exerts its neuroprotective effect via

mitochondrial uncoupling, which in turn reduces reactive oxygen species (ROS) and high molecular

weight αSyn species. In summary, we show that screening small molecules in a PD-relevant human model

system using a multi-phenotypical readout can identify modulators of not just a single, but multiple

disease relevant phenotypes.

Results

SNCA-4x mDA neurons have a distinct but modulable morphological profile

We differentiated patient-derived SNCA gene triplication and isogenic control iPSCs into mDA neurons

(SNCA-4x and SNCA-corr) for 30-37 days. These neurons showed the expected mDA neuron

morphology and no *de novo* copy number variations. Both cell lines expressed neuro-typical markers β-

tubulin III (90-94%), MAP2 (87-89%), and the dopaminergic neuron markers Tyrosine hydroxylase (TH)

(84-88%) and FOXA2 (84-90% in TH positive cells) (**Figure S1A-C**). Western blotting revealed

increased αSyn protein level and signs of αSyn aggregation in SNCA-4x mDA neurons after D70 (**Figure**

S1D). A JC-1 dye assay indicated reduced mitochondrial membrane potential in D35 SNCA-4x mDA

neurons (**Figure S1E**). For morphological profiling, mDA neurons were cultured until D36 in 384-well

plates and stained with Hoechst, and antibodies against TH, αSyn, and MAP2 (**Figure 1A**). Automated

segmentation and feature quantification were used to overcome challenges in assessing subtle phenotypes.

Image segmentation was performed on 16 fields per well and 127 morphological features were

determined and expressed as means per well. Features were calculated for the whole cell, nucleus,

cytoplasm or membrane and describe different aspects of the images (**Figure 1B**, **Table S4**). Texture

features capture the spatial distribution and arrangement of pixel intensities in an image. Context features

provide information about the spatial relationship between signals, for example the correlation between

image channels. Intensity features measure the fluorescence intensity and can provide insights into the

expression levels of proteins. Count and area/shape features quantify the number, size and morphology of

the cellular components. First, we examined basic morphological features of SNCA-4x and SNCA-corr

mDA neurons. SNCA-4x neurons had fewer live cells and approximately 50% increased αSyn protein

expression. In our study, we utilized Prostratin as a proof-of-concept molecule due to its proposed activity

in degrading αSyn. Previous research has demonstrated that specific phorbol esters, such as PEP005 or

Prostratin, can reduce αSyn through proteasomal degradation [7]. Based on these findings, we hypothesized

that Prostratin's activity would lead to an altered morphological profile in our experimental model.

Treatment with Prostratin indeed reduced dead cells and αSyn expression in SNCA-4x neurons to levels

similar to SNCA-corr neurons (**Figure 1C**). Neuronal morphological profiles before and after compound

treatment were created by scaling feature values and calculating mean values across wells (**Figure 1D**).

We used the mean absolute difference to find the features that differed the most between profiles.

Comparing SNCA-corr and SNCA-4x neurons, the radial αSyn intensity profile, which measures

fluorescence intensity as a function of distance from a central point, differed most. Prostratin-treated and

untreated SNCA-4x neurons also varied in number of nuclei, total MAP2 cellular surface, and αSyn

staining contrast (**Figure 1E**). These findings show that small molecular treatment can rescue

morphological features in mDA neurons which can be measured by morphological profiling.

Figure 1: SNCA-4x mDA neurons have a distinct but modulable morphological profile. (A) SNCA-4x and SNCA-corr mDA neurons were differentiated until D33, treated with 5 μM Prostratin or DMSO for 72 hours and immunofluorescently stained. Image channels were segmented for morphological feature quantification. (B)

Schematic overview of image data generation and morphological feature types. (C) Raw morphological feature values in control, and untreated and Prostratin treated SNCA-4x mDA neurons. Each datapoint represents technical replicate data from one 384-well plate well. Larger markers represent the medians each independent experiments. The boxplots show the median and the 2nd and 3rd quartiles of the data. Whiskers indicate 1.5X of the interquartile range. Welch's unequal variances t-test was performed using the medians of independent differentiation experiments. ns = not significant, * = p□<□0.05, ** = p□<□0.01, *** = p□<□0.001. (D) Scaled well and replicate median feature values aggregated into morphological profiles. Cosine similarity was used for clustering. (E) Top ten absolute mean differences between features of SNCA-4x and Control or SNCA-4x neurons treated with Prostratin. Means with standard deviations are shown. (F) Two first principal components (PCs) explaining most variation in the dataset after Principal Component Analysis (PCA). Each data point represents mean data from one well. (G) Absolute Pearson correlation of all features with the first ten PCs after PCA. Representative images illustrate cellular αSyn staining patterns. All data in this figure was generated in four independent experiments each containing at least 12 technical replicates per condition.

Next, we performed Principal Component Analysis (PCA) using the morphological profile data to visualize overall differences between compound-treated and untreated cells. The first two principal components (PCs) accounted for 57% of the variance and separated the tested cell lines and Prostratin-treated neurons, indicating significant differences between them (**Figure 1F**). The remaining PCs each accounted for 11% or less of the variance, suggesting lesser contribution to the overall variance (**Figure S1F**). To understand which features were mainly responsible for distinguishing the experimental categories, we calculated the absolute Pearson's correlation between the first ten PCs and each morphological feature. PC1 correlated mostly with αSyn signal texture-related features, indicating a different αSyn subcellular distribution pattern in the neurons. PC2 and PC3 correlated with signal radial distribution and nucleus-related features, similar to the results we obtained when comparing experimental conditions pairwise using the mean absolute difference (**Figure 1G**, **Figure S1G**). Manual inspection of images confirmed these findings, showing grainy αSyn localization outside the nucleus in SNCA-4x cells (**Figure 1G**). This indicated that radial distribution and texture features indeed capture αSyn localization

patterns. As a result, high-dimensional morphological profiling, by capturing a wide array of cellular

characteristics, allowed us to identify subtle differences between experimental conditions that may not be

apparent through visual inspection alone.

Compound modulation shifts a SNCA-4x towards a control morphological profile

Based on our observation that small molecular compounds, such as Prostratin, can partially rescue

SNCA-4x mDA neuron morphological profiles, we wondered whether more potent small molecules could

be identified using a morphological profile screening approach. We assembled a 1,020 compound library

which included 734 FDA-approved compounds, 435 of which can pass the blood-brain-barrier, 120

molecules targeting PD-relevant processes, and 166 diverse bioactive molecules (**Figure 2A**, **Table S3**).

Cryopreserved D30 mDA neurons were cultured in 384-well plates until D33, treated with 5 μM of each

compound for 72 hours and stained (**Figure 2B**). Each compound was tested in 4 replicates across 20

384-well plates, with 36 control wells per plate. Control conditions established baseline and maximum

effect profiles (SNCA-4x + DMSO and SNCA-corr + DMSO, respectively). Morphological profiles were

generated from 307,200 images and all data was aggregated per well (**Figure 2B**).

Figure 2: Compound modulation shifts a SNCA-4x towards a control morphological profile. (**A**) Overview about screening library composition and neuron treatment scheme. (**B**) Schematic overview of image data generation. Morphological features were calculated from imaging data to measure compound-induced morphological profile changes from a SNCA-4x towards a SNCA-corr-like profile. (**C**) Schematic of morphological

profile data processing workflow. **(D)** Overview of morphological feature types used for hit selection after removal

of redundant features using a Pearson correlation threshold of 0.95. **(E)** Morphological profile Mahalanobis

distances of compound treated mDA neurons relative to SNCA-4x and SNCA-corr morphological profiles. The

black dashed line indicates the selection window based on 4σ from the SNCA-4x median. Each datapoint represents

mean data from one well and for compounds the mean value of all four replicates is shown. **(F)** Compounds

inducing morphological profiles with the 10 shortest mean Mahalanobis distances to the median SNCA-corr

morphological profile. **(G)** Selected hits clustered by the Cosine similarity of their respective morphological

profiles. Compounds are color coded by attributed biological activity. (H) Tanimoto similarities of top 10 selected

compounds.

We aimed to identify compounds shifting SNCA-4x profiles towards SNCA-corr-like phenotypes.

Morphological profiles were Z-score normalized for comparability (**Figure 2C**, **Figure S2A**). Pearson's

correlation reduced the initial 127 features to 40 non-redundant ones, mostly describing cellular area,

shape, and TH and αSyn staining intensities (**Figure 2D**, **Figure S2B**). Plate quality was assessed using

the Strictly Standardized Mean Difference (SSMD) of the Mahalanobis distances between treatment

groups. All plates showed significant differences (**Figure S2C**). Linear Discriminant Analysis (LDA) was

used to compare each compound profile with the reference profiles. LDA achieved a classification

accuracy F1 score of 0.96 in a test data set and 0.93 ± 0.04 in a 10-fold cross-validation, indicating a high

ability to correctly predict whether compound-induced morphological profiles resemble one of the

reference classes (**Figure S2D**). Most compounds had no effect on the SNCA-4x profile, while some

induced SNCA-corr-like profiles (**Figure S2E**).

We aimed to identify compounds that could shift SNCA-4x profiles towards SNCA-corr-like phenotypes.

Morphological profiles were compared using the Mahalanobis distance, a metric that measures how

different two profile vectors are. We identified 28 compounds with a mean distance of at least 4 to the

SNCA-4x median profile (**Figure 2E**, **Table S1**). Of these, the 10 compounds with the shortest distance to

the SNCA-corr profile were considered to partially rescue the phenotype. Tyrphostin A9 and Alectinib

were the most similar to the SNCA-corr profile (**Figure 2F**). To validate our findings, we used Cosine

similarity, a metric that measures whether two vectors (or morphological profiles) are pointing in the

same direction. Nine out of the 10 compounds with the shortest Mahalanobis distance to SNCA-corr

grouped in the same cluster as SNCA-corr, supporting a distance and directional similarity between their

morphological profiles (**Figure 2F**, **Figure 2G**). Although the top hits' morphological profiles were

highly similar among each other and compared to the SNCA-corr profile, their chemical structure-based

Tanimoto similarities differed considerably (**Figure 2H**). We validated the effect of the identified

compounds on cellular phenotypes using data from the Joint Undertaking in Morphological Profiling

(JUMP) consortium, an effort to create compound-induced morphological profiles in U2OS cells by

CellPainting [30,31]. Also here, in a different cell type and using a different staining assay, we observed that

compounds with similar mode of action induced similar morphological profiles (**Figure S2F**). This

supports the notion that the identified compounds, while acting on different targets, might possibly acting

on related pathways leading to a similar phenotypic outcome. Tyrphostin A9 was originally developed as

an epidermal growth factor receptor (EGFR) inhibitor, but similar to other tyrphostins has been linked to

mitochondrial membrane fragmentation, reduced OXPHOS, and potentially neuroprotection via reduction

of ROS [32–35]. Since mitochondrial defects play an important role in PD pathology, we decided to further

characterize Tyrphostin A9's molecular mechanism in mDA neurons [3].

Tyrphostin A9 rescues the overall mDA neuron morphological profile in a dose dependent manner

To establish hit compound dose-response effects we focused on cell viability and αSyn level assessment

as well as on morphological profiling using six different compound concentrations ranging from 0.25-20

µM and a treatment duration of 72 hours (**Figure S3**, **Figure S4**). For Tyrphostin A9, we observed that

higher compound concentrations in SNCA-4x neurons resulted in a higher resemblance to SNCA-corr

neuron morphological features (**Figure 3A**). To quantify this observation, we calculated the Cosine

distance of the compound treated SNCA-4x morphological profile to the SNCA-4x + DMSO and SNCA-

corr morphological profiles as a function of compound concentration. Cosine distance can be defined as

1 – Cosine similarity. The intuition behind this is that if two vectors are perfectly the same then the

similarity is 1 (angle=0) and thus, the distance is 0 (1-1=0). We found that increasing compound

concentrations decreased the distance between SNCA-4x and SNCA-corr morphological profiles,

therefore increasing their similarity (**Figure 3B**). The opposite is true when SNCA-4x compound treated

neurons were compared to SNCA-4x DMSO-treated morphological profiles. Here, the Cosine distance

increased with increasing compound concentration, reflecting the change of phenotype towards the

SNCA-corr profile. Next, we investigated the effect of increasing compound concentrations on three

single morphological features: αSyn intensity, TH intensity and MAP2 neurite length (**Figure 3C**).

Tyrphostin A9 concentrations >1 µM decreased the nuclear and cytoplasmic αSyn staining intensity.

Tyrphostin A9 also positively impacted TH staining intensity. At higher concentrations we observed little

effects on MAP2 neurite length, indicating no acute toxicity.

215
216 **Figure 3: Tyrphostin A9 rescues the overall morphological profile in a dose dependent manner. (A)** Line plot

representation of morphological profiles of Tyrphostin A9 treated and untreated mDA neurons. D33 neurons were treated with 0.25-20 μM compound for 72 hours before fixation. **(B)** Cosine distances of SNCA-4x neurons as a function of compound concentration relative to DMSO-treated SNCA-4x and SNCA-corr neurons. **(C)** Dose-response curves for the three selected morphological features αSyn intensity, TH intensity, and MAP neurite length. Representative images of the same cell in each fluorescent channel are shown for DMSO and 5 μM compound treatment. Each compound concentration was tested at least in triplicate. Mean and 95% confidence interval are shown.

Multiple identified hits induce mitochondrial uncoupling and ROS reduction

Tyrphostin A9 has been explored *in vitro* for its roles as neuroprotective agent, possibly functioning as an oxidative phosphorylation uncoupler reducing ROS [32,33,36]. Mitochondrial uncoupling involves the decrease of the mitochondrial proton gradient disrupting the coupling between the electron transport chain and ATP synthesis. Whether Tyrphostin A9 can exert these effects in mDA neurons is currently unclear. We first assessed Tyrphostin A9's ability to modulate the mitochondrial membrane potential using the JC-1 dye. We treated D33 mDA neurons for 72 hours with increasing concentrations of Tyrphostin A9 (0.01 μM to 10 μM) and observed a decrease in red fluorescence for concentrations >0.5 μM, indicating a concentration dependent decrease in mitochondrial membrane potential (**Figure 4A**). Valinomycin was used as positive control inducing a very low membrane potential. While FCCP and Tyrphostin A9 lowered the mitochondrial membrane potential at the lowest doses, also other molecules with morphological profiles similar to Tyrphostin A9, such as Tinostamustine or VPS34 inhibitor 1, led to a decreased mitochondrial membrane potential (**Figure S5**). Since a reduction in mitochondrial membrane potential is often associated with decreased ROS levels, we used a dye-based ROS assay to measure ROS levels in mDA neurons upon compound exposure. Tyrphostin A9, at concentrations >0.5 μM reduced ROS down to the level of the positive control antioxidant N-acetyl-cysteine (NAC) (**Figure 4B**). Interestingly, also multiple other molecules with morphological profiles similar to Tyrphostin A9, such as Simvastatin, Trifluoperazine, or WAY-600 lowered neuronal ROS levels (**Figure 2G**, **Figure S6**). We hypothesized that Tyrphostin A9 might lower mitochondrial membrane potential and ROS due to its

function as protonophore facilitating proton transfer across membranes. We used a Rhod-FF assay which

measures intracellular Ca^{2+} levels and found only small baseline differences between SNCA-corr and

SNCA-4x mDA neurons, while the addition of Tyrphostin A9 to SNCA-4x mDA neurons increased

intracellular Ca^{2+} levels by more than 70% (**Figure 4C**). Addition of the known protonophore Ionomycin

further increased intracellular Ca^{2+} levels demonstrating that Tyrphostin A9 likely uncouples the

mitochondrial membrane potential via facilitating the entry of protons, thereby decreasing oxidative

phosphorylation and ROS levels. In neurons protonophores are also expected to disrupt the neuronal

membrane potential by decreasing the proton gradient, leading to a decrease in the driving force for the

generation of action potentials, which might negatively impact neuronal firing. Using a microelectrode

array (MEA) we measured the neuronal firing rate before and after Tyrphostin A9 treatment (**Figure 4D**).

Before treatment ca. 60-day old SNCA-corr mDA neurons showed approximately 2.5-fold higher firing

rates than SNCA-4x neurons. When treating the mDA neurons for up to 72 hours with 2.5 μM Tyrphostin

A9 we observed a sharp drop in firing already 30 minutes after treatment. After >24 hours of treatment a

slight recovery effect was observed and firing rates began to increase mildly. Taken together, our findings

suggest that the identified compounds, although affecting different targets, might be influencing related

pathways, resulting in a similar phenotypic outcome such as mitochondrial membrane potential and ROS

lowering.

260
Figure 4: Tyrphostin A9 acts as ROS-reducing mitochondrial uncoupler in mDA neurons. (A) mDA neurons

were treated with 0.1-10 μM Tyrphostin A9 for 72 hours and the mitochondrial membrane potential was measured

using the JC1 dye. JC1 fluoresces red at high membrane potential and green at lower membrane potentials. Three

independent experiments with at least three technical replicates each were performed. **(B)** mDA neurons were

treated with 0.5, 1, and 2.5 μM Tyrphostin A9 for 72 hours and the ROS level was determined using the CellROX

dye. The obtained signal was normalized to total protein content. Three independent experiments with at least three

technical replicates each were performed. **(C)** *Left panel:* Scheme illustrating the protonophore mode of action.

Right panel: mDA neurons were treated with 0.5, 1, and 2.5 μM Tyrphostin A9 for 72 hours and the intracellular

Calcium level was measured using the Rhod-FF dye for 250 seconds before and after the addition of Ionomycin. The

obtained signal was normalized to total protein content. Two independent experiments with at least three technical

replicates each were performed. **(D)** Neuronal activity was measured in ca. 60-day old mDA neurons using a

microelectrode array (MEA). The neuronal firing rate was assessed before and up to 72 hours after treatment with

2.5 μM Tyrphostin A9. Neuronal activity was measured in two independent experiments with at least three technical

replicates each. For all data, the mean and 95% confidence interval is shown.

275 **The Tyrphostin A9 structural analogue AG879 induces mitochondrial uncoupling**

276 Tyrphostins represent a large family of small molecules with various attributed neuroprotective modes of

277 action. Next to their role as mitochondrial uncouplers, tyrphostins have been described as antioxidants or

278 as having a positive effect on cellular glutathione levels [33]. We therefore compared a range of Tyrphostin

279 A9 structural analogues using various biochemical assays to establish a Tyrphostin A9 structure-function

280 relationship in mDA neurons (**Figure 5A**). FCCP is a potent mitochondrial uncoupler used during oxygen

281 consumption rate (OCR) measurements and facilitates proton transport across mitochondrial membranes.

282 This action collapses the proton gradient and disrupts the mitochondrial membrane potential. As a result,

283 the electron flow through the Electron Transport Chain (ETC) is uninhibited, allowing oxygen

284 consumption by complex IV to reach its maximum. The resulting OCR increase is used as an estimation

285 of the maximal respiratory capacity of the cell (**Figure 5B**, left panel). Next to its use as a control

286 compound, we also detected FCCP as one of the top hits during morphological profiling (**Figure 2F**). We

287 therefore compared Tyrphostin A9 and its structural analogues to FCCP using an OCR assay. We found

288 that, similar to FCCP, Tyrphostin A9 and its closest structural analogues AG1024 and AG879 led to a

289 strong increase in maximal respiratory capacity, highlighting their mitochondrial uncoupling activity

290 (**Figure 5B**, right panel). Next, we determined compound cytotoxicity by measuring Lactate

291 dehydrogenase (LDH) in the culture medium. LDH is a stable enzyme, present in all cell types and

292 rapidly released into the culture medium upon damage of the plasma membrane. None of the tested

293 compounds induced significantly more cell death than the DMSO control when used at 0.5-2.5 μM for 72

294 hours. Cell lysate was used as control corresponding to the maximum possible amount of released LDH

295 (**Figure 5C**). JC1 dye fluorescence was measured to determine the mitochondrial membrane potential.

296 Tyrphostin A9 and its most closely related structural analogue Tyrphostin AG879 decreased the

297 membrane potential as measured by the 547nm/488nm fluorescence ratio from 0.95 to approximately 0.2.

298 The second most similar analogue Tyrphostin AG1024 also decreased the membrane potential, although

299 to a lesser extent (**Figure 5D**). In line with the decreased membrane potential Tyrphostins A9 and AG879

also reduced ROS levels to ca. 80% of the DMSO baseline level (**Figure 5E**). Next, we assessed

glutathione levels. In healthy cells, both reduced glutathione (GSH) and its oxidized form (GSSG) are

present, with the reduced form (GSH) making up the majority, while the oxidized form (GSSG) increases

during oxidative stress. Although we expected an GSH increase due to ROS reduction, we observed that

Tyrphostins A9, AG879 and AG1024 decreased GSH ratio, possibly due to lower overall glutathione

synthesis after ROS decrease. Tyrphostin A9 lead to the strongest decrease of 48% at 2.5 μM (**Figure 5F**).

Mitochondrial uncoupling also strongly reduces cellular ATP production. We therefore tested the effects

of all compounds on mDA neuron ATP level. We found that the compounds with the strongest uncoupling

effects (Tyrphostins A9, AG1024, and AG879) reduced ATP levels the most (**Figure S7A**). mDA neuron

treatment with the mitochondrial complex I inhibitor Rotenone led to similarly low ATP levels as

Tyrphostin A9 treatment alone, indicating a near maximal Tyrphostin A9 uncoupling effect (**Figure S7B**).

Since close Tyrphostin A9 structural analogues such as AG879 and AG1024 reproduce its cellular effects,

the shared structural features may be critical for the molecule's function in mDA neurons.

Figure 5: Tyrphostin A9 structural analogues reduce mitochondrial membrane potential and ROS. (A) Molecular structures of Tyrphostin A9 and five selected structural analogues including their names, Compound ID (CID), and Tanimoto similarity coefficient relative to Tyrphostin A9. **(B)** Oxygen consumption rate (OCR), a proxy readout for mitochondrial respiration, was measured in D30 mDA neurons treated with 2.5 μM compound. (C-F)

D33 mDA neurons were treated with 0.5-2.5 μM of each compound for 72 hours. **(C)** Lactate dehydrogenase (LDH)

activity was measured in the culture medium to determine relative cytotoxicity compared to cellular lysate (=100%).

(D) Mitochondrial membrane potential was measured using the JC-1 dye. **(E)** ROS level was determined using the

ROS-ID detection kit and the obtained signal was normalized to total protein content. **(F)** Reduced glutathione

(GSH) was measured using monochlorobimane (MCB) and the obtained signal was normalized to total protein

content. Three independent experiments with at least three technical replicates were performed. The mean and 95%

confidence interval are shown. Dashed lines indicate the median values of the respective controls.

Tyrphostin A9 induces αSyn lowering

Our phenotypic compound screen revealed that mDA neuron αSyn levels were lowered by Tyrphostin A9

treatment (**Figure 3C**). Since immunofluorescent staining is not sensitive enough to differentiate between

αSyn monomers, oligomers, and larger structures such as fibrils, we performed Western blotting of

Tyrphostin A9 treated and untreated >D50 mDA neuron lysates (**Figure 6A**). We observed a 2-fold

reduction of both low molecular weight αSyn (≈15 kDa) as well as higher molecular weight αSyn species.

Most cellular proteins are selectively degraded by the proteasome via ubiquitination. Caspase and

chymotrypsin are part of the proteasomal protein degradation machinery, contributing to the proteolytic

activities of the proteasome. Using plate reader assays we therefore checked how caspase and

chymotrypsin activities are impacted by Tyrphostin A9 in mDA neurons. Both enzymes' activities were

not significantly different between SNCA-corr and SNCA-4x mDA neurons, while addition of Tyrphostin

A9 rather decreased (≈ 50%) or did not impact caspase- or chymotrypsin-related proteasomal degradation,

respectively (**Figure 6B**). As expected, the known proteasome inhibitor MG132 strongly decreased both

caspase and chymotrypsin activities in mDA neurons by >70%. Since Tyrphostin A9 does not positively

impact proteasomal protein degradation in mDA neurons, the observed lower αSyn levels after compound

treatment might result from an increased alternative protein degradation pathway.

Figure 6: Tyrphostin A9 decreases αSyn protein expression. (A) Anti-αSyn Western blotting of mDA neuron lysates. Before lysis >D50 mDA neurons were treated with 0.3 or 2.5 μM Tyrphostin A9 for 72 hours. Three independent neuronal differentiations were performed. Gel images are cropped. Full-length blots/gels are presented in Figure S8. **(B)** Caspase and Chemtrypsin activities were determined in approximately D30 mDA neurons treated with 1 or 2.5 μM Tyrphostin A9 or the proteasome inhibitor MG132 for 72 hours. Three independent experiments with at least three technical replicates each were performed. The boxplots show the median and the 2nd and 3rd quartiles of the data. Whiskers indicate 1.5X of the interquartile range. Welch's unequal variances t-test was performed using the technical replicates of all independent experiments. ns = not significant, * = p□<□0.05, ** = p□<□0.01, *** = p□<□0.001. **(C)** Scheme depicting the use of morphological profiling to identify compounds that

351 produce a healthy control-like (SNCA-corr) profile in mutant (SNCA-4x) mDA neurons. The identified compound

352 Tyrphostin A9 increases mitochondrial uncoupling and decreases ROS and αSyn levels which can explain the

353 rescued morphological profile.

Discussion

We used microscopy-based morphological profiling to characterize the effects of 1,020 FDA-approved or

PD pathobiology-related small molecules on the overall morphological appearance of human iPSC-

derived SNCA-4x mDA neurons. We scored compounds based on their potential to induce a

morphological profile resembling SNCA-corr mDA neurons (**Figure 2**). Phenotypic or morphological

profiling is increasingly used in iPSC-derived cells, primary cells or brain slice systems to study neuronal

diversity and to complement single cell RNA sequencing data [37–40]. Previously, we established

morphological profiling in LRRK2 G2019S mutated iPSC-derived mDA neurons and showed that

multiple LRRK2 inhibitors can partially rescue the neuronal morphological profile [8]. Drug screens in

mDA neuron model systems have been performed before, however either using only small screening

libraries, non-human cell systems such as zebrafish, or very young mDA neuron progenitors (<20 days of

differentiation) [41–44]. This study combines detailed morphological profiling in human mDA neurons with a

compound screen using a larger library (>1,000 compounds). Although morphological profiling has the

potential to identify completely novel targets, a limitation is that it typically provides little information

about the actual potential compound targets, requiring detailed target and mode of action deconvolution

studies.

In our study, we focused on mitochondrial uncoupling since two of our top scoring compounds (FCCP,

Tyrphostin A9) are annotated as protonophores and mitochondrial uncouplers. Originally developed as an

EGFR inhibitor, Tyrphostin A9 stood out due to its αSyn reducing and TH increasing effects in our

morphological profiling assay as well as the molecule's previously described role in neuroprotection

(**Figure 3**) [33–35]. We found that the molecule promotes mitochondrial uncoupling and reduces ROS in

mDA neurons in line with its previously described protonophore properties in other cell types (**Figure 4**)

[33,36,45]. Tyrphostins have been described as mitochondrial uncouplers for almost half a century, while

mitochondrial uncoupling as a neuroprotective strategy is more recent, but has faced several safety-related

hurdles during pre-clinical drug development [46–49]. In mDA neurons we made the observation that next to

its mitochondrial uncoupling effect, Tyrphostin A9 also decreased neuronal activity which relies on

charge-dependent action potentials (**Figure 4D**). Tyrphostin A9 did not permanently damage the neurons,

indicated by the gradual partial activity recovery over time, intact morphology, and low extracellular LDH

levels (**Figure 3**, **Figure 4D, Figure 5C**). Given that pace making is a critical physiological function of

mDA neurons in the *substantia nigra*, this observation raises concerns about potential detrimental effects

of Tyrphostin A9. Importantly, we do not claim that Tyrphostin A9 itself holds therapeutic potential.

However, here we show that morphological profiling in a PD-relevant human model system is able to

detect compound classes with potentially undervalued neuroprotective modes of action. Future research

should address the safety of mitochondrial uncoupling as a neuroprotective strategy and explore the link

between Tyrphostin A9-induced EGFR inhibition and neuroprotection. Recently the blocking of

mitochondrial complex I in pro-inflammatory microglia has been shown to protect the central nervous

system against neurotoxic damage in mice [50]. An attractive future neuroprotective therapeutic strategy

involving mitochondrial uncouplers might therefore involve non-cell autonomous mechanisms. The

molecular characterization of non-cell autonomous mechanisms in neurodegeneration is becoming

increasingly feasible by using more complex iPSC-derived model system involving multiple central

nervous system cell types in the form of co-culture experiments or organoids [51–53].

A surprising observation we made is Tyrphostin A9's ability to reduce the low and high molecular weight

αSyn levels in mDA neurons by about 50% (**Figure 3C**, **Figure 6**). We excluded a direct effect on caspase

or chymotrypsin related proteasomal degradation mechanisms since Tyrphostin A9 either did not impact

or even decreased these modes of protein degradation. Except for one study describing a role of

Tyrphostin AG1024 in accelerating the degradation of Retinoblastoma Protein (pRb) likely involving the

proteasome, tyrphostins have not been directly linked to enhanced protein degradation [54]. Next to

proteasomal protein degradation, the involvement of lysosomes in Parkinson's disease progression through two major autophagic pathways, chaperone-mediated autophagy and macroautophagy, and their central role in degrading αSyn is well documented [2]. A single study demonstrated that Tyrphostin AG490, via JAK2 inhibition, prevented the hormone-receptor complex Epo-EpoR from targeting lysosomes in the UT-7 human cell line, highlighting a possible link between tyrphostin action and lysosomal biology [55]. Although we did not elucidate the exact sequence of events in our mDA neuron model, oxidative stress can cause proteins to misfold, leading to proteostatic stress. Conversely, misfolded proteins can disrupt mitochondrial function, increasing ROS production and oxidative stress, leading to dopamine oxidation [3]. αSyn could play a key role in the harmful effects of oxidized dopamine. It has been shown that oxidized dopamine can alter and stabilize αSyn protofibrils *in vitro*, and an increased expression of αSyn in SH-SY5Y cells heightened their susceptibility to an excess of dopamine [56,57]. In PD patient-derived iPSC neurons, a rise in oxidized dopamine and a reduction in αSyn solubility has been observed [26]. Moreover, mitochondrial uncouplers can induce mitophagy which in turn could decrease the α-synuclein that is accumulated and aggregates in mitochondria [58–60]. Future work will need to establish the molecular connection between mitochondrial uncoupling and the increased αSyn degradation which we observed.

Material and Methods

Origin of iPSC lines and differentiation into mDA neurons

SNCA iPSC lines (originally named AST23 and AST23-2KO) were provided and generated by Dr. Tilo Kunath from The University of Edinburgh. The parental cell line AST23 was first described in Devine *et al.* [19]. The generation of the SNCA-corr cell line AST23-2KO is described in Chen *et al.* [61]. Both cell lines are deposited as EDi001-A and EDi001-A-5 in the hPSCreg repository, respectively (https://hpscreg.eu). iPSCs were cultured on Geltrex-coated (Thermo Fisher Scientific) dishes in StemMACS iPS-Brew XF (Miltenyi Biotech). The medium was changed daily, and cells were passaged twice a week using 0.5 mM EDTA in PBS (Thermo Fisher Scientific). Mycoplasma testing was performed twice per month. mDA

neurons were differentiated using a modified protocol based on Kriks *et al.* [62–64]. In summary, iPSCs were

initially placed on Geltrex-coated six-well plates or T75 flasks at a density of $2x10^5$ cells/cm2 in

StemMACS iPS-Brew XF with 10 μM Y-27632 (Hiss). The following day, the medium was changed to

KnockOut DMEM medium with KnockOut serum replacement (both from Thermo Fisher Scientific),

supplemented with 200 nM LDN19318 (Axon Medchem) and 10 μM SB431542 (Biozol) for dual

SMAD-inhibition. On the second day, 100 ng/mL Shh C24II (Miltenyi Biotech), 2 μM Purmorphamine

(Miltenyi Biotec), 100 ng/mL FGF8 (Peprotech), and 3 μM CHIR99021 (Miltenyi Biotec) were also

added to the medium. After a period of 5 days, the medium was gradually transitioned to Neurobasal

medium (Thermo Fisher Scientific), and SB431542 was removed from the medium. Starting from day 7,

cells were cultivated only in the presence of LDN19318 and CHIR99021. On day 11, cells were

transferred to Neurobasal/B27/L-glutamine medium with only CHIR99021. On day 13, cells were moved

onto Geltrex-coated dishes in Neurobasal/B27/L-glutamine medium with supplements of 20 ng/mL

BDNF, 20 ng/mL GDNF (both from Cell Guidance Sys.), 220 μM L-ascorbic-acid (Sigma-Aldrich), 10

μM DAPT (Axon Medchem), 1 ng/mL TGF-βIII (Peprotech), 0.5 mM dibutyryl-cAMP (Enzo Life

Sciences), and 10 μM Y-27632 (Hiss). Cells were kept in the same medium but without Y-27632. Around

days 23–25, cells were separated using StemPro Accutase (Thermo Fisher Scientific) and placed at a

density of $1.4x10^5$ cells/cm2 onto Geltrex-coated dishes. To remove non-neuronal cells, cultures were

exposed to 1 μg/mL Mitomycin C for 2 hours on day 26. On day 30, neuronal cultures were separated

using StemPro Accutase with 10 μM Y-27632 and singularized. Cells were counted and preserved at

$2.5x10^6$ cells/vial in CryoStor CS 10 tubes (Sigma-Aldrich).

Neuronal culture

30 days old mDA neurons were thawed and spun (400 g, 5 min, RT) in a basal medium (**Table S2**)

supplemented with a ROCK inhibitor (Tocris #1254). The cell pellets were then reconstituted in a

differentiation medium (**Table S2**) also supplemented with a ROCK inhibitor. 384-well plates (Perkin

Elmer, Phenoplate poly-D-lysine-coated #6057500) were prepared with a coating of 10 μg/ml laminin at

4°C overnight. Using trypan blue (Invitrogen, #T10282) and a Countess automated cell counter

(Invitrogen), cells were seeded in the 384-well plates at a density of 10×10^3 cells/well. Edge wells were

not used for seeding and were filled with PBS instead. Typically, the thawed cells were incubated at 37°C

and 5% CO2 for 6 days until they reached day 36, with changes in the differentiation medium occurring

at D33. The processes of plate coating, cell seeding, and medium changes were automated using an Bravo

Automated Liquid Handling Platform with 384ST head (Agilent) and a Multidrop Combi Dispenser

(ThermoFischer Scientific).

Compound library and compound treatment

1,020 small molecules were selected from a bioactive compound library (Selleckchem), based on PD-

relevant pathobiology, FDA approval status, and blood-brain-barrier penetrance (**Table S3**). *Prostratin*

treatment: Neurons were cultured until D33, treated with 5 μM Prostratin (Biotechne) or DMSO for 72

hours and immunofluorescently stained. *Primary screening:* Cryopreserved mDA D30 neurons were

cultured in 384-well plates until D33, treated with each compound at a concentration of 5 μM for a

duration of 72 hours, and subsequently stained. Each compound underwent testing in four technical

replicates. Compound wells were randomly assigned between technical replicates. 36 wells per plate were

designated as control wells. The control conditions were used to establish the baseline and maximum

effect profiles, specifically with SNCA-4x + DMSO serving as the baseline and SNCA-corr + DMSO as

the maximum effect profile. Compounds and DMSO were dispensed using a Bravo Automated Liquid

Handling Platform with 384ST head (Agilent). *Dose-response testing:* Identical to primary screening but

with six different compound concentrations ranging from 0.25-20 μM. *Biochemical assays:* >D30

neurons were treated with 0.5-2.5 μM compound for 72 hours. *Microelectrode array (MEA):* Ca. D60-old

mDA neurons were treated up to 72 hours with 2.5 μM Tyrphostin A9. *Western blotting:* >D50 mDA

neurons were treated with 0.3 and 2.5 μM Tyrphostin A9 for 5 days.

Immunofluorescent staining

Neurons were fixed using 4% PFA (EMS Euromedex, #15710) for a duration of 20 minutes, followed by

three PBS washes (Gibco, #14190136). The cells were then permeabilized and blocked using a solution of

10% FBS (Sigma, #F7524) and 0.1% Triton X-100 (Sigma, #T9284) in PBS for an hour. Primary

antibodies against TH, αSyn, and MAP2 (**Table S2**) were prepared in an antibody dilution buffer (PBS

with 5% FBS and 0.1% Triton X-100) and left to incubate with the cells at 4°C overnight. This was

followed by three PBS washes. Secondary antibodies and Hoechst (**Table S2**) were added to the cells in

the antibody dilution buffer and left for an hour at room temperature, followed by another three PBS

washes. Immunofluorescent staining was automated using an EL406 plate washer and dispenser (Biotek).

Image acquisition and feature extraction

Imaging was performed on a Yokogawa CV7000 microscope in scanning confocal mode using a dual

Nipkow disk and a cooled sCMOS camera with 2,560×2,160 pixels and a pixel size of 6.5 µm without

pixel binning. The system's CellVoyager software, the 405/488/561/640-nm solid laser lines, and a dry

40X objective were used to acquire 16 images in a 4×4 orientation from three 1 µm-separated Z-planes

from the center of each well. Wells were visited in a row-wise "zigzag" fashion. Maximum intensity

projections were performed. Image segmentation and feature extraction was performed with our in-house

software PhenoLink (https://github.com/Ksilink/PhenoLink). Image segmentation was performed on 4x4

stitched and illumination corrected images. Using intensity and size thresholds specific to each of the four

imaging channels (Hoechst, TH, αSyn, and MAP2) we extracted a total of 124 morphological features,

including count, texture, context, intensity and shape features (**Table S4**). We used location constraints to

identify subcellular compartments. For example, by subtracting the segmented nucleus from the whole

cell we obtained the cytoplasm or by calculating a 4 pixel-wide area around the outer edge of the cell we

estimated the location of the cell membrane.

Data processing & morphological profile generation

All data manipulation and transformation processes were carried out using Python's Pandas package [65].

All plots were generated using the Seaborn package [66]. All data processing and plotting can be reproduced

using the provided Python Jupyter Notebooks and accompanying raw data. Briefly, raw morphological

profiles were created per well by using the information from all measured 127 morphological features

(**Table S4**). First, per feature a mean feature value from all acquired 16 image fields per well was

calculated.   The   feature   values   were   subjected   to   a   standard   scaling   process   using

the StandardScaler function from the sklearn.preprocessing module in Python, resulting in the creation of

127-feature raw morphological profiles per well [67]. To determine the most defining morphological

features per treatment condition, the morphological profiles were grouped based on their experimental

replicates and treatments. The absolute differences in the mean values of the grouped data were calculated

and followed by the computation of the mean and standard deviation of these absolute differences. We

employed Principal Component Analysis (PCA) to compare morphological feature profiles across

biological conditions and treatments. PCA was performed with 10 components and the explained variance

ratio for each of the principal components was calculated. To measure how each original feature

contributed to the principal components, the correlation between the original features and the first 10

principal components was calculated.

Primary screening data processing and hit selection

The raw data was composed out of 307,200 images from 4,800 wells distributed over 20 384-well plates.

For all fluorescent channel images (Hoechst, TH, αSyn, and MAP2) we performed image segmentation

and feature extraction, giving rise to a raw 127-feature morphological profile. Per plate, extreme outlier

features were eliminated using a robust standard deviation-based method. This involved calculating the

absolute difference between each data point and the feature median, and retaining data that was within 4σ

of the Median Absolute Deviation (MAD). Next, data was normalized to allow comparisons between

plates. We applied a robust Z-score normalization, during which the original data was adjusted based on

its deviation from the median of the DMSO control data and was scaled by the MAD of the DMSO

control data. This method is robust to outliers as it uses median and MAD, which are less sensitive to

extreme values. Feature reduction was performed in two separate steps. Since the morphological profile

contains an overrepresentation of texture features, first the two most distinguishing texture features were

selected by calculating the z-factors for each feature, sorting them in descending order, and picking the

top two. In the second step, features with less than 95% pairwise Pearson correlation were retained. In

total, 40 features remained (**Figure S2B**). Plate quality was assessed using the Strictly Standardized Mean

Difference (SSMD) based on the 40-feature Mahalanobis distance of SNCA-4x + Prostratin-treated

control wells to SNCA-4x + DMSO control wells. SSMD values for all 20 plates were <-1, indicating a

good plate quality. We then calculated the 40-feature Mahalanobis distances of all SNCA-4x compound-

treated wells to the SNCA-4x + DMSO and SNCA-corr + DMSO control wells. We applied a

Mahalanobis distance threshold of >4 to identify compounds inducing morphological profiles different

from SNCA-4x + DMSO control profiles. Additionally, we selected compounds based on close

Mahalanobis distances to SNCA-corr + DMSO control profiles, thereby selecting compounds that induce

morphological profiles similar to healthy control neurons. All data processing and hit selection steps can

be reproduced by using the provided Python Jupyter Notebooks.

Evaluation of mitochondrial membrane potential

Mitochondrial membrane potential ($\Delta\psi$m) was estimated using the fluorescent probe JC-1 (ThermoFisher,

T3168). Briefly, 60×10^3 mDA neurons were seeded per well in a Geltrex-coated black, clear-bottom 96-

well cell culture plate and treated with selected compounds for three days. On neuronal maturation day

33, the neuronal medium was removed and cells were incubated with 100μL/well of fresh neuronal

medium containing 10 μM of JC-1 dye for 30 minutes at 37°C. After incubation, cells were washed three

times with pre-warmed DPBS (ThermoFisher, #14190094). Fluorescence was measured using a Cytation

5 (Bio-Tek) fluorescence plate reader, with excitation at 560 nm and emission at 610 nm for the

aggregated JC-1 and excitation at 500 nm and emission at 550 nm for the monomeric JC-1. Mitochondrial

membrane potential was estimated by calculating the ratio of aggregated to monomeric JC-1 fluorescent

signal. Statistical analyses were conducted on data from, at least, three independent biological replicates.

Measurement of intracellular ROS and glutathione levels

Intracellular ROS levels were measured using the ROS-ID Total ROS/Superoxide detection kit (Enzo,

Enz-51010), following the manufacturer's instructions. Briefly, 60×10^3 mDA neurons were seeded per

well in a Geltrex-coated black, clear bottom 96-well cell culture plate. Neurons were treated for three

days with the selected compounds. On neuronal maturation day 33, the neuronal medium was removed

and the cells were washed once with pre-warmed DPBS (ThermoFisher, #14190094). Cells were then

incubated with 100 μL/well of ROS Detection Solution for 1 hour at 37°C. Finally, fluorescence is

measured without removing the detection solution, using a Cytation 5 (Bio-Tek) fluorescence plate reader

(bottom reading at Ex=490 nm, Em=520 nm). Fluorescence intensity was normalized to total cell protein

levels, using the BCA Protein Detection Kit (ThermoFisher, #23227). Statistical analyses were conducted

on data from three independent biological replicates. Simultaneously, intracellular GSH levels were

measured using the monochlorobimane (MCB, AAT Bioquest, #76421-73-3) assay. In this procedure,

60×10^3 mDA neurons were seeded per well in a Geltrex-coated black, clear-bottom 96-well cell culture

plate and treated with selected compounds for three days. On neuronal maturation day 33, the neuronal

medium was removed, and the cells were washed once with pre-warmed DPBS (ThermoFisher,

#14190094). The cells were then incubated with 100 μL per well of a detection solution containing 50 μM

MCB for 1 hour a, with excitation at 360 nm and emission at 480 nm t 37°C. After incubation,

fluorescence was measured directly without removing the detection solution using a Cytation 5 (Bio-Tek)

fluorescence plate reader, with excitation at 360 nm and emission at 480 nm. The fluorescence intensity

readings were normalized to total cell protein levels, which were determined using the BCA Protein

Detection Kit (ThermoFisher, #23227). Statistical analyses were conducted on data from three

independent biological replicates.

Evaluation of cytotoxicity

Cytotoxicity was estimated using the lactate dehydrogenase (LDH) assay (Canvax, CA0020), following

the manufacturer's instructions. 60×10^3 mDA neurons were seeded per well in a Geltrex-coated black,

clear bottom 96-well cell culture plate. Neurons were treated for three days with the selected compounds.

On neuronal maturation day 33, 50 µL of media was obtained from each well and transferred to a 96-well

cell culture plate, in order to evaluate the release of LDH into the cell culture media. It was added to an

equal volume (50 µL) of mixed LSH assay buffer and dye solution and was incubated for 30 minutes, at

room temperature, protected from light. Finally, 50 µL Stop Solution was added per well and the

absorbance (OD) was measured using a Cytation 5 (Bio-Tek) plate reader (450 nm). LDH absorbance was

normalized to total cell protein levels, using the BCA Protein Detection Kit (ThermoFisher, #23227).

Statistical analyses were conducted on data from three independent biological replicates.

Measurement of intracellular ATP levels

Intracellular ATP levels were evaluated using the CellTiter-Glo 2.0 (Promega, G9241), according to the

product's instructions. Briefly, 70×10^3 mDA neurons were seeded per well in a Geltrex-coated black,

clear-bottom 96-well cell culture plate and treated with selected drugs or compounds for three days. On

neuronal maturation day 33, the neuronal medium was removed, and cells were washed once pre-warmed

DPBS (ThermoFisher, #14190094). Cells were incubated with 75 µL per well of HBSS and an equal

volume of the CellTiter-Glo 2.0 mix was added in each well. The plates were shaken for 2 minutes in RT,

followed by 10 minutes of incubation in RT. Luminescence was measured using a Cytation 5 (Bio-Tek)

plate reader. Luminescence intensity was normalized to total cell protein levels, using the BCA Protein

Detection Kit (ThermoFisher, #23227). Statistical analyses were conducted on data from three

independent biological replicates.

Analysis of mitochondrial respiration

Oxygen consumption rates (OCRs) were measured using the Seahorse XFe96 FluxPak (Agilent,

#102416-100) and the XF Cell Mito Stress Test Kit (Agilent, #103015-100), following the manufacturer's

instructions. For this experiment, 70×10^3 mDA neurons were seeded per well in a Geltrex-coated 96-well

Seahorse cell culture plate. On neuronal maturation day 33, the neuronal medium in each well was

replaced with 175 µl of Seahorse XF DMEM Medium (pH 7.4, Agilent, #103575-100) containing 25 mM

Glucose, 2 mM L-Glutamine, and 1 mM Sodium Pyruvate. The cells were incubated for 1 hour at 37°C

without CO2. Meanwhile, the sensor cartridge, pre-equilibrated overnight in XF Calibrant solution at

37°C without CO2, was prepared with standard mitochondrial inhibitors: Oligomycin (2.5 µM), FCCP or

tyrphostin analogs (2.5 µM), and a mixture of Rotenone and Antimycin A (1 µM each). After a further 30-

minute incubation at 37°C without CO2, the sensor cartridge and the cell culture plate were loaded into

the XFe96 Extracellular Flux Analyzer (Agilent). OCR was measured three times for basal respiration and

after each injection of mitochondrial toxins. Post-assay, the cell culture medium was removed, and total

protein content was determined using a BCA Protein Assay Kit (ThermoFisher, #23227). The OCR data

were analyzed with XFe Wave Pro software (Agilent), normalized to total cell protein levels (based on the

BCA assay), and exported using the Seahorse XF Cell Mito Stress Test Report Generator. Statistical

analyses were conducted on data from three independent biological replicates. The study calculated

various parameters related to oxidative phosphorylation, including basal respiration, proton leakage

across the inner mitochondrial membrane, ATP-coupled respiration, maximal respiration, and spare

respiratory capacity.

Multi-electrode array (MEA)

Cryopreserved 30 days old neurons were thawed in a water bath and centrifuged (400g, 5 min, RT) in

basal medium (Table S2) supplemented with ROCK inhibitor (Tocris, #1254). Cell pellets were

resuspended in differentiation medium (Table S2) supplemented with ROCK inhibitor. 48-well CytoView

MEA plates (Axion Biosystems, #M768-tMEA-48W) were coated with 15 µg/ml Polyethyleneimine

solution (Sigma, #P3143) for 1 hour at 37°C followed by 10 µg/ml Laminin coating for 2 hours at 37°C.

Using Tryphan Blue (Sigma, # T8154-20ML) and a Countess automated cell counter (Invitrogen), 80×10^3

cells/well were seeded in the CytoView plate. Cells were incubated at 37°C and 5% CO_2 for 26-33 days

with differentiation medium changes twice a week. Once a week, 1µg/ml Laminin was added to the fresh

media to maintain cell attachment. Neurons were then treated with compound for 72 hours. The electrical

activity was recorded on the Maestro Pro multiwell MEA (Axion Biosystems) using the AxIS Navigator

software (version 3.5.1, Axion Biosystems). Electrical activity was recorded for 10 minutes at a 12.5 kHz

sampling rate and a 5.5 standard deviation threshold level for action potential detection. Before plate

loading the device was allowed to equilibrate for approximately 30 minutes to 37°C and 5% CO2. Data

analyses were performed using the Neural Metric Tool (version 3.1.7, Axion Biosystems). Activity is

expressed as the mean firing rate across electrodes in a well.

SDS gel electrophoresis and Western blotting

$1.5x10^6$ mDA neurons were cultured for at least 50 days in Geltrex-coated 6-well plates with neuronal

media and treated with compound for five days. At day 50, the neuronal media was removed and cells

were washed with pre-warmed DPBS. DPBS was removed and cells were crosslinked by adding 1 mL

pre-warmed PBSx1 supplemented with 2 mM disuccinimidyl glutarate (DSG, ThermoFisher, #20593) and

protease inhibitors for 30 minutes at 37°C. Crosslinking was terminated by adding 50 mM Tris-HCl,

pH=7.5 for 15 minutes at room temperature. Cells were collected and centrifuged for 10 minutes at

1000xg at room temperature. The supernatant was removed and the cell pellet was lysed using lysis buffer

(100 mM NaCl, 5 mM EDTA, 50 mM Tris-Cl, pH 7.4, 50 mM NaF, containing 1% (v/v) Igepal CA-630

and supplemented with protease inhibitors). Cells were incubated for 30 minutes at 4°C with the lysis

buffer and sonicated three times. Total cellular protein was measured using the BCA Protein Detection Kit

(ThermoFisher, #23227) and 25 µg of cell lysate were electrophoresed at 100□V constant on NuPAGE 4–

12% Bis-Tris Midi gels (Invitrogen, NP0321BOX) with NuPage MES-SDS running buffer (Invitrogen,

NP0002). After electrophoreses, gels were electroblotted in nitrocellulose membranes, using a wet-

transfer approach. After electroblotting, the membranes were briefly rinsed in ultrapure water and

incubated in 4% paraformaldehyde/PBS for 30□min at room temperature, followed by blocking for 1

hour at room temperature in TBS-Tween 0.1% (v/v) supplemented with 5% (w/v) dry milk. After

blocking, membranes were incubated with primary antibodies against monomeric α-synuclein (BD

Bioscience, #610787), multimeric α-synuclein (Sigma, ABN2265M), and β-actin (CellSignaling, #3700)

or vinculin (Cell Signaling, #13901) at 4°C overnight. The next day, membranes were washed

3 × 10 min in PBS-Tween 0.1% and incubated with the corresponding secondary antibodies (anti-

Mouse HRP, ThermoFisher, #31430 and anti-Rabbit HRP, ThermoFisher, #A24537) at RT for 1 h in the

dark. Membranes were rinsed in PBS-Tween 0.1%, washed at least 3 × 10 min in PBS-Tween 0.1%,

incubated for 3 minutes with ECL western blotting substrate, and imaged on a LI-COR Odyssey XF

imaging system.

Measurement of intracellular calcium levels

Intracellular calcium levels were evaluated using the cell-permeant calcium indicator Rhod-FF AM

(Invitrogen, R23983). Briefly, $70x10^3$ mDA neurons were seeded per well in a Geltrex-coated black,

clear-bottom 96-well cell culture plate and treated with the selected compound for three days. On

neuronal maturation day 33, the neuronal medium was removed, and cells were washed with the pre-

warmed HBS buffer (145 mM NaCl, 5 mM KCl, 1 mM $CaCl_2$, 1 mM $MgCl_2$, 10 mM HEPES, 10mM

glucose) and incubated for 1 hour at room temperature, protected from light, with HBS buffer containing

0.1% (w/v) BSA and 5 μM Rhos-FF AM. After incubation, the cells were washed twice with 100μL per

well HBS and were incubated for 20 minutes at room temperature, protected from light, for dye de-

esterification. Fluorescence was measured 4 times, every 30 seconds, using a Cytation 5 (Bio-Tek)

fluorescence plate reader, with excitation at 550 nm and emission at 580 nm, at 28°C. After the first

measurement, HBS buffer supplemented with 10 μM of Ionomycin (Sigma, #19657) was added to the

cells, and the plate was measured again (8 times (8 times, every 30 seconds). Fluorescence intensity was

normalized to total cell protein levels, using the BCA Protein Detection Kit (ThermoFisher, #23227).

Statistical analyses were conducted on data from three independent biological replicates.

Evaluation of proteasomal activity

The proteasomal activity was evaluated by measuring the chymotrypsin-like and caspase-like activity.

The chymotrypsin-like activity was measured using the fluorogenic substrate Suc-Leu-Leu-Val-Tyr-AMC

(Enzo, BML-PB02-0005) and the caspase-like activity was measured by using the fluorogenic substrate

Z-Leu-Leu-Glu-AMC (Enzo, BML-ZW9345-0005). Briefly, $60x10^3$ mDA neurons were seeded per well

in a Geltrex-coated black, clear-bottom 96-well cell culture plate and treated with the selected compound

for three days. On neuronal maturation day 33, the neuronal medium was removed and cells were washed

with the pre-warmed assay buffer (20 mM Tris-HCl pH=7.5, 5 mM $MgCl_2$, 0.1 mM EDTA, 1 mM DTT, 1

mM ATP, pH = 7.4). After washing, cells are incubated for 30 minutes at 37°C with 100 μL per well of

assay buffer containing 100 μM of the respective fluorescent substrate. Finally, fluorescence was

measured using a Cytation 5 (Bio-Tek) fluorescence plate reader, with excitation at 380 nm and emission

at 460 nm, at 37°C. Fluorescence intensity was normalized to total cell protein levels, using the BCA

Protein Detection Kit (ThermoFisher, #23227). Statistical analyses were conducted on data from three

independent biological replicates.

Structural analogues

Compounds were directly obtained from Selleckchem (CIDs:

5614, 2044, 1838043, 135419190), Enamine (CID: 135407924), or custom synthesized by SynHet (CID:

3511020). For compound CID 3511020, we chose the abbreviation CEPPE based on its IUPAC name 2-

cyano-3-(3-ethoxy-4-propan-2-yloxyphenyl)prop-2-enethioamide. Chemical structures were plotted from

their Simplified molecular-input line-entry system (SMILES) strings using Python. The rdkit package was

used to create molecule objects from the SMILES strings and to compute 2D coordinates for these

molecules. Tanimoto coefficients were calculated using their CID code via the PubChem Score Matrix

service available at https://pubchem.ncbi.nlm.nih.gov/score_matrix/score_matrix.cgi.

References

1.   Wüllner, U. *et al.* The heterogeneity of Parkinson's disease. *J. Neural Transm.* **130**, 827–838 (2023).

2.   Poewe, W. *et al.* Parkinson disease. *Nat. Rev. Dis. Primer* **3**, 17013 (2017).

3.   Coukos, R. & Krainc, D. Key genes and convergent pathogenic mechanisms in Parkinson disease. *Nat. Rev.*

Neurosci. **25**, 393–413 (2024).

4.   Espay, A. J. *et al.* Biomarker⬚driven phenotyping in Parkinson's disease: A translational missing link in

disease⬚modifying clinical trials. *Mov. Disord.* **32**, 319–324 (2017).

5.   Krüger, R. *et al.* Classification of advanced stages of Parkinson's disease: translation into stratified treatments.

J. Neural Transm. **124**, 1015–1027 (2017).

6.   Prasuhn, J. & Brüggemann, N. Genotype-driven therapeutic developments in Parkinson's disease. *Mol. Med.*

27, 42 (2021).

7.   Laperle, A. H. *et al.* iPSC modeling of young-onset Parkinson's disease reveals a molecular signature of disease

and novel therapeutic candidates. *Nat. Med.* **26**, 289–299 (2020).

8.   Vuidel, A. *et al.* High-content phenotyping of Parkinson's disease patient stem cell-derived midbrain

dopaminergic neurons using machine learning classification. *Stem Cell Rep.* **17**, 2349–2364 (2022).

9.   D'Sa, K. *et al.* Prediction of mechanistic subtypes of Parkinson's using patient-derived stem cell models. *Nat.*

Mach. Intell. **5**, 933–946 (2023).

10. Bose, A., Petsko, G. A. & Studer, L. Induced pluripotent stem cells: a tool for modeling Parkinson's disease.

Trends Neurosci. **45**, 608–620 (2022).

11. Schiff, L. *et al.* Integrating deep learning and unbiased automated high-content screening to identify complex

disease signatures in human fibroblasts. *Nat. Commun.* **13**, 1590 (2022).

12. Cholerton, B. *et al.* Precision Medicine. *Am. J. Pathol.* **186**, 500–506 (2016).

13. Espay, A. J., Brundin, P. & Lang, A. E. Precision medicine for disease modification in Parkinson disease. *Nat.*

Rev. Neurol. **13**, 119–126 (2017).

14. Haston, K. M. & Finkbeiner, S. Clinical Trials in a Dish: The Potential of Pluripotent Stem Cells to Develop

Therapies for Neurodegenerative Diseases. *Annu. Rev. Pharmacol. Toxicol.* **56**, 489–510 (2016).

15. Polymeropoulos, M. H. *et al.* Mutation in the α-Synuclein Gene Identified in Families with Parkinson's

Disease. *Science* **276**, 2045–2047 (1997).

16. Krüger, R. *et al.* AlaSOPro mutation in the gene encoding α-synuclein in Parkinson's disease. *Nat. Genet.* **18**,

106–108 (1998).

17. Singleton, A. B. *et al.* α-Synuclein Locus Triplication Causes Parkinson's Disease. *Science* **302**, 841–841

(2003).

18. Chartier-Harlin, M.-C. *et al.* α-synuclein locus duplication as a cause of familial Parkinson's disease. *The*

Lancet **364**, 1167–1169 (2004).

19. Devine, M. J. *et al.* Parkinson's disease induced pluripotent stem cells with triplication of the α-synuclein locus.

Nat. Commun. **2**, 440 (2011).

20. Byers, B. *et al.* SNCA Triplication Parkinson's Patient's iPSC-derived DA Neurons Accumulate α-Synuclein

and Are Susceptible to Oxidative Stress. *PLoS ONE* **6**, e26159 (2011).

21. Mohamed, N.-V. *et al.* Midbrain organoids with an *SNCA* gene triplication model key features of

synucleinopathy. *Brain Commun.* **3**, fcab223 (2021).

22. Iannielli, A. *et al.* Modeling native and seeded Synuclein aggregation and related cellular dysfunctions in

dopaminergic neurons derived by a new set of isogenic iPSC lines with SNCA multiplications. *Cell Death Dis.*

13, 881 (2022).

23. Oliveira, L. M. A. *et al.* Elevated α-synuclein caused by SNCA gene triplication impairs neuronal

differentiation and maturation in Parkinson's patient-derived induced pluripotent stem cells. *Cell Death Dis.* **6**,

e1994–e1994 (2015).

24. Zambon, F. *et al.* Cellular α-synuclein pathology is associated with bioenergetic dysfunction in Parkinson's

iPSC-derived dopamine neurons. *Hum. Mol. Genet.* **28**, 2001–2013 (2019).

25. Heman-Ackah, S. M. *et al.* Alpha-synuclein induces the unfolded protein response in Parkinson's disease

SNCA triplication iPSC-derived neurons. *Hum. Mol. Genet.* **26**, 4441–4450 (2017).

26. Burbulla, L. F. *et al.* Dopamine oxidation mediates mitochondrial and lysosomal dysfunction in Parkinson's

disease. *Science* **357**, 1255–1261 (2017).

27. Angelova, P. R. *et al.* Alpha synuclein aggregation drives ferroptosis: an interplay of iron, calcium and lipid

peroxidation. *Cell Death Differ.* **27**, 2781–2796 (2020).

28. Tanudjojo, B. *et al.* Phenotypic manifestation of α-synuclein strains derived from Parkinson's disease and

multiple system atrophy in human dopaminergic neurons. *Nat. Commun.* **12**, 3817 (2021).

29. Chandrasekaran, S. N., Ceulemans, H., Boyd, J. D. & Carpenter, A. E. Image-based profiling for drug

discovery: due for a machine-learning upgrade? *Nat. Rev. Drug Discov.* **20**, 145–159 (2021).

30. Chandrasekaran, S. N. *et al.* *JUMP Cell Painting Dataset: Morphological Impact of 136,000 Chemical and*

Genetic Perturbations. http://biorxiv.org/lookup/doi/10.1101/2023.03.23.534023 (2023)

doi:10.1101/2023.03.23.534023.

31. Bray, M.-A. *et al.* Cell Painting, a high-content image-based assay for morphological profiling using

multiplexed fluorescent dyes. *Nat. Protoc.* **11**, 1757–1774 (2016).

32. Park, S. J. *et al.* A receptor tyrosine kinase inhibitor, Tyrphostin A9 induces cancer cell death through Drp1

dependent mitochondria fragmentation. *Biochem. Biophys. Res. Commun.* **408**, 465–470 (2011).

33. Sagara, Y., Ishige, K., Tsai, C. & Maher, P. Tyrphostins Protect Neuronal Cells from Oxidative Stress. *J. Biol.*

Chem. **277**, 36204–36215 (2002).

34. Ferreira, A. F. F. *et al.* Inhibition of TRPM2 by AG490 Is Neuroprotective in a Parkinson's Disease Animal

Model. *Mol. Neurobiol.* **59**, 1543–1559 (2022).

35. Dai, X. *et al.* Tyrphostin A9 protects axons in experimental autoimmune encephalomyelitis through activation

of ERKs. *Life Sci.* **294**, 120383 (2022).

36. Iaubasarova, I. R. *et al.* The mitochondria-targeted derivative of the classical uncoupler of oxidative

phosphorylation carbonyl cyanide m-chlorophenylhydrazone is an effective mitochondrial recoupler. *PLOS*

ONE **15**, e0244499 (2020).

37. McDiarmid, A. H. *et al.* Morphological profiling in human neural progenitor cells classifies hits in a pilot drug

screen for Alzheimer's disease. *Brain Commun.* **6**, fcae101 (2024).

38. Han, X. *et al.* Whole human-brain mapping of single cortical neurons for profiling morphological diversity and

stereotypy. *Sci. Adv.* **9**, eadf3771 (2023).

39. Peng, H. *et al.* Morphological diversity of single neurons in molecularly defined cell types. *Nature* **598**, 174–

181 (2021).

40. Kilfeather, P. *et al.* Single-cell spatial transcriptomic and translatomic profiling of dopaminergic neurons in

health, aging, and disease. *Cell Rep.* **43**, 113784 (2024).

41. Kim, G.-H. J. *et al.* A zebrafish screen reveals Renin-angiotensin system inhibitors as neuroprotective via

mitochondrial restoration in dopamine neurons. *eLife* **10**, e69795 (2021).

42. Cui, J., Carey, J. & Reijo Pera, R. A. Identification of DOT1L inhibitor in a screen for factors that promote

dopaminergic neuron survival. *Front. Aging Neurosci.* **14**, 1026468 (2022).

43. Tabata, Y. *et al.* T-type Calcium Channels Determine the Vulnerability of Dopaminergic Neurons to

778   Mitochondrial Stress in Familial Parkinson Disease. *Stem Cell Rep.* **11**, 1171–1184 (2018).

779   44. Yamaguchi, A. *et al.* Identifying Therapeutic Agents for Amelioration of Mitochondrial Clearance Disorder in

780   Neurons of Familial Parkinson Disease. *Stem Cell Rep.* **14**, 1060–1075 (2020).

781   45. Park, S. J. *et al.* A receptor tyrosine kinase inhibitor, Tyrphostin A9 induces cancer cell death through Drp1

782   dependent mitochondria fragmentation. *Biochem. Biophys. Res. Commun.* **408**, 465–470 (2011).

783   46. Terada, H. & Van Dam, K. On the, stoichiometry between uncouplers of oxidative phosphorylation and

784   respiratory chains. The catalytic action of SF 6847 (3,5-di-tert-butyl-4-hydroxybenzylidenemalononitrile).

785   *Biochim. Biophys. Acta BBA - Bioenerg.* **387**, 507–518 (1975).

786   47. Geisler, J. G., Marosi, K., Halpern, J. & Mattson, M. P. DNP, mitochondrial uncoupling, and neuroprotection: A

787   little dab'll do ya. *Alzheimers Dement.* **13**, 582–591 (2017).

788   48. Zorov, D. B. *et al.* Neuroprotective Potential of Mild Uncoupling in Mitochondria. Pros and Cons. *Brain Sci.*

789   **11**, 1050 (2021).

790   49. Terada, H., Fukui, Y., Shinohara, Y. & Ju-ichi, M. Unique action of a modified weakly acidic uncoupler without

791   an acidic group, methylated SF 6847, as an inhibitor of oxidative phosphorylation with no uncoupling activity:

792   Possible identity of uncoupler binding protein. *Biochim. Biophys. Acta BBA - Bioenerg.* **933**, 193–199 (1988).

793   50. Peruzzotti-Jametti, L. *et al.* Mitochondrial complex I activity in microglia sustains neuroinflammation. *Nature*

794   **628**, 195–203 (2024).

795   51. Gao, C., Jiang, J., Tan, Y. & Chen, S. Microglia in neurodegenerative diseases: mechanism and potential

796   therapeutic targets. *Signal Transduct. Target. Ther.* **8**, 359 (2023).

797   52. Park, D. S. *et al.* iPS-cell-derived microglia promote brain organoid maturation via cholesterol transfer. *Nature*

798   **623**, 397–405 (2023).

799   53. Zhang, W. *et al.* Microglia-containing human brain organoids for the study of brain development and pathology.

800   *Mol. Psychiatry* **28**, 96–107 (2023).

801   54. von Willebrand, M., Zacksenhaus, E., Cheng, E., Glazer, P. & Halaban, R. The tyrphostin AG1024 accelerates

802   the degradation of phosphorylated forms of retinoblastoma protein (pRb) and restores pRb tumor suppressive

803   function in melanoma cells. *Cancer Res.* **63**, 1420–1429 (2003).

55. Walrafen, P. *et al.* Both proteasomes and lysosomes degrade the activated erythropoietin receptor. *Blood* **105**,
600–608 (2005).

56. Conway, K. A., Rochet, J.-C., Bieganski, R. M. & Lansbury, P. T. Kinetic Stabilization of the α-Synuclein
Protofibril by a Dopamine-α-Synuclein Adduct. *Science* **294**, 1346–1349 (2001).

57. Junn, E. & Mouradian, M. M. Human α-Synuclein over-expression increases intracellular reactive oxygen
species levels and susceptibility to dopamine. *Neurosci. Lett.* **320**, 146–150 (2002).

58. Devi, L., Raghavendran, V., Prabhu, B. M., Avadhani, N. G. & Anandatheerthavarada, H. K. Mitochondrial
import and accumulation of alpha-synuclein impair complex I in human dopaminergic neuronal cultures and
Parkinson disease brain. *J. Biol. Chem.* **283**, 9089–9100 (2008).

59. Choi, M. L. *et al.* Pathological structural conversion of α-synuclein at the mitochondria induces neuronal
toxicity. *Nat. Neurosci.* **25**, 1134–1148 (2022).

60. Yin, S. *et al.* Nur77 increases mitophagy and decreases aggregation of α-synuclein by modulating the p-c-
Abl/p-PHB2 Y121 in α-synuclein PFF SH-SY5Y cells and mice. *Eur. J. Med. Chem.* **268**, 116251 (2024).

61. Chen, Y. *et al.* Engineering synucleinopathy☐resistant human dopaminergic neurons by CRISPR ☐mediated
deletion of the *SNCA* gene. *Eur. J. Neurosci.* **49**, 510–524 (2019).

62. Kriks, S. *et al.* Dopamine neurons derived from human ES cells efficiently engraft in animal models of
Parkinson's disease. *Nature* **480**, 547–551 (2011).

63. Ryan, S. D. *et al.* Isogenic Human iPSC Parkinson's Model Shows Nitrosative Stress-Induced Dysfunction in
MEF2-PGC1α Transcription. *Cell* **155**, 1351–1364 (2013).

64. Weykopf, B. *et al.* Induced pluripotent stem cell-based modeling of mutant LRRK2-associated Parkinson's
disease. *Eur. J. Neurosci.* **49**, 561–589 (2019).

65. McKinney, W. Data Structures for Statistical Computing in Python. *Proc. 9th Python Sci. Conf.* 56–61 (2010)
doi:10.25080/Majora-92bf1922-00a.

66. Waskom, M. L. seaborn: statistical data visualization. *J. Open Source Softw.* **6**, 3021 (2021).

67. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12**, 2825–2830 (2011).

Acknowledgements

This research was funded by grants from the Fond National de Recherche (FNR) within the BRIDGES program "MOTASYN" (No.12719684). Fraunhofer ITMP received funding from European Joint Program on Rare Diseases (German Federal Ministry of Education and Research (BMBF), 01GM2002B). Ksilink was supported by the Programme d'investissements d'avenir (PIA) by the Secrétariat général pour l'investissement (SGPI) of the French government. We thank Dr. Tilo Kunath (University of Edinburgh) for providing the AST23 and AST23-2KO iPSC lines. We are grateful for the constant discussions with various colleagues at Ksilink and the Luxembourg Centre for Systems Biomedicine (LCSB) which improved this work.

Author contributions

V.G. designed biochemical experiments, performed neuronal differentiation, biochemical assays, and data analysis; A.We. designed imaging experiments, performed neuronal culturing, imaging assay automation, and compound screening; L.C. developed and performed image analysis and data processing; D.H. performed MEA experiments; K.S. analyzed the MEA data; A.O. co-supervised data processing, P.A.B. co-initiated the study and contributed to grant writing; B.F.R.S. contributed to biochemical experiments; I.B. co-designed biochemical experiments; A.Wi., A.Z. and O.P. identified and provided Tyrphostin A9 structural analogues; R.K. co-initiated the study and supervised biochemical experiments; P.S. co-initiated the study and supported assay design; J.H.W. supervised the study, designed experiments, conceptualized and performed image analysis and data processing, and wrote the manuscript with input from all authors.

Data availability statement

The datasets supporting the conclusions of this article are available in the GitHub repository, https://github.com/Ksilink/Notebooks/tree/main/Neuro/PD_MorphProfileScreening. Raw data is available as CSV files which can be processed using the provided Python-based Jupyter Notebooks.

Competing interest statement

A.We., L.C., D.H., K.S., A.O., P.S., and J.H.W. are or were employed by Ksilink. P.A.B. is receiving or has received research grants from the FNR, Leir Foundation and William N. and Bernice E. Bumpus Foundation. R.K. is receiving or has received research grants from the Fonds National de Recherche (FNR) Luxembourg, Fondation Veuve-Metz-Tesch Luxembourg, the Leir Foundation, the Michael J. Fox Foundation for Parkinson□s Research (MJFF), the European Institute of Innovation and Technology (EIT Health), the Innovative Medicines Initiative (IMI) of the European Union and the European pharmaceutical industry, and the European Union's Horizon 2020 and Horizon Europe research and innovation programs, as well as personal speaker's honoraria and/or travel grants from Abbvie, Zambon and Medtronic; R.K. participated as PI or site-PI for industry sponsored clinical trials without receiving honoraria. All other authors declare no conflict of interest.

 # Supplementary Information

 Morphological profiling reveals neuroprotection via mitochondrial uncoupling

 in human dopaminergic neurons

 Gorgogietas *et al.*

Figure S1 (referring to Figure 1): Quality control of neuronal cell lines for morphological profiling. (A-B)

Immunofluorescent quantification of mDA SNCA-4x SNCA-corr neuron markers after differentiation D30. **(C)**

Karyotype comparison between source iPSCs and D30 neurons. **(D)** Western blot densiometric quantification of

αSyn signal from D70 mDA neuron lysates. The experiment was performed once. **(E)** Mitochondrial membrane

potential assessment by JC1 dye fluorescent signal change in D35 neurons. Each data point shows technical

replicates from a single experiment. The boxplots show the median and the 2nd and 3rd quartiles of the data.

Whiskers indicate 1.5X of the interquartile range. **(F)** Explained data variation by first ten principal components

(PCs) derived from principal components analysis. Pooled data from four independent experiments each containing

at least 12 technical replicates per condition was used for analysis. **(G)** Top ten Pearson correlated morphological

features for each of the first three PCs after PCA. Pooled data from four independent experiments each containing at

least 12 technical replicates per condition was used for analysis.

Figure S2 (referring to Figure 2): Primary screen quality assessment. (A) Raw and normalized αSyn staining

intensity and live cell ratio values for all 20 screened plates. Each data point corresponds to the median value per

well. **(B)** Selected features for hit identification based on <0.95 Pearson correlation. **(C)** Strictly standardized mean

difference (SSMD) values per plate. The SSMD was calculated based on the morphological profile Mahalanobis

distance between SNCA-4x + DMSO and SNCA-corr + DMSO conditions per plate (left panel) and between

SNCA-4x + DMSO and SNCA-4x + Prostratin conditions per plate (right panel). SSMD values <-1 indicate a good

plate quality (indicated by the dashed line). **(D)** Confusion matrix describing the per-class performance of the LDA

classifier on the test data set containing 20% of the overall data not seen during training (left panel). Cross validation

plot showing the LDA classifiers performance across 10 different slices of the full dataset (right panel). **(E)**

Supervised Linear Discriminant Analysis (LDA) classification of Control neurons, and SNCA-4x neurons treated

with either DMSO, Prostratin or screened compound. Each datapoint represents data from one well and for

compounds the mean value of all four replicates is shown. **(F)** CellPainting profiles obtained in U2OS cells of hits

selected during primary screening clustered by their Cosine similarity. Compounds are color coded by attributed

biological activity.

Figure S3 (referring to Figure 3): Dose-response assessment of cell viability and αSyn level. (A) Cell viability

assessment based on Hoechst-stained nuclear size and intensity. mDA neurons were treated with 0.25-20 μM

897   compound for 72 hours before fixation. **(B)** Immunostaining-based αSyn level assessment. Toxic compound

898   concentrations were excluded from the graph. Each compound concentration was tested at least in triplicate. The

899   mean and 95% confidence interval is shown.

900
901   **Figure S4 (referring to Figure 3): Dose-response assessment of compound effects on morphological profiles.**

902   Morphological profile Mahalanobis distances of compound treated mDA neurons relative to SNCA-4x and SNCA-

903   corr morphological profiles. The redline line indicates the 4σ window from the SNCA-4x median and blue line the

904   indicates the 2.5σ window from the SNCA-corr median. Dot sizes represent the used concentration. Each compound

905   concentration was tested at least in triplicate. Mean and 95% confidence interval are shown.

Figure S5 (referring to Figure 4): Dose-response assessment of compound effects on mitochondrial membrane potential. mDA neurons were treated with 0.1-10 μM of each compound for 72 hours and the mitochondrial membrane potential was measured using the JC1 dye. Three independent experiments with at least three technical

 replicates each were performed. The mean and 95% confidence interval is shown. Dashed lines indicate the means

911   of the indicated control treatments.

Figure S6 (referring to Figure 4): Dose-response assessment of compound effects neuronal ROS level. mDA

neurons were treated with 0.5, 1, and 2.5 μM of each compound for 72 hours and the ROS level was determined

using the ROS-ID Total ROS/Superoxide detection kit. The obtained signal was normalized to total protein content.

Three independent experiments with each at least three technical replicates were performed. The mean and 95%

confidence interval is shown. Dashed lines indicate the means of the indicated control treatments.

Figure S7 (referring to Figure 5): Tyrphostin A9 structural analogues with mitochondrial uncoupling activity

reduce cellular ATP levels. (A) SNCA-4x and SNCA-corr mDA neurons were differentiated until D33, treated with

DMSO, 2.5 μM Tyrphostin A9 or its structural analogues for 72 hours and ATP levels were determined in a

luminescence-based plate reader assay. **(B)** Same as in (A), but mDA neurons were simultaneously incubated with

compound and the mitochondrial complex I inhibitor Rotenone. Three independent differentiations with at least

three technical replicates each were performed. Large markers indicate biological replicate medians and small

markers represent all technical replicate data. The boxplots show the median and the 2nd and 3rd quartiles of the

data. Whiskers indicate 1.5X of the interquartile range. Welch's unequal variances t-test was performed using the

medians of all biological replicates. ns = not significant, * = p$\square$<$\square$0.05, ** = p$\square$<$\square$0.01, *** = p$\square$<$\square$0.001.

Figure S8 (referring to Figure 6): Full gel/blot images. Shown are raw uncropped images of membranes stained

with antibodies against αSyn and Vinculin.

Table S1: Supplementary dataset of identified compounds after morphological profiling including morphological

features.

Table S2: Key resource table.

Table S3: Supplementary dataset: Compound library overview.

Table S4: Supplementary dataset: Morphological feature overview.