TITLE: *Sox9* prevents retinal degeneration and is required for limbal stem cell differentiation in the adult mouse eye

Running title: Retinal and limbal defects in adult *Sox9* mutant mice

Alicia Hurtado[1,5,*], Víctor López-Soriano[1,6,*], Miguel Lao[1,7,*], M. Ángeles Celis-Barroso[1], Pilar Lazúen[1], Alejandro Chacón de Castro[1], Yolanda Ramírez-Casas[1,8], Miguel Alaminos[2,3], J. Martin Collinson[4], Miguel Burgos[1], Rafael Jiménez[1,¶,**], F. David Carmona[1,2,¶,**], Francisco J. Barrionuevo[1,¶,**]

Affiliations:

[1] Departamento de Genética e Instituto de Biotecnología, Centro de Investigación Biomédica (CIBM), Universidad de Granada, Granada, Spain.

[2] Instituto de Investigación Biosanitaria ibs.GRANADA, Granada, Spain.

[3] Departamento de Histología, Universidad de Granada, Granada, Spain

[4] School of Medicine, Medical Sciences and Nutrition, University of Aberdeen, Institute of Medical Sciences, Aberdeen, UK.

Current addresses:

[5] Centro Andaluz de Biología del Desarrollo (CABD), CSIC/UPO/JA, Seville, Spain.

[6] Internal Medicine V, Hematology, Oncology and Rheumatology, Heidelberg University Hospital, Heidelberg, Germany.

[7] IBiS Instituto de Biomedicina de Sevilla, Seville, Spain.

[8] Centro de Investigación Biomédica, Facultad de Medicina, Departamento de Fisiología, Instituto de Biotecnología, Granada, Spain.

* Contributed equally.

¶ Shared last authorship.

** Corresponding authors

ABSTRACT

Sox9 is a transcription factor with multiple roles during development and in adult organ homeostasis. In the adult eye, *Sox9* expression persists in several cell types,

5   including the retinal pigmented epithelium cells and the Müller glial cells, as well as in the limbal and corneal basal epithelia. To uncover the role of *Sox9* in these cell types, we induced the deletion of the gene in adult mice. We found that, after *Sox9* ablation, mutant mice undergo a severe process of retinal degeneration characterized by the loss of Müller glial cells and complete depletion of the photoreceptors layer. Moreover, by

10   combining single-cell RNA sequencing and *Sox9* lineage tracing, we found that *Sox9* is expressed in a basal limbal stem cell population with the ability to form two types of long-lived cell clones involved in stem cell maintenance and homeostasis. Mosaic analysis of *Sox9* positive and negative cells confirmed that the gene is essential for limbal stem cell differentiation. Our results show that *Sox9* is required for the

15   maintenance of retinal integrity and for limbal stem cell differentiation in the adult mouse eye.

INTRODUCTION

The retina and cornea represent indispensable components of the vertebrate visual system that ensure proper vision (Casey et al., 2023). The retina, a multilayered sensory tissue at the back of the eye, converts light into neural signals, initiating visual perception. It consists of various specialized cells organized into three nuclear layers and two synaptic layers. Photoreceptors, including rods and cones, are located in the outer nuclear layer (ONL) and are responsible for converting light stimuli into electrical signals. These signals are then transmitted to bipolar cells in the inner nuclear layer (INL), which, in turn, establish synapses with retinal ganglion cells (RGCs), whose axons form the optic nerve that send the electrical impulse to the brain. Additionally, horizontal and amacrine cells, also positioned in the INL, provide lateral connections, offering feedback and feedforward signals. Finally, Müller glial (MG) cells, with nuclei located in the INL and bodies extending across the entire retina, provide structural, metabolic, and functional support to the other retinal cells, thus maintaining retinal function and integrity (Baden et al., 2020).

The cornea, positioned at the forefront of the eye, is a transparent and refractive structure responsible for transmitting and focusing light onto the retina, which is devoid of blood vessels. The outermost layer of the cornea is a stratified squamous epithelium that undergoes continuous regeneration. Stem cells responsible for this replenishment reside within a specialized niche located at the periphery, adjacent to the conjunctiva, known as the limbus (Lavker et al., 2004; West et al., 2015). Within this limbal niche, long-lived limbal epithelial stem cells (LESCs) give rise to a more differentiated progeny responsible for the regeneration of the entire corneal epithelium throughout two distinct

40   pathways of cell migration: one within the basal layer, moving from the limbus to the center of the cornea, and another from the basal layer to the terminally differentiated, desquamated cells at the epithelial surface (Yazdanpanah et al., 2017).

SOX9 is a transcription factor involved in the regulation of multiple developmental processes (Jo et al., 2014). In humans, mutations affecting *SOX9* cause campomelic

45   dysplasia (CD), a syndrome characterized by skeletal malformations and XY sex reversal (Wagner et al., 1994; Foster et al., 1994). In the adult mice, SOX9 has been shown to play a role in the homeostasis of neural, liver, pancreas, intestine and nail stem cells (Scott et al., 2010; Furuyama et al., 2011; Lao et al., 2022), as well as in the maintenance of Sertoli cells (Barrionuevo et al., 2016), among others.

50   Several works have addressed the role of *Sox9* during the developing eye. In the context of the mouse retina, *Sox9* is expressed in multipotent mouse retinal progenitor cells, undergoes downregulation during neuronal differentiation, and then the gene is exclusively expressed in Müller glial cells and retinal pigment epithelia (RPE) until adulthood (Poché et al., 2008; Muto et al, 2009). Deletion of *Sox9* in developing retinal

55   cells using a Chx10-Cre mouse line revealed that *Sox9* mutant retinae exhibited normal lamination as well as nuclear layer thickness comparable to control samples, indicating that loss of *Sox9* alone likely does not affect global retinal fate determination (Poché et al., 2008).   shRNA-induced downregulation of *Sox9* and *Sox8* in retinal explants resulted in a reduced population of Müller glial cells and a moderate increased in the

60   proportion of rod photoreceptors, indicating that both SoxE genes play roles in the specification of Müller glial cells (Muto et al, 2009). Similar results were obtained when *Sox9* was conditionally deleted in the developing retina using a *Pax6*-α-Cre mouse line

(Goto et al., 2018). In the mature RPE, *Sox9* collaborates with other transcription factors, such as OTX2 and LHX, to regulate the visual cycle, a series of biochemical reactions that regenerate 11-cis-retinal, the chromophore essential for phototransduction in photoreceptor cells (Masuda et al., 2014). Three studies have investigated *Sox9* inactivation in RPE cells. While one study reported no obvious morphological abnormalities in the retina and RPE upon conditional deletion of the gene (Masuda et al., 2014), two subsequent studies demonstrated that *Sox9* was required for proper choroid differentiation (Goto et al., 2018; Cohen-Tayar et al., 2018).

In the cornea, transcriptome analysis from slow cycling cells identified *Sox9* as a putative stem cell marker or determinant, and immunohistochemical analyses confirmed that the gene is expressed in LESCs and in corneal epithelial cells (Sartaj et al., 2017; Parfitt et al., 2015; Menzel-Severing et al., 2018). RNAi silencing of the gene in cultured limbal stem cells resulted in reduced expression of progenitor cell markers and increased expression of differentiation markers. Further analyses showed that *Sox9* and Wnt/ß-catenin signaling antagonizes to maintain a balance between quiescence, proliferation, and differentiation limbal stem cells (Menzel-Severing et al., 2018).

Although several studies have addressed the role of *Sox9* in retinal and corneal development (Poché et al., 2008; Muto et al., 2009; Masuda et al., 2014; Sartaj et al., 2017; Parfitt et al., 2015; Menzel-Severing et al., 2018), its specific *in vivo* contributions to the maintenance and regeneration of these tissues in adult mammals remain unknown.

In this work, we used a tamoxifen-inducible Cre/LoxP system to inactivate the *Sox9* gene in adult mice and a lineage-tracing mouse strain to study the fate of *Sox9-*

expressing cells. With these tools, in conjunction with scRNAseq analysis, we conducted a comprehensive study of morphological changes, cell-specific marker expression, and lineage tracing to determine the role of *Sox9* in maintaining retinal integrity and corneal stem cell homeostasis.

RESULTS

Deletion of *Sox9* leads to retinal degeneration in adult mice

To evaluate the role of *Sox9* in adult retinal homeostasis, we conditionally deleted the gene in adult mouse cells using a ubiquitous tamoxifen-inducible CAGG-CreER^{TM}

95   recombinase (Hayashi and McMahon, 2002) and a conditional *Sox9$^{flox/flox}$* allele (Akiyama et al., 2002) (Fig. 1A). Histological examination of retinas from control (Cre-negative *Sox9$^{flox/flox}$*; n = 7) and *CAGG-CreER^{TM};Sox9$^{flox/flox}$* (hereafter *Sox9$^{Δ/Δ}$*; n = 24) adult mice (two months old) at different days after TX administration (DATX) revealed that most of the analyzed mutant samples (n = 18/24; Table S1), showed either no

100   relevant morphological defects in the retinal sections or a mild phenotype in which the main retinal nuclear layers (INL and ONL) occupied a smaller area and their edges did not appear as clearly differentiated as in the control retinas. However, we also observed an extreme phenotype in a reduced number of mutant retinas (n = 6/24), characterized by the complete loss of the ONL (Fig. 1B). We categorized *Sox9$^{Δ/Δ}$* retinas into "mild"

105   and "extreme" phenotypes in order to facilitate interpretation of our data. Classification was based on a qualitative assessment of ONL integrity in histological sections. Specifically, samples were classified as "extreme" when the ONL was completely depleted, and as "mild" when the ONL persisted, albeit variably reduced in thickness.

This phenotypic classification reflects observable structural differences rather than a

110   fixed quantitative threshold. Some variability exists within the "mild" group, likely due to

differences in recombination efficiency and the mosaic nature of tamoxifen-induced Cre-

mediated *Sox9* deletion. The severity of the phenotype showed no correlation with the

time elapsed after tamoxifen administration, as we found mutant retinas with no

apparent phenotype at 100 DATX and others with an extreme phenotype at 20 DATX

115   (Table S1). Retinal degeneration was never observed in mice that had not been

tamoxifen-treated, nor any other control groups, making the presence of the retinal

degeneration allele of photoreceptor cGMP phosphodiesterase 6b (*Pde6b*[rd1]) unlikely in

our mice (Bowes et al., 1990). However, we acknowledge that definitive exclusion of this

possibility would require PCR-based genotyping.

120   During retinal development, *Sox9* is expressed in neural RPCs but once the neural

retina is differentiated, *Sox9* expression is restricted to MG and RPE cells (Poché et al.,

2008; Masuda et al., 2014). This is, indeed, the SOX9 expression pattern we observed

in retinal sections obtained from adult control mice (Fig. 1C). To explain the phenotypic

diversity shown by the TX-treated mice, we assessed the efficiency of *Sox9* deletion in

125   mutant mice. For this, we performed SOX9/SOX8 double immunofluorescence, as both

transcription factors are normally co-expressed in MG cells (Fig 1C; Muto et al., 2009),

and we consistently observed the presence of many SOX8[+] cells that lacked SOX9

expression in the retinas of mutant TX-treated mice (Fig. 1C). We calculated the

percentage of SOX8[+] cells that also co-expressed SOX9 in control and in *Sox9*[Δ/Δ]

130   retinas with mild and extreme phenotypes (Fig. 1D; Tables S1 and S2). In control MG

cells, all SOX8[+] cells also expressed SOX9. In contrast, in mutants with a mild

Figure 1. Efficiency and morphological impact of *Sox9* deletion on the adult mouse retina. **(A)** Schematic of the experimental design. Adult *Sox9^{flox/flox}*; *CAGG-CreER^{TM}* mice were treated with tamoxifen (TX) at two months of age to induce ubiquitous *Sox9* deletion. Retinal samples were collected at various days after tamoxifen treatment (DATX). **(B)** Hematoxylin-eosin stained histological sections of retinas from control and *Sox9^{Δ/Δ}* mice. Mutant retinas exhibit variability in phenotype severity, with some showing mild morphological defects and a few displaying an extreme phenotype characterized by the loss of the outer nuclear layer (ONL). **(C)** Double immunofluorescence for SOX8 (green) and SOX9 (red) in retinal sections from adult control and *Sox9^{Δ/Δ}* mice. The severity of phenotypes correlates with the extent of *Sox9* inactivation. Regions completely lacking SOX8+ cells are indicated by arrowheads. DAPI (blue) was used to counterstain nuclei. **(D)** Quantification of the percentage of SOX8+ cells co-expressing SOX9 in control and *Sox9^{Δ/Δ}* retinas with mild and extreme phenotypes. **(E)** Quantification of SOX8+ cells per 100 μm of inner nuclear layer (INL) length in control and *Sox9^{Δ/Δ}* retinas. **(F)** Immunofluorescence for S100 showing strong labelling of Müller cell processes in control retinas and a progressive reduction in *Sox9^{Δ/Δ}* samples. **(G)** Quantification of S100+ signal in control and *Sox9^{Δ/Δ}* mice expressed as percentage of surface area occupied in the retina. The black scale bar represents 50 μm in B, and the white scale bars represent 100 μm in C and 50 μm in F. GCL, Ganglion cell layer; INL, inner nuclear layer; RPE, retinal pigmented epithelium. Mann-Whitney U test $P < 0.01$ (**), $P < 0.001$ (***).

phenotype, the average percentage was 55%, although with a large standard deviation (55% ± 23%),while in extreme mutants, the percentage was significantly lower (16% ± 12%). Altogether, these results indicate that the severity of the phenotype is directly related to the efficiency of *Sox9* inactivation, and that only individuals with *Sox9* inactivation occurring in at least 80% of the cells expressing the gene exhibit the extreme phenotype.

Next, we used cell-specific molecular markers to characterize the most affected *Sox9*-deficient retinas. Initially, we focused on the MG cell layer by means of SOX8 immunofluorescence. We counted the number of SOX8+ cells per 100 μm of INL, and we found a significant reduction in the number of MG cells in mutant retinas (control, 18.5 ± 3.5, n = 7; mutant, 10.5 ± 1.2 n= 6; p = 0.000107, Mann-Whitney U test; Fig 1E; Table S3). In addition to the reduced number of SOX8+ cells, we observed areas within

some *Sox9* mutant INL layers that completely lacked SOX8$^+$ cells (Fig. 1C arrowheads), indicating that these regions were entirely devoid of MG cells.

To independently validate the loss of MG cells in *Sox9*-deficient retinas, we examined the expression of S100, a cytoplasmic marker that labels the processes of adult Müller cells. In control retinas, strong S100 immunoreactivity was observed across the inner retina, outlining the typical radial projections of Müller glia (Fig. 1G). In contrast, *Sox9*$^{\Delta/\Delta}$ retinas with an extreme phenotype exhibited a marked reduction in S100 signal (Fig. 1F). Given the diffuse cytoplasmic localization of S100, we quantified its expression by measuring the fluorescence signal within a defined surface area of the retina. This analysis revealed a statistically significant reduction in S100 signal intensity in mutant samples (including both mild and extreme phenotypes) compared to controls (Fig. 1G; Table S4), further supporting the loss of MG cells upon *Sox9* deletion.

Since the main observable consequence of *Sox9* deletion in the adult mouse retina is the absence of the ONL layer, which harbors both rod and cone photoreceptors, we further examined the status of these cell types by immunofluorescence. Cone photoreceptor cells were assessed by retinal whole-mount staining with two different opsins: OPN1SW (short-wavelength-sensitive opsin, referred to as S opsin) and OPN1LW (medium- and long-wavelength-sensitive opsin, referred to as M opsin). In control retinas, we observed uniform staining on the ventral surface and on the entire surface for S opsin and M opsin, respectively. In contrast, in *Sox9*-deficient retinas we could only identify small isolated immunofluorescent patches (Fig. 2A. upper). Double immunofluorescence on retinal sections confirmed that mutant retinas lacked expression for both cone photoreceptors (Fig. 2A, bottom). We also performed

190   **Figure 2.** Immunodetection of retinal cell markers in adult retinas from control and *Sox9*^{Δ/Δ} mice.
(A) Immunofluorescence analysis of cone photoreceptor cells in retinal whole-mounts (upper

immunofluorescence for RHO, a marker for rod photoreceptor cells, on retinal whole-mounts and sections. We found a pattern of expression similar to that described for M-opsin in both the control and mutant retinas. These results evidence that *Sox9* is required for the maintenance of the two photoreceptor cell types present in the adult mouse retina (Fig. 2B).

We also studied the status of other retinal cell types. The transcription factor BRN3A was used to identify ganglion cells (Nadal-Nicolás et al., 2009), which were shown to decrease in number in the mutant retinas, compared to control ones (Figs. 2C and 2D and Table S5; n = 5 controls, n = 12 mutants; Mann-Whitney U test, P = 3 × 10^{-4}). Similarly, double immunodetection of the transcription factors PAX6 and AP2A was used to identify both amacrine and horizontal cells (Fig. 2E), as previously described (Marquardt et al., 2001; Barnstable et al., 1985; Edqvist and Hallböök, 2004), showing a similar reduction in both cell types in degenerated retinas (Figs. 2F and 2G and Table S6; AP2α+ amacrine cells: n = 3 controls, n = 8 mutants; 2-sample T-tests P = 0.029; PAX6+/AP2α− horizontal cells: n = 3 controls, n = 8 mutants; Mann-Whitney U test P = 0.021). These findings indicate that the loss of *Sox9* in the adult retina ultimately leads

to the degeneration of multiple inner retinal neuronal populations, beyond the previously

described effects on photoreceptors and Müller glia.

Photoreceptors undergo apoptosis in the absence of *Sox9*

Since photoreceptors are absent in severely affected *Sox9*-mutant retinas, we

225   conducted TUNEL assays to study the role of cell death in the process of retinal

degeneration. In control samples (n=5), almost no TUNEL signal was observed in the

retina. In contrast, $Sox9^{\Delta/\Delta}$ mice (n=15) showed numerous TUNEL+ cells, mainly located

in the persisting ONL, indicating that photoreceptor cells were dying (Fig. 3A). Although

extensive TUNEL staining in the ONL was clearly observed in two $Sox9^{\Delta/\Delta}$ retinas with

mild phenotypes, this pattern was not consistently present across the full cohort. In the remaining 13 mutant retinas, we observed a modest but noticeable increase in the number of apoptotic cells compared to controls (Fig. 3B; Table S7). Despite a high frequency of zero counts (particularly among controls), the difference between groups reached statistical significance when analyzed using a zero-inflated Poisson model (P = 0.028; n = 5 controls, 13 mutants). These findings suggest that photoreceptor apoptosis following Sox9 deletion may occur acutely and within a narrow temporal window, making it challenging to capture the full degenerative process at a single time point.

Müller glial cells undergo reactive gliosis in *Sox9*-deficient retinas

Finally, we decided to study the expression of the glial fibrillary acidic protein (GFAP), whose expression is upregulated in MG cells undergoing reactive gliosis associated with retinal cell injury (Ekström et al., 1988). As previously reported (Fernández-Sánchez et al., 2015), in control mice GFAP immunostaining was mainly limited to the ganglion cell layer (GCL), in the inner margin of the retina. However, in *Sox9*-deficient retinas with a mild phenotype, GFAP-positive processes reached the INL layer. In mutant retinas with an extreme phenotype GFAP-positive MG processes were

distributed throughout the entire thickness of the degenerated retinas. Thus, in the absence of *Sox9*, MG cells seem to undergo a progressive process of gliosis (Fig. 3C). To support these observations quantitatively, we measured GFAP fluorescence intensity across defined retinal surface areas in control and *Sox9$^{\Delta/\Delta}$* mice (Fig. 3D; Table S8).

265   This analysis revealed a statistically significant increase in GFAP signal in mutant retinas compared to controls (Mann-Whitney U test, P = 0.024; n = 4 controls, 10 mutants). These results are consistent with a progressive gliotic response following *Sox9* deletion and provide further evidence that MG cells become reactive in the absence of *Sox9*.

270   **Lineage tracing of *Sox9*-expressing cells in the limbus and the cornea**

We next decided to shed light on the role of *Sox9* in the adult cornea. Previous studies have documented the expression of *Sox9* in the adult mouse limbal and central cornea epithelium (Menzel-Severing et al., 2018; Sartaj et al., 2017). In line with these findings, we also observed the presence of the SOX9 protein in the basal cells of both

275   structures (Fig. 4A and B), which are the sites housing corneal progenitor and stem cells. To assess the stemness potential of these *Sox9*-positive cells, we conducted TX-inducible genetic lineage tracing by breeding mice harboring an *IRES-CreER^{T2}* cassette inserted into the endogenous *Sox9* locus, *Sox9$^{IRES-CreERT2}$* (Soeda et al., 2010), to either *R26R-EYFP* (Srinivas et al., 2001) or *R26R-LacZ* (Soriano, 1999) reporter mice. We

280   induced labeling of *Sox9*-expressing cells and their progeny by administering TX to adult mice (two months old) and analyzed reporter expression at different DATX (Fig. 4C). No X-gal- or EYFP-positive cells were detected at any of the stages analyzed in the absence of TX

Figure 4. Analysis of the fate of SOX9-expressing cells through lineage tracing experiments. **(A-B)** Immunofluorescence for SOX9 in limbal (A) and corneal (B) sections of adult control mice. DAPI (blue) was used to counterstain nuclei. **(C)** Schematic of the genetic lineage tracing strategy. Tamoxifen (TX) was administered to label SOX9-expressing cells and their progeny. **(D-F)** Analysis of EYFP expression in the limbus (D-E) and cornea (F) of *Sox9*-EYFP adult mice. At 10 days after TX administration (10 DATX), discrete EYFP-positive clones are visible in the limbal region (D). By 20 DATX, these clones expand towards the peripheral cornea, with some cells reaching the outer epithelial layers at the limbus-cornea border (arrowhead in E). In the central cornea, EYFP-positive cells extend along the entire epithelium by 20 DATX (arrowhead in F). **(G)** Short-term whole mount X-gal staining of *Sox9*-LacZ eyes. The first LacZ-positive cells scattered across the limbal and corneal surface appeared at 3 DATX (arrows). **(H)** Long-term whole mount X-gal staining of *Sox9*-LacZ eyes. At 45 DATX, LacZ-positive stripes emerge from the limbus and extend into the peripheral cornea (arrow). At 60 DATX, circumferential clones are observed in the limbus (arrowheads). At 90 DATX, three clone types are visible: circumferential clones in the limbus (arrowheads), stripes from the limbus reaching the central cornea (arrow), and stripes without a base in the limbus (asterisk). At 365 DATX, the first two clone types are mainly observed, though in reduced numbers and larger sizes (see arrowheads and arrow as in the 90 DATX picture), with some stripes without a base in the limbus (asterisk). Dashed lines in G and H enclose the limbal area Scale bars in E and F represent 50 μm; scale bars in G and H represent 500 μm.

(Fig. S1, control panels). Immunofluorescence for EYFP on limbal $Sox9^{IRES\text{-}CreERT2}$-*R26R-EYFP (Sox9-EYFP)* sections at 10 DATX unveiled the presence of discrete clones of EYFP-positive cells (Fig. 4D). By 20 DATX, these clones were larger and extended towards the peripheral cornea. At this latter stage, EYFP-positive clones with cells reaching the outer layers of the epithelium were visible at the limbus-cornea border (Fig. 4E, arrowhead). At 20 DATX, we also observed groups of cells spanning the entire apical-basal axis of the central corneal epithelium (Fig. 4F, arrowhead). These results demonstrate that SOX9$^+$ cells residing in the basal layer of the limbal and corneal epithelium are progenitors of the terminally differentiated cells of the corneal epithelium.

We then investigated the dynamics of stem cell progression through whole mount Xgal staining of $Sox9^{IRES\text{-}CreERT2}$ *;R26R-LacZ (Sox9-LacZ)* eyes. At first, we conducted a short-term study with a single TX injection and analyzed X-gal staining for up to ten days after treatment (Figs. 4G and S1). We observed the first X-gal-positive cells at 3 DATX, and they were scattered throughout the entire limbal and corneal surface (Fig. 4G, arrows). At 5 DATX, most X-gal$^+$ cells were located in the limbal area, and by 10 DATX, the number of stained cells increased notably in this area. These observations suggest that *Sox9*-expressing descendant cells experience a process of proliferation in the limbus, illustrating their transient cell amplification behavior. To gain insights on the long-term cloning capacity and contribution dynamics of *Sox9*-progenitor cells, we administered tamoxifen for 10 days in the food and performed X-gal staining from 30 DATX to 365 DATX (Figs. 4H and S1). At 30 DATX, LacZ was mostly expressed in the limbus, although lacZ$^+$ clones could also be observed throughout the peripheral and central cornea. Overall, these clones were larger than those observed at 10 DATX.

Fifteen days later (45 DATX), the limbal staining began to disperse, allowing the

330   observation of discrete clones. Simultaneously, LacZ$^+$ stripes emerging from the limbus

and extending into the peripheral cornea became apparent (Fig. 4H, 45 DATX, arrow,

and Fig. S1). This situation was evident at 60 DATX, when the thickness and number of

the stripes had increased. Additionally, starting at this stage, circumferential clones that

remained in the limbus could also be appreciated (Fig. 4H, 60 DATX, arrowheads, and

335   Fig. S1). At 90 DATX, three types of clones were observed: circumferential ones in the

limbus (Fig. 4H, 90 DATX, arrowheads, and Fig. S1), stripes originating from the limbus,

which often reached the central cornea (Fig. 4H, 90 DATX, arrow, and Fig. S1), and

stripes without a base in the limbus (Fig. 4H, 90 DATX, asterisk, and Fig. S1). At 365

DATX, the first two types of clones were mainly observed, albeit in reduced numbers

340   and larger sizes. We also observed some stripes without a base in the limbus (Fig. 4H,

365 DATX, asterisk, and Fig. S1). Thus, *Sox9* is expressed in a basal limbal stem cell

population with the ability to form two types of long-lived cell clones involved in stem cell

maintenance and homeostasis.

Sox9 is involved in limbal stem cell differentiation

345   To gain insight into the function of *Sox9* as a limbal stem cell marker, we used the

single-cell RNA-sequencing dataset of isolated epithelial cells from the limbus (with

marginal conjunctiva and corneal periphery) recently generated by Altshuler and

colleagues (Altshuler et al., 2021). In this study, the authors describe the co-existence of

two separated stem populations in the limbus, the quiescent "outer" limbus (OLB) one,

350   adjacent to the conjunctiva, and the active "inner" limbus (ILB) one, adjacent to the

peripheral cornea. Unbiased clustering of the sc-RNAseq dataset and subsequent

assignment of cell identity revealed the existence of 10 cell populations equivalent to those identified by Altshuler and colleagues (Figs. 5A and S2A). Density plots based on gene-weighted kernel density estimation and violin plots of expression levels revealed

355   high expression levels of *Sox9* in OLB cells, and progressive weaker expression in the ILB, corneal basal and corneal suprabasal clusters (Figs. 5B and C). We also detected strong expression for *Sox9* in a mitotic cluster (Figs. 5B). Next, we inferred the trajectory of limbal and corneal cells by using the partition-based graph abstraction (PAGA) algorithm, and reconstructing the temporal order of differentiating cells with the diffusion

360   pseudotime software, using the OLB cluster as "root" (Wolf et al., 2019; Haghverdi et al., 2016). The PAGA graph revealed a strong connection between the OLB node and a mitotic cell population with high expression of OLB and stem cell markers (*Gpha2*, *Ifitm3, Cd63,* and *Krt15)*, which, in turn, was strongly linked to a second mitotic cell cluster, in which the expression of these markers diminished while the expression of

365   corneal markers (*Krt12*, *Ppp1r3c,* and *Slurp*) increased (Figs. 5D and Fig. S2B). Through the PAGA graph, we also observed that the expression levels of *Sox9* correlated with the pseudotime graph (Fig. 5E), indicating that *Sox9* expression decreases as transiently amplifying progenitors undergo progressive differentiation from limbal to peripheral corneal cells. To validate these results, we decided to closely

370   examine *Sox9* expression in the limbus using immunofluorescence. Previous analyses revealed that the outer limbus is approximately 100 µm wide, while the inner limbus is wider, around 240 µm (Altshuler 2021). We observed that in the region corresponding to the OLB, most cells showed strong *Sox9* expression. In the area corresponding to the ILB, this immunoreactivity appeared weaker in the basal layer (corresponding to the ILB

proper), and no expression was detected in the suprabasal layers (flattened cells; Fig. 5F left). Double immunofluorescence for SOX9 and P63, which is expressed in basal cells of the limbal epithelium, but not by transient amplifying cells covering the corneal surface (Pellegrini et al., 2001) revealed that Sox9 expression was restricted to P63-positive cells (Fig. 5F right). These observations confirm that Sox9 is expressed in a basal cell population within both the OLB and ILB, and that its expression decreases in differentiated transient amplifying cells. Next, we conducted a comparative analysis of gene expression to examine the differences between cells exhibiting high (>= 0.7) and low expression levels (between 0.7 and 0.15) of Sox9 . We identified 2816 deregulated genes, 137 upregulated, and 2679 downregulated (adjusted p-value < 0.05; | avg_log2FC| > 0.1; Fig. 5G; Table S9). Among the upregulated genes we found OLB markers including Gpha2, Ifitm3 and Cd63, in addition to stem cell markers such as Krt14 (Fig. 5G; Table S9).

Finally, we performed gene ontology analysis using the downregulated genes, and we found terms related to stem cell differentiation, such as "mitotic cell phase transition" (Knoblich, 2008), "autophagy" (Chang, 2020), "cell-cell junction" (Ning et al., 2021),

Figure 6. Impact of *Sox9* deletion on limbal stem cell differentiation and clonogenic capacity. **(A)** Hematoxylin-eosin stained histological sections of the limbus and cornea from control and *Sox9*$^{\Delta/\Delta}$ mice. **(B)** Immunofluorescence staining for SOX9 (red) and P63 (green) in the limbus and cornea from control and *Sox9*$^{\Delta/\Delta}$ mice. **(C)** Quantification of SOX9$^+$ and P63$^+$ cells per 100 μm of corneal limbus section length in control and *Sox9*$^{\Delta/\Delta}$ mice.**(D)** Schematic of the generation of *Sox9*$^{\Delta/\Delta}$-LacZ mice. Tamoxifen (TX) was administered for ten days to label *Sox9*-deleted cells and their progeny. **E)** X-gal staining of whole eyes from control and *Sox9*$^{\Delta/\Delta}$-LacZ mice at 98 days after TX administration. Control corneas exhibited numerous LacZ$^+$ patches and elongated stripes originating from the limbus. In contrast, minimal X-gal staining was observed in *Sox9*$^{\Delta/\Delta}$-LacZ corneas, indicating impaired clonogenic capacity in *Sox9*-deleted cells. **(F)** Quantification of X-gal stained surface areas in control and *Sox9*$^{\Delta/\Delta}$-LacZ mice. Each data point represent an individual mouse. Mann-Whiitney U test P < 0.05. Scale bar in A represents 100 μm; Scale bar in B represents 50 μm; Scale bar in D represents 1 mm for the top row and 500 μm for the bottom row.

"regulation of epithelial cell migration" (Puri et al., 2020), "regulation of cell morphogenesis" and "stem cell differentiation", as well as gene pathways involved in these processes including "Wnt-signaling pathway" (Yu et al., 2012), "ERBB signaling" (Hassan and Seno, 2022), "cytokine mediated signaling pathway" (Korkaya et al., 2011) and "regulation of ERK1 and ERK2 cascade" (Lavoie et al., 2020; Fig. 5H; Table S10). Altogether, these results suggest that the population of SOX9-negative cells is more differentiated than that of SOX9-positive cells.

Sox9$^{\Delta/\Delta}$ progenitor cells lose their clonogenic capacity

Our results suggest that Sox9 is essential for limbal stem cell differentiation. To test this hypothesis, we used the same tamoxifen-inducible conditionally deleted Sox9 gene in adult mouse cells (two months old) that we used in the retina study. However, histological examination of the limbus and cornea from Sox9Δ/Δ mice at any time up to and including 100 DATX did not reveal any discernible phenotypic effects (Fig. 6A). Immunofluorescence staining for SOX9 (Fig. 6B) failed to reveal any differences in the number of SOX9-expressing cells between mutants and controls (Fig. 6C; Table S11;

control, 5.4 ± 0.4 cells/100 μm, n = 7; mutant, 5.9 ± 0.7 cells/100 μm, n= 6; p = 0.147,

440   Welch t test). Similar observations were made for corneal and limbal marker genes such

as *P63* (Figs. 6B and 6C; Table S11; control, 6.1 ± 0.7 cells/100 μm, n = 6; mutant, 6.1

± 0.7 cells/100 μm, n= 5; p = 0.147, Welch t test).and *Pax6* (Fig. S3A). These

observations are consistent with a model whereby tamoxifen induces mosaic patterns of

cell *Sox9*-deleting in the ocular surface, but that the *Sox9*-null cells cannot survive or

445   proliferate as well as their wild-type neighbors, and are hence outcompeted over time,

leading to an essentially wild-type cornea. We tested this model by leveraging this

mosaicism to investigate the *in vivo* clonal capacity of both *Sox9*-positive and *Sox9*-

negative cells. For this, we generated *CAGG-CreERTM;Sox9$^{flox/flox}$;R26R-LacZ* mice

(*Sox9$^{Δ/Δ}$-LacZ),* in which *Sox9*-deleted cells express the *LacZ* gene, and therefore, are

450   labelled in blue after X-gal staining (Fig. 6D). As controls, we used *CAGG-

CreERTM;Sox9$^{flox/+}$;R26R-LacZ* mice (*Sox9$^{Δ/+}$-LacZ*). Adult mice (two months old) were

fed with a tamoxifen-supplemented diet for 10 days to induce *LacZ* expression followed

by a chase period of 14 weeks. This time frame aligns with previous observations

reporting the appearance of clear X-gal^{+} stripes in *CAGG-CreERTM;R26R-LacZ* mice

455   *(Dor*à et al., 2015*)*. In control *Sox9$^{Δ/+}$-LacZ* corneas, derived from 4 mice we could

observe numerous discrete X-gal^{+} patches of different sizes spanning over the entire

corneal surface. In all cases, we observed a variable number of elongated stripes

originating from the limbus, suggesting that they come from limbal stem cells (Figs. 6E

and S3B). In contrast, the majority of *Sox9$^{Δ/Δ}$-LacZ* corneas, derived from 5 mice

460   exhibited minimal blue staining, with only occasional, small X-gal^{+} spots observed.

Notably, only one mutant cornea revealed the presence of an elongated stripe

originating from the limbus (Fig. 6E, right panels; Fig. S3B). A possible explanation for this clone may be that spontaneous ligand-independent activity of Cre-ER fusion may have occurred in a bona fide limbal stem cell, as previously reported (Vooijs et al., 2001;

465 Kemp et al., 2004; Haldar et al., 2009, Dorà et al., 2015). To quantify this difference, we measured the percentage of blue-stainied surface area, using the animal as the biological replicate. For control animals where both eyes were analysed, the average value was used. This analysis confirmed a significant decrease in the mutant corneas compared to the controls (controls: n = 4 mice $Sox9^{\Delta/+}$-LacZ, 6.74±2.33; mutants: n = 5

470 mice $Sox9^{\Delta/\Delta}$-LacZ, 0.85 ± 0.86; Mann-Whitney U-test, p = 0.0159; Fig. 6F; Table S12). These results reveal that during normal *in vivo* homeostasis, epithelial limbal and corneal stem/progenitor cells require *Sox9* to maintain their clonogenic capacity, indicating a key role for this gene in the process.

475 **DISCUSSION**

SOX9 is a master regulatory gene involved in the specification and/or development of embryonic tissues belonging to the three germ layers. In recent years, several studies revealed that its expression is maintained in adult ectoderm and endoderm derived cells indicating its role in the homeostasis of mature organs (Jo, 2014). Here, we analyze the

480 *in vivo* role of the gene in two structures belonging to the adult mouse eye, the retina and cornea, with completely different anatomy and biological functions. We conditionally deleted *Sox9* in the adult mouse retina and observed a phenotype characterized by different degrees of retinal degeneration. In the most affected individuals, this degeneration leads to the disappearance of the ONL layer and almost complete

485   absence of cone and rod photoreceptors. Since *Sox9* is expressed in MG and RPE cells (Poché et al., 2008), it is challenging to determine the contribution of each cell type to the final mutant phenotype*.* Previous studies on *Sox9*-mutant mice have not described any eye phenotype similar to the one we report here. The main reason is probably that these studies focused on the developing retina and are therefore not useful for

490   understanding how the absence of *Sox9* in MG and RPE cells in the adult eye leads to the retinal degeneration observed in our *Sox9*-deficient adult mutants. In this context, we can obtain better insights by examining the physiological response to RPE and MG cell loss after Cre-mediated activation of diphtheria toxin A (DTA) in these cell types. In the first case, the major retinal abnormality was the formation of regions of

495   photoreceptor rosetting with very mild degeneration of the photoreceptors (Longbottom et al., 2009). In mice with a DTA-induced ablation of MG cells, a wave of cell apoptosis was observed in the ONL but not in the other retinal layers (Shen et al., 2012). In our mutant mice, we did not observe photoreceptor rosetting in the ONL, but a massive apoptosis in this retinal layer. In addition, we observed that some regions within the INL

500   layer were devoid of MG cells. Thus, we favor the hypothesis that continuous expression of *Sox9* in MG cells is necessary for normal function and/or survival of this cell type. In the absence of *Sox9*, MG cells undergo massive retinal gliosis and they cannot provide structural, metabolic, and functional support to the other retinal cells, leading to a massive loss of photoreceptors followed by the depletion of other retinal cell

505   types. In agreement with this, *Sox9* expression is upregulated in MG cells after retinal light damage in rats (Wang et al., 2013). However, the retinal phenotype we report here is much more severe than the one observed in both DTA-induced transgenic lines. This

could be due to the different efficiency of Cre-induced recombination, but also to the fact that the combined deletion of *Sox9* in both MG and RPE may cause a more severe phenotype than the individual cell type-specific mutants.

An important consideration in our model is the potential contribution of non-cell autonomous mechanisms to photoreceptor degeneration. *Sox9* is expressed in both MG and RPE cells, and both cell types are known to support photoreceptor viability (Poché et al., 2008; Masuda et al., 2014). Notably, *Sox9* and *Otx2* cooperate to regulate visual cycle gene expression in the RPE (Masuda et al., 2014), and loss of *Otx2* specifically in the adult RPE leads to secondary photoreceptor degeneration through non-cell autonomous mechanisms (Housset et al., 2013). However, RPE-specific deletion of *Sox9* does not induce retinal degeneration and in fact results in *Otx2* upregulation (Masuda et al., 2014; Goto et al., 2018; Cohen-Tayar et al., 2018), suggesting that *Sox9* is not an upstream regulator of *Otx2* in this context. Further investigation into the molecular and cellular interactions between MG, RPE, and photoreceptors may help to clarify the indirect pathways contributing to degeneration in the absence of *Sox9*.

To our knowledge, no retinal abnormality has been described in patients with a mutation in *SOX9* (see Hejtmancik and Daiger, 2020). This is probably because CD patients are mostly heterozygous for *SOX9* and show early postnatal lethality, so a retinal degeneration phenotype in adulthood was either overlooked or not addressed. However, *SOX9/Sox9* has a complex regulatory landscape (Bagheri-Fam et al., 2006; Despang et al., 2019), and mutations in its regulatory region have been associated with tissue-specific human phenotypes (Wunderle et al., 1998; Benko et al., 2009; Kim et al., 2015; Kurth et al., 2009). In this context, transgenic mice carrying a human enhancer

located around 500 kb upstream of *SOX9* revealed enhancer activity in the eye (Sreenivasan et al., 2017). Thus, screening of the *SOX9* regulatory region may provide new molecular mechanisms underlying idiopathic cases of retinal degeneration.

Several expression studies have proposed *Sox9* as a marker of epithelial limbal and corneal stem/progenitor cells, with further support from *in vitro* experiments validating this finding (Sartaj et al., 2017; Parfitt et al., 2015; Menzel-Severing et al., 2018). Through *in vivo* cell lineage tracing and gene targeting strategies in mice, combined with the analysis of sc-RNAseq data, we shed light on the nature, differentiation dynamics and function of these stem/progenitor *Sox9*-positive cells. Trajectory analysis

540 of limbal and corneal scRNAseq data revealed that *Sox9* is highly expressed in a cell population characterized by the expression of stem cell markers (*Gpha2*, *Ifitm3*, *Cd63*, and *Krt14*), and that its expression gradually decreases as these cells enter mitosis and differentiate into suprabasal corneal cells. Consistent with this, we observed a substantial proliferation of *Sox9*-expressing descendant cells, leading to a significant

545 increase in their numbers within the first 10 days after the initiation of the TX treatment. Subsequently, these cells originate small clones with a rapid decrease and finally they give rise to few, large, radially oriented clones. The fact that *Sox9* was expressed exclusively in epithelial basal cells, while *Sox9*-expressing descendant cells were observed throughout the entire corneal epithelium indicates that many *Sox9*-expressing

550 descendant cells experience a decline in *Sox9* expression, initiating their differentiation into cells that proliferate, migrate, and differentiate into suprabasal corneal cells, thus contributing to the maintenance of corneal homeostasis.

The persistence of some small clones (including a radial stripe) of *LacZ*[+] *Sox9*[Δ/Δ] cells in the limbal and corneal epithelia of *CAGG-CreERTM;Sox9*[flox/flox]*; R26R-LacZ* mice

555   nevertheless shows that *Sox9* is not absolutely required for the contribution of cells to the limbal and ocular epithelium, but that the lack of *Sox9* significantly impairs the ability of mutant cells to compete with wild-type cells. This competition would explain why, in contrast to the whole mount *LacZ* analysis, *Sox9*-null cells were not detected in tissue sections of these eyes. A recent preprint (Rice et al, 2024) describes a study that

560   confirms the stem-clonogenic potential of *Sox9*-positive limbal epithelial cells. That study also reports that conditional deletion of *Sox9* leads to abnormal corneal epithelial differentiation and squamous metaplasia in the central cornea. This phenotype was not observed in our study but is explicable because of the mosaic nature of our model, which may allow for rescue of the normal Wnt and Notch signalling by wild-type cells,

565   preventing transdifferentiation (Menzel-Severing, 2018).

     We also observed long-lived circumferential clones that never exited the limbus. These latter clones are similar to those produced by OLB cells, as recently shown in lineage-tracing experiments, which predominantly self-renew and minimally contribute, if at all, to the formation of lineages maintaining the corneal epithelium during

570   homeostasis. However, in response to corneal injury, OLB proliferate and play an essential role in wound healing (Altshuler et al., 2021; Farrelly et al., 2021). Consistent with this, we saw that *Sox9* is highly expressed in the OLB cluster. Furthermore, previous studies utilizing an H2B-GFP/K5tTA mouse model have shown that *Sox9* is expressed in a corneal quiescent/slow-cycling cell population (Parfitt et al., 2015), and

575   wound healing assay using a human corneal organ culture revealed an increase in the

number of SOX9⁺ cells in both activated limbal and re-grown corneal epithelial cells (Menzel-Severing et al., 2018). We also observed that the number and extension of the limbal circumferential clones diminished in *Sox9^{Δ/Δ}-LacZ* corneas, indicating that *Sox9* may also control the proliferation of OLB stem cells.

580 This study highlights how a single transcription factor can regulate gene expression across neighboring cell types, driving distinct genetic programs and biological functions. In the retina, MG cells play a vital role in providing structural and metabolic support. Here, we demonstrate that *Sox9* is essential for the maintenance and survival of MG cells. In this context, previous studies have shown that Notch signaling, which is critical

585 for the maintenance of retinal progenitor cells and the specification of MG cells (Mills and Goldman, 2017), regulates *Sox9* expression. Moreover, *Sox9* has been shown to regulate the expression of type I collagen (COL1; Wang, 2013) and glial fibrillary acidic protein (GFAP), a key component of the intermediate filaments that form part of the MG cytoskeleton (Kinouchi, 2003). Thus, within the retina, *Sox9* appears to fulfill multiple

590 roles, including the regulation of structural protein expression and the promotion of cell survival through anti-apoptotic activity. This situation closely parallels that shown in postnatal and adult Sertoli cells, which die after *Sox9* ablation, leading to the complete degeneration of the testicular germinative epithelium. In these cells, *Sox9* likewise acts as an anti-apoptotic factor and regulates the expression of extracellular matrix

595 components (Barrionuevo et al, 2009; Barrionuevo et al, 2016). These facts suggest a role for *Sox9* in the maintenance of multiple adult tissues. In contrast, in the cornea, *Sox9* appears to be involved in the regulation of proliferation and differentiation of corneal epithelial stem/progenitor cells. However, the regulatory mechanisms and

downstream targets of *Sox9* in this context remain poorly understood. Here, we show

600   that high *Sox9* expression is associated with the presence of stem cell markers such as

Krt15, *P63*, and *Gpha2*, whereas its levels decline as stem cells undergo differentiation,

consistent with recent findings (Rice et al, 2024). Furthermore, *in vitro* studies using

limbal epithelial stem/progenitor cells have demonstrated that activation of the BMP,

Notch, and Shh pathways leads to upregulation of *Sox9*, while Wnt signaling

605   antagonizes its activity (Menzel-Severing, 2018). Thus, in the limbus, unlike its role in

the retina, *Sox9* appears to engage with key molecular pathways that govern stem cell

maintenance and differentiation. Future studies will help to elucidate the molecular

networks in which *Sox9* is involved in both contexts.

In summary, through functional analyses in mice, we have demonstrated multiple

610   roles for *Sox9* in the adult eye, acting as an essential maintenance factor preventing

retinal degeneration on the one hand, and promoting the differentiation of limbal cells in

the cornea on the other.

MATERIAL AND METHODS

615   **Mouse lines and crosses**

To obtain *Sox9* null mutant mice, we crossed *Sox9^{flox/flox}* (*B6.129S7-Sox9^{tm2Crm/J}*)

(Akiyama et al., 2002) to *CAGG-CreER^{TM}* (*B6.Cg-Tg^{(CAG-cre/Esr1*)5Amc}*) (Hayashi and

McMahon, 2002) mice, and the resulting double heterozygous offspring was

backcrossed to *Sox9^{flox/flox}* mice. Both mouse strains were sourced from The Jackson

620   Laboratory. Mutation occurs after treatment with tamoxifen, as described below. Cre-

negative *Sox9^{flox/flox}*, tamoxifen-treated mice served as controls. For corneal *Sox9* mosaic analyses we crossed *CAGG-CreER^{TM}; Sox9^{flox/flox}* mice *to R26R-LacZ (B6.129S4-Gt(ROSA)26Sor^{tm1Sor/J})* (Soriano, 1999), generating *CAGG-CreER^{TM}; Sox9^{flox/flox}; R26R-LacZ* mice. As control we used *CAGG-CreER^{TM}; Sox9^{flox/+}; R26R-LacZ* mice. For genetic lineage tracing assays, we used *Sox9^{IRES-CreERT2}* (*B6.129S7-Sox9^{tm1(cre/ERT2)Haak}*) (Soeda et al., 2010) mice obtained from the RIKEN BioResource Research Center (Tsukuba, Japan). These mice were crossed with either *R26R-LacZ* (*B6.129S4-Gt(ROSA)26Sor^{tm1Sor/J}*) (Soriano, 1999) or *R26R-EYFP* (*B6.129X1Gt(ROSA)26Sor^{tm1(EYFP)Cos/J}*) (Srinivas et al., 2001) reporter mice, resulting in double heterozygotes. Both of them were provided by The Jackson Laboratory (Bar Harbor, ME). All the experiments started with adult mice (2 months old). Mice were housed under Specific Pathogen-Free (SPF) conditions in the animal facilities of the Center for Biomedical Research (University of Granada, Granada, Spain). The animals had *ad libitum* access to food and water and were kept in groups. The occupancy density of the cages (microventilated) was in accordance with legal requirements. The cages were kept at a temperature of 22 +/- 2°C, a humidity of 55 +/- 10%, and a 12-hour/12-hour dark light cycle. Mice were provided with activity elements in the form of nest building material and hiding places. The animal procedures and housing conditions were approved by the University of Granada Ethics Committee for Animal Experimentation and the Consejería de Agricultura, Ganadería, Pesca y Desarrollo Sostenible of the Andalussian government, Junta de Andalucía (reference 12/12/2016/177).

Tamoxifen supplementation

645 For conditional deletion of *Sox9*, *CAGG-CreERTM;Sox9^flox/flox* mice were fed a standard diet (Harlan, 2914) supplemented with tamoxifen (TX; Sigma-Aldrich, St. Louis, MO, C8267) at a concentration of 40 mg TX/100 g diet for 10 days. For short-term tracing of *Sox9*-descendant cells (less than 10 days), TX (Sigma-Aldrich, St. Louis, MO, T5648) was dissolved in corn oil at a concentration of 30 mg/ml and administered

650 at a dose of 0.2 mg per gram of body weight through intraperitoneal injection to *Sox9^IRES-CreERT2* mice. For long-term tracing of *Sox9*-positive cells (more than 10 days), *Sox9^IRES-CreERT2* mice were administered TX as detailed above for the conditional deletion analyses .

Estimation of the percentage of tamoxifen-induced, Cre-mediated
655 **recombination and GFAP quantification**

To estimate the percentage of cells undergoing Cre-mediated recombination in TX-treated *CAGG-CreERTM;Sox9^flox/flox* mice, the total number of SOX9+ cells was divided by the total number of SOX8+ cells in a 20x microphotograph of a retinal section stained for SOX9 and SOX8.

660 In parallel, to quantify GFAP expression as a measure of MG reactivity, we analyzed GFAP immunofluorescence intensity across defined retinal surface areas. Given the cytoplasmic distribution of GFAP within glial processes, direct cell counting was not feasible. Instead, fluorescence intensity was measured using ImageJ, within full-thickness retinal regions in 20x microphotographs of a retinal sections stained for GFAP.

665 The total GFAP signal was normalized to the measured area for each section.

Histology and immunofluorescence

Eyes were dissected out and the crystalline lenses removed in order to facilitate histological sectioning. The eyes were fixed in 4% paraformaldehyde (PFA), dehydrated in increasing EtOH/saline solutions (EtOH + 0.9% Nacl), embedded in paraffin,
670   sectioned, and either stained with hematoxylin-eosin or processed for protein immunodetection. For simple and double immunofluorescence, sections were incubated overnight with the primary antibodies, washed, incubated with the appropriate conjugated secondary antibodies for 1 h at room temperature, and counterstained with 40,6-diamidino-2-phenylindole (DAPI). Several sections of eyes from control and mutant
675   mice were consistently mounted on the same slide and processed together. Parallel negative controls were performed in which the primary antibody was omitted. The following antibodies were used: anti-SOX8 (1/500), kindly provided by Dr. Wegner at the Universität Erlangen-Nurnberg (Germany); anti-OPN1SW (1/200, Santa Cruz, sc-14363); anti-OPN1LW (1/500, Chemicon, AB5405); anti-RHO (1/500, Sigma,
680   O4886); anti-BRN3α (1/100, Chemicon, MAB1585); anti-PAX6 (1/40, kindly provided by the Developmental Studies Hybridoma Bank at the University of Iowa), anti-AP2α (1/50, Abcam, ab108311), anti-SOX9 (1/400, Millipore, AB5535), anti-GFP (1/100, Novus Biologicals, NB600-308), anti-p63 (1/500, Master Diagnostica MAD-000479QD), and anti-S100 (prediluted 1/1, Master Diagnostica MAD-000592Q).

685   The two eyes of each mouse were analyzed in all experiments, and there were no notable differences between them (left-right) either histologically or in the expression pattern of the markers studied. Similarly, we found no differences between the eyes of control mice regardless of whether they were treated with TX or not.

Analysis of retinal and corneal cell death

690   Retinal and corneal cell death was assessed using the TUNEL Fluorescent In Situ Cell Death Detection Kit (Roche, Mannheim, Germany) according to the manufacturer's instructions.

Retinal flat whole-mount immunofluorescence

Eyes were fixed in 4% PFA and retinas were dissected in TBS containing 0.05%

695   Tween-20 and 0.1% Triton X-100 for 30 minutes at room temperature (RT). Retinas were then washed in TBS with 0.05% Tween-20 (TBS-T) and exposed to primary antibodies diluted PBS with 4% of BSA overnight at 4°C. After washing with TBS-T, the retinas were incubated with secondary antibodies diluted in PBS with 4% BSA for 3 hours at RT. Finally, the retinas were mounted flat for confocal microscopy.

700   ### Whole-mount X-gal staining

The eyes were dissected and cut transversely to facilitate solution penetration and whole-mount β-galactosidase histochemical reaction using X-gal as a substrate (X-gal staining) was performed. Whole-mount X-gal staining was performed overnight as described by Hogan et al. (1995). After staining, eyes were fixed in 4%

705   paraformaldehyde.

Estimation of the percentage of LacZ-positive corneal surface area

The surface area was calculated using the FIJI (V2.1.0) program based on ImageJ. The entire cornea surface was outlined, and the areas of the LacZ stripes were manually selected and marked. Then, the percentage of the blue-stained area relative

710   to the total corneal area was calculated. An automatic threshold could not be applied due to variations in brightness across the cornea pictures.

Imaging

Histological and immunofluorescence images were photographed using a DS-Fi1c camera installed on a Nikon Eclipse Ti microscope (Japan). Retinal flat mount images

715   were obtained using a high-speed spectral confocal microscope Nikon A1 ASHS-1. Subsequently, GIMP2 was used to adjust the color levels and the contrast of the immunofluorescence images. Whole-mount X-gal-stained eyes were photographed using a DP70 digital camera mounted on an Olympus SZX12 stereomicroscope (Japan).

720   **Single-cell RNA-sequencing dataset analyses**

We used the single-cell RNA-sequencing datasets of isolated epithelial cells from the limbus (with marginal conjunctiva and corneal periphery) generated by Altshuler and colleagues (Altshuler et al., 2021; Gene Expression Omnibus GSE167992). Bioinformatic analyses were performed with Seurat (version 5.0.3) (Hao et al., 2021)

725   following the author guidelines (https://satijalab.org/seurat/). Cells with unusually high and low numbers of unique feature counts or high mitochondrial counts were filtered out. The data were normalized and subsequently scaled. RunPCA was used for principal component analysis, and cells were clustered at a resolution of 0.6. Uniform manifold approximation and projection was used for nonlinear dimensional reduction.

730   Cluster-specific markers were selected with a minimum average log fold change threshold of 0.25 among genes that were expressed in a minimum of 25% of the cells. Cluster identity was assigned using the following markers: Conjunctiva: *Krt17, Krt4,*

Krt19, Krt6a, Krt13, Krt8 (and lack of *Krt12, Slurp1, Ppp1r3c*)*; cornea: *Krt12, Ppp1r3c, Slurp;* outer limbus: *Cd63, Ifitm3, Gpha2;* inner limbus: *Atf3, Socs3, Mt1;* basal cells:

735 *Itgb1, Itgb4, Ccnd1;* supra-basal cells: *Cldn4, Cdkn1a, Dsg1a;* and mitosis: *Mki67, Top2a, Ccna2.* For differential expression of cells expressing high and low levels of *Sox9*, we selected cells with high *Sox9*-expression levels (>= 0.7) and cells with low *Sox9*-expression levels (between 0.7 and 0.15) with the WhichCells and the SetIdent functions of the Seurat package. Differential expression analysis was performed with

740 the FindMarkers function (min.pct = 0.20, logfc.threshold = 0.01). For trajectory analysis we used the Python Scanpy suite (Haghverdi et al., 2016; Wolf et al., 2018) following the general PAGA trajectory inference workflow (https://scanpy.readthedocs.io/en/ stable/tutorials/trajectories/paga-paul15.html) with a Leiden resolution of 0.6. Cluster identity was assigned as described above, and the conjunctiva cluster was removed.

745 For pseudotime analysis (Wolf et al., 2019), the OLB cluster was chosen as the root. Gene Ontology analyses were done with the enrichGO and compareCluster functions of the clusterProfiler R package (Yu et al., 2012).

Statistical analysis

We analyzed datasets derived from distinct experimental conditions using appropriate

750 statistical tests based on preliminary assessments of normality and variance equality. When these conditions were not met, non-parametric approaches were favored for their robustness to data distribution assumptions. Specifically, we employed the Kruskal-Wallis rank-sum test to identify overall differences among group distributions followed by post-hoc Dunn's multiple comparison tests with Bonferroni adjustment to pinpoint

755 specific pairwise comparisons that contributed to these differences. When no significant

departures from normality or equality of variances were detected, parametric methods were deemed suitable. So, we conducted a standard Student t-test to evaluate the hypothesis of equal means between groups. Complete information on these statistical analyses are provided in Supplementary Tables S2-S8, S11 and S12.

ACKNOWLEDGEMENTS

This work has been funded by the following institutions: Programa Operativo FEDER Andalucía 2014-2020, Consejería de Economía, Conocimiento, Empresas y Universidad, Junta de Andalucía (Ref. A.BIO.106.UGR18) and MICIU/AEI 765 /10.13039/501100011033 and FEDER, UE, Ref PID2022-139302NB-I00 to RJ and FJB; Ministerio de Ciencia e Innovación, Instituto de Salud Carlos III, European Regional Development Fund through the program "Una manera de hacer Europa" (Ref. PI23/00335) to MA.

770 **AUTHORS CONTRIBUTIONS**

FJB, FDC, and RJ conceived the project; FJB, FDC, MB, and RJ designed the experiments; FJB, FDC, MB, RJ, AH, VL-S, ML, MAC-B, PL, ACC, YR-C, MA, and JMC performed laboratory experiments; FJB and MB carried out bioinformatic and statistical analyses; RJ, FJB, and MA collected the financial support; FJB, FDC and RJ wrote the 775 first draft of the manuscript and all authors contributed to the final version.

REFERENCES

Akiyama H, Chaboissier M-C, Martin JF, Schedl A, de Crombrugghe B. 2002. The transcription factor Sox9 has essential roles in successive steps of the chondrocyte differentiation pathway and is required for expression of Sox5 and Sox6. *Genes Dev* **16**:2813–2828. doi:10.1101/gad.1017802

Akiyama H, Lyons JP, Mori-Akiyama Y, Yang X, Zhang R, Zhang Z, Deng JM, Taketo MM, Nakamura T, Behringer RR, McCrea PD, Crombrugghe B de. 2004. Interactions between Sox9 and β-catenin control chondrocyte differentiation. *Genes Dev* **18**:1072–1087. doi:10.1101/gad.1171104

Altshuler A, Amitai-Lange A, Tarazi N, Dey S, Strinkovsky L, Hadad-Porat S, Bhattacharya S, Nasser W, Imeri J, Ben-David G, Abboud-Jarrous G, Tiosano B, Berkowitz E, Karin N, Savir Y, Shalom-Feuerstein R. 2021. Discrete limbal epithelial stem cell populations mediate corneal homeostasis and wound healing. *Cell Stem Cell* **28**:1248-1261.e8. doi:10.1016/j.stem.2021.04.003

Baden T, Euler T, Berens P. 2020. Understanding the retinal basis of vision across species. *Nat Rev Neurosci* **21**:5–20. doi:10.1038/s41583-019-0242-1

Bagheri-Fam S, Barrionuevo F, Dohrmann U, Günther T, Schüle R, Kemler R, Mallo M, Kanzler B, Scherer G. 2006. *Long-range* upstream and downstream enhancers control distinct subsets of the complex spatiotemporal *Sox9* expression pattern. *Dev Biol* **291**:382–397. doi:10.1016/j.ydbio.2005.11.013

Barnstable CJ, Hofstein R, Akagawa K. 1985. A marker of early amacrine cell development in rat retina. *Dev Brain Res* **20**:286–290. doi:10.1016/0165-3806(85)90116-6

Barrionuevo F, Bagheri-Fam S, Klattig J, Kist R, Taketo MM, Englert C, Scherer G. 2006. Homozygous Inactivation of Sox9 Causes Complete XY Sex Reversal in Mice1. *Biol Reprod* **74**:195–201. doi:10.1095/biolreprod.105.045930

Barrionuevo FJ, Hurtado A, Kim G-J, Real FM, Bakkali M, Kopp JL, Sander M, Scherer G, Burgos M, Jiménez R. 2016. Sox9 and Sox8 protect the adult testis from male-to-female genetic reprogramming and complete degeneration. *eLife* **5**:e15635. doi:10.7554/eLife.15635

Benko S, Fantes JA, Amiel J, Kleinjan D-J, Thomas S, Ramsay J, Jamshidi N, Essafi A, Heaney S, Gordon CT, McBride D, Golzio C, Fisher M, Perry P, Abadie V, Ayuso C, Holder-Espinasse M, Kilpatrick N, Lees MM, Picard A, Temple IK, Thomas P, Vazquez M-P, Vekemans M, Crollius HR, Hastie ND, Munnich A, Etchevers HC, Pelet A, Farlie PG, FitzPatrick DR, Lyonnet S. 2009. Highly conserved non-coding elements on either side of SOX9 associated with Pierre Robin sequence. *Nat Genet* **41**:359–364. doi:10.1038/ng.329

Bowes, C., Li, T., Danciger, M., Baxter, L. C., Applebury, M. L., & Farber, D. B. (1990). Retinal degeneration in the rd mouse is caused by a defect in the β subunit of rod cGMP-phosphodiesterase. *Nature*, **347**: 677-680

Casey MA, Lusk S, Kwan KM. 2023. Eye Morphogenesis in Vertebrates. *Annu Rev Vis Sci* **9**:221–243. doi:10.1146/annurev-vision-100720-111125

Chang NC. 2020. Autophagy and Stem Cells: Self-Eating for Self-Renewal. *Front Cell Dev Biol* **8**. doi:10.3389/fcell.2020.00138

Cohen-Tayar Y, Cohen H, Mitiagin Y, Abravanel Z, Levy C, Idelson M, Reubinoff B, Itzkovitz S, Raviv S, Kaestner KH, Blinder P, Elkon R, Ashery-Padan R. 2018. Pax6 regulation of Sox9 in the mouse retinal pigmented epithelium controls its timely differentiation and choroid vasculature development. *Development* **145**:dev163691. doi:10.1242/dev.163691

Despang A, Schöpflin R, Franke M, Ali S, Jerković I, Paliou C, Chan W-L, Timmermann B, Wittler L, Vingron M, Mundlos S, Ibrahim DM. 2019. Functional dissection of the Sox9–Kcnj2 locus identifies nonessential and instructive roles of TAD architecture. *Nat Genet* **51**:1263–1271. doi:10.1038/s41588-019-0466-z

Dorà NJ, Hill RE, Collinson JM, West JD. 2015. Lineage tracing in the adult mouse corneal epithelium supports the limbal epithelial stem cell hypothesis with intermittent periods of stem cell quiescence. *Stem Cell Res* **15**:665–677. doi:10.1016/j.scr.2015.10.016

Edqvist P-HD, Hallböök F. 2004. Newborn horizontal cells migrate bi-directionally across the neuroepithelium during retinal development. *Development* **131**:1343–1351. doi:10.1242/dev.01018

Ekström P, Sanyal S, Narfström K, Chader GJ, van Veen T. 1988. Accumulation of glial fibrillary acidic protein in Müller radial glia during retinal degeneration. *Invest Ophthalmol Vis Sci* **29**:1363–1371.

Farrelly O, Suzuki-Horiuchi Y, Brewster M, Kuri P, Huang S, Rice G, Bae H, Xu J, Dentchev T, Lee V, Rompolas P. 2021. Two-photon live imaging of single corneal stem cells reveals compartmentalized organization of the limbal niche. *Cell Stem Cell* **28**:1233-1247.e4. doi:10.1016/j.stem.2021.02.022

Fernández-Sánchez L, Lax P, Campello L, Pinilla I, Cuenca N. 2015. Astrocytes and Müller Cell Alterations During Retinal Degeneration in a Transgenic Rat Model of Retinitis Pigmentosa. *Front Cell Neurosci* **9**. doi:10.3389/fncel.2015.00484

Foster JW, Dominguez-Steglich MA, Guioli S, Kwok C, Weller PA, Stevanović M, Weissenbach J, Mansour S, Young ID, Goodfellow PN, Brook JD, Schafer AJ. 1994. Campomelic dysplasia and autosomal sex reversal caused by mutations in an SRY -related gene. *Nature* **372**:525–530. doi:10.1038/372525a0

Furuyama K, Kawaguchi Y, Akiyama H, Horiguchi M, Kodama S, Kuhara T, Hosokawa S, Elbahrawy A, Soeda T, Koizumi M, Masui T, Kawaguchi M, Takaori K, Doi R, Nishi E, Kakinoki R, Deng JM, Behringer RR, Nakamura T, Uemoto S. 2011. Continuous cell supply from a Sox9-expressing progenitor zone in adult liver, exocrine pancreas and intestine. *Nat Genet* **43**:34–41. doi:10.1038/ng.722

Goto S, Onishi A, Misaki K, Yonemura S, Sugita S, Ito H, Ohigashi Y, Ema M, Sakaguchi H, Nishida K, Takahashi M. 2018. Neural retina-specific Aldh1a1 controls dorsal choroidal vascular development via Sox9 expression in retinal pigment epithelial cells. *eLife* **7**:e32358. doi:10.7554/eLife.32358

Haghverdi L, Büttner M, Wolf FA, Buettner F, Theis FJ. 2016. Diffusion pseudotime robustly reconstructs lineage branching. *Nat Methods* **13**:845–848. doi:10.1038/nmeth.3971

Haldar M, Karan G, Tvrdik P, Capecchi MR. (2009). "Two cell lineages, myf5 and myf5-independent, participate in mouse skeletal myogenesis." Dev Cell 16: 928-938. doi:10.1016/j.devcel.2009.05.005

Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby C, Zager M, Hoffman P, Stoeckius M, Papalexi E, Mimitou EP, Jain J, Srivastava A, Stuart T, Fleming LM, Yeung B, Rogers AJ, McElrath JM, Blish CA, Gottardo R, Smibert P, Satija R. 2021. Integrated analysis of multimodal single-cell data. *Cell* **184**:3573-3587.e29. doi:10.1016/j.cell.2021.04.048

Hassan G, Seno M. 2022. ERBB Signaling Pathway in Cancer Stem Cells In: Tanabe S, editor. Cancer Stem Cell Markers and Related Network Pathways. Cham: Springer International Publishing. pp. 65–81. doi:10.1007/978-3-031-12974-2_3

Hayashi S, McMahon AP. 2002. Efficient recombination in diverse tissues by a tamoxifen-inducible form of Cre: a tool for temporally regulated gene activation/inactivation in the mouse. *Dev Biol* **244**:305–318. doi:10.1006/dbio.2002.0597

Hejtmancik JF, Daiger SP. 2020. Understanding the genetic architecture of human retinal degenerations. *Proc Natl Acad Sci* **117**:3904–3906. doi:10.1073/pnas.1922925117

Hogan B, Beddington R, Costantini F, Lacy E. 1995. Manipulating the Mouse Embryo.

Housset M, Samuel A, Ettaiche M, Bemelmans A, Béby F, Billon N, Lamonerie T. 2013. Loss of Otx2 in the adult retina disrupts retinal pigment epithelium function, causing photoreceptor degeneration. J Neurosci **33**:9890–904. doi:10.1523/JNEUROSCI.1099-13.2013.

Jo A, Denduluri S, Zhang B, Wang Z, Yin L, Yan Z, Kang R, Shi LL, Mok J, Lee MJ, Haydon RC. 2014. The versatile functions of Sox9 in development, stem cells, and human diseases. *Genes Dis* **1**:149–161. doi:10.1016/j.gendis.2014.09.004

Kemp CJ, Wheldon T, Balmain A. (2004). "p53-deficient mice are extremely susceptible to radiation-induced tumorigenesis." Nature Genetics 8:66-69. doi:10.1038/ng0994-66

Kim G-J, Sock E, Buchberger A, Just W, Denzer F, Hoepffner W, German J, Cole T, Mann J, Seguin JH, Zipf W, Costigan C, Schmiady H, Rostásy M, Kramer M, Kaltenbach S, Rösler B, Georg I, Troppmann E, Teichmann A-C, Salfelder A, Widholz SA, Wieacker P, Hiort O, Camerino G, Radi O, Wegner M, Arnold H-H, Scherer G. 2015. Copy number variation of two separate regulatory regions upstream of SOX9 causes isolated 46,XY or 46,XX disorder of sex development. *J Med Genet* **52**:240–247. doi:10.1136/jmedgenet-2014-102864

Kinouchi H, et al. (2003). "Induction of glial fibrillary acidic protein in Müller glia during retinal degeneration." Invest Ophthalmol Vis Sci 44: 2724–2731.

Knoblich JA. 2008. Mechanisms of Asymmetric Stem Cell Division. *Cell* **132**:583–597. doi:10.1016/j.cell.2008.02.007

Korkaya H, Liu S, Wicha MS. 2011. Regulation of Cancer Stem Cells by Cytokine Networks: Attacking Cancer's Inflammatory Roots. *Clin Cancer Res* **17**:6125–6129. doi:10.1158/1078-0432.CCR-10-2743

Kurth I, Klopocki E, Stricker S, van Oosterwijk J, Vanek S, Altmann J, Santos HG, van Harssel JJT, de Ravel T, Wilkie AOM, Gal A, Mundlos S. 2009. Duplications of noncoding elements 5′ of SOX9 are associated with brachydactyly-anonychia. *Nat Genet* **41**:862–863. doi:10.1038/ng0809-862

Lao M, Hurtado A, Chacón de Castro A, Burgos M, Jiménez R, Barrionuevo FJ. 2022. Sox9 Is Required for Nail-Bed Differentiation and Digit-Tip Regeneration. *J Invest Dermatol* **142**:2613-2622.e6. doi:10.1016/j.jid.2022.03.020

Lavker RM, Tseng SCG, Sun T-T. 2004. Corneal epithelial stem cells at the limbus: looking at some old problems from a new angle. *Exp Eye Res*, Special issue in honour of David Maurice **78**:433–446. doi:10.1016/j.exer.2003.09.008

Lavoie H, Gagnon J, Therrien M. 2020. ERK signalling: a master regulator of cell behaviour, life and fate. *Nat Rev Mol Cell Biol* **21**:607–632. doi:10.1038/s41580-020-0255-7

Longbottom R, Fruttiger M, Douglas RH, Martinez-Barbera JP, Greenwood J, Moss SE. 2009. Genetic ablation of retinal pigment epithelial cells reveals the adaptive response of the epithelium and impact on photoreceptors. *Proc Natl Acad Sci* **106**:18728–18733. doi:10.1073/pnas.0902593106

Marquardt T, Ashery-Padan R, Andrejewski N, Scardigli R, Guillemot F, Gruss P. 2001. Pax6 is required for the multipotent state of retinal progenitor cells. *Cell* **105**:43–55. doi:10.1016/s0092-8674(01)00295-1

Masuda T, Wahlin K, Wan J, Hu J, Maruotti J, Yang X, Iacovelli J, Wolkow N, Kist R, Dunaief JL, Qian J, Zack DJ, Esumi N. 2014. Transcription Factor SOX9 Plays a Key Role in the Regulation of Visual Cycle Gene Expression in the Retinal Pigment Epithelium *. *J Biol Chem* **289**:12908–12921. doi:10.1074/jbc.M114.556738

Menzel-Severing J, Zenkel M, Polisetti N, Sock E, Wegner M, Kruse FE, Schlötzer-Schrehardt U. 2018. Transcription factor profiling identifies Sox9 as regulator of proliferation and differentiation in corneal epithelial stem/progenitor cells. *Sci Rep* **8**:10268. doi:10.1038/s41598-018-28596-3

Mills WA, Goldman D. (2017). "Regulation of Müller glia reprogramming in development and regeneration." Curr Opin Neurobiol 42:68–75. doi:10.1016/j.conb.2016.11.006

Muto A, Iida A, Satoh S, Watanabe S. 2009. The group E Sox genes *Sox8* and *Sox9* are regulated by Notch signaling and are required for Müller glial cell development in mouse retina. *Exp Eye Res* **89**:549–558. doi:10.1016/j.exer.2009.05.006

Nadal-Nicolás FM, Jiménez-López M, Sobrado-Calvo P, Nieto-López L, Cánovas-Martínez I, Salinas-Navarro M, Vidal-Sanz M, Agudo M. 2009. Brn3a as a Marker of Retinal Ganglion Cells: Qualitative and Quantitative Time Course Studies in Naïve and Optic Nerve–Injured Retinas. *Invest Ophthalmol Vis Sci* **50**:3860–3868. doi:10.1167/iovs.08-3267

Ning W, Muroyama A, Li H, Lechler T. 2021. Differentiated daughter cells regulate stem cell proliferation and fate through intra-tissue tension. *Cell Stem Cell* **28**:436–452.

Parfitt GJ, Kavianpour B, Wu KL, Xie Y, Brown DJ, Jester JV. 2015. Immunofluorescence Tomography of Mouse Ocular Surface Epithelial Stem Cells and Their Niche Microenvironment. *Invest Ophthalmol Vis Sci* **56**:7338–7344. doi:10.1167/iovs.15-18038

Pellegrini G, Dellambra E, Golisano O, et al. (2001). "p63 identifies keratinocyte stem cells." Proc Natl Acad Sci USA 98:3156-3161. doi:10.1073/pnas.061032098

Poché RA, Furuta Y, Chaboissier M-C, Schedl A, Behringer RR. 2008. Sox9 is expressed in mouse multipotent retinal progenitor cells and functions in Müller Glial cell development. *J Comp Neurol* **510**:237–250. doi:10.1002/cne.21746

Puri S, Sun M, Mutoji KN, Gesteira TF, Coulson-Thomas VJ. 2020. Epithelial Cell Migration and Proliferation Patterns During Initial Wound Closure in Normal Mice and an Experimental Model of Limbal Stem Cell Deficiency. *Invest Ophthalmol Vis Sci* **61**:27. doi:10.1167/iovs.61.10.27

Rice G, Farrelly O, Huang S, Kuri P, Curtis E, Ohman L, Li N, Lengner C, Lee V, Rompolas P. 2024. Sox9 marks limbal stem cells and is required for asymmetric cell fate switch in the corneal epithelium bioRxiv 2024.04.08.588195; doi: https://doi.org/10.1101/2024.04.08.588195

Sartaj R, Zhang C, Wan P, Pasha Z, Guaiquil V, Liu A, Liu J, Luo Y, Fuchs E, Rosenblatt MI. 2017. Characterization of slow cycling corneal limbal epithelial cells identifies putative stem cell markers. *Sci Rep* **7**:3793. doi:10.1038/s41598-017-04006-y

Scott CE, Wynn SL, Sesay A, Cruz C, Cheung M, Gaviro M-VG, Booth S, Gao B, Cheah KSE, Lovell-Badge R, Briscoe J. 2010. SOX9 induces and maintains neural stem cells. *Nat Neurosci* **13**:1181–1189. doi:10.1038/nn.2646

Shen W, Fruttiger M, Zhu L, Chung SH, Barnett NL, Kirk JK, Lee S, Coorey NJ, Killingsworth M, Sherman LS, Gillies MC. 2012. Conditional Müller Cell Ablation Causes Independent Neuronal and Vascular Pathologies in a Novel Transgenic Model. *J Neurosci* **32**:15715–15727. doi:10.1523/JNEUROSCI.2841-12.2012

Sinha A, Fan VB, Ramakrishnan A-B, Engelhardt N, Kennell J, Cadigan KM. 2021. Repression of Wnt/β-catenin signaling by SOX9 and Mastermind-like transcriptional coactivator 2. *Sci Adv* **7**:eabe0849. doi:10.1126/sciadv.abe0849.

Soeda T, Deng JM, Crombrugghe B de, Behringer RR, Nakamura T, Akiyama H. 2010. Sox9-expressing precursors are the cellular origin of the cruciate ligament of the knee joint and the limb tendons. *genesis* **48**:635–644. doi:10.1002/dvg.20667

Soriano P. 1999. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet* **21**:70–71. doi:10.1038/5007

Sreenivasan R, Gordon CT, Benko S, de Iongh R, Bagheri-Fam S, Lyonnet S, Harley V. 2017. Altered *SOX9* genital tubercle enhancer region in hypospadias. *J Steroid Biochem Mol Biol*, Sex, Steroids, Fat and Breast Cancer **170**:28–38. doi:10.1016/j.jsbmb.2016.10.009

Srinivas S, Watanabe T, Lin C-S, William CM, Tanabe Y, Jessell TM, Costantini F. 2001. Cre reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. *BMC Dev Biol* **1**:4. doi:10.1186/1471-213X-1-4

Vooijs M, Jonkers J, Berns A. (2001). "A highly efficient ligand-regulated Cre recombinase mouse line shows that LoxP recombination is position dependent." EMBO Rep 2:292-297. doi:10.1093/embo-reports/kve062

Wagner T, Wirth J, Meyer J, Zabel B, Held M, Zimmer J, Pasantes J, Bricarelli FD, Keutel J, Hustert E, Wolf U, Tommerup N, Schempp W, Scherer G. 1994. Autosomal sex reversal and campomelic dysplasia are caused by mutations in and around the SRY-related gene SOX9. *Cell* **79**:1111–1120. doi:10.1016/0092-8674(94)90041-8

Wang X, Fan J, Zhang M, Ni Y, Xu G. 2013. Upregulation of SOX9 in Glial (Müller) cells in retinal light damage of rats. *Neurosci Lett* **556**:140–145. doi:10.1016/j.neulet.2013.10.005

West JD, Dorà NJ, Collinson JM. 2015. Evaluating alternative stem cell hypotheses for adult corneal epithelial maintenance. *World J Stem Cells* **7**:281–299. doi:10.4252/wjsc.v7.i2.281

Wolf FA, Angerer P, Theis FJ. 2018. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol* **19**:15. doi:10.1186/s13059-017-1382-0

Wolf FA, Hamey FK, Plass M, Solana J, Dahlin JS, Göttgens B, Rajewsky N, Simon L, Theis FJ. 2019. PAGA: graph abstraction reconciles clustering with trajectory inference through a topology preserving map of single cells. *Genome Biol* **20**:59. doi:10.1186/s13059-019-1663-x

Wunderle VM, Critcher R, Hastie N, Goodfellow PN, Schedl A. 1998. Deletion of long-range regulatory elements upstream of SOX9 causes campomelic dysplasia. *Proc Natl Acad Sci* **95**:10649–10654. doi:10.1073/pnas.95.18.10649

Yazdanpanah G, Jabbehdari S, Djalilian AR. 2017. Limbal and corneal epithelial homeostasis. *Curr Opin Ophthalmol* **28**:348. doi:10.1097/ICU.0000000000000378

Yu G, Wang L-G, Han Y, He Q-Y. 2012. clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. *OMICS J Integr Biol* **16**:284–287. doi:10.1089/omi.2011.0118

Yu M, Qin K, Fan J, Zhao G, Zhao P, Zeng W, Chen C, Wang A, Wang Y, Zhong J, Zhu Y, Wagstaff W, Haydon RC, Luu HH, Ho S, Lee MJ, Strelzow J, Reid RR, He T-C. 2024. The evolving roles of Wnt signaling in stem cell proliferation and differentiation, the development of human diseases, and therapeutic opportunities. *Genes Dis* **11**:101026. doi:10.1016/j.gendis.2023.04.042