Resting-state hyper- and hypo-connectivity in early schizophrenia: which tip of the iceberg should we focus on?

David Tomecek[1,2,3], Marian Kolenic[1,6], Barbora Rehak Buckova[1,3,4], Jaroslav Tintera[1,5], Filip Spaniel[1], Jiri Horacek[1,6,7], and Jaroslav Hlinka[1,3,*]

[1]National Institute of Mental Health, Klecany, Czech Republic
[2]Faculty of Electrical Engineering, Czech Technical University, Prague, Czech Republic
[3]Institute of Computer Science of the Czech Academy of Science, Prague, Czech Republic
[4]Donders Institute for Brain, Cognition and Behaviour, Nijmegen, Netherlands
[5]Institute of Clinical and Experimental Medicine, Prague, Czech Republic
[6]Third Faculty of Medicine, Charles University, Prague, Czech Republic
[7]Slovak Medical University, Bratislava, Slovak Republic

*Corresponding author: Jaroslav Hlinka

Head of the Department of Complex Systems
Deputy Director, Institute of Computer Science
Czech Academy of Sciences
Pod Vodarenskou vezi 2
Prague 8, 182 00, Czech Republic
Tel: (+420) 26605 3808
E-mail: hlinka@cs.cas.cz

Web: http://www.cs.cas.cz/hlinka/

January 19, 2026

Abstract

In this study, we explore the intricate landscape of brain connectivity in the early stages of schizophrenia, focusing on the patterns of hyper- and hypoconnectivity. Despite existing literature's support for altered functional connectivity (FC) in schizophrenia, inconsistencies and controversies persist regarding specific dysconnections.

Leveraging a large sample of 100 first-episode schizophrenia patients (42 females/58 males) and 90 healthy controls (50 females/40 males), we compare the functional connectivity across 90 brain regions of the Automated Anatomical Labeling atlas. We inspected the effects of medication and examined the association between FC changes and duration of untreated psychosis, duration of antipsychotic treatment, as well as symptom severity of the disorder. Our approach also includes a comparative analysis of three denoising strategies for functional magnetic resonance imaging data.

In patients, 15 region pairs exhibited increased FC, whereas 150 pairs showed reduced FC relative to controls. Despite this numerical asymmetry, the overall distribution of FC changes was relatively balanced: the median FC was not systematically shifted, indicating no global tendency toward either hyper- or hypoconnectivity. Notably, several FC alterations were significantly associated with variability in symptom severity and antipsychotic medication across patients.

Taken together, these results suggest a pattern of localized dysconnections embedded within an otherwise globally balanced change in connectivity profile in early schizophrenia. Importantly, this balance was substantially disrupted towards dominant observation of hypoconnectivity when less stringent denoising strategies were applied, with results increasingly dominated by hypoconnectivity, pointing to data preprocessing as a critical source of variability across studies.

1   Introduction

Although a neurodevelopmental hypothesis for schizophrenia complemented by neurodegenerative processes following the onset of psychosis is now well established, it is not clear how these processes are linked to the brain dysfunction underlying specific schizophrenia psychopathology (Dauvermann et al. 2017). The converging evidence from brain imaging studies indicates that these processes lead to disordered brain inter-regional functional connectivity (FC), and dysconnection represents the major candidate for the pathophysiological substrate of psychotic symptoms (Friston 2002). Concretely, the dysconnection concept suggests that the specific symptoms of schizophrenia can be described in terms of reduced or increased functional coupling between distinct brain regions (Friston & Frith 1995; Stephan et al. 2006). This concept is in line with earlier intuition referring to the title of the disease (i.e., "fragmented mind"), and it has been conceptualized in a neurobiological framework proposing a disruption of the anatomical

and functional connectivity between brain areas as the neurobiological correlate of altered information processing (Friston & Frith 1995). To quantify such connectivity disruption, the concept of functional connectivity, formalized as temporal covariation of neural signals between spatially disparate brain areas (Friston et al. 1993), is often used.

The functional dysconnection in schizophrenia has been reported to affect connections of a multitude of brain regions, such as the frontal lobe, including language areas (Backash et al. 2014; Benetti et al. 2009; Cole et al. 2011; Weinberger et al. 1994), sensory-motor cortex (Kaufmann et al. 2015), temporal and limbic structures (Crossley et al. 2009; Meyer-Lindenberg et al. 2005), and thalamus (Anticevic et al. 2014; Giraldo-Chica & Woodward 2017; Guller et al. 2012; Woodward & Heckers 2016; Woodward et al. 2012). Despite the fact that neuroimaging studies strongly support the role of altered FC in schizophrenia, these reported findings are highly inconsistent; specific regions associated with disconnection in schizophrenia still remain controversial, and a robust conclusion has not yet been obtained. Some of the probable reasons for these inconclusive results are the heterogeneity of clinical samples regarding the stage of illness (T. Li et al. 2016) and low sample sizes based on up to 40 participants on average (Dong et al. 2018). Indeed, a small sample size has been previously systematically documented to give rise to heterogeneous localized findings of white matter abnormalities in first-episode psychosis (Melicher et al. 2015). We further conjecture that variability in the data preprocessing and analysis approaches might also contribute to the variability of the observed results (Burgess et al. 2016). The majority of previous rs-fMRI studies in schizophrenia reported hypoconnectivity between brain regions (Dong et al. 2018; S. Li et al. 2019), but hyperconnectivity has also been repeatedly observed in various disease stages (T. Li et al. 2016; Dong et al. 2018; González-Vivas et al. 2019; Anticevic et al. 2015). Longitudinal studies of FES subjects reported (partial) normalization in terms of increased fMRI activation after antipsychotic treatment (González-Vivas et al. 2019). Another factor responsible for the heterogeneity of the findings is the methodological approaches focused on FC between a priori preselected regions, as contrasted with a whole-brain analysis based on the parcellation according to suitable anatomical or functional brain atlases. Given these principal differences, the reported findings are strongly influenced by the a priori choice of the regions of interest, which in some cases may not include all the pathophysiologically relevant areas. On the other hand, whole-brain studies may suffer from the fact that the correction for multiple comparisons could bring false negative results, specifically in the case of underpowered studies. Our study aimed to remediate some of the above-mentioned inconclusiveness and to utilize the whole-brain FC analysis on a large cohort of 100 first-episode schizophrenia (FES) patients, ensuring high homogeneity in terms of chronicity and exposure to medication. The connectivity was calculated between all 90 brain regions comprising the Automated Anatomical Labeling

(AAL) template image (Tzourio-Mazoyer et al. 2002) and compared with a group of 90 healthy controls. On the basis of the previous evidence, we hypothesized that schizophrenia would show decreased connectivity between the candidate brain regions. Furthermore, we tested whether FC changes were moderated by the effect of medication and whether they were linked to clinical factors such as positive and negative symptoms of schizophrenia. In addition, we evaluated two different denoising strategies of the fMRI data, as this step potentially affects the resulting FC measures (Burgess et al. 2016).

2 Methods

2.1 Study overview and samples

In total, 190 subjects participated in the study: 100 FES patients (mean age=28.78, SD=6.84, 42 females/58 males) and 90 healthy volunteers serving as controls (mean age=27.88, SD=6.83, 50 females/40 males). There were no significant differences between the patient and control samples in age and sex. In the patient group, at the time of the MRI scan, the average duration of untreated psychosis was 3.23 months (SD = 4.79), and the average duration of antipsychotic treatment was 2.29 months (SD = 4.55). The study design was approved by the local Ethics Committee of the Institute of Clinical and Experimental Medicine and the Psychiatric Center Prague. All subjects provided written informed consent after receiving a complete description of the study.

The FES patients were diagnosed according to ICD-10 criteria and structured MINI International Neuropsychiatric Interview (Sheehan et al. 1998). FES subjects were investigated during their first hospitalization in the Prague psychiatric hospitals with a catchment area of 1 million inhabitants. Patients were considered as FES if they fulfilled these criteria: a) first hospitalization for schizophrenia, and b) clinical interview identified first psychotic and/or prodromal symptoms of psychosis not earlier than 24 months ago (mean=5.90 months, SD=6.16).

The resting fMRI was performed at the initial stage of second-generation antipsychotic therapy (mean 10 weeks of medication at the time of rsfMRI). The mean dose of chlorpromazine equivalents (Woods 2003) was 388.87 mg (SD=235.44) per day. Psychometrics included the Positive and Negative Symptom Scale (PANSS) (Kay et al. 1987). Ninety healthy control subjects (HC) were recruited via a local advertisement; they had a similar socio-demographic background as the FES to whom they were matched by age and sex.

The HC had a slightly higher number of years of education than the FES (HC: mean=16.56, SD=2.33; FES: mean=13.51, SD=2.46; t=7.033, p<0.001). HC were evaluated with MINI (Sheehan et al. 1998) and were excluded if they had a lifetime

history of any psychiatric disorder or a family history of psychotic disorders. Other exclusion criteria for both groups included a history of seizures or significant head trauma, mental retardation, a history of substance dependence, and any MRI contraindications. Written informed consent was obtained from all participants.

2.2   fMRI data acquisition

Scanning was performed using a 3T Siemens Magnetom Trio MRI scanner located at the Institute of Clinical and Experimental Medicine in Prague, Czech Republic. Functional images were obtained using T2*-weighted gradient echo-planar imaging (GR-EPI) sequence with blood oxygenation level-dependent (BOLD) contrast with the following parameters: repetition time (TR)=2000 ms, echo time (TE)=30 ms, flip angle=70°, voxel size=3×3×3 mm, matrix size 48×64 voxels, 35 axial slices acquired continuously in sequential decreasing order covering the entire cerebrum, 400 functional volumes in total. Also, for anatomical reference, a three-dimensional T1-weighted high-resolution magnetization-prepared rapid gradient echo (MPRAGE) image was acquired from each participant using the following parameters: repetition time (TR)=2300 ms, echo time (TE)=4.63 ms, flip angle=10°, voxel size=1×1×1 mm, matrix size 256×256 voxels, 224 sagittal slices covering the entire brain.

2.3   Data preprocessing, brain parcellation, and FC analysis

Functional MRI is a neuroimaging method that is based on measuring blood oxygen level-dependent (BOLD) signal (Ulmer & Jansen 2020). One of the typical features of the fMRI data is the noise that is present in the raw BOLD signal (Biswal et al. 1995; Raichle 2015), which significantly limits the reliability of functional connectivity measures (Shirer et al. 2015). Typical artifacts, such as subject movements, arterial pulsation, respiration, and also the hardware of the MRI scanner itself, induce non-neural temporal correlations in the BOLD (Pamilo et al. 2015), and relatively sophisticated data preprocessing is warranted to maximize the level to which the functional connectivity estimates reflect the underlying neuronal dynamics.

First, the structural and functional images were converted from DICOM to NIFTI format using the dcm2niix tool (X. Li et al. 2016). The following preprocessing steps were performed with the functional data: functional realignment and unwarp; slice-timing correction; outlier identification; direct segmentation and normalization; and functional smoothing - pipeline labeled as "default preprocessing pipeline for volume-based analyses

(direct normalization to MNI-space)" in CONN toolbox (Whitfield-Gabrieli & Nieto-Castanon 2012).

The denoising steps included the regression of six head-motion parameters with their first-order temporal derivatives and five principal components of WM and CSF signals using a component-based noise correction method (CompCor) implemented in the CONN toolbox (Behzadi et al. 2007). Time series from defined regions of interest were additionally linearly detrended and finally filtered by a band-pass filter with cutoff frequencies 0.008-0.09 Hz. We shall refer to this preprocessing setup as the *stringent* denoising scheme.

As an alternative denoising pipeline, closer to the practice in some studies (Argyelan et al. 2014; Wang et al. 2015), we used a *moderate* denoising scheme including six head-motion parameters without their first-order derivatives and only the mean time series of WM and CSF. This pipeline was performed without explicit linear detrending. However, time series were also finally filtered by a band-pass filter with cutoff frequencies 0.008-0.09 Hz.

To isolate the impact of denoising, we conducted another analysis on the data that underwent the same default preprocessing pipeline in the CONN toolbox (Whitfield-Gabrieli & Nieto-Castanon 2012) as described above, while explicitly omitting all denoising procedures. This variant is referred to as the *raw*.

2.4   Analysis

For each subject, to quantify the whole-brain pattern of functional connectivity, we performed an ROI-to-ROI connectivity analysis by computing Pearson's correlation matrix among the regional mean time series extracted from 90 brain regions (excluding the cerebellar regions) of the Automated Anatomical Labeling (AAL) atlas (Tzourio-Mazoyer et al. 2002) using the CONN toolbox (Whitfield-Gabrieli & Nieto-Castanon 2012). The resulting connectivity matrices were represented as Fisher's z-transformed Pearson's correlation coefficients. Although participants were balanced in terms of age and sex, these variables were further included as nuisance covariates in the linear regression model to further minimize their potential influence on functional connectivity.

We used a two-sample Mann-Whitney U-test to evaluate differences between groups. Spearman's correlation was used to determine the relation between the four subscales of the PANSS scale (Kay et al. 1987), antipsychotic medication at the time of the MRI scan (measured by chlorpromazine equivalent), duration of untreated psychosis, duration of

antipsychotic treatment, and functional connectivity in patients. To increase the statistical analysis power, these analyses were limited to ROI pairs that showed a significant effect of disease (i.e., effect in the initial between-group comparison). The resulting p-values from each analysis were corrected for multiple testing by controlling the false discovery rate (FDR) (Benjamini & Hochberg 1995). Additional analyses with the moderate and raw denoising schemes were performed using the same analytical pipeline.

3   Results

3.1   The difference between healthy controls and schizophrenia patients

In the group comparison using the default, stringent denoising scheme, a difference was observed in a multitude of ROI pairs. 150 ROI pairs had significantly greater functional connectivity (FDR-corrected $p<0.05$) in patients, while healthy volunteers exhibited significantly greater (FDR-corrected $p<0.05$) functional connectivity in 15 ROI pairs compared to patients (Figure 1). Despite the numerical prevalence of observed hyperconnections in patients, healthy volunteers, and patients did not show a difference in the median functional connectivity (median FC in patients=0.0709, MAD=0.0182, median FC in controls=0.0653, MAD=0.0209, u=4007 (z-score=-0.8663), p=0.1932).

The most increased links in patients included a range of connections involving particularly the bilateral thalamus, right middle cingulate gyrus, right supramarginal gyrus, left middle temporal gyrus (see Table S1 for the full list of affected connections). In the opposite direction, the decreased links in patients comprised especially the left supplementary motor area, bilateral posterior cingulate gyrus, right superior temporal gyrus, and left middle cingulate gyrus (see Table S2 for the full list of affected connections). See Figure 2 for a graphical overview of the group comparison of healthy controls and patients, including its stability with respect to additional control for inter-subject variability in the amount of head motion; see Discussion for details.

3.2   Association between symptom severity, medication, duration of untreated psychosis, duration of antipsychotic treatment, and functional connectivity

We have observed a significant correlation between symptom severity measured by PANSS and functional connectivity, see Figure 3. To increase the power by focusing on the most relevant candidate connections, we used a masking matrix of p-values from a group comparison of healthy volunteers and patients to distinguish regions with

significantly greater functional connectivity in patients. PANSS positive scale significantly correlated with the functional connectivity between the right precentral gyrus and the right middle cingulate gyrus. PANSS negative scale significantly correlated with the functional connectivity between the right middle cingulate gyrus and the left lingual gyrus.

Figure 1: Difference between HC and FES; higher FC in healthy controls = blue, higher FC in patients = red, p<0.05 (FDR-corrected).

Figure 2: Difference between HC and FES. The significant differences after the p<0.05 FDR correction are shown in blue (higher FC in patients) and pink (higher FC in healthy controls). A) Default analysis; B) Analysis including additional inter-subject correction for the amount of motion; C) Analysis including additional inter-subject correction for the amount of motion, as well as rejecting outliers. See the Discussion section for details of the additional analysis shown in the B and C panels.

Figure 3: Association between antipsychotic medication, symptom severity, and functional connectivity. Antipsychotic medication significantly correlated with the functional connectivity of 12 ROI pairs (red links), PANSS Positive scale significantly correlated with the functional connectivity of one ROI pair (green link), and PANSS Negative scale significantly correlated with the functional connectivity of one ROI pair (blue link). Results with p<0.05 (FDR-corrected) are shown.

Further, we have observed a significant correlation between antipsychotic medication and functional connectivity. Regions with the most links included the bilateral thalamus, left

middle cingulate gyrus, right middle temporal gyrus, and right middle temporal pole; see Table 1 for the full list of connections.

We have also observed a significant correlation between antipsychotic medication and symptom severity measured by PANSS Total (Spearman r=0.33, p<0.001).

Conversely, we found no relationship between the duration of untreated psychosis, duration of antipsychotic treatment, and functional connectivity.

Region 1	Region 2	Spearman r
Antipsychotic medication		
Thalamus, Left	Middle Temporal Pole, Right	0.33
Superior Frontal Gyrus, Left	Rolandic Operculum, Right	0.32
Thalamus, Right	Middle Temporal Gyrus, Right	0.31
Thalamus, Right	Middle Temporal Pole, Right	0.31
Inferior Frontal Gyrus, Triangular Part, Left	Medial Orbital Frontal Cortex, Right	0.31
Lingual Gyrus, Right	Thalamus, Left	0.30
Lingual Gyrus, Left	Precuneus, Left	0.30
Paracentral Lobule, Left	Thalamus, Right	0.29
Precentral Gyrus, Right	Middle Cingulate Gyrus, Left	0.29
Thalamus, Left	Middle Temporal Gyrus, Right	0.27
Middle Cingulate Gyrus, Left	Postcentral Gyrus, Right	0.27
Paracentral Lobule, Right	Thalamus, Right	0.27
PANSS Positive		
Precentral Gyrus, Right	Middle Cingulate Gyrus, Right	0.35
PANSS Negative		
Middle Cingulate Gyrus, Right	Lingual Gyrus, Left	0.33

Table 1: Correlation between antipsychotic medication, PANSS Positive, PANSS Negative, and functional connectivity in patients; note that only p<0.05 (FDR-corrected) are reported.

3.3 Effect of data-denoising on functional connectivity

To evaluate the stability of the above-presented findings with respect to the strictness of denoising, we also performed our analysis with moderate and raw denoising settings. In the moderate denoising, healthy volunteers exhibited significantly higher (p<0.05 FDR-corrected) functional connectivity in 1098 ROI-pairs compared to patients. Conversely, 43 ROI pairs had significantly higher (p<0.05 FDR-corrected) functional connectivity in patients, see Figure 5B. The most increased links in patients included the bilateral thalamus, bilateral superior temporal gyrus, and left postcentral gyrus (see Table S3 for the full list of affected connections). In the opposite direction, the most decreased links in patients involved the right precentral gyrus, bilateral postcentral gyrus, left rolandic operculum, and the right paracentral lobule (see Table S4 for the full list of affected connections).

In stark contrast to the stringent denoising scheme, the median functional connectivity was greater in patients for the moderate denoising scheme. The median connectivity difference between healthy controls and patients in the moderate scheme was 0.0349 (median FC in patients=0.1388, MAD=0.0338, median FC in controls=0.1737, MAD=0.0477, u=5762 (z-score=-3.1351), p<0.001). Indeed, the disease effect on functional connectivity indices was clearly biased towards hypoconnectivity in the moderately denoised dataset, while there was a relatively balanced hypoconnectivity and hyperconnectivity (with a slight prevalence of hyperconnectivity) observed when using the default stringent denoising scheme, see Figure 4C.

In the case of moderate denoising, we found a significant correlation between functional connectivity and antipsychotic medication. Regions included the bilateral thalamus, bilateral rolandic operculum, and bilateral lingual gyrus (see Table S5 for the full list of affected connections).

In the raw denoising, healthy volunteers exhibited significantly higher (p<0.05 FDR-corrected) functional connectivity in 1281 ROI-pairs compared to patients; see Table S6 for the full list of affected connections. Conversely, no ROI pairs had significantly higher (p<0.05 FDR-corrected) functional connectivity in patients, see Figure 5A. The median connectivity difference between healthy controls and patients in the raw scheme was 0.0611 (median FC in patients=0.5025, MAD=0.1132, median FC in controls=0.5636, MAD=0.1190, u=5509 (z-score=-2.4224), p<0.01). The disease effect on functional connectivity indices was thus clearly biased towards hypoconnectivity in

the raw scheme, see Figure 4C. We also observed no significant relation between functional connectivity and any of the clinical variables.

Despite these vast differences in the overall shift of FC between the two populations, the spatial localization of the disease effect observed with the three denoising schemes was strongly correlated (raw and stringent: Spearman r=0.25, p<0.001; moderate and stringent: Spearman r=0.48, p<0.001), see Figure 4A and 4B respectively for the scatterplot across region pairs.

Figure 4: Comparison of the effect of disease on functional connectivity observed with the raw, moderate, and stringent denoising. A) Scatterplot of the median FC difference between patients and healthy controls in the raw and stringent scheme. B) Scatterplot of the median FC difference between patients and healthy controls in the moderate and stringent scheme. Black lines mark identity, and the red line corresponds to the best linear fit. Each dot corresponds to one pair of regions, showing the difference in their medians

in the two denoising schemes. C) Violin plot of the FC differences between healthy controls and patients in the raw, moderate, and stringent denoising schemes.

Figure 5: Difference between HC and FES for the A) raw and B) moderate denoising variants. The differences significant after the p<0.05 FDR correction are shown in blue (higher FC in patients) and pink (higher FC in healthy controls).

4   Discussion

Using the stringent denoising scheme, we observed that individuals with FES showed a pattern of focal hyperconnectivity in thalamic, basal ganglia, and cingulo-sensorimotor circuits, combined with cortical hypoconnectivity. This profile aligns with models of schizophrenia that conceptualize early psychosis as a disturbance of hierarchical cortico–subcortical integration, with thalamic and striatal systems exhibiting increased (possibly compensatory) reactivity against a background of impaired cortical coordination (Anticevic et al. 2015; Wengler et al. 2020). This hybrid pattern replicates prior work documenting thalamo–sensorimotor hyperconnectivity in schizophrenia (Anticevic et al. 2015; Woodward et al. 2012) and suggests an early imbalance in hierarchical predictive processing.

The most increased links in patients involved the bilateral thalamus, globus pallidus, bilateral middle cingulate gyrus, bilateral supramarginal gyrus, right superior occipital gyrus, and bilateral precuneus. These findings mirror the "thalamocortical dysconnectivity" framework proposed by Anticevic et al. (2014) and the results of meta-analyses reporting increased thalamic coupling to sensorimotor cortices in psychosis. According to these models, thalamic hyperconnectivity may reflect insufficient top-down regulation from prefrontal and hippocampal systems, leading subcortical hubs to over-signal or over-synchronize with cortical targets. Notably, a subset of the

hyperconnected edges identified here was also implicated in symptom severity and medication effects, suggesting that these subcortical–cingular–motor circuits may represent a core network phenotype of early schizophrenia.

In contrast, the most decreased links in patients involved circuits linking the supplementary motor area and precentral cortex with the hippocampus, caudate, and putamen. This pattern is consistent with meta-analytic evidence for widespread hypoconnectivity across fronto-parietal, hippocampal, and cortico-striatal networks in schizophrenia (Dong et al. 2018; S. Li et al. 2019). These networks support predictive processing, contextual integration of prior information, and precision-weighted gating of sensory signals (Friston 2005; Corlett et al. 2019). Their reduced functional integration in FES aligns with computational theories proposing that weakened cortical priors and impaired hippocampal-prefrontal coordination degrade the system's ability to maintain stable internal models and suppress noisy bottom-up prediction errors (Sterzer et al. 2018; Sigurdsson & Duvarci 2016). In this context, thalamic and striatal hyperconnectivity may arise as maladaptive attempts to stabilize cortical dynamics in the face of insufficient cortico-hippocampal control, a view that is compatible with theoretical work showing that hierarchical heterogeneity of cortical microcircuits shapes large-scale neural dynamics and stability (Demirtaş et al. 2019). Our findings also align with longitudinal evidence in antipsychotic-naive FES showing widespread baseline dysconnectivity and that antipsychotic exposure in antipsychotic-naive FES is associated with increased functional connectivity primarily between the thalamus and the rest of the brain (Chopra et al., 2021). A striking and unifying observation across all analyses was the role of the mid-cingulate cortex (MCC). MCC appeared as a major hyperconnected hub in patients, but also as the key node associated with both PANSS symptom dimensions and CPZ-equivalent antipsychotic exposure (see below). This convergence suggests MCC as the principal clinically relevant region in our dataset. The MCC is an integrator of salience, motor preparation, effort allocation, and executive control, functions that are highly disrupted in psychosis. Its involvement here strongly supports theories emphasizing abnormalities in salience signaling and motor relevance attribution in schizophrenia (Shackman et al. 2011; Hoffstaedter et al. 2014; Kapur 2003). This is further supported by studies demonstrating aberrant recruitment of the medial cingulate/ACC region during cognitive control tasks in schizophrenia during cognitive-control and motor-preparatory tasks in psychosis (Heckers et al. 2004).

Furthermore, brain hyperconnectivity correlated with both symptom severity as well as antipsychotic medication in subjects at their first episode of psychosis. The positive symptom dimension (PANSS Positive) was associated with FC between right precentral gyrus and right MCC, indicating enhanced coupling between motor cortex and a core salience/motor-control hub (Shackman et al. 2011; Du et al. 2019; Kapur 2003). This

pattern is highly compatible with Kapur's aberrant salience model and with contemporary models linking psychosis to dysregulated interactions between salience networks and motor/output systems. Conversely, the negative symptom dimension (PANSS Negative) was associated with functional connectivity between the right MCC and the left lingual gyrus in our sample, suggesting altered coupling between a cingulate salience/control hub and visual–perceptual cortex. While this specific MCC–lingual link is not a canonical finding, it is consistent with broader evidence that negative symptom dimensions relate to salience-network dysfunction and abnormal salience–default mode (and salience–sensory) communication in schizophrenia (Hare et al. 2019; Palaniyappan & Liddle 2012). These findings are consistent with the literature suggesting that negative symptom dimensions are associated with cingulate/salience-network dysfunction and with altered coupling between salience/control systems and sensory–perceptual networks, potentially contributing to impaired contextual appraisal.

This pattern most likely reflects a combination of illness severity, given that CPZ-equivalent dose is clinically titrated and therefore is often confounded with positive symptom severity in naturalistic samples (Troisi et al. 1997; Ringen et al. 2019), and medication-driven modulation of salience-related motor circuitry, in line with evidence showing antipsychotic effects on thalamo-cortical and cingulate networks (Chopra et al. 2021; Sarpal et al. 2015). Importantly, Chopra et al. demonstrated that DMN–limbic normalization can occur even without medication, whereas drug exposure specifically alters thalamo-cingulate circuits. Our findings extend this by showing that in a naturalistic FES sample, antipsychotic exposure was most strongly associated with the same motor–salience connections that also tracked positive symptom severity in our data. The overlap between CPZ-related and PANSS Positive-related connectivity further supports the interpretation that apparent medication effects may partly reflect underlying symptom burden.

It is worth mentioning some technical considerations of the present study. Several denoising strategies were discussed in recent papers. While a unique consensus has not been reached in the field, the implementation of a more stringent denoising scheme has become a standard practice (Power et al. 2014; Patel et al. 2014), motivating our default choice. To quantify functional connectivity, in line with common practice, we use Pearson's linear correlation coefficient, as it was shown to be sufficient under standard conditions, as with the current dataset (Hlinka et al. 2011). A possibly fruitful further direction would indeed lie in extension to mapping the consequences or analogies of our findings to other network characteristics previously shown to be altered in clinical high-risk and early-illness schizophrenia individuals, in particular to dynamic functional connectivity (Mennigen et al. 2019), and amplitude of low frequency fluctuations or graph-theoretical functional dysconnectivity characterization (Tang et al. 2019). On the

other side, utilizing the full FC matrix has shown an advantage in machine learning classification of schizophrenia above and beyond some advanced methods such as topological data analysis (Caputi et al. 2021).

There was a considerable difference between the stringent, moderate, and raw denoising schemes in the number of significant ROI pairs in each group comparison. Both raw and moderate denoising schemes resulted in a higher number of ROI pairs that showed significantly higher FC in healthy controls compared to patients (1281/0 in raw and 1098/43 in moderate), while in the stringent denoising scheme, the proportion changed as we observed a slightly higher number of ROI pairs exhibiting hyperconnectivity in patients (15/150). Note that the use of a substantially stricter denoising scheme than used in a range of studies in the literature may explain some of the deviation of our results with respect to the field. Despite the differences between denoising strategies, median connectivity matrices are strongly correlated within each group (the correlation of FC between raw and stringent was 0.5301 ($p<0.001$) in healthy individuals and 0.5681 ($p<0.001$) in patients, and between moderate and stringent was 0.8798 ($p<0.001$) in healthy individuals and 0.9181 ($p<0.001$) in patients. Also, the difference between the groups, i.e., the disease effect, is strongly correlated (across region pairs) between the denoising strategies (raw and stringent: Spearman $r=0.25$, $p<0.001$; moderate and stringent: Spearman $r=0.48$, $p<0.001$), see Figure 4A and 4B, respectively, for the scatterplot across region pairs. Although we found overlapping ROI-pairs between the stringent and moderate denoising that exhibited correlation with antipsychotic medication (namely, regions involving the bilateral thalamus, right middle temporal pole, right middle temporal gyrus, and right lingual gyrus), we found no relationship between functional connectivity and any of the selected clinical variables when the raw data were used, confirming that minimal standard preprocessing is needed to reduce noise in the data.

To ascertain that the results were not driven by any residual artifacts due to differences in head motion, we also carried out a control analysis where mean framewise displacement (FD), the aggregated measure of subject motion, was included as a covariate of no interest. The results again remained qualitatively equivalent, suggesting that the effect of motion on FC is relatively small compared to the effect of disease; see Figure 2B. The results remained largely unchanged after the additional removal of the 14 subjects who showed maximum head motion larger than 1.5×IQR (interquartile range), the Tukey criterion for detection of outliers (Tukey 1977); see Figure 2C. The difference in the number of ROI pairs with significantly changed FC could be due to lower degrees of freedom in the additional corrected analyses.

5   Conclusions

Our study using a whole-brain functional connectivity approach on a relatively large sample has shown a relatively balanced, yet spatially resolved, picture of hyperconnectivity and hypoconnectivity in first-episode psychosis while demonstrating a significant association between the affected connection strengths and clinical symptoms and cumulative dosage of antipsychotic medication. Our results support a model in which early schizophrenia combines cortical hypoconnectivity with subcortical and cingulo-sensorimotor hyperconnectivity, forming a hybrid dysconnectivity profile that is closely tied to symptoms and treatment. Importantly, we have shown that this balanced picture gets substantially skewed in the direction of dominantly observed hypo- or dysconnectivity when applying more moderate data-denoising schemes, while the relation to symptoms is rendered insignificant. While the results of all approaches are in overall qualitative agreement, including with localization of most commonly reported disease connectivity effects in the literature, taking into account the results obtained with the current stringent denoising may provide a stronger relation to clinical variables and contribute to the understanding of the diversity of previously reported results, as well as the interpretation of schizophrenia as a dysconnection disease in general.

6   Acknowledgments and Disclosures

The publication was supported by: ERDF-Project Brain dynamics, No. CZ.02.01.01/00/22_008/000464, Czech Health Research Council Project No. NU21-08-00432 and by the MSMT CR program Cooperatio, Neuroscience, Charles University.

This manuscript has been posted on a preprint server, BioRxiv, under doi: https://doi.org/10.1101/2024.09.20.613853.

Author contributions

FS, JH, and JHl are responsible for conceptualization; DT and JHl are responsible for methodology; DT is responsible for software; DT a JHl performed the validation; DT, JH, and JHl are responsible for the formal analysis; DT, MK, BRB, FS, JH, and JHl are responsible for investigation; JT, FS, and JHl provided the resources; DT, JT, FS, JH, and JHl are responsible for data curation; DT, MK, BRB, JH, and JHl wrote the original draft; DT, MK, BRB, JT, FS, JH, and JHl reviewed and edited the manuscript; DT is responsible for the visualization; JT, FS, JH, and JHl performed the supervision; JT, FS, and JHl administered the project; FS, JH and JHl are responsible for funding acquisition.

Financial disclosure

None reported.

Conflict of interest

The authors declare no potential conflict of interests.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

Anticevic, A., Cole, M. W., Repovs, G., Murray, J. D., Brumbaugh, M. S., Winkler, A. M., Savic, A., Krystal, J. H., Pearlson, G. D., & Glahn, D. C. (2014). Characterizing Thalamo-Cortical Disturbances in Schizophrenia and Bipolar Illness. Cerebral Cortex, 24(12), 3116–3130. https://doi.org/10.1093/cercor/bht165

Anticevic, A., Haut, K., Murray, J. D., Repovs, G., Yang, G. J., Diehl, C., McEwen, S. C., Bearden, C. E., Addington, J., Goodyear, B., Cadenhead, K. S., Mirzakhanian, H., Cornblatt, B. A., Olvet, D., Mathalon, D. H., McGlashan, T. H., Perkins, D. O., Belger, A., Seidman, L. J., … Cannon, T. D. (2015). Association of Thalamic Dysconnectivity and Conversion to Psychosis in Youth and Young Adults at Elevated Clinical Risk. JAMA Psychiatry, 72(9), 882. https://doi.org/10.1001/jamapsychiatry.2015.0566

Argyelan, M., Ikuta, T., DeRosse, P., Braga, R. J., Burdick, K. E., John, M., Kingsley, P. B., Malhotra, A. K., & Szeszko, P. R. (2014). Resting-State fMRI Connectivity Impairment in Schizophrenia and Bipolar Disorder. Schizophrenia Bulletin, 40(1), 100–110. https://doi.org/10.1093/schbul/sbt092

Backasch, B., Sommer, J., Klöhn-Saghatolislam, F., Müller, M. J., Kircher, T. T. J., & Leube, D. T. (2014). Dysconnectivity of the inferior frontal gyrus: Implications for an impaired self-other distinction in patients with schizophrenia. Psychiatry Research: Neuroimaging, 223(3), 202–209. https://doi.org/10.1016/j.pscychresns.2014.05.007

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. NeuroImage, 37(1), 90–101. https://doi.org/10.1016/j.neuroimage.2007.04.042

Benetti, S., Mechelli, A., Picchioni, M., Broome, M., Williams, S., & McGuire, P. (2009). Functional integration between the posterior hippocampus and prefrontal cortex is impaired in both first episode schizophrenia and the at risk mental state. Brain, 132(9), 2426–2436. https://doi.org/10.1093/brain/awp098

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B: Statistical Methodology, 57(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Biswal, B., Zerrin Yetkin, F., Haughton, V. M., & Hyde, J. S. (1995). Functional connectivity in the motor cortex of resting human brain using echo‑planar mri. Magnetic Resonance in Medicine, 34(4), 537–541. https://doi.org/10.1002/mrm.1910340409

Burgess, G. C., Kandala, S., Nolan, D., Laumann, T. O., Power, J. D., Adeyemo, B., Harms, M. P., Petersen, S. E., & Barch, D. M. (2016). Evaluation of Denoising Strategies to Address Motion-Correlated Artifacts in Resting-State Functional Magnetic Resonance Imaging Data from the Human Connectome Project. Brain Connectivity, 6(9), 669–680. https://doi.org/10.1089/brain.2016.0435

Caputi, L., Pidnebesna, A., & Hlinka, J. (2021). Promises and pitfalls of topological data analysis for brain connectivity analysis. NeuroImage, 238, 118245. https://doi.org/10.1016/j.neuroimage.2021.118245

Chopra, S., Francey, S. M., O'Donoghue, B., Sabaroedin, K., Arnatkeviciute, A., Cropley, V., Nelson, B., Graham, J., Baldwin, L., Tahtalian, S., Yuen, H. P., Allott, K., Alvarez-Jimenez, M., Harrigan, S., Pantelis, C., Wood, S. J., McGorry, P., & Fornito, A. (2021). Functional Connectivity in Antipsychotic-Treated and Antipsychotic-Naive Patients With First-Episode Psychosis and Low Risk of Self-harm or Aggression: A Secondary Analysis of a Randomized Clinical Trial. JAMA Psychiatry, 78(9), 994. https://doi.org/10.1001/jamapsychiatry.2021.1422

Cole, M. W., Anticevic, A., Repovs, G., & Barch, D. (2011). Variable Global Dysconnectivity and Individual Differences in Schizophrenia. Biological Psychiatry, 70(1), 43–50. https://doi.org/10.1016/j.biopsych.2011.02.010

Corlett, P. R., Horga, G., Fletcher, P. C., Alderson-Day, B., Schmack, K., & Powers, A. R. (2019). Hallucinations and Strong Priors. Trends in Cognitive Sciences, 23(2), 114–127. https://doi.org/10.1016/j.tics.2018.12.001

Crossley, N. A., Mechelli, A., Fusar‑Poli, P., Broome, M. R., Matthiasson, P., Johns, L. C., Bramon, E., Valmaggia, L., Williams, S. C. R., & McGuire, P. K. (2009). Superior temporal lobe dysfunction and frontotemporal dysconnectivity in subjects at risk of psychosis and in first‑episode psychosis. Human Brain Mapping, 30(12), 4129–4137. https://doi.org/10.1002/hbm.20834

Dauvermann, M. R., Lee, G., & Dawson, N. (2017). Glutamatergic regulation of cognition and functional brain connectivity: Insights from pharmacological, genetic and translational schizophrenia research. British Journal of Pharmacology, 174(19), 3136–3160. https://doi.org/10.1111/bph.13919

Demirtaş, M., Burt, J. B., Helmer, M., Ji, J. L., Adkinson, B. D., Glasser, M. F., Van Essen, D. C., Sotiropoulos, S. N., Anticevic, A., & Murray, J. D. (2019). Hierarchical Heterogeneity across Human Cortex Shapes Large-Scale Neural Dynamics. Neuron, 101(6), 1181-1194.e13. https://doi.org/10.1016/j.neuron.2019.01.017

Dong, D., Wang, Y., Chang, X., Luo, C., & Yao, D. (2018). Dysfunction of Large-Scale Brain Networks in Schizophrenia: A Meta-analysis of Resting-State Functional Connectivity. Schizophrenia Bulletin, 44(1), 168–181. https://doi.org/10.1093/schbul/sbx034

Du, X., Choa, F.-S., Chiappelli, J., Wisner, K. M., Wittenberg, G., Adhikari, B., Bruce, H., Rowland, L. M., Kochunov, P., & Hong, L. E. (2019). Aberrant Middle Prefrontal-Motor Cortex Connectivity Mediates Motor Inhibitory Biomarker in Schizophrenia. Biological Psychiatry, 85(1), 49–59. https://doi.org/10.1016/j.biopsych.2018.06.007

Friston, K. (2005). A theory of cortical responses. Philosophical Transactions of the Royal Society B: Biological Sciences, 360(1456), 815–836. https://doi.org/10.1098/rstb.2005.1622

Friston, K. J. (2002). Dysfunctional connectivity in schizophrenia. World Psychiatry: Official Journal of the World Psychiatric Association (WPA), 1(2), 66–71.

Friston, K. J., & Frith, C. D. (1995). Schizophrenia: A disconnection syndrome? Clinical Neuroscience (New York, N.Y.), 3(2), 89–97.

Friston, K. J., Frith, C. D., Liddle, P. F., & Frackowiak, R. S. J. (1993). Functional Connectivity: The Principal-Component Analysis of Large (PET) Data Sets. Journal of Cerebral Blood Flow & Metabolism, 13(1), 5–14. https://doi.org/10.1038/jcbfm.1993.4

Giraldo-Chica, M., & Woodward, N. D. (2017). Review of thalamocortical resting-state fMRI studies in schizophrenia. Schizophrenia Research, 180, 58–63. https://doi.org/10.1016/j.schres.2016.08.005

González-Vivas, C., Soldevila-Matías, P., Sparano, O., García-Martí, G., Martí-Bonmatí, L., Crespo-Facorro, B., Aleman, A., & Sanjuan, J. (2019). Longitudinal studies of functional magnetic resonance imaging in first-episode psychosis: A systematic review. European Psychiatry, 59, 60–69. https://doi.org/10.1016/j.eurpsy.2019.04.009

Guller, Y., Ferrarelli, F., Shackman, A. J., Sarasso, S., Peterson, M. J., Langheim, F. J., Meyerand, M. E., Tononi, G., & Postle, B. R. (2012). Probing Thalamic Integrity in Schizophrenia Using Concurrent Transcranial Magnetic Stimulation and Functional Magnetic Resonance Imaging. Archives of General Psychiatry, 69(7). https://doi.org/10.1001/archgenpsychiatry.2012.23

Hare, S. M., Ford, J. M., Mathalon, D. H., Damaraju, E., Bustillo, J., Belger, A., Lee, H. J., Mueller, B. A., Lim, K. O., Brown, G. G., Preda, A., Van Erp, T. G. M., Potkin, S. G., Calhoun, V. D., & Turner, J. A. (2019). Salience–Default Mode Functional Network Connectivity Linked to Positive and Negative Symptoms of Schizophrenia. Schizophrenia Bulletin, 45(4), 892–901. https://doi.org/10.1093/schbul/sby112

Heckers, S., Weiss, A. P., Deckersbach, T., Goff, D. C., Morecraft, R. J., & Bush, G. (2004). Anterior Cingulate Cortex Activation During Cognitive Interference in Schizophrenia. American Journal of Psychiatry, 161(4), 707–715. https://doi.org/10.1176/appi.ajp.161.4.707

Hlinka, J., Paluš, M., Vejmelka, M., Mantini, D., & Corbetta, M. (2011). Functional connectivity in resting-state fMRI: Is linear correlation sufficient? NeuroImage, 54(3), 2218–2225. https://doi.org/10.1016/j.neuroimage.2010.08.042

Hoffstaedter, F., Grefkes, C., Caspers, S., Roski, C., Palomero‑Gallagher, N., Laird, A. R., Fox, P. T., & Eickhoff, S. B. (2014). The role of anterior midcingulate cortex in cognitive motor control: Evidence from functional connectivity analyses. Human Brain Mapping, 35(6), 2741–2753. https://doi.org/10.1002/hbm.22363

Kapur, S. (2003). Psychosis as a State of Aberrant Salience: A Framework Linking Biology, Phenomenology, and Pharmacology in Schizophrenia. American Journal of Psychiatry, 160(1), 13–23. https://doi.org/10.1176/appi.ajp.160.1.13

Kaufmann, T., Skåtun, K. C., Alnæs, D., Doan, N. T., Duff, E. P., Tønnesen, S., Roussos, E., Ueland, T., Aminoff, S. R., Lagerberg, T. V., Agartz, I., Melle, I. S., Smith, S. M., Andreassen, O. A., & Westlye, L. T. (2015). Disintegration of Sensorimotor Brain Networks in Schizophrenia. Schizophrenia Bulletin, 41(6), 1326–1335. https://doi.org/10.1093/schbul/sbv060

Kay, S. R., Fiszbein, A., & Opler, L. A. (1987). The Positive and Negative Syndrome Scale (PANSS) for Schizophrenia. Schizophrenia Bulletin, 13(2), 261–276. https://doi.org/10.1093/schbul/13.2.261

Li, S., Hu, N., Zhang, W., Tao, B., Dai, J., Gong, Y., Tan, Y., Cai, D., & Lui, S. (2019). Dysconnectivity of Multiple Brain Networks in Schizophrenia: A Meta-Analysis of Resting-State Functional Connectivity. Frontiers in Psychiatry, 10, 482. https://doi.org/10.3389/fpsyt.2019.00482

Li, T., Wang, Q., Zhang, J., Rolls, E. T., Yang, W., Palaniyappan, L., Zhang, L., Cheng, W., Yao, Y., Liu, Z., Gong, X., Luo, Q., Tang, Y., Crow, T. J., Broome, M. R., Xu, K., Li, C., Wang, J., Liu, Z., … Feng, J. (2016). Brain-Wide Analysis of Functional Connectivity in First-Episode and Chronic Stages of Schizophrenia. Schizophrenia Bulletin, sbw099. https://doi.org/10.1093/schbul/sbw099

Li, X., Morgan, P. S., Ashburner, J., Smith, J., & Rorden, C. (2016). The first step for neuroimaging data analysis: DICOM to NIfTI conversion. Journal of Neuroscience Methods, 264, 47–56. https://doi.org/10.1016/j.jneumeth.2016.03.001

Melicher, T., Horacek, J., Hlinka, J., Spaniel, F., Tintera, J., Ibrahim, I., Mikolas, P., Novak, T., Mohr, P., & Hoschl, C. (2015). White matter changes in first episode psychosis and their relation to the size of sample studied: A DTI study. Schizophrenia Research, 162(1–3), 22–28. https://doi.org/10.1016/j.schres.2015.01.029

Mennigen, E., Fryer, S. L., Rashid, B., Damaraju, E., Du, Y., Loewy, R. L., Stuart, B. K., Calhoun, V. D., & Mathalon, D. H. (2019). Transient Patterns of Functional Dysconnectivity in Clinical High Risk and Early Illness Schizophrenia Individuals Compared with Healthy Controls. Brain Connectivity, 9(1), 60–76. https://doi.org/10.1089/brain.2018.0579

Meyer-Lindenberg, A. S., Olsen, R. K., Kohn, P. D., Brown, T., Egan, M. F., Weinberger, D. R., & Berman, K. F. (2005). Regionally Specific Disturbance of Dorsolateral

Prefrontal–Hippocampal Functional Connectivity in Schizophrenia. Archives of General Psychiatry, 62(4), 379. https://doi.org/10.1001/archpsyc.62.4.379

Palaniyappan, L., & Liddle, P. F. (2012). Does the salience network play a cardinal role in psychosis? An emerging hypothesis of insular dysfunction. Journal of Psychiatry and Neuroscience, 37(1), 17–27. https://doi.org/10.1503/jpn.100176

Pamilo, S., Malinen, S., Hotta, J., & Seppä, M. (2015). A correlation‑based method for extracting subject‑specific components and artifacts from group‑f MRI data. European Journal of Neuroscience, 42(9), 2726–2741. https://doi.org/10.1111/ejn.13034

Patel, A. X., Kundu, P., Rubinov, M., Jones, P. S., Vértes, P. E., Ersche, K. D., Suckling, J., & Bullmore, E. T. (2014). A wavelet method for modeling and despiking motion artifacts from resting-state fMRI time series. NeuroImage, 95, 287–304. https://doi.org/10.1016/j.neuroimage.2014.03.012

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI. NeuroImage, 84, 320–341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Raichle, M. E. (2015). The restless brain: How intrinsic activity organizes brain function. Philosophical Transactions of the Royal Society B: Biological Sciences, 370(1668), 20140172. https://doi.org/10.1098/rstb.2014.0172

Ringen, P. A., Reponen, E. J., Vedal, T. S. J., Andreassen, O. A., Steen, N. E., & Melle, I. (2019). Predictors for Antipsychotic Dosage Change in the First Year of Treatment in Schizophrenia Spectrum and Bipolar Disorders. Frontiers in Psychiatry, 10, 649. https://doi.org/10.3389/fpsyt.2019.00649

Sarpal, D. K., Robinson, D. G., Lencz, T., Argyelan, M., Ikuta, T., Karlsgodt, K., Gallego, J. A., Kane, J. M., Szeszko, P. R., & Malhotra, A. K. (2015). Antipsychotic Treatment and Functional Connectivity of the Striatum in First-Episode Schizophrenia. JAMA Psychiatry, 72(1), 5. https://doi.org/10.1001/jamapsychiatry.2014.1734

Shackman, A. J., Salomons, T. V., Slagter, H. A., Fox, A. S., Winter, J. J., & Davidson, R. J. (2011). The integration of negative affect, pain and cognitive control in the cingulate cortex. Nature Reviews Neuroscience, 12(3), 154–167. https://doi.org/10.1038/nrn2994

Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R., & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. The Journal of Clinical Psychiatry, 59 Suppl 20, 22-33;quiz 34-57.

Shirer, W. R., Jiang, H., Price, C. M., Ng, B., & Greicius, M. D. (2015). Optimization of rs-fMRI Pre-processing for Enhanced Signal-Noise Separation, Test-Retest Reliability, and Group Discrimination. NeuroImage, 117, 67–79. https://doi.org/10.1016/j.neuroimage.2015.05.015

Sigurdsson, T., & Duvarci, S. (2016). Hippocampal-Prefrontal Interactions in Cognition, Behavior and Psychiatric Disease. Frontiers in Systems Neuroscience, 9. https://doi.org/10.3389/fnsys.2015.00190

Stephan, K. E., Baldeweg, T., & Friston, K. J. (2006). Synaptic Plasticity and Dysconnection in Schizophrenia. Biological Psychiatry, 59(10), 929–939. https://doi.org/10.1016/j.biopsych.2005.10.005

Sterzer, P., Adams, R. A., Fletcher, P., Frith, C., Lawrie, S. M., Muckli, L., Petrovic, P., Uhlhaas, P., Voss, M., & Corlett, P. R. (2018). The Predictive Coding Account of Psychosis. Biological Psychiatry, 84(9), 634–643. https://doi.org/10.1016/j.biopsych.2018.05.015

Tang, Y., Zhou, Q., Chang, M., Chekroud, A., Gueorguieva, R., Jiang, X., Zhou, Y., He, G., Rowland, M., Wang, D., Fu, S., Yin, Z., Leng, H., Wei, S., Xu, K., Wang, F., Krystal, J. H., & Driesen, N. R. (2019). Altered functional connectivity and low-frequency signal fluctuations in early psychosis and genetic high risk. Schizophrenia Research, 210, 172–179. https://doi.org/10.1016/j.schres.2018.12.041

Troisi, A., Pasini, A., De Angelis, F., & Spalletta, G. (1997). Paranoid/Belligerence and Neuroleptic Dosage in Newly Admitted Schizophrenic Patients: Journal of Clinical Psychopharmacology, 17(2), 84–87. https://doi.org/10.1097/00004714-199704000-00003

Tukey, J. W. (1977). Exploratory data analysis (Repr.). Addison-Wesley.

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B., & Joliot, M. (2002). Automated Anatomical Labeling of Activations in SPM Using a Macroscopic Anatomical Parcellation of the MNI MRI Single-Subject Brain. NeuroImage, 15(1), 273–289. https://doi.org/10.1006/nimg.2001.0978

Ulmer, S., & Jansen, O. (Eds.). (2020). fMRI: Basics and Clinical Applications. Springer International Publishing. https://doi.org/10.1007/978-3-030-41874-8

Wang, H.-L. S., Rau, C.-L., Li, Y.-M., Chen, Y.-P., & Yu, R. (2015). Disrupted thalamic resting-state functional networks in schizophrenia. Frontiers in Behavioral Neuroscience, 9. https://doi.org/10.3389/fnbeh.2015.00045

Weinberger, D. R., Aloia, M. S., Goldberg, T. E., & Berman, K. F. (1994). The frontal lobes and schizophrenia. The Journal of Neuropsychiatry and Clinical Neurosciences, 6(4), 419–427. https://doi.org/10.1176/jnp.6.4.419

Wengler, K., Goldberg, A. T., Chahine, G., & Horga, G. (2020). Distinct hierarchical alterations of intrinsic neural timescales account for different manifestations of psychosis. eLife, 9, e56151. https://doi.org/10.7554/eLife.56151

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: A Functional Connectivity Toolbox for Correlated and Anticorrelated Brain Networks. Brain Connectivity, 2(3), 125–141. https://doi.org/10.1089/brain.2012.0073

Woods, S. W. (2003). Chlorpromazine Equivalent Doses for the Newer Atypical Antipsychotics. The Journal of Clinical Psychiatry, 64(6), 663–667. https://doi.org/10.4088/JCP.v64n0607

Woodward, N. D., & Heckers, S. (2016). Mapping Thalamocortical Functional Connectivity in Chronic and Early Stages of Psychotic Disorders. Biological Psychiatry, 79(12), 1016–1025. https://doi.org/10.1016/j.biopsych.2015.06.026

Woodward, N. D., Karbasforoushan, H., & Heckers, S. (2012). Thalamocortical Dysconnectivity in Schizophrenia. American Journal of Psychiatry, 169(10), 1092–1099. https://doi.org/10.1176/appi.ajp.2012.12010056