Biological oscillations without genetic oscillator or external forcing

Vincent Vandenbroucke [1,†] , Lucas Henrion [1,†] , and Frank Delvigne [1,*]

[1]Terra Research and Teaching Centre, Microbial Processes and Interactions (MiPI), Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium

[†]V.V. (Vincent Vandenbroucke) contributed equally to this work with L.H. (Lucas Henrion).

[*] Corresponding author. E-mail: F.Delvigne@uliege.be

Abstract

Oscillators are fundamental to biological systems, underpinning essential processes such as cell division, circadian rhythms, and developmental cycles. While both natural and synthetic genetic oscillators have been extensively studied, oscillatory behaviors in cells can also emerge without dedicated genetic circuits. In earlier work, we uncovered sustained oscillations in phenotypic switching across diverse cellular systems and gene circuits, occurring spontaneously, without external forcing and linked them to the induction of slow-growing phenotypes. In this study, we identify the conditions that give rise to such intrinsic phenotypic instabilities, leading to population-level oscillations. We develop and analytically solve a simplified mathematical model of a stress-induced phenotype, mapping the range of continuous culture conditions that trigger oscillatory gene expression. This instability range, predicted by the model, was experimentally validated in *Bacillus subtilis* cultures. Our findings reveal that oscillations can arise in the complete absence of genetic oscillators or external perturbations. Although demonstrated here for a stress response in continuous culture, this phenomenon may occur in any long-term cultivation where environmental feedback links an inducer to the cellular system, broadening the landscape of possible oscillatory behaviors in microbiology and synthetic biology.

Introduction

Oscillations in gene expression are traditionally attributed to dedicated genetic circuits, metabolic pathways or external forcing[1][5]. Here, we report sustained oscillations emerging in the complete absence of either, arising instead from the interplay between cellular burden and nutrient availability in a constant environment. Oscillatory behaviors are pervasive in biology, spanning multiple scales, from single cells to entire organisms, and regulating substantial portions of cellular metabolism[2][6]. Their study has revealed natural gene circuits acting as biological oscillators, such as the circadian clock in cyanobacteria[8] and gut microbiota[9][12], as well as the cell cycle in *E. coli* [14]. These discoveries have been complemented by synthetic designs[16][18], including the iconic repressillator, which generates spontaneous oscillations in gene expression through a triad of repressor proteins[19][23]. In general, biological oscillations arise from dedicated gene circuits [23][16], external periodic forcing[24], or a combination of both[25]. In all reported cases, oscillations under constant culture conditions were linked to an underlying genetic oscillator e.g., yeast respiratory cycles[3], bacterial cell cycle[7] and plankton synchronization[4].

Surprisingly, we previously observed that highly burdensome gene circuits can exhibit sustained oscillations in a simple chemostat[10], a continuous cultivation device initially designed to maintain constant growth conditions[26][27]. Strikingly, the two systems that displayed this behavior, the T7 expression system in *E. coli* BL21 and the sporulation program of *Bacillus subtilis*, lack network architectures typically associated with oscillatory dynamics. To uncover the origin of these instabilities, we focused on the *B. subtilis* system, which can be distilled into: i) Two discrete phenotypes linked to growth capacity, vegetative, non-fluorescent cells that grow, and sporulating, fluorescent cells that do not; ii) A single trigger, glucose, whose depletion both induces sporulation[11] and serves as the primary carbon source.

To uncover the origin of these unexpected oscillations, we developed a minimal mathematical model of the stress response in a chemostat, a system with constant nutrient influx, where stress is triggered when nutrient levels fall below a critical threshold. The model predicts that, even under the classically "stable" operating conditions of a chemostat, there exists a specific range of dilution rates in which no steady-state population-level stress response can be sustained. This challenges the long-standing assumption that chemostats inherently stabilize gene expression across microbial populations[26]. We experimentally tested this prediction using *Bacillus subtilis* 168 carrying a chromosomal *gfp* reporter under the control of the *spoIIE* promoter, SpoIIE being a key regulator

of asymmetric cell division during sporulation[13][15]. Cultures were grown at different dilution rates,

and sporulation activation was tracked at the single-cell level using a custom flow cytometry–

bioreactor interface[17][20]. Consistent with model predictions, we observed oscillatory sporulation

dynamics within the predicted operating range, manifesting as successive bursts of sporulating cells

sweeping through the population.

Results

A mathematical model predicts instabilities in gene expression for a specific range of dilution

rate.

To probe how oscillations can arise in the absence of a genetic oscillator, we constructed a minimal

ODE model capturing stress-induced dynamics in a cell population. The framework tracks two

phenotypic states: non-stressed cells (X_1) and stressed cells (X_2) (**Eqs. 1–5**). The stressed phenotype

grows more slowly, with a maximum growth rate $\mu_{max,2} = k\,\mu_{max,1}$ where $k \in [0, 1]$ and μ_{max} is the

maximal growth rate of the non-stressed biomass. Switching from X_1 to X_2 occurs at a constant rate

H when the substrate concentration S drops below a threshold K_I and is implemented via a step

function $H_1(S)$ (**Eq. 5**). Because stressed cells grow slowly and stress-related proteins are primarily

diluted through cell growth, the reverse transition from X_2 to X_1 is assumed negligible.

$$\frac{\mathrm{d}S}{\mathrm{d}t} = (S_{in} - S)D - \frac{\mu X_1}{Y_{X1}} - \frac{k\mu X_2}{Y_{X2}} \tag{1}$$

$$\frac{\mathrm{d}X_1}{\mathrm{d}t} = (\mu - D)X_1 - H_1(S)X_1 \tag{2}$$

$$\frac{\mathrm{d}X_2}{\mathrm{d}t} = (k\mu - D)X_2 + H_1(S)X_1 \tag{3}$$

$$\mu = \mu_{max}\frac{S}{K_S + S} \tag{4}$$

$$H_1(S) = \begin{cases} H \text{ if } S \leq K_I \\ 0 \text{ if } S > K_I \end{cases} \tag{5}$$

90 Here, S_{in} denotes the sugar concentration in the feed, D the dilution rate, Y_{Xi} the biomass yields of

91 each phenotype, and K_S the sugar concentration at half-maximum growth rate. Analytical solutions

of the model (see **SI Appendix**) reveal a critical range of dilution rates in which no steady-state equilibrium can be achieved among the state variables, rendering the system inherently unstable (**Fig. 1A**).

Within this instability window, two mutually exclusive conditions prevent convergence to equilibrium. First, the classical chemostat formulation, omitting the stress phenotype, fails because the residual substrate concentration S stabilizes below the threshold K_I, triggering the stress response. Second, the alternative scenario of continuous switching into the stressed state is also invalid, as the associated reduction in substrate uptake drives S above K_I, thereby suppressing the stress response.

Rather than resolving into either case, the system hovers near the switching threshold K_I, forcing cells to continually adjust to environmental fluctuations. This creates a negative feedback loop on population growth which, in combination with delayed phenotypic transitions, produces sustained oscillations analogous to those of the synthetic repressilator[23][19][28] (**Fig. 1B,C**). The appearance and persistence of these oscillations are expected to depend on both the delay magnitude and the fidelity of the step function in capturing the switching dynamics, while the dilution-rate range for their occurrence can be experimentally validated.

A. Solution space for S with respect to D and unstable region

B. Simulation of X and S over time in the unstable region

C. The cell population motif is analogous to a repressilator

Figure 1: Representation of the solution space of the stress-response model at steady state. **(A)** Model predictions reveal three distinct regimes. At high dilution rates, only the non-stressed phenotype is sustained, as the sugar concentration remains above the threshold K_I. At low dilution rates, the stressed phenotype dominates, with sugar concentrations falling below K_I. Between these extremes lies an instability window where no steady state exists, and oscillations can emerge. **(B)** Introducing a delay in the stress response generates sustained oscillations *in silico* within the unstable region. The example shown corresponds to *B. subtilis* sporulation dynamics with $D = 0.1$ h⁻¹ and a 0.5 h delay, although small parameter changes can markedly alter the oscillatory behavior. **(C)** Schematic analogy between the sporulation system in *B. subtilis* (right) and the synthetic repressilator (7) (left). Here, the stressed and non-stressed subpopulations, together with the extracellular sugar concentration, form a triple negative feedback loop. In the chemostat, oscillations arise at the level of reactor concentrations rather than intracellular protein levels: sporulation (or stress induction) reduces the number of unstressed cells, decreasing sugar uptake, raising sugar concentration above K_I, inhibiting sporulation, and allowing unstressed cells to expand again.

The instability region can be experimentally observed in chemostat for a given range of dilution rates

Model predictions were tested under continuous cultivation, a regime designed to maintain stable growth conditions through substrate (S) limitation, which can be tuned via the dilution rate (D). As a case study, we examined *Bacillus subtilis* sporulation, a stress response triggered when residual glucose concentration falls below a threshold K_I. Here, X_1 denotes vegetative (non-stressed) biomass and X_2 denotes stressed biomass (spores). Sporulation was monitored at the single-cell level using automated flow cytometry and a P$_{spoIIE}$*::gfp* transcriptional reporter, where *spoIIE* encodes a key regulator of asymmetric division during sporulation.

Using H, K_I, μ_{max}, and K_S values determined from batch cultures (**SI Appendix**), and assuming no growth for the stressed phenotype, the model predicts that at high D, with residual glucose above K_I, only X_1 persists. At low D, with residual glucose at or below K_I, both X_1 and X_2 are present, with X_2 (spores) periodically increasing in abundance. Due to the absence of growth in X_2 and a high switching rate from X_1, the model anticipates no regime of stable spore concentration.

Experimental observations aligned with these predictions: at high dilution rates, residual glucose remained above the switching threshold and sporulation was not activated, yielding exclusively X_1 biomass (**Fig. 2**). Reducing D caused residual glucose to fall below K_I, initiating bursts of sporulation from X_1 to X_2. These oscillatory transitions persisted for at least five residence times before gradually decreasing in amplitude.

A. Automated Flow Cytometry setup

2. Auto-Sampler

1. Continuous culture

3. Flow Cytometer

Side Scatter
Forward scatter
Fluorescence

B. Dilution rate sweep reveals instability in gene expression

D=0.32   D=0.13   D=0.10 h⁻¹

C. Confirmation of the instability

D.

Figure 2: **Mapping of cell population dynamics using automated flow cytometry reveals spontaneous oscillations in the**

activation of the *spoIIE* sysem during continuous cultivation of *B. subtilis*.

(A) Schematic of the automated flow cytometry setup. Samples are withdrawn directly from the bioreactor and analyzed at single-

cell   resolution   every   15   min.

(B) Time–fluorescence density plots, where brighter colors indicate higher cell counts. The black line marks the 75th percentile of

fluorescence; large oscillations are indicated by asterisks. In agreement with model predictions, gene expression dynamics vary with

the dilution rate (D): at $D = 0.32$ h⁻¹, cells remain vegetative, whereas lowering D to 0.13 h⁻¹ triggers sporulation bursts detected by

activation of the $P_{spoIIE}::gfp$ reporter. These bursts diminish in amplitude upon further reduction to $D = 0.10$ h⁻¹.

(C) Prolonged chemostat cultivation at $D = 0.11$ h⁻¹ confirmed the persistence of oscillatory sporulation.

(D) Differential fluorescence maps showing the increase (red) or decrease (blue) in cell counts per fluorescence bin between

successive time points, smoothed for clarity. Until ~40 h of continuous culture, clear waves of sporulation are observed. Color units

are arbitrary.

Discussion

Oscillatory behaviors in biology have long been associated with dedicated genetic oscillators, external forcing, or a combination of both[1,29,30]. From natural systems such as circadian clocks and cell cycles to synthetic designs like the repressilator[19], the prevailing assumption has been that sustained oscillations require an explicit genetic architecture to generate them. Our work challenges this paradigm by demonstrating that oscillations can arise spontaneously in constant environments, purely from the interplay between cellular physiology and environmental feedback.

Our results demonstrate that even gene networks lacking intrinsic oscillatory architecture can generate sustained oscillations in the absence of external forcing. These oscillations emerge from the interplay of two independent but complementary mechanisms. The first is the impact of gene expression on cellular growth. This phenomenon is widespread: many natural circuits, such as sporulation in *B. subtilis*, and synthetic systems, such as burdensome protein overproduction[21], can impose substantial growth penalties[22]. Similar effects are conceivable for genes affecting persistence traits, such as biofilm formation, where the "cost" to growth is offset by improved survival in specific environments[31].

The second requirement is environmental feedback, in which cells not only sense but also modify the concentration of their inducers. This feedback occurs when cells consume, degrade, or produce the relevant inducer, thereby reshaping their own stimulus landscape. In our case study, sporulation reduced glucose uptake, allowing residual glucose to rise above the sporulation threshold and thereby suppressing further induction in non-stressed cells. A similar mechanism was reported for a T7 expression system under lactose control[31], where growth inhibition from burdensome protein production increased residual glucose levels and suppressed induction via carbon catabolite repression.

Fundamentally, the oscillations observed here represent system-level circuits: emergent behaviors arising from population–environment interactions rather than from the regulatory topology of an isolated genetic network. Whereas genetic oscillators are defined by well-mapped gene–protein interactions[16,25], system-level oscillators depend on how those interactions couple to resource flows and environmental feedback[32,31,33]. The example we present is relatively simple, but more complex interactions, especially where multiple gene networks intersect through shared environmental variables, could produce rich and unexpected dynamical patterns.

188 The emergence of oscillations in the absence of an explicit genetic oscillator has several important

189 implications. First, it challenges the assumption, fundamental to chemostat theory[26 34], that constant

190 environmental conditions necessarily yield steady-state population structures and gene expression

191 profiles. Second, it suggests that similar instabilities may occur in other long-term cultures

192 whenever environmental feedback is tightly coupled to a phenotypic switch that significantly alters

193 resource consumption. This principle could extend to diverse microbial systems, from industrial

194 bioprocesses[35] to natural microbial communities[36 37], where environmental cues and physiological

195 states are dynamically linked.

Materials and Methods

Mathematical Model Solution

The dynamic model describing growth and stress-induced phenotypic switching was solved analytically, as detailed in the Supplementary Information (**SI Appendix**, "Stress model in continuous culture"). To this end, we first considered separately the regimes with or without switching toward the stressed phenotype and, subsequently, merged both solutions to obtain a complete analytical description. The approach was then generalized to exponential fed-batch conditions. Additional analyses regarding the model's domain of validity and biological assumptions are also provided in the SI.

The analytical solutions revealed that, for specific combinations of biological parameters (growth rates, switching kinetics) and operational conditions (dilution rate), steady-states do not exist, indicating transitions to oscillatory dynamics or population wash-out.

Model predictions were verified through numerical simulations of a continuous culture, performed in MATLAB using the delay differential equation solver *dde23*. Parameter values were selected based on experimental measurements obtained with our *Bacillus subtilis* strain, including a 30-min delay between the environmental stress threshold (substrate limitation) and activation or repression of the switch to the stressed phenotype.

Experimental Design

The objective of this study was to elucidate how regular gene-expression oscillations can emerge in the absence of dedicated oscillator motifs or engineered external feedback. We used *Bacillus subtilis* sporulation as a model system for stress-induced phenotypic transitions. To parameterize the model, we first experimentally quantified growth and switching characteristics, as described in the "Growth and switching parameter determination" section of the SI. Using these parameters and the analytical framework described above, we identified dilution-rate regimes predicted to either suppress sporulation or induce sustained oscillations. Chemostat experiments were then performed at representative dilution rates within these regimes. Sporulation dynamics were monitored using

our custom automated flow-cytometry setup, in combination with an engineered reporter strain producing GFP upon activation of the sporulation program. This setup enabled single-cell-resolved tracking of phenotypic transitions and direct validation of model predictions.

Strain and Growth media

All experiments were performed in minimal mineral media containing (in g/l): K_2HPO_4: 14.6; $NaH_2PO_4 \cdot 2H_2O$: 3.6; Na_2SO_4: 2; $(NH_4)_2SO_4$: 2.47; NH_4Cl: 0.5; $(NH_4)2H$-citrate: 1; glucose: 5, thiamine: 0.01, tryptophan: 0.05. The medium is supplemented with a trace element solution totaling 11 ml/l assembled from the following solutions (in g/l), 3/11 of $FeCl_3 \cdot 6H_2O$: 16.7, 3/11 of EDTA: 20.1, 2/11 of $MgSO_4$: 120 and 3/11 of a metallic trace element solution. The metallic trace element solution contains (in g/L): $CaCl_2 \cdot 2H_2O$: 0.74; $ZnSO_4 \cdot 7H_2O$: 0.18; $MnSO_4 \cdot H_2O$: 0.1; $CuSO_4 \cdot 5H_2O$: 0.1, $CoSO_4 \cdot 7H_2O$: 0.21. Both the trace element solution and the amino acids were filter sterilized (0.22 μm). To monitor the switching to the sporulating phenotype, *Bacillus subtilis* 168 was modified to express *gfpmut2* under the control of the SpoIIE promoter. The strain, *B. subtilis* 168 $P_{spoIIE:gfpmut2}$, was built by AmyE chromosomal integration of a cassette containing the promoter, gfpmut2 and the selection maker, kanamycin.

Cultivation Conditions

Precultures were started from single colonies picked from a LB plate and grown overnight. To avoid starting bioreactor cultivations with sporulating populations, the precultures were done in overfilled shake flasks (20 % of filled volume) without baffles to reach oxygen limitation before glucose. All bioreactor cultivations were performed at 37°C, pH 7 and aeration at 1 VVM. For growth and switching parameters determination, experiments were performed in DASBox mini - bioreactors (Eppendorf) with a stirring speed of 400 rpm and a cultivation volume of 160 ml. Continuous cultivations were performed in Bionet F1 bioreactor with a cultivation volume of 1L at an agitation speed of 1000 min^{-1}. The transition from batch to continuous was triggered once dissolved oxygen rose, signaling the transition from exponential to stationary phase.

Bioreactor cultivations were monitored with a custom-made sampling device (the Segregostat) that automatically draws a sample from the reactor and dilutes it with PBS before flow cytometry (FC) analysis (BD Accuri, C6 for the parameter determination and C6+ for the continuous cultures). A

total of 40,000 cells was analyzed in each sample where the FL1-A channel is used to visualize the GFP content.

Statistical Analysis

The switching and growth parameters were determined in triplicate cultivations. The cultivations to test the impact of dilution rates were performed in duplicates and the dilution rate leading to instability was confirmed one last time. During all cultivations, samples were automatically taken every 15 minutes and 40,000 cells were analyzed.

References

1. Tyson, J.J., Albert, R., Goldbeter, A., Ruoff, P., and Sible, J. (2008). Biological switches and clocks. J R Soc Interface *5 Suppl 1*, S1-8. https://doi.org/10.1098/rsif.2008.0179.focus.

2. Li, Z., and Yang, Q. (2018). Systems and synthetic biology approaches in understanding biological oscillators. Quant Biol *6*, 1–14. https://doi.org/10.1007/s40484-017-0120-7.

3. Henson, M.A. (2002). Cell population modelling of yeast glycolytic oscillations. Biochemical journal *368*, 433–446.

4. Massie, T.M., Blasius, B., Weithoff, G., Gaedke, U., and Fussmann, G.F. (2010). Cycles, phase synchronization, and entrainment in single-species phytoplankton populations. Proc Natl Acad Sci U S A *107*, 4236–4241. https://doi.org/10.1073/pnas.0908725107.

5. Goldbeter, A., and Yan, J. (2022). Multi-synchronization and other patterns of multi-rhythmicity in oscillatory biological systems. Interface Focus *12*, 20210089. https://doi.org/10.1098/rsfs.2021.0089.

6. Takhaveev, V., Özsezen, S., Smith, E.N., Zylstra, A., Chaillet, M.L., Chen, H., Papagiannakis, A., Milias-Argeitis, A., and Heinemann, M. (2023). Temporal segregation of biosynthetic processes is responsible for metabolic oscillations during the budding yeast cell cycle. Nat Metab *5*, 294–313. https://doi.org/10.1038/s42255-023-00741-x.

7. Tanouchi, Y., Pai, A., Park, H., Huang, S., Stamatov, R., Buchler, N.E., and You, L. (2015). A noisy linear map underlies oscillations in cell size and gene expression in bacteria. Nature *523*, 357–360. https://doi.org/10.1038/nature14562.

8. Cohen, S.E., and Golden, S.S. (2015). Circadian Rhythms in Cyanobacteria. Microbiol Mol Biol Rev *79*, 373–385. https://doi.org/10.1128/MMBR.00036-15.

9. Voigt, R.M., Forsyth, C.B., Green, S.J., Engen, P.A., and Keshavarzian, A. (2016). Circadian Rhythm and the Gut Microbiome. Int Rev Neurobiol *131*, 193–205. https://doi.org/10.1016/bs.irn.2016.07.002.

10. Henrion, L., Martinez, J.A., Vandenbroucke, V., Delvenne, M., Telek, S., Zicler, A., Grünberger, A., and Delvigne, F. (2023). Fitness cost associated with cell phenotypic switching drives population diversification dynamics and controllability. Nat Commun *14*, 6128. https://doi.org/10.1038/s41467-023-41917-z.

11. Dawes, I.W., and Mandelstam, J. (1970). Sporulation of Bacillus subtilis in continuous culture. J Bacteriol *103*, 529–535. https://doi.org/10.1128/jb.103.3.529-535.1970.

12. Bishehsari, F., Voigt, R.M., and Keshavarzian, A. (2020). Circadian rhythms and the gut microbiota: from the metabolic syndrome to cancer. Nat Rev Endocrinol *16*, 731–739. https://doi.org/10.1038/s41574-020-00427-4.

13. Wolf, D.M., Fontaine-Bodin, L., Bischofs, I., Price, G., Keasling, J., and Arkin, A.P. (2008). Memory in microbes: quantifying history-dependent behavior in a bacterium. PLoS One *3*, e1700. https://doi.org/10.1371/journal.pone.0001700.

14. Iuliani, I., Mbemba, G., Lagomarsino, M.C., and Sclavi, B. (2024). Direct single-cell observation of a key Escherichia coli cell-cycle oscillator. Sci Adv *10*, eado5398. https://doi.org/10.1126/sciadv.ado5398.

15. Fujita, M., and Losick, R. (2003). The master regulator for entry into sporulation in Bacillus subtilis becomes a cell-specific transcription factor after asymmetric division. Genes Dev *17*, 1166–1174. https://doi.org/10.1101/gad.1078303.

16. Stricker, J., Cookson, S., Bennett, M.R., Mather, W.H., Tsimring, L.S., and Hasty, J. (2008). A fast, robust and tunable synthetic gene oscillator. Nature *456*, 516–519. https://doi.org/10.1038/nature07389.

17. Sassi, H., Nguyen, T.M., Telek, S., Gosset, G., Grünberger, A., and Delvigne, F. (2019). Segregostat: a novel concept to control phenotypic diversification dynamics on the example of Gram-negative bacteria. Microbial Biotechnology *12*, 1064–1075. https://doi.org/10.1111/1751-7915.13442.

18. Din, M.O., Danino, T., Prindle, A., Skalak, M., Selimkhanov, J., Allen, K., Julio, E., Atolia, E., Tsimring, L.S., Bhatia, S.N., et al. (2016). Synchronized cycles of bacterial lysis for in vivo delivery. Nature *536*, 81–85. https://doi.org/10.1038/nature18930.

19. Elowitz, M.B., and Leibler, S. (2000). A synthetic oscillatory network of transcriptional regulators. Nature *403*, 335–338. https://doi.org/10.1038/35002125.

20. Delvigne, F., and Martinez, J.A. (2023). Advances in automated and reactive flow cytometry for synthetic biotechnology. Curr Opin Biotechnol *83*, 102974. https://doi.org/10.1016/j.copbio.2023.102974.

21. Borkowski, O., Ceroni, F., Stan, G.-B., and Ellis, T. (2016). Overloaded and stressed: whole-cell considerations for bacterial synthetic biology. Curr Opin Microbiol *33*, 123–130. https://doi.org/10.1016/j.mib.2016.07.009.

22. Klumpp, S., and Hwa, T. (2014). Bacterial growth: global effects on gene expression, growth feedback and proteome partition. Curr Opin Biotechnol *28*, 96–102. https://doi.org/10.1016/j.copbio.2014.01.001.

23.   Potvin-Trottier, L., Lord, N.D., Vinnicombe, G., and Paulsson, J. (2016). Synchronous long-term oscillations in a synthetic gene circuit. Nature *538*, 514–517. https://doi.org/10.1038/nature19841.

24.   Lugagne, J.-B., Sosa Carrillo, S., Kirch, M., Kohler, A., Batt, G., and Hersen, P. (2017). Balancing a genetic toggle switch by real-time feedback control and periodic forcing. Nature communications *8*, 1671. https://doi.org/10.1038/s41467-017-01498-0.

25.   Mondragón-Palomino, O., Danino, T., Selimkhanov, J., Tsimring, L., and Hasty, J. (2011). Entrainment of a population of synthetic genetic oscillators. Science *333*, 1315–1319. https://doi.org/10.1126/science.1205369.

26.   NOVICK, A., and SZILARD, L. (1950). Description of the chemostat. Science *112*, 715–716. https://doi.org/10.1126/science.112.2920.715.

27.   Wides, A., and Milo, R. Understanding the Dynamics and Optimizing the Performance of Chemostat Selection Experiments. arXiv:1806.00272.

28.   Liao, M.J., Din, M.O., Tsimring, L., and Hasty, J. (2019). Rock-paper-scissors: Engineered population dynamics increase genetic stability. Science *365*, 1045–1049. https://doi.org/10.1126/science.aaw0542.

29.   Mather, W., Bennett, M.R., Hasty, J., and Tsimring, L.S. (2009). Delay-induced degrade-and-fire oscillations in small genetic circuits. Phys Rev Lett *102*, 068105. https://doi.org/10.1103/PhysRevLett.102.068105.

30.   Novák, B., and Tyson, J.J. (2008). Design principles of biochemical oscillators. Nat Rev Mol Cell Biol *9*, 981–991. https://doi.org/10.1038/nrm2530.

31.   Tan, C., Marguet, P., and You, L. (2009). Emergent bistability by a growth-modulating positive feedback circuit. Nat Chem Biol *5*, 842–848. https://doi.org/10.1038/nchembio.218.

32.   Liao, C., Blanchard, A.E., and Lu, T. (2017). An integrative circuit-host modelling framework for predicting synthetic gene network behaviours. Nat Microbiol *2*, 1658–1666. https://doi.org/10.1038/s41564-017-0022-5.

33.   Piho, P., and Thomas, P. (2024). Feedback between stochastic gene networks and population dynamics enables cellular decision-making. Sci Adv *10*, eadl4895. https://doi.org/10.1126/sciadv.adl4895.

34.   Wong, B.G., Mancuso, C.P., Kiriakov, S., Bashor, C.J., and Khalil, A.S. (2018). Precise, automated control of conditions for high-throughput growth of yeast and bacteria with eVOLVER. Nat Biotechnol *36*, 614–623. https://doi.org/10.1038/nbt.4151.

35.   Enfors, S.O., Jahic, M., Rozkov, A., Xu, B., Hecker, M., Jurgen, B., Kruger, E., Schweder, T., Hamer, G., O'Beirne, D., et al. (2001). Physiological responses to mixing in large scale bioreactors. J Biotechnol *85*, 175–185.

36.   Rodríguez-Verdugo, A., Vulin, C., and Ackermann, M. (2019). The rate of environmental fluctuations shapes ecological dynamics in a two-species microbial system. Ecol Lett *22*, 838–846. https://doi.org/10.1111/ele.13241.

37.   Mancuso, C.P., Lee, H., Abreu, C.I., Gore, J., and Khalil, A.S. (2021). Environmental fluctuations reshape an unexpected diversity-disturbance relationship in a microbial community. Elife *10*. https://doi.org/10.7554/eLife.67175.

Acknowledgments

We thank Laurie Josselin, Juan Andres Martinez, Mélanie Grégoire and Mathéo Delvenne for commenting the article.

Funding: LH and VV are supported by a FRIA grant provided by the "Fonds de la Recherche Scientifique" FRS–FNRS from the Walloon region of Belgium.

Author contributions: V.V. contributed equally to this work with L.H. V.V. and L.H. developed the model, performed the experiments and drafted the manuscript. F.D. secured related fundings, wrote and reviewed the manuscript.

Competing interests: All other authors declare they have no competing interests.

Data and materials availability: The automated flow cytometry data and associated code have been deposited on https://zenodo.org/records/13803914). The data analysis code relies on the previously published toolbox available at https://gitlab.uliege.be/mipi/published-software/mbiomas-core.