Unravelling the Proteomic Landscape of Imipenem Resistance in *Pseudomonas*
***aeruginosa*: A Comparative Investigation between Clinical and Control Strains**

Marius Eidsaa[1], Animesh Sharma[3,4], Markus Janasch[1], Alicja Kuch[2], Anna Skoczyńska[2],
Francesca Di Bartolomeo[1*].

[1]Department of Biotechnology and Nanomedicine, SINTEF Industry, Trondheim 7034, Norway
[2]Department of Epidemiology and Clinical Microbiology, National Medicines Institute (NMI), Warsaw,
Poland
[3] Department of Clinical and Molecular Medicine, Norwegian University of Science and Technology (NTNU),
N-7489 Trondheim, Norway
[4] Proteomics and Modomics Experimental Core (PROMEC) at NTNU and the Central Norway Regional
Health Authority, Trondheim, Norway
12
13 *** Correspondence:**
14 Corresponding Author
15 francesca.dibartolomeo@sintef.no

16 **Keywords:** Antimicrobial Resistance ₁, *Pseudomonas aeruginosa* ₂, Label-free Quantitative
17 Proteomics ₃, antibiotic challenge ₄, imipenem ₅**.**

Highlights:
1: **Stable resistance mechanisms**: The clinical strain of *Pseudomonas aeruginosa* (ST235)
exhibited a stable proteomic profile across varying concentrations of imipenem, indicating pre-
established and efficient resistance mechanisms.
2: **Reactive proteomic response**: The control strain (*P. aeruginosa* ATCC 27853) showed a
broader and more reactive proteomic response, particularly in proteins related to membrane
transport, stress response, and biofilm formation.
3: **Uncharacterized proteins**: Significant upregulation of uncharacterized proteins in the
clinical strain suggests potential novel resistance mechanisms that warrant further investigation.
27

Abstract/Summary

The increasing prevalence of antimicrobial resistance (AMR) poses a significant challenge to global
health, particularly with bacterial pathogens such as *Pseudomonas aeruginosa*, a notorious cause of
nosocomial infections. This study focuses on the comparative proteomic analysis of an imipenem-
resistant strain of *P. aeruginosa*, a representative of world epidemic clone ST235, and a wildtype
control strain, *P. aeruginosa* ATCC 27853, in response to varying concentrations of imipenem. Using
label-free quantification (LFQ) and gene ontology (GO) enrichment analyses, we identified significant
differences in the proteomic responses between the two strains (data available via ProteomeXchange
with identifier PXD055744.). The clinical strain exhibited a stable proteomic profile across the
imipenem gradient, suggesting pre-established and efficient resistance mechanisms that do not require
extensive reconfiguration under antibiotic pressure. In contrast, the control strain showed a broader,
more reactive proteomic response, particularly in proteins associated with membrane transport, stress
response, and biofilm formation. Notably, uncharacterized proteins were significantly upregulated in

the clinical strain, indicating potential novel resistance mechanisms. These findings highlight the distinct strategies employed by the two strains, with the clinical strain's stable resistance mechanisms contrasting sharply with the control strain's reactive approach. The study underscores the importance of further research into the uncharacterized proteins that may play crucial roles in antibiotic resistance, potentially leading to new therapeutic targets in the fight against AMR.

1   Introduction

Figure 1 – *Graphical abstract. Setup and workflow used in the comparative proteomic analysis of Pseudomonas aeruginosa strains under varying concentrations of the antibiotic imipenem. The study involves a P. aeruginosa clinical strain (ST235) and a control strain (P. aeruginosa ATCC 27853), both subjected to increasing concentrations of imipenem based on their respective minimum inhibitory concentrations (MICs) with each condition replicated three times. Imipenem concentrations range from 1.6-16 mg/L and 0.2-2.0 mg/L for the clinical and control strains respectively, corresponding to 0.1x MIC, 0.25x MIC, 0.5x MIC and 1.0x MIC. The untreated control probes (no antibiotic) for each strain are indicated by* Control* *in the figure. Imipenem, a beta-lactam antibiotic, disrupts bacterial cell wall formation by binding to specific proteins in the bacterial cell membrane. The proteomic pipeline includes sample processing, mass spectrometry for label-free quantification, and data mining, leading to insights into differential protein expression and resistance mechanisms.* Part of the figure has been created with BioRender.com (Eidsaa, M. (2024) BioRender.com/w02t614).

Antimicrobial resistance (AMR) has emerged as a global health crisis, compromising the efficacy of traditional therapies against a diverse array of infections caused by bacteria, parasites, and fungi (Darby *et al.*, 2023). This intricate biological phenomenon is primarily fueled by the selective pressure imposed by the widespread and often indiscriminate utilization of antimicrobials, which fosters an environment conducive to the survival and proliferation of resistant strains. The consequent outcome is an escalating prevalence of hard-to-treat infections (Fair and Tor, 2014).

Particularly alarming is the resistance demonstrated by bacterial pathogens. These organisms harness mechanisms such as horizontal gene transfer, mutation and selection, and adaptive resistance, to rapidly

evolve and withstand antimicrobial agents (Davies and Davies, 2010). The rise of multi-drug resistant (MDR) and extensively drug-resistant (XDR) bacterial strains adds another layer of complexity to this scenario. These "superbugs" demonstrate resistance against multiple classes of antibiotics, posing serious therapeutic challenges and undermining patient outcomes (Magiorakos *et al.*, 2012; Aslam *et al.*, 2018).

Pseudomonas aeruginosa is an exemplar of such a resilient bacterial species. A notorious instigator of nosocomial infections worldwide, it demonstrates an exceptional ability to resist antibiotic therapies (Breidenstein, de la Fuente-Núñez and Hancock, 2011; Langendonk, Neill and Fothergill, 2021). Given the robust resistance of such strains, treatment often resorts to last-line antibiotics, such as those in the carbapenem class. Imipenem, a member of this class, has a broader antimicrobial spectrum and higher potency than other beta-lactam antibiotics. It operates by interfering with bacterial cell wall synthesis, thus exerting a bactericidal effect. This antibiotic is often seen as a therapeutic beacon against a variety of MDR bacterial infections (Lautenbach *et al.*, 2006; Xu *et al.*, 2020). However, the growing overreliance on imipenem has inadvertently catalyzed the emergence of imipenem-resistant *P. aeruginosa* strains, escalating the challenges associated with this formidable pathogen. A prime example of its adaptability is the clinical strain P. aeruginosa, recognized as a representative of world epidemic clone of sequence type ST235, owing to its extensive prevalence and heightened resistance to multiple antibiotics.

To effectively counter this rising tide of resistance, we must delve deeper into the molecular mechanisms underpinning the resistance of bacteria to potent antibiotics such as imipenem.

Proteomic analysis, in this regard, can prove instrumental by elucidating antibiotic resistance mechanisms at the protein level, as proteins are the ultimate executors of most cellular functions and interactions (Pérez-Llarena and Bou, 2016). Proteomics can offer invaluable insights into the adaptive responses of bacteria to antibiotic challenges. It achieves this by quantifying changes in protein expression, modifications, and interactions under different conditions. Proteome-wide label-free quantification, a technique frequently employed in proteomic analysis, allows for the comparison of protein abundance between resistant and susceptible strains, or between bacteria subjected to different antibiotic concentrations (Goodyear *et al.*, 2023)**.** As for this study, comparative proteomic studies serve as a potent investigative tool for understanding the profound and complex responses of bacteria to antibiotic exposure. They have an unprecedented ability to illuminate the molecular landscape where the dynamic interplay between antibiotics and bacteria unfolds. In the face of antibiotic stress, bacteria respond by expressing an array of proteins that enable them to neutralize the antibiotic's effect, remove the antibiotic from the bacterial cell, or modify the bacterial cell targets to evade the antibiotic's impact (Khodadadi *et al.*, 2020). Proteomic studies, through their ability to quantitatively analyze the entire complement of proteins expressed under different conditions, enable us to delineate this intricate molecular proteomics pool in detail. The proteomic response of bacteria to antibiotics, which includes the alterations in protein expression, interactions, and modifications, is part of a larger adaptive response of bacteria to survive under antibiotic stress. This response forms the "resistome" of bacteria, a term coined to describe the collection of all the antibiotic resistance genes in a bacterial cell or community (Wright, 2007). While the concept of the resistome is predominantly genomic in nature, the functional output of the resistome is embodied in the proteome. It is the proteins, being the

functional units of cells, that execute the instructions coded in the resistome, thereby manifesting the
resistance phenotype (Sulaiman and Lam, 2022).
In the present study, we utilize the power of proteomics to dissect the resistome in action, as we analyze
how two strains of *P. aeruginosa* - the representative of the world epidemic clone ST235 and the
control strain ATCC 27853 - respond to the antibiotic imipenem, scrutinizing these profiles across a
spectrum of escalating minimum inhibitory concentrations (MICs) of the antibiotic. This nuanced
approach will enable us to trace the dynamic shifts in protein expression and interactions as antibiotic
pressure intensifies. Additionally, comparing the two selected strains will yield valuable insights into
the proteomic signatures tied to the clinical strain's heightened resistance. Therefore, this research is
primed to enrich our understanding of *P. aeruginosa*'s proteomic responses and the resistance
mechanisms to imipenem. These insights may lead us towards potential solutions to the urgent and
ongoing crisis of antimicrobial resistance by identifying proteins or pathways that could serve as novel
targets for drug development or interventions that could disrupt the functionality of the resistome.
123
124 2   Materials and Methods
125 2.1   Strains

Pseudomonas aeruginosa 6378/09 strain (NMI collection number), hereinafter referred to as clinical
strain and *Pseudomonas aeruginosa* ATCC 27853 control (reference) strain, obtained from American
Type Culture Collection, were used for proteomic analysis. It is wildtype, imipenem-sensitive strain,
hereinafter referred to as control strain. The *P. aeruginosa* clinical strain was isolated from an oncology
patient from an abdominal swab. It is imipenem-resistant strain, associated with VIM-2 variant of
metallo-beta-lactamase (MBL) and belonged to world epidemic clone ST235, as previously described
(Urbanowicz *et al.*, 2021). ST235 has been the most frequent MDR *P. aeruginosa* clone globally,
associated with VIM/IMP-like MBLs (Oliver *et al.*, 2015).

134 2.2   Determination of the minimum inhibitory concentration (MIC)

Minimum inhibitory concentration (MIC) of imipenem was determined for both strains by the broth
microdilution method according to standard ISO 20776-1 (ISO 20776-1, 2020). The MIC was defined
to be the lowest concentration at which no visible growth (complete inhibition of growth) of bacteria
could be observed after incubation for 18±2 hours as recommended by the European Committee on
Antimicrobial Susceptibility Testing (EUCAST)(European Committee on Antimicrobial Susceptibility
Testing, 2024).

141 2.3   Cell culture and antibiotic challenge

The bacteria were grown in cation-adjusted Mueller-Hinton broth (Becton Dickinson, Sparks, MD,
USA) at 37 °C with shaking (120 rpm) to optical density OD_{600} = 1.0. For antibiotic challenge,
imipenem concentration of 0.1x MIC, 0.25x MIC, 0.5x MIC and 1x MIC of the particular strain was
used in a volume of 5 ml. Bacterial cultures without the antibiotic were considered as untreated
controls, **see Table 1**. Challenged bacterial cultures were then incubated with shaking (120 rpm) for 5
hours at 35 °C ± 2 °C. Each experiment was performed three times for both clinical and control strains
(3 biological replicates) with two technical replicates for each condition (antibiotic concentration).

After incubation the bacterial cell pellets were harvested in pre-cooled Falcon tubes by centrifugation
(5 min at 3,000 × g below 0 °C). Then, the cell pellets were washed twice with ice-cold 1 x PBS
(Sigma-Aldrich, Life Science, St. Louis, MO, USA). Finally, cell pellets were collected by
centrifugation (5 min at 3,000 × g below 0 °C), snap-frozen, and stored at −80 °C until further testing
(Liu *et al.*, 2016; Hashemi *et al.*, 2019; Jongers *et al.*, 2021).
154
155 **Table 1: Clinical strain (A) and control strain (B)**

A	*P. aeruginosa* clinical strain (ST235)	Control* **A1**	0.1x MIC **A2**	0.25x MIC **A3**	0.5x MIC **A4**	1.0x MIC **A5**
B	*P. aeruginosa* control strain (ATCC 27853)	Control* **B1**	0.1x MIC **B2**	0.25x MIC **B3**	0.5x MIC **B4**	1.0x MIC **B5**

156 *Note: All samples are in biological triplicates*
157 ** Bacterial cultures without the antibiotic (untreated controls)*
158

2.4   Label-Free Proteomics analyses

Proteins were quantified by processing the timsTOF-pro DDA MS-data using MaxQuant v2.0.3.0
(Tyanova, Temu and Cox, 2016). Namely, the following search parameters were used: enzyme
specified as trypsin with a maximum of two missed cleavages allowed; minimum peptide length of 7;
acetylation of protein N-terminal, oxidation of methionine, and deamidation of asparagine/glutamine
as dynamic post-translational modification. These were imported in MaxQuant which uses m/z and
retention time (RT) values to align each run against each other sample with a minute window match-
between-run function and 20 mins overall sliding window using a clustering-based technique. These
were further queried against the reference-proteome of *Pseudomonas aeruginosa* downloaded from
UniProt (UniProt Consortium, 2022) in May 2022 and MaxQuant's internal contaminants database
using Andromeda built into MaxQuant. Both Protein and peptide identifications false discovery rate
(FDR) was set to 1%, only unique peptides with high confidence were used for final protein group
identification. Peak abundances were extracted by integrating the area under the peak curve. Each
protein group abundance was normalized by the total abundance of all identified peptides for each run
and protein by calculated median summing all unique peptide-ion abundances for each protein using
label-free quantification (LFQ) algorithm (Cox *et al.*, 2014) with minimum unique peptide(s) ≥ 1. The
mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the
PRIDE (Perez-Riverol *et al.*, 2022) partner repository with the dataset identifier PXD055744
(Username: reviewer_pxd055744@ebi.ac.uk Password: XNtCwVbI4AIs).

2.5   Data processing and mining

Having obtained the label-free quantification (LFQ) intensity values for both the *Pseudomonas*
aeruginosa clinical strain and control strain, across the five different concentrations (as depicted in
Figure 1), we performed a series of data filtering, processing, and aggregation steps, including removal
of reverse hits and potential contaminants to ensure the integrity of our data. In the case of multiple
UniProt protein IDs per hits, the one with the highest curation status were used in the downstream
analysis when relevant. This led to a total of *N*=3811 proteins in our analysis, accounting for

approximately 60% of the chromosomal genes in an average *P. aeruginosa* strain (Mosquera-Rendón
et al., 2016).
For each strain and antibiotic concentration, we had three biological replicates (**Figure 1**). We
performed a standard *t*-test using the log2-transformed LFQ (log2LFQ) values for each protein ID to
obtain the associated *p*-values. These *p*-values were then corrected using the Benjamini-Hochberg
False Discovery Rate (FDR) procedure. This was done for all experiments with non-zero MIC, using
the 0.0x MIC as a control for each strain. This means that we compared LFQ values and performed *t*-
tests within strains only. Alongside the *t*-test, *p*-value and the FDR values, we used the median value
for LFQ intensities where applicable.
When comparing LFQ intensity values between experiments, we primarily used fold change relative
to the control probe (no antibiotic) to describe the difference in log2 space, expressed via log2LFQ
values. Proteins with LFQ intensity values of zero for both the control probe (no antibiotic) and 1.0x
MIC experiments were excluded. Proteins with values of zero in one experiment and non-zero in
another were assigned a large, finite fold-change value. When comparing across strains, we used the
median log2LFQ fold changes between the strains, as these values were already normalized with
respect to the 0.0x MIC control experiment for the respective strains.
Unless otherwise specified, the intra-strain protein LFQ intensity fold-change cutoff was set at 1.5, or
approximately +/- 0.5849 after log2 transformation. In the individual protein volcano plots an
uncorrected *p*-value cutoff of 0.1 was employed, with values above colored grey and values below
colored light red or blue, while FDR values below 0.1 are furthermore indicated by dark red or blue
color. These thresholds are not particularly conservative, which is deemed appropriate given the
explorative nature of the study. The individual protein volcano plot x-axis and y-axis ranges were fixed
to improve visual comparability, and log2 fold changes were visually capped at +/- 5 alongside *p*-
values below 1e-5.
To visualize the overall features of the data set, we calculated and used a Principal Component Analysis
(PCA), using standard scaling, and a correlation matrix (heatmap) of Spearman's correlation
coefficients between experiments. This provided an overview of the structure and relationships in the
data set (**Figure 2**).

2.6   Gene-ontology enrichment analysis

To discern significant systemic alterations in the proteomes within and across strains and antibiotic
concentrations, we conducted a Gene Ontology (GO) term enrichment analysis. This method is set-
based, meaning that groups of proteins are compared rather than individual proteins. Statistical tests
are performed by comparing a subset to the full background set. The GO term enrichment analysis
aims at finding enriched functional, process-related, and compartmental annotations associated with
each protein in the subset. Collectively, numerous individually insignificant protein-level changes can
be linked to statistically significant features on aggregate, cellular level.
The GO term enrichment analysis is agnostic to the subset selection method. Therefore, we employ
both FDR-value, uncorrected *p*-values, and no *p*-value thresholds when selecting proteins, in
conjunction with fold-change thresholds, at various parts of the analysis. We performed statistical tests
based on hypergeometric distributions on the subset selection, and subsequently corrected these using
the Benjamini-Hochberg FDR procedure, setting the FDR-value threshold to 0.1. Both over-

representation and under-representation were accounted for in the analyses. The choice of background,
i.e., the full set against which the subset is compared, is crucial for obtaining accurate results. For this,
we used all mapped and processed proteins with well-defined *t*-tests based on the three biological
replicates.
We collected GO term information for *Pseudomonas aeruginosa* using UniProt, release: 2024_03
(UniProt Consortium, 2023), and mapped the GO term IDs onto the Gene Ontology Knowledgebase
using AmiGO 2, version: 2.5.17 (Carbon and Mungall, 2024), aligning the information with the
UniProt IDs. Out of 3811 proteins, 3253 were associated with GO terms, and the remaining were
assigned a placeholder value to ensure their inclusion in the analyses, since the non-annotated proteins
may be of particular interest for further study.
In the results, we visualized the enriched GO terms using volcano plots. The x-coordinate of each GO
term represents the log2 value of its enrichment (blue indicates under-representation and red indicates
over-representation), and the range is fixed and visually capped at +/- 2.5. The y-coordinate represents
the FDR values associated with each GO term (not to be confused with the protein-level FDR-values).
The area of the GO-term circles is proportional to the number of proteins associated with the GO term
in the subset. All terms with a GO-term FDR value > 0.1 are indicated in grey and are excluded from
further analysis.

2.7   Resources and tools

Data analyses and visualizations were performed using Python 3.11, along with standard modules such
as numpy, pandas, scipy, matplotlib, and seaborn, among others. The *t*-tests and dose-correlation
statistics were conducted using R.

3   Results

Here, we present the results of our comparative proteomic investigation between the *Pseudomonas*
aeruginosa world epidemic clinical strain and the wildtype control strain under four increasing
concentrations (in mg/L): 1.6, 4, 8, 16 and 0.2, 0.5, 1.0, 2.0, which equates to 0.1x, 0.25x, 0.5x, and
1.0x MIC for the clinical and control strains respectively. The proteome data, acquired and processed
as described in the Materials and Methods section, comprised a total of *N*=3811 proteins. For each
protein, we calculated the LFQ intensity values across the five concentrations for both strains.
Subsequently, a *t*-test was performed across the biological replicates, using the control probe (no
antibiotic) experiments as controls, to derive protein-specific *p*-values.
256

257
258
Figure 2: *Visualization of the label-free quantification (LFQ) proteomics data sets, illustrating the impact of*
increasing imipenem concentration on the clinical (A1-5) and control (B1-5) strains. (a) Standard-scaled PCA
plot shown for the two most significant principal components, accounting for 49.8% and 17.7% of the total
variance. The imipenem-resistant clinical strain forms a compact cluster on the left-hand side, while the control
strain exhibits a spread on the right-hand side. This spread shows a clear gradient from low antibiotic
concentrations (lighter colors) at the top to high antibiotic concentrations (darker colors) at the bottom. (b)
Heatmap of Spearman's correlation coefficients indicating a similar trend as observed in the PCA plot. It
indicates a high overall correlation between the LFQ intensity scores, and particularly high correlations within
the clinical strain.

3.1 Results overview

Exploration of the overall trends in the data, as visualized in **Figure 2**, suggests that the protein
abundance in the clinical strain remains stable under the influence of imipenem, unlike the control
strain. The PCA plot (**Figure 2a**) reveals that the largest variance between the experiments is attributed
to the strains themselves (PC1, x-axis). The second largest factor of variance (PC2, y-axis) corresponds
to the increasing imipenem concentration, as evidenced by the gradual transition from control probe
(no antibiotic) in the top-right to 1.0x MIC in the bottom-right for the control strain. The heatmap
(**Figure 2b**) displays a highly intercorrelated block in the top-left corner, representing the clinical strain
experiments. The bottom-right corner shows a slight but clear gradient towards covariance among
experiments with similar imipenem concentrations. Notably, there is high similarity between the
strains, with correlation coefficients around 0.85 for the control probes (A1 and B1) and 0.80 at 1.0x
MIC (A5 and B5).

3.2 Proteomic changes within strains

Based on the overall results from **Figure 2**, we expect the significant fold changes for individual
proteins across MIC concentrations to be more pronounced for the control strain than the clinical strain.
The volcano plots in **Figure 3** confirms this prospect. Here, we have plotted the *N*=3811 proteins with

a fold change threshold of 1.5 for both the control (**Figure 3a**) and clinical (**Figure 3b**) strains at 1.0x
MIC, as compared with a *t*-test to 0.0x MIC (see Materials and Methods for details).
In these plots, an increase in fold change from 0.0x MIC to 1.0x MIC, i.e. up-regulation, is depicted
by red dots, while a decrease in fold change, i.e. down-regulation, is represented by blue dots. Each
dot corresponds to a protein. For both strains, we display the proteins with *p*-values above 0.1, and fold
change below the threshold, as grey, while light red and blue denote *p*-values below 0.1 and full
(darker) red and blue depict the proteins with FDR corrected *p*-values below 0.1. As expected, the
control strain shows a range of significant FDR values, while the clinical strain only has 8 proteins
fulfilling the criteria: AmpC (P24735), PA0868 (Q9I573), DDAH (Q9I4E3), Tse4 (Q9I069), PA2547
(Q9I0T7), PA0207 (Q9I6T1), VreR (G3XCU5) and PhzB2 (Q9S508). These proteins (depicted in the
top corners of **Figure 3b**) are further explored in **Figure 4** and **Table 2** and detailed results will be
presented in the following subsection.

296
Figure 3: Volcano plots illustrating the proteomic changes in Pseudomonas aeruginosa for 1.0x MIC relative
to control probe (no imipenem) for (a) the control strain (ATCC 27853) and (b) the clinical strain (ST235) with
p-values along the y-axis and log2 fold change along the x-axis. Proteins with a fold change < 1.5 in either
direction, or a p-value > 0.1 are depicted in grey. Light colors represent proteins with p-values < 0.1, but with
FDR corrected p-values > 0.1, while full colors represent FDR corrected p-values < 0.1. Significant upregulated
proteins are shown in red, while relevant downregulated proteins are shown in blue. Specific proteins of interest
*are indicated by their UniProt ID and constitute the union of the proteins in **Table 2** and **Table 6**.*

304
For the normalized 1.0x MIC control strain experiments, displayed in **Figure 3a**, there are *n*=601
proteins with FDR value below 0.1 which show a fold change greater than 1.5 (depicted in dark red
and blue), and *n*=1219 proteins adhere to the fold change threshold alone. The respective numbers for
the clinical strain are *n*=8 (where *n*=79 have an uncorrected *p*-value below 0.1) and *n*=358, as visualized
in **Figure 3b**.

3.3 Comparison of specific proteins

As mentioned in the previous section, 8 proteins passed the FDR value threshold of 0.1 for the clinical
strain, making them resistome candidates which are significantly involved in the antimicrobial
resistance in *Pseudomonas aeruginosa*. These proteins are shown in **Figure 4** and **Table 2**, where

Figure 4a-d show an overall down-regulation of the specific protein in the clinical strain (in green) for
increasing MIC, **Figure 4e-f** show an overall up-regulation for increasing MIC, while **Figure 4g-h**
show a particularly large difference between the strains, i.e., where the LFQ intensity in the clinical
strain is much lower than the control strain. The control strain (in orange) shows an overall trend
towards up-regulation for these 8 proteins, aside from the proteins in **Figure 4c** and **4d**. The standard
deviations, indicated by shaded green for the clinical strain and shaded orange for the control, vary.
Large standard deviations, such as for PA0868 (Q9I573) in **Figure 4d**, indicate high variability in
protein expression, while the high-LFQ intensity protein AmpC (P24735) in **Figure 4f** displays a clear
signal, suggesting consistent protein expression across replicates. The dose-correlation statistics were
calculated for all proteins for both strains, but since the number of measurements per protein is low,
the associated *p*-values are poor, potentially making the correlation values misleading. Consequently,
the specific dose correlations values will not be discussed here, but please refer to Supplementary
Tables S1 and S2 for details on clinical and control strains respectively.

327
Figure 4: Median LFQ intensity scores plotted against increasing fractions of MIC, ranging from no antibiotic
(zero concentration) to 1.0x MIC, for 8 proteins (a-h) passing the FDR corrected t-test for the Pseudomonas
aeruginosa clinical strain, as displayed in Figure 3b. The clinical strain is shown in green, the control strain is
shown in orange, and the shaded area depicts the associated standard deviations (n=3) among the replicates
with a linear interpolation between the measurements.

The proteins in **Figure 4** are described in more detail in **Table 2**, where a description and associated
gene ontologies are added (see Materials and Methods for details). These findings and their
implications for understanding antimicrobial resistance in *Pseudomonas aeruginosa* will be discussed
further in the Discussion section.

Table 2: Significant proteins in the clinical strain with associated GO terms

Fig. 4	UniProt ID	Gene	Protein description	Gene Ontology (GO)
(a)	Q9I069	tse4	Toxin protein Tse4	extracellular region [GO:0005576]; host cell membrane [GO:0033644]; membrane [GO:0016020]
(b)	Q9I4E3	DDAH	N(G),N(G)-dimethylarginine dimethylaminohydrolase	amino acid binding [GO:0016597]; dimethylargininase activity [GO:0016403]; metal ion binding [GO:0046872]; arginine metabolic process [GO:0006525]; citrulline metabolic process [GO:0000052]; positive regulation of nitric oxide biosynthetic process [GO:0045429]
(c)	Q9I0T7	PA2547	Probable transcriptional regulator	DNA-binding transcription factor activity [GO:0003700]; sequence-specific DNA binding [GO:0043565]; DNA-templated transcription [GO:0006351]
(d)	Q9I573	PA0868	RF_PROK_I domain-containing protein	aminoacyl-tRNA hydrolase activity [GO:0004045]; ribosome binding [GO:0043022]; translation release factor activity [GO:0003747]; rescue of stalled ribosome [GO:0072344]
(e)	G3XCU5	vreR	Sigma factor regulator, VreR	sigma factor antagonist activity [GO:0016989]
(f)	P24735	ampC	Beta-lactamase	outer membrane-bounded periplasmic space [GO:0030288]; beta-lactamase activity [GO:0008800]; antibiotic catabolic process [GO:0017001]; response to antibiotic [GO:0046677]
(g)	Q9S508	phzB2	Phenazine biosynthesis protein PhzB2	phenazine biosynthetic process [GO:0002047]
(h)	Q9I6T1	PA0207	Probable transcriptional regulator	DNA-binding transcription factor activity [GO:0003700]; sequence-specific DNA binding [GO:0043565]; DNA-templated transcription [GO:0006351]

3.4 GO term enrichment within strains

To better understand the biological implications of the significant up- and downregulated protein subsets (highlighted in red and blue in **Figure 3**), we conducted a GO term enrichment analysis as outlined in the Materials and Methods section. This analysis allows us to assess the biological relevance of our protein subsets, indicated by the enrichment of biological traits, when compared to the original full set of N=3811 proteins. By comparing high-fold-change proteins between 1.0x MIC and control probe (no imipenem), we can identify the biological functions and processes that the organism upregulates or downregulates. This comparison also provides p-values, which can furthermore be FDR corrected, indicating the statistical significance of these enrichments. I.e., there are two levels of (corrected) p-values at this stage: (i) protein-level p-values, based on t-tests for individual proteins, as depicted in **Figure 3**, and (ii) protein-subset level p-values, where enrichments of protein annotations in the subset compared to the full, original set are assessed via hypergeometric processes. Thus, a random selection of proteins, which does not reflect real biology, is expected to meet the protein-subset level, GO-term FDR threshold of 0.1, thus reducing the overall likelihood of false positive GO-term

enrichment scores at 10%. In the following, we present results based on the union of upregulated (red) and downregulated (blue) proteins, as depicted in **Figure 3**. GO term analysis was performed on up- and downregulated proteins separately, but these results were too similar to the union set, when combined, to warrant a separate discussion.

Figure 5: *Comparative GO term enrichment in Pseudomonas aeruginosa strains. The volcano plots depict the enrichment of GO terms in protein subsets from the control strain (a, c) and the clinical strain (b, d), with: (a) a protein-level FDR threshold of 0.1, (b) a protein-level p-value threshold of 0.1, and (c, d) no p-value threshold. All subsets adhere to a protein-level fold-change threshold of 1.5. The x-axis represents the log2 enrichment score, while the y-axis shows the transformed FDR. GO terms are represented by circles, where the size corresponds to the number of proteins linked to each GO term, visualizing the difference of significant proteins associated with the control strain (left) compared to the clinical strain (right). The circles are color-coded as light red for over-represented (enriched) and light blue for under-represented ("under-enriched") terms. **Table 3** and **Table 4** provide additional information on the terms labeled in capital letters for (a) and (d) respectively.*

The GO term enrichment analyses were conducted for both the control and clinical strains of *Pseudomonas aeruginosa*, as depicted in **Figure 5**. The protein subsets in **Figure 5a-d** were all subject to a fold change threshold of 1.5 per protein. Furthermore, **Figure 5a** has a protein-level FDR-value threshold of 0.1 applied to it, and **Figure 5b** an uncorrected protein-level *p*-value threshold of 0.1. The

reason for this choice is based on numbers, since the clinical strain only had 8 proteins adhering to the
FDR threshold, which is deemed too small for a sensible GO term analysis for comparative purposes.
Figures 5c-d has no additional *p*-value thresholds and is only subject to a fold-change threshold of 1.5.
The protein subsets in **Figures 5a** and **5b** are thus based on the *n*=601 and *n*=79 proteins, corresponding
to the dark red and dark blue dots in **Figure 3a**, and the light and dark red and blue dots in **Figure 3b**
respectively. In **Figures 5c**, and **5d**, on the other hand, the protein subsets comprise *n*=1219 and *n*=358
proteins respectively, equating to all the dots outside the dashed vertical fold-change threshold lines.
The significantly enriched GO terms, with hypergeometric subset enrichment FDR values < 0.1, are
indicated by circles color-coded as red for over-represented terms compared to what you would expect
from random processes (enrichment above 1) and blue for under-represented ones (enrichment below
1). The size of the circles indicates the number of proteins associated with each GO term, and the terms
are alphabetically labeled based on their significance (decreasing FDR values).
385
386   **Table 3: GO terms enriched in the control strain for protein *t*-test FDR < 0.1 and fold change > 1.5 (label**
387   **from Figure 5a)**

Label	GO ID	Enrichment	Size	FDR	GO term
🔴 A	GO:0042597	3.158	38	1.39e-09	periplasmic space
🔵 B	GO:0005829	0.431	35	3.57e-08	cytosol
🔴 C	GO:0030288	3.143	23	2.12e-05	outer membrane-bounded periplasmic space
🔵 D	GO:0003700	0.296	10	6.55e-05	DNA-binding transcription factor activity
🔵 E	GO:0003677	0.306	9	3.37e-04	DNA binding
🔵 F	GO:0005737	0.557	41	3.37e-04	cytoplasm
🔴 G	GO:1904680	5.381	7	3.82e-03	peptide transmembrane transporter activity
🔴 H	GO:0002049	3.758	11	3.82e-03	pyoverdine biosynthetic process
🔴 I	GO:0043190	2.54	19	6.25e-03	ATP-binding cassette (ABC) transporter complex
🔵 J	GO:0006355	0.441	15	7.81e-03	regulation of DNA-templated transcription
🔴 K	GO:0038023	3.417	10	0.0189	signaling receptor activity
🔴 L	GO:0015562	3.075	11	0.0258	efflux transmembrane transporter activity
🔴 M	GO:0006826	3.913	7	0.0483	iron ion transport
🔴 N	GO:0042938	5.125	5	0.0483	dipeptide transport
🔴 O	GO:0015419	6.15	4	0.0518	ABC-type sulfate transporter activity
🔴 P	GO:0005886	1.325	100	0.0568	plasma membrane
🔴 Q	GO:0015891	4.1	6	0.0594	siderophore transport
🔴 R	GO:0005615	3.28	8	0.0594	extracellular space
🔴 S	GO:0016020	1.394	68	0.0856	membrane
🔴 T	GO:0033103	3.69	6	0.0982	protein secretion by the type VI secretion system
🔴 U	GO:1990281	3.311	7	0.0982	efflux pump complex

388
Table 3 provides a list of the significantly enriched GO terms shown in **Figure 5a** with their respective
statistics. Noteworthy terms include proteins associated with various cellular components (e.g.,
periplasmic space, cytosol, cytoplasm, and membrane), transporter activities (peptide transmembrane,
efflux transmembrane, ABC-type sulfate and siderophore), DNA interactions, and specific processes

(e.g., pyoverdine biosynthesis and iron ion transport). The clinical strain where only the fold-change
threshold is applied, displayed in **Figure 5d** and **Table 4**, exhibits some of the same terms as in **Table**
3, albeit with lesser significance. A unique feature here is the enrichment of proteins without associated
GO term annotations, which, with an enrichment score of 1.52, is still significant due to the substantial
number of proteins in the "NOGO" pseudo-term as GO annotations are missing.

Table 4: GO terms enriched in the clinical strain with fold change > 1.5 (label from Figure 5d)

Label	GO ID	Enrichment	Size	FDR	GO term
● A	GO:0005737	0.386	17	1.51e-04	cytoplasm
● B	GO:0005524	0.336	12	1.56e-04	ATP binding
● C	GO:0042597	3.131	21	4.65e-04	periplasmic space
● D	GO:0005829	0.493	24	1.73e-03	cytosol
● E	NOGO	1.52	77	9.55e-03	No known GO terms

The corresponding results for **Figures 5b** and **5c** are presented in Supplementary Tables S3 and S4
respectively. The apparent overlap between these results and those listed in Tables 4 and 5 indicates
that our approach is robust and reliable, even when using different or no *p*-value thresholds.

3.5 Differential protein expression and GO term enrichment

To understand the proteomic landscape of imipenem resistance in *Pseudomonas aeruginosa*, we have
thus far delved into within-strain comparisons, specifically between the 1.0x MIC and control probe
(no antibiotic) experiments. We have also cross-compared these results between the clinical and control
strains. Now, we consider a more direct comparison across strains, by using the difference between the
normalized, log2-transformed LFQ values of the control and clinical strains as a measure of fold-
change (see Materials and Methods for details).
The volcano plots depicted in **Figures 6a** and **6c** illustrate fold-change thresholds of 1.5 and 8
respectively (equivalent to approximately +/- 0.585 and exactly +/- 3 in log2 space respectively). The
reason for these choices stems from **Figure 3**, where we wanted a comparable example (**Figure 6a**) as
well as a suitably sparse counter example, where only proteins with large differences in expression are
included (**Figure 6c**). Other thresholds could be chosen, but we found a >8-fold change threshold to
be a good compromise between sparsity and representativeness. The FDR corrected *p*-values displayed
on the y-axis are derived from the control strain *t*-test. These are included for visual comparability with
Figure 3 and are not considered in the GO term enrichment analysis displayed in **Figure 6b** and **6c.**

Figure 6: Direct comparison of protein expression between the control-probe-normalized control strain and
clinical strain for 1.0x MIC imipenem treatment. (a) and (c) are volcano plots displaying the normalized
differential protein intensity fold-change along the x-axis, with a fold-change threshold of 1.5 and 8,
respectively. Red dots have larger positive fold change in the control strain, from 1.0x MIC relative to control
probe (no antibiotic), compared to the clinical strain, and, correspondingly, larger negative fold change for the
blue dots. (b) and (d) are GO term volcano plots for the protein subset based on the non-grey proteins in (a)
and (b) respectively. The FDR values along the y-axis in (a) and (c) are derived from the control strain and
applied for visual comparison.

As **Figure 2** demonstrates, the clinical strain's overall protein expression is not significantly affected
by increasing MIC. Consequently, we anticipate that the normalized differential protein expression
will mirror the results observed in the control strain. This expectation is confirmed in **Figure 6b**, which
bears both visual and content similarities to **Figure 5c** (detailed information can be found in
Supplementary Table S5). A noteworthy change, however, is the enrichment of proteins associated
with magnesium ion binding.
A GO term enrichment analysis was conducted using the 146 proteins with fold change of more than
8, denoted by blue and red dots in **Figure 6c**. The results of this analysis are presented in **Figure 6c**

and further detailed in **Table 5**. The proteins involved are provided in Supplementary Table S6 (in the "IDs" column).

Table 5: GO terms enriched in differential protein expression (label from Figure 6d)

Label	GO ID	Enrichment	Size	FDR	GO term
● A	GO:0042597	6.158	18	4.02e-08	periplasmic space
● B	GO:0005829	0.152	3	1.92e-04	cytosol
● C	GO:0031460	10.126	4	0.0155	glycine betaine transport
● D	GO:0016989	10.126	4	0.0155	sigma factor antagonist activity
● E	GO:0043190	4.403	8	0.0155	ATP-binding cassette (ABC) transporter complex
● F	GO:0042121	8.438	5	0.0155	alginic acid biosynthetic process
● G	GO:0071475	12.658	3	0.0263	cellular hyperosmotic salinity response
● H	GO:0015838	12.658	3	0.0263	amino-acid betaine transport
● I	GO:0015226	12.658	3	0.0263	carnitine transmembrane transporter activity
● J	GO:1902495	25.315	2	0.0295	transmembrane transporter complex
● K	GO:0010438	25.315	2	0.0295	cellular response to sulfur starvation
● L	GO:0005576	4.467	6	0.0319	extracellular region
● M	NOGO	1.622	33	0.0405	No known GO terms
● N	GO:0030254	6.329	4	0.0429	protein secretion by the type III secretion system
● O	GO:0051131	16.877	2	0.0574	chaperone-mediated protein complex assembly
● P	GO:0015418	16.877	2	0.0574	ABC-type quaternary ammonium compound transporting activity
● Q	GO:0004130	12.658	2	0.0986	cytochrome-c peroxidase activity
● R	GO:0033644	12.658	2	0.0986	host cell membrane

Several of these terms are recognizable from protein subsets seen in e.g. **Table 3**, but there are also differences from earlier protein subsets. These include transport terms like glycine betaine, amino-acid betaine, and carnitine transmembrane transport. There are specific enzyme-related terms such as sigma factor antagonist and cytochrome-c peroxidase, along with chaperone-mediated protein complex assembly. The list also includes cellular responses to hyperosmotic salinity and sulfur starvation, protein secretion by the type III secretion system, alginic acid biosynthesis, and structures like the extracellular region and host cell membrane. It is also important to highlight the n=33 proteins not associated with any known GO terms, representing a significant over-representation. These proteins, along with other specific proteins, will be the focus of our investigation in the subsequent section.

3.6 Comparison of selected proteins with large differential protein expression between strains

In **Figure 6**, we visualized the differential protein expression comparing the difference in the clinical strain to that of the control strain, and in **Figure 6c**, we applied a threshold demanding this difference to be >8-fold. Here, we will delve into some of the proteins adhering to this threshold. To obtain a manageable number of proteins, we employed some criteria for selection: (i) low or negative correlation values (< 0.4) between clinical and control strain across the control probe (no antibiotic)

plus 4 imipenem concentrations (0.1x, 0.25x, 0.5x and 1.0x MIC), (ii) striking differences in expression, and/or (iii) part of significantly enriched GO terms in **Table 5**. This process led us to 26 proteins, 24 of which are shown in **Figure 7**. The remaining two proteins are Tse4 (Q9I069), already displayed in **Figure 4a**, and PpiC2 (Q9HWK5) which is highly expressed in the clinical strain, and comparably non-expressed in the control across all concentrations. The remaining 24 proteins are split so that **Figure 7a-o** has an overall downregulation for the control for increasing imipenem concentrations, while **Figure 7p-x** show an overall up-regulation for the control. The clinical strain displays more stability across varying concentrations, which is not surprising given the results shown in **Figure 2**. However, **Figures 7s** and **7w**, alongside **Figure 4a**, displaying the proteins Wzm (Q9HTB8), PA0101 (Q9I731) and Tse4 (Q9I069) respectively, also show clear down-regulation in the clinical strain for increasing concentrations, in addition to the up-regulation in the control strain. As in **Figure 4**, the standard deviations are shown as green (clinical) and orange (control) shading, where linear interpolation has been used between the measurements. While several proteins display large standard deviations for the clinical strain, most control strain trends are associated with comparably small standard deviations. This indicates a clear biological effect of increasing concentrations of imipenem for the *Pseudomonas aeruginosa* (ATCC 27853) control strain which can be related to these proteins and their associated functions and compartments. More information about the proteins in **Figure 7** can be found in **Table 6**, including their UniProt description (UniProt Consortium, 2023) and associated GO terms (Carbon and Mungall, 2024). Some proteins, PA2384 (Q9I195), PA4129 (Q9HWQ2), PA1469 (Q9HVV0), PA1034 (Q9I4U0), PA1404 (Q9I3U0) and PA0101 (Q9I731), in **Figure 7c**, **d**, **i**, **m**, **p** and **w** respectively, are uncharacterized and have thus no associated GO terms. These results will be further analyzed in the Discussion section, where we will shed light on their potential association with imipenem and antimicrobial activity.

Figure 7: Median LFQ intensity scores plotted against increasing fractions of the MIC, ranging from no
imipenem (zero concentration) to 1.0x MIC, for 24 proteins (a-x) with particularly large and/or interesting
differential protein expressions between the Pseudomonas aeruginosa clinical and control strains, as displayed
*by red and blue dots in **Figure 6c**. The clinical strain is shown in green, the control strain is shown in orange,*
and the shaded area depicts the associated standard deviations (n=3) among the replicates with a linear
interpolation between the measurements.

Table 6: Selected proteins with >8-fold difference between strains at 1.0x MIC

Fig. 7	UniProt ID	Gene	Protein description	Gene Ontology (GO)
(a)	Q9HVE1	PA4651	Probable pili assembly chaperone	outer membrane-bounded periplasmic space [GO:0030288]; cell wall organization [GO:0071555]
(b)	Q9HVS4	TRPBP	Tripeptide-binding protein	ATP-binding cassette (ABC) transporter complex [GO:0043190]; outer membrane-bounded periplasmic space [GO:0030288]; peptide transmembrane transporter activity [GO:1904680]; dipeptide transport [GO:0042938]; protein transport [GO:0015031]
(c)	Q9I195	PA2384	Uncharacterized protein	
(d)	Q9HWQ2	PA4129	Uncharacterized protein	
(e)	Q9I3T4	PA1410	Probable periplasmic spermidine/putrescine-binding protein	periplasmic space [GO:0042597]; polyamine binding [GO:0019808]; polyamine transport [GO:0015846]
(f)	Q9I7B9	PA0007	Cytochrome c domain-containing protein	cytochrome-c peroxidase activity [GO:0004130]; electron transfer activity [GO:0009055]; heme binding [GO:0020037]; metal ion binding [GO:0046872]
(g)	P95434	pscF	Type III needle protein PscF	cell surface [GO:0009986]; extracellular region [GO:0005576]; type III protein secretion system complex [GO:0030257]; protein secretion by the type III secretion system [GO:0030254]
(h)	Q9HYP6	PA3350	Flagella basal body P-ring formation protein FlgA	periplasmic space [GO:0042597]; bacterial-type flagellum assembly [GO:0044780]
(i)	Q9HVV0	PA4469	Uncharacterized protein	
(j)	Q03027	aprF	Alkaline protease secretion protein AprF	cell outer membrane [GO:0009279]; efflux pump complex [GO:1990281]; efflux transmembrane transporter activity [GO:0015562]; porin activity [GO:0015288]; secretion [GO:0046903]
(k)	Q9I609	nirN	Dihydro-heme d1 dehydrogenase	periplasmic space [GO:0042597]; electron transfer activity [GO:0009055]; heme binding [GO:0020037]; metal ion binding [GO:0046872]; oxidoreductase activity [GO:0016491]; heme biosynthetic process [GO:0006783]
(l)	P14532	ccpA	Cytochrome c551 peroxidase	periplasmic space [GO:0042597]; cytochrome-c peroxidase activity [GO:0004130]; electron transfer activity [GO:0009055]; heme binding [GO:0020037]; metal ion binding [GO:0046872]
(m)	Q9I4U0	PA1034	Uncharacterized protein	
(n)	Q9HWG5	pchG	Pyochelin biosynthetic protein PchG	nucleotide binding [GO:0000166]

(o)	G3XCY8	PA4514	Probable outer membrane receptor for iron transport	cell outer membrane [GO:0009279]; pore complex [GO:0046930]; porin activity [GO:0015288]; siderophore uptake transmembrane transporter activity [GO:0015344]; signaling receptor activity [GO:0038023]; iron ion transport [GO:0006826]; siderophore transport [GO:0015891]
(p)	Q9I3U0	PA1404	Uncharacterized protein	
(q)	Q9HXQ3	PA3741	HotDog ACOT-type domain-containing protein	cytoplasm [GO:0005737]; cytosol [GO:0005829]; long-chain fatty acyl-CoA binding [GO:0036042]; long-chain fatty acyl-CoA hydrolase activity [GO:0052816]; acyl-CoA metabolic process [GO:0006637]; fatty acid catabolic process [GO:0009062]
(r)	Q9HXC3	PA3890	Probable permease of ABC transporter	ATP-binding cassette (ABC) transporter complex [GO:0043190]; transmembrane transporter complex [GO:1902495]; carnitine transmembrane transporter activity [GO:0015226]; amino-acid betaine transport [GO:0015838]; cellular hyperosmotic salinity response [GO:0071475]; glycine betaine transport [GO:0031460]
(s)	Q9HTB8	wzm	Transport permease protein	ATP-binding cassette (ABC) transporter complex [GO:0043190]; inner membrane pellicle complex [GO:0070258]; ABC-type transporter activity [GO:0140359]; lipopolysaccharide transport [GO:0015920]; O antigen biosynthetic process [GO:0009243]; transmembrane transport [GO:0055085]
(t)	Q9I744	tssF1	Type VI secretion system component TssF1	
(u)	Q9I0U4	PA2540	Bifunctional alpha/beta hydrolase/class I SAM-dependent methyltransferase	membrane [GO:0016020]; lipase activity [GO:0016298]
(v)	P07874	algA	Alginate biosynthesis protein AlgA	GTP binding [GO:0005525]; mannose-1-phosphate guanylyltransferase (GTP) activity [GO:0004475]; mannose-6-phosphate isomerase activity [GO:0004476]; alginic acid biosynthetic process [GO:0042121]; GDP-mannose biosynthetic process [GO:0009298]
(w)	Q9I731	PA0101	Uncharacterized protein	
(x)	Q9HY69	alg44	Mannuronan synthase	Gram-negative-bacterium-type cell wall [GO:0009276]; periplasmic space [GO:0042597]; alginate synthase activity [GO:0047643]; cyclic-di-GMP binding [GO:0035438]; long-chain fatty acid-CoA ligase activity [GO:0004467]; alginic acid biosynthetic process [GO:0042121]; single-species biofilm formation [GO:0044010]

4   Discussion

The proteomic landscape of *Pseudomonas aeruginosa* under the influence of imipenem, a commonly used carbapenem antibiotic, has been the subject of investigation in this study revealing key insights into the molecular mechanisms of antimicrobial resistance. Our comparative analysis between the world epidemic clinical strain and the control strain has highlighted changes in the proteome that occur

in response to varying imipenem concentrations. Utilizing normalized label-free quantification (LFQ)
intensities, protein group level *t*-tests, and Gene Ontology (GO) enrichment analysis of protein subsets,
we have been able to delve deeper into the molecular mechanisms of imipenem resistance.

The observed differences in minimum inhibitory concentrations (MICs) between the clinical and
control strains (**Table 1; Figure 2**) underscore the clinical strain's enhanced resistance to imipenem,
reflecting its ability to survive under antibiotic pressures that typically inhibit or kill susceptible strains
(Leigue, Montiani-Ferreira and Moore, 2016). The varying antibiotic concentrations provide a suitable
framework to explore the adaptive proteomic responses of both strains, revealing molecular adaptations
associated with resistance.

In the clinical strain, eight proteins are identified as significant, mentioned in **Table 2**, each playing a
role in the resistance strategy and with wide-reaching implications for understanding the mechanisms
of imipenem resistance in *Pseudomonas aeruginosa*. The upregulation of beta-lactamase (AmpC,
P24735) is consistent with its well-documented role in hydrolyzing beta-lactam antibiotics, including
imipenem, and it is known to belong in a group of enzymes that provide multi-resistance to antibiotics
(Jacoby, 2009). Its overexpression in the clinical strain could be a direct response to the antibiotic
pressure, leading to the degradation of imipenem and thus, resistance (Berrazeg *et al.*, 2015; Barbier *et*
al., 2023; Elfadadny *et al.*, 2024). In addition, also the sigma factor regulator VreR (G3XCU5), another
upregulated protein, is likely involved in modulating the expression of genes that help the strain
withstand antibiotic-induced stress, contributing to its resilience (Matilla *et al.*, 2022).

Interestingly, Tse4 (Q9I069), a toxin protein associated with the Type VI secretion system, is
downregulated in the clinical strain. This could potentially be caused by protein degradation. While
RNA-seq data is not available for this study, future investigations comparing mRNA levels with protein
expression could provide further insights. However, this could suggest a strategic conservation of
resources, where the strain prioritizes essential survival functions over energy-intensive processes like
toxin production under antibiotic stress. The associated GO terms, such as "beta-lactamase activity"
(GO:0008800) and "sigma factor antagonist activity" (GO:0016989), highlight the biochemical
pathways enhanced in response to imipenem, pointing to a multifaceted resistance mechanism
involving both direct antibiotic degradation and stress response modulation. In contrast, the control
strain exhibits a broad activation of cellular pathways in response to increasing imipenem
concentrations.

Based on the Gene Ontology (GO) terms enriched in the *Pseudomonas aeruginosa* control strain
comparing 1.0x MIC to the (no imipenem) control probes (**Figure 3a and Table 3**), there are several
key processes of interest. The significant enrichment of GO terms related to membrane integrity and
transport processes, such as "periplasmic space" (GO:0042597) and "efflux transmembrane transporter
activity" (GO:0015562), suggests a defensive strategy focused on expelling the antibiotic and
maintaining membrane stability (Young *et al.*, 2019; Heywood and Lamont, 2020).

The upregulation of proteins involved in the "pyoverdine biosynthetic process" (GO:0002049) further
indicates an enhancement of iron acquisition systems, potentially as a countermeasure to the oxidative
stress imposed by imipenem. In addition, the production of pyoverdines, siderophores is essential for
iron acquisition, growth and survival by mediating biofilm formation and pathogenicity (Dell'Anno *et*
al., 2022). In addition to the broader activation of intracellular processes, as indicated by the
enrichment of GO terms like "cytosol" (GO:0005829) and "cytoplasm" (GO:0005737), there is

significant enrichment of terms related to the "ATP-Binding Cassette (ABC) Transporter Complex", which may play a role as an antibiotic efflux transporter (Lewis, Ween and McDevitt, 2012). Also noteworthy is the involvement of Type VI Protein Secretion System Complex, often linked to biofilm production as an antagonistic mechanism (Hespanhol, Nóbrega-Silva and Bayer-Santos, 2023). Interestingly, some studies suggest that imipenem can inhibit biofilm production, which is associated with Type VI Secretion System (de Sousa *et al.*, 2023). This further illustrates the control strain's strategy to activate a broad spectrum of defensive pathways to cope with antibiotic threat.

As reflected by the results from **Table 4** (**Figure 5d**), the proteomic response in the clinical strain is more targeted, it involves fewer GO terms, reflecting stable resistance mechanism that does not require extensive proteasomal reconfigurations. The enrichment of "periplasmic space" (GO:0042597) and "cytosol" (GO:0005829) suggests that the clinical strain, like the control strain, emphasizes membrane integrity and cytoplasmic processes. However, the clinical strain does not show a broad spectrum of activated pathways, suggesting either pre-existing or well-adapted mechanisms that require little alteration in response to imipenem. What is remarkable, is the significant enrichment of uncharacterized proteins, indicated by the "NOGO" term, suggesting that the clinical strain may use novel or less-characterized proteins to its resistance strategy, e.g. by proteins like PA2384 (Q9I195) or PA4129 (Q9HWQ2) in **Figures 7c** and **7d**, to be further discussed below. This is indicative of potential areas where further research could uncover new aspects of antimicrobial resistance. If confirmed, and further investigated, the role of these proteins might underscore novel proteomic profiles linked to adaptation to antibiotic stress.

A direct comparison of protein expression between the clinical and control strains at 1.0x MIC reveal a clear difference in their strategies to cope with imipenem (**Table 5**, **Figure 6d**). The control strain shows substantial upregulation of proteins involved in membrane and transport processes, reflecting a reactive and broad-spectrum response to antibiotic pressure. The significant enrichment of GO terms like "glycine betaine transport" (GO:0031460) and "ABC transporter complex" (GO:0043190) suggests that the control strain activates mechanisms to manage osmotic stress and expel imipenem, respectively. Glycine betaine is known to protect cells against osmotic stress (Zhao *et al.*, 2018), and its increased transport activity could be part of the control strain's attempt to mitigate the effects of imipenem-induced stress. In contrast, the clinical strain shows enrichment in oxidative stress management pathways, such as "cytochrome-c peroxidase activity" (GO:0004130), highlighting its focus on specific oxidative stress management rather than broad reactive changes. The presence of "extracellular region" (GO:0005576) enrichment in the clinical strain suggests it may involve extracellular defense mechanisms, potentially modifying or interacting with its environment to resist the antibiotic. The pronounced differences in protein expression and overall cellular responses to imipenem between the clinical and control strains of *Pseudomonas aeruginosa* are further elucidated by the analysis of selected proteins with >8-fold differential expression, as detailed in **Table 6** and visualized in **Figure 7.** The clinical strain's ability to maintain a relatively stable proteomic profile across varying concentrations of imipenem contrasts starkly with the control strain's more reactive and fluctuating proteomic responses, which likely reflect a broad-spectrum attempt to mitigate antibiotic stress. Uncharacterized proteins, such as PA2384 (Q9I195) and PA4129 (Q9HWQ2), exhibit significant upregulation in the clinical strain, suggesting these proteins may be integral to its resistance mechanisms. The consistent expression of these proteins, even under high antibiotic pressure, indicates

a potentially critical role in the clinical strain's adaptation and survival. This upregulation, coupled
with the stable expression of well-characterized proteins like AprF (Q03027) and AlgA (P07874),
underscores the clinical strain's reliance on pre-established resistance mechanisms, which seem to
operate efficiently without the need for extensive proteomic reconfiguration in response to imipenem.
In contrast, the control strain demonstrates significant proteomic shifts, particularly in proteins
associated with membrane transport and stress response, such as PA4514 (G3XCY8) and Wzm
(Q9HTB8). These shifts reflect the strain's reactive adjustments to counteract the effects of imipenem,
emphasizing a less specialized and more variable approach to resistance. The marked increase in
expression of alginate biosynthesis proteins in the control strain at higher MICs also suggests an
attempt to enhance biofilm formation as a defensive measure, a strategy that the clinical strain does not
appear to rely on as heavily, likely due to its already optimized biofilm-related defenses.
These observations highlight the distinct proteomic strategies employed by the clinical and control
strains. The clinical strain's more stable and efficient resistance mechanisms, which involve both
characterized and novel proteins, contrast with the control strain's broader and more reactive defense
responses. These findings suggest that the uncharacterized proteins in the clinical strain could be targets
for future research aimed at uncovering novel antimicrobial resistance mechanisms and developing
targeted therapies.

5 Conclusion

This study provides a comprehensive comparative analysis in *Pseudomonas aeruginosa* of the
proteomic responses of a clinical strain, representative of the world epidemic clone (*P. aeruginosa*
ST235), and a control strain (*P. aeruginosa* ATCC 27853) to varying concentrations of imipenem. The
clinical strain's ability to maintain a stable proteomic profile, even under high antibiotic pressure,
underscores its reliance on pre-established resistance mechanisms. Both well-characterized and
uncharacterized proteins have been found to be involved in the clinical strain's adaptation and survival.
This stability contrasts sharply with the control strain's more reactive and fluctuating proteomic
responses, which reflect a broad attempt to mitigate antibiotic stress. The control strain's significant
proteomic shifts, particularly in proteins associated with membrane transport and stress response,
highlight its less specialized and more variable approach to resistance. These findings highlight the
distinct proteomic strategies employed by the clinical and control strains. The clinical strain's efficient
and stable resistance mechanisms, involving both characterized and novel proteins, contrast with the
control strain's broader and more reactive defense responses. The identification of uncharacterized
proteins in the clinical strain as potential resistome targets for future research underscores the
importance of further investigation into novel antimicrobial resistance mechanisms. This could lead to
the development of targeted therapies, providing new avenues in the fight against antimicrobial
resistance.
612

Declaration of Interest

The authors declare no competing interests.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT-4 (via Microsoft Copilot) for proofreading and suggestions for figure plotting in Python. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Author Contributions

FDB, ME conceived and designed the study. ME, ASh, AK, ASk performed the experiments. ME, ASh, FDB analyzed the data. ME, ASh, FDB wrote the manuscript. All authors discussed the results and contributed to the final manuscript. FDB: Conceptualization, Project administration, Supervision. ME, ASh, AK: Methodology, Investigation. ME, ASh: Formal analysis, Data curation. ME, FDB: Writing – original draft. All authors have read and agreed to the published version of the manuscript.

Acknowledgments

This manuscript was developed under the project "Yeast-based biosensors for the specific and accessible detection of pathogens and antimicrobial resistance" (Acronym: AntiRYB, Reference Number: JPIAMR_2019_P08) from the Antimicrobial Resistance Joint Programming Initiative (JPI AMR) and the Research Council of Norway (RCN project number 311248). The work performed at the National Medicines Institute was funded by the National Science Centre, Poland under JPI-EC-AMR Call 2019 (NCN project number 2019/01/Y/NZ6/00006). MS-data is handled under Notur/NorStore Project NN9036K/NS9036K by PROMEC, a member of the National Network of Advanced Proteomics Infrastructure (NAPI), which is funded by the RCN INFRASTRUKTUR-program (295910).

Data Availability Statement

The datasets generated for this study can be found via ProteomeXchange with identifier PXD055744 (Username: reviewer_pxd055744@ebi.ac.uk Password: XNtCwVbI4AIs).

Supplementary Material

The supplementary material includes the following tables, which are provided as XLSX files:

Supplementary Table S1: Dose-correlation statistics for control strain.
Supplementary Table S2: Dose-correlation statistics for clinical strain.
Supplementary Table S3: Tabular GO-term enrichment results corresponding to **Figure 5b**.
Supplementary Table S4: Tabular GO-term enrichment results corresponding to **Figure 5c**.
Supplementary Table S5: Tabular GO-term enrichment results corresponding to **Figure 6b**.
Supplementary Table S6: List of protein IDs associated with GO terms in **Table 5**.

References:

Aslam, B. *et al.* (2018) 'Antibiotic resistance: a rundown of a global crisis', *Infection and Drug Resistance*, 11, pp. 1645–1658. Available at: https://doi.org/10.2147/IDR.S173867.

Barbier, F. *et al.* (2023) 'Rationale and evidence for the use of new beta-lactam/beta-lactamase inhibitor combinations and cefiderocol in critically ill patients', *Annals of Intensive Care*, 13(1), p. 65. Available at: https://doi.org/10.1186/s13613-023-01153-6.

Berrazeg, M. *et al.* (2015) 'Mutations in β-Lactamase AmpC Increase Resistance of Pseudomonas aeruginosa Isolates to Antipseudomonal Cephalosporins', *Antimicrobial Agents and Chemotherapy*, 59(10), pp. 6248–6255. Available at: https://doi.org/10.1128/AAC.00825-15.

Breidenstein, E.B.M., de la Fuente-Núñez, C. and Hancock, R.E.W. (2011) 'Pseudomonas aeruginosa: all roads lead to resistance', *Trends in Microbiology*, 19(8), pp. 419–426. Available at: https://doi.org/10.1016/j.tim.2011.04.005.

Carbon, S. and Mungall, C. (2024) 'Gene Ontology Data Archive'. Zenodo. Available at: https://doi.org/10.5281/ZENODO.12173881.

Cox, J. *et al.* (2014) 'Accurate Proteome-wide Label-free Quantification by Delayed Normalization and Maximal Peptide Ratio Extraction, Termed MaxLFQ', *Molecular & Cellular Proteomics*, 13(9), pp. 2513–2526. Available at: https://doi.org/10.1074/mcp.M113.031591.

Darby, E.M. *et al.* (2023) 'Molecular mechanisms of antibiotic resistance revisited', *Nature Reviews Microbiology*, 21(5), pp. 280–295. Available at: https://doi.org/10.1038/s41579-022-00820-y.

Davies, J. and Davies, D. (2010) 'Origins and evolution of antibiotic resistance', *Microbiology and molecular biology reviews: MMBR*, 74(3), pp. 417–433. Available at: https://doi.org/10.1128/MMBR.00016-10.

Dell'Anno, F. *et al.* (2022) 'Novel Insights on Pyoverdine: From Biosynthesis to Biotechnological Application', *International Journal of Molecular Sciences*, 23(19), p. 11507. Available at: https://doi.org/10.3390/ijms231911507.

Elfadadny, A. *et al.* (2024) 'Antimicrobial resistance of Pseudomonas aeruginosa: navigating clinical impacts, current resistance trends, and innovations in breaking therapies', *Frontiers in Microbiology*, 15, p. 1374466. Available at: https://doi.org/10.3389/fmicb.2024.1374466.

European Committee on Antimicrobial Susceptibility Testing (2024) *Broth microdilution - EUCAST reading guide v 5.0*. Available at: https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/MIC_testing/Reading_guide_BMD_v_5.0_2024.pdf.

Fair, R.J. and Tor, Y. (2014) 'Antibiotics and bacterial resistance in the 21st century', *Perspectives in Medicinal Chemistry*, 6, pp. 25–64. Available at: https://doi.org/10.4137/PMC.S14459.

Goodyear, M.C. *et al.* (2023) 'Quantitative proteomics reveals unique responses to antimicrobial treatments in clinical *Pseudomonas aeruginosa* isolates', *mSystems*. Edited by J.E. Elias, 8(5), pp. e00491-23. Available at: https://doi.org/10.1128/msystems.00491-23.

Hashemi, M.M. *et al.* (2019) 'Proteomic Analysis of Resistance of Gram-Negative Bacteria to
Chlorhexidine and Impacts on Susceptibility to Colistin, Antimicrobial Peptides, and Ceragenins',
Frontiers in Microbiology, 10, p. 210. Available at: https://doi.org/10.3389/fmicb.2019.00210.

Hespanhol, J.T., Nóbrega-Silva, L. and Bayer-Santos, E. (2023) 'Regulation of type VI secretion
systems at the transcriptional, posttranscriptional and posttranslational level', *Microbiology*, 169(8).
Available at: https://doi.org/10.1099/mic.0.001376.

Heywood, A. and Lamont, I.L. (2020) 'Cell envelope proteases and peptidases of *Pseudomonas
aeruginosa* : multiple roles, multiple mechanisms', *FEMS Microbiology Reviews*, 44(6), pp. 857–
873. Available at: https://doi.org/10.1093/femsre/fuaa036.

'ISO 20776-1:2020. Clinical laboratory testing and in vitro diagnostic test systems—Susceptibility
testing of infectious agents and evaluation of performance of antimicrobial susceptibility test devices.
Part 1: Broth micro-dilution reference method for testing the in vitro activity of antimicrobial agents
against rapidly growing aerobic bacteria involved in infectious diseases.' (2020). International
Organization for Standardization (ISO).

Jacoby, G.A. (2009) 'AmpC β-Lactamases', *Clinical Microbiology Reviews*, 22(1), pp. 161–182.
Available at: https://doi.org/10.1128/CMR.00036-08.

Jongers, B. *et al.* (2021) 'Proteomic response of P. aeruginosa to sub-MIC antibiotics', in. *31st
European Congress of Clinical Microbiology and Infection Diseases (ECCMID)*, Online.

Khodadadi, Ehsaneh *et al.* (2020) 'Proteomic Applications in Antimicrobial Resistance and Clinical
Microbiology Studies', *Infection and Drug Resistance*, 13, pp. 1785–1806. Available at:
https://doi.org/10.2147/IDR.S238446.

Langendonk, R.F., Neill, D.R. and Fothergill, J.L. (2021) 'The Building Blocks of Antimicrobial
Resistance in Pseudomonas aeruginosa: Implications for Current Resistance-Breaking Therapies',
Frontiers in Cellular and Infection Microbiology, 11, p. 665759. Available at:
https://doi.org/10.3389/fcimb.2021.665759.

Lautenbach, E. *et al.* (2006) 'Imipenem Resistance Among *Pseudomonas aeruginosa* Isolates Risk
Factors for Infection and Impact of Resistance on Clinical and Economic Outcomes', *Infection
Control & Hospital Epidemiology*, 27(9), pp. 893–900. Available at: https://doi.org/10.1086/507274.

Leigue, L., Montiani-Ferreira, F. and Moore, B.A. (2016) 'Antimicrobial susceptibility and minimal
inhibitory concentration of Pseudomonas aeruginosa isolated from septic ocular surface disease in
different animal species', *Open Veterinary Journal*, 6(3), pp. 215–222. Available at:
https://doi.org/10.4314/ovj.v6i3.9.

Lewis, V.G., Ween, M.P. and McDevitt, C.A. (2012) 'The role of ATP-binding cassette transporters
in bacterial pathogenicity', *Protoplasma*, 249(4), pp. 919–942. Available at:
https://doi.org/10.1007/s00709-011-0360-8.

Liu, X. *et al.* (2016) 'Proteomic response of methicillin-resistant S. aureus to a synergistic
antibacterial drug combination: a novel erythromycin derivative and oxacillin', *Scientific Reports*,
6(1), p. 19841. Available at: https://doi.org/10.1038/srep19841.

Magiorakos, A.-P. *et al.* (2012) 'Multidrug-resistant, extensively drug-resistant and pandrug-resistant
bacteria: an international expert proposal for interim standard definitions for acquired resistance',
Clinical Microbiology and Infection: The Official Publication of the European Society of Clinical
Microbiology and Infectious Diseases, 18(3), pp. 268–281. Available at:
https://doi.org/10.1111/j.1469-0691.2011.03570.x.

Matilla, M.A. *et al.* (2022) 'Virulence Induction in Pseudomonas aeruginosa under Inorganic
Phosphate Limitation: a Proteomics Perspective', *Microbiology Spectrum*. Edited by G. Rampioni,
10(6), pp. e02590-22. Available at: https://doi.org/10.1128/spectrum.02590-22.

Mosquera-Rendón, J. *et al.* (2016) 'Pangenome-wide and molecular evolution analyses of the
Pseudomonas aeruginosa species', *BMC Genomics*, 17(1), p. 45. Available at:
https://doi.org/10.1186/s12864-016-2364-4.

Oliver, A. *et al.* (2015) 'The increasing threat of Pseudomonas aeruginosa high-risk clones', *Drug*
Resistance Updates: Reviews and Commentaries in Antimicrobial and Anticancer Chemotherapy,
21–22, pp. 41–59. Available at: https://doi.org/10.1016/j.drup.2015.08.002.

Pérez-Llarena, F.J. and Bou, G. (2016) 'Proteomics As a Tool for Studying Bacterial Virulence and
Antimicrobial Resistance', *Frontiers in Microbiology*, 7, p. 410. Available at:
https://doi.org/10.3389/fmicb.2016.00410.

Perez-Riverol, Y. *et al.* (2022) 'The PRIDE database resources in 2022: a hub for mass spectrometry-
based proteomics evidences', *Nucleic Acids Research*, 50(D1), pp. D543–D552. Available at:
https://doi.org/10.1093/nar/gkab1038.

de Sousa, T. *et al.* (2023) 'Determination of Antimicrobial Resistance and the Impact of Imipenem +
Cilastatin Synergy with Tetracycline in Pseudomonas aeruginosa Isolates from Sepsis',
Microorganisms, 11(11), p. 2687. Available at: https://doi.org/10.3390/microorganisms11112687.

Sulaiman, J.E. and Lam, H. (2022) 'Proteomics in antibiotic resistance and tolerance research:
Mapping the resistome and the tolerome of bacterial pathogens', *Proteomics*, 22(8), p. e2100409.
Available at: https://doi.org/10.1002/pmic.202100409.

Tyanova, S., Temu, T. and Cox, J. (2016) 'The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics', *Nature Protocols*, 11(12), pp. 2301–2319. Available at:
https://doi.org/10.1038/nprot.2016.136.

UniProt Consortium (2022) 'Pseudomonas aeruginosa (strain PAO1) reference proteome'. Available
at: https://www.uniprot.org/proteomes/UP000002438.

UniProt Consortium (2023) 'UniProt: the Universal Protein Knowledgebase in 2023', *Nucleic Acids*
Research, 51(D1), pp. D523–D531. Available at: https://doi.org/10.1093/nar/gkac1052.

Urbanowicz, P. *et al.* (2021) 'Molecular and genomic epidemiology of VIM/IMP-like metallo-β-
lactamase-producing *Pseudomonas aeruginosa* genotypes in Poland', *Journal of Antimicrobial*
Chemotherapy, 76(9), pp. 2273–2284. Available at: https://doi.org/10.1093/jac/dkab188.

Wright, G.D. (2007) 'The antibiotic resistome: the nexus of chemical and genetic diversity', *Nature*
Reviews Microbiology, 5(3), pp. 175–186. Available at: https://doi.org/10.1038/nrmicro1614.

Xu, Y. *et al.* (2020) 'Mechanisms of Heteroresistance and Resistance to Imipenem in Pseudomonas
aeruginosa', *Infection and Drug Resistance*, Volume 13, pp. 1419–1428. Available at:
https://doi.org/10.2147/IDR.S249475.

Young, K. *et al.* (2019) 'In vitro studies evaluating the activity of imipenem in combination with
relebactam against Pseudomonas aeruginosa', *BMC Microbiology*, 19(1), p. 150. Available at:
https://doi.org/10.1186/s12866-019-1522-7.

Zhao, G. *et al.* (2018) 'Betaine in Inflammation: Mechanistic Aspects and Applications', *Frontiers in*
Immunology, 9, p. 1070. Available at: https://doi.org/10.3389/fimmu.2018.01070.