Medicago truncatula CORYNE **regulates inflorescence meristem branching, nutrient signaling,**

and arbuscular mycorrhizal symbiosis

Johnathan Orosz[1], Erica Xinlei Lin[1]*, Penelope Lindsay[2]*, Mikayla Kappes[3], Sagar Bashyal[1,3,4],

Hasani Everett[1], Chandan Kumar Gautam[1], David Jackson[2], Lena Maria Müller[1,3#]

* equal contribution

[#] corresponding author

[1]Department of Biology, University of Miami, Coral Gables, FL, USA

[2]Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, USA

[3]The Salk Institute for Biological Studies, La Jolla, CA, USA

[4]School of Biological Sciences, University of California San Diego, La Jolla, CA 92093

Present addresses:

E.X.L.: Stowers Institute for Medical Research, Kansas City, MO 64110

H.E.: Systems, Synthetic, and Physical Biology, Rice University, Houston, TX 77005

C.K.G.: Department of Microbiological Sciences, North Dakota State University, Fargo, ND 58108

Abstract

The CLAVATA signaling pathway regulates plant development and plant-environment

interactions. CLAVATA signaling consists of mobile, cell-type or environment-specific

CLAVATA3/ESR-related (CLE) peptides, which are perceived by a receptor complex consisting of

leucine-rich repeat receptor-like kinases such as CLAVATA1 and receptor-like proteins such as

CLAVATA2, which often functions with the pseudokinase CORYNE (CRN). CLAVATA signaling has

been extensively studied in various plant species for its role in meristem maintenance and in

legumes for modulating root interactions with nitrogen-fixing rhizobia. Some signaling proteins

involved in development and nodulation, including CLAVATA1, also regulate plant interactions

with mutualistic arbuscular mycorrhizal (AM) fungi. However, our knowledge on AM symbiosis

regulation by CLAVATA signaling remains limited and only a handful of genetic regulators have

been identified. Here we report that *Medicago truncatula CRN* controls inflorescence meristem

branching and negatively regulates root interactions with AM fungi. *MtCRN* functions partially

independently of the AM autoregulation signal *MtCLE53*. Transcriptomic data revealed that *crn*

roots display signs of perturbed signaling related to a range of responses to biotic and abiotic

factors, suggesting that *MtCRN* plays various roles in plant development and interactions with

the environment.

Introduction

Cell-cell communication is critical for plant development and cellular responses to internal and

external cues. CLAVATA (CLV) signaling, which consists of mobile CLAVATA3/ESR-related (CLE)

peptide hormones and CLE receptor complexes, regulates many aspects of plant development

and responses to the environment (Bashyal *et al.*, 2023; Lindsay *et al.*, 2023). CLE receptor

complexes consist of leucine-rich repeat receptor-like kinases (LRR-RLKs) with an extracellular

LRR ligand-binding domain and an intracellular kinase domain (e.g., CLAVATA1, CLV1, or BARELY

ANY MERISTEM, BAM1-3), receptor-like proteins (e.g., CLAVATA2, CLV2) containing an

extracellular ligand-binding domain but lack a kinase domain, and intracellular regulators like

the membrane-bound pseudokinase CORYNE (CRN) (Hazak and Hardtke, 2016). In Arabidopsis

and maize, CLV2 and CRN are considered a CLE perception unit (Bleckmann *et al.*, 2009; Je *et al.*,

2018). CLV2-CRN can form complexes with CLV1 or BAMs, but also function independently as a

parallel pathway (Guo *et al.*, 2010; Somssich *et al.*, 2016*b*; Hazak *et al.*, 2017). Similarly, the

Medicago truncatula ortholog of CLV1 (termed SUNN in this species), MtCLV2 and MtCRN can

interact with each other (Crook *et al.*, 2016). While the cellular function of CRN is still elusive,

co-expression assays in *Nicotiana benthamiana* leaves suggest that the short extracellular

domain of AtCRN is important for stabilizing the CRN-CLV2 complex at the plasma membrane,

whereas the intracellular kinase domain is important for transmitting CLE peptide signals

(Somssich *et al.*, 2016*a*).

Mutations in CLV receptors and ligands disrupt meristem organization, often leading to enlarged vegetative, inflorescence and floral shoot meristems (Lindsay *et al.*, 2023). These meristem defects in turn impact floral organization and fruit development, altering the number of flowers produced in an inflorescence, the number of floral organs that are produced, or the arrangement and size of the fruit. The impact of a particular CLV mutation on meristem organization depends on the specific gene mutated, the nature of the mutation (allele), and the plant species; this phenomenon is thought to be a result of differences in compensatory mechanisms in meristem maintenance among plant species (Rodriguez-Leal *et al.*, 2019). To date, maize and Arabidopsis are the only plant species in which *crn* mutants have been described. In both species, shoot and inflorescence meristems are enlarged. Arabiopsis *crn* mutants produce more carpels per flower, and some flowers make stamens without anthers (Müller *et al.*, 2008). Depending on the temperature, *crn* floral primordia and flowers are prematurely terminated (Jones *et al.*, 2021). Maize *crn* mutants produce fasciated ears, with a disordered arrangement of kernel rows and larger kernel row number (Je *et al.*, 2018). CLV signaling also plays roles in root development, including maintenance of the root apical meristem and protophloem differentiation. *AtCRN* is expressed in multiple cell types in the root, including stele, endodermis, and quiescent center cells of the meristem (Somssich *et al.*, 2016*a*; Hazak *et al.*, 2017). Phloem-expressed AtCRN was shown to be required for the perception of root-active CLE peptides (Hazak *et al.*, 2017). Despite its importance for CLE signaling not all Arabidopsis *crn* mutant alleles show a phenotype characterized by longer primary roots (Müller *et al.*, 2008; Somssich *et al.*, 2016*a*; Hazak *et al.*, 2017), suggesting *CRN* by itself is not required for root development.

In addition to plant development, CLAVATA signaling pathways also modulate plant interactions with the environment, which includes responses to abiotic and biotic factors such as host responses to pathogenic or beneficial microbes (Bashyal *et al.*, 2023). Mutualistic plant interactions with rhizobia during nodulation or *Glomeromycotina* fungi during arbuscular mycorrhiza (AM) symbiosis are regulated by various receptor-ligand signaling pathways (Roy and Müller, 2022), including CLV signaling. In both symbioses, the host plant provides microbial

symbionts with photosynthetically fixed carbon in exchange for mineral nutrients. While

rhizobia supply the host with nitrogen (N), AM fungi provide a range of benefits including N,

phosphorus (P), and water (Smith and Reid, 2008). Both symbioses are governed by negative

regulatory pathways based on plant nutrient (N or P) status and prior microbial colonization

(autoregulation); these pathways are mediated by nutrient- or symbiosis-induced CLE peptides,

respectively (Roy and Müller, 2022). Autoregulation of nodulation (AON) is a systemic negative

feedback loop that has been studied in numerous legume species (Wang *et al.*, 2018). In *M.*

truncatula, AON is regulated by root-derived, nodule-induced CLE peptides (MtCLE12,

MtCLE13), which negatively regulate nodule number in a MtSUNN- and MtCRN-dependent

manner (Schnabel *et al.*, 2005; Crook *et al.*, 2016; Nowak *et al.*, 2019). Grafting experiments

revealed that MtSUNN and MtCRN act in the shoot (Schnabel *et al.*, 2005; Crook *et al.*, 2016);

however, both genes are also expressed in roots (Schnabel *et al.*, 2012; Crook *et al.*, 2016),

suggesting various functions across tissues and organs. Autoregulation of mycorrhizal symbiosis

(AOM) is regulated by the CLE peptide MtCLE53, which is induced in the vascular tissue of

colonized roots and negatively regulates AM fungal colonization by repressing biosynthesis of

strigolactones (Müller *et al.*, 2019; Karlo *et al.*, 2020, Preprint). Strigolactones are produced by

host plants in response to P and N starvation and promote AM symbiosis (Yoneyama *et al.*,

2007; Liu *et al.*, 2011; Foo *et al.*, 2013; Marro *et al.*, 2022). In the current AOM model, AM-

induced *MtCLE53* down-regulates strigolactone biosynthesis and consequently reduces

subsequent AM fungal root colonization (Müller *et al.*, 2019).

Like AON, AOM depends on the LRR-RLK SUNN. Plants defective in *SUNN* display a

hypermycorrhizal phenotype with elevated root colonization levels and arbuscule numbers and

are insensitive to *MtCLE53* overexpression (Morandi *et al.*, 2000; Müller *et al.*, 2019; Karlo *et al.*,

2020, Preprint). Despite many similarities, AOM is less understood than AON. Although AON

and AOM signal through the same LRR-RLK SUNN, the upstream-acting CLE peptides and

downstream signaling appear to at least partially differ between the two symbioses (Mortier *et*

al., 2010; Müller *et al.*, 2019; Karlo *et al.*, 2020, Preprint; Chaulagain *et al.*, 2023*b,a*).

Furthermore, recent evidence from tomato (*Solanum lycopersicum*) suggests that AM symbiosis

is regulated by multiple, at least partially independent CLV signaling pathways: grafting

experiments revealed that while SlFAB (ortholog of CLV1) acts in the root, SlCLV2 is part of two

distinct signaling pathways controlling AM symbiosis from the shoot and the root, respectively

(Wang *et al.*, 2021). In addition, the AM-induced SlCLE11 peptide acts as a negative regulator of

AM symbiosis but functions independently of SlFAB or SlCLV2 (Wulf *et al.*, 2024). By contrast,

supernodulating pea mutants in *CLV2* (*sym28*), did not show a hypermycorrhizal phenotype,

suggesting species-specific differences in symbiosis regulation (Morandi *et al.*, 2000). In

addition, in contrast to *MtCLE53* over-expression, *sunn* itself (as well as mutants in the tomato

and pea orthologs) does not exhibit mis-regulated strigolactone biosynthesis (Foo *et al.*, 2014;

Müller *et al.*, 2019; Karlo *et al.*, 2020, Preprint; Wang *et al.*, 2021), suggesting additional

signaling components must be involved in MtCLE53-SUNN-mediated AOM signaling that impact

strigolactone biosynthesis.

In addition to autoregulation, AM symbiosis is tightly regulated by the availability of soil mineral

nutrients to the plant. High N or P concentrations in the substrate can suppress AM symbiosis

development (Abbott *et al.*, 1984). Although P is the predominant element plants acquire

through AM fungi, N starvation can promote AM symbiosis even under high P (Nouri *et al.*,

2014). In *M. truncatula*, P-induced *MtCLE33* acts as a negative regulator of AM symbiosis via a

SUNN-dependent pathway although SUNN itself is not required for P repression of AM

symbiosis (Müller *et al.*, 2019). Similarly, work in tomato revealed that regulation of AM by CLV

receptors is independent of P but depends on plant N status (Wang *et al.*, 2021). Despite

substantial progress in recent years, our knowledge on AM symbiosis regulation by CLV signaling

is incomplete and fragmented across plant species. Similar to meristem development, CLV

signaling may be differently wired across plant species with different nutrient uptake strategies

and capacities for symbiotic interactions.

Here, we report *MtCRN* as a regulator of *M. truncatula* inflorescence branching. In addition, we

find that *MtCRN* acts as a negative regulator of AM symbiosis in *M. truncatula*. Our results

further suggest that, while *MtCRN* contributes to AOM, it also regulates AM symbiosis via

additional pathways. Transcriptomics data suggest that *crn* mutants display broad alterations in

response to biotic and abiotic factors, indicating that MtCRN may integrate multiple signaling

pathways with distinct outcomes.

Results

***M. truncatula* CLAVATA signaling regulates inflorescence meristem branching and flowering**

time but is dispensable for root development

Mutants in CLV signaling have previously been described to be impaired in meristem

maintenance, with mutants often displaying inflorescence or stem fasciation (Clark *et al.*, 1997).

Arabidopsis *CRN* specifically has been reported to modulate inflorescence meristem termination

and flower outgrowth under stress (Müller *et al.*, 2008; Jones *et al.*, 2021). Although legume CLV

mutants have been extensively studied in the context of plant-microbe symbioses, reports on

developmental phenotypes are rare. Notable exceptions are pea *sym28* mutants, which display

stem fasciation and produce excess leaves and flowers per node (Krusell *et al.*, 2011), and

GmCLV1A, which regulates shoot branching as well as flower number and development, and

appears to have functionally diversified from its symbiosis-regulating paralog *GmNARK* (Mirzaei

et al., 2017).

To understand if CLV signaling is also involved in shoot development of other legumes, we

investigated above-ground developmental phenotypes of *M. truncatula* mutants carrying a

homozygous Tnt1 insertion within the coding sequence of the pseudokinase *MtCRN* (Crook *et*

al., 2016). Above-ground development, including time to flowering, appeared to be delayed in

crn when compared to R108 wild-types (Supplementary Fig.1). Stem fasciation, which is

characteristic for many CLV pathway mutants in other species (Somssich *et al.*, 2016*b*), was not

observed. *M. truncatula* typically produces one solitary flower per inflorescence unit (Fig. 1A, B)

(Benlloch *et al.*, 2003). However, we noted that *crn* mutants displayed two flowers per peduncle

with high frequency: 91% of 8-week old *crn* (n= 11 inflorescences) produced two flowers (often

with an additional third bud appearing from the same inflorescence, Fig. 1C, D, Supplementary

Fig. 1), whereas R108 wildtype plants of the same age and grown under the same growth

conditions only produced a single flower per peduncle (n=14 inflorescences, Fig. 1A, B,

Supplementary Fig. 1). Notably, similar inflorescence phenotypes (76% of inflorescences, n=25)

were detected in *M. truncatula* mutants defective in the *CLV1* ortholog *SUNN* (*sunn-5*;

Supplementary Fig. 1), suggesting both *SUNN* and *MtCRN* regulate inflorescence meristem

branching. Although *crn* mutants produced two flowers per peduncle, ultimately only a single

seed was produced per inflorescence as one of the two flowers typically appeared to terminate

prematurely. In Arabidopsis, *crn* mutants show increased numbers of floral organs and

increased inflorescence meristem size (Müller *et al.*, 2008). However, manual dissection of

flowers as well as scanning electron microscopy analysis of inflorescences revealed that floral

organ numbers and inflorescence width of *M. truncatula crn* did not differ from those in R108

wildtypes (Figure 1 A-G). This indicates that, unlike Arabidopsis and maize, meristem branching

rather than stem cell proliferation is altered in *M. truncatula crn*.

In roots, CLV signaling is critical for root meristem maintenance, lateral root emergence, and

protophloem differentiation (Hazak and Hardtke, 2016). We set out to investigate if *M.*

truncatula crn mutants play a role in root development by analyzing root system architecture of

12-day old *crn* and R108 seedlings grown on agar plates. *M. truncatula crn* mutants did not

display strong defects in root system development in our experimental growth conditions. We

did not detect differences in primary root length, lateral root number, lateral root length, or

lateral root density (lateral roots per cm primary root length) between *crn* and R108

(Supplementary Figure 1). This is reminiscent of the phenotypes previously reported in

Arabidopsis, where *crn* mutants do not show altered root development although *CRN* is

required for the perception of root-active CLE peptides (Müller *et al.*, 2008; Hazak *et al.*, 2017).

***MtCRN* negatively regulates AM symbiosis**

M. truncatula CRN was previously described as a negative regulator of nodulation symbiosis in

concert with SUNN (Crook *et al.*, 2016; Nowak *et al.*, 2019). AON, AOM, and nutrient regulation

of AM symbiosis share multiple common components, including CLV1, but the role of CRN in

AM symbiosis regulation has not been investigated. Because tomato *clv2* mutants are

hypermycorrhizal (Wang *et al.*, 2018), CRN-CLV2 are commonly considered a signaling unit

(Müller *et al.*, 2008), and MtCRN can interact with the AOM regulator SUNN as well as MtCLV2

(Crook *et al.*, 2016), we hypothesized that *MtCRN* may also regulate root interactions with AM

fungi. Indeed, *crn* mutants displayed significantly increased overall *R. irregularis* root

colonization levels relative to the R108 wildtype (Figure 2A). A detailed analysis revealed that

crn mutants also show higher arbuscule density (i.e., number of arbuscules per mm colonized

root in a defined region at the hyphopodium) than wildtype roots, whereas vesicle density

(vesicles/mm colonized roots) did not differ between the wild-type and the mutant (Figure 2B-

E). These results are consistent with the hypothesis that MtCRN acts as a negative regulator of

AM symbiosis.

***MtCRN* functions partially independent of the AOM signal MtCLE53**

Autoregulation of nodulation (AON) and mycorrhizal symbiosis (AOM) negatively regulates

nodulation and AM symbiosis, respectively, based on existing symbioses. *MtCRN* has been

implicated in AON and acts downstream of the nodulation-induced CLE peptides MtCLE12 and

MtCLE13 (Nowak *et al.*, 2019). We previously reported that ectopic over-expression of the *M.*

truncatula AOM signal *MtCLE53* reduced overall root length colonization in a SUNN-dependent

manner (Müller *et al.*, 2019). To test if *MtCRN* also acts down-stream of AOM signals, we over-

expressed *MtCLE53* under the control of the constitutively active Cauliflower mosaic virus 35S

promoter (*35S::MtCLE53*) as well as a control construct (*35S::GUS*) in roots of *crn* mutants and

R108 wild-type controls. Similar to what was previously reported for the *M. truncatula* A17

accession (Müller *et al.*, 2019; Karlo *et al.*, 2020, Preprint), ectopic over-expression of *MtCLE53*

roots also reduced overall AM fungal root length colonization in R108. Interestingly, unlike *sunn*,

which is fully insensitive to *MtCLE53* over-expression suggesting the LRR-RLK acts in the same

pathway as the peptide hormone (Müller *et al.*, 2019), ectopic *MtCLE53* overexpression reduced

AM fungal root length colonization also in *crn* (Fig. 3). However, colonization reduction in

35S::MtCLE53 relative to *35S::GUS*-expressing roots is weaker in *crn* (29%) than in the wild-type

background (41%). The partial sensitivity of *crn* to *MtCLE53* over-expression suggests that

MtCRN is involved in *MtCLE53* signaling but that the AM symbiosis phenotype of *crn* may also

be influenced by additional, *MtCLE53*-independent pathways.

Transcriptomic analysis of *M. truncatula crn* mutants reveals global changes in stress

response, nutrient homeostasis, and environment interactions

To understand the molecular mechanisms causing the elevated AM fungal root colonization

levels in *crn* mutants, we sequenced the transcriptomes of 3 *R. irregularis*-colonized and 4

mock-inoculated *crn* and R108 wild-type root systems, respectively (Supplementary table 1). As

expected, we observed major transcriptomic changes induced by *R. irregularis* colonization in

R108 wild-type roots (434 up-regulated and 266 down-regulated genes, here defined as

differentially expressed genes with a log2 fold-change >1 or <-1, respectively, and padj<0.05;

Supplementary figure 2, Supplementary figure 3). Comparing *R. irregularis*-colonized *crn* mutant

roots with mock-inoculated *crn* roots, we identified 816 up- and 325-downregulated genes

(Supplementary figure 2, Supplementary figure 4). In mock-inoculated *crn* roots relative to

mock-inoculated wild-type roots, we identified 1506 up- and 647 down-regulated genes

(Supplementary figure 2, Supplementary figure 5). When comparing differentially expressed

genes in *R. irregularis*-colonized *crn* roots with *R. irregularis*-colonized wild-type roots, we

identified 1061 up- and 594-downregulated genes (Supplementary figure 2, Supplementary

figure 6, Supplementary table 2). Because *crn* roots are more colonized than wild-type roots,

these observed transcriptomic differences may be caused by genotype differences and/or

differences in AM fungal colonization levels. We performed hierarchical clustering analysis of all

differentially expressed genes in *R. irregularis*-inoculated *crn* roots relative to *R. irregularis*-

inoculated wild-type roots to identify clusters of differentially expressed genes with similar

expression patterns and to distinguish differentially expressed genes solely influenced by

colonization levels from those regulated by *MtCRN*.

Differentially expressed genes were grouped into 6 clusters (Fig. 4A, Supplementary table 2).

Clusters 1, 3 and 6 contained genes that are regulated by AM fungal root colonization, and

whose general expression pattern changes were similar in *crn* and wild-type roots. However, the

degree of up- or down-regulation differs between the genotypes, likely caused by the

differential colonization levels in the mutant compared to the wildtype (Fig. 2 A-E).

Cluster 1 genes are induced by AM symbiosis and are generally more highly expressed in *R.*

irregularis-colonized *crn* roots relative to colonized wild-type roots (Fig. 4A). GO analysis of the

genes in cluster 1 revealed, among others, an enrichment of genes associated with GO terms

related to symbiosis ("AM association", "response to symbiotic fungus", "nodulation"), nitrate

assimilation, as well as genes associated with the biosynthesis and transport of secondary

metabolites and hormones known to regulate AM symbiosis (e.g., carotenoids, gibberellic acid,

brassinosteroids; Fig. 4B, Supplementary table 3). This cluster contains well-studied AM marker

genes including but not limited to *MtPT4*, *MtRAM1*, *MtRAM2,* or *MtSTR*, which function in

bidirectional nutrient exchange between the symbiotic partner and whose expression is known

to be correlated with AM colonization levels (Serrano *et al.*, 2024).

Clusters 3 and 6 contain genes that are down-regulated by AM fungal root colonization; AM-

mediated down-regulation is more pronounced in *crn* roots than in wildtypes. While no

significant GO terms were found in cluster 3 (11 genes), in cluster 6 we detected enrichment of

GO terms related to cell wall biogenesis and remodeling (Fig. 4D, Supplementary table 3) as well

as ethylene signaling. Notably, this GO term contained 5 transcription factors of the

APETALA2/ETHYLENE RESPONSE FACTOR (AP2/ERF) family; however, AP2/ERF transcription

factors have a more role in plant stress signaling, and, apart from ethylene, also often respond

to the hormones abscisic acid (ABA), gibberellic acid (GA), brassinosteroids (BR), and cytokinin

(Xie *et al.*, 2019). To our knowledge, no symbiosis-related function has been reported for any of

the AP2/ERF transcription factors in cluster 6.

Cluster 2 contains 549 genes that are up-regulated in *crn* roots relative to wild-type roots,

regardless of AMF colonization status (Fig. 4A). Many genes in this cluster appear to play roles in

plant responses to abiotic and biotic stresses: The GO term with the highest fold-enrichment

(61-fold) in this cluster is saponin biosynthetic process (Fig. 4C, Supplementary table 3).

Saponins are secondary metabolites with antioxidant activity that play various roles in plant-

microbe interactions, including plant defense against pathogens and herbivores (Mugford and

Osbourn, 2012; Chen *et al.*, 2014). Soy saponins have been identified in root exudates as

modulators of plant interactions with rhizosphere microbes (Tsuno *et al.*, 2018; Fujimatsu *et al.*,

2020). In addition, 12 peroxidases associated with the GO terms "hydrogen peroxide catabolic

process" and "cellular oxidant detoxification" were enriched in cluster 2. Peroxidases are major

regulators of cellular reactive oxygen species (ROS) homeostasis (Passardi *et al.*, 2005). ROS play

various roles in plants, including response to pathogens and beneficial microbes, including AM,

but also development, signaling, and cell growth (Wojtaszek, 1997; Fester and Hause, 2005;

Huang *et al.*, 2019). The GO term "cellular modified amino acid metabolic process" was

enriched in cluster 2; this GO term contained (among others) 4 genes annotated as glutathione

S-transferases, which also have antioxidant activity and are involved in ROS scavenging and

cellular detoxification during stress responses (Gullner *et al.*, 2018). Together, this data suggests

that cellular oxidative stress may be attenuated in *crn* through increased ROS scavenging.

Conversely, *crn* roots display an enrichment of GO terms associated with lipid metabolism ("lipid

biosynthetic process", "fatty acid metabolic process", "monocarboxylic acid biosynthetic

process") and oxylipin biosynthesis. The latter contained four genes encoding 9-lipoxigenases

(*9-LOX*), and one gene annotated as *ALPHA-DIOXYGENASE 1 (αDOX1)*. 9-oxylipins play a role in

cellular signaling, defense, and abiotic stress response (Vicente *et al.*, 2012), and their up-

regulation suggests that *crn* mutants may have a heightened response to biotic stress even in

the absence of microbes. In line with this hypothesis, the GO term "defense response" was also

enriched among the genes in cluster 2 (Fig. 4C); this GO term contained several genes encoding

NUCLEOTIDE-BINDING SITE LEUCINE-RICH REPEAT (NBS-LRR) family proteins, which can directly

bind to microbial effectors and thus play critical roles in plant immune responses to pathogens

(Marone *et al.*, 2013). This suggests that, despite signatures of attenuated oxidative stress,

other components of the plant response system to biotic stresses are up-regulated in *crn*.

Furthermore, we detected an enrichment of genes associated with the GO term

"transmembrane transport" among the *crn*-upregulated genes in cluster 2. Interestingly, these

included, among others, genes encoding transporters for carbon products such as sugar

(*MtESL16.1/MtESL2.4, MtESL3.4, MtESL6.4, MtSTP3.2, MtSTP5.1/MtSTP5.2*), which may

indicate that carbon flux is altered in *crn*. In addition, we identified putative transporters for

nitrate (*MtNPF2.11, MtNPF5.13, MtNPF5.5, MtNPF5.10*), ammonium (*Medtr4g017820*),

phosphate (*MtPT2, MtPT9*), metals (*MtMTP8*), potassium (*Medtr4g099260*), and zinc (*MtZIP6*)

among the genes in cluster 2 (Supplementary table 2, Supplementary table 3), suggesting mis-

regulation of various nutrient uptake or homeostasis pathways in the mutant (further discussed

below).

Cluster 4 contains genes that are down-regulated in *crn* roots relative to wild-type roots with

little impact of symbiosis status (Fig. 4A). In this group of 391 genes, only the GO term 'plant-

type secondary cell wall biogenesis' was enriched (Supplementary table 3). Intriguingly, all 13

genes associated with this GO term in cluster 4 are annotated as FASCICLIN-LIKE

ARABINOGALACTAN PROTEIN (FLA). FLAs are extracellular glycoproteins involved in a plethora

of functions, including development, growth, and environmental responses (He *et al.*, 2019),

but to our knowledge have not been studied in the context of AM symbiosis or CLAVATA

signaling. However, genes encoding arabinogalactan proteins as well as other cell wall-related

genes have been shown to be down-regulated also in a *A. thaliana crn* mutant (Pallakies and

Simon, 2014). This suggests a conserved role of CRN in cell wall organization, which is critical for

its function in growth and development.

Cluster 5 contains 14 genes that are induced in R108 *R. irregularis*-colonized roots relative to

mock-inoculated roots but show the opposite pattern in *crn* background (down-regulated in *R.*

irregularis-colonized roots relative to mock-inoculated *crn* roots) (Fig. 4A, Supplementary table

3). No significantly enriched GO terms were detected in cluster 5.

N and P homeostasis signaling is perturbed in *crn* roots

Our cluster analysis suggests an up-regulation of multiple mineral nutrient transporters in *crn*,

regardless of root colonization status (see above, Supplementary table 3). This suggests that in

crn mutants, nutrient starvation and/or homeostasis signaling pathways may be mis-regulated.

Because P and N are major regulators of AM symbiosis and their homeostasis is regulated by

CLAVATA signaling pathways in various plant species (Bashyal *et al.*, 2023), we examined genes

associated with P and N homeostasis more closely in our dataset. Interestingly, 16 genes

putatively associated with N homeostasis were regulated by *MtCRN* regardless of root

colonization status, and 11 additional ones were found to be synergistically regulated by *MtCRN*

and AM status of the plant (Fig. 5A-B, Supplementary table 4). Similarly, 8 genes associated with

P homeostasis were mis-regulated in *crn* relative to R108 without additional regulation by root

colonization status, and an additional 3 genes were regulated by *MtCRN* and AM symbiosis (Fig.

5C-D, Supplementary table 4). This data suggests a major perturbance in nutrient homeostasis

signaling in *crn* even in the absence of symbiosis. Because nutrient starving plants initiate more

interactions with beneficial microbes (Nouri *et al.*, 2014), a mis-regulation of nutrient

homeostasis and/or starvation signaling may promote AM fungal root colonization in *crn*.

Expression of early symbiosis signaling genes is mis-regulated in *crn*

Consistent with an increase in AM fungal root colonization, we observed elevated gene

expression of known AM-induced genes in *R. irregularis*-colonized *crn* roots relative to colonized

wild-type controls (see above). This was the case for well-studied genes involved in arbuscule

accommodation and symbiotic nutrient exchange (e.g., *MtPT4*, *MtRAM2*, *MtSTR*), and arbuscule

degeneration (e.g., *MtMYB1*, *MtCP3*, *MtTGL*) (Figure 6A and B, Supplementary table 5).

Expression of these genes did not differ when mock-inoculated *crn* roots were compared to

mock-inoculated R108 roots.

Conversely, expression of many genes involved in pre-symbiotic signaling was perturbed in

mock-inoculated *crn* roots relative to mock-inoculated wildtype roots, suggesting these genes

may be direct or indirect targets of *MtCRN* signaling: We found that the expression of genes

involved in strigolactone biosynthesis and secretion (*MtD27*, *MtCCD7*, *MtCCD8*, *MtPDR1a*) not

only respond to AM fungal root colonization in *crn* and R108, but are also differentially

regulated in *crn* mock-inoculated roots relative to mock-inoculated wild-type controls (Fig. 6A

and B). In addition, *MtCCD8b* and *MtMAX1a*, orthologs of genes involved in strigolactone

biosynthesis in other species, as well as *MtMAX2* and *MtD14*, involved in strigolactone signaling

and perception, respond to AM fungal root colonization and are also differentially regulated in

mock-inoculated *crn* roots relative to controls. Similar trends were observed for the known

transcriptional regulators governing strigolactone biosynthesis *NSP1* and *NSP2.* Because the

production of AM-symbiosis promoting strigolactones is regulated by P and N status of the plant

(Yoneyama *et al.*, 2007, 2012; Liu *et al.*, 2011; Foo *et al.*, 2013; Marro *et al.*, 2022), an up-

regulation of the strigolactone biosynthesis pathway may be a consequence of the apparent

mis-regulation of N and P homeostasis signaling in *crn* roots (Fig. 5A-D).

In addition, expression of several genes involved in the common symbiosis signaling pathway

(CSSP) appear to be directly or indirectly regulated by *MtCRN* (Fig. 6A-B). The CSSP is triggered

by cellular perception of N-acetylglucosamine (GlcNAc)-derived (lipo)chitooligosaccharide

molecules derived from AM fungi or rhizobia (myc-LCOs/COs or nod-LCOs, respectively), and is

required for cellular entry of symbiotic microbes at the onset of symbiotic interactions (Oldroyd,

2013). *DMI2*, which encodes a putative co-receptor perceiving microbial signals, was found to

be up-regulated in mock-inoculated *crn* roots when compared to R108. Likewise, *DMI3*, *IPD3*,

and *VPY*, which regulate cellular signal transduction and symbiosis initiation in response to

microbial LCO perception (MacLean *et al.*, 2017; Lindsay *et al.*, 2022) function downstream of

MtCRN signaling. In addition, *DMI3* and *VPY* expression was also found to be responsive to AM

symbiosis (Fig. 6A). Because the CSSP pathway is not only required for the establishment of AM

but also nodulation symbiosis, and because *crn* was reported to have a hypermycorrhizal

phenotype (this study) and a hypernodulation phenotype in *M. truncatula* (Crook *et al.*, 2016),

we hypothesize that *crn* mutants are primed for symbiotic interactions even in the absence of

microbes, potentially due to an up-regulation of the symbiosis accommodation program as a

result of perturbations in N and P starvation signaling (Li *et al.*, 2022). In line with this

hypothesis, we found that some genes encoding lysin motif (LysM) RLKs involved in the

recognition of nod- and myc-LCOs (e.g., *MtLYK8* and *NFP*, but not *MtLYK3*) are up-regulated in

crn roots even in the absence of a microbe (Fig. 6 C-D, supplementary table 1, supplementary

table 6) (Limpens *et al.*, 2003; Smit *et al.*, 2007; Zhang *et al.*, 2024). Intriguingly, 10 additional

LysM RLKs or receptor-like proteins (LYK, LYR, LYM) that can bind LCOs but have no known role in

nodulation or symbiosis (Buendia *et al.*, 2018) were also up-regulated in *crn* mutants relative to

wildtypes (both in *R. irregularis*- or mock-inoculated treatments), suggesting recognition of

diverse microbial patterns may be altered in *crn*.

***Rhizophagus irregularis* transcriptionally responds to colonizing *crn* roots**

To investigate R*. irregularis* responses to host genotype, we mapped our transcriptome reads to

the *R. irregularis* DAOM197198 v2.0 genome (Yildirir *et al.*, 2022). About 1.2-3.8 % of the reads

mapped to the fungal genome in colonized R108 and *crn* roots (supplementary table 1). We

found 19 genes to be up-, and 10 genes to be down-regulated in *crn* roots relative to wild-type

roots (logFC>1 or <-1, padj<0.05; Fig. 7A, supplementary table 7). Among the functionally

annotated genes down-regulated in *R. irregularis* colonizing *crn* roots relative to wild-type roots

(logFC<-1, padj<0.05), we identified multiple transcripts putatively involved in cellular signaling,

including two protein tyrosine kinases, a C2H2 zinc finger transcription factor, and one SAM-

domain containing protein, which mediate protein-protein interactions and have many roles,

including cellular signaling (Kim, 2003). This suggests that *R. irregularis* modulates signaling in

response to colonizing *crn* roots; however, because AM fungi are functionally not well

understood, we cannot predict which signaling pathways may be impacted.

Among the *R. irregularis* genes significantly up-regulated in *crn* roots relative to wild-type roots

(logFC>1, padj<0.05), we identified multiple transcripts that suggest fungal N metabolism might

be perturbed in *crn* (Fig. 7A). Differentially regulated genes associated with fungal N metabolism

include an argininosuccinate synthase (*ASS*), which is involved in the biosynthesis of arginine, a

critical component of N storage in AM fungi (Cruz *et al.*, 2007). In addition, among the

significantly up-regulated genes we found a transcript annotated as *NmrA (Nitrogen metabolic*

regulation A)-like protein, which in other fungi is part of a protein complex that functions in

transcriptional regulation of N metabolism based on N availability (Andrianopoulos *et al.*, 1998;

Stammers, 2001). Both *ASS* and *NmrA* have been predominantly detected in the extraradical

mycelium of AM fungi where N is taken up (Dumas-Gaudot *et al.*, 2004; Tian *et al.*, 2010). Other

known genes involved in *R. irregularis* N metabolism tended to be up-regulated in *crn* roots

relative to wildtypes (Fig. 7B), although the adjusted p-value did not fall below the stringent

significance criteria described above. This included the ammonium transporters *GintAMT1* and

GintAMT3, but not *GintAMT2*, two glutamine synthetases (*GS1, GS2*) and a N transporter (*NT*)

(Fellbaum *et al.*, 2012) (Fig. 7B).

Interestingly, a plasma membrane H^+-transporting ATPase HA1 with suggested roles in fungal P

sensing (Xie *et al.*, 2022) was also observed among the *R. irregulari*s genes significantly up-

regulated in *crn* roots (Fig. 7A). This observation prompted us to examine other transcripts

known to be involved in fungal in P homeostasis and transport (Kikuchi *et al.*, 2014, 2016; Xie *et*

al., 2022). Although none of the known P homeostasis or transport genes were differentially

regulated in *crn* vs. R108 roots, some displayed trends toward increased expression in *crn*

relative to R108 (Fig. 7C). This included, among others, the $Pi:H^+$ symporter RiPT thought to be

involved in fungal P uptake from the environment (Harrison and van Buuren, 1995; Maldonado-

Mendoza *et al.*, 2001) and the Pi stress marker PHO2 (Austin and Mayer, 2020).

In addition, our *R. irregularis* transcriptome displays signs of increased chitin biosynthesis when

colonizing *crn* roots (Fig. 7A, supplementary table 7). Among the *R. irregularis* genes

significantly up-regulated in *crn* relative to R108 roots (logFC>1, padj<0.05), we identified a gene

encoding a UDP-N-acetylglucosamine diphosphorylase, which catalyzes the formation of

GlcNAc, and a chitin synthase, which catalyzes chitin formation from GlcNAc building blocks (Fig.

7A) (Chen *et al.*, 2022). Notably, GlcNAc are also building blocks of LCO and CO molecules

critical for pre-symbiotic communication with the host. Myc-LCO and CO production is

promoted by host strigolactones (Genre *et al.*, 2013); biosynthesis genes for these

phytohormones are induced in *crn* roots relative to wildtypes (Fig. 6), which could be linked to

the observed up-regulation of GlcNAc in *R. irregularis*. However, while fungal LCO and CO

molecules initiate CSSP signaling in the host upon recognition by plant LysM-type RLKs, they

also have a more general function in regulating fungal growth and development (Rush *et al.*,

2020). In addition, a member of the SGNH hydrolase/GDSL lipase family was up-regulated.

Although the specific function of this gene is unknown, SGNH hydrolases catalyze a variety of

carbon-modifying reactions (Anderson *et al.*, 2022), which may be related to fungal growth and

development within the host plant.

Together, our fungal transcriptome dataset suggests that *crn* roots results stimulate *R.*

irregularis N and P metabolism, growth and development. Notably, it is possible that some of

these changes are an indirect consequence of higher fungal colonization levels observed in the

mutant; however, the lack of differential regulation of *R. irregularis* housekeeping genes

between *crn* and wild-type roots (Supplementary table 8) suggests that the observed gene

expression differences are not solely caused by increased fungal biomass in *crn* roots. While

more research is required, our data supports the hypothesis that the *R. irregularis* differentially

responds to as-yet-unknown signals derived from *crn* and wild-type roots.

Discussion

CLV signaling has been reported to modulate various aspects of plant development and

responses to the environment (Bashyal *et al.*, 2023; Lindsay *et al.*, 2023). Here we describe *M.*

truncatula CRN as a central regulator of meristem branching and root interactions with AM

fungi.

Although legume mutants in the CLV signaling pathway have been well-studied in multiple plant

species in the context of symbiosis regulation, an effect on meristem development has only

been described in soybean and pea (Krusell *et al.*, 2011; Mirzaei *et al.*, 2017). Soybean encodes

two CLV1 copies, one of which controls symbiosis and the other controls meristem development

(Mirzaei *et al.*, 2017). Our work contributes to a better understanding of legume inflorescence

meristem patterning by revealing a role for *MtCRN* and *SUNN* in this process. Although the

molecular mechanisms remain to be determined, several hints in our data and that of others

point toward an intersection of auxin and CLV in *M. truncatula* meristem signaling: Auxin is an

important regulator of meristem identity, and *AtCRN* was found to act upstream of auxin to

493 control floral primordia outgrowth in Arabidopsis (Jones et al., 2021). The *M. truncatula* auxin

494 transport mutant *smooth leaf margin1* has pleiotropic developmental effects, including multiple

495 flowers per inflorescence that are reminiscent of the *crn* flower phenotype reported here (Zhou

496 et al, 2011), suggesting that *MtCRN* may control auxin-mediated outgrowth in *M. truncatula*

497 floral development similar to what was found in Arabidopsis (Jones et al., 2021). In addition,

498 previous studies reported that *SUNN* regulates shoot-to-root auxin transport to control N

499 response (Jin *et al.*, 2012) and that auxin signaling, among other hormonal pathways, is mis-

500 regulated in *sunn* (Karlo *et al.*, 2020, Preprint), further supporting the hypothesis that CLV

501 signaling may act upstream of auxin also in the context of *M. truncatula* meristem maintenance.

502 Like in Arabidopsis, legume meristem signaling may be controlled by multiple, functionally

503 redundant pathways and/or buffering. Consequently, phenotypes may only be apparent in

504 certain mutant alleles or growth conditions. Because these buffering mechanisms appear

505 distinct among plant species, this may explain why inflorescence meristem branching is

506 primarily affected in Medicago *crn* mutants, resembling the phenotype in other legumes (Krusell

507 *et al.*, 2011; Mirzaei *et al.*, 2017), but inflorescence meristem size is affected in maize and

508 Arabidopsis *crn* mutants (Rodriguez-Leal et al., 2019, Hazak and Hardtke, 2016, Je et al., 2016).

510 CLV signaling has been previously described to be critical for autoregulation of AM symbiosis

511 (AOM), but our knowledge on the molecular mechanisms remains limited. We previously

512 reported the up-regulation of three *MtCLE* genes in *R. irregularis*-inoculated roots, one of which

513 was functionally characterized as a member of the AOM pathway (*MtCLE53*) (Müller *et al.*,

514 2019; Karlo *et al.*, 2020, Preprint). *MtCLE53* requires *SUNN* to regulate AM symbiosis (Crook *et*

515 *al.*, 2016; Müller *et al.*, 2019; Karlo *et al.*, 2020, Preprint). Because MtCRN and SUNN can

516 interact and are required to perceive nodulation-associated CLE peptides during AON (Crook *et*

517 *al.*, 2016; Nowak *et al.*, 2019), we hypothesized that MtCRN may also be required for MtCLE53

518 signaling together with SUNN. However, the data presented here suggests that the negative

519 effect of *MtCLE53* overexpression on AM symbiosis is only partially dependent on *MtCRN*. One

520 possible explanation for our observation is that MtCLE53 can signal through MtCRN-SUNN but

521 also parallel, MtCRN-independent pathway(s). Such parallel pathways and compensation

mechanisms are common in CLAVATA signaling modules (Pallakies and Simon, 2014; Nimchuk,

2017; Je *et al.*, 2018; Rodriguez-Leal *et al.*, 2019; Selby and Jones, 2023). This also suggests that

additional, *MtCLE53*-independent pathways contribute to the elevated AM fungal root

colonization observed in *crn*. MtCRN can interact not only with CLV1 (SUNN) but also other

receptors, including MtCLV2 (Crook *et al.*, 2016); therefore it is possible that CRN is involved in

the signal transduction from multiple input signals. Multiple CLV signaling pathways also appear

to govern AM symbiosis in tomato (*Solanum lycopersicum*); for example, SlCLV2 was reported to

regulate AM symbiosis through two distinct local and systemic signaling pathways (Wang *et al.*,

2021; Wulf *et al.*, 2024). Previously published *MtCRN* promoter-reporter analyses revealed that

MtCRN is expressed in all root cell types with higher expression in the vascular tissue (Crook *et*

al., 2016). *SUNN*, *MtCLE53*, and other symbiosis autoregulation regulators such as *MtRDN1* and

MtDAR, are also expressed in the vasculature (Schnabel *et al.*, 2012, 2024; Müller *et al.*, 2019;

Serrano *et al.*, 2024). Therefore, it is possible that a MtCLE53-SUNN-MtCRN autoregulatory

module regulates AM symbiosis through long-distance signaling in the vascular tissue, whereas

MtCRN expressed in other root cell types may have a distinct signaling function to regulate AM

symbiosis in concert with other partners.

Our transcriptomics experiment revealed that *M. truncatula crn* mutants display signatures of

abiotic stress, most notably nutrient starvation as suggested by a transcriptional up-regulation

of N, P and other nutrient transporters. It is possible that perpetually heightened nutrient

starvation signals promote AM symbiosis in *crn* independently of AOM. Nutrient-regulated CLE

peptides have been identified and CLV receptors have been implicated in various plant

responses to nutrient availability (including P and N) (Bashyal *et al.*, 2023); thus, it is tempting to

speculate that signaling of nutrient-induced CLE peptides may be perturbed in *crn*. Intriguingly,

R. irregularis also displays signs of disrupted N and P metabolism as indicated by an up-

regulation of *NmrA and ASS*. The proteins encoded by these genes are predicted to function in

the extracellular mycelium of AM fungi to mediate N utilization and incorporation into arginine

(Andrianopoulos *et al.*, 1998; Cruz *et al.*, 2007; Fellbaum *et al.*, 2012). Arginine plays a key role

in N translocation from the extra- to the intraradical mycelium in AM fungi (Cruz *et al.*, 2007;

Fellbaum *et al.*, 2012). In addition, *R. irregularis HA1* expression was increased when colonizing

crn roots; *RiHA1* was previously implicated in P signaling and uptake by the extraradical

mycelium, where it powers Pi:H$^+$ symporters to promote phosphate uptake (Ezawa and Saito,

2018; Xie *et al.*, 2022). Together with our host transcriptomics data, this finding suggests that

crn roots may represent a greater sink for N and P than wild-types and consequently stimulate N

and P uptake in the colonizing *R. irregularis* symbiont (Cruz *et al.*, 2007). Further research is

required to better understand the impact of host genotype and host CLV signaling on symbiont

metabolism.

In the plant, N and P starvation results in the upregulation of genes involved in the biosynthesis

of strigolactones, which in turn act as positive regulators of AM symbiosis (Akiyama *et al.*, 2005;

Yoneyama *et al.*, 2007; Liu *et al.*, 2011; Foo *et al.*, 2013, 2014; Marro *et al.*, 2022). Intriguingly,

our data reveal that many of these genes are upregulated in *crn* relative to wildtype controls,

which may be a direct consequence of the apparent perturbations in N and P signaling in *crn*.

Strigolactone biosynthesis is also the target of AOM, and the elevated expression of

strigolactone biosynthesis genes may also be caused by an at least partial disruption of AOM. In

either case, upregulation of genes involved in the symbiosis-promoting strigolactones in *crn* can

explain the elevated root colonization levels in the mutant. Together, our functional and

transcriptomic data suggests that MtCRN may act upstream of strigolactone biosynthesis

regulation in response to MtCLE53 and/or N and P signaling. Intriguingly, up-regulation of

strigolactone biosynthesis genes have not been observed in CLV1 mutants in *M. truncatula*

(*sunn*), pea (*nark*) or tomato (*fab*) (Foo *et al.*, 2014; Karlo *et al.*, 2020, Preprint; Wang *et al.*,

2021), suggesting that CRN (but not SUNN/FAB) may modulate cross-talk between N and P

homeostasis, MtCLE53-mediated autoregulation, and AM fungal root colonization.

Apart from genes involved in nutrient homeostasis and strigolactone biosynthesis, our

transcriptome analysis also suggests that *crn* roots display an upregulation of genes associated

with the CSSP. The CSSP is activated upon perception of diffusible LCO or CO signals derived

from AM fungi or rhizobia and is required for cellular entry of the microbial symbionts. Because

crn has previously been described as a regulator of nodulation symbiosis, an effect of *crn* on

CSSP gene expression may contribute not only to the hypermycorrhizal phenotype described

here, but also to the hypernodulation phenotype described previously (Crook *et al.*, 2016). It

was reported previously in rice that transcription of certain CSSP genes (including orthologs of

MtVPY, DMI2, DMI3, IPD3, which we found here to be regulated by *crn*) is regulated by the plant

phosphate starvation response (Das *et al.*, 2022). In addition, expression of LysM-type RLKs and

LysM receptor proteins as well as plant responses to LCO/CO perception depend on N and P

status of the plant and are greatly increased upon nutrient starvation (Li *et al.*, 2022). We

therefore hypothesize that the perturbed N and P signaling in *crn* promotes AM symbiosis by

increasing the expression of genes involved in early symbiosis signaling (strigolactone

biosynthesis genes, CSSP genes, LysM-RLKs), which facilitates colonization by AM fungi upon

contact. It has been established that nutrient starvation determines the outcome of plant-

microbe interactions by fine-tuning plant immune responses, potentially in order to facilitate

symbiotic interactions (Paries and Gutjahr, 2023). Consistent with this model, our dataset

suggests attenuated oxidative stress in *crn* relative to wildtypes; oxidative stress is a hallmark of

responses to abiotic factors but also plant immunity. However, we also detected up-regulation

of plant defense genes, suggesting *crn* displays additional signatures of enhanced response to

biotic stress. Complex biotic stress responses have been described before for *crn*: For example,

A. thaliana crn mutants display increased resistance to *Heterodera schachtii* cyst nematodes as

well as certain bacterial and oomycete pathogens (*Ralstonia solanacearum* and

Hyaloperonospora arabidopsidis, respectively), while being more susceptible to *Botrytis*

necrotrophic fungi or *Pseudomonas syringae* bacterial infection (Replogle *et al.*, 2011;

Hanemian *et al.*, 2016). Together with our data, this suggests that coordination or cross-talk of

multiple combined abiotic and/or biotic stress responses, rather than overall stress signaling,

may be impaired in *crn*.

In summary, our findings suggest that *MtCRN* plays multiple, likely independent, roles in plant

development, nutrient homeostasis, and plant interactions with the environment. Based on

data from other plant species, where different combinations of CLV1-type receptors, CLV2, and

CRN are responsible for the perception of diverse CLE signals in a context- and cell-type-specific

manner (Whitewoods, 2021), we propose that *MtCRN* acts downstream of diverse CLE peptides

in the context of meristem development, nutrient homeostasis, and plant-microbe interactions.

More research is needed to identify additional context-dependent CLE signals and receptors

that provide specificity to the various signaling pathways *MtCRN* is involved in.

Materials and methods

Plant materials and growth conditions

Homozygous *M. truncatula crn* (NF3436) and *sunn-5* (NF2262) mutants, both in R108

background, were a gift from Dr. Julia Frugoli (Clemson University, SC, USA) and were previously

described (Crook *et al.*, 2016; Nowak *et al.*, 2019; Thomas and Frugoli, 2024).

For AM inoculation experiments, seeds were extracted from seed pods and scarified using

sandpaper or by incubating the seeds in concentrated sulfuric acid for 7 minutes. Seeds were

washed 5 times in autoclaved distilled H2O and surface-sterilized with 10% Bleach for 7

minutes. Seeds were rinsed again 5 times in autoclaved distilled H2O and incubated at 4°C for

48 hours. Imbibed seeds were germinated on germination paper (Anchor paper) for 3-5 days at

room temperature before transplanting into substrate. Plants were grown as previously

described (Chaulagain *et al.*, 2023*a*). In brief, germinated seedlings were transplanted into SC10

cone-tainers (Steuwe and Sons) filled with washed and autoclaved substrate mix consisting of

play sand and fine vermiculite (1:1 ratio). For inoculated samples, 250 *Rhizophagus irregularis*

DAOM197198 spores (Premier Tech, Canada) were placed onto a sand layer 5 cm below the

surface of the substrate in each cone-tainer. For mock-inoculated samples, an equal amount of

distilled water was added onto the sand layer. Plants were grown in a growth chamber under a

16 hour light (24°C)/8 hour dark (22°C) regime at 40% relative humidity. For standard

experiments, all cones were fertilized twice per week with 20 ml half-strength Hoagland's

fertilizer supplied with 20μM potassium phosphate. Unless otherwise noted, all plants were

harvested 5-6 weeks after inoculation. To enrich for *R. irregularis* colonization, we sampled 5

cm-long root sections centered around the inoculated region. For the transcriptomics

experiment, whole root systems were harvested.

Transient root transformation

Transient root transformations of *M. truncatula* R108 and *crn* plants with *35S:GUS* and

35S:MtCLE53 constructs were performed as described previously (Boisson-Dernier *et al.*, 2001;

Müller *et al.*, 2019). Chimeric plants were inoculated with *R. irregularis* as described above and

harvested 6.5 weeks after planting into cone-tainers.

Root system phenotyping

Seeds were extracted from seed pods and sterilized as described above. Seeds were then

distributed on the bottom of a glass petri dish using water surface tension (Floss *et al.*, 2013).

Glass petri plates were placed upside down in a 28°C incubator and seedlings were allowed to

germinate overnight. With this method, we were able to obtain seedlings with straight roots.

Seedlings were then transferred to 12x12 cm square petri dishes filled with modified Fahraeus

(F) medium supplied with 4 g/L Gelzan CM agar (Sigma-Aldrich) (Floss *et al.*, 2013). Plates were

placed at a 45° angle in a growth chamber under a 16 hour light (24°C)/8 hour dark (22°C)

regime. The bottom 2/3 of the plates was wrapped in aluminum foil to protect the roots from

light. Pictures of seedlings were taken every 3 days for 12 days starting with the day of transfer

to agar plates (day 0). Root system parameters (primary root length, lateral root length, lateral

root number, lateral root density) were analyzed using the SmartRoot package in Fiji (Lobet *et*

al., 2011; Schindelin *et al.*, 2012) for each timepoint.

AM fungus staining, microscopy, and quantification of AM fungal colonization

WGA-Alexafluor488 staining was performed by incubating the roots in 20% KOH for 3-4 days at

65°C. Roots were rinsed 3-5 times with distilled water and incubated in 1xPBS overnight.

Cleared and washed roots were stained with 0.2 mg/ml WGA-Alexafluor488 (Invitrogen) in

1xPBS for at least 6 hours. Stained roots were visualized using a Leica Thunder Imager or a Leica

M205 FA stereomicroscope. Root length colonization was assessed using the grid-line method

(McGonigle *et al.*, 1990). Arbuscule and vesicle density was quantified as previously described

(Müller *et al.*, 2020; Chaulagain *et al.*, 2023*a*). In brief, a minimum of 5 z-stacks each of

randomly selected colonized root sections from at least 3 independent root systems were

imaged that encompassed all arbuscules in that area. Roots were positioned so that the

hyphopodium was at the center of the image to ensure infections were at a comparable

developmental stage. Imaged arbuscules were counted using the Fiji Image Analysis Package

(Schindelin *et al.*, 2012), which was also used to accurately measure the imaged root length.

Arbuscule density (arbuscules/mm) was calculated by dividing the number of arbuscules by the

root length. Confocal images were obtained using a Leica SP5 confocal microscope.

Transcriptomics

For each genotype (*crn* and R108), 6-week-old mock- (n=4 per genotype) and *R. irregularis*-

colonized root systems (n=3 per genotype) were harvested as described above. WGA-

Alexafluor488 staining was performed on a subset of the samples to confirm colonization. RNA

was extracted from the remaining samples as previously described (Javot *et al.*, 2011). Library

preparation and transcriptome sequencing was performed by Novogene Co, Ltd. Samples were

enriched for mRNA using poly(A) capture. Paired-end sequencing (PE150) was performed on the

Illumina Novaseq X platform. Sequencing raw data is deposited in the NCBI Sequence Read

Archive (SRA) (BioProject PRJNA1140176). Filtered reads were mapped against the *M.*

truncatula Mt4.0v1 genome (downloaded from Phytozome 13) and *R. irregularis* DAOM

RhiirDAOM197198 v2.0 (downloaded from the JGI Mycocosm database) genome using HiSAT2

(Tang *et al.*, 2014; Kim *et al.*, 2019; Yildirir *et al.*, 2022). Normalized read counts were used to

calculate differential gene expression with DESeq2 (Love *et al.*, 2014). For each pairwise

comparison, log2-transformed fold changes were obtained for each mapped gene. Differential

gene expression was considered significant if the p-value adjusted for multiple comparisons

using Benjamini-Hochberg method (padj) was below 0.05. Heatmaps were constructed using

FPKM-normalized read counts; rows were clustered using Pearson correlation, columns using

Spearman correlation. To identify gene groups with similar expression patterns across samples,

hierarchical clustering was performed on FPKM-normalized read counts averaged for each

genotype/treatment. The number of clusters (k=6) was determined using the sum of squared

error method. GO term analysis for biological process was performed on

https://geneontology.org/; visualization of the obtained GO term enrichment results was

performed in R.

Scanning electron microscopy

For inflorescence phenotyping, plants were grown in 6-inch azalea pots in soil (Promix) under a

16 hour light/8 hour dark regime at 20-22°C and watered with distilled water as needed. Axils

from 8-week-old plants were dissected with 30 gauge insulin syringe needles to reveal

inflorescence meristems under a stereoscope. Meristems and the surrounding tissue were

mounted on stubs using double-sided tape and imaged in a Jeol JCM-7000 benchtop scanning

electron microscope (SEM) with the following conditions: high vacuum, 15 kV, high probe

current, analysis mode. Meristems from SEM images were categorized based on (Benlloch *et al.*,

2003) and measured using Fiji.

Statistical analyses

Statistical significance between two experimental groups was assessed using two-sided

Student's t-test. In graphs, statistical significances between two groups are represented as

follows: ***$p<0.001$; **$p<0.01$; *$p<0.05$. Statistical significance between multiple experimental

groups was calculated with ANOVA, followed by Tukey's HSD test for multiple comparisons.

Pairwise comparisons were considered statistically significant if $p<0.05$. Significant differences

are denoted with different letters. The % reduction of root length colonization shown in Fig. 3

was calculated separately for each genotype by the following formula: ((sample $RLC_{35S:CLE53}$ –

average $RLC_{35S:GUS}$)/average $RLC_{35S:GUS}$)*100. The % reduction values for each genotype were

then compared to each other using student's t-test. See paragraphs 'transcriptomics' for

additional details on the statistics used to analyze RNA sequencing data. Graphs show data from

individual experiments. All experiments were repeated at least twice with similar results.

Acknowledgements

We thank Julia Frugoli (Clemson University, SC) for kindly gifting us the *crn* and *sunn-5* mutant

lines. We also thank Dr. Yerisf Torres Ascurra for helpful discussions and critically reading the

manuscript.

Author contributions

729 L.M.M. conceived the project. J.O., E.X.L, M.K., S.B., H.E., C.K.G. and L.M.M. performed all

730 experiments except for the inflorescence meristem analyses, which were performed by P.L.L.

731 L.M.M. and D.J. acquired funding for the project. L.M.M. wrote the manuscript with input from

732 all co-authors.

Conflict of interest

735 None to declare.

Funding

738 This work was partially funded by USDA-NIFA award 2022-67013-42820 to L.M.M. and an

739 American Society of Plant Biologists (ASPB) Summer Undergraduate Research Fellowships

740 (SURF) fellowship to E.X.L. P.L.L. and D.J. were funded by USDA-NIFA award 2020-67013-30909.

Data availability

744 Raw data from the transcriptome sequencing has been deposited on the NCBI SRA database

745 (Bioproject ID: PRJNA1140176).

Figures and figure legends

Figure 1: *M. truncatula CRN* **impacts inflorescence development.** A) Close-up of R108 wild-type flower. 100% of inflorescences produced a solitary flower in our growth conditions. B) Dissected wild-type flower. Std, standard petal; fused keel and 2 wing petals; c+s, carpel and stamen. C) Close-up of inflorescence produced by *crn*. Note the additional undeveloped bud (arrowhead). In our growth conditions, 91% of *crn* produced two flowers per inflorescence, the remaining 9% produced solitary flowers similar to the one shown in (A). D) Dissected *crn* flower. Flower organ number and morphology appeared similar to the R108 wildtype shown in (B). (A-D) Scale bars: 0.5 mm. E) Scanning electron microscopy images of R108 and *crn* inflorescence meristems. In *M. truncatula*, the primary inflorescence meristem (I_1) produces a secondary inflorescence meristem (I_2) and leaf primordium (L). The secondary meristem will in turn produce a floral meristem (F), which will differentiate floral organ primordia. Top panels show stage five floral meristems, in which all floral organ primordia have been established. Floral organ primordia size and number are unaltered in *crn*, but there is a second stage 5 floral meristem, indicated with an asterisk. The bottom panel shows a range of floral meristem stages in R108 and *crn*. In *crn*, two I_2 meristems are present at the same time, indicated by an asterisk. C, carpel primordium,

k, keel petal, s_{ab}, abaxial sepal, st_p, non-fused vexillary stamen, st_s, antesepal stamen, v_x, vexillum, Br, bract. Scale bar: 100 μm. F) Width of R108 and *crn* meristems in different floral meristem stages. No effect of the *crn* mutation was observed on meristem width in any developmental stage assessed. I_2, secondary inflorescence meristem, F, floral meristem, in which sepal primordia begin to form, stage 4, in which common primordia that will form stamens and petals are formed, stage 5, in which all floral organ primordia have been established. G) Number of stamen and petal floral organ primordia in stage 5 meristems. N.s., no significant difference between R108 and *crn* (Student's t-test).

Figure 2: *M. truncatula crn* mutants display elevated levels of colonization by the AM fungus *R. irregularis*. A) *crn* mutants diplay higher levels of overall root length colonization relative to R108 wild-type controls. B) *crn* mutants show higher arbuscule density (number of arbuscules per mm colonized root length) relative to R108 wild-type controls. C) Vesicle density (vesicles per mm colonized root length) does not differ between R108 and *crn*. D) Representative image of an *R. irregularis*-colonized R108 wildtype root. *R. irregularis* was stained with WGA-Alexafluor488 (green). Dashed line: outline of the root. E) Representative image of an *R. irregularis* (green) colonized root section in *crn*. Dashed line: outline of the root. A-C) Bar plots show averages, error bars the standard deviation, points represent individual measurements (i.e., n=4 individual root systems per genotype in A) and n=19 (R108) and n=25 (*crn*) imaged root sections in B) and C), derived from the same root systems as shown in A). Graphs display data from one experiment; this experiment was repeated twice with similar results. ** p<0.01 (t-test). Scale bar in D), E): 100 μm.

Figure 3: AM fungal root length colonization of *crn* mutants partially depend on *MtCLE53*.

Graph depicts average root length colonization (%) of R108 and *crn* plants transformed with the *35S::GUS (GUS)* or the *35S::MtCLE53 (CLE53)* construct. *MtCLE53* over-expression reduces *R. irregularis* root colonization in R108 and *crn*; however, this suppressive effect is stronger in R108 than in *crn*. For each genotype, the % of colonization reduction upon *MtCLE53* over-expression is shown. The bar graph shows average values, error bars denote standard deviation. Black dots indicate individual datapoints (biological replicates; n=3-5 replicates per sample). This graph displays data from one experiment; the experiment was repeated twice with similar results. Different letters denote significant differences (p<0.05) after ANOVA (p=2.12*10^{-5}) followed by Tukey's HSD test for pairwise comparisons.

Figure 4: Hierarchical cluster analysis of differentially expressed genes in *R. irregularis*-colonized and mock-inoculated *crn* and R108 wild-type roots. A) 1655 genes that are differentially expressed between *R. irregularis*-colonized *crn* and R108 wild-type roots (log2 fold-change >1 or <-1, padj<0.05) were selected for cluster analysis to identify gene groups with similar expression patterns. Genes were clustered using Pearson correlation. Heatmap shows averages of n=4 root samples each for *crn* and R108 mock-inoculated, and n=3 samples each for *crn* and R108 *R. irregularis*-inoculated (*R.i.*). Gene lists for each cluster can be found in table S2. Line graphs display normalized relative expression changes across samples for each individual gene in a cluster (grey) and average changes in color. B) Biological functions GO terms enriched in cluster 1, C) Biological functions GO terms enriched in cluster 2, D) Biological functions GO

terms enriched in cluster 6. Note: Cluster 4 has one significantly enriched GO term "plant-type secondary cell wall biogenesis", which is not shown in this figure. Supplementary tables 2 and 3 contains a list of genes for each cluster and the associated GO terms, respectively. DXP, 1-deoxy-d-xylulose 5-phosphate. FDR, false discovery rate.

Figure 5: Differential regulation of N and P homeostasis genes in *crn* mutants. A) Heatmap for genes annotated as *NITRATE TRANSPORTER (NRT), NITRATE AND PEPTIDE TRANSPORTER FAMILY (NPF), NITRATE ASSIMILATION RELATED (NAR), AMMONIUM TRANSPORTER (AMT),* and *NITRATE REDUCTASE (NR)* with putative roles in *M. truncatula* N metabolism. B) Venn diagram showing the number of putative N homeostasis genes responding to AM symbiosis, *crn*, or both (padj<0.05). C) Venn diagram showing the number of putative P homeostasis genes responding to AM symbiosis, *crn*, or both (padj<0.05). D) Heatmap for genes annotated as *PHOSPHATE TRANSPORTER (PT), PHOSPHATE1 (PHO), PHOSPHATE STARVATION RESPONSE (PHR* and *PHR-*

LIKE, PHRL), or *SYG1/PHO81/XPR1 family (SPX)* with putative roles in P homeostasis and transport. A and D) Heatmaps show averages of n=4 root samples each for *crn* and R108 mock-inoculated, and n=3 samples each for *crn* and R108 *R. irregularis*-inoculated (*R.i.*). AM-regulated genes are highlighted in grey, genes differentially regulated in *crn* vs. R108 with no effect of colonization status in light green. Genes responsive to AM and *crn* are highlighted in dark green. B and C) 'AM-regulated' is defined as genes with differential regulation (padj<0.05) between *R.irregularis*-colonized and mock-inoculated R108 and/or *crn* roots, respectively. '*crn*-regulated' is defined as genes with differential regulation (padj<0.05) in mock-inoculated *crn* relative to mock-inoculated R108 roots. Transcript data used to produce these heatmaps can be found in Supplementary table 4.

Figure 6: Differential regulation of AM symbiosis regulators in *crn* mutants. A) Heatmaps for known genes involved in pre-symbiotic signaling, the common symbiosis signaling pathway (CSSP) shared between AM and nodulation symbiosis, arbuscule accommodation, and arbuscule degradation. B) Venn diagram showing the number of AM regulatory genes from panel A) responding to AM symbiosis, *crn*, or both. C) Venn diagram showing the number of LysM-RLK or LysM receptor protein-encoding genes (LYK, LYR, LYM) responding to AM symbiosis, *crn*, or both. D) Heatmap for LysM-RLK genes in mock- and *R. irregularis*-inoculated wildtype (R108) and *crn* roots. A and D) Heatmaps show averages of n=4 root samples each for *crn* and R108 mock-

inoculated, and n=3 samples each for *crn* and R108 *R. irregularis*-inoculated (*R.i.*). AM-regulated genes are highlighted in grey, genes differentially regulated in *crn* vs. R108 with no effect of colonization status in light green. Genes responsive to AM and *crn* are highlighted in dark green. B and C) 'AM-regulated' is defined as genes with differential regulation (padj<0.05) between *R. irregularis*-colonized and mock-inoculated R108 and/or *crn* roots, respectively. '*crn*-regulated' is defined as genes with differential regulation (padj<0.05) in mock-inoculated *crn* relative to mock-inoculated R108 roots. Transcript data used to produce these heatmaps can be found in supplementary tables 5 and 6.

Figure 7: Host genotype-dependent regulation of *R. irregularis* gene expression. A)

Differentially expressed *R. irregularis* genes (log2 fold change >1 or <-1, padj<0.05) in *crn* and

R108 roots. B) Heatmap of genes involved in fungal N metabolism. C) Heatmap of genes involved in fungal P metabolism. Significance: ***adjusted p-value (padj)<0.05, **padj<0.01, *padj<0.05; if no symbol is shown, fungal gene expression does not significantly differ between host genotypes. Transcript data used to produce these heatmaps can be found in supplementary table 7. Data for the same Argininosuccinate synthase (*ASS*) gene is shown in panels A) and B); data for the same H$^+$-transporting ATPase (*HA1*) gene is shown in panels A) and C).

References

Abbott LK, Robson AD, de Boer G. 1984. THE EFFECT OF PHOSPHORUS ON THE FORMATION OF HYPHAE IN SOIL BY THEVESICULAR-ARBUSCULAR MYCORRHIZAL FUNGUS, *GLOMUS FASCICULATUM*. The new phytologist **97**, 437–446.

Akiyama K, Matsuzaki K-I, Hayashi H. 2005. Plant sesquiterpenes induce hyphal branching in arbuscular mycorrhizal fungi. Nature **435**, 824–827.

Anderson AC, Stangherlin S, Pimentel KN, Weadge JT, Clarke AJ. 2022. The SGNH hydrolase family: a template for carbohydrate diversity. Glycobiology **32**, 826.

Andrianopoulos A, Kourambas S, Sharp JA, Davis MA, Hynes MJ. 1998. Characterization of the Aspergillus nidulans nmrA gene involved in nitrogen metabolite repression. Journal of bacteriology **180**, 1973–1977.

Austin S, Mayer A. 2020. Phosphate homeostasis – A vital metabolic equilibrium maintained through the INPHORS signaling pathway. Frontiers in microbiology **11**.

Bashyal S, Gautam CK, Müller LM. 2023. CLAVATA signaling in plant–environment interactions. Plant physiology doi: 10.1093/plphys/kiad591.

Benlloch R, Navarro C, Beltrán J, Cañas LA. 2003. Floral development of the model legume Medicago truncatula: ontogeny studies as a tool to better characterize homeotic mutations. Sexual plant reproduction **15**, 231–241.

Bleckmann A, Weidtkamp-Peters S, Seidel CAM, Simon R. 2009. Stem cell signaling in Arabidopsis requires CRN to localize CLV2 to the plasma membrane. Plant physiology **152**, 166–176.

Boisson-Dernier A, Chabaud M, Garcia F, Bécard G, Rosenberg C, Barker DG. 2001. Agrobacterium rhizogenes-transformed roots of Medicago truncatula for the study of nitrogen-

fixing and endomycorrhizal symbiotic associations. Molecular plant-microbe interactions: MPMI **14**, 695–700.

Buendia L, Girardin A, Wang T, Cottret L, Lefebvre B. 2018. LysM Receptor-Like Kinase and LysM Receptor-Like Protein Families: An Update on Phylogeny and Functional Characterization. Frontiers in plant science **9**, 539.

Chaulagain D, Schnabel E, Crook A, Bashyal S, Müller LM, Frugoli J. 2023*a*. A mutation in mediator subunit MED16A suppresses nodulation and increases arbuscule density in Medicago truncatula. Journal of plant growth regulation doi: 10.1007/s00344-023-10993-2.

Chaulagain D, Schnabel E, Lin EX, Garcia RR, Noorai RE, Müller LM, Frugoli JA. 2023*b*. *TML1*AND*TML2*SYNERGISTICALLY REGULATE NODULATION BUT NOT ARBUSCULAR MYCORRHIZA IN*MEDICAGO TRUNCATULA*. bioRxiv.

Chen W, Cao P, Liu Y, *et al.* 2022. Structural basis for directional chitin biosynthesis. Nature **610**, 402–408.

Chen Y, Miao Y, Huang L, Li J, Sun H, Zhao Y, Yang J, Zhou W. 2014. Antioxidant activities of saponins extracted from Radix Trichosanthis: an in vivo and in vitro evaluation. BMC complementary and alternative medicine **14**, 86.

Clark SE, Williams RW, Meyerowitz EM. 1997. The CLAVATA1 gene encodes a putative receptor kinase that controls shoot and floral meristem size in Arabidopsis. Cell **89**, 575–585.

Crook AD, Schnabel EL, Frugoli JA. 2016. The systemic nodule number regulation kinase SUNN in Medicago truncatula interacts with MtCLV2 and MtCRN. The Plant journal: for cell and molecular biology **88**, 108–119.

Cruz C, Egsgaard H, Trujillo C, Ambus P, Requena N, Martins-Loução MA, Jakobsen I. 2007. Enzymatic evidence for the key role of arginine in nitrogen translocation by arbuscular mycorrhizal fungi. Plant physiology **144**, 782–792.

Das D, Paries M, Hobecker K, Gigl M, Dawid C, Lam H-M, Zhang J, Chen M, Gutjahr C. 2022. PHOSPHATE STARVATION RESPONSE transcription factors enable arbuscular mycorrhiza symbiosis. Nature communications **13**, 1–12.

Dumas-Gaudot E, Valot B, Bestel-Corre G, Recorbet G, St-Arnaud M, Fontaine B, Dieu M, Raes M, Saravanan RS, Gianinazzi S. 2004. Proteomics as a way to identify extra-radicular fungal proteins from Glomus intraradicesâ RiT-DNA carrot root mycorrhizas. FEMS microbiology ecology **48**, 401–411.

Ezawa T, Saito K. 2018. How do arbuscular mycorrhizal fungi handle phosphate? New insight into fine-tuning of phosphate metabolism. The new phytologist **220**, 1116–1121.

Fellbaum CR, Gachomo EW, Beesetty Y, Choudhari S, Strahan GD, Pfeffer PE, Kiers ET, Bücking H. 2012. Carbon availability triggers fungal nitrogen uptake and transport in arbuscular mycorrhizal symbiosis. Proceedings of the National Academy of Sciences of the United States of America **109**, 2666–2671.

Fester T, Hause G. 2005. Accumulation of reactive oxygen species in arbuscular mycorrhizal roots. Mycorrhiza **15**, 373–379.

Floss DS, Schmitz AM, Starker CG, Gantt JS, Harrison MJ. 2013. Gene Silencing in Medicago truncatula Roots Using RNAi. Methods in molecular biology (Clifton, N.J.). Legume Genomics. Totowa, NJ: Humana Press, 163–177.

Foo E, Ferguson BJ, Reid JB. 2014. The potential roles of strigolactones and brassinosteroids in the autoregulation of nodulation pathway. Annals of botany **113**, 1037–1045.

Foo E, Yoneyama K, Hugill CJ, Quittenden LJ, Reid JB. 2013. Strigolactones and the regulation of pea symbioses in response to nitrate and phosphate deficiency. Molecular plant **6**, 76–87.

Fujimatsu T, Endo K, Yazaki K, Sugiyama A. 2020. Secretion dynamics of soyasaponins in soybean roots and effects to modify the bacterial composition. Plant direct **4**.

Genre A, Chabaud M, Balzergue C, *et al.* 2013. Short-chain chitin oligomers from arbuscular mycorrhizal fungi trigger nuclear Ca2+ spiking in Medicago truncatula roots and their production is enhanced by strigolactone. The New phytologist **198**, 190–202.

Gullner G, Komives T, Király L, Schröder P. 2018. Glutathione S-Transferase Enzymes in Plant-Pathogen Interactions. Frontiers in plant science **9**.

Guo Y, Han L, Hymes M, Denver R, Clark SE. 2010. CLAVATA2 forms a distinct CLE-binding receptor complex regulating Arabidopsis stem cell specification. The Plant journal: for cell and molecular biology **63**, 889–900.

Hanemian M, Barlet X, Sorin C, *et al.* 2016. Arabidopsis CLAVATA1 and CLAVATA2 receptors contribute to Ralstonia solanacearum pathogenicity through a miR169-dependent pathway. The new phytologist **211**, 502–515.

Harrison MJ, van Buuren ML. 1995. A phosphate transporter from the mycorrhizal fungus Glomus versiforme. Nature **378**, 626–629.

Hazak O, Brandt B, Cattaneo P, Santiago J, Rodriguez-Villalon A, Hothorn M, Hardtke CS. 2017. Perception of root-active CLE peptides requires CORYNE function in the phloem vasculature. EMBO reports **18**, 1367–1381.

Hazak O, Hardtke CS. 2016. CLAVATA 1-type receptors in plant development. Journal of experimental botany **67**, 4827–4833.

He J, Zhao H, Cheng Z, Ke Y, Liu J, Ma H. 2019. Evolution analysis of the fasciclin-like arabinogalactan proteins in plants shows variable fasciclin-AGP domain constitutions. International journal of molecular sciences **20**, 1945.

Huang H, Ullah F, Zhou D-X, Yi M, Zhao Y. 2019. Mechanisms of ROS regulation of plant development and stress responses. Frontiers in plant science **10**.

Javot H, Penmetsa RV, Breuillin F, Bhattarai KK, Noar RD, Gomez SK, Zhang Q, Cook DR, Harrison MJ. 2011. Medicago truncatula mtpt4 mutants reveal a role for nitrogen in the regulation of arbuscule degeneration in arbuscular mycorrhizal symbiosis. The Plant journal: for cell and molecular biology **68**, 954–965.

Je BI, Xu F, Wu Q, Liu L, Meeley R, Gallagher JP, Corcilius L, Payne RJ, Bartlett ME, Jackson D. 2018. The CLAVATA receptor FASCIATED EAR2 responds to distinct CLE peptides by signaling through two downstream effectors. eLife **7**, e35673.

Jin J, Watt M, Mathesius U. 2012. The autoregulation gene SUNN mediates changes in root organ formation in response to nitrogen through alteration of shoot-to-root auxin transport. Plant physiology **159**, 489–500.

Jones DS, John A, VanDerMolen KR, Nimchuk ZL. 2021. CLAVATA signaling ensures reproductive development in plants across thermal environments. Current biology: CB **31**, 220-227.e5.

Karlo M, Boschiero C, Landerslev KG, Blanco GS, Wen J, Mysore KS, Dai X, Zhao PX, de Bang TC. 2020. The CLE53–SUNN genetic pathway negatively regulates arbuscular mycorrhiza root colonization in Medicago truncatula.

Kikuchi Y, Hijikata N, Ohtomo R, Handa Y, Kawaguchi M, Saito K, Masuta C, Ezawa T. 2016. Aquaporin-mediated long-distance polyphosphate translocation directed towards the host in arbuscular mycorrhizal symbiosis: application of virus-induced gene silencing. The new phytologist **211**, 1202–1208.

Kikuchi Y, Hijikata N, Yokoyama K, Ohtomo R, Handa Y, Kawaguchi M, Saito K, Ezawa T. 2014. Polyphosphate accumulation is driven by transcriptome alterations that lead to near-synchronous and near-equivalent uptake of inorganic cations in an arbuscular mycorrhizal fungus. The new phytologist **204**, 638–649.

Kim C. 2003. SAM domains: uniform structure, diversity of function. Trends in biochemical sciences **28**, 625–628.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nature biotechnology **37**, 907–915.

Krusell L, Sato N, Fukuhara I, *et al.* 2011. The *Clavata2* genes of pea and *Lotus japonicus* affect autoregulation of nodulation. The Plant journal: for cell and molecular biology **65**, 861–871.

Li X-R, Sun J, Albinsky D, *et al.* 2022. Nutrient regulation of lipochitooligosaccharide recognition in plants via NSP1 and NSP2. Nature communications **13**, 1–15.

Limpens E, Franken C, Smit P, Willemse J, Bisseling T, Geurts R. 2003. LysM Domain Receptor Kinases Regulating Rhizobial Nod Factor-Induced Infection. Science **302**, 630–633.

Lindsay PL, Ivanov S, Pumplin N, Zhang X, Harrison MJ. 2022. Distinct ankyrin repeat subdomains control VAPYRIN locations and intracellular accommodation functions during arbuscular mycorrhizal symbiosis. Nature communications **13**, 5228.

Lindsay P, Swentowsky KW, Jackson D. 2023. Cultivating potential: Harnessing plant stem cells for agricultural crop improvement. Molecular plant doi: 10.1016/j.molp.2023.12.014.

Liu W, Kohlen W, Lillo A, *et al.* 2011. Strigolactone biosynthesis in Medicago truncatula and rice requires the symbiotic GRAS-type transcription factors NSP1 and NSP2. The Plant cell **23**, 3853–3865.

Lobet G, Pagès L, Draye X. 2011. A novel image-analysis toolbox enabling quantitative analysis of root system architecture. Plant physiology **157**, 29–39.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology **15**, 550.

MacLean AM, Bravo A, Harrison MJ. 2017. Plant Signaling and Metabolic Pathways Enabling Arbuscular Mycorrhizal Symbiosis. The Plant cell **29**, 2319–2335.

Maldonado-Mendoza IE, Dewbre GR, Harrison MJ. 2001. A phosphate transporter gene from the extra-radical mycelium of an arbuscular mycorrhizal fungus *Glomus intraradices* is regulated in response to phosphate in the environment. Molecular plant-microbe interactions: MPMI **14**, 1140–1148.

Marone D, Russo M, Laidò G, De Leonardis A, Mastrangelo A. 2013. Plant nucleotide binding site–leucine-rich repeat (NBS-LRR) genes: Active guardians in host defense responses. International journal of molecular sciences **14**, 7302–7326.

Marro N, Lidoy J, Chico MÁ, Rial C, García J, Varela RM, Macías FA, Pozo MJ, Janoušková M, López-Ráez JA. 2022. Strigolactones: New players in the nitrogen–phosphorus signalling interplay. Plant, cell & environment **45**, 512–527.

McGonigle PT, Miller MH, Evans DG, Fairchild GL, Swan JA. 1990. A new method which gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi. The New phytologist **150**, 495–501.

Mirzaei S, Batley J, El-Mellouki T, Liu S, Meksem K, Ferguson BJ, Gresshoff PM. 2017. Neodiversification of homeologous CLAVATA1-like receptor kinase genes in soybean leads to distinct developmental outcomes. Scientific reports **7**.

Morandi D, Sagan M, Prado-Vivant E, Duc G. 2000. Influence of genes determining supernodulation on root colonization by the mycorrhizal fungus Glomus mosseae in Pisum sativum and Medicago truncatula mutants. Mycorrhiza **10**, 37–42.

Mortier V, Den Herder G, Whitford R, Van de Velde W, Rombauts S, D'Haeseleer K, Holsters M, Goormachtig S. 2010. CLE peptides control Medicago truncatula nodulation locally and systemically. Plant physiology **153**, 222–237.

Mugford ST, Osbourn A. 2012. Saponin synthesis and function. Isoprenoid Synthesis in Plants and Microorganisms. New York, NY: Springer New York, 405–424.

Müller R, Bleckmann A, Simon R. 2008. The receptor kinase CORYNE of Arabidopsis transmits the stem cell-limiting signal CLAVATA3 independently of CLAVATA1. The plant cell **20**, 934–946.

Müller LM, Campos-Soriano L, Lévesque-Tremblay V, Bravo A, Daniels DA, Pathak S, Park H-J, Harrison MJ. 2020. Constitutive Overexpression of RAM1 Leads to an Increase in Arbuscule Density in Brachypodium distachyon. Plant physiology **184**, 1263.

Müller LM, Flokova K, Schnabel E, Sun X, Fei Z, Frugoli J, Bouwmeester HJ, Harrison MJ. 2019. A CLE–SUNN module regulates strigolactone content and fungal colonization in arbuscular mycorrhiza. Nature plants **5**, 933–939.

Nimchuk ZL. 2017. CLAVATA1 controls distinct signaling outputs that buffer shoot stem cell proliferation through a two-step transcriptional compensation loop. PLoS genetics **13**, e1006681.

Nouri E, Breuillin-Sessoms F, Feller U, Reinhardt D. 2014. Phosphorus and nitrogen regulate arbuscular mycorrhizal symbiosis in petunia hybrida. PloS one **9**, e90841.

Nowak S, Schnabel E, Frugoli J. 2019. The Medicago truncatula CLAVATA3-LIKE CLE12/13 signaling peptides regulate nodule number depending on the CORYNE but not the COMPACT ROOT ARCHITECTURE2 receptor. Plant signaling & behavior **14**, 1598730.

Oldroyd GED. 2013. Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. Nature reviews. Microbiology **11**, 252–263.

Pallakies H, Simon R. 2014. The CLE40 and CRN/CLV2 signaling pathways antagonistically control root meristem growth in Arabidopsis. Molecular plant **7**, 1619–1636.

Paries M, Gutjahr C. 2023. The good, the bad, and the phosphate: regulation of beneficial and detrimental plant–microbe interactions by the plant phosphate status. The new phytologist **239**, 29–46.

Passardi F, Cosio C, Penel C, Dunand C. 2005. Peroxidases have more functions than a Swiss army knife. Plant cell reports **24**, 255–265.

Replogle A, Wang J, Bleckmann A, Hussey RS, Baum TJ, Sawa S, Davis EL, Wang X, Simon R, Mitchum MG. 2011. Nematode CLE signaling in Arabidopsis requires CLAVATA2 and CORYNE. The Plant journal: for cell and molecular biology **65**, 430–440.

Rodriguez-Leal D, Xu C, Kwon C-T, *et al.* 2019. Evolution of buffering in a genetic circuit controlling plant stem cell proliferation. Nature genetics **51**, 786–792.

Roy S, Müller LM. 2022. A rulebook for peptide control of legume-microbe endosymbioses. Trends in plant science **27**, 870–889.

Rush TA, Puech-Pagès V, Bascaules A, *et al.* 2020. Lipo-chitooligosaccharides as regulatory signals of fungal growth and development. Nature communications **11**.

Schindelin J, Arganda-Carreras I, Frise E, *et al.* 2012. Fiji: an open-source platform for biological-image analysis. Nature methods **9**, 676–682.

Schnabel E, Bashyal S, Corbett C, Kassaw T, Nowak S, Rosales-García RA, Noorai RE, Müller LM, Frugoli J. 2024. The Defective in Autoregulation (DAR) gene of Medicago truncatula encodes a protein involved in regulating nodulation and arbuscular mycorrhiza. BMC plant biology **24**, 766.

Schnabel E, Journet E-P, de Carvalho-Niebel F, Duc G, Frugoli J. 2005. The Medicago truncatula SUNN gene encodes a CLV1-like leucine-rich repeat receptor kinase that regulates nodule number and root length. Plant molecular biology **58**, 809–822.

Schnabel E, Karve A, Kassaw T, Mukherjee A, Zhou X, Hall T, Frugoli J. 2012. The*M. truncatula SUNN*gene is expressed in vascular tissue, similarly to*RDN1*, consistent with the role of these nodulation regulation genes in long distance signaling. Plant signaling & behavior **7**, 4–6.

Selby R, Jones DS. 2023. Complex peptide hormone signaling in plant stem cells. Current opinion in plant biology **75**, 102442.

Serrano K, Bezrutczyk M, Goudeau D, Dao T, O'Malley R, Malmstrom RR, Visel A, Scheller HV, Cole B. 2024. Spatial co-transcriptomics reveals discrete stages of the arbuscular mycorrhizal symbiosis. Nature plants **10**, 673–688.

Smit P, Limpens E, Geurts R, Fedorova E, Dolgikh E, Gough C, Bisseling T. 2007. Medicago LYK3, an entry receptor in rhizobial nodulation factor signaling. Plant physiology **145**, 183–191.

Smith SE, Reid D. 2008. *Mycorrhizal Symbiosis*. Academic Press.

Somssich M, Bleckmann A, Simon R. 2016*a*. Shared and distinct functions of the pseudokinase CORYNE (CRN) in shoot and root stem cell maintenance of Arabidopsis. Journal of experimental botany **67**, 4901–4915.

Somssich M, Je BI, Simon R, Jackson D. 2016*b*. CLAVATA-WUSCHEL signaling in the shoot meristem. Development **143**, 3238–3248.

Stammers DK. 2001. The structure of the negative transcriptional regulator NmrA reveals a structural superfamily which includes the short-chain dehydrogenase/reductases. The EMBO journal **20**, 6619–6626.

Tang H, Krishnakumar V, Bidwell S, *et al.* 2014. An improved genome release (version Mt4.0) for the model legume Medicago truncatula. BMC genomics **15**, 312.

Thomas J, Frugoli J. 2024. Mutation of BAM2 rescues the sunn hypernodulation phenotype in Medicago truncatula, suggesting that a signaling pathway like CLV1/BAM in Arabidopsis affects nodule number. Frontiers in plant science **14**.

Tian C, Kasiborski B, Koul R, Lammers PJ, Bücking H, Shachar-Hill Y. 2010. Regulation of the nitrogen transfer pathway in the arbuscular mycorrhizal symbiosis: Gene characterization and the coordination of expression with nitrogen flux. Plant physiology **153**, 1175–1187.

Tsuno Y, Fujimatsu T, Endo K, Sugiyama A, Yazaki K. 2018. Soyasaponins: A New Class of Root Exudates in Soybean (Glycine max). Plant & cell physiology **59**, 366–375.

Vicente J, Cascón T, Vicedo B, García-Agustín P, Hamberg M, Castresana C. 2012. Role of 9-lipoxygenase and α-dioxygenase oxylipin pathways as modulators of local and systemic defense. Molecular plant **5**, 914–928.

Wang C, Reid JB, Foo E. 2018. The Art of Self-Control - Autoregulation of Plant-Microbe Symbioses. Frontiers in plant science **9**, 988.

Wang C, Velandia K, Kwon C-T, Wulf KE, Nichols DS, Reid JB, Foo E. 2021. The role of CLAVATA signalling in the negative regulation of mycorrhizal colonization and nitrogen response of tomato. Journal of experimental botany **72**, 1702–1713.

Whitewoods CD. 2021. Evolution of CLE peptide signalling. Seminars in cell & developmental biology **109**, 12–19.

Wojtaszek P. 1997. Oxidative burst: an early plant response to pathogen infection. The Biochemical journal **322**, 681–692.

Wulf K, Sun J, Wang C, *et al.* 2024. The role of CLE peptides in the suppression of mycorrhizal colonization of tomato. Plant & cell physiology **65**, 107–119.

Xie X, Lai W, Che X, Wang S, Ren Y, Hu W, Chen H, Tang M. 2022. A SPX domain-containing phosphate transporter from *Rhizophagus irregularis* handles phosphate homeostasis at symbiotic interface of arbuscular mycorrhizas. The new phytologist **234**, 650–671.

Xie Z, Nolan TM, Jiang H, Yin Y. 2019. AP2/ERF transcription factor regulatory networks in hormone and abiotic stress responses in Arabidopsis. Frontiers in plant science **10**.

Yildirir G, Sperschneider J, Malar C M, Chen ECH, Iwasaki W, Cornell C, Corradi N. 2022. Long reads and Hi-C sequencing illuminate the two-compartment genome of the model arbuscular mycorrhizal symbiont *Rhizophagus irregularis*. The new phytologist **233**, 1097–1107.

Yoneyama K, Xie X, Kim HI, Kisugi T, Nomura T, Sekimoto H, Yokota T, Yoneyama K. 2012. How do nitrogen and phosphorus deficiencies affect strigolactone production and exudation? Planta **235**, 1197–1207.

Yoneyama K, Xie X, Kusumoto D, Sekimoto H, Sugimoto Y, Takeuchi Y, Yoneyama K. 2007. Nitrogen deficiency as well as phosphorus deficiency in sorghum promotes the production and exudation of 5-deoxystrigol, the host recognition signal for arbuscular mycorrhizal fungi and root parasites. Planta **227**, 125–132.

Zhang J, Sun J, Chiu CH, *et al.* 2024. A receptor required for chitin perception facilitates arbuscular mycorrhizal associations and distinguishes root symbiosis from immunity. Current biology: CB **34**, 1705-1717.e6.