1 Limited population structure but signals of recent selection in introduced African Fig
2 Fly *(Zaprionus indianus)* in North America
3

4 Priscilla A. Erickson[1]*, Alyssa Bangerter[2], Ansleigh Gunter[1], Nikolaos T. Polizos[3], and Alan O.
5 Bergland[2]
6

7   1. University of Richmond, Richmond, Virginia
8   2. University of Virginia, Charlottesville, Virginia
9   3. University of Miami, Coral Gables, Florida
10   *Corresponding author; perickso@richmond.edu*

12 Running title
13

14 Genomics of invasive *Zaprionus indianus*
15

16 Abstract
17

Invasive species have devastating consequences for human health, food security, and the
environment. Many invasive species adapt to new ecological niches following invasion, but
little is known about the early steps of adaptation. Here we examine population genomics of a
recently introduced drosophilid in North America, the African Fig Fly, *Zaprionus indianus*. This
species is likely intolerant of subfreezing temperatures and recolonizes temperate
environments yearly. We generated a new chromosome-level genome assembly for *Z.
indianus*. Using resequencing of over 200 North American individuals collected over four years
in temperate Virginia, plus a single collection from subtropical Florida, we tested for signatures
of recolonization, population structure, and adaptation within invasive populations. We show
founding populations are sometimes small and contain close genetic relatives, yet temporal
population structure and differentiation of populations is mostly absent across recurrent
recolonization events. Although we find limited signals of genome-wide spatial or temporal
population structure, we identify haplotypes on the X chromosome that are repeatedly
differentiated between Virginia and Florida populations. These haplotypes show signatures of
natural selection and are not found in African populations. We also find evidence for several
large structural polymorphisms segregating within North America populations and show X
chromosome evolution in invasive populations is strikingly different from the autosomes. These
results show that despite limited population structure, populations may rapidly evolve genetic
differences early in an invasion. Further uncovering how these genomic regions influence
invasive potential and success in new environments will advance our understanding of how
organisms evolve in changing environments.

40 Article Summary
41

42 Invasive species (organisms that have been moved outside their natural range by human
43 activities) can cause problems for both humans and the environment. We studied the genomes

44 of over 200 individuals of a newly invasive fruit fly in North America, the African Fig Fly. We
45 found genetic evidence that these recently introduced flies may be evolving in their new
46 environments, which could make them stronger competitors and more likely to become pests.
47
48 **Introduction**
49
Understanding how species expand and adapt to new environments in an era of climate
change and global commerce is central to controlling the spread of disease (Altizer et al. 2013;
Hoberg and Brooks 2015), to maintaining crop security (Oerke 2006; Sutherst et al. 2011) and
to preserving biodiversity (Bellard et al. 2012). Many organisms are moving to new, previously
unoccupied ranges at rates that continue to accelerate (Ricciardi 2007; Seebens et al. 2015;
Seebens et al. 2017; Platts et al. 2019; Sardain et al. 2019) due to changes in climate and
habitat as well as anthropogenic introductions. Genetic adaptation to new environments may
allow some vulnerable organisms to survive in new habitats but may also permit potentially
harmful organisms to expand even further (Clements and Ditommaso 2011). The past two
decades have produced a wealth of studies characterizing the genetic and genomic basis of
adaptation in a variety of organisms, from experimental populations of microbes (Good et al.
2017; Nguyen Ba et al. 2019; Johnson et al. 2021) to natural populations of eukaryotes
(Hancock et al. 2011; Jones et al. 2018; Barrett et al. 2019; Lovell et al. 2021; Schluter et al.
2021). Recent and ongoing invasions offer the opportunity to study rapid evolution and
adaptation to new environments in nearly real-time (Koch et al. 2020; Pélissié et al. 2022;
Parvizi et al. 2023; Soudi et al. 2023). Recently, genomics has helped trace the history and
sources of many well-known invasions (Pélissié et al. 2022; Picq et al. 2023) and shown that
genetic divergence and even local adaptation are common in invasive populations that have
been established for decades or even centuries (Ma et al. 2020; Stuart et al. 2021; Li et al.
2023). However, much remains unknown about the genetic mechanisms that allow invasive
organisms to colonize and thrive in new environments. A better understanding of adaptive
pathways in invasion may assist in predicting the success of invasions and controlling their
outcomes.
73
The African Fig Fly, *Zaprionus indianus*, serves as a unique model to study how
invasion history and local environment influence patterns of genetic variation. The ongoing,
recurrent invasion of *Z. indianus* in North America offers a premier opportunity to study the
possibility of rapid genetic changes following invasion. The *Zaprionus* genus arose in Africa but
Z. indianus was first described in India in 1970 (Gupta 1970), where it has adapted to a range
of environments (da Mata et al. 2010). It is one of the most ecologically diverse drosophilids in
Africa; its ability to utilize up to 80 different food sources (Yassin and David 2010) and its
generation time of as few as ~13 days (Nava et al. 2007) likely fueled its spread around the
world. In 1999, it was first detected in Brazil (Vilela 1999), where it subsequently spread and
caused major damage to fig and berry crops as well as native fruit species (Leão and Tldon
2004; Oliveira et al. 2013; Roque et al. 2017; Zanuncio-Junior et al. 2018; Allori Stazzonelli et
al. 2023). It was later found in Mexico and Central America in 2002-2003 (Markow et al. 2014)
and eventually Florida in 2005 (Linde et al. 2006). In 2011-2012, its range expanded

northwards in eastern North America (Joshi et al. 2014; Timmeren and Isaacs 2014; Pfeiffer et al. 2019) and eventually reached as far north as Ontario (Renkema et al. 2013) and Minnesota (Holle et al. 2018). It has also recently been found in the Middle East, Europe, and Hawaii (Parchami-Araghi et al. 2015; Kremmer et al. 2017; Willbrand et al. 2018), suggesting that the invasion is ongoing. *Z. indianus* can damage fig and berry crops (Pfeiffer et al. 2019; Allori Stazzonelli et al. 2023), increasing concerns about its pest potential in its expanding range.

Despite its global success, *Z. indianus* males are sterile below 15 °C, making cold temperatures a limiting factor to their success (Araripe et al. 2004). Within the temperate environment of Virginia, the species exhibits strong seasonal fluctuations in abundance (Rakes et al. 2023). First detection in Virginia is usually in June or July, weeks after the appearance of other overwintering Drosophilids, and population sizes climb dramatically through the late summer and early fall, when it often dominates the drosophilid community in temperate orchards. Typically, the peak in early to mid-September is followed by a dip in abundance and then a second peak in October, suggesting a seasonal component to reproduction or fluctuations in factors influencing *Z. indianus'* relative fitness. However, despite its early post-colonization success, it does not appear to survive temperate winters; *Z. indianus* populations became undetectable in Virginia by early December (Rakes et al. 2023). In locations in Minnesota, Kansas, and the northeastern US, *Z. indianus* has been detected one year and then not the next, suggesting that the populations are not permanently established, but are extirpated by cold and re-introduced by stochastic dispersal processes (Holle et al. 2018; Gleason et al. 2019; Rakes et al. 2023). Therefore, *Z. indianus* likely repeatedly invades temperate environments and evolves for several generations in these new habitats, offering an opportunity to recurrently study the genetic impacts of invasion and post-colonization adaptation across multiple years of sampling.

Genetic *s*tudies *of Z. indianus* are limited but provide important context to understand its worldwide invasion. The invasion of North America likely resulted from separate founding events on the East and West coasts (Commar et al. 2012). Comeault *et al* (2020) showed that North American populations are genetically distinct from those from Africa. Invasive populations of *Z. indianus* have an approximately 30% reduction in genetic diversity relative to ancestral African populations (Comeault et al. 2020), though invasive populations of *Z. indianus* maintain levels of genetic diversity that are often higher than those of non-invasive congeners. Despite the loss of diversity, *Z. indianus* is extremely successful in temperate habitats (Rakes et al. 2023). Further studies demonstrated that genetically distinct populations from eastern and western Africa likely admixed prior to a single colonization of the Americas (Comeault et al. 2021). How the high degree of genetic diversity in invasive populations influences the potential for ongoing evolution in North America, which is in a critical early stage of invasion, remains understudied.

Here, we assembled and annotated a chromosome-level genome assembly for *Z. indianus* and used the newly improved genome to answer several questions with the whole genome sequences of over 200 North American flies collected from three locations over four

130 years. First, do recolonizing North American *Z. indianus* populations demonstrate spatial or
131 temporal population structure and if so, do specific regions of the genome have an outsized
132 contribution to population structure? Second, is the invasion and recolonization history
133 recapitulated in population genetic data? And third, do temperate populations show signatures
134 of selection relative to native and tropical invasive populations?
135
136 **Materials and Methods**
137
138 *Hi-C based genome scaffolding*
139
An inbred line was generated from flies originally captured from Carter Mountain Orchard, VA
(37.9913° N, 78.4721° W) in 2018. Wild caught flies were reared in the lab for approximately
one year prior to initiating isofemale lines. The offspring of the isofemale lines were propagated
through 10 rounds of full-sib mating. The resulting lines were then passaged for approximately
one additional year in the lab and the most vigorous remaining line ("24.2") was chosen for
sequencing.
146
3rd instar larvae from a single inbred line were snap frozen in liquid nitrogen and sent to
Dovetail corporation (now Cantata Bio, Scotts Valley, CA) for chromatin extraction, Hi-C
sequencing and genome scaffolding. Briefly, chromatin was fixed in place with formaldehyde in
the nucleus and then extracted. Fixed chromatin was digested with DNAse I, chromatin ends
were repaired and ligated to a biotinylated bridge adapter followed by proximity ligation of
adapter containing ends. After proximity ligation, crosslinks were reversed and the DNA
purified. Purified DNA was treated to remove biotin that was not internal to ligated fragments.
Sequencing libraries were generated using NEBNext Ultra enzymes and Illumina-compatible
adapters. Biotin-containing fragments were isolated using streptavidin beads before PCR
enrichment of each library. The library was sequenced on an Illumina HiSeqX platform to
produce approximately 30x sequence coverage.
158
The input *de novo* assembly was the *Z. indianus* "RCR04" PacBio assembly (assembly
ASM1890459v1) from Kim et al. (2021). This assembly and Dovetail OmniC library reads
were used as input data for HiRise, a software pipeline designed specifically for using
proximity ligation data to scaffold genome assemblies (Putnam et al. 2016). Dovetail OmniC
library sequences were aligned to the draft input assembly using *bwa* (Li and Durbin 2009).
The separations of Dovetail OmniC read pairs mapped within draft scaffolds were analyzed by
HiRise to produce a likelihood model for genomic distance between read pairs, and the model
was used to identify and break putative misjoins, to score prospective joins, and make joins
above a threshold. See Figure S1 for link density histogram of scaffolding data.
168
169 *Annotation*
170
Repeat families found in the genome assemblies of *Z. indianus* were identified de novo and
classified using the software package *RepeatModeler* v. 2.0.1 (Flynn et al. 2020).

RepeatModeler depends on the programs *RECON* v. 1.08 (Bao and Eddy 2002) and
RepeatScout v. 1.0.6 (Price et al. 2005) for the de novo identification of repeats within the
genome. The custom repeat library obtained from *RepeatModeler* was used to discover,
identify and mask the repeats in the assembly file using *RepeatMasker* v. 4.1.0 (Smit et al.
2015).
178
RNA sequencing was conducted on 3 replicates of 3rd instar larva and 3 replicates of
mixed stage pupa that were snap frozen in liquid nitrogen. RNA extraction and sequencing
was performed by GeneWiz (South Plainfield, NJ). New larval and pupal RNAseq reads were
combined with adult RNA sequencing from Comeault et al. (2020) for annotation. Coding
sequences from *D. grimshawi, D. melanogaster*, *D. pseudoobscura, D. virilis, Z. africanus, Z.*
indianus, Z. tsacasi and *Z. tuberculatus* (Kim et al. 2021) were used to train the initial *ab initio*
model for *Z. indianus* using the *AUGUSTUS* software v. 2.5.5 (Keller et al. 2011). Six rounds of
prediction optimization were done with the software package provided by *AUGUSTUS*. The
same coding sequences were also used to train a separate *ab initio* model for *Z. indianus*
using *SNAP* (version 2006-07-28) (Korf 2004). RNAseq reads were mapped onto the genome
using the *STAR* aligner software (version 2.7) (Dobin et al. 2013) and intron hints generated
with the bam2hints tools within *AUGUSTUS*. *MAKER* v. 3.01.03 (Cantarel et al. 2008), SNAP
and AUGUSTUS (with intron-exon boundary hints provided from RNAseq) were then used to
predict for genes in the repeat-masked reference genome. To help guide the prediction
process, Swiss-Prot peptide sequences from the UniProt database were downloaded and used
in conjunction with the protein sequences from *D. grimshawi, D. melanogaster, D.*
pseudoobscura, D. virilis, Z. africanus, Z. indianus, Z. tsacasi and *Z. tuberculatus* to generate
peptide evidence in the *MAKER* pipeline. Only genes that were predicted by both SNAP and
AUGUSTUS were retained in the final gene sets. To help assess the quality of the gene
prediction, AED scores were generated for each of the predicted genes as part of the *MAKER*
pipeline. Genes were further characterized for their putative function by performing a BLAST
search of the peptide sequences against the UniProt database. tRNA were predicted using the
software *tRNAscan-SE* v. 2.05 (Chan and Lowe 2019). Transcriptome completeness was
assessed with *BUSCO* v. 4.0.5 (Manni et al. 2021) using the eukaryota_odb10 list of 255
genes.
204
205   *Wild fly collections*
206
Flies were collected by aspiration and netting from Carter Mountain Orchard, VA (37.9913° N,
78.4721° W) in 2017-2020 and from Hanover Peach Orchard, VA (37.5694° N, 77.2660° W) in
2019-2020. Flies were sampled from Coral Gables, FL (25.7239° N, 80.2802° W) in June 2019
using traps baited with bananas, oranges, yeast, and red wine. Flies were frozen in 70%
ethanol at -20°C (2017-2018) or dry at -80 °C (2019-2020) prior to sequencing. Collections
performed in July and August were called "early season." In 2019, the earliest collections were
not made until September (typically when *Z. indianus* abundance peaks, Rakes *et al.* 2023),
and were assigned "mid-season." Collections from October and November were called "late
season." For some analyses, the mid-season collection and early collections were combined,

as they were the first collections available each year. See Table S1 for the number of
individual flies sequenced from each location and timepoint.
218
219   *Individual whole genome sequencing*
220
The sex of each wild-caught fly was recorded, then DNA was extracted from individual flies
using the DNAdvance kit (Beckman Coulter, Indianapolis, IN) in 96 well plates, including an
additional RNAse treatment step. DNA concentration was measuring using the QuantIT kit
(Invitrogen, Waltham, MA) and purified DNA was diluted to 1 ng/µL. Libraries were prepared
from 1 ng of genomic DNA using a reduced-volume dual-barcoding Nextera (Illumina, San
Diego, CA) protocol as previously described (Erickson et al. 2020). The libraries were
quantified using the QuantIT kit and equimolar ratios of each individual DNA were combined
for sequencing. The pooled library was size-selected for 500 bp fragments using a BluePippin
gel cassette (Sage Sciences, Beverly, MA). The pooled libraries were sequenced in one
Illumina NovaSeq 6000 lane using paired-end, 150 bp reads by Novogene (Sacramento, CA).
231
Existing raw reads from *Z. indianus* collections from North America, South America, and
Africa (Comeault et al. 2020; Comeault et al. 2021) were downloaded from the SRA from
BioProject number PRJNA604690. These samples were combined with the new sequence
data and processed together with the same mapping and SNP-calling pipeline. Overlapping
paired-end reads were merged with *BBMerge* v. 38.92 (Bushnell et al. 2017). Reads were
mapped to the genome assembly described above using *bwa mem* v. 0.7.17 (Li and Durbin
2009). Bam files for merged and unmerged reads were combined, sorted and de-duplicated
with *Picard* v. 2.26.2 (https://github.com/broadinstitute/picard).
240
We next used *Haplotype Caller* from *GATK* v. 4.2.0.0 (McKenna et al. 2010) to generate
a gVCF for each individual. We built a GenomicsDBI database for each scaffold, then used this
database to genotype each gVCF. We used *GATK's* hard filtering options to filter the raw
SNPs based on previously published parameters (*--filter-expression "QD < 2.0 || FS > 60.0 ||*
SOR > 3.0 || MQ < 40.0 || MQRankSum < -12.5 || ReadPosRankSum < -8.0") (Comeault et al.
2020). We then removed SNPs within 20 bp of an indel from the output and removed all SNPs
in regions identified by *RepeatMasker*. We analyzed several measures of individual and SNP
quality using *VCFtools* v. 0.1.17 (Danecek et al. 2011). We removed 16 individuals with mean
coverage < 7X or over 10% missing genotypes. Next, we filtered SNPs with mean depths <10
or > 50 across all samples. We removed individual genotypes supported by 6 or fewer reads or
with more than 100 reads to produce a final VCF with 5,185,389 SNPs and 2,099,147 non-
singleton SNPs. See Table S1 for the final number of individuals included in the analysis from
each population. See Figure S2 for the average SNP depth per sampling time and location.
254
255   *Sex chromosome and Muller element identification*
256
samtools v. 1.12 (Li et al. 2009) was used to measure coverage and depth of mapped reads
from individual sequencing. This analysis revealed that the five main scaffolds (all over 25 Mb

in length) had a mean depth of ~16X coverage in both males and females in our dataset,
except for scaffold 3, which had ~16X coverage in females but ~8X coverage in males,
suggesting it is the X chromosome (Figure S3). Some of the previously sequenced samples
had no sex recorded, so we used the ratio of X chromosome reads (scaffold 3) to autosome
(scaffolds 1, 2, 4 and 5) reads to assign sexes to those individuals. Individuals with a ratio
greater than 0.8 were assigned female, and ratios less than 0.8 were assigned male (Figure
S4). For two known-sex individuals, the sex recorded prior to sequencing did not match the
sex based on coverage; for those two samples we used the coverage-based sex assignment
for analyses. We used D-GENIES (Cabanettes and Klopp 2018) to create dot-plots comparing
the *Z. indianus* and *D. melanogaster* genome (BDGP6.46, downloaded from ensemble.org) to
confirm the sex chromosome identification and assign Muller elements to *Z. indianus*
autosomes (Figure S5, Table S2). Five additional scaffolds had lengths over 1 Mb. Scaffold 8
is the dot chromosome (Muller element F) based on sequence comparison to *D. melanogaster*
(Figure S5) and had similar coverage to the autosomes (Figure S3). Scaffolds 6,7,9, and 10
had reduced coverage (Figure S3) and contain mostly repetitive elements. Downstream SNP
calling and population genetic analysis included the five large scaffolds (named chromosomes
1-5) and excluded all smaller scaffolds.
276
277   *Related individuals*
278
Preliminary exploration of population genetic data indicated that some individual samples may
be close relatives. For downstream analyses, we used the–*king-cutoff 0.0625* argument in
Plink v. 2.0 (Chang et al. 2015) to generate a list of unrelated individuals. This filtering
removed 21 individuals from the dataset. To quantify relatedness between all individuals, we
used the function *snpgdsibdKING* in *SNPRelate* v. 1.38.0 (Zheng et al. 2012) to determine the
kinship coefficients and probability of zero identity by descent for pairs of individuals using
autosomal SNPs. We used thresholds established in Thornton *et al.* (2012) to classify
relatedness between individuals.
287
288   *Population structure and F_{ST}*
289
We conducted principal components analysis using the R package *SNPRelate* v. 1.38.0
(Zheng et al. 2012) in R v. 4.1.1 (R Core Team) using a vcf that excluded singleton SNPs. We
LD pruned SNPs with a minor allele frequency of at least 0.05 using *SNPgdsLDpruning* with an
LD threshold of 0.2 and then calculated principal components with *snpgdsPCA* using all four
autosomes. For subsequent analyses, we repeated the LD pruning within subsets of the data
(North America only, or Carter Mountain, VA only). We also calculated principal components
using individual chromosomes; for the X chromosome, only females were used in the analysis.
We used t-tests and one-way ANOVAs followed by Tukey post-hoc tests to compare PC
values between sampling locations and time points.
299
300   We used *Plink* v. *1.9* (Purcell et al. 2007; Chang et al. 2015) to LD prune VCF files with
301   parameters (*--indep-pairwise 1000 50 0.2*) and used *ADMIXTURE* v. 1.3.0 (Alexander and

302 Lange 2011) to evaluate population structure for each chromosome separately. For the X
303 chromosome, only females were used. We tested up to k=10 genetic clusters and used cross-
304 validation analysis to choose the optimal k for each chromosome separately.
305
306   We calculated F_{ST} between Florida samples and early season Virginia samples using
307 the *snpgdsFST* function in *SNPRelate* for all SNPs with a minor allele frequency > 0.01. For
308 the X chromosome, only females were used in F_{ST} calculations to ensure diploid genotypes.
309 We used the same function to calculate genome-wide, pairwise F_{ST} between all Virginia
310 collections using autosomal SNPs.
311
312 *Testing for structural variants*
313
We used *smoove* v. 0.2.6 (Pedersen et al. 2020) to identify and genotype insertions, deletions,
and rearrangements in the paired-end sequencing data from all individuals as described in the
documentation. We also used linkage disequilibrium (LD) of randomly sampled SNPs from
each chromosome to visually inspect for linkage due to potential inversions. We generated a
list of SNPs segregating in each focal population with no missing genotypes and randomly
sampled 4,000 SNPs from each chromosome. We used the *snpgdsLDMat* function in
SNPRelate to calculate LD between all pairs of SNPs. LD heatmaps were created with the
ggLD package (https://github.com/mmkim1210/ggLD).
322
323 *Estimation of historic population sizes*
324
We used *smc++* v. 1.15.4 (Terhorst et al. 2017) to estimate historic population sizes for several
subpopulations of individuals using autosomal genotypes. We used individuals from each
African location and used the earliest sampling available for each year and Virginia orchard.
We used *vcf2smc* to prepare the input files for each autosome separately. We assigned each
individual as the "distinguished individual" and ran the analysis using all possible combinations
of distinguished individual as described in (Bemmels et al. 2021). We used 10-fold cross
validation to estimate final model parameters with the option (*-cv –folds 10)*. We assumed a
generation time of 0.08 years (~12 generations per year) based on Nava *et al*. (2007), which
assumes year-round reproduction in tropical regions. We note that for Virginia populations
experiencing temperate conditions in recent years, 12 generations per year is likely an
overestimate due to the shortened breeding season.
336
337 *Selection scan*
338
We used *WhatsHap* v. 1.7 (Patterson et al. 2015) to perform read-based phasing of the full vcf
including singletons. To polarize the vcf for the genome wide selection scan relative to the
invasion, we reassigned the reference allele of the phased vcf as the allele that was most
common across all African individuals sequenced in previous studies. We calculated allele
frequencies using all African samples in *SNPRelate*, then used *vcf-info-annotator*
(https://vatools.readthedocs.io/en/latest/index.html) to assign the "ancestral" allele in the INFO

column. Lastly we used *bcftools* v. 1.13 (Danecek et al. 2021) to make simplified vcfs
containing only the GT and AA fields for each chromosome separately.
347
We used the R package *rehh* v. 3.2.2 (Gautier and Vitalis 2012) to conduct the selection
scan using integrated haplotype homozygosity score. We split samples into four possible
populations (Africa, Florida, all North America, Virginia only) and conducted the scans
separately for each population using phased, polarized vcfs for each individual chromosome.
We used the *haplo2hh*, *scan,* and *ihh2ihs* functions to implement the scan. For the X
chromosome, we only used a single haplotype for each male in the dataset to avoid double
counting haploid genotypes. Haplotypes under selection were visualized by plotting all SNPs
with IHS > 5. LD between candidate SNPs was calculated in *SNPRelate*.
356
357 *Genetic diversity statistics*
358
Because we obtained variable sequencing coverage within and across populations (Figure S2)
we used software designed for low coverage and missing data to analyze population genetic
statistics in genomic windows. We used *pixy* v. 1.2.5 (Korunes and Samuk 2021) to calculate
Pi, F_{ST} and D_{XY} in 5 kb windows. Samples were grouped by collection location and year or by
collection location for different analyses. We used *ANGSD* v. 0.941 (Korneliussen et al. 2014)
to calculate Tajima's D. We first calculated genotype likelihoods from the bam files using
arguments *-doSaf* and *-GL*. We then calculated Tajima's D and theta using the folded site
frequency spectrum across 5 kb windows with 5 kb steps as described in ANGSD
documentation.
368
369 *Data management and plotting*
370
371 We used the R packages *foreach* (Microsoft and Weston 2017) and *data.table* (Dowle and
372 Srinivasan 2019) for data management and manipulation and used *ggplot2* (Wickham 2016)
373 for all plotting. The *ggpubfigs* (Steenwyk and Rokas 2021) and *viridis* (Garnier 2018) packages
374 were used for color palettes.
375
376 **Results and Discussion**
377
378 *Genome assembly and annotation*
379
High quality genome assemblies and annotations are a critical component of tracking and
controlling invasive species and understanding the potential evolution of invasive species in
invaded ranges (Matheson and McGaughran 2022). We conducted Hi-C based scaffolding of a
previously sequenced *Z. indianus* genome (Kim et al. 2021) to achieve a chromosome-level
assembly. There were 1,014 scaffolds with an N50 of 26.6 Mb, an improvement from an N50
of 4.1-6.8 Mb in previous assemblies (Kim et al. 2021). The five main chromosomes (Figure
S1, named in order of size from largest to smallest) varied in length from 25.7 to 32.3 Mb (total
length of five main scaffolds = 146,062,119 bp), in agreement with *Z. indianus* karyotyping

(Gupta and Kumar 1987; Campos et al. 2007). Chromosome 3 was identified as the sex
chromosome using sequencing coverage of known-sex individuals (Figure S3, S4) and
sequence comparison dot-plots (Figure S5). See Table S2 for assignment of *Z. indianus*
chromosomes to Muller elements based on alignment to the *D. melanogaster* genome.
392
The annotation using RNAseq from larvae, pupae, and adults predicted 13,162
transcripts and 13,075 proteins, with 93% of 255 benchmarking universal single copy orthologs
(BUSCO) genes (Simão et al. 2015) identified as complete and an additional 1.2% of BUSCO
genes identified as fragmented. This transcriptome-based completeness estimate is lower than
the genome-based estimate of 99% complete (Kim et al. 2021) but is in line with other
arthropod genomes (Feron and Waterhouse 2022). Within the 5 main scaffolds, 24.6% of
sequences were repetitive; within the entire assembly including all smaller scaffolds, 41% were
repetitive. The five main chromosomes contain 11,327 predicted mRNAs (87% of all
predicted), including 99.5% of all complete BUSCO genes. This improved genome resource
will be valuable for future evolutionary studies *of Z. indianus*, which is becoming an
increasingly problematic pest in some regions of the world (Allori Stazzonelli et al. 2023).
404
405 *Limited spatial or temporal population structure in North American Z. indianus*
406
To study spatial and temporal patterns of genetic variation in the seasonally repeated invasion
of *Z. indianus*, we resequenced ~220 individuals collected from two orchards in Virginia
(Charlottesville and Richmond) from 2017-2020, as well as one population collected from
Miami, Florida in 2019. Because temperate locations such as Virginia are thought to be
recolonized by *Z. indianus* each year (Pfeiffer et al. 2019; Rakes et al. 2023), we sampled both
early in the season (~July-August) and late in the season (~October-November) in each year
to capture the founding event, population expansion, and potential adaptation to the temperate
environment.
415
We were first interested in studying geographic and temporal variation in population
structure in North American populations of *Z. indianus*. For this analysis, we incorporated
previous sequencing data from the Western Hemisphere and Africa (Comeault et al. 2020).
While previous studies have shown limited structure within North America (Comeault et al.
2020; Comeault et al. 2021), we wanted to test for structure using deeper sampling within
introduced locations and with greater temporal resolution across the *Z. indianus* growing
season (Rakes et al. 2023). As shown previously, in an autosome-wide principal component
analysis, PC1 separated Western Hemisphere and African samples (Figure 1A; t-test: t =
78.92, df = 36, p < 2 x 10^{-16}). However, with the increased sample size of North American flies
relative to previous studies, PC2 separated North American samples into two clusters,
explaining 8% of total variation. To focus on potential structure within invasive North American
samples, we excluded the African samples and recalculated principal components. This
analysis revealed little genome-wide differentiation of North American populations collected
from different locations (Figure 1B; ANOVA P > 0.05 for PC1 and PC2), though the samples

did fall into three large groups based on PC1, which may be indicative of structural variation (Li and Ralph 2019); see below.

***Figure 1: Principal component analysis of individual Z. indianus** from this study and previous studies using autosomal SNPs.* *(A) All unrelated individuals (n=247), color coded by continent/locale of collection. (B). All unrelated North American individuals (n=190), color coded by collection site; HPO and CM are two orchards in Virginia; Northeast refers to samples from NY, NJ, and PA. (C) All unrelated individuals from Carter Mountain, Virginia (n=110), color coded by year of collection. For each analysis, only the individuals shown in the plot were included in the PC calculation.*

North American samples clustered in single-chromosome PCA for chromosomes 1, 2 and 5, but these clusters generally did not correspond to sampling locations (Figure S6; PC2 separated the two Virginia orchards for chromosome 5; Tukey P = 0.01). Interestingly, PC1 separated Florida from both Virginia orchards for the X chromosome (Tukey P < 0.01 for each comparison), suggesting some degree of genetic differentiation on the X. Visual inspection of plots of PC3 and PC4 did not indicate additional geographic population structure (results not shown). The overall lack of genome- or chromosome-wide geographic population structure suggests that there is not a high degree of genetic differentiation between eastern North American populations spread over a latitudinal transect (~1600 km) encompassing distinct climates, but some localized patterns of population structure may exist on the X chromosome. Many invasive species evolve complex population structures in the invaded range due to a combination of bottlenecks, founder effects and rapid local adaptation (Koch et al. 2020; Atsawawaranunt et al. 2023; García-Escudero et al. 2023). On the other hand, some invasive species have more homogenous populations across widespread invaded ranges in eastern North America (Friedline et al. 2019; Barrett et al. 2023). A high rate of migration between orchards (occurring naturally or due to human-mediated transport) or large founding population sizes could result in a lack of geographic differentiation between populations.

We next hypothesized that founder effects during each recolonization event might lead to unique genetic compositions of temperate populations sampled in different years (Uller and

Leimu 2011). We calculated principal components using only samples collected from Carter
Mountain, VA in 2017-2020. Surprisingly, in these samples, we saw no evidence of population
structure between years across the genome (Figure 1C; ANOVA $P > 0.05$ for PC1 and PC2) or
on individual chromosomes, except for chromosome 4, which showed subtle separation of
some years (Figure S6; Tukey $P < 0.05$ for PC1: 2018 vs. 2019 and PC2: 2017 vs. 2019).
These data suggest that the founding fly populations in Virginia are relatively homogeneous
each year at a genome-wide scale. This result is consistent with the lack of spatial population
structure and likewise could indicate large founding populations or ongoing migration.
Alternatively, the Virginia population could be permanently established with little genetic
differentiation year-to-year, though this possibility is not supported by field data (Rakes et al.,
2023).
474
We used ADMIXTURE (Alexander and Lange 2011) to test for population structure
using individuals from Africa, Florida, and the two focal Virginia orchards, calculating the most
likely number of genetic clusters for each chromosome separately. Consistent with the PCA,
the four autosomes each produced between two to four genetic groups, but there was no
apparent geographic population structure, aside from African samples mostly belonging to
different clusters from all North American samples for each chromosome (Figure 2). Notably,
for chromosomes 1 and 2, many individuals showed ~50% ancestry assignment to different
clusters, which could reflect genotypes for large structural rearrangements (see below). For the
X chromosome, using females only, we identified structure within African samples as
previously described (Comeault et al. 2020; Comeault et al. 2021) and a total of five genetic
clusters within North American populations, including one of the African genetic groups which
was found in Florida (Figure 2 third row; see orange grouping). X chromosomes have smaller
effective population sizes in species with XY sex determination systems and often experience
more extreme loss of genetic diversity upon population contraction (Ellegren 2009). The
complex population structure seen on the X chromosome may be the result of this small
population size or caused by selection on X-linked variants in different environments.
491

Figure 2: Admixture analysis of individual* Z. indianus *chromosomes from
different locations. *Each column is an individual, and colors represent*
assignment to distinct genetic clusters. The most likely number of genetic
clusters for each chromosome (k) was obtained with cross-validation analysis
and is shown at right. For chromosome 3, the X chromosome, only female flies
were used for admixture analysis, resulting in reduced sample size. FL=Miami,
Florida, VA-HPO = Richmond, VA, VA-CM = Charlottesville, VA. African
sequences represent five geographic locations and are taken from Comeault et
al. (2020 & 2021).

Structural polymorphism

The clustering of samples in the single-chromosome PCA (Figure S6), combined with many
individuals showing ~50% assignment to genetic clusters (Figure 2), suggested that large
structural variants may be segregating in *Z. indianus* (Li and Ralph 2019; Nowling et al. 2020).
Analysis of paired-end sequencing data with *smoove* provided evidence of two large
rearrangements on chromosome 1 located at 7.1 and 9.1 Mb; the genotypic combinations for
these variants largely correlate with the clustering of samples in the PCA (Figure S7; PC1
correlation with variant at 7.1Mb: P = 2 x 10^{-9}; PC1 correlation with variant at 9.1Mb: P = 3 x
10^{-5}; PC2 correlation with variant at 9.1Mb: P = 0.001). Since chromosome 1 is the longest
chromosome in our assembly, these rearrangements likely correspond to the complex *In(IV)EF*
polymorphism, made up of two overlapping inversions (Ananina et al. 2007). *smoove* did not

identify large structural variants on chromosomes 2 or 5 whose genotypes correlated to the
PCA clusters.

To look for evidence of structural variants via depressed recombination rates, we
examined linkage disequilibrium (LD) from 4,000 randomly sampled SNPs on each
chromosome. In North American samples, we discovered large blocks of LD spanning
substantial portions of chromosomes 1, 2, 3, and 5 (Figure S8), potentially indicative of
inversions (Fang et al. 2012; da Silva et al. 2019). However, there was no evidence of long-
distance LD in these regions in the African samples (Figure S8). *smoove* did not identify
inversions that corresponded to the sizes and locations of these linkage blocks. These results
support the read-based evidence of a complex rearrangement on chromosome 1 (Figure S7)
and suggest inversions on chromosomes 1, 2, 3, and 5 are segregating in North America but
are relatively rare in Africa. Given the relative chromosome sizes in the genome assembly, the
linkage blocks on chromosome 2 and 5 likely correspond to *In(V)B* and *In(II)*A, respectively
(Ananina et al. 2007). The X chromosome has three described inversions in *Z. indianus*
(Ananina et al. 2007), which may explain to the complex pattern of linkage observed in North
American samples and the population structure observed for the X chromosome within North
America (Figure 2). Major chromosomal polymorphisms are known to be important for local
adaptation and phenotypic divergence in a wide variety of species (Joron et al. 2011; Küpper
et al. 2016; Lee et al. 2016; Huang et al. 2020; Nunez et al. 2024), including inversions that
facilitate invasive phenotypes (Galludo et al. 2018; Tepolt and Palumbi 2020; Tepolt et al.
2022; Ma et al. 2024). These inversions may have been present at low frequency in the
bottlenecked population that founded *Z. indianus* populations in the Western Hemisphere, but
then experienced subsequent selection in the invaded range. Alternatively, these
polymorphisms may have arisen in a currently undescribed population and then been
introduced to the Western Hemisphere.

Recolonization, bottlenecks and seasonal dynamics in Z. indianus

Invasive species typically experience a genetic bottleneck due to small founding population
sizes (Barrett 2015; Estoup et al. 2016). We hypothesized that North American populations
would show reduced effective population size (N_e) relative to African populations, and that
Virginia populations would show a further, more recent reduction in N_e relative to Florida
populations as the result of a secondary population bottleneck upon temperate recolonization.
Our prediction was correct with respect to Africa vs North America: African populations show
historical fluctuations but population sizes typically in the range of ~ 10^5-10^7 individuals.
Interestingly, introduced populations in North America demonstrate population sizes that
increased, decreased, then increased again in the past ~500 years. Comeault et al. (2021)
suggested that introduced populations in the Americas are derived from a historically admixed
population composed of both East African and West African flies, and the historic expansion of
introduced populations might correspond to this admixture event. The subsequent drop in
population size to 10^4-10^5 may then reflect a bottleneck following colonization of Brazil in the
late 1990s (Yassin et al. 2008), followed by a rebound as introduced populations expanded.

559
560
***Figure 3: Demographic effects of bottlenecks in* Z. indianus *populations** A)*
Population history reconstruction with smc++ *using autosomal genotypes.*
Introduced-Florida flies were collected in Miami in 2019. Introduced-Virginia flies
were collected in the early-mid season (June-September) from two Virginia
orchards in 2017-2020 (n=5 populations grouped by orchard and year). Native
populations are distinct African populations (Kenya, Zambia, Senegal-Forest,
Senegal-Desert, and Sao Tome [Comeault et al 2020]). B) Kinship and
probability of zero identity by descent for pairs of individual flies from the same
collection location and season within North America calculated with autosomal
SNPs. C) Kinship coefficients for pairs of individual flies collected at Carter
Mountain Orchard, Virginia, as a function of the number of days between
sampling. Relatedness was assigned according to thresholds from (Thornton et
al. 2012).
574

Overall, the ancestral population sizes for Virginia and Florida were quite similar, and
our prediction of reduced recent population sizes in Virginia relative to Florida was not well-
supported. The minimum population sizes for Florida and Virginia (10^4-10^5) are larger than
expected for a single small colonization event. Field data suggest founding populations in
orchards are small and then rapidly expand (Rakes et al. 2023), suggesting that these large
population sizes could be caused by ongoing gene flow from the source population after
colonization, which is consistent with the lack of temporal population structure. Given our
limited sample sizes and potential differences in the number of generations per year in

temperate and subtropical environments, detecting fine-scale differences in very recent
population fluctuations may be beyond the detection ability of the software; *smc++* becomes
less accurate at timescales less than ~133 generations (Patton et al. 2019). Alternatively, the
Virginia populations may be admixed populations reflecting individuals from multiple sources,
producing larger effective population sizes than would otherwise be expected if recolonization
occurs from a single source population undergoing a bottleneck. Admixture and gene flow ae
important factors fueling genetic diversity and invasiveness in introduced species
(McGaughran et al. 2024) and could potentially contribute to *Z. indianus'* local success
following each recolonization event.

We additionally tested for bottlenecks by looking for inbreeding, which might be a
product of small founding populations. Using two measures of genetic similarity, we discovered
many pairs of related flies in our dataset (Figure 3B). Most dramatically, many flies collected in
2018 appeared to be close relatives (Figure S9). In collections from late July and early August
2018, 26 pairs of close relatives involving 13 individual flies were collected. Of those, 21 pairs
of relatives were collected on different days, suggesting the relatedness was not solely a
sampling artifact due to collecting closely related flies in the same microhabitat of the orchard.
The effect of this apparent bottleneck was sometimes retained throughout the growing season,
as a pair of full sibs was sampled 77 days apart in 2018, two pairs of second-degree relatives
were sampled over 110 days apart in 2018, and two pairs of third-degree relatives were
sampled 140 days apart in 2017 (Figure 3C). Given that *Z. indianus* are collected in small
numbers early in the season (Rakes et al. 2023) and 2017 and 2018 had particularly early
captures (Table S1), we suggest small founding population size followed by inbreeding could
produce individuals sampled distantly in time that still show close genetic similarity.
Alternatively, flies may live for a relatively long time or have slower generations in the wild,
allowing us to capture close relatives separated by longer time periods. However, we note that
the same pattern was not seen in every year of our collections, suggesting that colonization
dynamics might differ dramatically from year to year, which is expected if recolonization occurs
due to chance events each year.

The founder effect could generate temporal population structure by creating populations
that were more similar within a year than between years, creating a positive relationship
between F_{ST} and the elapsed time between collections (Bergland et al. 2014). We tested this
prediction with samples collected from Carter Mountain, Virginia over four years, and there
was no relationship between F_{ST} and the time between sampling (linear model, df=17, P=0.9,
Figure S10). This lack of temporal differentiation is consistent with the PCA and the relatively
large minimum population sizes previously described and could be produced by ongoing gene
flow that eliminates any signal of a founder effect and inbreeding. This finding is distinct from
trends observed in *D. melanogaster,* which experiences a strong overwintering bottleneck and
shows temporal patterns of differentiation (Bergland et al. 2014; Nunez et al. 2024).

Repeated differentiation between Florida and Virginia populations

Despite the lack of genome-wide differentiation between different North American locales, we
were interested in testing whether specific regions of the genome might differ between
populations given environmental differences: Virginia has a temperate, seasonal climate with a
relatively limited variety cultivated produce, and southern Florida is subtropical with an
abundance and diversity of fruits throughout the year. Other factors such as diseases,
insecticide use, and competing species may also differ widely between locales. In the absence
of genome-wide population structure, genomic regions differentiated between these locations
are candidates for local adaptation. We conducted a SNP-level F_{ST} analysis comparing all flies
collected in Florida to those collected in the early season in Virginia over four years. We
observed elevated F_{ST} throughout much of the X chromosome, with a pronounced peak at 690
kb (Figure 4A). This peak was observed when comparing the Florida collection to Virginia
collections from both Charlottesville and Richmond across all four years of Virginia sampling
both early and late in the season (Figure S11), suggesting that this differentiation is maintained
through recurrent rounds of recolonization, potentially via local adaptation. Alternatively, this
region could correspond to alleles that directly promote dispersal and/or invasion (Weinig et
al. 2007) and are found at higher frequency in invaded populations. One limitation of our
sampling strategy is that we have only a single year of sampling in Florida; additional data will
be needed to determine whether the genetic composition of this population (and differentiation
from Virginia) remains steady across multiple years. However, assuming that this result is not
an artifact related to the Florida sample, an alternative possible explanation for the repeated
differentiation seen between Florida and Virginia is that Virginia is recolonized by a source
population that is genetically distinct from the southern Florida population we sampled here.
Regardless, the finding implies localized genetic structure across a latitude gradient in North
American.
650
651

652
Figure 4: Signals of selection in temperate Z. indianus populations. *A)*
Genome-wide SNP-level F_{ST} comparing individual flies sampled in Florida (n=26)
to all flies sampled in the early season in Virginia (n=123), color-coded by
chromosome. Only females were used for the X chromosome (chromosome 3,
green). B) Integrated haplotype homozygosity score (IHS) using all flies collected
in Virginia. C) Zoomed in view of SNP-level FST between Florida and Virginia on
chr 3: 0-2Mb. D) IHS for the X chromosome (chromosome 3:0-2 Mb) calculated
separately for flies from Africa, Florida, all North America, and Virginia. Dashed
line indicates IHS = 5 to facilitate comparisons between populations. E-F)
Extended haplotype homozygosity for the two alleles of the SNP with highest F_{ST}
(E; chr 3: 689841) and highest IHS (F; chr 3: 973443), calculated using all
haplotypes from Virginia.

665
666 *Genomic signals of differentiation and selection*

667
The elevated F_{ST} seen on the X chromosome raised the intriguing possibility that some genetic
variation could potentially be under selection in temperate environments (Virginia) relative to
subtropical Florida. We phased the paired-end sequencing data and calculated extended
haplotype homozygosity (EHH) and IHS (integrated haplotype homozygosity score) using all
Virginia individuals to look for long, shared haplotypes that can be signatures of selective
sweeps (Sabeti et al. 2007). As in the F_{ST} analysis, we observed a region on chromosome 3

that stood out in this analysis with many SNPs with IHS > 5; this region overlapped with the
F_{ST} peak (Figure 4, B-D). The peak F_{ST} SNP was approximately 300 kb away from the peak
IHS SNP. We then repeated the IHS analysis using flies from Africa, Florida, and all North
America (Virginia + Florida + Comeault (2020) locations) to determine whether this signature
was unique to temperate populations. There was no signal of elevated IHS in African flies
(Figure 4D, 1st row), suggesting this selective signature is unique to invasive populations.
Further, this region showed a less substantial IHS peak when analyzing flies collected in
Florida (Figure 4D, 2nd row) but was prominent when examining Virginia flies or all North
American flies (Figure 4D, 3rd-4th rows), suggesting the signal of the selective sweep is
primarily driven by individuals collected in temperate environments. Both the peak IHS SNP
and the peak F_{ST} SNP showed evidence of long extended haplotypes characteristic of sweeps
(Figure 4E-F). These results further support the possibility that this locus is advantageous to
invasive potential or survival in temperate habitats.
687
We investigated this region of the genome by examining linkage disequilibrium (LD) and
haplotype structure of the 400 SNPs with a Virginia IHS > 5 (Figure 5A). We discovered this
region spanning ~700 kb has several large haplotype blocks in temperate North American
samples (Figure 5C-D) and in Florida (Figure S12B), but these same haplotypes are not found
in Africa (Figure 5C, Figure S12A), suggesting they are unique to introduced populations. In
invasive copepods, haplotypes under selection in the invasive range are ancestral
polymorphisms under balancing selection in the native range (Stern and Lee 2020). A similar
situation was found for a balanced inversion polymorphism that fuels invasion in invasive crabs
(Tepolt and Palumbi 2020; Tepolt et al. 2022). However, ancestral polymorphism selected in
the invaded range does not appear to be the case in *Z. indianus*, as the haplotypes from North
America were not found in any African flies. These novel haplotypes could be new mutations
or derived due to hybridization/introgression from another species or divergent population;
hybridization can be an important evolutionary force in invasive species (Ellstrand and
Schierenbeck 2000; Fournier and Aron 2021). The *Zaprionus* genus shows signals of historic
introgression, though *Z. indianus* was not directly implicated in a previous analysis (Suvorov et
al. 2022). Therefore, two major haplotypes not found in Africa contribute to the differentiation of
Florida and Virginia populations, though the source of these haplotypes remains to be
determined. Though we focus on one genomic region, we note that most of the X chromosome
shows elevated IHS scores (Figure 3B), and many SNPs on the X show $F_{ST} > 0.25$ between
Virginia and Florida (Figure 3A). This observation is in line with the findings of Comeault et al
2021, who showed that many X-linked scaffolds showed signs of selection in invasive
populations and is likely related to the presence of several inversions on this chromosome. We
also note that our approach would not detect sweeps involving multiple alleles from standing
variation (soft sweeps; (Messer and Petrov 2013; Garud et al. 2015)), which could be an
important potential component of *Z. indianus* evolution given the high levels of genetic diversity
found even in invasive populations (Avalos et al. 2017).
714

Figure 5: Major haplotypes on the X chromosome with signals of selection and differentiation. *Only SNPs with IHS > 5 in Virginia (n=400) are shown in this figure for clarity; scale at top shows physical positions of SNPs, which are equally spaced in panels A-D. A) FST of individual SNPs comparing Florida and*

     Virginia populations (see Figure 4). B) IHS for individual SNPs. C) Haplotypes:
722   *each horizontal row shows genotypes for a single haploid chromosome phased*
723   *with read-backed phasing. Blue indicates the allele more common in African*
724   *populations and green is the other allele. Missing genotypes are shown in gray.*
725   *D) LD (R^2) for these SNPs in all North American flies, excluding Florida.*
726

To explore population genetic signals around this highly divergent region of the X
chromosome, we broadly grouped flies into three populations: Africa, Florida, and Virginia and
calculated population genetic statistics in 5 kb non-overlapping windows for all females. This
analysis confirmed two regions of relatively high F_{ST} between Florida and Virginia, though they
are separated by a region of nearly zero differentiation within North America, as measured by
F_{ST}, D_{xy}, and nucleotide diversity (Figure 6A-C, ~700-800kb). Virginia and Florida are both
highly differentiated from Africa in this region, and it has negative Tajima's D in both Florida
and Virginia (Figure 6D), potentially indicating recovery from a selective sweep in North
America. The region of no divergence may represent a selective sweep of a haplotype that
existed on two different genetic backgrounds that were subsequently favored in Florida and
Virginia, producing a high degree of genetic differentiation in the surrounding sequences. The
region with a potential sweep in North America contains ~6 genes, including the gene *yin/opt1*,
which is important for absorption of dietary peptides in *D. melanogaster* (Roman et al. 1998).
Allelic differences between African and invasive range flies in this gene could be involved in
adaptation to new diets in new environments.
742
743

Figure 6: Population genetic statistics for the region surrounding a

selected haplotypes on chromosome 3 (250-1,1000 kb). *All statistics were*

calculated for 5 kb, non-overlapping windows. Black points at the top indicate the

locations of the 400 SNPs shown in Figure 5. A) F_{ST} comparing combinations of

flies from Africa, Virginia (both focal orchards combined) and Florida. B) Absolute

nucleotide divergence (D_{xy}) for the same comparisons. C) Nucleotide diversity

(π) for each population. D) Tajima's D for the three populations. E) Average

sequencing depth per window relative to the mean depth for the entire

chromosome. Relative depths were averaged for all individuals in each

population. See Figure S14 for whole-genome analysis of the same statistics.

To confirm the patterns described above were not driven by genome assembly issues, we
also examined the normalized depth of sequencing coverage relative to the chromosome
average and discovered a region of variable coverage that overlaps the region of high F_{ST}
between Florida and Virginia and is immediately adjacent to the region with zero divergence
between Florida and Virginia (Figure 6E, Figure S13). This region (~600kb-700kb) has low
coverage in Africa. In Florida and Virginia coverage varies from 0.5X -1X throughout the
region. *smoove* identified paired-end evidence for a 52 kb duplication in this region, but
genotype calls showed frequencies were similar in Virginia, Florida, and Africa. Combined with
the sequencing depth data, these findings suggest copy number variation of these loci might
contribute to the Florida-Virginia divergence, though long-read sequencing will likely be
required to resolve the sequence variation. The region of elevated FST between Florida and
Virginia and variable copy number contains several genes with neuronal and metabolic
functions, offering exciting possibilities for future studies of the potential functional basis of this
geographic divergence.
769
For comparison, we also examined the same five population signals genome-wide
(Figure S14) and observed that the X chromosome is an outlier in many regards. Divergence
between Africa and North American samples is greater on the X chromosome (Figure S14A).
As previously described (Comeault et al. 2020; Comeault et al. 2021), the X chromosome has
reduced genetic diversity relative to autosomes, especially in invaded populations (Figure
S14B-C). Tajima's D is negative across the genome for African flies and mostly positive in
North American autosomes, indicative of a strong bottleneck in North American flies. However,
Tajima's D fluctuates between strongly positive and strongly negative in North American
populations along the X chromosome (Figure S14D). This finding, combined with complex
patterns of genetic ancestry on the X (Figure 2) and many regions with high haplotype
homozygosity (Figure 4), suggest complex evolutionary dynamics on the X that warrant further
investigation. These findings agree with Comeault et al. 2021, who found that regions under
selection on the X chromosome typically showed higher divergence between invasive
populations and African populations. Further global sampling and sequencing of X
chromosomes with long reads to resolve inversion genotypes and CNVs may offer insight
towards the role of X-linked genes in fueling the ongoing invasion of *Z. indianus*.
786
787
788   **Conclusions**
789
In addition to posing economic, health, and environmental threats, invasive species also
serve as outstanding models for studying rapid evolution in new environments. Here we report
an improved genome assembly and annotation for *Z. indianus*, an introduced drosophilid that
is thought to repeatedly recolonize temperate environments each year and is a potential crop
pest. We use it for a preliminary assessment of potential rapid evolution and genetic variation
in the early stages of invasion. We show that recolonization is likely a stochastic process
resulting in different evolutionary dynamics in different years, even within a single orchard. This
finding demonstrates broad sampling is important for invasive species that are repeatedly

introduced or have multiple introduced populations that may undergo different evolutionary
trajectories in different years or different locations. While some founding populations may be
small, several population genetic patterns we observe could be explained by ongoing gene
flow with the source population, or between temperate populations following recolonization,
suggesting gene flow that spreads and maintains favorable alleles could be an important
component in *Z. indianus*'s widespread success, as it is for many invasive species (Díez-del-
Molino et al. 2013; Medley et al. 2015; Arredondo et al. 2018). Demographic simulations and
additional whole genome data will be required to better describe the recent histories of and
potential gene flow between invasive populations and to infer colonization routes within North
America.
808
Though we find limited population structure across space or time in introduced North
American populations, we find a region on the X chromosome that may have experienced a
selective sweep in North America followed by separate sweeps in Virginia and Florida.
Studying how genetic variation in this region of the genome influences survival in temperate
environments will be an important direction of future research. We additionally find that the X
chromosome has an unusually complex evolutionary history in *Z. indianus*. It may have several
segregating inversions and CNVs, has strong signatures of selection, and shows regions of
high divergence both between African and North American populations and within North
America. Specifically, long-read sequencing strategies will be important to understand likely
inversions both on the X and throughout the *Z. indianus* genome that are common in the
invaded range. Large inversions can link together adaptive alleles and are often important
drivers of evolution in rapidly changing environments (Thompson and Jiggins 2014), so these
regions will be important to track over larger spatial and temporal scales in future studies.
822
These results underscore the complexity of genetic dynamics during invasions and the
need for further studies to explore the adaptive potential and ecological impacts of *Z. indianus*
in its invasive range. *Z. indianus* provides a unique system in which we can study independent
invasion events across multiple years and locations. One limitation of our study is sample size
for each year and location: our ability to estimate allele frequencies or detect subtle changes in
allele frequencies across time or space is limited. Sampling strategies that incorporate more
individuals, such as pooled sequencing (Bergland et al. 2014; Kapun et al. 2021; Machado et
al. 2021; Nunez et al. 2024), will be required to detect these more subtle changes, if they
occur, and to understand how they may contribute to rapid adaptation to new environments.
The recurrent nature of *Z. indianus* colonization may also offer insight towards the predictability
of rapid evolution of invasive species.
834
835 **Data Availability**
836
New individual sequencing data has been deposited in the SRA under project number #
PRJNA991922. RNA sequencing from larval and pupal samples, and larval Hi-C data used for
scaffolding are deposited under the same project number. The genome sequence has been
deposited at DDBJ/ENA/GenBank under the accession JAUIZU000000000. The metadata for

all sequencing samples (including date and location of collection); the annotation information
for transcripts, proteins and repeats; and VCFs of SNPs and structural variants have been
deposited to Dryad: https://doi.org/10.5061/dryad.q2bvq83v3. All code to reproduce analyses
has been deposited to Zenodo via Dryad. All code for analysis is also available at:
https://github.com/ericksonp/Z.indianus_individual_sequencing/tree/main
846
847 Temporary reviewer dryad link:
848 http://datadryad.org/stash/share/6td1mtLMrbgLL6IgyaEtpmqK4cigV0Vl2Hhqw9Aspvo
849

850 **Acknowledgments**
851
The authors acknowledge The University of Richmond's High Performance Computer
(https://data.richmond.edu/About-HPC-at-UR/index.html) for providing computational
resources that contributed to the results reported herein. We particularly thank George
Flanagin for technical support. Preliminary analyses were conducted using resources provided
by Research Computing at The University of Virginia (https://rc.virginia.edu). We thank the
owners and managers of Carter Mountain Orchard and Hanover Peach Orchard for graciously
allowing us to collect flies on their properties.
859

860 **Funding**
861
862 This work was funded by award #61-1673 from the Jane Coffin Childs Memorial Fund for
863 Medical Research (to PAE), NIH NIGMS award # R15GM146208 (to PAE), NSF BIO-DEB
864 (EP) award # 2145688 (to AOB), NIH NIGMS award # R35GM119686 (to AOB), and startup
865 funds from the University of Richmond to PAE.
866

867 **Author contributions**
868
869 PAE, AB, AOB: conceptualization; PAE, AOB: funding; PAE, AG: investigation; PAE, AB, NP:
870 resources; PAE: methodology, formal analysis, visualization, writing-original draft; PAE, AOB:
871 writing-reviewing and editing
872

873
References
875

Alexander DH, Lange K. 2011. Enhancements to the ADMIXTURE algorithm for individual
ancestry estimation. *BMC Bioinformatics* [Internet] 12:246. Available from:
https://doi.org/10.1186/1471-2105-12-246

Allori Stazzonelli E, Funes CF, Corral Gonzalez MN, Gibilisco SM, Kirschbaum DS. 2023.
Population fluctuation and infestation levels of Zaprionus indianus Gupta (Diptera:
Drosophilidae) in berry crops of northwestern Argentina | International Society for
Horticultural Science. *Acta Horticultura* [Internet]. Available from:
http://www.actahort.org/books/1381/1381_19.htm

Altizer S, Ostfeld RS, Johnson PTJ, Kutz S, Harvell CD. 2013. Climate Change and Infectious
Diseases: From Evidence to a Predictive Framework. *Science* 341:514–519.

Ananina G, Rohde C, David JR, Valente VLS, Klaczko LB. 2007. Inversion polymorphism and
a new polytene chromosome map of Zaprionus indianus Gupta (1970) (Diptera:
Drosophilidae). *Genetica* 131:117–125.

Araripe LO, Klaczko LB, Moreteau B, David JR. 2004. Male sterility thresholds in a tropical
cosmopolitan drosophilid, Zaprionus indianus. *Journal of Thermal Biology* [Internet]
29:73–80. Available from:
http://www.sciencedirect.com/science/article/pii/S0306456503000950

Arredondo TM, Marchini GL, Cruzan MB. 2018. Evidence for human-mediated range
expansion and gene flow in an invasive grass. *Proceedings of the Royal Society B:
Biological Sciences* [Internet] 285:20181125. Available from:
https://royalsocietypublishing.org/doi/full/10.1098/rspb.2018.1125

Atsawawaranunt K, Ewart KM, Major RE, Johnson RN, Santure AW, Whibley A. 2023. Tracing
the introduction of the invasive common myna using population genomics. *Heredity*
[Internet] 131:56–67. Available from: https://www.nature.com/articles/s41437-023-
00621-w

Avalos A, Pan H, Li C, Acevedo-Gonzalez JP, Rendon G, Fields CJ, Brown PJ, Giray T,
Robinson GE, Hudson ME, et al. 2017. A soft selective sweep during rapid evolution of
gentle behaviour in an Africanized honeybee. *Nat Commun* [Internet] 8:1550. Available
from: https://www.nature.com/articles/s41467-017-01800-0

Bao Z, Eddy SR. 2002. Automated De Novo Identification of Repeat Sequence Families in
Sequenced Genomes. *Genome Res.* [Internet] 12:1269–1276. Available from:
https://genome.cshlp.org/content/12/8/1269

Barrett CF, Corbett CW, Thixton-Nolan HL. 2023. A lack of population structure characterizes
the invasive Lonicera japonica in West Virginia and across eastern North America1,2.
tbot [Internet] 150:455–466. Available from: https://bioone.org/journals/the-journal-of-
the-torrey-botanical-society/volume-150/issue-3/TORREY-D-23-00007.1/A-lack-of-
population-structure-characterizes-the-invasive-Lonicera-japonica/10.3159/TORREY-D-
23-00007.1.full

Barrett RDH, Laurent S, Mallarino R, Pfeifer SP, Xu CCY, Foll M, Wakamatsu K, Duke-Cohan
JS, Jensen JD, Hoekstra HE. 2019. Linking a mutation to survival in wild mice. *Science*
363:499–504.

Barrett SCH. 2015. Foundations of invasion genetics: the Baker and Stebbins legacy.
Molecular Ecology [Internet] 24:1927–1941. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.13014

Bellard C, Bertelsmeier C, Leadley P, Thuiller W, Courchamp F. 2012. Impacts of climate
change on the future of biodiversity. *Ecology Letters* 15:365–377.

Bemmels JB, Mikkelsen EK, Haddrath O, Colbourne RM, Robertson HA, Weir JT. 2021.
Demographic decline and lineage-specific adaptations characterize New Zealand kiwi.
Proc Biol Sci 288:20212362.

Bergland AO, Behrman EL, O'Brien KR, Schmidt PS, Petrov DA. 2014. Genomic Evidence of
Rapid and Stable Adaptive Oscillations over Seasonal Time Scales in Drosophila. *PLoS*
Genet [Internet] 10:e1004775. Available from:
http://dx.doi.org/10.1371/journal.pgen.1004775

Bushnell B, Rood J, Singer E. 2017. BBMerge – Accurate paired shotgun read merging via
overlap. *PLOS ONE* 12:e0185056.

Cabanettes F, Klopp C. 2018. D-GENIES: dot plot large genomes in an interactive, efficient
and simple way. *PeerJ* [Internet] 6:e4958. Available from: https://peerj.com/articles/4958

Campos SRC, Rieger TT, Santos JF. 2007. Homology of polytene elements between
Drosophila and Zaprionus determined by in situ hybridization in Zaprionus indianus.
Genet Mol Res 6:262–276.

Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, Holt C, Alvarado AS, Yandell M.
2008. MAKER: An easy-to-use annotation pipeline designed for emerging model
organism genomes. *Genome Res.* [Internet] 18:188–196. Available from:
https://genome.cshlp.org/content/18/1/188

Chan PP, Lowe TM. 2019. tRNAscan-SE: Searching for tRNA genes in genomic sequences.
Methods Mol Biol [Internet] 1962:1–14. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6768409/

Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. 2015. Second-generation
PLINK: rising to the challenge of larger and richer datasets. *Gigascience* 4:7.

Clements DR, Ditommaso A. 2011. Climate change and weed adaptation: can evolution of
invasive plants lead to greater range expansion than forecasted? *Weed Research*
51:227–240.

Comeault AA, Kautt AF, Matute DR. 2021. Genomic signatures of admixture and selection are
shared among populations of Zaprionus indianus across the western hemisphere.
Molecular Ecology 30:6193–6210.

Comeault AA, Wang J, Tittes S, Isbell K, Ingley S, Hurlbert AH, Matute DR. 2020. Genetic
Diversity and Thermal Performance in Invasive and Native Populations of African Fig
Flies. *Molecular Biology and Evolution* 37:1893–1906.

Commar LS, Galego LG da C, Ceron CR, Carareto CMA. 2012. Taxonomic and evolutionary
analysis of Zaprionus indianus and its colonization of Palearctic and Neotropical
regions. *Genet Mol Biol* 35:395–406.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics*
[Internet] 27:2156–2158. Available from: https://doi.org/10.1093/bioinformatics/btr330

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools.
Gigascience 10:giab008.

Díez-del-Molino D, Carmona-Catot G, Araguas R-M, Vidal O, Sanz N, García-Berthou E,
García-Marín J-L. 2013. Gene Flow and Maintenance of Genetic Diversity in Invasive
Mosquitofish (Gambusia holbrooki). *PLOS ONE* [Internet] 8:e82501. Available from:
https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0082501

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* [Internet]
29:15–21. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3530905/

Dowle M, Srinivasan A. 2019. data.table: Extension of `data.frame`. Available from:
https://CRAN.R-project.org/package=data.table

Ellegren H. 2009. The different levels of genetic diversity in sex chromosomes and autosomes.
Trends in Genetics [Internet] 25:278–284. Available from:
https://www.sciencedirect.com/science/article/pii/S0168952509000900

Ellstrand NC, Schierenbeck KA. 2000. Hybridization as a stimulus for the evolution of
invasiveness in plants? *Proceedings of the National Academy of Sciences* [Internet]
97:7043–7050. Available from: https://www.pnas.org/doi/abs/10.1073/pnas.97.13.7043

Erickson PA, Weller CA, Song DY, Bangerter AS, Schmidt P, Bergland AO. 2020. Unique
genetic signatures of local adaptation over space and time for diapause, an ecologically
relevant complex trait, in Drosophila melanogaster. *PLOS Genetics* 16:e1009110.

Estoup A, Ravigné V, Hufbauer R, Vitalis R, Gautier M, Facon B. 2016. Is There a Genetic
Paradox of Biological Invasion? *https://doi.org/10.1146/annurev-ecolsys-121415-*
032116 [Internet]. Available from:
https://www.annualreviews.org/doi/abs/10.1146/annurev-ecolsys-121415-032116

Fang Z, Pyhäjärvi T, Weber AL, Dawe RK, Glaubitz JC, González J de JS, Ross-Ibarra C,
Doebley J, Morrell PL, Ross-Ibarra J. 2012. Megabase-Scale Inversion Polymorphism in
the Wild Ancestor of Maize. *Genetics* [Internet] 191:883–894. Available from:
https://doi.org/10.1534/genetics.112.138578

Feron R, Waterhouse RM. 2022. Assessing species coverage and assembly quality of rapidly
accumulating sequenced genomes. *GigaScience* [Internet] 11:giac006. Available from:
https://doi.org/10.1093/gigascience/giac006

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. 2020.
RepeatModeler2 for automated genomic discovery of transposable element families.
Proceedings of the National Academy of Sciences [Internet] 117:9451–9457. Available
from: https://www.pnas.org/doi/full/10.1073/pnas.1921046117

Fournier D, Aron S. 2021. Hybridization and invasiveness in social insects — The good, the
bad and the hybrid. *Current Opinion in Insect Science* [Internet] 46:1–9. Available from:
https://www.sciencedirect.com/science/article/pii/S2214574521000018

Friedline CJ, Faske TM, Lind BM, Hobson EM, Parry D, Dyer RJ, Johnson DM, Thompson LM,
Grayson KL, Eckert AJ. 2019. Evolutionary genomics of gypsy moth populations
sampled along a latitudinal gradient. *Molecular Ecology* [Internet] 0. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.15069

Galludo M, Canals J, Pineda-Cirera L, Esteve C, Rosselló M, Balanyà J, Arenas C, Mestres F.
2018. Climatic adaptation of chromosomal inversions in Drosophila subobscura.
Genetica [Internet] 146:433–441. Available from: https://doi.org/10.1007/s10709-018-
0035-x

García-Escudero CA, Tsigenopoulos CS, Manousaki T, Tsakogiannis A, Marbà N, Vizzini S,
Duarte CM, Apostolaki ET. 2023. Population genomics unveils the century-old invasion
of the Seagrass Halophila stipulacea in the Mediterranean Sea. *Mar Biol* [Internet]
171:40. Available from: https://doi.org/10.1007/s00227-023-04361-7

Garnier S. 2018. viridis: Default Color Maps from "matplotlib." Available from: https://CRAN.R-
project.org/package=viridis

Garud NR, Messer PW, Buzbas EO, Petrov DA. 2015. Recent Selective Sweeps in North
American Drosophila melanogaster Show Signatures of Soft Sweeps. *PLOS Genetics*
[Internet] 11:e1005004. Available from:
https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1005004

Gautier M, Vitalis R. 2012. rehh: an R package to detect footprints of selection in genome-wide
SNP data from haplotype structure. *Bioinformatics* 28:1176–1177.

Gleason JM, Roy PR, Everman ER, Gleason TC, Morgan TJ. 2019. Phenology of Drosophila
species across a temperate growing season and implications for behavior. *PLOS ONE*
14:e0216601.

Good BH, McDonald MJ, Barrick JE, Lenski RE, Desai MM. 2017. The dynamics of molecular
evolution over 60,000 generations. *Nature* 551:45–50.

Gupta JP. 1970. Description of a new species of Phorticella zaprionus (Drosophilidae) from
India. *Proceedings of the Indian National Science Academy* 36B:62–70.

Gupta JP, Kumar A. 1987. Cytogenetics of Zaprionus indianus Gupta (Diptera: Drosophilidae):
Nucleolar organizer regions, mitotic and polytene chromosomes and inversion

polymorphism. *Genetica* [Internet] 74:19–25. Available from: https://doi.org/10.1007/BF00055090

Hancock AM, Brachi B, Faure N, Horton MW, Jarymowycz LB, Sperone FG, Toomajian C, Roux F, Bergelson J. 2011. Adaptation to Climate Across the Arabidopsis thaliana Genome. *Science* 334:83–86.

Hoberg EP, Brooks DR. 2015. Evolution in action: climate change, biodiversity dynamics and emerging infectious disease. *Philosophical Transactions of the Royal Society B: Biological Sciences* 370:20130553.

Holle SG, Tran AK, Burkness EC, Ebbenga DN, Hutchison WD. 2018. First Detections of Zaprionus indianus (Diptera: Drosophilidae) in Minnesota. *ents* 54:99–102.

Huang K, Andrew RL, Owens GL, Ostevik KL, Rieseberg LH. 2020. Multiple chromosomal inversions contribute to adaptive divergence of a dune sunflower ecotype. *Molecular Ecology* [Internet] 29:2535–2549. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.15428

Johnson MS, Gopalakrishnan S, Goyal J, Dillingham ME, Bakerlee CW, Humphrey PT, Jagdish T, Jerison ER, Kosheleva K, Lawrence KR, et al. 2021. Phenotypic and molecular evolution across 10,000 generations in laboratory budding yeast populations.Verstrepen KJ, Wittkopp PJ, Verstrepen KJ, Hodgins-Davis A, editors. *eLife* 10:e63910.

Jones MR, Mills LS, Alves PC, Callahan CM, Alves JM, Lafferty DJR, Jiggins FM, Jensen JD, Melo-Ferreira J, Good JM. 2018. Adaptive introgression underlies polymorphic seasonal camouflage in snowshoe hares. *Science* 360:1355–1358.

Joron M, Frezal L, Jones RT, Chamberlain NL, Lee SF, Haag CR, Whibley A, Becuwe M, Baxter SW, Ferguson L, et al. 2011. Chromosomal rearrangements maintain a polymorphic supergene controlling butterfly mimicry. *Nature* [Internet] 477:203–206. Available from: http://www.nature.com/nature/journal/v477/n7363/full/nature10341.html

Joshi NK, Biddinger DJ, Demchak K, Deppen A. 2014. First report of Zaprionus indianus (Diptera: Drosophilidae) in commercial fruits and vegetables in Pennsylvania. *J. Insect Sci.* 14:259.

Kapun M, Nunez JCB, Bogaerts-Márquez M, Murga-Moreno J, Paris M, Outten J, Coronado-Zamora M, Tern C, Rota-Stabelli O, Guerreiro MPG, et al. 2021. Drosophila Evolution over Space and Time (DEST) - A New Population Genomics Resource. *bioRxiv* [Internet]:2021.02.01.428994. Available from: https://www.biorxiv.org/content/10.1101/2021.02.01.428994v1

Keller O, Kollmar M, Stanke M, Waack S. 2011. A novel hybrid gene prediction method employing protein multiple sequence alignments. *Bioinformatics* [Internet] 27:757–763. Available from: https://doi.org/10.1093/bioinformatics/btr010

Kim BY, Wang JR, Miller DE, Barmina O, Delaney E, Thompson A, Comeault AA, Peede D, D'Agostino ER, Pelaez J, et al. 2021. Highly contiguous assemblies of 101 drosophilid

genomes.Coop G, Wittkopp PJ, Sackton TB, editors. *eLife* [Internet] 10:e66405. Available from: https://doi.org/10.7554/eLife.66405

Koch JB, Dupuis JR, Jardeleza M-K, Ouedraogo N, Geib SM, Follett PA, Price DK. 2020. Population genomic and phenotype diversity of invasive Drosophila suzukii in Hawaiʻi. *Biol Invasions* [Internet] 22:1753–1770. Available from: https://doi.org/10.1007/s10530-020-02217-5

Korf I. 2004. Gene finding in novel genomes. *BMC Bioinformatics* [Internet] 5:59. Available from: https://doi.org/10.1186/1471-2105-5-59

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: Analysis of Next Generation Sequencing Data. *BMC Bioinformatics* [Internet] 15:356. Available from: https://doi.org/10.1186/s12859-014-0356-4

Korunes KL, Samuk K. 2021. pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of missing data. *Molecular Ecology Resources* [Internet] 21:1359–1368. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/1755-0998.13326

Kremmer L, David J, Borowiec N, Thaon M, Ris N, Poirie M, Gatti J-L. 2017. The African fig fly Zaprionus indianus: a new invasive pest in France? *Bulletin of Insectology* 70:57–62.

Küpper C, Stocks M, Risse JE, dos Remedios N, Farrell LL, McRae SB, Morgan TC, Karlionova N, Pinchuk P, Verkuil YI, et al. 2016. A supergene determines highly divergent male reproductive morphs in the ruff. *Nat Genet* [Internet] 48:79–83. Available from: http://www.nature.com/ng/journal/v48/n1/full/ng.3443.html

Leão BFD, Tldon R. 2004. Newly invading species exploiting native host-plants: the case of the African Zaprionus indianus (Gupta) in the Brazilian Cerrado (Diptera, Drosophilidae). *Annales de la Société entomologique de France (N.S.)* 40:285–290.

Lee YW, Fishman L, Kelly JK, Willis JH. 2016. A Segregating Inversion Generates Fitness Variation in Yellow Monkeyflower (Mimulus guttatus). *Genetics* [Internet] 202:1473–1484. Available from: http://www.genetics.org/content/202/4/1473

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* [Internet] 25:2078–2079. Available from: https://doi.org/10.1093/bioinformatics/btp352

Li H, Peng Y, Wang Y, Summerhays B, Shu X, Vasquez Y, Vansant H, Grenier C, Gonzalez N, Kansagra K, et al. 2023. Global patterns of genomic and phenotypic variation in the invasive harlequin ladybird. *BMC Biol* [Internet] 21:141. Available from: https://doi.org/10.1186/s12915-023-01638-7

Li H, Ralph P. 2019. Local PCA Shows How the Effect of Population Structure Differs Along the Genome. *Genetics* [Internet] 211:289–304. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6325702/

Linde K van der, Steck GJ, Hibbard K, Birdsley JS, Alonso LM, Houle D. 2006. FIRST RECORDS OF ZAPRIONUS INDIANUS (DIPTERA: DROSOPHILIDAE), A PEST SPECIES ON COMMERCIAL FRUITS FROM PANAMA AND THE UNITED STATES OF AMERICA. *flen* 89:402–404.

Lovell JT, MacQueen AH, Mamidi S, Bonnette J, Jenkins J, Napier JD, Sreedasyam A, Healey A, Session A, Shu S, et al. 2021. Genomic mechanisms of climate adaptation in polyploid bioenergy switchgrass. *Nature*:1–7.

Ma L, Cao L-J, Hoffmann AA, Gong Y-J, Chen J-C, Chen H-S, Wang X-B, Zeng A-P, Wei S-J, Zhou Z-S. 2020. Rapid and strong population genetic differentiation and genomic signatures of climatic adaptation in an invasive mealybug. *Diversity and Distributions* [Internet] 26:610–622. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/ddi.13053

Ma L-J, Cao L-J, Chen J-C, Tang M-Q, Song W, Yang F-Y, Shen X-J, Ren Y-J, Yang Q, Li H, et al. 2024. Rapid and Repeated Climate Adaptation Involving Chromosome Inversions following Invasion of an Insect. *Molecular Biology and Evolution* [Internet] 41:msae044. Available from: https://doi.org/10.1093/molbev/msae044

Machado HE, Bergland AO, Taylor R, Tilk S, Behrman E, Dyer K, Fabian DK, Flatt T, González J, Karasov TL, et al. 2021. Broad geographic sampling reveals the shared basis and environmental correlates of seasonal adaptation in Drosophila.Nordborg M, Wittkopp PJ, Nordborg M, editors. *eLife* [Internet] 10:e67577. Available from: https://doi.org/10.7554/eLife.67577

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution* [Internet] 38:4647–4654. Available from: https://doi.org/10.1093/molbev/msab199

Markow TA, Hanna G, Riesgo-Escovar JR, Tellez-Garcia AA, Richmond MP, Nazario-Yepiz NO, Laclette MRL, Carpinteyro-Ponce J, Pfeiler E. 2014. Population genetics and recent colonization history of the invasive drosophilid Zaprionus indianus in Mexico and Central America. *Biol Invasions* 16:2427–2434.

da Mata RA, Tidon R, Côrtes LG, De Marco P, Diniz-Filho JAF. 2010. Invasive and flexible: niche shift in the drosophilid Zaprionus indianus (Insecta, Diptera). *Biol Invasions* 12:1231–1241.

Matheson P, McGaughran A. 2022. Genomic data is missing for many highly invasive species, restricting our preparedness for escalating incursion rates. *Sci Rep* [Internet] 12:13987. Available from: https://www.nature.com/articles/s41598-022-17937-y

McGaughran A, Dhami MK, Parvizi E, Vaughan AL, Gleeson DM, Hodgins KA, Rollins LA, Tepolt CK, Turner KG, Atsawawaranunt K, et al. 2024. Genomic Tools in Biological Invasions: Current State and Future Frontiers. *Genome Biology and Evolution* [Internet] 16:evad230. Available from: https://doi.org/10.1093/gbe/evad230

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,
Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a MapReduce
framework for analyzing next-generation DNA sequencing data. *Genome Res.*
20:1297–1303.

Medley KA, Jenkins DG, Hoffman EA. 2015. Human-aided and natural dispersal drive gene
flow across the range of an invasive mosquito. *Molecular Ecology* [Internet] 24:284–
295. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.12925

Messer PW, Petrov DA. 2013. Population genomics of rapid adaptation by soft selective
sweeps. *Trends in Ecology & Evolution* [Internet] 28:659–669. Available from:
https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347(13)00207-3

Microsoft, Weston S. 2017. foreach: Provides Foreach Looping Construct for R. Available
from: https://CRAN.R-project.org/package=foreach

Nava DE, Nascimento AM, Stein CP, Haddad ML, Bento JMS, Parra JRP. 2007. Biology,
thermal requirements, and estimation of the number of generations of Zaprionus
indianus (Diptera: Drosopholidae) for the main fig producing regions of Brazil. *flen*
90:495–501.

Nguyen Ba AN, Cvijović I, Rojas Echenique JI, Lawrence KR, Rego-Costa A, Liu X, Levy SF,
Desai MM. 2019. High-resolution lineage tracking reveals travelling wave of adaptation
in laboratory yeast. *Nature* 575:494–499.

Nowling RJ, Manke KR, Emrich SJ. 2020. Detecting inversions with PCA in the presence of
population structure. *PLOS ONE* [Internet] 15:e0240429. Available from:
https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0240429

Nunez JCB, Lenhart BA, Bangerter A, Murray CS, Mazzeo GR, Yu Y, Nystrom TL, Tern C,
Erickson PA, Bergland AO. 2024. A cosmopolitan inversion facilitates seasonal
adaptation in overwintering Drosophila. *Genetics* [Internet] 226:iyad207. Available from:
https://doi.org/10.1093/genetics/iyad207

Oerke E-C. 2006. Crop losses to pests. *The Journal of Agricultural Science* 144:31–43.

Oliveira CM, Auad AM, Mendes SM, Frizzas MR. 2013. Economic impact of exotic insect pests
in Brazilian agriculture. *Journal of Applied Entomology* 137:1–15.

Parchami-Araghi M, Gilasian E, Keyhanian A. 2015. Olive infestation with Zaprionus indianus
Gupta (Dip.: Drosophilidae) in northern Iran: a new host record and threat to world olive
production. *Drosophila Information Service* 98:60–61.

Parvizi E, Dhami MK, Yan J, McGaughran A. 2023. Population genomic insights into invasion
success in a polyphagous agricultural pest, Halyomorpha halys. *Molecular Ecology*
[Internet] 32:138–151. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.16740

Patterson M, Marschall T, Pisanti N, van Iersel L, Stougie L, Klau GW, Schönhuth A. 2015.
WhatsHap: Weighted Haplotype Assembly for Future-Generation Sequencing Reads. *J
Comput Biol* 22:498–509.

Patton AH, Margres MJ, Stahlke AR, Hendricks S, Lewallen K, Hamede RK, Ruiz-Aravena M, Ryder O, McCallum HI, Jones ME, et al. 2019. Contemporary Demographic Reconstruction Methods Are Robust to Genome Assembly Quality: A Case Study in Tasmanian Devils. *Molecular Biology and Evolution* [Internet] 36:2906–2921. Available from: https://doi.org/10.1093/molbev/msz191

Pedersen BS, Layer R, Quinlan AR. 2020. smoove: structural-variant calling and genotyping with existing tools.

Pélissié B, Chen YH, Cohen ZP, Crossley MS, Hawthorne DJ, Izzo V, Schoville SD. 2022. Genome Resequencing Reveals Rapid, Repeated Evolution in the Colorado Potato Beetle. *Molecular Biology and Evolution* [Internet] 39:msac016. Available from: https://doi.org/10.1093/molbev/msac016

Pfeiffer DG, Shrader ME, Wahls JCE, Willbrand BN, Sandum I, van der Linde K, Laub CA, Mays RS, Day ER. 2019. African Fig Fly (Diptera: Drosophilidae): Biology, Expansion of Geographic Range, and Its Potential Status as a Soft Fruit Pest. *J Integr Pest Manag* [Internet] 10. Available from: https://academic.oup.com/jipm/article/10/1/20/5514212

Picq S, Wu Y, Martemyanov VV, Pouliot E, Pfister SE, Hamelin R, Cusson M. 2023. Range-wide population genomics of the spongy moth, Lymantria dispar (Erebidae): Implications for biosurveillance, subspecies classification and phylogeography of a destructive moth. *Evolutionary Applications* [Internet] 16:638–656. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/eva.13522

Platts PJ, Mason SC, Palmer G, Hill JK, Oliver TH, Powney GD, Fox R, Thomas CD. 2019. Habitat availability explains variation in climate-driven range shifts across multiple taxonomic groups. *Scientific Reports* 9:15039.

Price AL, Jones NC, Pevzner PA. 2005. De novo identification of repeat families in large genomes. *Bioinformatics* [Internet] 21:i351–i358. Available from: https://doi.org/10.1093/bioinformatics/bti1018

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker PIW, Daly MJ, et al. 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* 81:559–575.

Putnam NH, O'Connell BL, Stites JC, Rice BJ, Blanchette M, Calef R, Troll CJ, Fields A, Hartley PD, Sugnet CW, et al. 2016. Chromosome-scale shotgun assembly using an in vitro method for long-range linkage. *Genome Res* [Internet] 26:342–350. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4772016/

R Core Team. R: A language and environment for statistical   computing. Available from: http://www.R-project.org/

Rakes LM, Delamont M, Cole C, Yates JA, Blevins LJ, Hassan FN, Bergland AO, Erickson PA. 2023. A small survey of introduced Zaprionus indianus (Diptera: Drosophilidae) in orchards of the eastern United States. *Journal of Insect Science* [Internet] 23:21. Available from: https://doi.org/10.1093/jisesa/iead092

Renkema JM, Miller M, Fraser H, Légaré J-P, Hallett RH. 2013. First records of *Zaprionus*
indianus Gupta (Diptera: Drosophilidae) from commercial fruit fields in Ontario and
Quebec, Canada. *The Journal of the Entomological Society of Ontario* [Internet] 144.
Available from: https://journal.lib.uoguelph.ca/index.php/eso/article/view/3745

Ricciardi A. 2007. Are modern biological invasions an unprecedented form of global change?
Conserv Biol 21:329–336.

Roman G, Meller V, Wu KH, Davis RL. 1998. The opt1 gene ofDrosophila melanogaster
encodes a proton-dependent dipeptide transporter. *American Journal of Physiology-Cell*
Physiology [Internet] 275:C857–C869. Available from:
https://journals.physiology.org/doi/full/10.1152/ajpcell.1998.275.3.C857

Roque F, Matavelli C, Lopes PHS, Machida WS, Von Zuben CJ, Tidon R. 2017. Brazilian Fig
Plantations Are Dominated by Widely Distributed Drosophilid Species (Diptera:
Drosophilidae). *Annals of the Entomological Society of America* 110:521–527.

Sabeti PC, Varilly P, Fry B, Lohmueller J, Hostetter E, Cotsapas C, Xie X, Byrne EH, McCarroll
SA, Gaudet R, et al. 2007. Genome-wide detection and characterization of positive
selection in human populations. *Nature* [Internet] 449:913–918. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2687721/

Sardain A, Sardain E, Leung B. 2019. Global forecasts of shipping traffic and biological
invasions to 2050. *Nature Sustainability* 2:274–282.

Schluter D, Marchinko KB, Arnegard ME, Zhang H, Brady SD, Jones FC, Bell MA, Kingsley
DM. 2021. Fitness maps to a large-effect locus in introduced stickleback populations.
PNAS [Internet] 118. Available from: https://www.pnas.org/content/118/3/e1914889118

Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE, Jeschke JM, Pagad S, Pyšek P,
Winter M, Arianoutsou M, et al. 2017. No saturation in the accumulation of alien species
worldwide. *Nature Communications* 8:14435.

Seebens H, Essl F, Dawson W, Fuentes N, Moser D, Pergl J, Pyšek P, Kleunen M van, Weber
E, Winter M, et al. 2015. Global trade will accelerate plant invasions in emerging
economies under climate change. *Global Change Biology* 21:4128–4140.

da Silva VH, Laine VN, Bosse M, Spurgin LG, Derks MFL, van Oers K, Dibbits B, Slate J,
Crooijmans RPMA, Visser ME, et al. 2019. The Genomic Complexity of a Large
Inversion in Great Tits. *Genome Biology and Evolution* [Internet] 11:1870–1881.
Available from: https://doi.org/10.1093/gbe/evz106

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics [Internet] 31:3210–3212. Available from:
https://doi.org/10.1093/bioinformatics/btv351

Smit A, Hubley R, Green P. 2015. RepeatMasker Open-4.0. Available from:
http://www.repeatmasker.org

Soudi S, Crepeau M, Collier TC, Lee Y, Cornel AJ, Lanzaro GC. 2023. Genomic signatures of local adaptation in recent invasive Aedes aegypti populations in California. *BMC Genomics* [Internet] 24:311. Available from: https://doi.org/10.1186/s12864-023-09402-5

Steenwyk JL, Rokas A. 2021. ggpubfigs: Colorblind-Friendly Color Palettes and ggplot2 Graphic System Extensions for Publication-Quality Scientific Figures. *Microbiology Resource Announcements* [Internet] 10:10.1128/mra.00871-21. Available from: https://journals.asm.org/doi/10.1128/MRA.00871-21

Stern DB, Lee CE. 2020. Evolutionary origins of genomic adaptations in an invasive copepod. *Nat Ecol Evol* [Internet] 4:1084–1094. Available from: https://www.nature.com/articles/s41559-020-1201-y

Stuart KC, Cardilini APA, Cassey P, Richardson MF, Sherwin WB, Rollins LA, Sherman CDH. 2021. Signatures of selection in a recent invasion reveal adaptive divergence in a highly vagile invasive species. *Molecular Ecology* [Internet] 30:1419–1434. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.15601

Sutherst RW, Constable F, Finlay KJ, Harrington R, Luck J, Zalucki MP. 2011. Adapting to crop pest and pathogen risks under a changing climate. *WIREs Climate Change* 2:220–237.

Suvorov A, Kim BY, Wang J, Armstrong EE, Peede D, D'Agostino ERR, Price DK, Waddell PJ, Lang M, Courtier-Orgogozo V, et al. 2022. Widespread introgression across a phylogeny of 155 *Drosophila* genomes. *Current Biology* [Internet] 32:111-123.e5. Available from: https://www.sciencedirect.com/science/article/pii/S0960982221014962

Tepolt CK, Grosholz ED, de Rivera CE, Ruiz GM. 2022. Balanced polymorphism fuels rapid selection in an invasive crab despite high gene flow and low genetic diversity. *Molecular Ecology* [Internet] 31:55–69. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.16143

Tepolt CK, Palumbi SR. 2020. Rapid Adaptation to Temperature via a Potential Genomic Island of Divergence in the Invasive Green Crab, Carcinus maenas. *Front. Ecol. Evol.* [Internet] 8. Available from: https://www.frontiersin.org/articles/10.3389/fevo.2020.580701

Terhorst J, Kamm JA, Song YS. 2017. Robust and scalable inference of population history from hundreds of unphased whole-genomes. *Nat Genet* [Internet] 49:303–309. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5470542/

Thompson MJ, Jiggins CD. 2014. Supergenes and their role in evolution. *Heredity* [Internet] 113:1–8. Available from: http://www.nature.com/hdy/journal/v113/n1/full/hdy201420a.html

Thornton T, Tang H, Hoffmann TJ, Ochs-Balcom HM, Caan BJ, Risch N. 2012. Estimating kinship in admixed populations. *Am J Hum Genet* 91:122–138.

Timmeren SV, Isaacs R. 2014. Drosophila suzukii in Michigan vineyards, and the first report of Zaprionus indianus from this region. *Journal of Applied Entomology* 138:519–527.

Uller T, Leimu R. 2011. Founder events predict changes in genetic diversity during human-
mediated range expansions. *Global Change Biology* [Internet] 17:3478–3485. Available
from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2486.2011.02509.x

Vilela C. 1999. Is Zaprionus indianus Gupta, 1970 (Diptera, Drosophilidae) currently colonizing
the Neotropical region? *Drosophila Information Service* 82:37–39.

Weinig C, Brock MT, Dechaine JA, Welch SM. 2007. Resolving the genetic basis of
invasiveness and predicting invasions. *Genetica* [Internet] 129:205–216. Available from:
https://doi.org/10.1007/s10709-006-9015-7

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis.

Willbrand B, Pfeiffer D, Leblanc L, Yassin A. 2018. First Report of African Fig Fly, Zaprionus
indianus Gupta (Diptera: Drosophilidae), on the Island of Maui, Hawaii, USA, in 2017
and Potential Impacts to the Hawaiian Entomofauna. *Proceedings of the Hawaiian*
Entomological Society 50:55–65.

Yassin A, Capy P, Madi-Ravazzi L, Ogereau D, David JR. 2008. DNA barcode discovers two
cryptic species and two geographical radiations in the invasive drosophilid Zaprionus
indianus. *Molecular Ecology Resources* [Internet] 8:491–501. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1471-8286.2007.02020.x

Yassin A, David J. 2010. Revision of the Afrotropical species of Zaprionus (Diptera,
Drosophilidae), with descriptions of two new species and notes on the internal
reproductive structures and immature stages. *ZooKeys* 51:33–72.

Zanuncio-Junior JS, Fornazier MJ, Andreazza F, Culik MP, Mendonça L de P, Oliveira EE,
Martins D dos S, Fornazier ML, Costa H, Ventura JA. 2018. Spread of Two Invasive
Flies (Diptera: Drosophilidae) Infesting Commercial Fruits in Southeastern Brazil. *flen*
101:522–525.

Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. 2012. A high-performance
computing toolset for relatedness and principal component analysis of SNP data.
Bioinformatics 28:3326–3328.

1328