Lifetime fitness benefits of breeding site fidelity and low costs of inbreeding permit inbreeding tolerance in an avian cooperative breeder

Shailee S. Shah[1,2*,] Jennifer Diamond[3], Sahas Barve[4], Elissa J. Cosgrove[5], Reed Bowman[4], John W. Fitzpatrick[2,6], Nancy Chen[1*]

[1] Department of Biology, University of Rochester, Rochester, NY 14627, USA
[2] Cornell Lab of Ornithology, Cornell University, Ithaca, NY 14850, USA
[3] Department of Evolution and Ecology, University of California, Davis, Davis, CA 95616, USA
[4] Avian Ecology Program, Archbold Biological Station, Venus, FL 33960, USA
[5] Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14850, USA
[6] Department of Ecology & Evolutionary Biology, Cornell University, Ithaca, NY 14850, USA

* Correspondence: shailee.shah@cornell.edu, nancy.chen@rochester.edu

Acknowledgments: We thank Archbold Biological Station and its staff for continued support and management of the study area; all interns, students, and postdocs who have worked on collecting and managing the long-term data; the Cornell Statistical Consulting Unit for statistical support; V. Sclater for GIS support; and F. Romero, J. Summers, D. Seidman, and other members of the Chen Lab at the University of Rochester for invaluable feedback on previous versions of this manuscript.

Funding: This work was supported by National Science Foundation (NSF) Grants DEB0855879 and DEB1257628 and by the Cornell Lab of Ornithology Athena Fund. S.S.S was supported by NIH grant 1R35GM133412 to N.C. and NSF Postdoctoral Research Fellowship in Biology 2305705.

1 **Title:** Lifetime fitness benefits of breeding site fidelity and low costs of inbreeding permit inbreeding
2 tolerance in an avian cooperative breeder
3
4 **Abstract**
5

Inbreeding, or breeding with close relatives, often decreases individual fitness, but mate choice has been
shown to increase inbreeding risk in some species. Inbreeding is more likely in species with limited
dispersal, such as cooperative breeders where non-parental individuals—often offspring from previous
broods—provide parental care and frequently breed close to home. We leverage 32 years of data from a
population of Florida Scrub-Jays (*Aphelocoma coerulescens*), an avian cooperative breeder, to investigate
whether mate choice, and its lifetime fitness outcomes, affects inbreeding tolerance. We find that both sexes
show shorter dispersal distances than expected under a random mating scenario. While female-biased
dispersal mitigates inbreeding risk when pairing for the first time, limited movement away from the
breeding territory by both sexes for later pairings (following mate death or divorce) results in higher
relatedness between mates than expected under random mating. Males paired with close relatives and
females that move longer distances experience equivalently reduced lifetime reproductive success. Fitness
benefits of breeding site fidelity and relatively low fitness costs of inbreeding thus permit inbreeding
tolerance in Florida Scrub-Jays. Understanding how and why mate choice contributes to inbreeding is
crucial for informed conservation action for endangered species like the Florida Scrub-Jay.

20
21 **Keywords:** Mate choice, inbreeding tolerance, site fidelity, cooperative breeding, Florida Scrub-Jay

Main Text

Introduction

Matings among relatives, or inbreeding, can have severe negative fitness consequences such as increased susceptibility to parasites [1], lower birth rates and offspring survival [2,3], and increased risk of mortality during severe weather events [4]. In wild populations, inbreeding can be the result of two non-exclusive factors. First, in populations that are small and isolated, mating with relatives might be inevitable even when mating is random [5]. Second, mate choice may be driven by factors that increase fitness despite the risk of inbreeding, such as phenotype matching or limited dispersal [6]. In the latter case, while inbreeding tolerance may persist without severe population-level impacts in large, well-connected populations, it may exacerbate inbreeding depression in isolated population fragments [7]. Understanding how mate choice may result in increased instances of inbreeding in species of conservation concern is thus vital for informed conservation action.

Moving away from their natal area—*i.e.*, dispersal—is the primary mechanism by which animals avoid mating with kin [8,9]. However, dispersal may be limited in some species, such that when individuals become breeders, they often choose mates that are geographically close [8,10]. While limited dispersal can lead to fitness benefits such as better access to resources [11], higher survival [12], and nepotism [13], it can also increase kin competition [14] and the likelihood of mating with a close relative [15–17]. Similarly, once individuals establish a breeding territory, any subsequent dispersal also may be geographically limited even when pairing with a new mate [8]. Such site fidelity may have fitness benefits due to familiarity and reduced competitive interactions [18–20], but also may increase inbreeding risk in spatially structured populations where limited dispersal results in the formation of kin neighborhoods [21]. Nevertheless, if the fitness benefits of limited dispersal outweigh the costs of inbreeding at the individual level, the risk of inbreeding may be tolerated [6,22]. Conversely, additional behaviors such as kin avoidance—where individuals can recognize and avoid mating with kin—and sex-biased dispersal—where one sex disperses farther than the other—may prevent inbreeding even in species with limited dispersal [6].

Previous studies investigated whether species with limited dispersal exhibit inbreeding tolerance or avoidance behaviors [7,23–28], and even quantified the fitness tradeoffs between dispersal distance and kinship in mate choice [23]. However, few such studies have incorporated measures of lifetime fitness. This matters because reproductive strategies over an individual's lifetime, especially in long-lived species such as many cooperative breeders [29,30], may compensate for any fitness costs associated with inbreeding or dispersal. Thus, we must examine lifetime fitness to obtain a comprehensive understanding of the individual fitness outcomes underlying mate choice that may increase inbreeding risk. Additionally, the importance of dispersal distance in mate choice may differ based on whether an individual is breeding for the first time ("first pairing") versus an experienced breeder pairing with a new mate ("later pairing") [8]. Because most studies only examine mate choice during first pairing, mechanisms such as site fidelity that may limit dispersal or increase inbreeding risk at later life stages [8] are ignored. Here, we leverage 32 years of individual-level data on dispersal, mating, reproductive success, and survival from a continuously monitored population of an avian cooperative breeder, the Florida Scrub-Jay (*Aphelocoma coerulescens*), to investigate whether inbreeding risk is elevated due to mate choice throughout an individual's lifetime, and the resulting lifetime fitness outcomes.

Fewer than 6,000 individuals of the Federally Threatened Florida Scrub-Jay remain, located in numerous, widely isolated populations and restricted to fire-maintained, xeric oak scrub habitats of the Florida peninsula [31]. Their preferred habitat is usually saturated with year-round territories, each occupied by a socially monogamous dominant breeding pair, accompanied often by subordinate non-breeders of both sexes [32]. Subordinates are primarily offspring from previous broods that delay dispersal and remain in their natal territory for one or more years, despite being physiologically capable of breeding at one year of

age [32]. Most individuals become breeders by dispersing to replace dead breeders on existing territories,
though direct inheritance or "budding" of natal territories sometimes occurs [32]. Females disperse farther
than males, though neither sex typically disperses very far [33,34]. Once they become breeders, Florida
Scrub-Jays remain breeders until their death, pairing with a new mate after their current mate dies [32], or
in rare cases when the mate is still alive ("divorce") [35]. We focus on a population of Florida Scrub-Jays
that has been intensively studied at Archbold Biological Station in central Florida (27.10°N, 81.21°W) since
1969 [32]. Immigration into the Archbold population from neighboring populations has declined since
1995, leading to a corresponding increase in inbreeding and associated fitness costs [3].
81
To investigate how mate choice impacts inbreeding risk, we first examined how the relatedness of Florida
Scrub-Jays to potential mates varied with distance from both their natal and any subsequent breeding
territories. Given fine-scale genetic isolation-by-distance in this population [34], we predicted that potential
mates closer to both their natal and breeding territories would be relatively closely related for both sexes,
and more so for females since males are the more philopatric sex [33]. Second, we examined the extent to
which dispersal distance, relatedness, or both, correspond with mate choice. Previous work in this study
population found individuals that dispersed shorter distances had higher fitness [36], and inbred individuals
have lower fitness [3]. Thus, we predicted that both dispersal distance and relatedness of mates would
deviate from expectations under a random mating scenario, with average dispersal distance shorter, and
relatedness of mates lower (owing to inbreeding avoidance), than expected under a random mating scenario.
92
Next, to investigate the relative fitness costs and benefits involved in mate choice patterns, we quantified
the impact of distance moved and relatedness between mates on lifetime reproductive success and breeder
survival. We predicted that for both sexes, longer dispersal distances for both first and later pairings would
result in lower lifetime reproductive success and survival [36]. Further, we predicted that if we found no
evidence of kin avoidance in our mate choice analysis, the fitness cost of relatedness between mates would
be equivalent to or lower than the cost of long-distance movement (suggesting inbreeding tolerance).
Finally, we examined whether Florida Scrub-Jays employ other reproductive strategies, such as divorce, to
minimize the costs of inbreeding. Since lifetime reproductive success is directly proportional to the number
of annual breeding attempts [31], Florida Scrub-Jays may favor early breeding with a close relative over
not breeding at all, but then may switch to less related mates when an opportunity arises. Thus, we predicted
that pairs with more closely related mates would be more likely to dissolve because of divorce than death
of a mate. Ultimately, by linking the effect of distance and kinship on mate choice with individual lifetime
fitness outcomes, we investigated how mate choice that increases inbreeding risk can persist in a population.
106
107   **Methods**
108
109   *Data collection*
110
Florida Scrub-Jays at Archbold Biological Station have been continuously monitored since 1969, with the
study tract expanded to its current size (mean ± sd = 914.04 ± 143.44 ha) by 1990 [31]. All birds are banded
with unique combinations of color bands. Throughout the breeding season (early March to early June), all
nests are located and monitored for fledging or failure [31]. All territories are mapped in April, with territory
boundaries assigned at locations of territorial disputes between neighboring groups [31]. In addition to nest
monitoring, the population is censused each month, resulting in detailed data on dispersal and survival of
both fledglings and adults. Pairs are socially monogamous, with very low rates of extra pair-paternity (<1%)
[37], allowing accurate parentage inference from field observations alone. Furthermore, previous work
verified genetic parentage of offspring born between ~1989-1991, 1994-1995, and 1999-2013 [3]. Using
these detailed data, we have accumulated a 15-generation pedigree of known-age, known-parentage
individuals with precise measurements of lifespan and reproductive success. For this study, we used data
from 1990-2021, which included on average (± SD) 71.4 (± 14.7) territories (area mean ± sd =12.98 ± 6.94
ha) per year. All analyses were conducted using R version 4.2.2 [38]. Fieldwork was approved by the

Institutional Animal Care and Use Committees at Cornell University (IACUC 2010-0015) and Archbold
Biological Station (AUP-006-R) under permits from the US Fish and Wildlife Service (TE824723-8, TE-
117769), the US Geological Survey (banding permits 07732, 23098), and the Florida Fish and Wildlife
Conservation Commission (LSSC-10-00205).
128
We first measured observed dispersal distances and calculated relatedness to mate for both male and female
breeders in all new breeding pairs formed between 1990 and 2021. New breeding pairs include pairs formed
by two first-time breeders, one first-time and one experienced breeder, or two experienced breeders. For
first-time breeders, we called the event a "first pairing" and for experienced breeders, a "later pairing". The
first pairing for one mate is thus not always the first pairing for the other mate (*e.g.*, an experienced male
could repair with a female who is a first-time breeder, and *vice versa*). New pairs might occupy a new
territory, which may be inherited by one of the two breeders or formed as an extension of their natal territory
in a process called "budding" [31]. Alternatively, a new pair might form on a non-natal territory already
occupied by one of the two breeders, where a breeding vacancy is typically formed due to the death of a
breeder, but occasionally via divorce [31]. Thus, both first and later pairings can occur without any
movement away from the currently occupied territory by at least one of the two breeders. When the focal
individual remained on its natal territory and became a breeder, formed a territory by budding, or remained
on its non-natal breeding territory, we quantified distance moved as 0 m. To obtain accurate measures of
distance moved, we excluded any pairings that involved movement that started or ended outside of the
study tract. We also excluded any movement that did not end with the individual attaining a breeding
position (*i.e.*, staging, a strategy adopted by about 28% of yearlings [39]). For staging individuals that
eventually became breeders, we considered it a first pairing event with movement from their staging to
breeding territory. Additionally, using distance from natal to breeding territory as distance moved for first
pairing did not change our results (see *SI*).
148
In accordance with previous work in this system, we used the observed number of territories crossed as the
most biologically relevant measure of distance moved [36]. The number of territories crossed was highly
correlated with distance moved in meters (Pearson's $r = 0.91$; [36]). To calculate the number of territories
crossed by a jay for a new pairing, we drew a straight-line movement trajectory from the centroid of its
previous territory to the centroid of its new breeding territory. We then used the *st_intersects* function from
the R package *sf* [40] to calculate the number of territories (*i.e.*, polygons) that each line intersected. For
cases where the function returned a distance ≥ 1 territory away, but the territory name did not change from
its natal or previous breeding territory, we manually corrected distances using territory maps (see *SI*). This
occurred in 84 out of 1690 [5%] of movement events due to minor inconsistencies in territory naming
conventions, where territories were named based on the dominant male breeder rather than territory
location, and stochastic drift in territory boundaries. We also calculated the kinship coefficient between
mates for all new breeding pairs using the population pedigree with the *kinship* function in the R package
kinship2 [10]. With no inbreeding, the kinship coefficient ranges from 0 to 0.5, such that 0.5 is an
individual's kinship coefficient with itself; 0.25 is the average kinship coefficient between siblings and
between parent and offspring; 0.125 is the average kinship coefficient between half-siblings, grandparents-
grandchildren, and aunt/uncle-niece/nephew pairs; and 0.0625 is the average kinship coefficient between
first cousins [41]. Pedigree-based inbreeding estimates are correlated with genomic-based estimates in our
study population [3].
167
168 *Data analysis*
169
First, to examine how relatedness to potential mates varies with distance, we constructed a dataset of all
possible mate pairings for each year and each sex separately. We only included breeders that formed new
breeding pairs in a given year (*i.e.*, individuals that were both available as potential mates and successful at
finding new mates), which comprised both first-time and experienced breeders. For a given year, we paired
all available females with each available male and vice versa (*N*: Male, First pairing = 421; Male, Later

pairing = 249; Female, First pairing = 391; Female, Later pairing = 256 instances). We calculated the
potential distance that each focal individual would have to travel to realize this pairing as the distance
between the individual's territory in the previous year to the potential mate's territory in the current year.
For simplicity, we assumed that the potential mate would retain its territory. We also calculated the kinship
coefficient for all potential pairings as described above, and classified potential mates to whom the kinship
coefficient of the focal individual was ≥ 0.0625 (at least as related as first cousins, Fig. S2) as close relatives.
We then fit generalized linear mixed models (GLMMs) with whether the potential mate was a close relative
(1 = yes, 0 = no) as the dependent variable, a binomial error structure, distance to mate as a fixed effect,
and year and focal individual ID as random effects. We fit separate models for each sex and pairing type
(N: Male, First pairing = 12,155; Male, Later pairing = 11,837; Female, First pairing = 10,938; Female,
Later pairing = 11,000 possible pairings).
186
Second, we investigated the effects of distance to mate and relatedness on mate choice. To quantify
deviations of distance and average relatedness of mates from expectations under a random mating scenario,
we randomly permuted all available mates of the opposite sex among all available mates of the focal sex
for each year. We then calculated the distance and kinship coefficient to mate using the same methods as
above for each randomly generated pair. We ran 1,000 iterations per year and calculated the mean distance
and kinship coefficient to mate for each year for each focal sex and pairing type. We then compared these
expected distributions of mean distances and kinship coefficients to observed means by calculating
empirical p-values [42]. Specifically, we quantified the proportion of simulations that had a mean distance
to mate that was higher than the observed mean (S_{high}), and the proportion of simulations that had a mean
kinship coefficient to mate that was lower than the observed mean (S_{low}). Empirical p-values to test whether
observed means were significantly higher than expected under a random mating model were calculated as
$(S_{high}+1)/(N+1)$, and empirical p-values to test whether observed means were significantly lower than
expected under a random mating model were calculated as $(S_{low}+1)/(N+1)$, where N is the total number of
simulations.
201
Third, to examine how distance and relatedness to mate impacts lifetime fitness of Florida Scrub-Jays, we
calculated lifetime reproductive success for all breeders as the total number of offspring produced over an
individual's lifetime that survived to adulthood, *i.e.*, age 1 year, the stage at which Florida Scrub-Jays
become physiologically capable of breeding [32,43–45]. Since the mean growth rate of our study
population is close to 1 (0.998) [46], we deemed LRS an appropriate metric of individual fitness [47]. For
this analysis, we restricted our dataset to individuals from cohorts from which all members born within the
study tract were last observed before 2021 to eliminate bias towards short-lived individuals [48,49]. The
final dataset included individuals born between 1988-2006 and in 2009 (N: Males =196, Females =122).
210
We then fit two separate GLMMs by sex with negative binomial error distributions (to account for
overdispersion), lifetime reproductive success as the dependent variable, and distance moved for first
pairing, weighted kinship coefficient to mates, and breeding lifespan as fixed effects (N: Males = 196,
Females = 122). Fixed effects were not correlated (Table S1). We calculated the weighted kinship
coefficient by multiplying the relatedness of the focal individual to each of their mates by the number of
years they were paired with that mate and summing the resulting values. We defined breeding lifespan as
the number of years from becoming a breeder to death. Random effects included natal year to account for
variation between birth cohorts. Further, to specifically examine the effect of moving away from the
breeding territory, we also fit two additional GLMMs with the same error structure, dependent variable,
and random effects, with total distance moved for later pairings, weighted kinship coefficient to mates, and
breeding lifespan as fixed effects using a truncated dataset that only included individuals that had switched
mates in their lifetimes (N: Male = 92, Female = 51). All GLMMs were fit using the function *glm.nb* in the
R package *MASS* [50].
224

Next, to examine the effect of distance moved on breeder survival, we fit two Cox proportional hazards
models, one for each sex, with distance moved for first pairing as a fixed effect and natal year as a random
effect using the *Surv* and *coxme* functions in the R packages *survival* [51] and *coxme* [52]. For the subset
of individuals that switched mates at least once in their lifetime (*N*: Male = 92, Female = 51), we also
examined the effect of territory loss (*i.e.*, distance moved for later pairings ≥ 1 territory away) on breeder
survival. We fit Cox proportional hazards models with loss of territory (1 = territory lost, 0 = territory
retained) as a fixed effect and natal year as a random effect.

232
Finally, to investigate whether Florida Scrub-Jays compensated for the impact of inbreeding on
reproductive success via divorce, we fit GLMMs with the cause of pair dissolution (1 = divorce, 0 = mate
death) as the dependent variable; the kinship coefficient to mate as a fixed effect; year and individual ID as
random effects; and a binomial error structure. We also fit models with a categorical fixed effect of kinship
between mates (1 = first-order relative, 0 = not a first-order relative) to account for cases where opposite-
sex offspring might inherit dominant breeding positions, resulting in mother-son or father-daughter pairings
which may be more likely to end in divorce [53]. We fit separate GLMMs by sex and pairing type using
the function *glmer* in the R package *lme4* [54]. Fixed effects for all models were standardized using *z* scores
[55], and the variance inflation factor was < 2 for all fixed effects in all GLMMs [56].

242
243 **Results**

244
The number of available mates ranged from 14-42 females and 13-45 males across our study period with
an average (± SD) of 26.2 (± 7.70) males and 25.2 (± 7.44) females available as mates per year. These
available mates accounted for, on average, 36% of all breeders per year. Of the individuals included in our
lifetime fitness analysis, 42% of females (51 out of 122) and 47% of males (92 out of 196) switched mates
at least once in their lifetime (mean ± SD number of mate switching instances per lifetime: Males = 0.83 ±
1.17, range: 0-6; Females = 0.73 ± 1.04, range: 0-4). Females, on average, moved farther than males, though
more so for first pairings (Male = 1.05 ± 1.54, Female = 2.04 ± 2.61 territories crossed; *t*-test t = 6.52, df =
631.23, P < 0.001; [33]) than later pairings (Male = 0.34 ± 0.91, Female = 0.62 ± 1.29 territories crossed;
t-test t = 3.73, df = 765.87, P < 0.001). In 212 out of 869 (24%) instances of later pairings, the focal
individual (60% of whom were females) moved one or more territories away from its established breeding
territory. Furthermore, we found that out of all mate switching events observed, ~18% (*N*: Males = 64 of
187, Females = 31 of 171 events) were due to divorce, not mate death, a similar proportion to that found
previously [32].

258
The kinship coefficient to mate ranged from 0 (unrelated) to 0.25 (*i.e.,* either siblings or parent-offspring).
Across all years, only 44 out of 1443 (3%) of observed breeding pairs were comprised of individuals that
were at least as closely related as first cousins (*i.e.*, had a kinship coefficient ≥ 0.0625). As predicted, we
found that both sexes were more closely related to available opposite-sex breeders at shorter geographical
distances on both their natal and breeding territories, with the greatest effect size for females on their natal
territories (Males, Natal: *Z* = -11.40, *P* < 0.001; Females, Natal: *Z* = -14.79, *P* < 0.001; Males, Breeding: *Z*
= -8.49, *P* < 0.001; Females, Breeding: *Z* = -10.38, *P* < 0.001; Fig. 1).

266
We found that dispersal distance, more than relatedness to mate, impacted mate choice decisions in Florida
Scrub-Jays. When compared to a random mating scenario, both sexes moved significantly shorter distances
within the study tract than expected under random mating for both first and later pairings (empirical p-value
≤ 0.05 for both), even though females moved farther than males in the first pairing stage (Fig. 3A). We
found no evidence of kin avoidance: the kinship coefficient to mates for both sexes was never lower than
predicted by the random mating model (*P* > 0.05 across all categories). Mates of later pairings were more
closely related than expected by chance for both sexes (Males: *P* = 0.001; Females: *P* = 0.001; Fig. 3B),
but not for first pairings (Males: *P* = 0.26; Females: *P* = 0.30; Fig. 3B).

Next, we quantified the lifetime fitness consequences of different mate choice decisions. We found that a
higher weighted kinship coefficient to mates over an individual's lifetime resulted in significantly fewer
offspring that survived to adulthood for males—such that in the relatively rare cases that males breed with
closely related females for the majority of their lifetime, their LRS is close to zero (All breeding males: Z
= -2.93, P = 0.003, Only males that switched mates at least once: Z = -0.29, P = 0.09; Fig. 3B, D; Table
1)—but not females (All breeding females: Z = 0.57, P = 0.57, Only females that switched mates at least
once: Z = -0.29, P = 0.09; Fig. 3B, D; Table 1). Conversely, females, but not males, that moved farther
distances for later pairings had significantly fewer offspring that survived to adulthood (Females: Z = -2.42,
P = 0.02, Males: Z = -0.88, P = 0.38; Fig. 3CA, Table 1), with a similar trend for dispersal distance at first
pairing, though the effect was not significant (Females: Z = -1.74, P = 0.08, Males: Z = -0.27, P = 0.78;
Fig. 3A, Table 1). Moreover, as we predicted, breeding lifetime—*i.e.*, the number of years an individual
spends as a breeder—had the largest effect on lifetime reproductive success for both sexes: longer breeding
lifetimes resulted in more total offspring surviving to adulthood (Table S1-2). For both sexes, we found
that neither distance moved for first pairing (Male: Z = -0.01, P = 0.99; Female: Z = 0.92, P = 0.36) nor
dispersal away from an established breeding territory in the event of mate switching (Male: Z = 1.02, P =
0.31; Female: Z = -0.25, P = 0.81) impacted breeder survival.
292
Finally, the kinship coefficient to their mate had no effect on the likelihood of divorce in any scenario
(Males, First pairing: Z = 0.47, P = 0.64; Males, Later pairing: Z = -0.03 P = 0.98; Females, First pairing:
Z = -0.14, P = 0.89; Females, Later pairing: Z = 0.66, P = 0.51), nor was divorce more likely in cases of
pairing with first-order relatives (Males, First pairing: Z = 0.63, P = 0.53; Males, Later pairing: Z = -0.03,
P = 0.98; Females, First pairing: No instances of divorce when paired with a first-order relative; Females,
Later pairing: Z = 0.57, P = 0.57). However, only 0.01% (13 out of 1443) of all breeding pairs were first-
order relatives.
300
301   **Discussion**
302
In many animals, the costs of inbreeding avoidance, such as traversing a long distance to find an unrelated
mate, might outweigh the benefits of behaviors that can increase the likelihood of inbreeding, such as
limited dispersal [22]. Inbreeding may instead be tolerated, even while behaviors such as sex-biased
dispersal and kin recognition may evolve to prevent it [6,57,58]. In cooperatively breeding species, where
offspring often delay dispersal from their natal territory for one or more years and frequently breed near
home, fitness benefits of familiarity with the surrounding area or nepotism, or both, can lead to extremely
limited dispersal [8]. Additionally, since cooperative breeding typically arises in habitats saturated with
territories, breeders may show extreme site fidelity when pairing with a new mate [8]. Thus, especially
among long-lived species, both sexes may exhibit little propensity for movement away from their natal and
breeding territories, thereby increasing the likelihood of pairing with closely related mates. Here, we
investigated the impact of dispersal distance and relatedness on mate choice in an avian cooperative breeder,
the Florida Scrub-Jay. This species exhibits limited dispersal for both first and subsequent pairings. We
found that only fidelity to their breeding territory, rather than limited movement for the first pairing, resulted
in higher relatedness between mates than expected by random chance. Moreover, we found that both pairing
with close relatives and moving a greater distance from breeding territory were correlated with lower
lifetime reproductive success, although the effects varied by sex. However, breeder survival did not vary
with natal or breeding dispersal distance. Overall, our results suggest that although sex-biased natal
dispersal acts as an inbreeding avoidance mechanism in Florida Scrub-Jays, the fitness benefits of
remaining on established breeding territories may outweigh the costs of breeding with close relatives,
resulting in inbreeding tolerance.
323
Not surprisingly for a generally philopatric species, we found that Florida Scrub-Jays are more closely
related to geographically closer potential mates on both their natal and breeding territories. Previous work
has shown that when pairing for the first time, Florida Scrub-Jay females move farther distances than males

[32,33,34]. This sex-bias in distance moved for first pairing is reflected in our finding that females are more
closely related to potential mates (*i.e.*, males) closer to their natal territory than males are to potential mates
(*i.e.*, females), while both sexes are significantly more related to potential mates at dispersal distances of ≤
4 territories away from their natal territory than potential mates >4 territories away. Further, even around
the breeding territory, both sexes were more closely related to potential mates at closer geographical
distances. Thus, in agreement with previous work examining genetic isolation-by-distance in this species
[34], and similar to other communal or cooperatively breeding species [59,60], Florida Scrub-Jays exhibit
a viscous population structure such that inbreeding risk is elevated at shorter dispersal distances.
335
Surprisingly, we found that despite elevated inbreeding risk at shorter dispersal distances and evidence of
inbreeding depression in our study population [3], mate choice was driven primarily by distance to mate.
We compared observed means of distance moved and kinship coefficient to mate to those expected under
a random mating scenario. We used a similar approach to simulating a random mating scenario as previous
studies of breeding populations of territorial birds where most, if not all, individuals were uniquely marked
[7,23,26]. However, contrary to previous studies, we differentiated between instances of first and later
pairings. We found that neither sex at either pairing stage exhibited significantly lower than expected
kinship coefficient to mates, indicating no evidence of inbreeding avoidance. Although we found that the
distances moved for first and later pairings are both lower than expected under a random mating scenario
for both sexes, female bias in distance moved for first pairing prevents higher relatedness between breeders
in first pairings than expected by random chance. We found no sex bias in distance moved for later pairings,
which likely leads to the observed higher relatedness to mate than expected under a random mating scenario.
Thus, our results suggest that fidelity to the breeding territory, rather than limited natal dispersal, is the
most frequent cause of inbreeding in Florida Scrub-Jays. To our knowledge, this is the first study to examine
the role of breeding site fidelity on inbreeding risk. The factors influencing mate choice for later pairings
might be drastically different from mate choice for first pairings [8]. Future studies on mate choice and
inbreeding patterns in this and other species would do well to take later life stages into account.
353
Significant philopatry, which has been found in both social and asocial species, can be driven by various
factors such as better access to resources and protection from predators due to prior local experience [11],
lower competition due to nepotism [61,62], and high physiological costs of dispersal [63]. Although we
expected to find a pattern of kin avoidance given that inbred Florida Scrub-Jay offspring have a lower
probability of survival [3], theoretical work suggests that if the frequency and/or cost of inbreeding is low,
inbreeding avoidance mechanisms—such as avoiding mating with kin—may not evolve [22]. Inbreeding—
especially with close relatives—is relatively rare in our study population, with only 3% of all breeding pairs
resulting in consanguineous matings. Further, the cost threshold for the evolution of inbreeding avoidance
can be fairly high, especially in the absence of numerous viable mates [22]. Since our results suggest that
geographical distance might constrain the pool of viable mates in our study population, kin discrimination
and avoidance may lead to further loss of breeding opportunities and thus may not be adaptive. Finally,
breeders may employ reproductive strategies that minimize the fitness costs of inbreeding over their lifetime
in more nuanced ways not yet detected in our study.
367
Finally, we investigated how lifetime fitness varies with dispersal distance and relatedness to mates in
Florida Scrub-Jays. As predicted, we found a negative effect of both dispersal distance and relatedness on
lifetime reproductive success, but the effects varied by sex. The number of offspring that survived to
adulthood was lower for males that mated with closely related females for longer proportions of their
breeding lifetime. On the other hand, relatedness to mate did not significantly predict female lifetime
reproductive success. Instead, females that moved longer distances for their later pairings had lower lifetime
fitness. This finding is contrary to other studies on avian species where breeding dispersal increases lifetime
reproductive success, generally because dispersers settle on territories of higher quality than their previous
territory [64–67]. Most of those studies are on species that migrate seasonally, which could decrease the
cost of acquiring a new territory at the start of the breeding season. However, Canada Jays (*Perisoreus*

canadensis), which, similar to Florida Scrub-Jays, defend year-round territories and cache food—which
could lead to higher costs of acquiring new territories and a greater benefit of site fidelity to a familiar
territory—have been found to have higher lifetime fitness when they engage in breeding dispersal [67].
Though unlike our much more stable study population [46], the study population for the Canada Jay study
is declining, which may lead to more breeding vacancies and increased fitness as a result of breeding
dispersal [67]. Additionally, we did not find any evidence of lower male or female breeder survival with
higher distance moved for first or later pairings. These results contradict previous studies in this population
that found that both sexes had lower lifetime reproductive success and annual survival at greater dispersal
distances, with a stronger effect in males compared to females [36]. However, the previous study included
immigrants from outside the study tract in its analysis of lifetime fitness. We excluded immigrants because
in the context of mate choice, we were unable to account for all possible mates within their natal population,
nor accurately measure distances moved.
390
Overall, while effects differed by sex, our results suggest a trade-off in Florida Scrub-Jays between nearly
equivalent costs of mating with close relatives and moving farther distances away from their natal and
breeding territories. We note that our analysis likely underestimates the cost of moving farther distances,
since it does not include individuals that die in transit or otherwise fail to establish as breeders and also
excludes dispersers out of the study tract [68]. Additionally, the relationship between dispersal distance and
lower lifetime reproductive success may be a result of lower-quality individuals dispersing farther distances
rather than a cost of dispersal itself [69,70]. Thus, since the cost of inbreeding does not outweigh the higher
fitness experienced with limited dispersal, the population seems to exhibit inbreeding tolerance in
accordance with theoretical expectations [22]. Our results are similar to those found in a range of other
animal taxa—including insects such as the field cricket (*Teleogryllus commodus*) [71] and the parasitoid
wasp *Venturia canescens* [72]; birds such as the Eurasian blue tit (*Cyanistes caeruleus*) [24], great tit (*Parus
major*) [27], medium ground finch (*Geospiza fortis*) [26], collared flycatcher (*Ficedula albicollis*) [15], and
red-cockaded woodpecker (*Leuconotopicus borealis*) [23]; and mammals such as reindeer (*Rangifer
tarandus*) [73], grey wolf (*Canis lupus*) [74], and arctic fox (*Vulpes lagopus*) [74]—where higher than
expected levels of inbreeding are tolerated despite some level of inbreeding depression. Interestingly, we
found no evidence that Florida Scrub-Jays are more likely to divorce closely related mates, suggesting that
breeding with close relatives is not simply a response to lack of mate availability [17]. Further, since
reproductive success increases with pair experience in Florida Scrub-Jays [32], mate switching due to
divorce may prove costly. While we did not find evidence of preference for closely related mates, breeding
with kin is also thought to confer kin selection benefits that may additionally contribute to inbreeding
tolerance in some species [22,75].
412
In summary, we show that a mate choice behavior that maximizes individual lifetime fitness—*i.e.*,
relatively short-distance dispersal—contributes to some inbreeding in an avian cooperative breeder.
Importantly, we show that it is site fidelity at the later pairing stage, not limited dispersal away from the
natal territory, that drives higher inbreeding than expected under random mating. Despite documented
inbreeding depression, we find no evidence of inbreeding avoidance. Instead, we find that individual
lifetime fitness costs of inbreeding are low enough for inbreeding tolerance and the persistence of limited
dispersal. On a population level, inbreeding tolerance in mate choice might exacerbate inbreeding
depression due to increasing population isolation. Our study underscores the importance of examining the
costs and benefits underlying mate choice at all life stages to better understand the drivers of inbreeding in
species of conservation concern. From a conservation standpoint, since large populations tend to be less
inbred, our findings also underscore the profound importance of protecting large, contiguous populations
in which successful long-distance dispersal is possible, thereby reducing levels of inbreeding to tolerable
levels over the long term.
426

References

1. Coltman DW, Smith JA, Bancroft DR, Pilkington J, MacColl ADC, Clutton-Brock TH, Pemberton JM. 1999 Density-dependent variation in lifetime breeding success and natural and sexual selection in Soay rams. *Am. Nat.* **154**, 730–746. (doi:10.1086/303274)

2. Szulkin M, Garant D, Mccleery RH, Sheldon BC. 2007 Inbreeding depression along a life-history continuum in the great tit. *Journal of Evolutionary Biology* **20**, 1531–1543. (doi:10.1111/j.1420-9101.2007.01325.x)

3. Chen N, Cosgrove EJ, Bowman R, Fitzpatrick JW, Clark AG. 2016 Genomic consequences of population decline in the endangered Florida scrub-jay. *Curr. Biol.* **26**, 2974–2979. (doi:10.1016/j.cub.2016.08.062)

4. Keller L, Reeve HK. 1994 Partitioning of reproduction in animal societies. *Trends Ecol. Evol.* **9**, 98–102. (doi:10.1016/0169-5347(94)90204-6)

5. Keller LF, Waller DM. 2002 Inbreeding effects in wild populations. *Trends Ecol. Evol.* **17**, 230–241. (doi:10.1016/S0169-5347(02)02489-8)

6. Nichols HJ. 2017 The causes and consequences of inbreeding avoidance and tolerance in cooperatively breeding vertebrates. *J. Zool.* **303**, 1–14. (doi:10.1111/jzo.12466)

7. Billing AM *et al.* 2012 Evidence of inbreeding depression but not inbreeding avoidance in a natural house sparrow population. *Molecular Ecology* **21**, 1487–1499. (doi:10.1111/j.1365-294X.2012.05490.x)

8. Greenwood PJ, Harvey PH. 1982 The natal and breeding dispersal of birds. *Annu. Rev. Ecol. Syst.* **13**, 1–21. (doi:10.1146/annurev.es.13.110182.000245)

9. Szulkin M, Stopher KV, Pemberton JM, Reid JM. 2013 Inbreeding avoidance, tolerance, or preference in animals? *Trends Ecol. Evol.* **28**, 205–211. (doi:10.1016/j.tree.2012.10.016)

10. Clutton-Brock TH, Lukas D. 2012 The evolution of social philopatry and dispersal in female mammals. *Mol. Ecol.* **21**, 472–492. (doi:10.1111/j.1365-294X.2011.05232.x)

11. Pärt T. 1994 Male philopatry confers a mating advantage in the migratory collared flycatcher, *Ficedula albicollis*. *Animal Behaviour* **48**, 401–409. (doi:10.1006/anbe.1994.1254)

12. Griesser M, Nystrand M, Ekman J. 2006 Reduced mortality selects for family cohesion in a social species. *Proceedings of the Royal Society B: Biological Sciences* **273**, 1881–1886. (doi:10.1098/rspb.2006.3527)

13. Thünken T, Bakker TCM, Baldauf SA, Kullmann H. 2007 Active inbreeding in a cichlid fish and its adaptive significance. *Current Biology* **17**, 225–229. (doi:10.1016/j.cub.2006.11.053)

14. West SA, Pen I, Griffin AS. 2002 Cooperation and competition between relatives. *Science* **296**, 72–75. (doi:10.1126/science.1065507)

15. Pärt T. 1996 Problems with testing inbreeding avoidance: the case of the collared flycatcher. *Evolution* **50**, 1625–1630. (doi:10.1111/j.1558-5646.1996.tb03934.x)

16. Daniels SJ. 1997 Female dispersal and inbreeding in the red-cockaded woodpecker. Virginia
Polytechnic Institute and State University.

17. Szulkin M, Sheldon BC. 2008 Dispersal as a means of inbreeding avoidance in a wild bird population.
Proc. R. Soc. B. **275**, 703–711. (doi:10.1098/rspb.2007.0989)

18. Beck KB, Loretto M-C, Bugnyar T. 2020 Effects of site fidelity, group size and age on food-caching
behaviour of common ravens, *Corvus corax*. *Animal Behaviour* **164**, 51–64.
(doi:10.1016/j.anbehav.2020.03.015)

19. Schmidt KA. 2014 Site fidelity curbs sequential search and territory choice: a game theoretical
approach. *Funct. Ecol.* **28**, 1494–1503. (doi:10.1111/1365-2435.12291)

20. Byrne ME, Cohen BS, Collier BA, Chamberlain MJ. 2022 Nest site fidelity and nesting success of
female wild turkeys. *Wildlife Society Bulletin* **46**, e1279. (doi:10.1002/wsb.1279)

21. Hamilton WD. 1964 The genetical evolution of social behaviour. I. *J. Theor. Biol.* **7**, 1–16.
(doi:10.1016/0022-5193(64)90038-4)

22. Kokko H, Ots I. 2006 When not to avoid inbreeding. *Evol.* **60**, 467–475. (doi:10.1111/j.0014-
3820.2006.tb01128.x)

23. Daniels SJ, Walters JR. 2000 Inbreeding depression and its effects on natal dispersal in red-cockaded
woodpeckers. *The Condor* **102**, 482–491. (doi:10.1093/condor/102.3.482)

24. Foerster K, Valcu M, Johnsen A, Kempenaers B. 2006 A spatial genetic structure and effects of
relatedness on mate choice in a wild bird population. *Molecular Ecology* **15**, 4555–4567.
(doi:10.1111/j.1365-294X.2006.03091.x)

25. Frommen JG, Bakker TCM. 2006 Inbreeding avoidance through non-random mating in sticklebacks.
Biology Letters **2**, 232–235. (doi:10.1098/rsbl.2005.0432)

26. Gibbs HL, Grant PR. 1989 Inbreeding in Darwin's medium ground finches (Geospiza fortis). *Evolution*
43, 1273–1284. (doi:10.1111/j.1558-5646.1989.tb02574.x)

27. Szulkin M, Zelazowski P, Nicholson G, Sheldon BC. 2009 Inbreeding avoidance under different null
models of random mating in the great tit. *Journal of Animal Ecology* **78**, 778–788.

28. Wheelwright NT, Mauck RA. 1998 Philopatry, natal dispersal, and inbreeding avoidance in an island
population of savannah sparrows. *Ecology* **79**, 755–767. (doi:10.1890/0012-
9658(1998)079[0755:PNDAIA]2.0.CO;2)

29. Beauchamp G. 2014 Do avian cooperative breeders live longer? *Proceedings of the Royal Society B:
Biological Sciences* **281**, 20140844. (doi:10.1098/rspb.2014.0844)

30. Arnold KE, Owens IPF. 1998 Cooperative breeding in birds: a comparative test of the life history
hypothesis. *Proc. R. Soc. Lond. B* **265**, 739–745. (doi:10.1098/rspb.1998.0355)

31. Fitzpatrick JW, Bowman R. 2016 Florida scrub-jays: Oversized territories and group defense in a fire-
maintained habitat. In *Cooperative Breeding in Vertebrates* (eds WD Koenig, JL Dickinson), pp. 77–
96. Cambridge University Press. (doi:10.1017/CBO9781107338357.006)

32. Woolfenden GE, Fitzpatrick JW. 1984 *The Florida Scrub Jay: Demography of a Cooperative-breeding Bird*. Princeton University Press.

33. Suh YH, Pesendorfer MB, Tringali A, Bowman R, Fitzpatrick JW. 2020 Investigating social and environmental predictors of natal dispersal in a cooperative breeding bird. *Behav. Ecol.* **31**, 692–701. (doi:10.1093/beheco/araa007)

34. Aguillon SM, Fitzpatrick JW, Bowman R, Schoech SJ, Clark AG, Coop G, Chen N. 2017 Deconstructing isolation-by-distance: The genomic consequences of limited dispersal. *PLoS Genet.* **13**, e1006911. (doi:10.1371/journal.pgen.1006911)

35. Marzluff JM, Woolfenden GE, Fitzpatrick John W, Balda RP. 1996 Breeding partnerships of two New World jays. In *Partnerships in Birds: The Study of Monogamy* (eds JM Black, M Hulme), p. 0. Oxford University Press. (doi:10.1093/oso/9780198548614.003.0007)

36. Fitzpatrick JW, Woolfenden GE, Bowman R. 1999 Dispersal distance and its demographic consequences in the Florida Scrub-jay. In *Proceedings of the International Ornithological Congress*,

37. Townsend AK, Bowman R, Fitzpatrick JW, Dent M, Lovette IJ. 2011 Genetic monogamy across variable demographic landscapes in cooperatively breeding Florida scrub-jays. *Behavioral Ecology* **22**, 464–470. (doi:10.1093/beheco/arq227)

38. R Core Team. 2022 R: A Language and Environment for Statistical Computing.

39. Suh YH, Bowman R, Fitzpatrick JW. 2022 Staging to join non-kin groups in a classical cooperative breeder, the Florida scrub-jay. *J. Anim. Ecol.* **91**, 970–982. (doi:10.1111/1365-2656.13669)

40. Pebesma E. 2018 Simple features for R: Standardized support for spatial vector data. *The R Journal* **10**, 439. (doi:10.32614/RJ-2018-009)

41. Malécot G. 1948 *Les mathématiques de l'hérédité*. Paris: Masson.

42. Davison AC, Hinkley DV. 1997 *Bootstrap Methods and Their Application*. Cambridge University Press.

43. Mumme RL, Bowman R, Pruett MS, Fitzpatrick JW. 2015 Natal territory size, group size, and body mass affect lifetime fitness in the cooperatively breeding Florida Scrub-Jay. *Auk* **132**, 634–646. (doi:10.1642/AUK-14-258.1)

44. McGowan KJ, Woolfenden GE. 1990 Contributions to fledgling feeding in the Florida Scrub-Jay. *Journal of Animal Ecology* **59**, 691–707. (doi:10.2307/4889)

45. Schoech SJ, Mumme RL, Wingfield JC. 1996 Delayed breeding in the cooperatively breeding Florida scrub-jay (Aphelocoma coerulescens): inhibition or the absence of stimulation? *Behav Ecol Sociobiol* **39**, 77–90. (doi:10.1007/s002650050269)

46. Summers J, Cosgrove EJ, Bowman R, Fitzpatrick JW, Chen N. 2024 Density dependence maintains long-term stability despite increased isolation and inbreeding in the Florida Scrub-Jay. *Ecology Letters* **27**, e14483. (doi:10.1111/ele.14483)

47. Brommer JE, Merilä J, Kokko H. 2002 Reproductive timing and individual fitness. *Ecology Letters* **5**,
802–810. (doi:10.1046/j.1461-0248.2002.00369.x)

48. Barve S, Riehl C, Walters EL, Haydock J, Dugdale HL, Koenig WD. 2021 Lifetime reproductive
benefits of cooperative polygamy vary for males and females in the acorn woodpecker (Melanerpes
formicivorus). *Proceedings of the Royal Society B: Biological Sciences* **288**, 20210579.
(doi:10.1098/rspb.2021.0579)

49. Shah SS, Rubenstein DR. 2022 Prenatal environmental conditions underlie alternative reproductive
tactics that drive the formation of a mixed-kin cooperative society. *Sci. Adv.* **8**, eabk2220.
(doi:10.1126/sciadv.abk2220)

50. Venables W, Ripley B. 2002 *Modern Applied Statistics with S*. 4th edn. New York: Springer. See
https://www.stats.ox.ac.uk/pub/MASS4/.

51. Therneau T. 2022 A Package for Survival Analysis in R.

52. Therneau T. 2022 coxme: Mixed Effects Cox Models. R package.

53. Cockburn A, Osmond HL, Mulder RA, Green DJ, Double MC. 2003 Divorce, dispersal and incest
avoidance in the cooperatively breeding superb fairy-wren *Malurus cyaneus* . *J. Anim. Ecol.* **72**, 189–
202. (doi:10.1046/j.1365-2656.2003.00694.x)

54. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using lme4. *J.*
Stat. Softw. **67**, 1–48. (doi:10.18637/jss.v067.i01)

55. Schielzeth H. 2010 Simple means to improve the interpretability of regression coefficients. *Methods*
Ecol. Evol. **1**, 103–113. (doi:10.1111/j.2041-210X.2010.00012.x)

56. Fox J, Monette G. 1992 Generalized Collinearity Diagnostics. *Journal of the American Statistical*
Association **87**, 178–183.

57. Pusey A, Wolf M. 1996 Inbreeding avoidance in animals. *Trends Ecol. Evol.* **11**, 201–206.
(doi:10.1016/0169-5347(96)10028-8)

58. Leedale AE, Simeoni M, Sharp SP, Green JP, Slate J, Lachlan RF, Robinson EJH, Hatchwell BJ. 2020
Cost, risk, and avoidance of inbreeding in a cooperatively breeding bird. *Proceedings of the National*
Academy of Sciences **117**, 15724–15730. (doi:10.1073/pnas.1918726117)

59. Double MC, Peakall R, Beck NR, Cockburn A. 2005 Dispersal, philopatry, and infidelity: dissecting
local genetic structure in superb fairy-wrens (*Malurus cyaneus*). *Evol.* **59**, 625–635.

60. Leedale AE, Sharp SP, Simeoni M, Robinson EJH, Hatchwell BJ. 2018 Fine-scale genetic structure
and helping decisions in a cooperatively breeding bird. *Mol. Ecol.* **27**, 1714–1726.
(doi:10.1111/mec.14553)

61. Kokko H, Ekman J. 2002 Delayed dispersal as a route to breeding: territorial inheritance, safe havens,
and ecological constraints. *Am. Nat.* **160**, 468–484. (doi:10.1086/342074)

62. Green JP, Hatchwell BJ. 2018 Inclusive fitness consequences of dispersal decisions in a cooperatively breeding bird, the long-tailed tit (*Aegithalos caudatus*). *Proc. Natl. Acad. Sci. U.S.A.* **115**, 12011–12016. (doi:10.1073/pnas.1815873115)

63. Maag N, Cozzi G, Bateman A, Heistermann M, Ganswindt A, Manser M, Clutton-Brock T, Ozgul A. 2019 Cost of dispersal in a social mammal: body mass loss and increased stress. *Proc. R. Soc. B.* **286**, 20190033. (doi:10.1098/rspb.2019.0033)

64. Payne RB, Payne LL. 1993 Breeding dispersal in indigo buntings: Circumstances and consequences for breeding success and population structure. *The Condor: Ornithological Applications* **95**, 1–24. (doi:10.2307/1369382)

65. Forero MG, Donázar JA, Blas J, Hiraldo F. 1999 Causes and consequences of territory change and breeding dispersal distance in the black kite. *Ecology* **80**, 1298–1310. (doi:10.1890/0012-9658(1999)080[1298:CACOTC]2.0.CO;2)

66. Winkler DW, Wrege PH, Allen PE, Kast TL, Senesac P, Wasson MF, Llambías PE, Ferretti V, Sullivan PJ. 2004 Breeding dispersal and philopatry in the tree swallow. *The Condor: Ornithological Applications* **106**, 768–776. (doi:10.1093/condor/106.4.768)

67. Fuirst M, Strickland D, Norris DR. 2021 Breeding dispersal in a resident boreal passerine can lead to short- and long-term fitness benefits. *Ecosphere* **12**. (doi:10.1002/ecs2.3747)

68. Bonte D *et al.* 2012 Costs of dispersal. *Biol. Rev.* **87**, 290–312. (doi:10.1111/j.1469-185X.2011.00201.x)

69. Fuirst M, Strickland D, Freeman NE, Sutton AO, Ryan Norris D. 2023 Early-life sibling conflict in Canada jays has lifetime fitness consequences. *Proceedings of the Royal Society B: Biological Sciences* **290**, 20221863. (doi:10.1098/rspb.2022.1863)

70. Hui C, Roura-Pascual N, Brotons L, Robinson RA, Evans KL. 2012 Flexible dispersal strategies in native and non-native ranges: environmental quality and the 'good–stay, bad–disperse' rule. *Ecography* **35**, 1024–1032. (doi:10.1111/j.1600-0587.2012.07697.x)

71. Jennions MD, Hunt J, Graham R, Brooks R. 2004 No evidence for inbreeding avoidance through postcopulatory mechanisms in the black field cricket, teleogryllus commodus. *Evolution* **58**, 2472–2477. (doi:10.1111/j.0014-3820.2004.tb00876.x)

72. Collet M, Amat I, Sauzet S, Auguste A, Fauvergue X, Mouton L, Desouhant E. 2020 Insects and incest: Sib-mating tolerance in natural populations of a parasitoid wasp. *Molecular Ecology* **29**, 596–609. (doi:10.1111/mec.15340)

73. Holand Ø, Askim KR, Røed KH, Weladji RB, Gjøstein H, Nieminen M. 2006 No evidence of inbreeding avoidance in a polygynous ungulate: the reindeer (Rangifer tarandus). *Biology Letters* **3**, 36–39. (doi:10.1098/rsbl.2006.0575)

74. Geffen E *et al.* 2011 Kin encounter rate and inbreeding avoidance in canids. *Molecular Ecology* **20**, 5348–5358. (doi:10.1111/j.1365-294X.2011.05358.x)

75. Puurtinen M. 2011 Mate choice for optimal (k)inbreeding. *Evolution* **65**, 1501–1505. (doi:10.1111/j.1558-5646.2010.01217.x)

607 **Figures**

609
Figure 1. For both male (blue) and female (orange) Florida Scrub-Jays, mates at shorter distances from
both their natal ("First Pairings") and breeding territories ("Later Pairings") are more likely to be close
relatives. Lines indicate model fits, shaded areas 95% confidence intervals, and boxplots indicate
distribution of raw data, with points indicating outliers. We defined close relatives as opposite sex
individuals that are at least as closely related as first cousins to the focal individuals (1 = close relative, 0 =
not close relative). We included all instances where individuals become breeders for the first time that year
or switched mates (*N*: Male, First pairing = 421; Male, Later pairing = 249; Female, First pairing = 391;
Female, Later pairing = 256 instances).

618
619
Figure 2. Observed and expected (under random mating) (A) dispersal distance and (B) relatedness to mate
for male (blue) and female (orange) Florida Scrub-Jays at both the first (unfilled bars) and later (filled bars)
pairing stages. Histograms show the distributions of expected means under 1000 simulations of random
mating. The observed population means across all 32 years of our study period are indicated by vertical
lines (solid = significant difference, dashed = no significant difference from expectations under a random
mating scenario). Observed distances moved were significantly lower than expected for all four categories,
while kinship coefficient to mate was higher than expected for both sexes at the later, but not first, pairing
stages.

629
Figure 3. Predicted model fits investigating the effects (solid line = significant, dashed line = not
significant; shaded area = 95% confidence interval) of (A) Distance moved for first pairing, and (B)
Weighted kinship coefficients to mates on lifetime reproductive success (LRS) for Florida Scrub Jay male
(blue, triangles) and female (orange, circles) breeders with lifetime data between 1990 and 2021 (*N*: Males
= 196, Females = 122), and (C) Total distance moved for later pairings, and (D) Weighted kinship
coefficients to mates for the subset of breeders that switched mates at least once in their lifetime (*N*: Male
= 92, Female = 51). LRS is quantified as the number of offspring surviving to adulthood. Breeding lifespan
(not shown) was also included as a fixed effect in the model, since the longer a breeder lives the more
chances it has to accrue greater LRS. Females had reduced LRS with greater total distance moved for later
pairings, while males showed a relatively similar reduction in LRS with higher kinship to mates over their
lifetime. Points are jittered both horizontally and vertically for better visual discrimination.
641

Table 1. Results of generalized linear mixed models estimating the effect of distance moved for first pairings, later pairings, relatedness to mate, and breeding lifespan on lifetime reproductive success (*i.e.*, number of offspring surviving to adulthood) of Florida Scrub-Jays breeders. Significant effects are in bold.

Dataset	Sex	Fixed Effect	Estimate	Std Err	z-value	p-value
All breeders with lifetime data between 1990-2021	Male (*N*=196)	**Intercept**	0.40	0.09	4.69	**<0.001**
		Distance moved for first pairing	-0.03	0.11	-0.27	0.78
		Weighted kinship coefficient to mates	**-0.39**	**0.13**	**-2.93**	**0.003**
		Breeding Lifespan	**0.94**	**0.07**	**13.01**	**<0.001**
	Female (*N*=122)	**Intercept**	**0.65**	**0.10**	**6.61**	**<0.001**
		Distance moved for first pairing	-0.14	0.08	-1.74	0.08
		Weighted kinship coefficient to mates	0.05	0.08	0.57	0.57
		Breeding Lifespan	**0.86**	**0.10**	**9.00**	**<0.001**
Breeders with lifetime data between 1990-2021 that switched mates at least once	Male (*N*=92)	**Intercept**	**1.18**	**0.08**	**14.62**	**<0.001**
		Total distance moved for later pairings	-0.09	0.10	-0.88	0.38
		Weighted kinship coefficient to mates	**-0.29**	**0.09**	**-3.13**	**<0.001**
		Breeding Lifespan	**0.60**	**0.07**	**8.22**	**<0.001**
	Female (*N*=51)	**Intercept**	**1.13**	**0.11**	**9.94**	**<0.001**
		Total distance moved for later pairings	**-0.28**	**0.12**	**-2.42**	**0.02**
		Weighted kinship coefficient to mates	0.06	0.16	0.36	0.72
		Breeding Lifespan	**0.75**	**0.12**	**6.46**	**<0.001**