Lifetime fitness benefits of short-distance dispersal are associated with inbreeding tolerance despite multiple inbreeding avoidance mechanisms

Shailee S. Shah[1,2*,] Jennifer Diamond[3], Sahas Barve[4], Elissa J. Cosgrove[5], Reed Bowman[4], John W. Fitzpatrick[2,6], Nancy Chen[1,7*]

[1] Department of Biology, University of Rochester, Rochester, NY 14627, USA
[2] Cornell Lab of Ornithology, Cornell University, Ithaca, NY 14850, USA
[3] Department of Evolution and Ecology, University of California, Davis, Davis, CA 95616, USA
[4] Avian Ecology Program, Archbold Biological Station, Venus, FL 33960, USA
[5] Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14850, USA
[6] Department of Ecology & Evolutionary Biology, Cornell University, Ithaca, NY 14850, USA
[7] Department of Ecology & Evolutionary Biology, University of California, Los Angeles, Los Angeles, CA 90095, USA

* Correspondence: shailee.shah@cornell.edu, popgenchen@ucla.edu

Acknowledgments: We thank Archbold Biological Station and its staff for continued support and management of the study area; all interns, students, and postdocs who have worked on collecting and managing the long-term data; the Cornell Statistical Consulting Unit for statistical support; V. Sclater for GIS support; and F. Romero, J. Summers, D. Seidman, and other members of the Chen Lab at the University of Rochester for invaluable feedback on previous versions of this manuscript.

Funding: This work was supported by National Science Foundation (NSF) Grants DEB0855879 and DEB1257628 and by the Cornell Lab of Ornithology Athena Fund. S.S.S was supported by NIH grant 1R35GM133412 to N.C. and NSF Postdoctoral Research Fellowship in Biology 2305705.

Title: Lifetime fitness benefits of short-distance dispersal are associated with inbreeding tolerance despite

multiple inbreeding avoidance mechanisms

Abstract

Breeding with relatives often decreases individual fitness but can be mitigated via inbreeding avoidance

mechanisms such as dispersal, mate-switching, and kin avoidance. Disentangling the effects of multiple

mechanisms is challenging without detailed individual-level data and incomplete without considering all

life stages. We leveraged 32 years of data from a population of Florida Scrub-Jays (*Aphelocoma*

coerulescens) to investigate how dispersal and mate choice contribute to inbreeding risk across individuals'

lifetimes and quantify the associated fitness outcomes. We find that when mating is random, inbreeding is

two-fold higher when dispersal is limited, but adding sex-biased dispersal significantly reduces inbreeding

risk. Limited movement away from the breeding territory for later pairings (site fidelity) contributes more

to higher-than-expected levels of inbreeding than limited natal dispersal. Within observed dispersal

constraints, however, inbreeding is lower than expected under random mating, suggesting Florida Scrub-

Jays actively avoid mating with relatives, especially first-order kin. Finally, males paired with close

relatives and females who move farther for later pairings experience equivalently lower lifetime

reproductive success. Breeders who move farther for first pairings have lower survival. Overall, our results

show how variation in fitness tradeoffs and mate choice behavior across life stages affects inbreeding in a

long-lived, social species.

Keywords: dispersal**,** mate choice, inbreeding tolerance, inbreeding avoidance, site fidelity, lifetime

fitness, Florida Scrub-Jay

Main Text

Introduction

Matings among relatives, or inbreeding, have been shown to sometimes have negative fitness consequences in both experimental and observational studies (Jiménez et al., 1994; Keller et al., 1994; Coltman et al., 1999; Chen et al., 2016; Szulkin et al., 2007; C. W. Fox et al., 2008). Consequently, it is commonly assumed that inbreeding depression will lead to the evolution of reproductive strategies for inbreeding avoidance such as dispersal, mate switching (*i.e.*, "divorce"), and kin avoidance via mechanisms of kin recognition (Pusey & Wolf, 1996). However, evidence for such strategies is mixed (de Boer et al., 2021), in part because disentangling inbreeding avoidance strategies operating at different stages of mate choice is challenging (Szulkin et al., 2013).

In most species, moving away from the natal area (*i.e.*, natal dispersal) is a passive mechanism that adequately facilitates inbreeding avoidance (Perrin & Goudet, 2001). However, natal dispersal may be limited in some species, such that individuals are unable or disinclined to move far from their place of birth (Hamilton, 1964), often due to fitness benefits such as better access to resources (Pärt, 1994), higher survival (Griesser et al., 2006), and nepotism (Thünken et al., 2007). In such species, when individuals become breeders, they often choose mates that are geographically close (Greenwood & Harvey, 1982; Clutton-Brock & Lukas, 2012), though inbreeding risk can be mitigated if one sex disperses farther than the other (*i.e.*, sex-biased dispersal) (Greenwood & Harvey, 1982; Li & Kokko, 2019; Lawson Handley & Perrin, 2007). If natal dispersal is limited to some extent in both sexes, as is the case in many social species, active inbreeding avoidance mechanisms via mate choice, such as kin recognition and divorce, may additionally mitigate inbreeding risk (Cockburn et al., 2003; Nichols, 2017; Galezo et al., 2022; Leedale et al., 2024). Kin recognition for inbreeding avoidance in such cases could be largely based on early-life association via either geographic proximity or group membership (Riehl & Stern, 2015). However, active inbreeding avoidance behavior can also be costly if individuals have to forego breeding or spend more energy and incur higher fitness costs to find unrelated mates (Bengtsson, 1978; Kokko & Ots, 2006). Theory

suggests that if the fitness costs of inbreeding avoidance strategies outweigh the fitness costs of inbreeding,

inbreeding may instead be tolerated (Kokko & Ots, 2006; Nichols, 2017).

Inbreeding avoidance strategies may vary at different life stages. For example, moving away from

the breeding territory for any subsequent pairings may carry an increased mortality risk (Daniels & Walters,

2000a), whereas site fidelity (*i.e.*, remaining at or close to the same territory across breeding seasons) can

lead to fitness benefits due to familiarity and reduced competitive interactions (Schmidt, 2014; Beck et al.,

2020; Byrne et al., 2022). In a kin-structured population, with short natal dispersal distances, such site

fidelity for later pairings can also increase inbreeding risk (Nelson-Flower et al., 2012; van Dijk et al.,

2015), but if the fitness benefits of site fidelity outweigh the costs of inbreeding during later pairings,

inbreeding tolerance may be higher for individuals at later life stages. Further, since distance moved for

first pairing tends to be longer than distance moved for later pairings (Paradis et al., 1998), mate choice at

later life stages could potentially contribute more to inbreeding. Disentangling inbreeding avoidance

strategies thus requires examining events across an individual's lifetime, not just at the natal dispersal and

first mate choice stage.

A rigorous test of inbreeding tolerance or avoidance requires five key components: (i) a study

system with sufficient variation in relatedness between potential mates, the ability to identify both (ii) actual

and (iii) potential mates, (iv) accurate measures of relatedness, and (v) an accounting of mate choice events

occurring throughout different stages of an individual's lifetime (Szulkin et al., 2013). Though previous

studies of inbreeding have incorporated one or several of these components, to date no single study meets

all the requirements (Gibbs & Grant, 1989; Wheelwright & Mauck, 1998; Daniels & Walters, 2000b;

Foerster et al., 2006; Frommen & Bakker, 2006; Szulkin et al., 2009; Billing et al., 2012; Galezo et al.,

2022). Additionally, most studies lack measures of lifetime fitness. Reproductive strategies over an

individual's lifetime, especially in long-lived species, may compensate for any short-term fitness costs

associated with inbreeding or inbreeding avoidance strategies (Van de Casteele et al., 2003). Thus, we need

more studies that explicitly consider dispersal and mate choice decisions throughout an individual's lifetime

and lifetime fitness consequences of different strategies to better understand how dispersal and mate choice

decisions contribute to inbreeding risk in natural populations.

Here, we leverage 32 years of individual-level data on dispersal, mating, survival, and reproductive

success from a continuously monitored population of Florida Scrub-Jays (*Aphelocoma coerulescens*) to test

for contributions of dispersal and mate choice across an individual's lifetime to inbreeding risk and quantify

the associated lifetime fitness outcomes. Florida Scrub-Jays are avian cooperative breeders that defend

year-round territories in a social group comprising a dominant breeding pair and non-breeding subordinates,

the majority being first-order kin, that remain as helpers for one or more years before dispersing to breed

(Woolfenden & Fitzpatrick, 1984). Though females disperse farther than males, neither sex typically

disperses more than 2.5 km away (or the equivalent of five territories) (Suh et al., 2020), such that the

relatedness of potential mates within a population range from parents/full-siblings to unrelated immigrants

(Aguillon et al., 2017). Florida Scrub-Jays pair with a new mate after their current mate dies (Woolfenden

& Fitzpatrick, 1984), or in rare cases when the mate is still alive (*i.e.*, divorce) (Marzluff et al., 1996). Here,

we focus on a population of Florida Scrub-Jays that has been intensively studied at Archbold Biological

Station in central Florida since 1969 (Woolfenden & Fitzpatrick, 1984). Detailed monitoring data from this

population provides the unique ability to identify all actual and potential mates, calculate accurate

relatedness measures using a single, extensive pedigree, incorporate mate choice events throughout

individuals' lifetimes, and quantify lifetime fitness in terms of reproductive success and survival

(Fitzpatrick & Bowman, 2016). Moreover, previous work has found evidence of inbreeding depression in

Florida Scrub-Jays, suggesting that inbreeding avoidance strategies may be adaptive in this species (Chen

et al., 2016), though long-distance dispersal may also have associated fitness costs (Fitzpatrick et al., 1999).

We performed a series of analyses to quantify the effects of dispersal and mate choice decisions on

inbreeding risk across sexes and life stages. First, we compared the observed distance moved for pairing

and levels of inbreeding (relatedness to mate) to simulated means generated under three random mating

scenarios: (i) no constraints on dispersal distance, (ii) limited dispersal distance with existing sex bias

included, and (iii) limited dispersal distance with any sex bias removed. As limited dispersal is expected to

increase inbreeding risk, we predicted that observed inbreeding will be higher than expected under the

unconstrained model (scenario i). Further, we predicted that if, when dispersal is limited, sex-biased

dispersal helps reduce inbreeding risk, simulated relatedness to mates will be lower in the model with sex-

biased dispersal (scenario ii) than in the model without (scenario ii). In fact, if sex-biased dispersal is

sufficient to overcome increased inbreeding risk due to limited dispersal, simulated relatedness to mates

should be equivalent between the sex-biased, limited dispersal model and unconstrained dispersal model.

Finally, we predicted that in addition to passive mechanisms, if Florida Scrub-Jays actively avoid mating

with kin, observed inbreeding should be lower than expected under the limited, sex-biased dispersal model.

If such kin avoidance is facilitated by kin recognition based on familiarity of individuals within the natal

social group, the observed proportion of first-order kin mates should be lower than that expected under the

limited, sex-biased dispersal model. Additionally, we tested whether divorce is another mate choice strategy

to mitigate inbreeding risk. We predicted that pairs with more closely related mates will be more likely to

dissolve because of divorce than death of a mate.

113   Finally, to investigate the relative fitness costs and benefits associated with inbreeding and

inbreeding avoidance strategies, we quantified the impact of distance moved and inbreeding on breeder

survival and lifetime reproductive success (LRS). We predicted that, for both sexes, both inbreeding and

longer dispersal distances at all life stages would result in lower LRS and survival. Further, we predicted

that if we found no evidence of active inbreeding avoidance, the fitness cost of inbreeding will be lower

than the cost of long-distance dispersal. Conversely, if individuals are actively avoiding inbreeding, the

fitness costs of inbreeding will be higher than or equivalent to the cost of long-distance dispersal.

Ultimately, we delineate passive and active mechanisms of inbreeding avoidance, and quantify the

associated fitness outcomes, to provide the first complete picture of how dispersal patterns and mate choice

contribute to—or mitigate—inbreeding risk in a natural population experiencing inbreeding depression.

Methods

Data collection

Florida Scrub-Jays at Archbold Biological Station (27.10°N, 81.21°W) have been continuously monitored

since 1969, with the study tract expanded to its current size (mean ± sd = 914.04 ± 143.44 ha) by 1990

(Woolfenden & Fitzpatrick, 1984). All birds are banded with unique combinations of color bands.

Throughout the breeding season (early March to early June), all nests are located and monitored for fledging

or failure. All territories are mapped in April, with territory boundaries assigned at locations of territorial

disputes between neighboring groups. In addition to nest monitoring, the population is censused each

month, resulting in detailed data on dispersal and survival of both fledglings and adults. Pairs are socially

monogamous, with very low rates of extra pair-paternity (<1%) (Townsend et al., 2011), allowing accurate

parentage inference from field observations alone. Furthermore, previous work verified genetic parentage

of offspring born between ~1989-1991, 1994-1995, and 1999-2013 (Chen et al., 2016). Using these detailed

data, we have accumulated a 15-generation pedigree of known-age, known-parentage individuals with

precise measurements of lifespan and reproductive success. For this study, we used data from 1990-2021,

which included on average (± SD) 71.4 (± 14.7) territories (area mean ± sd =13.24 ± 6.37 ha) per year.

Fieldwork was approved by the Institutional Animal Care and Use Committees at Cornell University

(IACUC 2010-0015) and Archbold Biological Station (AUP-006-R) under permits from the US Fish and

Wildlife Service (TE824723-8, TE-117769), the US Geological Survey (banding permits 07732, 23098),

and the Florida Fish and Wildlife Conservation Commission (LSSC-10-00205).

Observed mates

Our dataset of actual mates included all new breeding pairs formed between 1990 and 2021 (*N* = 1077).

New breeding pairs include pairs formed by two first-time breeders, one first-time and one experienced

breeder, or two experienced breeders. For first-time breeders, we called the event a "first pairing" and for

experienced breeders, a "later pairing". The first pairing for one mate is thus not always the first pairing for

the other mate (*e.g.*, an experienced male could repair with a female who is a first-time breeder, and vice

versa). New pairs might occupy a new territory, which may be inherited by one of the two breeders or

formed as an extension of their natal territory in a process called "budding" (Woolfenden & Fitzpatrick,

1984; Fitzpatrick & Bowman, 2016). Alternatively, a new pair might form on a non-natal territory already

occupied by one of the two breeders, where a breeding vacancy is typically formed due to the death of a

breeder, but occasionally via divorce " (Woolfenden & Fitzpatrick, 1984; Fitzpatrick & Bowman, 2016).

Thus, both first and later pairings can occur without any movement away from the currently occupied

territory by at least one of the two breeders.

Random mating models

To delineate passive and active mechanisms of inbreeding avoidance, we constructed three random mating

scenarios with (i) no constraints on dispersal distance, (ii) limited dispersal distance with existing sex bias

included, and (iii) limited dispersal distance without sex bias (Figs. S1-S2). For the unconstrained model,

we defined potential mates for all individuals of one sex in a given year as all individuals of the opposite

sex from our dataset of observed mates (*i.e.*, individuals that were both available as potential mates and

successful at finding new mates) (*N* focal individuals: mean ± SD = 25.4 ± 7.06; *N* available mates per

individual: mean ± SD = 25.5 ± 7.01; see *SI* for more details). For the limited dispersal with sex-biased

dispersal model, we limited available mates to opposite sex individuals at the observed dispersal distance

of each individual (*N* available mates per individual: mean ± SD = 1.37 ± 0.63). Finally, for the limited

dispersal but no sex bias model, we randomly shuffled observed dispersal distances values across sexes,

but within life stages (*i.e.*, first or later pairing). We then limited available mates for each focal individual

to opposite sex individuals at the assigned dispersal distance (*N* available mates per individual: mean ± SD

= 1.39 ± 0.68). For each focal individual we then randomly assigned an available mate for each year. These

random assignments were permuted 1000 times. In dispersal constrained models, individuals that did not

move away from their territory but paired with a new mate in a given year were assigned the dispersal

distance value of their new mate. While both mates may move territories to form a new breeding pair (107

out of 1077 [10%] observations), our simulations are sex-specific and assume that only the focal individual

moved to the territory occupied by the potential mate.

Distance moved for pairing

We measured the distance moved for pairing as the number of territories crossed, which is the most

biologically relevant measure of distance moved (Fitzpatrick et al., 1999; Greenwood & Harvey, 1982).

The number of territories crossed was highly correlated with distance moved in meters (Pearson's $r = 0.91$;

(Fitzpatrick et al., 1999)). We drew a straight-line movement trajectory from the centroid of an individual's

previous territory to the centroid of its new breeding territory. We then used the *st_intersects* function from

the R package *sf* (Pebesma, 2018) to calculate the number of territories (i.e., polygons) that each line

intersected (see *SI* for more details). When the focal individual remained on its natal territory and became

a breeder, formed a territory by budding, or remained on its non-natal breeding territory, we quantified

distance moved as 0 m.

To obtain accurate measures of distance moved, we excluded any pairings that involved movement

that started or ended outside of the study tract. We also excluded any movement that did not end with the

individual attaining a breeding position (*i.e.*, staging, a strategy adopted by about 28% of yearlings (Suh et

al., 2022)). For staging individuals that eventually became breeders, we used movement from their staging

to breeding territory to calculate distance moved for first pairing. However, using distance from natal to

breeding territory as distance moved for first pairing did not change our results (see *SI*, Fig. S3, Table S1).

Relatedness to mate

To quantify levels of inbreeding, we calculated the kinship coefficient between mates for all observed and

simulated breeding pairs using the population pedigree with the kinship function in the R package *kinship2*

(Sinnwell et al., 2014). With no inbreeding, the kinship coefficient ranges from 0 to 0.5, such that 0.5 is an

individual's kinship coefficient with itself; 0.25 is the average kinship coefficient between siblings and

between parents and offspring (first-order kin); 0.125 is the average kinship coefficient between half-

siblings, grandparents-grandchildren, and aunt/uncle-niece/nephew pairs (second-order kin); and 0.0625 is

the average kinship coefficient between first cousins (third-order kin) (Malécot, 1948). Pedigree-based

inbreeding estimates are correlated with genomic-based estimates in our study population, confirming that

our pedigree is complete enough to provide accurate measures of inbreeding and relatedness (Chen et al.,

2016).

Lifetime fitness metrics

We assessed breeder survival using monthly census and capture records and calculated breeding lifespan

as the number of years from becoming a breeder to death. We calculated lifetime reproductive success

(LRS) for all breeders as the total number of offspring produced over an individual's lifetime that survived

to adulthood, *i.e.*, age 1 year, the stage at which Florida Scrub-Jays become physiologically capable of

breeding (McGowan & Woolfenden, 1990; Mumme et al., 2015; Schoech et al., 1996; Woolfenden &

Fitzpatrick, 1984). Since the mean growth rate of our study population is close to 1 (0.998) (Summers et

al., 2024), we deemed LRS an appropriate metric of individual fitness (Brommer et al., 2002). For all fitness

analyses, we restricted our dataset to individuals from cohorts from which all members born within the

study tract were last observed before 2021 to eliminate bias towards short-lived individuals (Barve et al.,

2021; Shah & Rubenstein, 2022). The final dataset for all fitness analyses included individuals born

between 1988-2006 and in 2009 (N: Males = 197, Females = 120).

Data analysis

All analyses were conducted using R version 4.2.2 (R Core Team, 2022). Fixed effects for all models were

standardized using z scores (Schielzeth, 2010), and the variance inflation factor was < 2 for all fixed effects

in all GLMMs (J. Fox & Monette, 1992).

Effects of dispersal and mate choice on inbreeding: To quantify the contributions of dispersal and mate

choice behavior to levels of inbreeding, we compared observed population means of inbreeding (measured

as kinship coefficient to mates), aggregated across years, to the expected distributions generated by our

random mating simulations using empirical *p*-values (Davison & Hinkley, 1997). Specifically, we

quantified the proportion of simulations that had a mean value lesser than or equal (S_{low}) or greater than or

equal (S_{high}) to the observed mean. Empirical *p*-values were then calculated as ($S_{high/low}$+1)/(N+1), where N

was the total number of simulations (*i.e.,* 1000). For all random mating scenarios, we tested whether the

observed kinship coefficient and proportion of first-order kin pairs was greater or less than expected by

random chance across sexes and pairing types. We also compared the simulated kinship coefficients

between the limited dispersal models with and without sex-bias using *t*-tests, with Bonferroni corrections

to *p*-values to account for multiple testing.

Divorce as an inbreeding avoidance strategy: To investigate whether Florida Scrub-Jays compensated for

the impact of inbreeding on reproductive success via divorce, we fit GLMMs with the cause of pair

dissolution (1 = divorce, 0 = mate death) as the dependent variable, the kinship coefficient to mate as a

fixed effect, year and individual ID as random effects, and a binomial error structure. We also fit models

with a categorical fixed effect of kinship between mates (1 = first-order relative, 0 = not a first-order

relative) to account for cases where opposite-sex offspring might inherit dominant breeding positions,

resulting in mother-son or father-daughter pairings that may be more likely to end in divorce (Cockburn et

al., 2003). We fit separate GLMMs by sex and pairing type using the function *glmer* in the R package *lme4*

(*N*: Males, First pairing = 194; Males, Later pairing = 187; Females, First pairing = 163; Females, Later

pairing = 169 pairings) (Bates et al., 2015).

Lifetime fitness outcomes: First, we examined the effect of distance moved on breeder survival. We found

that fitting Cox proportional hazard models with distance moved for first pairing as a fixed effect violated

the proportional hazards assumption for both sexes (Males: χ^2 = 4.33, df = 1, P = 0.04; Females: χ^2 = 4.21,

df = 1, P = 0.04), suggesting that the effect of distance moved for first pairing on the hazard rate varies over

time. To account for this time-varying effect, we fit stratified Cox proportional hazards models with two

time intervals: ≤ 4 years and > 4 years from the start of breeding (Zhang et al., 2018). We tested varying

thresholds for splitting the analysis time into intervals and chose a cut time of 4 years based on visual

assessment of variation in the hazard ratio over time (see *SI*, Table S2, Fig. S4) (Zhang et al., 2018). We

used the *Surv* and *coxph* function from the R package *survival* (Therneau, 2022) to fit separate models by

sex of breeder, and incorporated natal year as a random effect (N: Males = 197, Females = 120). For the

subset of individuals who switched mates at least once in their lifetime (N: Males = 91, Females = 50), we

also examined the effect of territory loss (*i.e.*, total distance moved for later pairings ≥ 1 territory away) on

breeder survival. We fit Cox proportional hazards models with loss of territory (1 = territory lost, 0 =

territory retained) as a fixed effect and natal year as a random effect. The proportional hazards assumption

was met for these models (Males: χ^2 = 1.17, df = 1, P = 0.28; Females: χ^2 = 0.27, df = 1, P = 0.61).

Next, to examine how distance moved for pairing and inbreeding impacts lifetime fitness of Florida

Scrub-Jays, we fit two separate GLMMs by sex with negative binomial error distributions (to account for

overdispersion), LRS as the dependent variable, and distance moved for first pairing, weighted relatedness

to mates, and breeding lifespan as fixed effects (N: Males = 197, Females = 120). We calculated weighted

kinship coefficient to mates for all breeders across their lifetimes by multiplying the kinship coefficient of

the focal individual to each of their mates by the number of years they were paired with that mate and

summing the resulting values. Breeding lifespan was included in the model to account for the fact that the

more breeding seasons a breeder experiences, the more chances it has at successful reproduction

(Fitzpatrick & Bowman, 2016). We also included natal year as a random effect to account for variation

between birth cohorts.

Finally, to specifically examine the effect of moving away from the breeding territory, we also fit

two additional GLMMs with the same error structure, dependent variable, and random effects, with total

distance moved for later pairings, weighted kinship coefficient to mates, and breeding lifespan as fixed

effects using a truncated dataset that only included individuals that had switched mates in their lifetimes

(N: Males = 91, Females = 50). All GLMMs were fit using the function glmer.nb in the R package MASS

(Venables & Ripley, 2002). Model fits were checked using quantile-quantile plots of residuals (Fig. S5).

Fixed effects were not correlated (Table S3).

Results

Observed dispersal and mate choice patterns

The annual number of new breeding pairs ranged from 17-47 pairs with an average (± SD) of 29.2 (± 8.03) new pairs per year (including both first and later pairings). Of the individuals included in our lifetime fitness analysis, 42% of females (50 out of 120) and 46% of males (91 out of 197) had later pairings at least once in their lifetime. Surprisingly, only about half of all mates in later pairings were naïve breeders for both sexes (Males: 218 out of 427 [51%], Females: 213 out of 421 [51%] instances). Thus, not all later pairings were a result of subordinates moving in to fill a breeding vacancy—nearly half of such cases were a result of the mate also re-pairing because of death of, or divorce from, their previous mate. However, for breeders that retained their breeding territory for later pairings, ≥ 90% of new mates were from ≤ 2 territories away (Fig. S6). Furthermore, we found that out of all later pairing events observed, 18% were due to divorce, not mate death (Males = 33 out of 187; Females = 31 out of 169 pairings).

Females, on average, moved farther than males, though the effect size was greater for first pairings than later pairings (see *SI*, Fig. S1). In 197 out of 848 (23%) instances of later pairings the focal individual (62% of whom were females) moved one or more territories away from its established breeding territory. The kinship coefficient (*i.e.*, relatedness) to mate ranged from 0 (unrelated) to 0.25 (*i.e.*, either full siblings or parent-offspring) (Fig. S2). Across all years, only 43 out of 1077 (4%) of observed new breeding pairs were comprised of individuals who were at least as closely related as first cousins (*i.e.*, had a kinship coefficient ≥ 0.0625) and only 13 out of 1077 (1.2%) of observed breeding pairs comprised first-order relatives (7 mother-son, 3 father-daughter, and 3 full sibling pairs; Fig. S7). Relatedness to potential mates was significantly higher at closer geographical distances (see *SI*, Fig. S8).

Passive and active mechanisms of inbreeding and inbreeding avoidance

We found that limited dispersal is an important component of the mating pattern increasing inbreeding risk in Florida Scrub-Jays. Under a model of random mating with limited but not sex-biased dispersal, mean

relatedness to mates was two-fold or more higher for both sexes for all life stages than expected under a

model of random mating with unconstrained dispersal (Males, First Pairings: $t_{637757} = 68.12$, $P < 0.001$;

Females, First pairings: $t_{565627} = 72.94$, $P < 0.001$; Males, Later pairings: $t_{472441} = 82.40$, $P < 0.001$, Females,

Later pairings: $t_{503923} = 77.13$, $P < 0.001$; Fig. 1A-D). Sex-biased dispersal does help mitigate inbreeding

risk: the expected level of inbreeding across both sexes was significantly lower (by $\geq 50\%$) when we added

sex-biased dispersal for both sexes for first pairings (Males: $t_{527991} = -44.39$, $P < 0.001$; Females: $t_{520491} = -$

46.64, $P < 0.001$, Fig. 1A-B) and for females for later pairings ($t_{597723} = -13.39$, $P < 0.001$, Fig.1C), but the

pattern was reversed for males for later pairings ($t_{585009} = 34.9$, $P < 0.001$, Fig. 1D).

As predicted, we found that, compared to an unconstrained random mating scenario, both sexes

moved significantly shorter distances within the study tract ($P \leq 0.05$ for both, Fig. S9). Observed

relatedness to mates was higher than expected by chance under unconstrained random mating for both sexes

at all life stages, except males for first pairings where the difference approached significance (Males, First

pairings: $P = 0.07$; Females, First pairings: $P = 0.05$; Males, Later pairings: $P = 0.001$; Females, Later

pairings: $P = 0.003$; Fig. 1A-D). However, observed mean relatedness to mates remained significantly lower

across all sexes and life stages than expected under the limited, sex-biased dispersal scenario (Males, First

pairings: $P = 0.005$; Females, First pairings: $P < 0.001$; Males, Later pairings: $P = 0.03$; Females, Later

pairings: $P = 0.04$; Fig. 1A-D), indicating additional active inbreeding avoidance behavior.

We also found that observed proportions of pairing between first-order kin were not significantly

different from those expected under the unconstrained model (Males: $P = 0.43$, Females: $P = 0.32$, Fig.

1E-F) but were significantly lower than those expected under the limited, sex-biased dispersal model in

both sexes for first pairings (Males: $P < 0.001$, Females: $P = 0.008$, Fig. 1E-F). Conversely, for later

pairings, the observed proportions of first-order kin were significantly higher than expected under the

unconstrained model (Males: $P = 0.05$; Females: $P = 0.04$, Fig. 1G-H), and, for males, not significantly

different from expectations under the limited, sex-biased dispersal model (Males: $P = 0.28$, Females: $P =$

0.008, Fig. 1G-H). For second- and third-order kin mates, patterns were more variable across sexes and

dispersal types (Fig. S10). Overall, both sexes exhibited higher proportions of non-kin mates than expected

under the limited, sex-biased dispersal model for first pairings, but not for later pairings (Males, First

pairings: $P = 0.009$, Females, First pairings: $P = 0.04$, Males, Later pairings: $P = 0.10$, Females, Later

pairings: $P = 0.17$; Fig. S10).

Divorce as an inbreeding avoidance strategy

Relatedness to their mate had no effect on likelihood of divorce for either sex at either life stage (Males,

First pairing: $Z = 0.42$, $P = 0.68$; Males, Later pairing: $Z = 0.42$, $P = 0.68$; Females, First pairing: $Z = -0.84$,

$P = 0.40$; Females, Later pairing: $Z = 0.76$, $P = 0.45$), nor was divorce more likely in cases of pairing with

first-order relatives (Males, First pairing: $Z = 1.87$, $P = 0.06$; Males, Later pairing: $Z = 0.22$, $P = 0.83$;

Females, First pairing: No instances of divorce when paired with a first-order relative; Females, Later

pairing: $Z = 0.52$, $P = 0.60$).

Lifetime fitness outcomes

For both sexes, breeding lifespan was affected by distance moved for first pairing: longer distance moved

was associated with higher mortality in the first four years after breeding onset (Males: $Z = 2.16$, $P = 0.03$;

Females: $Z = 3.36$, $P = 0.001$), but the relationship was reversed in subsequent years (Males: $Z = -2.07$, P

$= 0.04$; Females: $Z = -2.02$, $P = 0.04$) (Fig. S11). Though the median breeding lifespan for both sexes was

4 years, 37% of individuals lived past their fourth year as a breeder (Males: 73 out of 197, Females: 45 out

of 120 individuals). We obtained qualitatively similar results when we varied the interval split times

between 5-7 years post breeding (see *SI*, Table S2; upper quartile of breeding lifespan: Males = 6, Females

= 7 years, Fig. S12). However, dispersal away from an established breeding territory in the event of later

pairing did not impact breeder survival for either sex (Males: $Z = -0.37$, $P = 0.71$; Females: $Z = 0.76$, $P =$

0.45).

We found that a higher weighted kinship coefficient to mates over an individual's lifetime was

associated with significantly fewer offspring that survived to adulthood for males (all breeding males: $Z =$

-2.83, $P = 0.005$; only males that switched mates at least once: $Z = -0.28$, $P = 0.09$; Fig. 2B, D; Table 1),

but not females (all breeding females: $Z = 0.31$, $P = 0.76$; only females that switched mates at least once:

$Z = -0.09$, $P = 0.97$; Fig. 2B, D; Table 1). The few males who bred with closely related females for the

majority of their lifetime had LRS close to zero (Fig. 2B, D), but we did not observe the same pattern in

females. Females, but not males, who moved farther distances for later pairings had significantly fewer

offspring who survived to adulthood (Males: $Z = -0.64$, $P = 0.52$, Females: $Z = -2.62$, $P = 0.01$; Fig. 2C,

Table 1). After controlling for the significant negative effect of breeding lifespan—which is lower for

individuals that move longer distances for first pairings—on lifetime reproductive success (Males: $Z = 0.93$,

$P < 0.001$, Females: $Z = 0.86$, $P < 0.001$; Table 1), we did not find an additional effect of distance moved

for first pairing on lifetime reproductive success for either sex (Males: $Z = -0.29$, $P = 0.77$, Females: $Z = -$

0.29, $P = 0.77$; Fig. 2A, Table 1).

Discussion

Testing for inbreeding avoidance using observational datasets can be difficult since the underlying

mechanisms can operate at different stages of mate choice (Szulkin et al., 2013) and their relative strength

can vary at different life stages (Nelson-Flower et al., 2012). Here, we leverage data on pedigree

relationships, dispersal, mate pairings, and fitness from a long-term, continuously monitored population of

Florida Scrub-Jays to disentangle the contribution of passive and active inbreeding avoidance strategies to

inbreeding risk across different life stages and quantify the associated lifetime fitness outcomes. First, using

random mating simulations with varying dispersal constraints, parametrized with empirical data, we show

that limited dispersal is expected to increase inbreeding by at least two-fold, and that sex-biased dispersal

plays an important role in mitigating this risk, especially at the first pairing stage. Though theoretical models

have shown this effect of sex-biased dispersal, to our knowledge this is one of the first empirical

demonstrations (see Blyton et al., 2015). Second, we find that observed inbreeding levels are higher than

expected under an unconstrained random mating model. Within the observed, sex-biased dispersal

limitations, however, observed relatedness to mates is lower than expected by chance for both sexes across

both life stages, suggesting that though Florida Scrub-Jays are not more likely to divorce when paired with

close relatives, they actively avoid pairing with them in the first place. This pattern can be largely explained

by avoidance of first-order kin during first pairings for both sexes and life stages, except males during later

pairings. The nearly two-fold higher propensity of males to remain on their breeding territory, compared to

females, for later pairings likely increases inbreeding risk. Finally, we find that dispersing farther, at both

life stages, and pairing with close relatives is associated with lower fitness, suggesting that Florida Scrub-

Jays balance the tradeoffs of dispersal and inbreeding such that some level of inbreeding is tolerated, in

accordance with theoretical expectations (Kokko & Ots, 2006). Overall, our results show how variation in

fitness tradeoffs and mate choice behavior across life stages affects inbreeding risk in a long-lived, social

species.

We find that Florida Scrub-Jays exhibit a degree of inbreeding tolerance due to limited dispersal at

all life stages. Although elevated risk of inbreeding and associated inbreeding depression should select for

longer dispersal distances, limited dispersal is found in both social and asocial species. Individuals that do

not disperse far may gain better access to resources and protection from predators due to prior local

experience (Pärt, 1994) or higher habitat quality (Suh et al., 2020), face lower competition due to nepotism

(Green & Hatchwell, 2018; Kokko & Ekman, 2002), and not incur the high physiological costs of dispersal

(Maag et al., 2019). As such, limited dispersal has been shown to increase levels of inbreeding in various

taxa (Keller et al., 1994; Costantini et al., 2007; Lubin et al., 2009; Lloyd et al., 2018; De Ro et al., 2021;

Ceresa et al., 2024). Similarly, our results suggest that limited natal dispersal in Florida Scrub-Jays results

in higher levels of inbreeding than expected under a model of unconstrained dispersal. We also considered

instances of all subsequent pairings with new mates at later life stages. We found that high site fidelity, *i.e.*

limited movement away from the breeding territory for later pairings, also results in higher levels of

inbreeding than expected under a scenario of unconstrained dispersal. In fact, for both sexes, the difference

between observed relatedness to mates and its expected distribution under random mating with

unconstrained dispersal is an order of magnitude higher for later pairings than first pairings. While studies

have rarely examined the contribution of dispersal patterns in later life stages to inbreeding, site fidelity to

lekking areas in red deer (*Cervus elaphus*) lead to inbreeding by increasing the chances of males mating

with females of the same matriline (Stopher et al., 2012). Thus, neglecting later stages of life excludes

demographic processes that can significantly contribute to inbreeding, especially in long-lived species.

Sex-biased dispersal is a passive mechanism that can mitigate inbreeding risk. The difference in

expected relatedness to mate under a limited dispersal scenario with and without sex-bias provides clear

evidence of the importance of sex-biased dispersal in lowering inbreeding risk in our study population

during the first pairing stage. However, Florida Scrub-Jay females still show considerable philopatry at this

stage, only dispersing on average ~2 territories away for first pairings. The likelihood of encountering a

closely related potential mate is thus still significantly higher within the average dispersal distance of both

sexes. These results add to a growing body of literature showing that sex-biased dispersal in social

vertebrates is not enough to mitigate inbreeding risk due to the opposing fitness benefits of remaining in

the same social group as, or close to, kin (Daniels & Walters, 2000b; Riehl & Stern, 2015; Kingma et al.,

2017; Morrison et al., 2023).

The sex-bias in distance moved for later pairings was less pronounced, but males were still twice

as likely as females to remain on their breeding territory for later pairings. A similar pattern of sex bias in

breeding territory fidelity is found in other avian taxa, though the underlying causes and fitness

consequences may differ (Harvey et al., 1979; Clarke et al., 1997). In Florida Scrub-Jays, females may be

more likely to lose their breeding territory when they switch mates (later pairing) as a result of social

dominance dynamics or avoidance of inbreeding with first-order kin if their son is more likely to inherit the

territory than daughters are for male breeders (Woolfenden & Fitzpatrick, 1984). Thus, site fidelity in males

may face stronger selective pressures resulting in reduced variation. Moreover, contrary to our predictions,

removing the sex bias in distance moved for later pairings, *i.e.*, relaxing the strong skew in males towards

site fidelity, decreased, rather than increased, inbreeding risk in males by 50% (though this decrease is

countered by a 17% increase in mean inbreeding for females). This result suggests that fitness tradeoffs

may shift over time in long-lived species, influencing dispersal patterns and inbreeding risk. Future studies

should investigate dispersal dynamics during later life stages in more detail in this and other cooperative

breeding, non-migratory species.

While passive mechanisms of inbreeding avoidance do not entirely mitigate inbreeding risk in

Florida Scrub-Jays, we found evidence of active kin avoidance given observed dispersal limitations. For

both sexes during first pairings, the proportion of observed first-order kin mates—*i.e.*, parent-offspring or

sibling—was significantly lower than expected under the limited dispersal, sex-biased model and was not

significantly different from expectations under the unconstrained model. These results suggest that active

inbreeding avoidance in Florida Scrub-Jays during first pairing might be based on strong avoidance of close

relatives based on familiarity. Florida Scrub-Jays live in cooperative breeding groups where the majority

of helpers are offspring from previous broods that delay dispersal for one or more years to act as helpers

(Woolfenden & Fitzpatrick, 1984). Thus, first-order kin have ample opportunity to build familiarity.

Additionally, in a cooperatively breeding species like the Florida Scrub-Jays, kin recognition may be more

likely to evolve, since it could serve an additional purpose in facilitating behaviors that result in indirect

fitness benefits from helping kin (Nichols, 2017). Similar patterns of kin avoidance have been shown in

some social species such as yellow baboons (*Papio cynocephalus*) (Galezo et al., 2022), mountain gorillas

(*Gorilla beringei beringei*) (Morrison et al., 2023), acorn woodpeckers (*Melanerpes formicivorus*)

(Haydock et al., 2001), long-tailed tits (*Aegithalos caudatus*) (Hatchwell et al., 2000; Leedale et al., 2020),

and superb fairy-wrens (*Malurus cyaneus*) (Cockburn et al., 2003).

Interestingly, we found that the pattern of kin avoidance was weaker for later pairings, with the

observed proportion of first-order kin mates being significantly higher than expected under the

unconstrained model for both sexes, and not significantly different from the limited, sex-biased dispersal

model for males. This result further suggests that, with a shift in fitness tradeoffs, inbreeding tolerance may

increase for Florida Scrub-Jays at later life stages. Future research into the mechanisms underlying sexual

selection and mate choice at all life stages in Florida Scrub-Jays, which still remain unknown, would help

clarify mechanisms of inbreeding avoidance while accounting for other factors underlying mate choice.

Finally, though divorce is a common active inbreeding avoidance strategy in birds (Hatchwell et

al., 2000; Cockburn et al., 2003; Hidalgo Aranzamendi et al., 2016), we did not find a similar pattern in

Florida Scrub-Jays. However, pairs of close relatives are relatively rare in our study population, with only

4% of all breeding pairs resulting in consanguineous matings and only 1.2% resulting in matings between

first-order relatives. Active kin avoidance may reduce the likelihood of first-order kin pairs forming in the

first place. Further, since reproductive success increases with pair experience in Florida Scrub-Jays

(Woolfenden & Fitzpatrick, 1984), mate switching due to divorce may prove costly.

Finally, we found that Florida Scrub-Jays experience fitness costs with longer distances moved at

all life stages. We found that breeder survival decreased with farther natal dispersal distances, though this

effect was limited to the first few years post breeding onset. For individuals with a breeding lifespan of >4

years, survival was greater for breeders that had moved farther for their first pairing, potentially because

individuals of both sexes that disperse from their natal territory at a younger age generally disperse farther

(Suh et al., 2020). We found that distance moved for first pairing did not affect LRS for either sex after

accounting for breeding lifespan, suggesting no further effect of dispersal distance on fitness beyond

reducing breeding lifespan. These results are consistent with previous work in this species and underscore

the importance of breeders survival on the lifetime fitness and contribution to population growth in Florida

Scrub-Jays (Fitzpatrick et al., 1999; Summers et al., 2024). Similarly, in long-lived species across different

taxa, breeder survivorship, rather than fecundity, has been shown to have a greater effect on individual

fitness and population growth (Brommer, 2000; Sæther & Bakke, 2000; McAdam et al., 2007).

Surprisingly, we did not find an effect of the loss of the breeding territory on breeder survival in

either sex. Instead, females that moved longer distances for their later pairings had lower LRS. Though we

did not find a similar pattern in males, this may be due to lower variation in their distance moved for later

due to their high breeding site fidelity. We note that the relationship between distance moved for first and

later pairings and lower survival and LRS may be a result of lower-quality individuals dispersing farther

and losing breeding territories rather than a cost of dispersal itself (Fuirst et al., 2023; Hui et al., 2012).

Additionally, since helpers contribute to nest success to some extent in Florida Scrub-Jays (Woolfenden &

Fitzpatrick, 1984; Mumme, 1992), the observed decline in LRS could also be a result of variation in helper

number with distance moved. Future studies should examine the intrinsic and extrinsic factors influencing

variation in reproductive success with distance moved in more detail.

The lifetime fitness costs of engaging in inbreeding were evident only in males, who had fewer

offspring that survived to adulthood when they mated with closely related females for higher proportions

of their breeding lifetime. Overall, while effects differed by sex, our results suggest that in Florida Scrub-

Jays the cost of inbreeding does not outweigh the fitness costs experienced with longer distances moved at

all breeding stages, resulting in a degree of inbreeding tolerance in accordance with theoretical expectations

(Kokko & Ots, 2006). Our results are similar to those found in a range of other animal taxa, including

insects (Jennions et al., 2004; Collet et al., 2020); birds (Gibbs & Grant, 1989; Pärt, 1996; Daniels &

Walters, 2000b; Foerster et al., 2006; Szulkin et al., 2009), and mammals (Holand et al., 2006; Geffen et

al., 2011), where higher levels of inbreeding than expected under an unconstrained random mating scenario

are tolerated despite some level of inbreeding depression. Interestingly, our results show that LRS only

approached zero for the few males that bred with closely related females for the majority of their lifetimes.

A similar pattern of sharply reduced hatch failure in closely related pairs was found in a previous study

(Chen et al., 2016). Combined, these results suggest that inbreeding costs may be sharply elevated for close

kin, but inbreeding with more distant relatives is not costly enough to outweigh the benefits of limited

dispersal in Florida Scrub-Jays. We note that our analysis likely underestimates the cost of moving farther

distances, since it does not include individuals who die in transit or otherwise fail to establish as breeders

as well as dispersers out of the study tract (Bonte et al., 2012).

In summary, we use data from a long-term, continuously monitored population with detailed

individual life history data to disentangle passive and active mechanisms of inbreeding avoidance and

quantify their fitness outcomes across the lifetime of a long-lived, cooperatively breeding species. Our

results corroborate predictions from theoretical models, demonstrating how sex-biased dispersal can

mitigate inbreeding risk and how, at the same time, in social species with fitness benefits of limited

dispersal, such passive mechanisms of inbreeding avoidance do not ensure complete mitigation of

inbreeding risk. As in other social species, we find some evidence of active inbreeding avoidance through

avoidance of first-order, likely familiar, kin. However, we also find empirical evidence to support

predictions from theoretical studies that inbreeding tolerance can persist when the fitness costs of

inbreeding do not outweigh the costs of inbreeding avoidance. Interestingly, we find that the degree of

inbreeding avoidance differs across life stages. Our study underscores the importance of examining the

costs and benefits underlying mate choice at all life stages to better disentangle the mechanisms underlying

inbreeding in natural populations.

References
Aguillon, S. M., Fitzpatrick, J. W., Bowman, R., Schoech, S. J., Clark, A. G., Coop, G., & Chen, N. (2017).

Deconstructing isolation-by-distance: The genomic consequences of limited dispersal. *PLOS*

Genetics, *13*(8), e1006911. https://doi.org/10.1371/journal.pgen.1006911

Barve, S., Riehl, C., Walters, E. L., Haydock, J., Dugdale, H. L., & Koenig, W. D. (2021). Lifetime

reproductive benefits of cooperative polygamy vary for males and females in the acorn woodpecker

(*Melanerpes formicivorus*). *Proceedings of the Royal Society B: Biological Sciences*, *288*(1957),

20210579. https://doi.org/10.1098/rspb.2021.0579

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4.

Journal of Statistical Software, *67*, 1–48. https://doi.org/10.18637/jss.v067.i01

Beck, K. B., Loretto, M.-C., & Bugnyar, T. (2020). Effects of site fidelity, group size and age on food-

caching behaviour of common ravens, *Corvus corax*. *Animal Behaviour*, *164*, 51–64.

https://doi.org/10.1016/j.anbehav.2020.03.015

Bengtsson, B. O. (1978). Avoiding inbreeding: At what cost? *Journal of Theoretical Biology*, *73*(3), 439–

444. https://doi.org/10.1016/0022-5193(78)90151-0

Billing, A. M., Lee, A. M., Skjelseth, S., Borg, Å. A., Hale, M. C., Slate, J., Pärn, H., Ringsby, T. H.,

Sæther, B.-E., & Jensen, H. (2012). Evidence of inbreeding depression but not inbreeding

avoidance in a natural house sparrow population. *Molecular Ecology*, *21*(6), 1487–1499.

https://doi.org/10.1111/j.1365-294X.2012.05490.x

Blyton, M. D. J., Banks, S. C., & Peakall, R. (2015). The effect of sex-biased dispersal on opposite-sexed

spatial genetic structure and inbreeding risk. *Molecular Ecology*, *24*(8), 1681–1695.

https://doi.org/10.1111/mec.13149

Bonte, D., Van Dyck, H., Bullock, J. M., Coulon, A., Delgado, M., Gibbs, M., Lehouck, V., Matthysen, E.,

Mustin, K., Saastamoinen, M., Schtickzelle, N., Stevens, V. M., Vandewoestijne, S., Baguette, M.,

Barton, K., Benton, T. G., Chaput-Bardy, A., Clobert, J., Dytham, C., … Travis, J. M. J. (2012).

Costs of dispersal. *Biological Reviews*, *87*(2), 290–312. https://doi.org/10.1111/j.1469-185X.2011.00201.x

Brommer, J. E. (2000). The evolution of fitness in life-history theory. *Biological Reviews*, *75*(3), 377–404. https://doi.org/10.1017/S000632310000551X

Brommer, J. E., Merilä, J., & Kokko, H. (2002). Reproductive timing and individual fitness. *Ecology Letters*, *5*(6), 802–810. https://doi.org/10.1046/j.1461-0248.2002.00369.x

Byrne, M. E., Cohen, B. S., Collier, B. A., & Chamberlain, M. J. (2022). Nest site fidelity and nesting success of female wild turkeys. *Wildlife Society Bulletin*, *46*(2), e1279. https://doi.org/10.1002/wsb.1279

Ceresa, F., Brambilla, M., Kvist, L., Vitulano, S., Pes, M., Tomasi, L., Pedrini, P., Bettega, C., Anderle, M., Hilpold, A., & Kranebitter, P. (2024). Restricted dispersal and inbreeding in a high-elevation bird across the 'sky islands' of the European Alps. *Journal of Biogeography*, *51*(5), 853–868. https://doi.org/10.1111/jbi.14787

Chen, N., Cosgrove, E. J., Bowman, R., Fitzpatrick, J. W., & Clark, A. G. (2016). Genomic consequences of population decline in the endangered Florida scrub-jay. *Current Biology*, *26*(21), 2974–2979. https://doi.org/10.1016/j.cub.2016.08.062

Clarke, A. L., Sæther, B.-E., & Røskaft, E. (1997). Sex Biases in Avian Dispersal: A Reappraisal. *Oikos*, *79*(3), 429–438. https://doi.org/10.2307/3546885

Clutton-Brock, T. H., & Lukas, D. (2012). The evolution of social philopatry and dispersal in female mammals. *Molecular Ecology*, *21*(3), 472–492. https://doi.org/10.1111/j.1365-294X.2011.05232.x

Cockburn, A., Osmond, H. L., Mulder, R. A., Green, D. J., & Double, M. C. (2003). Divorce, dispersal and incest avoidance in the cooperatively breeding superb fairy-wren *Malurus cyaneus* . *Journal of Animal Ecology*, *72*(2), 189–202. https://doi.org/10.1046/j.1365-2656.2003.00694.x

Collet, M., Amat, I., Sauzet, S., Auguste, A., Fauvergue, X., Mouton, L., & Desouhant, E. (2020). Insects and incest: Sib-mating tolerance in natural populations of a parasitoid wasp. *Molecular Ecology*, *29*(3), 596–609. https://doi.org/10.1111/mec.15340

Coltman, D. W., Pilkington, J. G., Smith, J. A., & Pemberton, josephine M. (1999). Parasite-mediated

selection against inbred soay sheep in a free-living island populaton. *Evolution*, *53*(4), 1259–1267.

https://doi.org/10.1111/j.1558-5646.1999.tb04538.x

Costantini, F., Fauvelot, C., & Abbiati, M. (2007). Fine-scale genetic structuring in Corallium rubrum:

Evidence of inbreeding and limited effective larval dispersal. *Marine Ecology Progress Series*, *340*,

109–119. https://doi.org/10.3354/meps340109

Daniels, S. J., & Walters, J. R. (2000a). Between-year breeding dispersal in red-cockaded woodpeckers:

Multiple causes and estimated cost. *Ecology*, *81*(9), 2473–2484. https://doi.org/10.1890/0012-

9658(2000)081%255B2473:BYBDIR%255D2.0.CO;2

Daniels, S. J., & Walters, J. R. (2000b). Inbreeding depression and its effects on natal dispersal in red-

cockaded woodpeckers. *The Condor*, *102*(3), 482–491. https://doi.org/10.1093/condor/102.3.482

Davison, A. C., & Hinkley, D. V. (1997). *Bootstrap Methods and Their Application*. Cambridge University

Press.

de Boer, R. A., Vega-Trejo, R., Kotrschal, A., & Fitzpatrick, J. L. (2021). Meta-analytic evidence that

animals rarely avoid inbreeding. *Nature Ecology & Evolution*, *5*(7), 949–964.

https://doi.org/10.1038/s41559-021-01453-9

De Ro, A., Vanden Broeck, A., Verschaeve, L., Jacobs, I., T'Jollyn, F., Van Dyck, H., & Maes, D. (2021).

Occasional long-distance dispersal may not prevent inbreeding in a threatened butterfly. *BMC*

Ecology and Evolution, *21*(1), 224. https://doi.org/10.1186/s12862-021-01953-z

Fitzpatrick, J. W., & Bowman, R. (2016). Florida scrub-jays: Oversized territories and group defense in a

fire-maintained habitat. In W. D. Koenig & J. L. Dickinson (Eds.), *Cooperative Breeding in*

Vertebrates (1st ed., pp. 77–96). Cambridge University Press.

https://doi.org/10.1017/CBO9781107338357.006

Fitzpatrick, J. W., Woolfenden, G. E., & Bowman, R. (1999). Dispersal distance and its demographic

consequences in the Florida Scrub-jay. *Proceedings of the International Ornithological Congress*.

Foerster, K., Valcu, M., Johnsen, A., & Kempenaers, B. (2006). A spatial genetic structure and effects of relatedness on mate choice in a wild bird population. *Molecular Ecology*, *15*(14), 4555–4567. https://doi.org/10.1111/j.1365-294X.2006.03091.x

Fox, C. W., Scheibly, K. L., & Reed, D. H. (2008). Experimental evolution of the genetic load and its implications for the genetic basis of inbreeding depression. *Evolution*, *62*(9), 2236–2249. https://doi.org/10.1111/j.1558-5646.2008.00441.x

Fox, J., & Monette, G. (1992). Generalized Collinearity Diagnostics. *Journal of the American Statistical Association*, *87*(417), 178–183.

Frommen, J. G., & Bakker, T. C. M. (2006). Inbreeding avoidance through non-random mating in sticklebacks. *Biology Letters*, *2*(2), 232–235. https://doi.org/10.1098/rsbl.2005.0432

Fuirst, M., Strickland, D., Freeman, N. E., Sutton, A. O., & Ryan Norris, D. (2023). Early-life sibling conflict in Canada jays has lifetime fitness consequences. *Proceedings of the Royal Society B: Biological Sciences*, *290*(1997), 20221863. https://doi.org/10.1098/rspb.2022.1863

Galezo, A. A., Nolas, M. A., Fogel, A. S., Mututua, R. S., Warutere, J. K., Siodi, I. L., Altmann, J., Archie, E. A., Tung, J., & Alberts, S. C. (2022). Mechanisms of inbreeding avoidance in a wild primate. *Current Biology*, *32*(7), 1607-1615.e4. https://doi.org/10.1016/j.cub.2022.01.082

Geffen, E., Kam, M., Hefner, R., Hersteinsson, P., Angerbjörn, A., Dalèn, L., Fuglei, E., Norèn, K., Adams, J. R., Vucetich, J., Meier, T. J., Mech, L. D., VonHOLDT, B. M., Stahler, D. R., & Wayne, R. K. (2011). Kin encounter rate and inbreeding avoidance in canids. *Molecular Ecology*, *20*(24), 5348–5358. https://doi.org/10.1111/j.1365-294X.2011.05358.x

Gibbs, H. L., & Grant, P. R. (1989). Inbreeding in Darwin's medium ground finches (Geospiza fortis). *Evolution*, *43*(6), 1273–1284. https://doi.org/10.1111/j.1558-5646.1989.tb02574.x

Green, J. P., & Hatchwell, B. J. (2018). Inclusive fitness consequences of dispersal decisions in a cooperatively breeding bird, the long-tailed tit (*Aegithalos caudatus*). *Proceedings of the National Academy of Sciences*, *115*(47), 12011–12016. https://doi.org/10.1073/pnas.1815873115

Greenwood, P. J., & Harvey, P. H. (1982). The natal and breeding dispersal of birds. *Annual Review of*

Ecology and Systematics, *13*(1), 1–21. https://doi.org/10.1146/annurev.es.13.110182.000245

Griesser, M., Nystrand, M., & Ekman, J. (2006). Reduced mortality selects for family cohesion in a social

species. *Proceedings of the Royal Society B: Biological Sciences*, *273*, 1881–1886.

https://doi.org/10.1098/rspb.2006.3527

Hamilton, W. D. (1964). The genetical evolution of social behaviour. I. *Journal of Theoretical Biology*,

7(1), 1–16. https://doi.org/10.1016/0022-5193(64)90038-4

Harvey, P. H., Greenwood, P. J., & Perrins, C. M. (1979). Breeding Area Fidelity of Great Tits (Parus

major). *Journal of Animal Ecology*, *48*(1), 305–313. https://doi.org/10.2307/4115

Hatchwell, B. J., Russell, A. F., Ross, D. J., & Fowlie, M. K. (2000). Divorce in cooperatively breeding

long–tailed tits: A consequence of inbreeding avoidance? *Proceedings of the Royal Society B:*

Biological Sciences, *267*(1445), 813–819. https://doi.org/10.1098/rspb.2000.1076

Haydock, J., Koenig, W. D., & Stanback, M. T. (2001). Shared parentage and incest avoidance in the

cooperatively   breeding   acorn   woodpecker.   *Molecular   Ecology*,   *10*(6),   1515–1525.

https://doi.org/10.1046/j.1365-294X.2001.01286.x

Hidalgo Aranzamendi, N., Hall, M. L., Kingma, S. A., Sunnucks, P., & Peters, A. (2016). Incest avoidance,

extrapair paternity, and territory quality drive divorce in a year-round territorial bird. *Behavioral*

Ecology, *27*(6), 1808–1819. https://doi.org/10.1093/beheco/arw101

Holand, Ø., Askim, K. R., Røed, K. H., Weladji, R. B., Gjøstein, H., & Nieminen, M. (2006). No evidence

of inbreeding avoidance in a polygynous ungulate: The reindeer (*Rangifer tarandus*). *Biology*

Letters, *3*(1), 36–39. https://doi.org/10.1098/rsbl.2006.0575

Hui, C., Roura-Pascual, N., Brotons, L., Robinson, R. A., & Evans, K. L. (2012). Flexible dispersal

strategies in native and non-native ranges: Environmental quality and the 'good–stay, bad–

disperse' rule. *Ecography*, *35*(11), 1024–1032. https://doi.org/10.1111/j.1600-0587.2012.07697.x

Jennions, M. D., Hunt, J., Graham, R., & Brooks, R. (2004). No evidence for inbreeding avoidance through postcopulatory mechanisms in the black field cricket, *Teleogryllus commodus*. *Evolution*, *58*(11), 2472–2477. https://doi.org/10.1111/j.0014-3820.2004.tb00876.x

Jiménez, J. A., Hughes, K. A., Alaks, G., Graham, L., & Lacy, R. C. (1994). An experimental study of inbreeding depression in a natural habitat. *Science*, *266*(5183), 271–273. https://doi.org/10.1126/science.7939661

Keller, L. F., Arcese, P., Smith, J. N. M., Hochachka, W. M., & Stearns, S. C. (1994). Selection against inbred song sparrows during a natural population bottleneck. *Nature*, *372*(6504), Article 6504. https://doi.org/10.1038/372356a0

Kingma, S. A., Komdeur, J., Burke, T., & Richardson, D. S. (2017). Differential dispersal costs and sex-biased dispersal distance in a cooperatively breeding bird. *Behavioral Ecology*, *28*(4), 1113–1121. https://doi.org/10.1093/beheco/arx075

Kokko, H., & Ekman, J. (2002). Delayed dispersal as a route to breeding: Territorial inheritance, safe havens, and ecological constraints. *The American Naturalist*, *160*(4), 468–484. https://doi.org/10.1086/342074

Kokko, H., & Ots, I. (2006). When not to avoid inbreeding. *Evolution*, *60*(3), 467–475. https://doi.org/10.1111/j.0014-3820.2006.tb01128.x

Lawson Handley, L. J., & Perrin, N. (2007). Advances in our understanding of mammalian sex-biased dispersal. *Molecular Ecology*, *16*(8), 1559–1578. https://doi.org/10.1111/j.1365-294X.2006.03152.x

Leedale, A. E., Simeoni, M., Sharp, S. P., Green, J. P., Slate, J., Lachlan, R. F., Robinson, E. J. H., & Hatchwell, B. J. (2020). Cost, risk, and avoidance of inbreeding in a cooperatively breeding bird. *Proceedings of the National Academy of Sciences*, *117*(27), 15724–15730. https://doi.org/10.1073/pnas.1918726117

Leedale, A. E., Vullioud, P., Seager, D., Zöttl, M., Glauser, G., & Clutton-Brock, T. (2024). Kin recognition
for incest avoidance in Damaraland mole-rats, Fukomys damarensis. *Proceedings of the Royal*
Society B: Biological Sciences, *291*(2033), 20241138. https://doi.org/10.1098/rspb.2024.1138

Li, X.-Y., & Kokko, H. (2019). Sex-biased dispersal: A review of the theory: Sex-biased dispersal: a review
of the theory. *Biological Reviews*, *94*(2), 721–736. https://doi.org/10.1111/brv.12475

Lloyd, M. W., Tumas, H. R., & Neel, M. C. (2018). Limited pollen dispersal, small genetic neighborhoods,
and biparental inbreeding in Vallisneria americana. *American Journal of Botany*, *105*(2), 227–240.
https://doi.org/10.1002/ajb2.1031

Lubin, Y., Birkhofer, K., Berger-Tal, R., & Bilde, T. (2009). Limited male dispersal in a social spider with
extreme inbreeding. *Biological Journal of the Linnean Society*, *97*(2), 227–234.
https://doi.org/10.1111/j.1095-8312.2009.01190.x

Maag, N., Cozzi, G., Bateman, A., Heistermann, M., Ganswindt, A., Manser, M., Clutton-Brock, T., &
Ozgul, A. (2019). Cost of dispersal in a social mammal: Body mass loss and increased stress.
Proceedings of the Royal Society B: Biological Sciences, *286*(1896), 20190033.
https://doi.org/10.1098/rspb.2019.0033

Malécot, G. (1948). *Les mathématiques de l'hérédité*. Masson.

Marzluff, J. M., Woolfenden, G. E., Fitzpatrick, John W., & Balda, R. P. (1996). Breeding partnerships of
two New World jays. In J. M. Black & M. Hulme (Eds.), *Partnerships in Birds: The Study of*
Monogamy (p. 0). Oxford University Press. https://doi.org/10.1093/oso/9780198548614.003.0007

McAdam, A. G., Boutin, S., Sykes, A. K., & Humphries, M. M. (2007). Life histories of female red squirrels
and their contributions to population growth and lifetime fitness. *Écoscience*, *14*(3), 362–369.
https://doi.org/10.2980/1195-6860(2007)14%255B362:LHOFRS%255D2.0.CO;2

McGowan, K. J., & Woolfenden, G. E. (1990). Contributions to fledgling feeding in the Florida Scrub-Jay.
Journal of Animal Ecology, *59*(2), 691–707. https://doi.org/10.2307/4889

Morrison, R. E., Ndayishimiye, E., Stoinski, T. S., & Eckardt, W. (2023). Multiple mechanisms for

inbreeding avoidance used simultaneously in a wild ape. *Proceedings of the Royal Society B:*

Biological Sciences, *290*(2009), 20231808. https://doi.org/10.1098/rspb.2023.1808

Mumme, R. L. (1992). Do helpers increase reproductive success? *Behavioral Ecology and Sociobiology*,

31(5), 319–328. https://doi.org/10.1007/BF00177772

Mumme, R. L., Bowman, R., Pruett, M. S., & Fitzpatrick, J. W. (2015). Natal territory size, group size, and

body mass affect lifetime fitness in the cooperatively breeding Florida Scrub-Jay. *The Auk*, *132*(3),

634–646. https://doi.org/10.1642/AUK-14-258.1

Nelson-Flower, M. J., Hockey, P. A. R., O'Ryan, C., & Ridley, A. R. (2012). Inbreeding avoidance

mechanisms: Dispersal dynamics in cooperatively breeding southern pied babblers. *Journal of*

Animal Ecology, 876–883.

Nichols, H. J. (2017). The causes and consequences of inbreeding avoidance and tolerance in cooperatively

breeding vertebrates. *Journal of Zoology*, *303*(1), 1–14. https://doi.org/10.1111/jzo.12466

Paradis, E., Baillie, S. R., Sutherland, W. J., & Gregory, R. D. (1998). Patterns of natal and breeding

dispersal in birds. *Journal of Animal Ecology*, *67*(4), 518–536. https://doi.org/10.1046/j.1365-

2656.1998.00215.x

Pärt, T. (1994). Male philopatry confers a mating advantage in the migratory collared flycatcher, *Ficedula*

albicollis. *Animal Behaviour*, *48*(2), 401–409. https://doi.org/10.1006/anbe.1994.1254

Pärt, T. (1996). Problems with testing inbreeding avoidance: The case of the collared flycatcher. *Evolution*,

50(4), 1625–1630. https://doi.org/10.1111/j.1558-5646.1996.tb03934.x

Pebesma, E. (2018). Simple features for R: Standardized support for spatial vector data. *The R Journal*,

10(1), 439. https://doi.org/10.32614/RJ-2018-009

Perrin, N., & Goudet, J. (2001). Inbreeding, kinship, and the evolution of natal dispersal. In J. Clobert, E.

Danchin, A. A. Dhondt, & J. D. Nichols (Eds.), *Dispersal* (pp. 123–142). Oxford University Press.

https://iris.unil.ch/handle/iris/62054

Pusey, A., & Wolf, M. (1996). Inbreeding avoidance in animals. *Trends in Ecology & Evolution*, *11*(5),

201–206. https://doi.org/10.1016/0169-5347(96)10028-8

R Core Team. (2022). *R: A Language and Environment for Statistical Computing.* [Computer software]. R

Foundation for Statistical Computing.

Riehl, C., & Stern, C. A. (2015). How cooperatively breeding birds identify relatives and avoid incest: New

insights into dispersal and kin recognition. *BioEssays*, *37*(12), 1303–1308.

https://doi.org/10.1002/bies.201500120

Sæther, B.-E., & Bakke, Ø. (2000). Avian life history variation and contribution of demographic traits to

the population growth rate. *Ecology*, *81*(3), 642–653. https://doi.org/10.1890/0012-

9658(2000)081%255B0642:ALHVAC%255D2.0.CO;2

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients. *Methods in*

Ecology and Evolution, *1*(2), 103–113. https://doi.org/10.1111/j.2041-210X.2010.00012.x

Schmidt, K. A. (2014). Site fidelity curbs sequential search and territory choice: A game theoretical

approach. *Functional Ecology*, *28*(6), 1494–1503. https://doi.org/10.1111/1365-2435.12291

Schoech, S. J., Mumme, R. L., & Wingfield, J. C. (1996). Delayed breeding in the cooperatively breeding

Florida scrub-jay (Aphelocoma coerulescens): Inhibition or the absence of stimulation? *Behavioral*

Ecology and Sociobiology, *39*(2), 77–90. https://doi.org/10.1007/s002650050269

Shah, S. S., & Rubenstein, D. R. (2022). Prenatal environmental conditions underlie alternative

reproductive tactics that drive the formation of a mixed-kin cooperative society. *Science Advances*,

8(8), eabk2220. https://doi.org/10.1126/sciadv.abk2220

Sinnwell, J. P., Therneau, T. M., & Schaid, D. J. (2014). The kinship2 R Package for Pedigree Data. *Human*

Heredity, *78*(2), 91–93. https://doi.org/10.1159/000363105

Stopher, K. V., Nussey, D. H., Clutton-brock, T. H., Guinness, F., Morris, A., & Pemberton, J. M. (2012).

Re-mating across years and intralineage polygyny are associated with greater than expected levels

of inbreeding in wild red deer. *Journal of Evolutionary Biology*, *25*(12), 2457–2469.

https://doi.org/10.1111/j.1420-9101.2012.02626.x

Suh, Y. H., Bowman, R., & Fitzpatrick, J. W. (2022). Staging to join non-kin groups in a classical

cooperative breeder, the Florida scrub-jay. *Journal of Animal Ecology*, *91*(5), 970–982.

https://doi.org/10.1111/1365-2656.13669

Suh, Y. H., Pesendorfer, M. B., Tringali, A., Bowman, R., & Fitzpatrick, J. W. (2020). Investigating social

and environmental predictors of natal dispersal in a cooperative breeding bird. *Behavioral Ecology*,

31(3), 692–701. https://doi.org/10.1093/beheco/araa007

Summers, J., Cosgrove, E. J., Bowman, R., Fitzpatrick, J. W., & Chen, N. (2024). Density dependence

maintains long-term stability despite increased isolation and inbreeding in the Florida Scrub-Jay.

Ecology Letters, *27*(12), e14483. https://doi.org/10.1111/ele.14483

Szulkin, M., Garant, D., Mccleery, R. H., & Sheldon, B. C. (2007). Inbreeding depression along a life-

history continuum in the great tit. *Journal of Evolutionary Biology*, *20*(4), 1531–1543.

https://doi.org/10.1111/j.1420-9101.2007.01325.x

Szulkin, M., Stopher, K. V., Pemberton, J. M., & Reid, J. M. (2013). Inbreeding avoidance, tolerance, or

preference in animals? *Trends in Ecology & Evolution*, *28*(4), 205–211.

https://doi.org/10.1016/j.tree.2012.10.016

Szulkin, M., Zelazowski, P., Nicholson, G., & Sheldon, B. C. (2009). Inbreeding avoidance under different

null models of random mating in the great tit. *Journal of Animal Ecology*, *78*(4), 778–788.

https://www.jstor.org/stable/27696427

Therneau, T. (2022). *A Package for Survival Analysis in R* (Version R package version 3.4-0) [Computer

software]. https://CRAN.R-project.org/package=survival

Thünken, T., Bakker, T. C. M., Baldauf, S. A., & Kullmann, H. (2007). Active inbreeding in a cichlid fish

and its adaptive significance. *Current Biology*, *17*(3), 225–229.

https://doi.org/10.1016/j.cub.2006.11.053

Townsend, A. K., Bowman, R., Fitzpatrick, J. W., Dent, M., & Lovette, I. J. (2011). Genetic monogamy

across variable demographic landscapes in cooperatively breeding Florida scrub-jays. *Behavioral*

Ecology, *22*(3), 464–470. https://doi.org/10.1093/beheco/arq227

Van de Casteele, T., Galbusera, P., Schenck, T., & Matthysen, E. (2003). Seasonal and lifetime reproductive
consequences of inbreeding in the great tit Parus major. *Behavioral Ecology*, *14*(2), 165–174.
https://doi.org/10.1093/beheco/14.2.165

van Dijk, R. E., Covas, R., Doutrelant, C., Spottiswoode, C. N., & Hatchwell, B. J. (2015). Fine-scale
genetic structure reflects sex-specific dispersal strategies in a population of sociable weavers
(Philetairus socius). *Molecular Ecology*, *24*(16), 4296–4311. https://doi.org/10.1111/mec.13308

Venables, W., & Ripley, B. (2002). *Modern Applied Statistics with S* (4th ed.). Springer.
https://www.stats.ox.ac.uk/pub/MASS4/

Wheelwright, N. T., & Mauck, R. A. (1998). Philopatry, natal dispersal, and inbreeding avoidance in an
island population of savannah sparrows. *Ecology*, *79*(3), 755–767. https://doi.org/10.1890/0012-
9658(1998)079%255B0755:PNDAIA%255D2.0.CO;2

Woolfenden, G. E., & Fitzpatrick, J. W. (1984). *The Florida Scrub Jay: Demography of a Cooperative-*
breeding Bird. Princeton University Press.

Zhang, Z., Reinikainen, J., Adeleke, K. A., Pieterse, M. E., & Groothuis-Oudshoorn, C. G. M. (2018).
Time-varying covariates and coefficients in Cox regression models. *Annals of Translational*
Medicine, *6*(7), 121. https://doi.org/10.21037/atm.2018.02.12

  Figures and Tables

Figure 1. Relatedness to mate and proportion of first-order kin mates observed and expected under three

different models of random mating in Florida Scrub-Jays at two life stages (first and later pairings) and for

both sexes (male and female). The observed population means across all 32 years of our study period are

indicated by black vertical lines. Histograms show the distributions of expected means under 1000

simulations of random mating in a scenario with unconstrained dispersal (purple), limited and sex-biased

dispersal (red), and limited dispersal without sex-bias (yellow) (filled bars = distribution significantly

different; unfilled bars = distribution not significantly different from the observed mean). Means of

distributions are indicated by arrows.

Figure 2. Predicted model fits investigating the effects (solid line = significant, dashed line = not significant; shaded area = 95% confidence interval) of (A) Distance moved for first pairing, and (B) Weighted kinship coefficients to mates on lifetime reproductive success (LRS) for Florida Scrub Jay male (blue, triangles) and female (orange, circles) breeders with lifetime data between 1990 and 2021 (*N*: Males = 197, Females = 120), and (C) Total distance moved for later pairings, and (D) Weighted kinship coefficients to mates for the subset of breeders that switched mates at least once in their lifetime (*N*: Male = 91, Female = 50). Females had lower LRS with greater total distance moved for later pairings, while males showed similarly lower LRS with higher kinship to mates over their lifetime. Points are jittered both horizontally and vertically for better visual discrimination.

806   **Table 1.** Results of generalized linear mixed models estimating the effect of distance moved for first

807   pairings, later pairings, relatedness to mate, and breeding lifespan on lifetime reproductive success (*i.e.*,

808   number of offspring surviving to adulthood) of Florida Scrub-Jays breeders. Significant effects are in bold.

Dataset	Sex	Fixed Effect	Estimate	Std Err	z-value	p-value
All breeders with lifetime data between 1990-2021	Male (*N*=197)	**Intercept**	0.47	0.08	5.86	**<0.001**
		Distance moved for first pairing	-0.09	0.09	-0.94	0.35
		Weighted kinship coefficient to mates	-0.34	0.12	-2.83	**0.005**
		Breeding Lifespan	0.93	0.07	13.75	**<0.001**
	Female (*N*=120)	**Intercept**	0.70	0.10	7.38	**<0.001**
		Distance moved for first pairing	-0.12	0.08	-1.43	0.15
		Weighted kinship coefficient to mates	0.03	0.08	0.31	0.76
		Breeding Lifespan	0.86	0.10	8.86	**<0.001**
Breeders with lifetime data between 1990-2021 that switched mates at least once	Male (*N*=91)	**Intercept**	1.25	0.08	14.90	**<0.001**
		Total distance moved for later pairings	-0.06	0.09	-0.64	0.52
		Weighted kinship coefficient to mates	-0.28	0.09	-3.05	**0.002**
		Breeding Lifespan	0.60	0.07	8.63	**<0.001**
	Female (*N*=50)	**Intercept**	1.21	0.10	12.15	**<0.001**
		Total distance moved for later pairings	-0.27	0.10	-2.62	**0.01**
		Weighted kinship coefficient to mates	-0.01	0.14	-0.04	0.97
		Breeding Lifespan	0.83	0.10	7.93	**<0.001**

810