1 **Lifetime fitness benefits of breeding site fidelity and low costs of inbreeding permit inbreeding**
2 **tolerance in an avian cooperative breeder**

3 Shailee S. Shah[1,2]*, Jennifer Diamond[3], Sahas Barve[4], Elissa J. Cosgrove[5], Reed Bowman[4], John W.
4 Fitzpatrick[2,6], Nancy Chen[1]*
5
[1] Department of Biology, University of Rochester, Rochester, NY 14627, USA
[2] Cornell Lab of Ornithology, Cornell University, Ithaca, NY 14850, USA
[3] Department of Evolution and Ecology, University of California, Davis, Davis, CA 95616, USA
[4] Avian Ecology Program, Archbold Biological Station, Venus, FL 33960, USA
[5] Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14850, USA
[6] Department of Ecology & Evolutionary Biology, Cornell University, Ithaca, NY 14850, USA
12
* Correspondence: shailee.shah@cornell.edu, nancy.chen@rochester.edu

Acknowledgments: We thank Archbold Biological Station and its staff for continued support and
management of the study area; all interns, students, and postdocs who have worked on collecting and
managing the long-term data; the Cornell Statistical Consulting Unit for statistical support; V. Sclater for
GIS support; and F. Romero, J. Summers, D. Seidman, and other members of the Chen Lab at the University
of Rochester for invaluable feedback on previous versions of this manuscript.
20
21 **Funding:** This work was supported by National Science Foundation (NSF) Grants DEB0855879 and
22 DEB1257628 and by the Cornell Lab of Ornithology Athena Fund. S.S.S was supported by NIH grant
23 1R35GM133412 to N.C. and NSF Postdoctoral Research Fellowship in Biology 2305705.

Abstract

Inbreeding, or breeding with close relatives, often decreases individual fitness, but mate choice in many species can increase inbreeding risk. Inbreeding is more likely in species with limited dispersal, such as cooperative breeders where non-parental individuals—often offspring from previous broods—provide parental care and frequently breed close to home. We leverage 32 years of data from a population of Florida Scrub-Jays (*Aphelocoma coerulescens*), an avian cooperative breeder, to investigate whether mate choice, and its lifetime fitness outcomes, affects inbreeding tolerance. We find that both sexes show a stronger preference for geographically closer rather than less related mates. While female-biased dispersal mitigates inbreeding risk when pairing for the first time, limited movement away from the breeding territory by both sexes for later pairings (following mate death or divorce) results in higher relatedness between mates than expected under random mating. Males paired with close relatives and females that move longer distances experience equivalently reduced lifetime reproductive success. Fitness benefits of breeding site fidelity and relatively low fitness costs of inbreeding thus permit inbreeding tolerance in Florida Scrub-Jays. Understanding how and why mate choice contributes to inbreeding is crucial for informed conservation action for endangered species like the Florida Scrub-Jay.

Keywords: Mate choice, inbreeding tolerance, site fidelity, cooperative breeding, Florida Scrub-Jay

Main Text

Introduction

Matings among relatives, or inbreeding, can have severe negative fitness consequences such as increased susceptibility to parasites [1], lower birth rates and offspring survival [2,3], and increased risk of mortality during severe weather events [4]. In wild populations, inbreeding can be the result of two non-exclusive factors. First, in populations that are small and isolated, mating with relatives might be inevitable even when mating is random [5]. Second, mate choice may be driven by factors that increase fitness despite the risk of inbreeding, such as phenotype matching or limited dispersal [6]. In the latter case, while inbreeding tolerance may persist without severe population-level impacts in large, well-connected populations, it may exacerbate inbreeding depression in isolated population fragments [7]. Understanding how mate choice may result in increased instances of inbreeding in species of conservation concern is thus vital for informed conservation action.

Moving away from their natal area—*i.e.*, dispersal—is the primary mechanism by which animals avoid mating with kin [8,9]. However, dispersal may be limited in some species, such that when individuals become breeders, they often choose mates that are geographically close [8,10]. While limited dispersal can lead to fitness benefits such as better access to resources [11], higher survival [12], and nepotism [13], it also can increase the likelihood of mating with a close relative [14–16]. Similarly, once individuals establish a breeding territory, any subsequent dispersal also may be geographically limited even when pairing with a new mate [8]. Such site fidelity may have fitness benefits due to familiarity and reduced competitive interactions [17–19], but also may increase inbreeding risk in spatially structured populations where limited dispersal results in the formation of kin neighborhoods [20]. Nevertheless, if the fitness benefits of limited dispersal outweigh the costs of inbreeding at the individual level, inbreeding may be tolerated [6,21]. Conversely, additional behaviors such as kin avoidance—where individuals can recognize and avoid mating with kin—and sex-biased dispersal—where one sex disperses farther than the other—may prevent inbreeding even in species with limited dispersal [6].

Previous studies have investigated whether species with limited dispersal exhibit inbreeding tolerance or avoidance behaviors [7,22–27], and even quantified the fitness tradeoffs between dispersal distance and kinship in mate choice [22]. However, few such studies have incorporated measures of lifetime fitness. This matters because reproductive strategies over an individual's lifetime, especially in long-lived species such as many cooperative breeders [28,29], may compensate for any fitness costs associated with inbreeding or dispersal. Thus, we must examine lifetime fitness to obtain a comprehensive understanding of the individual fitness outcomes underlying mate choice that may increase inbreeding risk. Additionally, the importance of dispersal distance in mate choice may differ based on whether an individual is breeding for the first time ("first pairing") versus an experienced breeder pairing with a new mate ("later pairing") [8]. Because most studies only examine mate choice during first pairing, mechanisms such as site fidelity that may limit dispersal or increase inbreeding risk at later life stages [8] are ignored. Here, we leverage 32 years of individual-level data on dispersal, mating, reproductive success, and survival from a continuously monitored population of an avian cooperative breeder, the Florida Scrub-Jay (*Aphelocoma coerulescens*), to investigate whether inbreeding risk is elevated due to mate choice, and the resulting fitness outcomes throughout an individual's lifetime.

Fewer than 6,000 individuals of the Federally Threatened Florida Scrub-Jay remain in numerous, widely isolated populations restricted to fire-maintained, xeric oak scrub habitats of the Florida peninsula [30]. Their preferred habitat is usually saturated with year-round territories, each occupied by a socially monogamous dominant breeding pair, accompanied often by subordinate non-breeders of both sexes [31]. Subordinates are primarily offspring from previous broods that delay dispersal and remain in their natal territory for one or more years, despite being physiologically capable of breeding at one year of age [31].

Most individuals become breeders by dispersing to replace dead breeders on existing territories, though
direct inheritance or "budding" of natal territories sometimes occurs [31]. Females disperse farther than
males, though neither sex typically disperses very far [32,33]. Once they become breeders, Florida Scrub-
Jays remain breeders until their death, pairing with a new mate if their current mate dies [31], or in rare
cases when the mate is still alive ("divorce") [34]. We focus on a population of Florida Scrub-Jays that has
been intensively studied at Archbold Biological Station in central Florida (27.10°N, 81.21°W) since 1969
[31]. Immigration into the Archbold population from neighboring populations has declined since 1995,
leading to a corresponding increase in inbreeding and associated fitness costs [3].

To investigate how mate choice affects inbreeding risk, we first examined how the relatedness of Florida
Scrub-Jays to potential mates varied with distance from both their natal and any subsequent breeding
territories. Given fine-scale genetic isolation-by-distance in this population [33], we predicted that potential
mates closer to both their natal and breeding territories would be relatively closely related for both sexes,
and more so for females since males are the more philopatric sex [32]. Second, we examined the extent to
which dispersal distance, relatedness, or both, correspond with mate choice. Previous work in this study
population found individuals that dispersed shorter distances had higher fitness [35], and inbred individuals
have lower fitness [3]. Thus, we predicted that pairings with shorter dispersal distances and lower
relatedness between mates would be more common, because these traits would be adaptively favored.
Third, we tested whether dispersal distance and average relatedness of mates deviated from expectations
under random mating. We predicted that average dispersal distance would be shorter than expected, and
that the relatedness of mates would be lower than that expected under a random mating scenario (i.e., owing
to inbreeding avoidance).

Next, to investigate the relative fitness costs and benefits involved in mate choice patterns, we quantified
the impact of distance moved and relatedness between mates on lifetime reproductive success and survival.
We predicted that for both sexes, longer dispersal distances for both first and later pairings would result in
lower lifetime reproductive success and survival [35]. Further, we predicted that if we found no evidence
of kin avoidance in our mate choice analysis, the fitness cost of relatedness between mates would be
equivalent to or lower than the cost of long-distance movement (suggesting inbreeding tolerance). Finally,
we examined whether Florida Scrub-Jays employ other reproductive strategies, such as divorce, to
minimize the costs of inbreeding. Since lifetime reproductive success increases with the number of annual
breeding attempts [30], Florida Scrub-Jays may favor early breeding with a close relative over not breeding
at all, but then may switch to less related mates when an opportunity arises. Thus, we predicted that pairs
with more closely related mates would be more likely to dissolve because of divorce than death of a mate.
Ultimately, by linking the effect of distance and kinship on mate choice with individual lifetime fitness
outcomes, we sought evidence for how mate choice that increases inbreeding risk can persist in a
population.

Methods

Data collection

Florida Scrub-Jays at Archbold Biological Station have been continuously monitored since 1969, with the
study tract expanded to its current size (mean ± sd = 914.04 ± 143.44 ha) by 1990 [30]. All birds are banded
with unique combinations of color bands. Throughout the breeding season (mid-March to early June), all
nests are located and monitored for fledging or failure [30]. All territories are mapped in April, with territory
boundaries assigned at locations of territorial disputes between neighboring groups [30]. In addition to nest
monitoring, the population is censused each month, resulting in detailed data on dispersal and survival of
both fledglings and adults. Pairs are socially monogamous, with very low rates of extra pair-paternity [36],
allowing accurate parentage inference from field observations alone. Furthermore, previous work verified
genetic parentage of offspring born between ~1989-1991, 1994-1995, and 1999-2013 [3]. Using these

detailed data, we have accumulated a 15-generation pedigree of known-age, known-parentage individuals
with precise measurements of lifespan and reproductive success. For this study, we used data from 1990-
2021, which included on average (± SD) 71.4 (± 14.7) territories (area mean ± sd =12.98 ± 6.94 ha) per
year. All analyses were conducted using R version 4.2.2 [37]. Fieldwork was approved by the Institutional
Animal Care and Use Committees at Cornell University (IACUC 2010-0015) and Archbold Biological
Station (AUP-006-R) under permits from the US Fish and Wildlife Service (TE824723-8, TE-117769), the
US Geological Survey (banding permits 07732, 23098), and the Florida Fish and Wildlife Conservation
Commission (LSSC-10-00205).
152
We first calculated observed dispersal distances and relatedness to mate for both male and female breeders
in all new breeding pairs formed between 1990 and 2021. New breeding pairs include pairs formed by two
first-time breeders, one first-time and one experienced breeder, as well as two experienced breeders. For
first-time breeders, we called the event "first pairing" and for experienced breeders, "later pairing". New
pairs might occupy a new territory, which may be inherited by one of the two breeders or formed as an
extension of their natal territory in a process called "budding" [30]. Alternatively, a new pair might form
on a non-natal territory already occupied by one of the two breeders, where a breeding vacancy typically
formed due to the death of a breeder, but occasionally via divorce [30]. Thus, both first and later pairings
can occur without any movement away from the currently occupied territory by at least one of the two
breeders. When the focal individual remained on its natal territory and became a breeder, formed a territory
by budding, or remained on its non-natal breeding territory, we quantified distance moved as 0 m. To obtain
accurate measures of distance moved, we excluded any pairings that involved movement that started or
ended outside of the study tract. We also excluded any movement that did not end with the individual
attaining a breeding position (*i.e.*, staging, a strategy adopted by about 28% of yearlings [38]). For staging
individuals that eventually became breeders, we considered it a first pairing event with movement from
their staging to breeding territory.
169
In accordance with previous work in this system, we used the observed number of territories crossed as the
most biologically relevant measure of distance moved [35]. The number of territories crossed was highly
correlated with distance moved in meters (Pearson's $r = 0.91$; [35]). To calculate the number of territories
crossed by a jay for a new pairing, we drew a straight-line movement trajectory from the centroid of its
previous territory to the centroid of its new breeding territory. We then used the *st_intersects* function from
the R package *sf* [39] to calculate the number of territories (*i.e.*, polygons) that each line intersected. For
cases where the function returned a distance ≥ 1 territory away, but the territory name did not change from
its natal or previous breeding territory (84 out of 1690 [5%] of movement events), we manually corrected
distances using territory maps (see *SI*). We also calculated the kinship coefficient between mates for all
new breeding pairs using the population pedigree with the *kinship* function in the R package *kinship2*
[10]. With no inbreeding, the kinship coefficient ranges from 0 to 0.5, such that 0.5 is an individual's
kinship coefficient with itself; 0.25 is the average kinship coefficient between siblings and between parent
and offspring; 0.125 is the average kinship coefficient between half-siblings, grandparents-grandchildren,
and aunt/uncle-niece/nephew pairs; and 0.0625 is the average kinship coefficient between first cousins [40].
Pedigree-based inbreeding estimates are correlated with genomic-based estimates in our study population
[3].
186
187 *Data analysis*
188
First, to examine how relatedness to potential mates varies with distance, we constructed a dataset of all
possible mate pairings for each year and each sex separately. We only included breeders that formed new
breeding pairs in a given year (*i.e.*, individuals that were both available as potential mates and successful at
finding new mates), which comprised both first-time and experienced breeders. For a given year, we paired
all available females with each available male and vice versa (*N*: Male, First pairing = 421; Male, Later
pairing = 249; Female, First pairing = 391; Female, Later pairing = 256 instances). We calculated the

potential distance that each focal individual would have to travel to realize this pairing as the distance
between the individual's territory in the previous year to the potential mate's territory in the current year.
For simplicity, we assumed that the potential mate would retain its territory. We also calculated the kinship
coefficient for all potential pairings as described above, and classified potential mates to whom the kinship
coefficient of the focal individual was ≥ 0.0625 (at least as related as first cousins) as close relatives. We
then fit generalized linear mixed models (GLMMs) with whether the potential mate was a close relative (1
= yes, 0 = no) as the dependent variable, a binomial error structure, distance to mate as a fixed effect, and
year and focal individual ID as random effects. We fit separate models for each sex and pairing type (N:
Male, First pairing = 12155; Male, Later pairing = 11837; Female, First pairing = 10938; Female, Later
pairing = 11000 possible pairings).

Second, using the same dataset of all possible breeding pairs, we investigated the effect of distance to mate
and relatedness on mate choice. We classified each potential pairing as "realized" (*i.e.*, observed that year)
or "not realized" (*i.e.*, not observed that year). We then fit four separate GLMMs, one for each category,
with whether a potential pairing was realized as the dependent variable and kinship coefficient and distance
moved as fixed effects. We included year and focal individual ID as random effects and used a binomial
error structure. We repeated this analysis twice, once with the full dataset and once with the dataset
truncated to only include potential mates within a distance of four territories (N: Male, First pairing = 5202;
Male, Later pairing = 5013; Female, First pairing = 4525; Female, Later pairing = 4686 possible pairings),
since beyond this distance the likelihood of pairing with close relatives approaches zero (Fig. 1A). We fit
GLMMs using the function *glmer* in the R package *lme4* [41].

Third, to quantify deviations of distance and average relatedness of mates from expectations under a random
mating scenario, we randomly permuted all available mates of the opposite sex among all available mates
of the focal sex for each year. We then calculated the distance and kinship coefficient to mate using the
same methods as above for each randomly -generated pair. We ran 1,000 iterations per year and calculated
the mean distance and kinship coefficient to mate for each year for each focal sex and pairing type. We then
compared these expected distributions of mean distances and kinship coefficients to observed means by
calculating empirical p-values. Specifically, we quantified the proportion of simulations that had a mean
distance to mate that was higher than the observed mean, and the proportion of simulations that had a mean
kinship coefficient to mate that was lower than the observed mean.

Fourth, to examine how distance and relatedness to mate impacts lifetime fitness of Florida Scrub-Jays, we
calculated three metrics of lifetime reproductive success for all breeders: the total number of offspring
produced over an individual's lifetime that survived to day 30, day 90, and day 300, corresponding
respectively to the ages at which Florida Scrub-Jays become physiologically capable of sustained flight,
nutritional independence, and breeding [31,42–44]. Survival to these early life stages is governed by a
varying degree of interactions between social and environmental factors, such as group and territory size,
and offspring condition [42]. For this analysis, we restricted our dataset to individuals from cohorts from
which all members born within the study tract were last observed before 2021 to eliminate bias towards
short-lived individuals [45,46]. The final dataset included individuals born between 1988-2006 and in 2009
(N: Males =196, Females =122).

We then fit six separate GLMMs (two sexes × three fitness metrics), with negative binomial error
distributions (to account for overdispersion), lifetime reproductive success as the dependent variable, and
distance moved for first pairing, weighted kinship coefficient to mates, and breeding lifespan as fixed
effects (N: Males = 196, Females = 122). We calculated the weighted kinship coefficient by multiplying
the relatedness of the focal individual to each of their mates by the number of years they were paired with
that mate and summing the resulting values. We defined breeding lifespan as the number of years from
becoming a breeder to death. Random effects included natal year to account for variation between birth
cohorts. Further, to specifically examine the effect of moving away from the breeding territory, we also fit

six additional GLMMs with the same error structure, dependent variables, and random effects, with total
distance moved for later pairings, weighted kinship coefficient to mates, and breeding lifespan as fixed
effects with a truncated dataset that only included individuals that had switched mates in their lifetimes (N:
Male = 92, Female = 51). All GLMMs were fit using the function *glm.nb* in the R package *MASS* [47].

Next, to examine the impact of distance moved on breeder survival, we fit two Cox proportional hazards
models, one for each sex, with distance moved for first pairing as the fixed effect and natal year as a random
effects using the *Surv* and *coxme* functions in the R packages *survival* [48] and *coxme* [49]. For the subset
of individuals that switched mates at least once in their lifetime (N: Male = 92, Female = 51), we also
examined the effect of territory loss (*i.e.*, distance moved for later pairings ≥ 1 territory away) on breeder
survival. We fit Cox proportional hazards models with loss of territory (1 = territory lost, 0 = territory
retained) as the fixed effect and natal year as random effects.

Finally, to investigate whether Florida Scrub-Jays compensated for the impact of inbreeding on
reproductive success via divorce, we fit GLMMs with the cause of pair dissolution (1 = divorce, 0 = mate
death) as the dependent variable; the kinship coefficient to mate as a fixed effect; year and individual ID as
random effects; and a binomial error structure. We fit separate GLMMs by sex and pairing type using the
function *glmer* in the R package *lme4* [41]. Fixed effects for all models were standardized using z scores
[50], and variance inflation factor was < 2 for all fixed effects in all GLMMs [51].

Results

The number of available mates ranged from 14-42 females and 13-45 males across our study period with
an average (± SD) of 26.2 (± 7.70) males and 25.2 (± 7.44) females available as mates per year. These
available mates accounted for, on average, 36% of all breeders per year. Of the individuals included in our
lifetime fitness analysis, 42% of females (51 out of 122) and 47% of males (92 out of 196) switched mates
at least once in their lifetime (mean ± SD number of mate switching instances per lifetime: Males = 0.83 ±
1.17, range: 0-6; Females = 0.73 ± 1.04, range: 0-4). Females, on average, moved farther than males, though
more so for first pairings (Male = 1.05 ± 1.54, Female = 2.04 ± 2.61 territories crossed; *t*-test t = 6.52, df =
631.23, P < 0.001; [32]) than later pairings (Male = 0.34 ± 0.91, Female = 0.62 ± 1.29 territories crossed;
t-test t = 3.73, df = 765.87, P < 0.001). In 212 out of 869 (24%) instances of later pairings, the focal
individual (60% of whom were females) moved one or more territories away from its established breeding
territory. Furthermore, we found that out of all mate switching events observed, ~18% (N: Males = 64 of
187, Females = 31 of 171 events) were due to divorce, not mate death, a similar proportion to that found
previously [31]. The kinship coefficient to mate ranged from 0 (unrelated) to 0.25 (*i.e.,* either siblings or
parent-offspring). Across all years, only 44 out of 1443 (3%) of observed breeding pairs were comprised
of individuals that were at least as closely related as first cousins (*i.e.*, had a kinship coefficient ≥ 0.0625).
As predicted, we found that both sexes were more closely related to available opposite-sex breeders at
shorter geographical distances on both their natal and breeding territories, with the greatest effect size for
females on their natal territories (Males, Natal: Z = -11.40, P < 0.001; Females, Natal: Z = -14.79, P <
0.001; Males, Breeding: Z = -8.49, P < 0.001; Females, Breeding: Z = -10.38, P < 0.001; Fig. 1).

We found that both distance and relatedness to mate influence mate choice decisions in the Florida Scrub-
Jay. The likelihood of pairing with a potential mate increased at shorter geographical distances for both
sexes at both life stages (Males, First pairing: Z = -23.06, P < 0.001; Females, First pairing: Z = -19.86, P
< 0.001; Males, Later pairing: Z = -24.12, P < 0.001; Females, Later pairing: Z = -24.43, P < 0.001; Fig.
2B, D), and males at the first pairing and females at the both life stages showed evidence of discriminating
against more related mates (Males, First pairing: Z = -4.14, P < 0.001; Females, First pairing: Z = -3.27, P
= 0.001; Males, Later pairing: Z = -0.50, P = 0.62; Females, Later pairing: Z = -2.56, P = 0.01; Fig. 2A).
This pattern of the influence of relatedness to mate on mate choice remained the same when limited to mate
options within four territories away, beyond which the likelihood of pairing with close relatives approaches

zero (Fig. 1) (Males, First pairing: $Z = -4.41$, $P < 0.001$; Females, First pairing: $Z = -4.35$, $P < 0.001$; Males,
Later pairing: $Z = -0.49$, $P = 0.62$; Females, Later pairing: $Z = -2.83$, $P = 0.005$; Fig. 2C). Nevertheless, the
effect of weighted kinship coefficient to mate remained considerably smaller than the effect of geographical
distance on mate choice across sexes and pairing types (Fig. 2).

When compared to a random mating scenario, both sexes moved significantly shorter distances within the
study tract than expected under random mating for both first and later pairings (empirical p-value ≤ 0.05
for both), even though females moved farther than males in the first pairing stage (Fig. 3A). The kinship
coefficient to mates for both sexes was never lower than predicted by the random mating model ($P > 0.05$
across all categories). Mates from later pairings were more closely related than expected by chance for both
sexes (Males: $P = 0$; Females: $P = 0$; Fig. 3B), but not for first pairings (Males: $P = 0.25$; Females: $P =$
0.30; Fig. 3B).

Finally, we quantified the fitness consequences of different mate choice decisions using three metrics of
lifetime reproductive success. We found that a higher weighted kinship coefficient to mates over an
individual's lifetime resulted in significantly fewer offspring that survived to the independence and yearling
stages for males, but not females (Fig. 4B, Table S1). This effect extended to offspring surviving to the
fledgling stage when only males that switched mates at least once in their lifetime were considered (Fig.
5B, Table S2). Conversely, females, but not males, that moved farther from their natal territory for their
first pairing had significantly fewer offspring that survived to independence (Fig. 4A, Table S1), and those
that moved farther distances for later pairings had significantly fewer offspring that survived to the
independence and yearling stage (Fig. 5A, Table S2).

Moreover, as we predicted, breeding lifetime—*i.e.*, the number of years an individual spends as a breeder—
had the largest effect on lifetime reproductive success for both sexes: longer breeding lifetimes resulted in
more total offspring that survived to all three life stages (Table S1-2). For both sexes, we found that neither
distance moved for first pairing (Male: $Z = -0.01$, $P = 0.99$; Female: $Z = 0.92$, $P = 0.36$) nor dispersal away
from an established breeding territory in the event of mate switching (Male: $Z = 1.02$, $P = 0.31$; Female: Z
$= -0.25$, $P = 0.81$) impacted breeder survival. The kinship coefficient to their mate had no effect on the
likelihood of divorce in any scenario (Males, First pairing: $Z = 0.47$, $P = 0.64$; Males, Later pairing: $Z = -$
0.03 $P = 0.98$; Females, First pairing: $Z = -0.14$, $P = 0.89$; Females, Later pairing: $Z = 0.66$, $P = 0.51$).

Discussion

In many animals, the costs of inbreeding avoidance, such as traversing a long distance to find an unrelated
mate, might outweigh the benefits of behaviors that increase the likelihood of inbreeding, such as limited
dispersal [21]. Inbreeding may instead be tolerated, and behaviors such as sex-biased dispersal and kin
recognition may evolve to prevent inbreeding [6,52,53]. In cooperatively breeding species, where offspring
often delay dispersal from their natal territory for one or more years and often breed near home, the fitness
benefits of familiarity with the surrounding area and nepotism can lead to extremely limited dispersal [8].
Additionally, since cooperative breeding typically arises in habitats saturated with territories, breeders may
show extreme site fidelity when pairing with a new mate [8]. Both sexes may thus exhibit little propensity
for movement away from their natal and breeding territories, thus increasing the likelihood of pairing with
closely related mates. Here, we investigated the impact of dispersal distance and relatedness on mate choice
in an avian cooperative breeder, the Florida Scrub-Jay. We found that dispersal distance had a greater
impact on mate choice than relatedness. Next, we examined how limited dispersal, and its fitness
consequences, drives inbreeding in this species. We found that fidelity to their breeding territory, rather
than limited movement away from the natal territory, resulted in higher relatedness between mates than
expected by random chance. Moreover, we found that both pairing with close relatives and moving a greater
distance from the natal and breeding territories had a negative impact on lifetime reproductive success but
not survival, though the effects varied by sex. Overall, our results suggest that although sex-biased natal

dispersal acts as an inbreeding avoidance mechanism in Florida Scrub-Jays, the fitness benefits of remaining on established breeding territories may outweigh the costs of breeding with close relatives, resulting in inbreeding tolerance.

We found that Florida Scrub-Jays are more closely related to geographically closer potential mates on both their natal and breeding territories. Previous work has shown that when pairing for the first time, Florida Scrub-Jay females move farther distances than males [32,33]. This sex-bias in distance moved for first pairing is reflected in our finding that females are more closely related to potential mates (*i.e.*, males) closer to their natal territory than males are to potential mates (*i.e.*, females), though both sexes are significantly more related to potential mates at dispersal distances of ≤ 4 territories away from their natal territory than potential mates > 4 territories away. Further, even around the breeding territory, both sexes were more closely related to potential mates at closer geographical distances. Thus, in agreement with previous work examining genetic isolation-by-distance in this species [33], and similar to other communal or cooperatively breeding species [54,55], Florida Scrub-Jays exhibit a viscous population structure such that inbreeding risk is elevated at shorter dispersal distances.

Surprisingly, we found that despite elevated inbreeding risk at shorter dispersal distances and evidence of inbreeding depression in our study population [3], mate choice was driven primarily by distance to mate, with a very small, if any, effect of relatedness to mate. Both sexes were more likely to pair with a geographically closer mate when breeding for the first time or switching mates as an experienced breeder. This degree of philopatry, which has been found in both social and asocial species, can be driven by various factors such as better access to resources and protection from predators due to prior local experience [11], lower competition due to nepotism [56,57], and high physiological costs of dispersal [58]. Although we expected to find a pattern of kin avoidance given that inbred Florida Scrub-Jay offspring have a lower probability of survival [3], we found a relatively small negative effect of kinship on mate choice. Theoretical work suggests that if the frequency and/or cost of inbreeding is low, inbreeding avoidance mechanisms—such as avoiding mating with kin—may not evolve [21]. Further, the cost threshold for the evolution of inbreeding avoidance can be fairly high, especially in the absence of many viable mates [21]. Since our results suggest that geographical distance might constrain the pool of viable mates in our study population, kin discrimination and avoidance may lead to further loss of breeding opportunities, and thus may not be adaptive. Finally, breeders may employ reproductive strategies that minimize the fitness costs of inbreeding over their lifetime. Our results do indicate that males breeding for the first time and females switching mates discriminate against closely related mates to some extent, suggesting that relatedness to mate may be a factor in mate choice in a more nuanced way that merits further study.

To further understand the extent to which limited movement away from natal and breeding territories in mate choice leads to inbreeding, we compared observed means of distance moved and kinship coefficient to mate to those expected under a random mating scenario. We used a similar approach to simulating a random mating scenario as previous studies of breeding populations of territorial birds where most, if not all, individuals were uniquely marked [7,22,25]. However, contrary to previous studies, we differentiated between instances of first and later pairings. We found that neither sex at either pairing stage exhibited significantly lower than expected kinship coefficient to mates. Thus, there was no evidence of inbreeding avoidance. Though we found that the distances moved for first and later pairings are both lower than expected under a random mating scenario for both sexes, female bias in distance moved for first pairing prevents higher relatedness between breeders in first pairings than expected by random chance. We found no sex bias in distance moved for later pairings, which likely leads to the observed higher relatedness to mate than expected under a random mating scenario. Thus, our results suggest that fidelity to the breeding territory, rather than limited natal dispersal, is a driver of inbreeding in Florida Scrub-Jays. To our knowledge, this is the first study to examine the role of breeding site fidelity on inbreeding risk. The factors driving mate choice for later pairings might be drastically different from mate choice for first pairings [8].

398 Future studies on mate choice and inbreeding patterns in this and other species should thus take later life
399 stages into account.
400
Finally, we investigated the lifetime fitness outcomes of dispersal distance and relatedness to mates in
Florida Scrub-Jays. As predicted, we found a negative effect of both dispersal distance and relatedness on
lifetime reproductive success, but the effects varied by sex and were only apparent at later offspring survival
stages. This result is consistent with previous work in Florida Scrub-Jays which found that the effect of
inbreeding on offspring survival increased with number of life stages considered [3]. The total number of
offspring fledged—the standard metric of reproductive success in birds—was only significantly lower for
males with higher weighted kinship coefficient to mates when the data excluded males with only one mate
in their lifetime. However, the number of offspring that survived to independence and yearling stages was
consistently lower for males that mated with closely related females for longer proportions of their breeding
lifetime. On the other hand, relatedness to mate did not significantly affect female lifetime reproductive
success. Instead, females that moved longer distances for both their first and later pairings had lower
lifetime fitness (measured as the total number of independent or yearling offspring produced). We did not
find any evidence of lower male or female breeder survival with higher distance moved for first or later
pairings. These results are contrary to previous work in this population that found that both sexes had lower
lifetime reproductive success and annual survival at greater dispersal distances, with a stronger effect in
males compared to females [35]. However, unlike the previous study, we did not include immigrants from
outside of the study tract in our analysis of lifetime fitness, since, in the context of mate choice, we are not
able to account for all possible mates within their natal population nor accurately measure distance moved.
419
Overall, though effects differed by sex, our results suggest a trade-off in Florida Scrub-Jays between nearly
equivalent costs of mating with close relatives and moving farther distances away from their natal and
breeding territories. We note that our analysis likely underestimates the cost of moving farther distances,
since it does not include individuals that die in transit or fail to establish as breeders and also excludes
dispersers out of the study tract [59]. Thus, since the cost of inbreeding does not outweigh the benefit of
limited dispersal, the population seems to exhibit inbreeding tolerance in accordance with theoretical
expectations [21]. Our results are similar to that found in a range of other animal taxa—including insects
such as the field cricket (*Teleogryllus commodus*) [60] and the parasitoid wasp *Venturia canescens* [61];
birds such as the Eurasian blue tit (*Cyanistes caeruleus*) [23], great tit (*Parus major*) [26], medium ground
finch (*Geospiza fortis*) [25], collared flycatcher (*Ficedula albicollis*) [14], and red-cockaded woodpecker
(*Leuconotopicus borealis*) [22]; and mammals such as reindeer (*Rangifer tarandus*) [62], grey wolf (*Canis
lupus*) [63], and arctic fox (*Vulpes lagopus*) [63]—where higher than expected levels of inbreeding are
tolerated despite some level of inbreeding depression. Interestingly, we found no evidence that Florida
Scrub-Jays are more likely to divorce closely related mates, suggesting that breeding with close relatives is
not simply a response to lack of mate availability [16]. Further, since reproductive success increases with
pair experience in Florida Scrub-Jays [31], mate switching due to divorce may prove costly. While we did
not find evidence of preference for closely related mates, breeding with kin is also thought to confer kin
selection benefits that may additionally contribute to inbreeding tolerance in some species [21,64].
438
In summary, we show that a mate choice that maximizes individual lifetime fitness—*i.e.*, relatively short-
distance dispersal—contributes to some inbreeding in an avian cooperative breeder. Importantly, we show
that it is site fidelity at the later pairing stage, not limited dispersal away from the natal territory, that drives
higher inbreeding than expected under random mating. Despite documented inbreeding depression, we find
no evidence of inbreeding avoidance. Instead, we find that individual lifetime fitness costs of inbreeding
are low enough for inbreeding tolerance and the persistence of limited dispersal. On a population level,
inbreeding tolerance in mate choice might exacerbate inbreeding depression due to increasing population
isolation. Our study underscores the importance of examining the costs and benefits underlying mate choice
at all life stages to better understand the drivers of inbreeding in species of conservation concern.
448

References

449
450

1. Coltman DW, Smith JA, Bancroft DR, Pilkington J, MacColl ADC, Clutton-Brock TH, Pemberton JM.
1999 Density-dependent variation in lifetime breeding success and natural and sexual selection in Soay
rams. *Am. Nat.* **154**, 730–746. (doi:10.1086/303274)

2. Szulkin M, Garant D, Mccleery RH, Sheldon BC. 2007 Inbreeding depression along a life-history
continuum in the great tit. *Journal of Evolutionary Biology* **20**, 1531–1543. (doi:10.1111/j.1420-
9101.2007.01325.x)

3. Chen N, Cosgrove EJ, Bowman R, Fitzpatrick JW, Clark AG. 2016 Genomic consequences of
population decline in the endangered Florida scrub-jay. *Curr. Biol.* **26**, 2974–2979.
(doi:10.1016/j.cub.2016.08.062)

4. Keller L, Reeve HK. 1994 Partitioning of reproduction in animal societies. *Trends Ecol. Evol.* **9**, 98–
102. (doi:10.1016/0169-5347(94)90204-6)

5. Keller LF, Waller DM. 2002 Inbreeding effects in wild populations. *Trends Ecol. Evol.* **17**, 230–241.
(doi:10.1016/S0169-5347(02)02489-8)

6. Nichols HJ. 2017 The causes and consequences of inbreeding avoidance and tolerance in cooperatively
breeding vertebrates. *J. Zool.* **303**, 1–14. (doi:10.1111/jzo.12466)

7. Billing AM *et al.* 2012 Evidence of inbreeding depression but not inbreeding avoidance in a natural
house sparrow population. *Molecular Ecology* **21**, 1487–1499. (doi:10.1111/j.1365-
294X.2012.05490.x)

8. Greenwood PJ, Harvey PH. 1982 The natal and breeding dispersal of birds. *Annu. Rev. Ecol. Syst.* **13**,
1–21. (doi:10.1146/annurev.es.13.110182.000245)

9. Szulkin M, Stopher KV, Pemberton JM, Reid JM. 2013 Inbreeding avoidance, tolerance, or preference
in animals? *Trends Ecol. Evol.* **28**, 205–211. (doi:10.1016/j.tree.2012.10.016)

10. Clutton-Brock TH, Lukas D. 2012 The evolution of social philopatry and dispersal in female mammals.
Mol. Ecol. **21**, 472–492. (doi:10.1111/j.1365-294X.2011.05232.x)

11. Pärt T. 1994 Male philopatry confers a mating advantage in the migratory collared flycatcher, *Ficedula*
albicollis. *Animal Behaviour* **48**, 401–409. (doi:10.1006/anbe.1994.1254)

12. Griesser M, Nystrand M, Ekman J. 2006 Reduced mortality selects for family cohesion in a social
species. *Proceedings of the Royal Society B: Biological Sciences* **273**, 1881–1886.
(doi:10.1098/rspb.2006.3527)

13. Thünken T, Bakker TCM, Baldauf SA, Kullmann H. 2007 Active inbreeding in a cichlid fish and its
adaptive significance. *Current Biology* **17**, 225–229. (doi:10.1016/j.cub.2006.11.053)

14. Pärt T. 1996 Problems with testing inbreeding avoidance: the case of the collared flycatcher. *Evolution*
50, 1625–1630. (doi:10.1111/j.1558-5646.1996.tb03934.x)

15. Daniels SJ. 1997 Female dispersal and inbreeding in the red-cockaded woodpecker. Virginia
Polytechnic Institute and State University.

16. Szulkin M, Sheldon BC. 2008 Dispersal as a means of inbreeding avoidance in a wild bird population.
Proc. R. Soc. B. **275**, 703–711. (doi:10.1098/rspb.2007.0989)

17. Beck KB, Loretto M-C, Bugnyar T. 2020 Effects of site fidelity, group size and age on food-caching
behaviour of common ravens, *Corvus corax*. *Animal Behaviour* **164**, 51–64.
(doi:10.1016/j.anbehav.2020.03.015)

18. Schmidt KA. 2014 Site fidelity curbs sequential search and territory choice: a game theoretical
approach. *Funct. Ecol.* **28**, 1494–1503. (doi:10.1111/1365-2435.12291)

19. Byrne ME, Cohen BS, Collier BA, Chamberlain MJ. 2022 Nest site fidelity and nesting success of
female wild turkeys. *Wildlife Society Bulletin* **46**, e1279. (doi:10.1002/wsb.1279)

20. Hamilton WD. 1964 The genetical evolution of social behaviour. I. *J. Theor. Biol.* **7**, 1–16.
(doi:10.1016/0022-5193(64)90038-4)

21. Kokko H, Ots I. 2006 When not to avoid inbreeding. *Evol.* **60**, 467–475. (doi:10.1111/j.0014-
3820.2006.tb01128.x)

22. Daniels SJ, Walters JR. 2000 Inbreeding depression and its effects on natal dispersal in red-cockaded
woodpeckers. *The Condor* **102**, 482–491. (doi:10.1093/condor/102.3.482)

23. Foerster K, Valcu M, Johnsen A, Kempenaers B. 2006 A spatial genetic structure and effects of
relatedness on mate choice in a wild bird population. *Molecular Ecology* **15**, 4555–4567.
(doi:10.1111/j.1365-294X.2006.03091.x)

24. Frommen JG, Bakker TCM. 2006 Inbreeding avoidance through non-random mating in sticklebacks.
Biology Letters **2**, 232–235. (doi:10.1098/rsbl.2005.0432)

25. Gibbs HL, Grant PR. 1989 Inbreeding in Darwin's medium ground finches (Geospiza fortis). *Evolution*
43, 1273–1284. (doi:10.1111/j.1558-5646.1989.tb02574.x)

26. Szulkin M, Zelazowski P, Nicholson G, Sheldon BC. 2009 Inbreeding avoidance under different null
models of random mating in the great tit. *Journal of Animal Ecology* **78**, 778–788.

27. Wheelwright NT, Mauck RA. 1998 Philopatry, natal dispersal, and inbreeding avoidance in an island
population of savannah sparrows. *Ecology* **79**, 755–767. (doi:10.1890/0012-
9658(1998)079[0755:PNDAIA]2.0.CO;2)

28. Beauchamp G. 2014 Do avian cooperative breeders live longer? *Proceedings of the Royal Society B:
Biological Sciences* **281**, 20140844. (doi:10.1098/rspb.2014.0844)

29. Arnold KE, Owens IPF. 1998 Cooperative breeding in birds: a comparative test of the life history
hypothesis. *Proc. R. Soc. Lond. B* **265**, 739–745. (doi:10.1098/rspb.1998.0355)

30. Fitzpatrick JW, Bowman R. 2016 Florida scrub-jays: Oversized territories and group defense in a fire-
maintained habitat. In *Cooperative Breeding in Vertebrates* (eds WD Koenig, JL Dickinson), pp. 77–
96. Cambridge University Press. (doi:10.1017/CBO9781107338357.006)

31. Woolfenden GE, Fitzpatrick JW. 1984 *The Florida Scrub Jay: Demography of a Cooperative-breeding
Bird*. Princeton University Press.

32. Suh YH, Pesendorfer MB, Tringali A, Bowman R, Fitzpatrick JW. 2020 Investigating social and
environmental predictors of natal dispersal in a cooperative breeding bird. *Behav. Ecol.* **31**, 692–701.
(doi:10.1093/beheco/araa007)

33. Aguillon SM, Fitzpatrick JW, Bowman R, Schoech SJ, Clark AG, Coop G, Chen N. 2017
Deconstructing isolation-by-distance: The genomic consequences of limited dispersal. *PLoS Genet.* **13**,
e1006911. (doi:10.1371/journal.pgen.1006911)

34. Marzluff JM, Woolfenden GE, Fitzpatrick John W, Balda RP. 1996 Breeding partnerships of two New
World jays. In *Partnerships in Birds: The Study of Monogamy* (eds JM Black, M Hulme), p. 0. Oxford
University Press. (doi:10.1093/oso/9780198548614.003.0007)

35. Fitzpatrick JW, Woolfenden GE, Bowman R. 1999 Dispersal distance and its demographic
consequences in the Florida Scrub-jay. In *Proceedings of the International Ornithological Congress*,

36. Townsend AK, Bowman R, Fitzpatrick JW, Dent M, Lovette IJ. 2011 Genetic monogamy across
variable demographic landscapes in cooperatively breeding Florida scrub-jays. *Behavioral Ecology* **22**,
464–470. (doi:10.1093/beheco/arq227)

37. R Core Team. 2022 R: A Language and Environment for Statistical Computing.

38. Suh YH, Bowman R, Fitzpatrick JW. 2022 Staging to join non-kin groups in a classical cooperative
breeder, the Florida scrub-jay. *J. Anim. Ecol.* **91**, 970–982. (doi:10.1111/1365-2656.13669)

39. Pebesma E. 2018 Simple features for R: Standardized support for spatial vector data. *The R Journal*
10, 439. (doi:10.32614/RJ-2018-009)

40. Malécot G. 1948 *Les mathématiques de l'hérédité*. Paris: Masson.

41. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using lme4. *J.*
Stat. Softw. **67**, 1–48. (doi:10.18637/jss.v067.i01)

42. Mumme RL, Bowman R, Pruett MS, Fitzpatrick JW. 2015 Natal territory size, group size, and body
mass affect lifetime fitness in the cooperatively breeding Florida Scrub-Jay. *Auk* **132**, 634–646.
(doi:10.1642/AUK-14-258.1)

43. McGowan KJ, Woolfenden GE. 1990 Contributions to fledgling feeding in the Florida Scrub-Jay.
Journal of Animal Ecology **59**, 691–707. (doi:10.2307/4889)

44. Schoech SJ, Mumme RL, Wingfield JC. 1996 Delayed breeding in the cooperatively breeding Florida
scrub-jay (Aphelocoma coerulescens): inhibition or the absence of stimulation? *Behav Ecol Sociobiol*
39, 77–90. (doi:10.1007/s002650050269)

45. Barve S, Riehl C, Walters EL, Haydock J, Dugdale HL, Koenig WD. 2021 Lifetime reproductive
benefits of cooperative polygamy vary for males and females in the acorn woodpecker (Melanerpes
formicivorus). *Proceedings of the Royal Society B: Biological Sciences* **288**, 20210579.
(doi:10.1098/rspb.2021.0579)

46. Shah SS, Rubenstein DR. 2022 Prenatal environmental conditions underlie alternative reproductive
tactics that drive the formation of a mixed-kin cooperative society. *Sci. Adv.* **8**, eabk2220.
(doi:10.1126/sciadv.abk2220)

47. Venables W, Ripley B. 2002 *Modern Applied Statistics with S*. 4th edn. New York: Springer. See
https://www.stats.ox.ac.uk/pub/MASS4/.

48. Therneau T. 2022 A Package for Survival Analysis in R.

49. Therneau T. 2022 coxme: Mixed Effects Cox Models. R package.

50. Schielzeth H. 2010 Simple means to improve the interpretability of regression coefficients. *Methods*
Ecol. Evol. **1**, 103–113. (doi:10.1111/j.2041-210X.2010.00012.x)

51. Fox J, Monette G. 1992 Generalized Collinearity Diagnostics. *Journal of the American Statistical*
Association **87**, 178–183.

52. Pusey A, Wolf M. 1996 Inbreeding avoidance in animals. *Trends Ecol. Evol.* **11**, 201–206.
(doi:10.1016/0169-5347(96)10028-8)

53. Leedale AE, Simeoni M, Sharp SP, Green JP, Slate J, Lachlan RF, Robinson EJH, Hatchwell BJ. 2020
Cost, risk, and avoidance of inbreeding in a cooperatively breeding bird. *Proceedings of the National*
Academy of Sciences **117**, 15724–15730. (doi:10.1073/pnas.1918726117)

54. Double MC, Peakall R, Beck NR, Cockburn A. 2005 Dispersal, philopatry, and infidelity: dissecting
local genetic structure in superb fairy-wrens (*Malurus cyaneus*). *Evol.* **59**, 625–635.

55. Leedale AE, Sharp SP, Simeoni M, Robinson EJH, Hatchwell BJ. 2018 Fine-scale genetic structure
and helping decisions in a cooperatively breeding bird. *Mol. Ecol.* **27**, 1714–1726.
(doi:10.1111/mec.14553)

56. Kokko H, Ekman J. 2002 Delayed dispersal as a route to breeding: territorial inheritance, safe havens,
and ecological constraints. *Am. Nat.* **160**, 468–484. (doi:10.1086/342074)

57. Green JP, Hatchwell BJ. 2018 Inclusive fitness consequences of dispersal decisions in a cooperatively
breeding bird, the long-tailed tit (*Aegithalos caudatus*). *Proc. Natl. Acad. Sci. U.S.A.* **115**, 12011–
12016. (doi:10.1073/pnas.1815873115)

58. Maag N, Cozzi G, Bateman A, Heistermann M, Ganswindt A, Manser M, Clutton-Brock T, Ozgul A.
2019 Cost of dispersal in a social mammal: body mass loss and increased stress. *Proc. R. Soc. B.* **286**,
20190033. (doi:10.1098/rspb.2019.0033)

59. Bonte D *et al.* 2012 Costs of dispersal. *Biol. Rev.* **87**, 290–312. (doi:10.1111/j.1469-
185X.2011.00201.x)

60. Jennions MD, Hunt J, Graham R, Brooks R. 2004 No evidence for inbreeding avoidance through
postcopulatory mechanisms in the black field cricket, teleogryllus commodus. *Evolution* **58**, 2472–
2477. (doi:10.1111/j.0014-3820.2004.tb00876.x)

61. Collet M, Amat I, Sauzet S, Auguste A, Fauvergue X, Mouton L, Desouhant E. 2020 Insects and incest:
Sib-mating tolerance in natural populations of a parasitoid wasp. *Molecular Ecology* **29**, 596–609.
(doi:10.1111/mec.15340)

62. Holand Ø, Askim KR, Røed KH, Weladji RB, Gjøstein H, Nieminen M. 2006 No evidence of inbreeding avoidance in a polygynous ungulate: the reindeer (Rangifer tarandus). *Biology Letters* **3**, 36–39. (doi:10.1098/rsbl.2006.0575)

63. Geffen E *et al.* 2011 Kin encounter rate and inbreeding avoidance in canids. *Molecular Ecology* **20**, 5348–5358. (doi:10.1111/j.1365-294X.2011.05358.x)

64. Puurtinen M. 2011 Mate choice for optimal (k)inbreeding. *Evolution* **65**, 1501–1505. (doi:10.1111/j.1558-5646.2010.01217.x)

601   **Figures**
602
603

604
Figure 1. Both male (blue) and female (orange) Florida Scrub-Jays are significantly more likely to mate
with a close relative if they move shorter distances from both their natal ("First Pairings") and breeding
territories ("Later Pairings"). Lines indicate model fits and shaded areas 95% confidence intervals. We
defined close relatives as opposite sex individuals that are at least as closely related as first cousins to the
focal individuals (1 = close relative, 0 = not close relative). We included all instances where individuals
become breeders for the first time that year or switched mates (*N*: Male, First pairing = 421; Male, Later
pairing = 249; Female, First pairing = 391; Female, Later pairing = 256 instances).
612
613

615
Figure 2. The predicted effect of relatedness and distance to mate (solid line = significant, dashed line =
not significant effect) on mate choice in both sexes (blue = male, orange = female) of Florida Scrub-Jays
with (A-B) the full dataset from our study tract (*N*: Male, First pairing = 12155; Male, Later pairing =
11837; Female, First pairing = 10938; Female, Later pairing = 11000 possible pairings) and (C-D) only
including potential mates located four or fewer territories away (*N*: Male, First pairing = 5202; Male, Later
pairing = 5013; Female, First pairing = 4525; Female, Later pairing = 4686 possible pairings). While
relatedness to mate significantly lowered likelihood of pairing for males at the first pairing and females at
the later pairing stage, with the effect extending to females at the first pairing stage when choice of mate
was limited to ≤ 4 territories away, the effect of relatedness to mate was minimal compared to that of
distance to mate. Both sexes were more likely to pair with mates at shorter distances for both first and later
pairings. Shaded areas around predicted fits depict 95% confidence intervals.

627
628
Figure 3. Observed and expected (under random mating) (A) dispersal distance and (B) relatedness to mate
for male (blue) and female (orange) Florida Scrub-Jays at both the first (filled bars) and later (unfilled bars)
pairing stages. Histograms show the distributions of expected means under 1000 simulations of random
mating. The observed population means across all 32 years of our study period are indicated by vertical
lines (solid = significant difference, dashed = no significant difference from expectations under a random
mating scenario). Observed distances moved were significantly lower than expected for all four categories,
while kinship coefficient to mate was higher than expected for both sexes at the later, but not first, pairing
stages.

637
Figure 4. Predicted model fits investigating the effects (solid line = significant, dashed line = not
significant) of (A) Distance moved for first pairing, and (B) Weighted kinship coefficients to mates on three
metrics of lifetime reproductive success (LRS)—number of offspring surviving to fledgling, independence,
and yearling stages—for males (blue) and females (orange) (N: Males = 196, Females = 122). Breeding
lifespan (not shown) was also included as a fixed effect in the model, since the longer a breeder lives the
more chances it has to accrue greater LRS. Females had reduced LRS with increasing distance moved for
first pairing, while males showed a relatively similar reduction in LRS with increased kinship to mates over
their lifetime.
646

647
Figure 5. Predicted model fits investigating the effects (solid line = significant, dashed line = not
significant) of (A) Total distance moved for later pairings, and (B) Weighted kinship coefficients to mates
on three metrics of lifetime reproductive success (LRS)—number of offspring surviving to fledgling,
independence, and yearling stages—for males (blue) and females (orange) that paired with more than one
mate in their lifetime (*N*: Males = 92, Females = 51). As in the previous model, breeding lifespan (not
shown) was also included as a fixed effect since the longer a breeder lives the more chances it has to accrue
more LRS. Females had reduced LRS with increasing distance moved for later pairings, while males
showed a relatively similar reduction in LRS with increased kinship to mates over their lifetime.
656
657