Thermal inactivation spectrum of influenza A H5N1 virus in raw milk

Mohammed Nooruzzaman[1], Lina M. Covaleda[1], Nicole H. Martin[2], Katherine Koebel[1], Renata

Ivanek[1], Samuel D. Alcaine[2], Diego G. Diel[1, *]

[1]Department of Population Medicine and Diagnostic Sciences, College of Veterinary Medicine,

Cornell University, Ithaca, NY, USA.

[2]Department of Food Science, College of Agriculture and Life Sciences, Cornell University,

Ithaca, NY, USA.

* Corresponding author: Diego G. Diel; Email: dgdiel@cornell.edu.

Abstract

The spillover of highly pathogenic avian influenza (HPAI) H5N1 virus to dairy cows and shedding of high amounts of infectious virus in milk raised public health concerns. Here, we evaluated the decay and thermal stability spectrum of HPAI H5N1 virus in raw milk. For the decay studies, HPAI H5N1 positive raw milk was incubated at different temperatures and viral titers and the thermal death time D-values were estimated. We then heat treated HPAI H5N1 virus positive milk following different thermal conditions including pasteurization and thermization conditions. Efficient inactivation of the virus was observed in all tested conditions, except for thermization at 50°C 10 min. Utilizing a submerged coil system with temperature ramp up times that resemble commercial pasteurizers, we showed that the virus was rapidly inactivated by pasteurization and most thermization conditions. These results provide important insights on the food safety measures utilized in the dairy industry.

Introduction

Highly pathogenic avian influenza (HPAI) H5N1 virus clade 2.3.4.4b was first detected in the United States on February 8, 2022[1]. Since then, the virus has been detected in over 1,170 poultry flocks in 48 states resulting in the death or culling of more than 100 million birds[1]. A unique feature observed throughout the HPAI H5N1 panzootic, is the frequent spillover of the virus to wild terrestrial and aquatic mammalian species with at least 24 species confirmed positive to date[2,3], including important livestock species such as goats and dairy cows [4–6]. Spillover of HPAI H5N1 virus to dairy cows and detection of high infectious viral loads and shedding in milk from infected cows[6,7] raised public health concerns given the potential for zoonotic spillover of the virus. As of September 10, 2024, there have been 15 confirmed human cases in the USA since the beginning of the HPAI H5N1 panzootic in 2022[8], 14 of those documented since the reported spillover of the virus to dairy cows in March 2024. Importantly, four of the 2024 human cases have been linked to occupational exposures to dairy cows, nine were linked to exposures to poultry, while, the most recent case reported by CDC on September 6, 2024, in Missouri - a state where the virus has not yet been detected in dairy cows – is the first human case without a known occupational exposure to infected animals[8].

Thermal treatment of milk or pasteurization is the main procedure used by the dairy industry to ensure safety of dairy products and extend their shelf-life by reducing and destroying non-pathogenic and pathogenic microorganisms[9]. Currently, in the USA two pasteurization conditions are approved by the U.S. Food and Drug Administration (FDA) for milk including low temperature long time (LTLT) treatment at or above 63°C for 30 min and high temperature short time (HTST) treatment at or above 72°C for 15 sec. Early studies reporting the detection of HPAI H5N1 viral RNA in retail milk and other dairy products, such as cheese, butter and ice cream[10,11], coupled with laboratory studies suggesting that heat treatment of raw HPAI H5N1 positive milk at 72°C for 15 sec, the most common pasteurization method used in the USA, did not completely inactivate the virus[12] heightened public health concerns about the potential risk of human infections through milk. Confirmation that no infectious virus was present in retail milk products subjected to pasteurization eased some of the initial public health concerns[11]. However, the public health risk posed by consumption of raw milk or milk subjected to subpasteurization conditions, a process known as thermization which is milder than pasteurization and involves heating to a

temperature typically between 50-68°C for about 15 to 30 sec to up to 10 min, remains unknown.
Notably, while thermization reduces the number of non-sporeforming pathogens and
psychrotrophic spoilage bacteria, it does not destroy all pathogenic microorganisms present in
milk[13,14].

Here, we investigated the decay of HPAI H5N1 virus in milk and evaluated the efficacy of different
heat treatment conditions including the FDA approved pasteurization conditions and several
thermization conditions on inactivation of HPAI H5N1. The work provides a comprehensive
overview of the thermal inactivation spectrum of HPAI H5N1 virus in raw milk.

Results

Highly pathogenic avian influenza (HPAI) H5N1 virus presents long term stability in raw
milk at 4ºC but is rapidly inactivated at higher temperatures. We studied the decay of HPAI
H5N1 virus in raw milk at different temperatures. For this, we incubated HPAI H5N1 virus positive
raw milk collected from infected cows or normal raw milk spiked with HPAI H5N1 virus at 4°C,
20°C, 30°C and 37°C and collected sequential samples daily between days 1-7 and weekly
thereafter until day 56 (8 weeks). HPAI H5N1 virus decay over time was assessed by viral titrations
performed in bovine uterine epithelial cells (Cal-1). Samples that were toxic or negative in the cell
culture titration assay were subjected to confirmatory virus isolation in embryonated chicken eggs
(ECEs). Virus titers were used to estimate the thermal death time values (D-values) in each studied
temperature. At 4°C, there was about 4 log reduction in infectious virus titers from 5.97 ± 0.14 log
$TCID_{50}$/mL on day 0 to 2.05 ± 0.43 log $TCID_{50}$/mL on day 42 **(Fig. 1a)**. Samples from days 49
and 56 were toxic to cell culture, but when inoculated in ECEs both samples were confirmed to
contain infectious HPAI H5N1 virus (**Extended Data Table 1**), demonstrating the long-term
stability of the virus in raw milk stored 4°C. The D-value of HPAI at 4°C was estimated to be
10.74 days.

At 20°C, we observed 3 log reduction in infectious virus titers within seven days, with
titers dropping from 6.88 ± 0.14 log $TCID_{50}$/mL on day 0 to 3.88 ± 0.14 log $TCID_{50}$/mL on day 7
(Fig. 1b). Samples collected on days 14 and onward were toxic to cell culture and, thus, were
subjected to inoculation in ECEs. One out of three milk samples collected on day 14 led to embryo
mortality, which was confirmed to be caused by HPAI H5N1 virus by hemagglutination (HA)
assay (**Extended Data Table 1**). No embryo mortality was observed in milk samples collected on

day 21 and the allantoic fluids collected from inoculated ECEs were HA negative, indicating virus inactivation in milk stored at 20°C at day 21 (**Extended Data Table 1**). The estimated D-value of HPAI H5N1 virus in milk stored at 20°C was 2.23 days.

At 30°C, we detected a rapid decrease in infectious HPAI H5N1 virus titers in milk, which dropped from 5.88 ± 0.14 log $TCID_{50}$/mL on day 0 to 1.63 ± 0.29 log $TCID_{50}$/mL on day 5 resulting in about 4-log reduction in infectious virus titers **(Fig. 1c)**. No infectious virus was detected in milk samples from day 6 and onward when we used both cell culture and ECEs to assess virus infectivity **(Fig. 1c, Extended Data Table 1)**. The estimated D-value of HPAI H5N1 virus in milk stored at 30°C was 1.16 days.

At 37°C, we observed a drastic drop in HPAI H5N1 virus titers, with a 3-log reduction in virus infectivity within the first day of incubation, decreasing from 5.88 ± 0.14 log $TCID_{50}$/mL on day 0 to 2.88 ± 0.14 log $TCID_{50}$/mL on day 1 **(Fig. 1d)**. From day 2 and onwards, no infectious virus was detected when samples were inoculated in cell culture and ECEs **(Fig. 1d, Extended Data Table 1)**. The estimated D-value of HPAI H5N1 virus in milk stored at 37°C was 0.34 days. Together these results show that HPAI H5N1 virus presents long term stability in raw milk stored at 4°C, with viral titers decreasing rapidly at temperatures above 30°C.

Thermal treatment of raw HPAI H5N1 virus positive milk following thermization conditions and FDA approved pasteurization results in efficient virus inactivation. Initially, we evaluated the efficacy of thermization at 60°C for 10 min and the FDA approved pasteurization conditions (63°C for 30 min and 72°C for 15 sec), on the inactivation of HPAI H5N1 virus in raw milk collected from clinically affected cows or in raw milk spiked with HPAI H5N1 virus utilizing a thermocycler. Following heat treatment, milk samples were subjected to virus inoculation and titrations in Cal-1 cells and ECEs. While robust virus replication was detected by immunofluorescence (IFA) in cells inoculated with control non-heat-treated milk samples, no evidence of virus infectivity in raw milk subjected to heat treatment at 60°C for 10 min and at 63°C for 30 min or 72°C for 15 sec was observed in Cal-1 cells (**Fig. 2a**).

We then quantified viral RNA by qRT-PCR and determined infectious virus titers in control and heat-treated milk samples using cell culture and ECEs. Similar levels of viral RNA were detected in control and in heat-treated milk samples (**Fig. 2b**). Importantly, viral titrations revealed that non-heat-treated raw milk clinical samples had a virus titer of 6.47 ± 0.29 log $TCID_{50}$/mL and

7.47 ± 0.29 log EID_{50}/mL, while no infectious virus was detected after heat treatment of the raw

milk clinical samples, in cell culture or ECEs, respectively (**Fig. 2c-2d**).

We further assessed the thermal inactivation of HPAI H5N1 in normal raw milk spiked

(1:10 ratio) with raw milk positive for HPAI H5N1 collected from clinically affected cows. We

followed the same heat treatment conditions described above utilizing a thermocycler. Similar to

the results observed in raw milk from clinical cows, robust HPAI H5N1 virus replication was

observed in Cal-1 cells inoculated with non-heat-treated spiked milk samples, whereas no virus

replication was observed after heat treatment at 60°C for 10 min or at 63°C for 30 min or 72°C for

15 sec (**Fig. 2e**). While the viral RNA levels remained stable after heat treatment (**Fig. 2f**),

complete virus inactivation was observed in spiked milk samples following heat treatment (**Fig.**

2g-2h). These results suggest that thermization at 60°C for 10 min and pasteurization at 63°C for

30 min or 72°C for 15 sec efficiently led to HPAI H5N1 virus inactivation in raw milk.

Evaluation of thermization conditions on HPAI H5N1 virus inactivation in raw milk. We next

evaluated the efficacy of three additional thermization conditions on HPAI H5N1 virus inactivation

in raw milk. For this, raw normal milk was spiked with a HPAI H5N1 virus stock amplified in Cal-

1 cells (1:20 ratio) and subjected to thermal treatment at 50°C for 10 min, 63°C for 22 sec and

69°C for 22 sec utilizing a thermocycler. Inoculation of the control and heat-treated samples in

Cal-1 cells revealed robust HPAI H5N1 virus replication in control non-treated raw spiked milk

samples. Additionally, HPAI H5N1 virus replication was detected by IFA in cells inoculated with

raw spiked milk heat treated at 50°C for 10 min, however, the proportion of infected cells and the

intensity of staining was lower compared to the control non-treated raw milk samples (**Fig. 3a**).

No virus replication was detected in cells inoculated with raw HPAI-spiked milk samples heated

at 63°C for 22 sec and 69°C for 22 sec, suggesting efficient virus inactivation under these thermal

conditions.

Quantification of HPAI H5N1 virus RNA by qRT-PCR revealed similar viral RNA levels

in control non-treated and all three heat-treated raw spiked milk samples (**Fig. 3b**). Viral titrations

have shown that heat treatment of raw HPAI-spiked milk at 50°C for 10 min led to 1.5-2 log

reduction in infectious virus titers, which decreased from 4.94 ± 0.21 to 3.19 ± 0.81 log $TCID_{50}$/mL

or 4.87 ± 0.13 to 2.87 ± 0.35 log EID_{50}/mL after heat treatment (**Fig. 3c-d**). No infectious virus

was detected in raw HPAI-spiked milk samples heat treated at 63°C for 22 sec and 69°C for 22

sec. These results show that while thermization at 50°C for 10 min does not completely inactivate HPAI H5N1 virus in raw milk, higher temperatures of 63 and 69°C efficiently and rapidly inactivate the virus in raw milk.

Validation of thermal inactivation of HPAI H5N1 virus in raw milk using a submerged coil system. To more closely simulate the thermal treatment conditions (e.g. circulating hot water heat source and rapid ramp up temperatures) to which raw fluid milk is subjected to during commercial pasteurization in the dairy industry, we developed a submerged coil heating system to heat treat milk samples (**Fig. 4a**). The system holds 15 mL of milk within a coil and is heated with a circulating water bath. The submerged coil heating system is furnished with four thermal sensors that are connected to a data logger to continuously monitor and record the milk temperature as it flows through four segments of the coil (**Fig. 4a**). The first coil segment (capacity of 4.2 mL) contains two portions, including one external coil component carrying the sample injection port and a temperature sensor (T1) (**Fig. 4a**) to monitor the temperature of the milk at the injection port (**Fig. 4b**); and one submerged coil component (**Fig. 4a**) which preheats the milk sample to the desired temperature within 7-10 sec (**Fig. 4b**) as it flows through this portion of the coil. The temperature of the milk that flows through the pre-heating coil segment is recorded by thermal sensor T2 (**Fig. 4a** and **4b**). The second (capacity 5.5 mL) and third (capacity 5.7 mL) coil segments are completely submerged in water and hold the sample at the target temperature for the intended time. The thermal sensor T3 monitors the milk temperature in the holding coil segment (**Fig. 4a** and **4b**). The fourth coil segment (capacity of 1.5 mL) consists of the exit loop which dispenses the milk in a collection tube after the heat treatment. Cooling of the milk after thermal treatment was achieved by placing the collection tubes on ice for 10 min, immediately after sample collection.

To validate the results obtained with the thermocycler, we evaluated the thermal inactivation of HPAI in raw milk using the submerged coil system following the FDA approved pasteurization conditions at 63°C for 30 min and 72°C for 15 s, and four thermization conditions at 50°C for 10 min, 60°C for 10 min, 63°C for 22s and 69°C for 22s. For each thermal treatment condition, 15 mL of raw milk spiked with HPAI H5N1 virus (1:20 ratio) were injected into the submerged coil heating system and subjected to heat treatment at the desired temperatures and times. Temperature monitoring as recorded by the thermal sensors data loggers following each

treatment confirmed that the milk samples were subjected to the heat treatment at the target
temperatures (**Fig. 4b, Extended Data Fig. 1**). Initially, virus infectivity was monitored by
inoculation of control and thermal treated spiked milk samples in Cal-1 cells. As shown in **Fig. 4c**,
virus replication was detected in control non-treated samples and in milk samples subjected to heat
treatment at 50°C for 10 min, as evidenced by positive viral NP staining in inoculated cells. Similar
to the results obtained with the experiments conducted in the thermocycler, a lower number of NP
positive cells were observed in cells inoculated with milk samples subjected to heat treatment at
50°C for 10 min, when compared to control non-treated samples (**Fig. 4c**). No evidence of virus
replication was detected in HPAI-spiked milk samples heat treated at 60°C for 10 min, 63°C for
22 s, 63°C for 30 min, 69°C for 22 s and 72°C for 15 s.

Next, we quantified viral RNA and infectious virus in all control and thermal treated
samples. The qRT-PCR results revealed comparable levels of viral RNA in both heat-treated and
non-treated milk samples (**Fig. 4d**). Virus quantifications in Cal-1 cells or ECEs demonstrated viral
titers of 4.66 ± 0.19 log $TCID_{50}$/mL and 4.54 ± 0.03 log EID_{50}/mL in untreated milk samples,
which decreased by one log in milk samples heat treated at 50°C for 10 min, with titers of $3.69 \pm$
0.38 log $TCID_{50}$/mL and 3.5 log EID_{50}/mL (**Fig. 4e-f**). Notably, no infectious virus titers were
detected in milk samples heat treated at 60°C for 10 min, 63°C for 22 s, 63°C for 30 min, 69°C for
22 s and 72°C for 15 s. These results corroborate the findings of our studies using a thermocycler
and demonstrate the efficacy of a broad spectrum of thermal treatment conditions on inactivation
of HPAI H5N1 virus in raw milk.

Kinetics of HPAI H5N1 virus inactivation by thermal treatment in raw milk. We evaluated
the kinetics of thermal inactivation of HPAI H5N1 virus in raw milk at different temperatures over
time using the submerged coil system. For this, raw HPAI-spiked milk was subjected to the thermal
treatment and samples were collected at different time points as follows: 50°C - 5, 10, 15 20, 25,
and 30 min; 60°C - 5, 10, 25, 20, 30, and 60 sec; 63°C - 5, 10, 15, 20, and 30 sec; and 72°C - 5, 7,
10, 12, and 15 sec. Viral RNA and infectious virus were quantified by qRT-PCR and virus titrations
(cells and ECEs), respectively. Viral RNA levels detected in untreated and heat-treated milk
samples were comparable in all temperatures evaluated in the study (**Fig. 5a, d, g, j**). At 50°C, we
observed a gradual decrease in infectious virus titers from 4.88 ± 0.14 to 2.30 ± 0.25 log
$TCID_{50}$/mL within 25 min heat treatment with virus titers falling under the titration detection limit

between the 25- and 30-min time point (**Fig. 5b**). Similar results were observed in ECEs with viral virus titers decreasing from 4.58 ± 0.07 to 1.17 ± 1.01 log EID_{50}/mL within 25 min (**Fig. 5c**). Of note, one out of three milk samples heat treated at 50°C for 30 min showed embryo mortality and HA activity, indicating the presence of low amounts of infectious virus in this sample. The D-value of HPAI H5N1 at 50°C was estimated at 10.11 min. Notably, rapid virus inactivation of HPAI H5N1 was observed in raw milk samples heat treated at 60°C (**Fig. 5e, 5f**), 63°C (Fig. **5h, 5i**) and 72°C (**Fig. 5k, 5l**), with heating for 5 sec leading to complete virus inactivation at these temperatures. These results demonstrate that HPAI H5N1 virus is stable for at least 30 min at 50°C in raw milk. Importantly, however, efficient and rapid inactivation (within five seconds) of HPAI H5N1 virus is achieved in raw milk heat treated at temperatures of 60°C or above.

Discussion

The high tropism of HPAI H5N1 virus and its replication in milk secreting epithelial cells of the mammary gland in dairy cows, lead to high viral loads and shedding (from $10^{4.4}$ to up $10^{8.8}$ $TCID_{50}.mL^{-1}$) in milk[6,7,15]. Despite the widespread use of milk pasteurization by the dairy industry, the high viral loads detected in milk from infected cows raised major public health concerns. Although it has been shown that both high (HPAI)- and low pathogenic avian influenza (LPAI) viruses can be inactivated by pasteurization of various poultry products including fat-free egg products, allantoic fluid and plasma[16], and milk coproduct concentrated lactose[17], in the beginning of the dairy outbreak there was no data demonstrating the efficacy of pasteurization on HPAI H5N1 virus in dairy products. Initial laboratory studies evaluating inactivation of HPAI in raw milk revealed that the most used FDA approved milk pasteurization condition (72°C for 15 sec) markedly reduced viral loads in raw milk, however, low amounts of infectious virus were still detected in heat treated samples[12,18]. Here we evaluated the stability and determined the decay of HPAI H5N1 virus in raw milk stored at different temperatures over time and investigated the efficacy and kinetics of inactivation of HPAI H5N1 virus following the two FDA approved pasteurization- as well as various thermization (subpasteurization) conditions widely used by the dairy industry.

Results from our decay studies revealed long-term stability of HPAI H5N1 in raw clinical milk samples collected from HPAI H5N1 infected cows and stored at 4°C (refrigeration temperature), with infectious virus being detected for up to 56 days (8 weeks) in these samples.

These results corroborate findings of a recent study in which infectious HPAI H5N1 was recovered

for up to 5 weeks from milk samples spiked with the virus[12], highlighting the potential public

health risk posed by consumption of raw milk and other raw milk derived products such as raw

milk cheeses which must be aged for 60 days at 4°C[19] prior to human consumption. Survival of

HPAI H5N1 virus in raw milk incubated at 4°C for 56 days with residual titers of about 3.5-3.8 log

$EID_{50}.ml^{-1}$ as shown here (**Extended Data Fig. 1a**), underscore the need for further studies to

determine whether the aging process efficiently inactivates HPAI H5N1 virus in raw milk cheeses.

The decay studies performed here provide important information regarding the sensitivity of HPAI

H5N1 to higher temperatures. Incubation of raw clinical milk from infected cows at 20°C resulted

in inactivation of the virus within 21 days (3 weeks), whereas rapid virus decay was observed at

30°C (6 days, <1 week) and 37°C (2 days) in spiked milk samples. Therefore, thermal treatment

of milk from infected cows to temperatures between 30-37°C, could potentially be utilized to

inactivate HPAI H5N1 virus in raw milk prior to its disposal in affected farms. This would

minimize the risk of environmental contamination and further virus spread.

Although the grade A pasteurized milk ordinance (PMO)[20] requires that milk from sick

animals is segregated on the farm, FDA and USDA found PCR positive shelf milk samples[11],

indicating that the virus found its way into bulk raw milk making it into grocery stores. Importantly,

follow up testing of these samples to determine virus infectivity have shown that no infectious

virus was present in these PCR positive samples[11], which is likely a result of pasteurization. While

early laboratory studies have shown that heat treatment of HPAI H5N1 spiked milk samples to

conditions that mimic FDA approved pasteurization conditions (63°C for 30 min and 72°C for 15

sec) results in drastic reduction in viral titers, residual virus infectivity was still detected when

these samples were subjected to the thermal treatment at 72°C for 15 sec[12,18]. Results here,

however, using raw milk from HPAI infected cows or raw milk spiked with HPAI H5N1 virus,

show that the two FDA approved pasteurization conditions efficiently inactivated the virus

resulting in up to 7.3 log EID_{50} reduction in virus infectivity. These findings corroborate a study

conducted by USDA and FDA which showed that HTST in a lab-scale continuous flow pasteurizer

efficiently inactivated HPAI H5N1 in raw milk[21]. Together, these findings demonstrate that

pasteurization is highly effective in reducing the load of HPAI H5N1 virus in milk.

We further extended our thermal inactivation studies to include four thermization

conditions that are frequently used in dairy industry for production of some cheeses[13,14]. Three out

of four thermization conditions tested including 60°C for 10 min, 63°C for 22 s and 69°C for 22 s

resulted in inactivation of HPAI in raw milk. Thermization of raw milk at 50°C for 10 min,

however, only partially reduced HPAI H5N1 virus titers (0.83-2.42 EID_{50} log reduction). Together

these results demonstrate that HPAI H5N1 is highly sensitive to inactivation in raw milk subjected

to thermal treatment at temperatures on and above 60°C. This was confirmed by our inactivation

kinetics studies. Based on our calculated D-value at 50°C, it would take 50.55 minutes to achieve

a 5-log reduction in HPAI H5N1 loads in contaminated raw milk. In FDA's survey of bulk milk

tanks, they observed and average 3.5 log EID_{50} load, with a range of 1.3 to 6.3 log EID_{50}[21]. A 10°C

increase in the treatment temperature to 60°C resulted in rapid virus inactivation within 5 seconds

of treatment, and while the inactivation was too fast to capture shorter-time intervals, the

approximated D-value is 1.04s (1/ (4.8 log EID_{50}/5)), assuming a linear decay over 5 seconds,

highlighting the efficacy of thermal treatment in ensuring safety of milk and other dairy products

derived from pasteurized milk. Additionally, several of the thermization conditions tested in our

controlled experimental settings proved effective against HPAI H5N1, suggesting the use of

thermized milk treated at 60°C and above for 15-20 seconds to produce cheeses would reduce the

risk of human exposure to products with infectious HPAI H5N1.

In summary, our study provides a comprehensive overview of the thermal inactivation

spectrum of HPAI H5N1 virus in raw milk, demonstrating the efficacy of thermal treatment

including thermization and pasteurization conditions on inactivation of HPAI H5N1 virus in raw

milk.

Materials and Methods

Cells and virus

Bovine uterine epithelial cells (Cal-1, developed in house at the Virology Laboratory at AHDC)

were cultured in minimal essential medium (MEM, Corning Inc., Corning, NY) supplemented with

10% fetal bovine serum (FBS) and penicillin-streptomycin (ThermoFisher Scientific, Waltham,

MA; 10 $U.mL^{-1}$ and 100 $\mu g.mL^{-1}$, respectively). The HPAI H5N1 TX2/24

(A/Cattle/Texas/063224-24-1/2024, genotype B3.13, GISAID accession number:

EPI_ISL_19155861) isolated from the milk of infected dairy cattle in Texas, USA[6] was used in

the decay and pasteurization studies. The virus stock was propagated and titrated in Cal-1 cells.

Biosafety and biosecurity

All work involving handling and propagation of HPAI was performed following strict biosafety measures in AHDC BSL3 laboratories of College of Veterinary Medicine, Cornell University.

Milk samples

Normal raw milk samples were obtained from Teaching Dairy at Cornell University and used as negative control as well as the matrix to be spiked with HPAI H5N1 virus during heat treatment studies. Raw clinical milk samples used in the decay and thermal treatment studies were collected by field veterinarians from a clinically affected dairy farm in Texas, United States[6]. A total of 15 milk samples were pooled and used in decay study (Fig. 1a-b) and heat treatment studies (Fig. 2).

Thermal decay of HPAI H5N1 virus in milk

For thermal decay studies, raw clinical milk samples from HPAI infected dairy cows and/or HPAI-spiked raw milk samples were used. For raw clinical milk samples, 15 IAV M RT-PCR positive milk samples from HPAI infected dairy cows from Texas were pooled and 45 vials of 1 ml aliquots were prepared. The raw clinical milk samples were incubated at 4°C and 20°C. For HPAI-spiked samples, raw normal milk samples obtained from Cornell Teaching Dairy were spiked with HPAI H5N1 TX2/24 (stock titer 2×10^7 log $TCID_{50}$/mL) at 1:20 dilution to achieve a final titer of 6 log $TCID_{50}$/mL and 45 vials of 1 ml aliquots were prepared. The HPAI-spiked milk samples were then incubated at 30°C and 37°C. For all temperatures, sequential samples (n=3 per sampling point) were collected on days 0, 1, 2, 3, 4, 5, 6, 7, 14, 21, 28, 35, 42, 49 and 56 and stored at -80°C for virus quantifications.

Thermal inactivation of HPAI in raw milk using a thermocycler

Initially, the thermal treatment of raw milk to assess inactivation of HPAI H5N1 was performed using a thermocycler. Both pooled raw clinical milk samples from HPAI infected cows and HPAI-spiked raw milk samples were used. For HPAI-spiked samples, raw normal milk obtained from the Cornell Teaching Dairy were spiked with HPAI H5N1 TX2/24 (stock titer 2×10^7 log $TCID_{50}$/mL) at 1:20 ratio to achieve a target titer of 6 log $TCID_{50}$/mL. For heat treatment, we used combinations of six different temperatures and holding times that are used for thermal processing of milk and milk products in dairy industry. We tested two pasteurization conditions, low temperature long

time (LTLT) at 63°C 30 min and high temperature short time (HTST) at 72°C 15 s as well as four

thermization conditions of 50°C for 10 min, 60°C for 10 min, 63°C for 22 sec and 69°C for 22 sec.

For heat treatment, 100 µL/tube of raw clinical milk or HPAI-spiked raw normal milk samples

were added in 8-tube PCR strip. A total of 4 strips (3.2 mL milk) were used per replicate. Following

heat treatment in a thermocycler (BioRad T100 Thermal Cycler), milk samples were chilled

immediately on ice for 10 min. The milk samples were pooled, three aliquots of 1 ml volume were

prepared for each replicate and stored at -80°C until further testing.

Heat treatment of milk using a submerged coil heating system

To more closely simulate pasteurization conditions, we developed a submerged coil heating

system. The stainless-steel coil tubing (FITOK 316/316L Stainless Steel Seamless Tubing, 1/8"

OD × 0.028" Wall, Coil) system has four segments (Fig. 4a) that are furnished with four

temperature sensors (Quick Disconnect Thermocouples with Miniature Connectors, Omega

Engineering Inc., USA) at each segment. The first segment consists of two injection ports for

sample and water and is partly submerged into a water bath. The second and third loops are

completely submerged in water and hold the desired temperature immediately after the sample

injection. The fourth segment is the exit loop for sample collection and is partly submerged into

the water. The thermal sensors are connected to a data logger (HH378, Data Logger/K, J, E, T,

Type Thermometer, Omega Engineering Inc, USA) and temperatures are recorded throughout the

heat treatment process. A circulating water bath (Cole-Parmer EW-16101-89: Unheated Water

Bath, Fisher Scientific) furnished with Immersion Circulators (VWR) was used to heat the coils.

About 15 mL of milk samples were injected into the sample port by using a 20 mL syringe. After

incubation at the desired temperature and time, milk samples (1-5 mL) were pushed out of the coil

system by injecting sterile water in the system. Each milk sample was collected into a sterile tube.

Immediately after collection, the milk samples were chilled on ice for 10 min, aliquoted and stored

at -80°C. After each round, the coil was cleaned by flushing the coil with a 1.5% w/v solution of

alkaline detergent (Alcanox, Sigma-Aldrich) followed by 1% w/v solution of acid detergent

(Citranox, Alconox) and finally passing several coil-full volumes of sterile water and by passing

the air several times.

RNA extraction and RT-PCR

Viral RNA from milk samples was extracted using the IndiMag Pathogen kit (INDICAL
Bioscience) on the IndiMag 48s automated nucleic acid extractor (INDICAL Bioscience, Leipzig,
Germany), and the rRT-PCR was performed using the Path-ID™ Multiplex One-Step RT-PCR Kit
(Thermo Fisher, Waltham, MA, USA) and primers and probes targeting the M gene under the
following conditions: 15 min at 48ºC, 10 min at 95 °C, 40 cycles of 15 s at 95 °C for denaturation
and 60 s at 60 °C. A standard curve was prepared using RNA extracted from HPAI H5N1 TX2/24
spiked milk samples. Serial 10-fold dilutions of the stock virus ($2x10^7$ log $TCID_{50}$/mL) were
prepared in raw normal milk samples for RNA extraction followed by RT-PCR as described above.
The Ct values were used to estimate the viral RNA copy number in the tested milk samples using
relative quantification method.

Virus titration in cell culture ($TCID_{50}$)

The infectious viral loads in milk samples were quantified by viral titrations in Cal-1 cells. For
this, serial 10-fold dilutions of milk samples were prepared in MEM and inoculated into Cal-1
cells in 96-well plates. Each dilution was inoculated in quadruplicate wells. At 48h post-
inoculation, culture supernatant was aspirated, and cells were fixed with 3.7% formaldehyde
solution for 30 min at RT and subjected to IFA using the anti-NP (HB65) mouse monoclonal
antibody, followed by secondary anti-mouse-Alexa-594 incubation. The 50% tissue culture
infectious dose ($TCID_{50}$) was determined using end-point dilutions and the Spearman and Karber's
method and expressed as $TCID_{50}.mL^{-1}$.

Virus titration in embryonated chicken eggs (EID_{50})

For virus titration in eggs, 9-days-old embryonated chicken eggs were used. Serial 10-fold
dilutions of milk samples were prepared in phosphate buffered saline (PBS) supplemented with
antibiotic-antimycotic (anti-anti 100X, Thermo Fisher Scientific, USA) at 2:1 ratio. The eggs were
candled to mark the air sac on the shell, sanitized with 70% ethanol and a hole was drilled in the
eggshell. One hundred µL of the diluted milk sample was injected in quadruplicate into the chorio-
allantoic sac route and sealed with glue. Eggs were candled daily for four days, and dead embryos
were chilled overnight before collection. After 4 days post-inoculation, all surviving embryos were
chilled for 24 hours, and allantoic fluids were collected. All allantoic fluids were tested by
hemagglutination assay (HA) using 0.5% chicken RBC. Finally, the 50% egg infectious dose
(EID_{50}) was calculated by the Reed and Muench method.

Hemagglutination assay (HA)

For the HA, 100 µL allantoic fluid was taken into the wells of 1st row of a 96-well U-bottom plate. 50 µL of PBS was added to the wells of 2nd, 3rd and 4th rows. Three 2-fold dilutions (1:2, 1:4 and 1:8) of the allantoic fluids were prepared in PBS. Then, 50 µL of 0.5% chicken red blood cells (RBC) were added to each well and incubated at room temperature for 30 min. Lack of RBC button formation indicates positive HA reaction.

Statistical analysis

Graphs were prepared using GraphPad Prism 10 software. Linear regression analysis was performed to obtain the thermal death time D-values.

Acknowledgements

The work was funded by the New York State Department of Agriculture and Markets (award no. CM04068HM). The authors would like to thank the Cornell EH&S and Biosafety teams for their help in setting up the protocols and procedures to conduct work with HPAI H5N1 virus at the Cornell BSL-3 facilities and the Cornell Teaching Dairy for proving the raw milk samples. The authors would further like to acknowledge the efforts of Alan Bitar for fabrication of the submerged coil system for use in this study.

Author contributions

Conceptualization: DGD; Methodology: MN, LMC, KK, RI, NHM, SDA, DGD; Software: MN, KK; Validation: MN, LMC, KK; Formal analysis: MN, LMC, KK, RI, DGD; Investigation: MN, LMC; Resources: NHM, SDA, DGD; Data Curation: MN, LMC, KK; Writing - Original Draft: MN, DGD; Writing - Review & Editing: MN, LMC, KK, NHM, RI, SDA, DGD; Visualization: MN; Supervision: DGD; Project administration: DGD, Funding acquisition: NHM, SDA, DGD.

References

1. USDA. 2022-2023 Confirmations of Highly Pathogenic Avian Influenza in Commercial and Backyard Flocks. *HPAI 2022/2023 Confirmed Detections* Preprint at https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/avian/avian-influenza/hpai-2022/2022-hpai-commercial-backyard-flocks (2023).

2. APHIS-USDA. *Detections of Highly Pathogenic Avian Influenza in Mammals*. (2024).

3. Elsmo, E. J. *et al.* Highly Pathogenic Avian Influenza A(H5N1) Virus Clade 2.3.4.4b Infections in Wild Terrestrial Mammals, United States, 2022. *Emerg Infect Dis* **29**, (2023).

4. AVMA. *Goat in Minnesota Tests Positive for HPAI. Available at: Https://Www.Avma.Org/News/Goat-Minnesota-Tests-Positive-Hpai.* (2024).

5. Burrough, E. R. *et al.* Highly Pathogenic Avian Influenza A(H5N1) Clade 2.3.4.4b Virus Infection in Domestic Dairy Cattle and Cats, United States, 2024. *Emerg Infect Dis* **30**, (2024).

6. Caserta, L. C. *et al.* Spillover of highly pathogenic avian influenza H5N1 virus to dairy cattle. *Nature* (2024) doi:10.1038/s41586-024-07849-4.

7. Halwe, N. J. *et al.* Outcome of H5N1 clade 2.3.4.4b virus infection in calves and lactating cows. *bioRxiv* 2024.08.09.607272 (2024) doi:10.1101/2024.08.09.607272.

8. Centers for Disease Control and Prevention. H5 Bird Flu: Current Situation. (2024).

9. Dash, K. K. *et al.* A comprehensive review on heat treatments and related impact on the quality and microbial safety of milk and milk-based products. *Food Chemistry Advances* **1**, 100041 (2022).

10.   FDA. *Updates on Highly Pathogenic Avian Influenza (HPAI). Available at: Https://Www.Fda.Gov/Food/Alerts-Advisories-Safety-Information/Updates-Highly-Pathogenic-Avian-Influenza-Hpai*. (2024).

11.   Spackman, E. *et al.* Characterization of highly pathogenic avian influenza virus in retail dairy products in the US. *J Virol* **98**, (2024).

12.   Guan, L. *et al.* Cow's Milk Containing Avian Influenza A(H5N1) Virus — Heat Inactivation and Infectivity in Mice. *New England Journal of Medicine* **391**, 87–90 (2024).

13.   Deeth, H. C. Heat Treatment of Milk: Pasteurization (HTST) and thermization (LTLT). in *Encyclopedia of Dairy Sciences* 645–654 (Elsevier, 2022). doi:10.1016/B978-0-12-818766-1.00133-1.

14.   Deeth, H. Optimum Thermal Processing for Extended Shelf-Life (ESL) Milk. *Foods* **6**, 102 (2017).

15.   Nelli, R. K. *et al.* Sialic Acid Receptor Specificity in Mammary Gland of Dairy Cattle Infected with Highly Pathogenic Avian Influenza A(H5N1) Virus. *Emerg Infect Dis* **30**, (2024).

16.   Pitino, M. A., O'Connor, D. L., McGeer, A. J. & Unger, S. The impact of thermal pasteurization on viral load and detectable live viruses in human milk and other matrices: a rapid review. *Applied Physiology, Nutrition, and Metabolism* **46**, 10–26 (2021).

17.   Kwon, T. *et al.* Bovine Highly Pathogenic Avian Influenza Virus Stability and Inactivation in the Milk Byproduct Lactose. *Viruses* **16**, 1451 (2024).

18.   Kaiser, F. *et al.* Inactivation of Avian Influenza A(H5N1) Virus in Raw Milk at 63°C and 72°C. *New England Journal of Medicine* **391**, 90–92 (2024).

19.   21 CFR Part 133. Cheeses and Related Cheese Products. *CFR* (2024).

20.   USPHS/FDA. *Grade A Pasteurized Milk Ordinance (PMO)*. (2023).

21.   Spackman, E. *et al.* Inactivation of highly pathogenic avian influenza virus with high

temperature short time continuous flow pasteurization and virus detection in bulk milk

tanks. *medRxiv* (2024) doi:10.1101/2024.07.01.24309766.

Extended Data Table 1. Isolation of HPAI H5N1 virus in raw clinical milk collected from HPAI infected cows and HPAI-spiked raw normal milk incubated at different temperatures by using embryonated chicken eggs.

	Cal-1	Chicken embryos				Cal-1	Chicken embryos		
4°C D49		Dilutions	Death	HA (+)	**4°C D56**		Dilutions	Death	HA (+)
R1	Toxicity	1:10	2/3	2/3	R1	Toxicity	1:10	3/3	3/3
		1:10^2	1/3	1/3			1:10^2	3/3	3/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
R2	Toxicity	1:10	2/3	2/3	R2	Toxicity	1:10	3/3	3/3
		1:10^2	1/3	1/3			1:10^2	3/3	3/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
R3	Toxicity	1:10	3/3	3/3	R3	Toxicity	1:10	3/3	3/3
		1:10^2	3/3	3/3			1:10^2	3/3	3/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
20°C D14					**20°C D21**				
R1	Toxicity	1:10	1/3	1/3	R1	Toxicity	1:10	1/3	0/3
		1:10^2	0/3	0/3			1:10^2	0/3	0/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
R2	Toxicity	1:10	1/3	0/3	R2	Toxicity	1:10	2/3	0/3
		1:10^2	0/3	0/3			1:10^2	0/3	0/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
R3	Toxicity	1:10	1/3	0/3	R3	Toxicity	1:10	1/3	0/3
		1:10^2	0/3	0/3			1:10^2	0/3	0/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
30°C D6					**30°C D7**				
R1	Toxicity	1:10	0/3	0/3	R1	Toxicity	1:10	0/3	0/3
		1:10^2	0/3	0/3			1:10^2	0/3	0/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
R2	Toxicity	1:10	0/3	0/3	R2	Toxicity	1:10	0/3	0/3
		1:10^2	0/3	0/3			1:10^2	0/3	0/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
R3	Toxicity	1:10	0/3	0/3	R3	Toxicity	1:10	0/3	0/3
		1:10^2	0/3	0/3			1:10^2	0/3	0/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
37°C D2					**37°C D3**				
R1	Toxicity	1:10	0/3	0/3	R1	Toxicity	1:10	0/3	0/3
		1:10^2	0/3	0/3			1:10^2	0/3	0/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
R2	Toxicity	1:10	0/3	0/3	R2	Toxicity	1:10	0/3	0/3
		1:10^2	0/3	0/3			1:10^2	0/3	0/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3
R3	Toxicity	1:10	0/3	0/3	R3	Toxicity	1:10	0/3	0/3
		1:10^2	0/3	0/3			1:10^2	0/3	0/3
		1:10^3	0/3	0/3			1:10^3	0/3	0/3

Note: D, days; R, replicates; HA, hemagglutination assay; +, positive

Fig. 1. Kinetics HPAI H5N1 virus decay in milk. Decay of HPAI H5N1 virus in raw clinical milk
collected from HPAI infected cows incubated at 4°C (**a**) and 20°C (**b**) for 56 days. Decay of HPAI
H5N1 virus in HPAI-spiked (1:20 ratio) raw normal milk and incubated at 30°C (**c**) and 37°C (**d**)
for 56 days. The virus titers were determined by end point titrations in Cal-1 cells (Log $TCID_{50}.mL^{-1}$
^{1}). Data represents observations (dots) from three replicates (n=3) overlayed with the mean
(horizontal line) and ± SEM (whiskers).
8

Fig. 2. Effect of low temperature thermization and FDA approved pasteurization conditions on HPAI H5N1 infectivity in raw milk. **a-d.** Thermal inactivation of HPAI H5N1 virus in raw clinical milk samples collected from HPAI infected cows. **a.** Immunofluorescence (IFA) images of Cal-1 cells inoculated with control (normal and clinical) or heat-treated clinical milk samples. Raw clinical milk samples were collected from HPAI H5N1 infected cows and heat treated at the indicated temperatures and times utilizing a thermocycler. **b.** HPAI H5N1 viral RNA loads in control or heat-treated raw clinical milk samples as determined by qRT-PCR (n = 3). **c.** HPAI H5N1 virus titrations in Cal-1 cells (n = 3). **d.** HPAI H5N1 virus titrations in embryonated chicken eggs

(n = 3). **e-h.** Thermal inactivation of HPAI H5N1 virus in raw normal milk spiked (1:10 ratio) with
milk samples collected from HPAI infected cows using a thermocycler. **e.** IFA images of Cal-1
cells inoculated with control (normal and clinical) or heat-treated spiked milk samples. **f.** HPAI
H5N1 viral RNA loads in control or heat-treated raw spiked milk samples as determined by qRT-
PCR (n = 3). **g.** HPAI H5N1 virus titrations in Cal-1 cells (n = 3). **h.** HPAI H5N1 virus titrations
in embryonated chicken eggs (n = 3). Red fluorescence indicates detection of the viral NP protein.
Images are representative of three independent experiments (n = 3). The virus titers were
determined by end point dilutions in Cal-1 cells and in embryonated chicken eggs and expressed
as tissue culture infectious dose 50 per mL ($TCID_{50}.mL^{-1}$) and egg infectious dose 50 per ml
($EID_{50}.mL^{-1}$), respectively. Data represents observations (dots) from three independent
experiments (n=3) overlayed with the mean (horizontal line) and ± SEM (whiskers).
29

Fig. 3. Evaluation of thermization conditions on inactivation of HPAI H5N1 virus in raw milk. Thermal inactivation of HPAI H5N1 virus in raw normal milk spiked (1:20 ratio) with HPAI H5N1 using a thermocycler. **a.** IFA images of Cal-1 cells inoculated with control or heat-treated HPAI-spiked raw normal milk samples. **b.** HPAI H5N1 viral RNA loads in control or heat-treated HPAI-spiked raw milk samples as determined by qRT-PCR (n = 3). **c.** HPAI H5N1 virus titrations in Cal-1 cells (n = 3). **d.** HPAI H5N1 virus titrations in embryonated chicken eggs (n = 3). Red fluorescence indicates detection of the viral NP protein. Images are representative of three independent experiments (n = 3). The virus titers were determined by end point dilutions in Cal-1 cells and in embryonated chicken eggs and expressed as tissue culture infectious dose 50 per mL ($TCID_{50}.mL^{-1}$) and egg infectious dose 50 per ml ($EID_{50}.mL^{-1}$), respectively. Data represents observations (dots) from three independent experiments (n=3) overlayed with the mean (horizontal line) and ± SEM (whiskers).

Fig. 4. Thermal inactivation of HPAI H5N1 in virus-spiked milk samples using a submerged coil heating system. **a**. Schematic representation of the submerged coil heating system. **b.** Representative plot showing temperature changes over time in milk samples injected in the submerged coil system during heat treatment at 72°C for 15 sec. Data are presented as the mean ± SEM (whiskers) of 3 independent experiments and a line connecting the means. **c-f.** Thermal inactivation of HPAI H5N1 virus in raw normal milk spiked (1:20 ratio) with HPAI H5N1 using a submerged coil heating system. **c.** IFA images of Cal-1 inoculated with control or heat-treated HPAI-spiked raw normal milk samples. **d.** HPAI H5N1 viral RNA loads in control or heat-treated HPAI-spiked raw milk samples as determined by qRT-PCR (n = 3). **e.** HPAI H5N1 virus titrations in Cal-1 cells (n = 3). **f.** HPAI H5N1 virus titrations in embryonated chicken eggs (n = 3). Red fluorescence indicates detection of the viral NP protein. Images are representative of three

55   independent experiments (n = 3). The virus titers were determined by end point dilutions in Cal-1
56   cells and in embryonated chicken eggs and expressed as tissue culture infectious dose 50 per mL
57   ($TCID_{50}.mL^{-1}$) and egg infectious dose 50 per ml ($EID_{50}.mL^{-1}$), respectively. Data represents
58   observations (dots) from three independent experiments (n=3) overlayed with the mean (horizontal
59   line) and ± SEM (whiskers).
60

Fig. 5. Kinetics of HPAI H5N1 virus inactivation in milk using submerged coil heating system.
Raw normal milk was spiked with HPAI H5N1 at 1:20 ratio and heat treated at 50°C (**a-c**), 60°C
(**d-f**), 63°C (**g-i**) and 72°C (**j-l**) with a submerged coil heating system. Sequential samples were
collected at the indicated time points and analyzed by qRT-PCR and virus titration in cell culture
and embryonated chicken eggs. The virus titers were determined by end point titration in Cal-1

67   cells ($TCID_{50}.mL^{-1}$) and embryonated chicken eggs ($EID_{50}.mL^{-1}$). Data represents observations
68   (dots) from three independent experiments (n=3) overlayed with the mean (horizontal line) and $\pm$
69   SEM (whiskers).
70

Extended Data Fig. 1. Line graph showing temperature changes over time in injected milk samples during heat treatment in the submerged coil heating apparatus at 50°C for 10 min (a), 60°C for 10 min (b), 63°C for 22 s (c), 63°C for 30 min (d) and 69°C for 22 s (e). Data are presented as mean ± SEM of 3 independent experiments.