Pollinator visitation pattern and differential reproductive compatibilities

contribute to the unequal morph abundances in a nocturnal ginger from Western

Ghats, India

Saket Shrotri[1] and Vinita Gowda[1]

[1]Tropical Ecology and Evolution (TrEE) Lab, Department of Biological Sciences, Indian

Institute of Science Education and Research (IISER) Bhopal, Bhopal, Madhya Pradesh, India,

462066.

Corresponding author: Vinita Gowda.

Email: gowdav@iiserb.ac.in

ORCiD:

Saket Shrotri: 0000-0002-4596-9451

Vinita Gowda: 0000-0001-8533-0014

Summary

• Polymorphism, the presence of discrete floral morphs within a population, is often

maintained by pollinator-mediated or frequency-dependent selection. While

pollinators often act on visual displays, this study investigates hawkmoth-mediated

selection in the nocturnal *Curcuma caulina* (Zingiberaceae), where visual cues play a

minimal role. We explore the impact of non-visual traits, such as morph abundance,

nectar rewards, and reproductive compatibility, on the maintenance of bract color

polymorphism.

• We measured floral traits, nectar rewards, and abundance of morphs in the natural

population, and quantified reproductive compatibility using hand pollination

treatments. We next used path analysis to test the effect of morph abundance on

reproductive fitness via floral and ecological traits.

• We identified tri-modal morph abundance with two common morphs and one rare

morph. Morphs differed in nectar traits and reproductive compatibility but not in

floral morphologies. Rare morphs had high-energy nectar, higher pollinator revisits

and self-pollination, and lower seed output, while common morphs had low-energy

nectar, lower revisits, higher outcrossing and fitness.

31 •  Morph abundance in *C. caulina* is shaped by an interplay between nectar rewards,

32  pollinator attraction, and compatibility. We conclude that in perennial rhizomatic

33  species understanding historical selection regimes and both abiotic and biotic factors

34  are necessary to characterize floral polymorphism.

35 **Keywords**

36 Floral color polymorphism, Frequency-dependent selection, Hawkmoth pollination, Path

37 analysis, Reproductive fitness, Self-incompatibility

38 **Introduction**

39  In plants, floral color polymorphism (henceforth referred to as FCP) is a common trait

40 that is often under pollinator-mediated selection. The emergence of FCP in plants has been

41 attributed to two factors - mutation events (Kellenberger *et al.*, 2019) and genetic

42 recombination (Stebbins, 2019). However, its maintenance in a population is often explained

43 by spatiotemporal variation in both biotic and abiotic selection regimes such as balancing

44 selection, frequency-dependent selection, selection of pleiotropic traits, mating strategies,

45 heterozygous advantage, and neutral or stochastic processes (Narbona *et al.*, 2018; Vaidya *et*

46 *al.*, 2018; Sapir *et al.*, 2021).

47  In plants with FCP, some of the key floral characters that are variable and are under

48 positive selection by the pollinators are floral color (Waser & Price, 1981, 1985; Jones &

49 Reithel, 2001; Gigord *et al.*, 2001; Dormont *et al.*, 2010; Ortiz *et al.*, 2015), flower shape or

50 size (GÓmez, 2000; Majetic *et al.*, 2007, 2009; Barazani *et al.*, 2019), nectar characteristics

51 (Melendez-Ackerman & Campbell, 1998; Carlson & Holsinger, 2013), number of open

52 flowers (Dormont et al., 2010; M. Eckhart et al., 2006), and flowering time (Brunet *et al.*,

53 2015). Almost all of these studies note that pollinators show selective preference for specific

54 traits that are associated with floral color, hence influencing selection differentials among the

55 floral color morphs. For example, in *Lobularia maritima*, pollinators show preferential

56 visitations towards one of the morphs, however, it is not the floral color, but floral display

57 size associated with the white morph that is the target of pollinator-mediated selection

58 (Gomez, 2000). Similarly, traits such as plant height and flower production in *Hesperis*

59 *matronalis* (Majetic et al., 2007; 2009), and floral nectar and flower number in *Protea aurea*

60 (Carlson and Holsinger, 2013) have been found to be associated with floral color and targets

61 for pollinator-mediated selection of FCP.

In the above-mentioned studies, selection is measured on the trait in an individual and
morph abundance is often the outcome of selection processes. However, despite the
recognition that the relative abundance of morphs within a population can influence selection,
morph abundance as a critical structural feature of a polymorphic species has rarely been
investigated. Narbona *et al.* (2018) reported that nearly all of the Mediterranean species with
FCP have unequal morph frequencies at the population-level, while some species exhibited
temporal fluctuations in morph frequencies (Dormont *et al.*, 2010; Tang & Huang, 2010;
Tang *et al*., 2016). Morph frequency influences pollinator discrimination (Smithson and
Macnair, 1996; Eckhart et al., 2006; Dormont *et al.*, 2010; Ortiz *et al.*, 2015; Wenzell *et al.*,
2023) and can maintain FCP via frequency-dependent selection (FDS) through disassortative
mating (i.e., negative frequency-dependent visitation; Gigord et al. 2001; Gegear and Laverty
2005; Benadi and Gegear 2018) and adaptive foraging influencing assortative mating
(positive frequency-dependent visitation; Jones and Reithel, 2001; Eckhart et al., 2006).
Therefore, morph abundances within a population can modulate not only pollinator visitation
rates (Dormont *et al.*, 2010; Ortiz *et al.*, 2015; Wenzell *et al.*, 2023), but also outcrossing
rates (Fry & Rauher, 1997; Subramaniam & Rauher, 2000; Holsinger, 2000), and ultimately
the reproductive fitness of the morphs (Gigord *et al.*, 2001; Lau & Galloway, 2004; Koski *et
al*., 2020).

In contrast, when FCPs are not driven by pollinator-mediated selection, the floral
color morphs are either associated with differential reproductive compatibilities, such as in
geophytes. Here polymorphic forms within a population can be maintained by features such
as clonal reproduction. For example, in *Iris lutescens* a novel floral morph has been shown to
gain a selective advantage because of its higher clonality (Thompson & Eckert, 2004) even
when the pollinators did not favour it (Imbert *et al.*, 2014). Further, rare morph may be
maintained in a population if it is self-compatible and autogamous, despite it being under
pollination limitation as observed in *Ipomoea purpurea* (Fry & Rauher, 1997; Subramaniam
& Rauher, 2000) and in *Lysimachia arvensis* (Arista *et al.*, 2013; Ortiz *et al.*, 2015). Thus,
inter-morph differences in physiological compatibilities can play a critical role in the
maintenance of rare morphs within a population, through fitness differences among the
morphs.

Within the Asian tropics, several genera within the family Zingiberaceae (gingers) are
known to have a diversity of floral polymorphisms ranging from stylar polymorphism to
color polymorphisms. Gingers are a dominant, perennial understory herb with clonal

reproduction, and both diurnal as well as nocturnal flowering. While stylar polymorphism is

common in the genera *Alpinia* and *Amomum* (Raju *et al.*, 2004; Takano *et al.*, 2009; Ren *et*

al., 2022), FCP is common across several genera in the family Zingiberaceae such as

Achasma, *Aframomum*, *Boesenbergia*, *Curcuma*, *Globba*, *Hedychium* and *Roscoea* (Poulsen

& Lock, 1997; Harris *et al.*, 2000; Leong-Skornicková *et al.*, 2007; Mood *et al.*, 2013;

Yadav, 2024; Yadav and Gowda, 2024). In genera like *Alpinia* and *Roscoea*, floral

polymorphism, regardless of the type, has been shown to be critical for plant-pollinator

interactions and speciation (Takano *et al.*, 2009; Paudel *et al.*, 2019). Among the

polymorphic gingers, the taxonomically diverse genus *Curcuma* is known for its floral color

variation (Leong-Skornicková *et al.*, 2007), including the nocturnal species *C. caulina* (Fig.

S1).

In the tropics, nocturnal plant species are often pollinated by crepuscular and

nocturnal pollinators such as long-tongued hawkmoths (Johnson & Raguso, 2016; Hahn &

Brühl, 2016; Johnson *et al.*, 2017; Liu & Sun, 2019), however very little is known about their

floral preferences, ability to discriminate floral traits, especially floral colors, and pollination

efficiency (Weiss, 2001; Goyret *et al.*, 2008). In a nocturnal ginger *C. caulina*, we noted both

the presence of FCP and unequal morph abundance within a population in Western Ghats,

India. In this study, we tested whether floral trait differences, pollinator visitation differences,

and reproductive differences suggest that the morphs are under pollinator-mediated and

frequency-dependent selection. To the best of our knowledge, the maintenance of FCP has

not been studied in any exclusively nocturnally flowering plant species and we address the

following questions:

1. Do morphs share similar floral traits and outcrossing rates?

2. Does morph abundance affect pollinator visitations and reproductive fitness?

3. How are rare morphs maintained in a polymorphic population?

To explain the unequal morph abundance at the population level, we finally tested the

significance of the four traits - plant abundance, nectar quality, pollinator visitation rate, and

selfing/outcrossing rates, on the reproductive fitness of the morphs using path analysis.

Materials and Methods

Study species and study site

Curcuma caulina (Zingiberaceae) is a rhizomatous, mass-flowering, perennial herb

that flowers between July to October (Fig. S1). The plant is 30–50 cm tall with a single shoot

bearing a terminal inflorescence which consists of 10-18 fertile 'lateral bracts' that bear

flowers (Fig. S1). Flowers are born in cincinni (1–3 flowers per cincinnus), are nocturnal

(anthesis at 18:00), and last 12–14 hours only. The stigma is funnel-shaped and projects

beyond the anther, and fruits mature within the bracts. *C. caulina* is endemic to the northern

Western Ghats of India and this study was carried out on Kaas Plateau (कास पठार;

17°43'12.36" N, 73°49'28.0956" E, elevation 1225m, Fig. S1), Satara, Maharashtra, India

between 2019 and 2022. Kaas Plateau is an elevated ferricrete outcrop (red laterite crust) and

a UNESCO Natural World Heritage site recognized for its floral diversity and high

endemicity (Watve, 2013).

Bract color polymorphism and variation in floral morphology and nectar rewards

color of the lateral bracts of *C. caulina* individuals varied within the population from

predominantly green to predominantly red colored inflorescence (Fig. S2a). To determine the

frequency of these variants, we used randomly placed ten transects, each measuring 10 x 4

m². Within each transect, frequency of all bract color variants were identified and counted.

The extent of the reddish-pink coloration (0% to 100%) on the first or basal lateral bract was

used to classify the variants and this was also measured using handheld spectrophotometer

(Ocean Optics; Fig. S2a, b). This method identified six variants with continuous bract color

variation within the population (Fig. S2c). However, we identified only three variants (0%,

60%, and 100%) as morphs based on Ford's criteria for polymorphic forms (Ford, 1945; Fig.

S2). These three morphs were named based on the coloration of their lateral and coma bracts:

the 0% variant as Green-White (GW), the 60% variant as Green-Red (GR), and the 100%

variant as Red-White (RW; Supp Fig. 1a). The distribution of these morphs was unequal,

with GR morphs being the most common (~ 44% occurrence), followed by GW (~ 29%) and

RW (13%). The remaining three bract color variants (15%, 30%, and 45%) were extremely

rare, each constituting less than 5% of the population (Fig. S2c).

To morphologically characterise the three morphs, we measured a total of 22

vegetative and floral characters using digital vernier calliper and rulers (n=30 per morph;

character list in Table S1). To assess morphological variations among the morphs, we ran a

non-metric multidimensional scaling (nMDS) analysis with Bray–Curtis dissimilarity using

the VEGAN package (Dixon, 2003) in R (4.3.0; R core Team, 2021). We used the first three

dimensions, to construct three-dimensional (3D) plots, which were used to visualise both

intra-morph and inter-morph variations. The statistical difference among the overlapping

clusters (morphs) was measured using the ANOSIM test, where the Bray–Curtis dissimilarity

matrix with 1000 permutations was used (VEGAN package; Dixon, 2003).

We measured nectar volume and concentration in 106 bagged flowers (32 to 38

unopened buds per morph) between 18:00 and 18:30 hrs by destructive sampling. The nectar

volume (μl), concentration (% sucrose), and nectar energy (cal) were compared among the

three morphs. Nectar volume was measured using calibrated glass microcapillary tubes

(Drummond Scientific Company), and the sucrose concentration in nectar was determined

using a handheld refractometer (ERMA BRIX 0~55%). The brix sucrose % value was then

converted into sugar (mg) per 1 ml of nectar using the table given in the CRC Handbook of

Chemistry and Physics (Lide, 2004) and as outlined by Bolten *et al.* (1979). This value was

used to calculate the amount of sugar in a given volume of nectar, and it was converted into

calories (1 mg of sucrose = 4 calories as given in Heinrich, 1975). Differences between the

morphs were tested using the Kruskal-Wallis test followed by Dunn's test as a post-hoc; the

significance was set at $p<0.05$; analysis was carried out in R (4.3.0; R Core Team, 2021).

Natural pollinator visitations

To determine if pollinators exhibit differential visitation towards a particular morph,

pollinator visits were monitored by night vision CCTV cameras (Ae Zone CCTV IR

day/night vision USB camera) between 18:00 hrs and 06:00 hrs. These observations were

carried out in 18 randomly selected observation plots on the plateau, where each plot

consisted of three to six individuals. All morphs were observed for a total of 648 hrs (i.e., 216

hrs per morph). Cameras were positioned approximately 1 to 1.5 m away from the focal

plants such that all inflorescences were visible in their entirety in the field of view. The

recordings were manually scored, where the total number of visits by each pollinator and the

total number of flowers present in the plot were measured. We calculated the pollinator

visitation rate as the total number of visits per flower per hour for each morph. The visitation

rates were summed for every hour between 18:00 to 06:00 hrs in order to record the temporal

shift in pollinator visitation rate. To compute the visitation rate to each morph within the

entire population, we simulated visitation rates for each morph using observed data in a

Monte Carlo simulation in R. We computed the visitation rate for each morph at the

population level as follows: a) We generated 10,000 values (with replacement) for the two

variables - pollinator visitation rates (PVR) and morph abundance using sampling with

replacement from the observed data set (98 values for PVR and 30 values for morph

abundance). b) We next calculated the product of the two variables (PVR x morph

abundance) and computed the mean and variance for this simulated data set (Fig. 1c, Z axis).

This value represents the expected PVR for each morph in the population, given the recorded

plot-level variation in morph abundance in the study area.

Natural pollinator visitation rates (number of visits per flower per hour) were

compared among the morphs on a temporal scale using the circular statistical analysis in

ORIANA (v. 4.02 Kovach Computing Services; Kovach, 2011). Temporal differences in

visitation rates between the morphs were tested using the Mardia-Watson Wheeler test (or

uniform score test; Zar, 1999). Since the pollinator observation plots were not always located

within the morph-census transects, the pollinator visitation rates were standardized for morph

frequency using Monte Carlo Simulation (n = 10,000). The morph frequency and pollinator

visitation rates for each morph were utilized to estimate the distribution of their product for

the respective morphs. To test for inter-morph differences in pollinator visitation rates we

used the peak visitation rates from 19:00 hrs to 22:00 hrs, which accounted for ~85% of all

pollinator visits. We used the Kruskal-Wallis test followed by Dunn's test as a post-hoc test

at $p<0.05$. All statistical analyses were carried out in R (4.3.0; R Core Team, 2021).

Physiological compatibility and natural fruit set

To comprehensively evaluate the physiological self-compatibility and cross-

compatibility rates in the three floral color morphs, we carried out a total of 15 hand-

pollination treatments. Each morph was treated separately, and the pollination treatments

included three autogamy, three self-pollination, and nine cross-pollinations that included

three intra-morph cross-pollinations and six inter-morph cross-pollinations. Thus, a total of

471 hand pollinations were carried out (with n ≥ 30 per treatment) where the three intra-

morph cross-pollination treatments were: GW x GW, GR x GR, and RW x RW (male x

female morph) and will be interchangeably referred to as 'cross-pollination' treatments. Inter-

morph cross-pollination treatments were categorized into three pairs to investigate the effect

of male and female parents in each treatment. These crosses are (male x female) identified as

pair a: GW x GR and GR x GW, pair b: GW x RW and RW x GW, and pair c: GR x RW and

RW x GR.

Autogamy treatment included unmanipulated, bagged flowers that were checked for

fruit set after four weeks. For cross-pollination treatments, flowers were emasculated and

mixed pollen grains from 10-15 individuals were manually transferred on marked flowers.

Flowers were observed for fruit set in ~4 weeks and percent success rate in reproduction was

measured for each treatment using the number of fruits per treatment as well as the number of

seeds per fruit (seed count). We statistically tested the reproductive success rates between

self- and cross-pollination treatments for each morph using chi-square tests. Similar

comparative analyses were also carried out for the seed counts from self- and cross-

pollination treatments using the Kruskal-Wallis test followed by Dunn's post-hoc test with

significance set at $p < 0.05$.

We quantified the reproductive success of natural pollination events within the study

population at the end of the flowering season in mid-October by measuring fruit set in ≥ 500

individuals per morph and seed counts in ≥ 50 per morph. For each inflorescence, we

estimated the total number of flowers per inflorescence by multiplying the number of lateral

bracts with the number of flowers estimated in a cincinnus. Fruit-set per inflorescence was

then calculated as the ratio of the number of matured fruits to the total number of flowers

estimated for a particular inflorescence. We used Kruskal-Wallis test to compare both natural

fruit set and natural seed count among morphs. Dunn's post-hoc test was used with a

significance set at $p < 0.05$.

Path analysis to test traits effects on the reproductive fitness of the morphs

We used path analysis to understand the direct and indirect effects of abundance of a morph

on its natural fitness as described by Mitchell (1993). Path analysis was tested for the entire

Kaas population of *C. caulina* as well as for each morph independently (n = 300 individuals;

i.e., 100 per morph) in order to assess the net effect and important selection regimes within

the population and within each morph. Three indirect effects included in the path analyses

were- total nectar reward, pollinator visitation rates, and physiological compatibilities. Values

for the total nectar reward was normalized with morph abundance and of natural fitness with

seed-set using Monte Carlo simulations (n = 10,000). Path analysis was performed in R

(4.3.0; R Core Team, 2021) using the *lavaan* package (Rosseel, 2012) and path estimates

were computed using 10,000 iterations in the bootstrapping procedure. Estimates of model fit

were computed using Chi-squared goodness of fit, Comparative Fit Index (CFI) and Tucker-

Lewis Index (TLI). A nonsignificant chi-squared value indicates that a model is not

significantly different from the observed correlations in the data and is therefore a good fit

whereas CFI and TLI indices above 0.9 are considered to be a good fit (Mitchell, 2001).

Finally, we tested the hypothetical relationship between reproductive fitness and relative

abundance using a regression model which was not part of the main path analysis.

Results

Floral (bract) color polymorphism and morphological variability

The abundance of the three morphs differed within the population. The GR morph (98.6 ± 7.2

individuals; ~44 %) and GW morph (75.7 ± 7.2 individuals; ~29 %) were the most common

morphs; their frequencies were not significantly different from each other (Kruskal-Wallis $\chi 2$

= - 0.692, df = 1, p = 0.2446). The rarest of the three morphs was the Red-White morph (RW)

with 24.2 ± 2.0 individuals (~13 %) in the population, and the RW morph had significantly

fewer individuals when compared to both GR and GW (RW vs GR, Kruskal-Wallis $\chi 2$ = -

2.427, df = 1, p = 0.0076 and RW vs GW, Kruskal-Wallis $\chi 2$ = 1.736, df = 1, p = 0.0413

respectively). All other bract color variants totaled to ~14% in the population and thus were

extremely rare. The non-metric multidimensional scaling (nMDS) analysis of the 22 floral

and vegetative morphometric characters (Table S1; Fig. S2) failed to form distinct clusters

separating the morphs by their morphological traits. Further, the Analysis of Similarities

(ANOSIM) test (p = 0.4835) and the associated ANOSIM R-value (ANOSIM R = - 0.0038)

also indicated no statistical significance in the differentiation of these clusters for each

morph. Since we detected polymorphism in bract colors and not floral colors, henceforth, the

term color polymorphism refers to only bract color polymorphism in *C. caulina*.

Variability in nectar traits and hawkmoth visitation rates

Nectar is the primary pollinator reward in *C. caulina*, and there was no significant

difference between the total nectar rewards (assessed in calories as mean ± s.e.) of the two

common morphs GR (16.74 ± 1.47) and GW (18.04 ± 2.2). However, the total nectar reward

was highest for the rare RW morph (31.31 ± 2.25) due to both its higher nectar volume and

concentration (Fig. 1b; Table S2; Kruskal-Wallis $\chi 2$ = 33.53, df = 2, p < 0.0001). Flowers of

C. caulina were visited by two species of hawkmoths with long probosces: *Agrius convolvuli*

(~12.4 cm) and *Hippotion rafflesii* (~4.6 cm, Fig. S3; Sphingidae). During foraging, the

moths insert their proboscis into the floral tube, hover over the flower, and the sticky pollen is

carried on the proboscis of *A. convolvuli* (Fig. S3) or on the head in the case of *H. rafflesii*.

We did not observe the curling of the proboscis during foraging, as noted by Smith et al.

(2022). The average time spent by each pollinator per flower was ~ 1.5 seconds for *A.*

convolvuli and ~ 2.5 seconds for *H. rafflesii*, and the rupture in anther theca could be used as

a visual feature of flowers visited by hawkmoths.

The circular statistical analysis of visitation rates across the anthesis time showed no

temporal differences in pollinator visitations among the three morphs (Rayleigh Test; p <

0.0001; Fig. S4). The mean pollinator visitation time, denoted by the circular mean vector, μ

was noted to be between 21:00 to 22:00 for all morphs (Fig. S4). All three morphs showed

the highest hawkmoth activity during the first few hours of anthesis, between 19:00 hrs to

23:00 hrs, which gradually decreased until there was no activity by 06:00 hrs the following

morning (Fig. S4). Pollinator visitation rates of the three morphs during the peak visitation

hours were statistically different from each other with RW morph showing the highest

visitation rate (mean number of visits per flower per hour ± s.e.; 5.14 ± 0.96), followed by the

GW morph (2.8 ± 0.59), and GR morph (2.19 ± 0.3; Fig. 2c; Kruskal-Wallis chi-square =

6.52; df = 2; p = 0.03839; Table S3). However, the visitation rate standardized for morph

abundance showed a significantly lower visitation rate (Kruskal-Wallis chi-square = 3.77; df

= 2; p = 0.01511) to the RW morph (mean number of visits per hour ± s.e.; 119.94 ± 10.75),

than the common GR (175.15 ±17.29) and GW morphs (171.39 ± 18.72; Fig. 1d;).

From video recordings of pollinator visitations, we also observed that hawkmoths

revisited the same flowers within an RW morph (~ 10-12 times), once they discovered this

rare but nectar-rich inflorescence. In contrast, the two common morphs GR and GW with low

nectar rewards received lower visitation rates, and pollinators rarely revisited the flowers.

Thus within-inflorescence visits were higher in the rare RW morph compared to the

negligible revisits observed in the common GR and GW morphs.

Fig. 1. Three bract color morphs of *C. caulina*, their nectar rewards energy, and natural pollinator visitation rates. a. From left, GW, GR, and RW morphs of *C. caulina*, b. The mean (± s.e.) nectar energy (cal), c. Bar plot represents pollinator visitation rates (mean ± s.e.) measured for each morph in the observation plots (y-axis) and simulated pollinator visitation rates (mean ± s.e.) calculated for the population shown by red dots. Different alphabets indicate significant differences between morphs ($p < 0.05$).

Physiological compatibility and natural fitness vary among the morphs

From a total of 478 hand-pollination treatments, which consisted of autogamous, selfing, and outcrossing treatments, only 108 treatments set fruits (Fig. 2a). None of the autogamous treatments set fruits. Selfing treatments failed to set fruits exclusively in the GR morph, whereas the GW and RW morphs demonstrated leaky self-compatibility with success rates of 6.67% and 9.68%, respectively. Fruit-set success in selfed treatments was 5.8%, which was significantly lower ($\chi2 = 31.299$, $p < 0.0001$; Table S4) when compared with fruit-set success in the intra-morph cross (48.38%) and inter-morph cross (58.4%). The GR morph

demonstrated the highest fruit set in the intra-morph outcrossed treatment with a success rate

of 46.67%, followed by the RW morph at 27.03% and the GW morph at 18.75% (Fig. 2a).

In the inter-morph crosses, we noted donor-specific differences in the GW and RW

morphs. The donor specificity was related to whether a morph was used as a male parent or a

female parent in the inter-morph cross-pollination treatments. In crosses where GW morph

was the male parent (i.e., pollen donor), overall lower success rates (pair a: GW x GR,

24.24% and in pair b: GW x RW, 25%) were recorded when compared to crosses where it

was used as a female parent, that is pollen recipient (in pair a: GR x GW, 43.33% and in pair

b: RW x GW, 44.44%, Fig. 2a). In the last inter-morph treatment (pair c) where RW and GR

morphs were crossed, fruit set rates were 37.84% when RW was the female morph and

46.67% when RW was the male morph.

Among all hand-pollination treatments, self-pollination treatments exhibited the

lowest seed count (mean number of seeds per matured fruit ± s.e.; 8.63 ± 3.15), followed by

intra-morph crosses (16.93 ± 1.8) and then inter-morph crosses (18.63 ± 1.2). In the self-

pollination treatment, the GR morph exhibited a significantly lower seed count (zero)

compared to the GW morph (14.5 ± 6.5) and RW morph (13.33 ± 4.37; Kruskal-Willis $\chi2 =$

5.2499, df = 2, p = 0.07244; Fig. 2b; Table S5 and S6). In the intra-morph cross-pollination

treatment, the seed counts varied significantly among the morphs. The GW morph exhibited

the lowest seed count (10.72 ± 2.7) compared to the GR morph (20.71 ± 2.8) and the RW

morph (15.7 ± 2.7; Kruskal-Willis $\chi2$ = 5.2499, df = 2, p = 0.07244; Fig. 2b; Table S6). Seed

count in inter-morph cross-pollination treatments was not significantly different among the

three pairs (Fig. 2b; Table S6). Finally, the mean total fruit set per plant in open pollination

treatments was recorded to be equal in all three morphs (Fig. 3a). However, the mean seed

count per fruit was significantly different among the morphs (Kruskal-Wallis $\chi2$ = 36.3222,

df = 2, p < 0.0001) with the GR morph being the highest (23.7 ± 1.5), followed by GW (20.3

± 1.2) and RW (11.5 ± 1.4; Fig. 3b). We also noted that the seed counts recorded in the open

(natural) pollination and self-pollination treatments of RW were similar (Fig. 2a and 3a).

Fig. 2. Physiological compatibility between the three floral color morphs of *C. caulina*. a.

Heatmap illustrates the percent success rate (values inside box) from 478 hand-pollination

from 15 treatments. The male morphs (pollen source) are displayed on the x-axis, while the

female morphs (stigma identity) are shown on the y-axis. b. Mean (± s.e.) seed count from the

pollination treatments in the three morphs of *C. caulina*. The male donors (pollen source) are

displayed on the x-axis while the boxes represent female recipients (stigma identity).

Significant differences are shown by different alphabets.

Fig. 3. Natural fitness of the three morphs of *C. caulina* measured as A. mean (± s.e.)
reproductive success (% success) of open pollination treatments and B. mean (± s.e.) seed
count of the successful fruits from those open pollination treatments. Small letters indicate
significant differences between female morphs. Results of the statistical test between morphs
are given in Table S11.

Associating morph abundance and floral traits

In the path analyses, the five variables resulted in one direct path between abundance
and reproductive fitness (e), seven indirect paths (a to d and f to h), one correlation path
between abundance and type of compatibility (i), and one path on the effect of reproductive
fitness on relative abundance (j; Fig. 4). The Chi-squared goodness of fit test, CFI, and TLI
scores identified a good fit for all four models (*C. caulina*, GR, GW and RW morph; Table
1). In the case of the GW morph, the model fit as estimated by TLI values was not good (TLI
= -0.65), and we did not find any significant relationships between the traits as well as for the
net effect. We found that both the direct effect (path e in Fig. 4) and the net effect of morph
abundance on reproductive fitness were positive and statistically significant in three (*C.
caulina*, GR morph, and RW morph) of the four path analyses models (Table 1).

In the species-specific model, *C. caulina* in KAS showed a positive and significant
effect of only morph abundance on total nectar reward (a), self-compatibility on reproductive
fitness (d'), and finally, reproductive fitness on relative morph abundance (j). The correlation
between morph abundance and self-compatibility was also noted to be positive (i''). For the
within-morph path analyses, significant effects among indirect effects were observed only in

the GR morph, although positive effects were present across several paths for the other two

morph-specific models (GW and RW). In the GR morph, the following paths were significant

- morph abundance and total nectar reward (a), total nectar reward and reproductive fitness

(g), cross-compatibility and reproductive fitness (d''), and the regression of reproductive

fitness on morph abundance (j). The direct effect between relative morph abundance and

reproductive fitness (e), the regression of the reverse effect, that is, between reproductive

fitness and abundance (j) was significant and positive in RW but negative and not significant

for GW. We failed to detect any significant relationships between the other traits in the RW

morph, although notably, the effect of self-compatibility on reproductive fitness was positive,

whereas the effect of cross-compatibility on reproductive fitness was negative but not

significant (path d; Table 1).

Fig. 4. Path diagrams illustrating the effect of morph abundance on reproductive fitness,

analysed a) as species-specific model, b) for the common GR morph and c) for the rare RW

morph. One-headed arrows (paths a—h and j) indicate regression, while the two-headed

arrow (path i) indicates correlation. Thick arrows denote significant effects (P < 0.05), with

the strength of regression or correlation specified adjacent to them. Path IDs for all direct and

indirect effects are provided, and their strengths can be referenced in Table 1.

Table 1: Results of path analysis for all four models. Bold values represent the statistical

significance. Path IDs are the same as fig. 4.

		For *C. caulina*	GR morph	GW morph	RW morph
Model fit:	Chi-square statistics	0.052	0.037	0.006	0.664
	p-value (Chi-square)	**0.82**	**0.847**	**0.94**	**0.415**
	CFI value	**1**	**1**	1	**1**
	TLI value	**1.332**	**7.138**	-0.65	**1.494**

Regression:	**Path ID**				
Reproductive fitness ~ Relative abundance	e	**2.324**	**8.64**	-3.21	**12.241**
Total nectar reward	g	-0.122	**-1.94**	0.416	-0.63
Pollinator visitation rates	f	0.004	0.048	0.019	0.08
Self-compatibility	d'	**3.077**	NA	1.922	8.672
Cross-compatibility	d''	0.092	**0.428**	0.121	-0.341
Pollinator visitation rates ~ Relative abundance	h	1.058	-8.15	-2.84	-3.509
Total nectar reward	b	0.56	1.221	0.664	-1.327
Self-compatibility ~ Pollinator visitation rates	c'	-0.004	NA	-0.004	< 0.001
Cross-compatibility ~	c''	-0.013	-0.038	-0.013	-0.024

Pollinator visitation rates					
Total nectar reward ~ Relative abundance	a	**0.214**	**1.808**	-0.156	-0.112
Relative abundance ~ Reproductive fitness	j	**2.461**	**6.800**	-2.871	**10.942**
Correlation					
Relative abundance ~~ Self-compatibility	i'	**-42.15**	NA	-5.488	-0.303
Relative abundance ~~ Cross-compatibility	i''	214.939	-18.372	35.678	62.978
Net Total Effect		2.302	4.814	-3.219	12.045

Discussion

Floral color polymorphism is well-documented in various plant species, although the empirical assessment of the morphological, physiological, and ecological variations among floral color morphs in relation to their population densities remains underexplored (Elam & Linhart, 1988; Lau & Galloway, 2004; Caruso *et al.*, 2010; Berardi *et al.*, 2016; Koski *et al.*, 2020; Jacquemyn & Brys, 2020). In the tropics, species from the order Zingiberales (Costaceae, Heliconiaceae, Cannaceae and Marantaceae; Maas-Van De Kamer et al. 2016; Temeles *et al.*, 2000; Gowda & Kress, 2013) and the family Zingiberaceae (*Hedychium, Globba,* and *Curcuma*; Leong-Skornicková *et al.*, 2007; Saryan *et al.*, 2020) exhibit intra-specific floral and bract color polymorphisms along with pollinator-mediated selection of floral traits. This has prompted hypotheses regarding 'one pollinator-one color morph' specialization in several Zingiberales species (Temeles et al., 2013; Meléndez-Ackerman et al. 2005), though empirical validation has been limited to a few species, such as the Caribbean *H. caribaea* (Temeles et al. 2013) and the trans-Himalayan *Roscoea purpurea* and *R. tumjensis* (Paudel et al. 2018). This study reports findings on plant-pollinator interactions in the nocturnal color polymorphic *C. caulina*, endemic to the Western Ghats.

Based on our comprehensive characterization of the polymorphic *C. caulina* from Kaas plateau, we identified morph abundance to be tri-modal, with two common morphs (GR and GW) and one rare morph (RW) in the study population. Flowers of *C. caulina* displayed

classic moth-pollination syndromes (Johnson *et al.*, 2017), such as nocturnal opening, white

color, long floral tube with copious nectar, and flowers exposed from the inflorescence. True

to its syndrome, we discovered that *C. caulina* is exclusively pollinated by nocturnal

hawkmoths. However, in this nocturnal ginger-moth interaction, the concept of color

polymorphism is atypical. We found no morph-specific variations in floral morphologies,

such as floral tube length, floral size, or floral or inflorescence color and size that are

typically associated with pollinator-mediated selection in plants with floral color

polymorphism (Meléndez☐Ackerman and Campbell 1998; Gomez 2000; Majetic et al. 2007;

Milano et al. 2016). Instead, we found that the three morphs of *C. caulina* only differed in

their nectar traits, reproductive compatibility, and fitness (question 1), and pollinator

visitation was not shaped by morph abundance (question 2). We recorded an elaborate

pollinator-mediated frequency-dependent selection (FDS) only in one trait, nectar rewards.

We propose that the lack of variation in floral shape and size may be because these specific

floral traits have already undergone a stabilizing selection (Waser and Price 1981) by the

nocturnal hawk-moth pollinators, while nectar traits may still be under selection by

pollinators (Carlson and Holsinger 2013) or some abiotic factors (Plos et al. 2023; Malovrh et

al. 2024). Our failure to detect the selection of any morphs based on the color of their lateral

bracts suggests that while color acts as a morphological identifier, other physiological or

ecological traits linked to bract color may experience a direct pollinator-mediated or

frequency-dependent selection (question 3).

442   While color may not be under direct pollinator-mediated selection, in the path

analyses, we identified a few ecological and floral traits that showed a direct or indirect effect

on the differential reproductive fitness of the morphs. The first trait we identified is morph

abundance, which showed a strong direct effect on the morph's reproductive fitness (path e)

in both species-specific and moth-species models (Fig. 4). We noted that while visitation

rates were high for RW in our experimental setups, the simulated pollinator visitation

estimates for all RW in the population showed lower visitation rates simply because RW is

rare in the population (Fig. 1c Y axis vs Z axis). In contrast, we recorded fewer visits to the

common GR morph in the experimental setups, but we noted higher visitation estimates in

the simulation, since GR is common in the population. The contrasting pattern between the

visitation rate recorded per morph (Fig. 1c, Y axis) in our experimental set ups and

simulation results (Z axis) for the same morph shows that morph frequency does influence

the total visits received by each morph in this population.

Physiological compatibility is the second trait affecting reproductive fitness (path d)

with cross-compatibility having a significant effect only in the GR morph-specific model

(Fig. 4b) and self-compatibility showing a significant effect in the overall species-specific

model (Fig. 4a). The identification of self-incompatibility and enhanced reproductive fitness

exclusively in the common GR morph, alongside the absence of high reproductive fitness in

the self-compatible rare RW morph—despite higher pollinator visitation rates—further

indicates that the contrasting morph-specific compatibility plays an important role in the

differential reproductive assurance in the common GR and the rare RW morphs. Our

discovery of self-compatibility in the rare morph also aligns with studies demonstrating that

self-compatibility in floral color morphs is advantageous when the morph is rare in the

population (Holsinger, 2000) and when the morph is under pollen or pollinator limitation

(Subramaniam and Rusher 2000; Arista et al. 2013; Ortiz et al. 2015; Fry and Rausher 2017).

Physiological compatibility in *C. caulina* was noted to be labile, especially evident in

the hand-pollination treatments. For instance, intra-morph crossing success was morph-

specific, and inter-morph crossing success was donor-specific, meaning crossing success

depended on which morph was used as a pollen donor or pollen recipient (Fig. 2a). Although

such morph-specific compatibility differences are known only from a few studies, this pattern

has been mostly reported from annual plants (Wolfe and Seller 1997; Jorgenson et al. 2006;

Kosi and Galloway 2020) and rarely from perennials (Stanton et al. 1989). However, it is

premature to assume that the origin and maintenance of this phenomenon are comparable

between these contrasting life forms. While morph-specific compatibility can contribute to

the maintenance of polymorphism under both negative or positive frequency-dependent

selection in annuals, in perennial species like *C. caulina*, this is hard to determine (Jesson and

Barrett 2002). This is because, in perennial species, it is possible that different morphs

evolved under distinct evolutionary timeframes, resulting in an extant population composed

of relic individuals from different selection regimes. Thus, we cannot entirely dismiss the fact

that the different compatibility traits of the color morphs of *C. caulina* may have originated

under different historical selection regimes and different ecological conditions.

Nectar reward is the third trait that is variable among morphs, has an indirect

significant relationship with morph abundance, and affects reproductive fitness. Morph-

specific variability in nectar traits within a population has been suggested to show a balancing

selection in *I. purpurea* (Fry and Rausher, 1997; Subramaniam and Rausher, 2000). In *C.

caulina*, morph-specific nectar variability may also result from balancing selection, although

the mechanism is fairly complex and involves multiple traits. While high nectar energy in the

rare morph and low nectar energy in the common morph resembles negative FDS, the

resulting reproductive fitness contradicts the expected outcomes. We observed lower

reproductive fitness in the rare morph despite high pollinator visitation rates, whereas the

common morph showed higher fruit and seed sets despite lower pollinator visitation rates.

Therefore, although nectar quality predicts higher pollinator visitations, nectar alone did not

determine reproductive fitness in *C. caulina*. However, it does ensure pollinator visitations,

especially in the rare morph.

So, what contributes to the observed disparity in morph abundances in *C. caulina* at

Kaas? We propose a multi-trait model involving an interplay between nectar reward,

pollinator attraction, and compatibility type (self vs. cross), which influences the reproductive

success of different morphs and subsequently affects their population densities. For example,

the common GR morph ensures greater reproductive success despite its lower nectar energy

and lower pollinator visitation rate, since its higher abundance allows for a quicker detection

by pollinators. This is complemented by self-incompatibility, which enhances fruit and seed

set, promoting higher abundance of the common morph (Fig. 4b, Table 1). Conversely, the

rare RW morph, while less likely to be discovered by pollinators, benefits from its high-

energy nectar that encourages twice as many pollinator visits compared to the common

morph. However, video recordings of individual visits revealed that RW flowers received

repeated visits from the same moth, which can increase self-pollination (Harder *et al.*, 2004;

M. Eckhart *et al.*, 2006; Ishii & Harder, 2006). While this ensures reproductive success in the

self-compatible rare morph (Fig. 4c, Table 1), it comes at the cost of an overall reduced

reproductive fitness in naturally pollinated RW individuals (Fig. 2a and 3a). In contrast, we

observed a higher frequency of inter-individual switches (from video recordings, SS and VG)

in the common GR and GW morphs, likely facilitating outcrossing, as confirmed by higher

seed set in the open pollinated GR and GW morphs despite lower pollinator visits.

Population shifts in morph ratios are known in annual plants that are under FDS.

While we elucidate ecological mechanisms for the maintenance of rare morphs in *C. caulina*

in the Kaas population, we did not record transitions in morph ratios or elimination of the rare

morph in at least 15 years of our ecological studies at the Kaas plateau (unpublished data,

collected by authors). This may be because change in morph frequencies will often go

undetected in perennial rhizomatous species that also propagate vegetatively (as reported in

heterostylous species: Eckert and Barrett, 1993; Eckert et al., 1999; Barrett, 2015 and in

dioecious species: Irish and Nelson, 1989; Petzold et al., 2013). Therefore, in the light of

vegetative propagation whose importance cannot be discounted in gingers (Záveská et al.

2012; Skopalíková et al., 2023), we suggest that ecological factors identified in this study can

only explain morph maintenance via sexual reproduction, and vegetative reproduction can

reduce the intensity of frequency-dependent selection (Barrett and Harder 1996; Jesson and

Barrett 2002).

Color morphs have often been associated with abiotic-stress-related anthocyanin

production (Warren & Mackenzie, 2001; Tang & Huang, 2010; Berardi *et al.*, 2013, 2016;

Narbona *et al.*, 2018; Sapir *et al.*, 2021). A question rarely addressed in perennial plants that

show color polymorphism is whether the different color morphs represent historical stress

regimes in the population. We reiterate that the different color morphs and traits linked with

them, that is, morph abundance, physiological compatibility, and nectar traits, may represent

morphs selected under different abiotic stress regimes. Therefore, while we elucidate three

ecological mechanisms that maintain variable morph ratios in *C. caulina*, we do not discount

the role of balancing selection via abiotic factors also shaping polymorphism in *C. caulina*.

We aim to test this hypothesis under a population genetic framework since homologous

mutations among morphs will be key in establishing the genetic identities of the morphs

within and across populations.

In summary, *C. caulina* from the Kaas population demonstrates trait variations and

differential selection among its color morphs, which also results in their differential

reproductive fitness, which explains their highly skewed morph distribution. The role of

environmental factors in the maintenance of FCPs has been discussed in a few perennial and

rhizomatous plants (Twyford *et al.*, 2018; Vaidya *et al.*, 2018; Streinzer *et al.*, 2019; Tang *et*

al., 2016) and they suggest that FCPs are common in regions with higher climatic

fluctuations and high diversity of herbaceous species. Interestingly, these are also habitat

features of the sky-islands of the Western Ghats, where the Kaas plateau is located. Our

results from *C. caulina* reveal physiological and ecological factors that differentiate morphs

within a polymorphic perennial species and the interplay between them. Although some of

these factors can be examined through an ecological lens, perennial plants carry the burden of

the past, necessitating the consideration of both abiotic and biotic histories to fully

comprehend polymorphism in plants.

Acknowledgements

Authors thank the Ministry of Human Resource Development (MHRD) and the National

Geographic Society for research funds to VG; Heliconia Society Internationals for funds to

SS; IISER Bhopal for the infrastructure; DST for the INSPIRE fellowship to SS. We also

thank the PCCF, Maharashtra Forest Department, DFO of Satara Division (Forest

Department), and the Joint Forest Management Committee (JFMC), Kaas, Satara for

permitting our fieldwork. We thank Mr. Dilip Shrotri and Azad G for assistance during night

work, Amal, Rahul, Viraj, Asawari, Najla, and Sooraj for their assistance during various data

collections, including morphometry and Susnata and Sukhraj for assistance during

physiological compatibility experiments during the fieldwork. Finally, we thank all fellow

TrEE lab members for their valuable advice, discussions, and support.

Competing interests

564   The authors have no conflicts of interest to declare.

Author contributions

566   SS and VG conceived and designed the study. SS conducted the experiments and collected

567   the field data. SS performed the statistical analyses with inputs from VG. SS drafted the

568   manuscript; VG provided conceptual advice and edited the manuscript. All authors gave final

569   approval for publication.

Data availability

571   The entire dataset and analysis codes used in this study are accessible from the authors upon

572   reasonable request.

References

Arista M, Talavera M, Berjano R, Ortiz PL. 2013. Abiotic factors may explain the

geographical distribution of flower color morphs and the maintenance of color polymorphism

in the scarlet pimpernel. *The Journal of Ecology* 101: 1613–1622.

Barazani O, Erez T, Ogran A, Hanin N, Barzilai M, Dag A, Shafir S. 2019. Natural variation

in flower color and scent in populations of *Eruca sativa* (Brassicaceae) affects pollination

behavior of honey bees. *Journal of Insect Science* 19: 6.

Barrett, S. C. (2015). Influences of clonality on plant sexual reproduction. *Proceedings of the*

National Academy of Sciences, *112*(29), 8859-8866.

Benadi, G., & Gegear, R. J. (2018). Adaptive foraging of pollinators can promote pollination
of a rare plant species. *The American Naturalist*, *192*(2), E81-E92.

Berardi AE, Fields PD, Abbate JL, Taylor DR. 2016. Elevational divergence and clinal
variation in floral color and leaf chemistry in *Silene vulgaris*. *American Journal of Botany*
103: 1508–1523.

Berardi AE, Frey F, Denton EM, Wells JH. 2013. Betalain color morphs exhibit differential
growth, defensive ability, and pollen tube growth rates in *Mirabilis jalapa* (Nyctaginaceae).
International Journal of Plant Sciences 174: 1229–1238.

Bolten AB, Feinsinger P, Baker HG, Baker I. 1979. On the calculation of sugar concentration
in flower nectar. *Oecologia* 41: 301–304.

Brunet J, Thairu MW, Henss JM, Link RI, Kluever JA. 2015. The effects of flower, floral
display, and reward sizes on bumblebee foraging behavior when pollen is the reward and
plants are dichogamous. *International Journal of Plant Sciences* 176: 811–819.

Carlson JE, Holsinger KE. 2013. Direct and indirect selection on floral pigmentation by
pollinators and seed predators in a color polymorphic South African shrub. *Oecologia* 171:
905–919.

Caruso C, Scott SL, Wray JC, Walsh CAJ. 2010. Pollinators, herbivores, and the maintenance
of flower color variation: A case study with *Lobelia siphilitica*. *International Journal of Plant*
Sciences 171: 1020–1028.

Dixon P. 2003. VEGAN, a package of R functions for community ecology. *Journal of*
vegetation science 14: 927–930.

Dormont L, Delle-Vedove R, Bessière J-M, Hossaert-Mc Key M, Schatz B. 2010. Rare
white-flowered morphs increase the reproductive success of common purple morphs in a
food-deceptive orchid. *The New Phytologist* 185: 300–310.

Eckert, C. G., & Barrett, S. C. (1993). Clonal reproduction and patterns of genotypic diversity
in Decodon verticillatus (Lythraceae). *American Journal of Botany*, *80*(10), 1175-1182.

Eckert, C. G., Dorken, M. E., & Mitchell, S. A. (1999). Loss of sex in clonal populations of a
flowering plant, Decodon verticillatus (Lythraceae). *Evolution*, *53*(4), 1079-1092.

Elam DR, Linhart YB. 1988. Pollination and seed production in *Ipomopsis Aggregata*: Differences among and within flower color morphs. *American Journal of Botany* 75: 1262–1274.

Ford EB. 1945. Polymorphism. *Biological Reviews of the Cambridge Philosophical Society* 20: 73–88.

Fry, J. D., & Rausher, M. D. (1997). Selection on a floral color polymorphism in the tall morning glory (Ipomoea purpurea): transmission success of the alleles through pollen. *Evolution*, *51*(1), 66-78.

Fry JD, Rausher MD. 1997. Selection on a floral color polymorphism in the tall morning glory (*Ipomoea purpurea*): Transmission success of the alleles through pollen. *Evolution* 51: 66–78.

Gegear, R. J., & Laverty, T. M. (2005). Flower constancy in bumblebees: a test of the trait variability hypothesis. *Animal Behaviour*, *69*(4), 939-949.

Gigord LD, Macnair MR, Smithson A. 2001. Negative frequency-dependent selection maintains a dramatic flower color polymorphism in the rewardless orchid *Dactylorhiza sambucina* (L.) Soo. *Proceedings of the National Academy of Sciences, USA* 98: 6253–6255.

GÓmez G. 2000. Phenotypic selection and response to selection in *Lobularia maritima*: importance of direct and correlational components of natural selection. *Journal of Evolutionary Biology* 13: 689–699.

Gowda V, Kress WJ. 2013. A geographic mosaic of plant-pollinator interactions in the Eastern Caribbean islands. *Biotropica* 45: 224–235.

Goyret J, Pfaff M, Raguso RA, Kelber A. 2008. Why do *Manduca sexta* feed from white flowers? Innate and learnt color preferences in a hawkmoth. *Die Naturwissenschaften* 95: 569–576.

Hahn M, Brühl CA. 2016. The secret pollinators: an overview of moth pollination with a focus on Europe and North America. *Arthropod-Plant Interactions* 10: 21–28.

Harder, L.D., Jordan, C.Y., Gross, W.E. and Routley, M.B., 2004. Beyond floricentrism: the pollination function of inflorescences. *Plant Species Biology*, *19*(3), pp.137-148.

Harris DJ, Poulsen AD, Frimodt-Møller C, Preston J, Cronk QCB. 2000. Rapid radiation in
Aframomum (Zingiberaceae): Evidence from nuclear ribosomal DNA internal transcribed
spacer (ITS) sequences. *Edinburgh Journal of Botany* 57: 377–395.

Heinrich, B., 1975. Bee flowers: a hypothesis on flower variety and blooming times.
Evolution, 325-334.

Holsinger, K.E., 2000. Reproductive systems and evolution in vascular plants. *Proceedings of*
the National Academy of Sciences, *97*(13), pp.7037-7042.

Imbert E, Wang H, Conchou L, Vincent H, Talavera M, Schatz B. 2014. Positive effect of the
yellow morph on female reproductive success in the flower color polymorphic *Iris lutescens*
(Iridaceae), a deceptive species. *Journal of Evolutionary Biology* 27: 1965–1974.

Irish, E. E., & Nelson, T. (1989). Sex determination in monoecious and dioecious plants. *The*
plant cell, *1*(8), 737.

Ishii H, Harder L. 2006. The size of individual *Delphinium* flowers and the opportunity for
geitonogamous pollination. *Functional Ecology* 20: 1115–1123.

Jacquemyn H, Brys R. 2020. Lack of strong selection pressures maintains wide variation in
floral traits in a food-deceptive orchid. *Annals of Botany* 126: 445–453.

Jesson, L. K., & Barrett, S. C. (2002). Enantiostyly in Wachendorfia (Haemodoraceae): the
influence of reproductive systems on the maintenance of the polymorphism. *American*
Journal of Botany, *89*(2), 253-262.

Johnson SD, Moré M, Amorim FW, Haber WA, Frankie GW, Stanley DA, Coccuci AA,
Raguso RA. 2017. The long and the short of it: a global analysis of hawkmoth pollination
niches and interaction networks. *Functional Ecology* 31: 101–115.

Johnson SD, Raguso RA. 2016. The long-tongued hawkmoth pollinator niche for native and
invasive plants in Africa. *Annals of Botany* 117: 25–36.

Jones KN, Reithel JS. 2001. Pollinator-mediated selection on a flower color polymorphism in
experimental populations of *Antirrhinum* (Scrophulariaceae). *American Journal of Botany* 88:
447–454.

Jorgensen, T. H., Petanidou, T., & Andersson, S. (2006). The potential for selection on pollen
colour dimorphisms in Nigella degenii: morph-specific differences in pollinator visitation,
fertilisation success and siring ability. *Evolutionary Ecology*, *20*, 291-306.

Kellenberger RT, Byers KJRP, De Brito Francisco RM, Staedler YM, LaFountain AM,
Schönenberger J, Schiestl FP, Schlüter PM. 2019. Emergence of a floral color polymorphism
by pollinator-mediated overdominance. *Nature Communications* 10: 63.

Koski, M. H., & Galloway, L. F. (2020). Geographic variation in floral color and reflectance
correlates with temperature and colonization history. *Frontiers in plant science*, *11*, 991.

Koski MH, Berardi AE, Galloway LF. 2020. Pollen color morphs take different paths to
fitness. *Journal of Evolutionary Biology* 33: 388–400.

Kovach, W.L., 2011. Oriana–circular statistics for windows, ver. 4. *Kovach Computing*
Services, Pentraeth, Wales, UK.

Lau JA, Galloway LF. 2004. Effects of low-efficiency pollinators on plant fitness and floral
trait evolution in *Campanula americana* (Campanulaceae). *Oecologia* 141: 577–583.

Leong-Skornickóvá J, Sída O, Jarolímová V, Sabu M, Fér T, Trávníček P, Suda J. 2007.
Chromosome numbers and genome size variation in Indian species of *Curcuma*
(Zingiberaceae). *Annals of Botany* 100: 505–526.

Lide, D.R. ed., 2004. *CRC handbook of chemistry and physics* (Vol. 85). CRC press.

Liu C-Q, Sun H. 2019. Pollination in *Lilium sargentiae* (Liliaceae) and the first confirmation
of long☐tongued hawkmoths as a pollinator niche in Asia. *Journal of Systematics and*
Evolution 57: 81–88.

M. Eckhart V, S. Rushing N, M. Hart G, D. Hansen J. 2006. Frequency–dependent pollinator
foraging in polymorphic *Clarkia xantiana* ssp. *xantiana* populations: implications for flower
color evolution and pollinator interactions. *Oikos* 112: 412–421.

Maas-Van De Kamer, H., Maas, P. J. M., Wieringa, J. J., & Specht, C. D. (2016). Monograph
of African Costaceae. *Blumea-Biodiversity, Evolution and Biogeography of Plants*, *61*(3),
280-318.

Majetic CJ, Raguso RA, Ashman TL. 2009. The sweet smell of success: floral scent affects
pollinator attraction and seed fitness in *Hesperis matronalis*. *Functional Ecology* 23: 480–
487.

Majetic CJ, Raguso RA, Tonsor SJ, Ashman T-L. 2007. Flower color-flower scent
associations in polymorphic *Hesperis matronalis* (Brassicaceae). *Phytochemistry* 68: 865–
874.

Malovrh, K., Bavcon, J., Križman, M., & Ravnjak, B. (2024). Effect of Abiotic Factors on
Nectar Quality and Secretion of Two Early Spring Species, Galanthus nivalis L. and
Helleborus niger L. Diversity, 16(8), 469.

Meléndez-Ackerman, E. J., Speranza, P., Kress, W. J., Rohena, L., Toledo, E., Cortés, C., ...
& Soltis, D. (2005). Microevolutionary processes inferred from AFLP and morphological
variation in Heliconia bihai (Heliconiaceae). *International Journal of Plant Sciences*, *166*(5),
781-794.

Melendez-Ackerman E, Campbell DR. 1998. Adaptive significance of flower color and inter-
trait correlations in an *Ipomopsis* hybrid zone. *Evolution* 52: 1293.

Milano, E. R., Kenney, A. M., & Juenger, T. E. (2016). Adaptive differentiation in floral
traits in the presence of high gene flow in scarlet gilia (Ipomopsis aggregata). *Molecular*
Ecology, *25*(23), 5862-5875.

Mitchell, R. J. (1993). Adaptive significance of Ipomopsis aggregata nectar production:
observation and experiment in the field. *Evolution*, *47*(1), 25-35.

Mitchell, R. J. (2001). Path analysis. Design and analysis of ecological experiments.

Mood, J.D., Prince, L.M., Veldkamp, J.F. and Dey, S., 2013. The history and identity of
Boesenbergia longiflora (Zingiberaceae) and descriptions of five related new taxa. *Gard.*
Bull. Singapore, *65*(1), 47-95.

Narbona, E., Wang, H., Ortiz, P.L., Arista, M. and Imbert, E., 2018. Flower color
polymorphism in the Mediterranean Basin: occurrence, maintenance and implications for
speciation. *Plant Biology*, *20*, 8-20.

Ortiz PL, Berjano R, Talavera M, Rodríguez-Zayas L, Arista M. 2015. Flower color

polymorphism in *Lysimachia arvensis*: How is the red morph maintained in Mediterranean

environments? *Perspectives in Plant Ecology, Evolution and Systematics* 17: 142–150.

Paudel, B. R., Kessler, A., Shrestha, M., Zhao, J. L., & Li, Q. J. (2019). Geographic isolation,

pollination syndromes, and pollinator generalization in Himalayan Roscoea

spp.(Zingiberaceae). *Ecosphere*, *10*(11), e02943.

Paudel BR, Kessler A, Shrestha M, Zhao JL, Li Q. 2019. Geographic isolation, pollination

syndromes, and pollinator generalization in Himalayan *Roscoea* spp. (Zingiberaceae).

Ecosphere 10: e02943.

Petzold, A., Pfeiffer, T., Jansen, F., Eusemann, P., & Schnittler, M. (2013). Sex ratios and

clonal growth in dioecious Populus euphratica Oliv., Xinjiang Prov., Western China. *Trees*,

27, 729-744.

Plos, C., Stelbrink, N., Römermann, C., Knight, T. M., & Hensen, I. (2023). Abiotic

conditions affect nectar properties and flower visitation in four herbaceous plant species.

Flora, 303, 152279.

Poulsen AD, Lock JM. 1997. New species and new records of Zingiberaceae and Costaceae

from Tropical East Africa. *Kew Bulletin* 52: 601–616.

Raju A, Rao S, Ezradanam V, Zafar R, Kalpana P, Kiran P. 2004. The Hawkmoth,

Macroglossum gyrans and its interaction with some plant species at Visakhapatnam. *Zoos'*

print journal 19: 1595–1598.

R Core Team. (2023). *R: A language and environment for statistical computing* [Software]. R

Foundation for Statistical Computing. https://www.R-project.org/

Ren S, Vitasse Y, Chen X, Peichl M, An S. 2022. Assessing the relative importance of

sunshine, temperature, precipitation, and spring phenology in regulating leaf senescence

timing of herbaceous species in China. *Agricultural and Forest Meteorology* 313: 108770.

Rosseel Y. 2012. Lavaan: An R package for structural equation modeling. *Journal of*

Statistical Software 48: 1–36.

Sapir Y, Gallagher MK, Senden E. 2021. What maintains flower color variation within

populations? *Trends in Ecology & Evolution* 36: 507–519.

Saryan P, Gupta S, Gowda V. 2020. Species complex delimitations in the genus *Hedychium*:

a machine learning approach for cluster discovery. *Applications in Plant Sciences* 8: e11377.

Skopalíková, J., Leong‑Škorničková, J., Šída, O., Newman, M., Chumová, Z., Zeisek, V., ...

& Záveská, E. (2023). Ancient hybridization in Curcuma (Zingiberaceae)—Accelerator or

brake in lineage diversifications?. *The Plant Journal*, *116*(3), 773-785.

Smith GP, Davidowitz G, Raguso RA, Bronstein JL. 2022. Proboscis curling in a pollinator

causes extensive pollen movement and loss. *Ecological Entomology* 47: 234–241.

Smithson, A., & Macnair, M. R. (1997). Negative frequency‑dependent selection by

pollinators on artificial flowers without rewards. *Evolution*, *51*(3), 715-723.

Stanton, M. L., Snow, A. A., Handel, S. N., & Bereczky, J. (1989). The impact of a

flower‑color polymorphism on mating patterns in experimental populations of wild radish

(Raphanus raphanistrum L.). *Evolution*, *43*(2), 335-346.

Stebbins GL. 2019. Chapter VI. Isolation and the Origin of Species. In: Chapter VI. Isolation

and the Origin of Species. *Columbia University Press*, 189–250.

Streinzer M, Roth N, Paulus HF, Spaethe J. 2019. Color preference and spatial distribution of

glaphyrid beetles suggest a key role in the maintenance of the color polymorphism in the

peacock anemone (*Anemone pavonina*, Ranunculaceae) in Northern Greece. *Journal of*

Comparative Physiology A 205: 735–743.

Subramaniam B, Rausher MD. 2000. Balancing selection on a floral polymorphism.

Evolution 54: 691–695.

TAKANO, A., Julius, A. and Mohamed, M., 2009. First report of flexistyly in *Plagiostachys*

(Zingiberaceae). *Acta Phytotaxonomica et Geobotanica*, *60*(1), pp.56-59.

Tang X, Liu H, Zhang C, Wang Y, Zhang X, Yi Y. 2016. Flower color polymorphism in

G*eranium nepalense* (Geraniaceae): Adaptation to non-pollinator agents. *Polish Journal of*

Ecology 64: 526–533.

Tang X-X, Huang S-Q. 2010. Fluctuating selection by water level on gynoecium color

polymorphism in an aquatic plant. *Annals of Botany* 106: 843–848.

Temeles, E. J., Pan, I. L., Brennan, J. L., & Horwitt, J. N. (2000). Evidence for ecological
causation of sexual dimorphism in a hummingbird. *Science*, *289*(5478), 441-443.

Temeles EJ, Rah YJ, Andicoechea J, Byanova KL, Giller GSJ, Stolk SB, Kress WJ. 2013.
Pollinator-mediated selection in a specialized hummingbird-*Heliconia* system in the Eastern
Caribbean. *Journal of Evolutionary Biology* 26: 347–356.

Thompson FL, Eckert CG. 2004. Trade-offs between sexual and clonal reproduction in an
aquatic plant: experimental manipulations vs. phenotypic correlations. *Journal of*
Evolutionary Biology 17: 581–592.

Twyford AD, Caola AM, Choudhary P, Raina R, Friedman J. 2018. Loss of color
pigmentation is maintained at high frequency in a Monkey flower population. *The American*
Naturalist 191: 135–145.

Vaidya P, McDurmon A, Mattoon E, Keefe M, Carley L, Lee C-R, Bingham R, Anderson JT.
2018. Ecological causes and consequences of flower color polymorphism in a self-pollinating
plant (*Boechera stricta*). *The New Phytologist* 218: 380–392.

Warren J, Mackenzie S. 2001. Why are all color combinations not equally represented as
flower-color polymorphisms? *The New Phytologist* 151: 237–241.

Waser NM, Price MV. 1981. Pollinator choice and stabilizing selection for flower color in
Delphinium nelsonii. *Evolution* 35: 376–390.

Waser NM, Price MV. 1985. The effect of nectar guides on pollinator preference:
experimental studies with a montane herb. *Oecologia* 67: 121–126.

Watve A. 2013. Status review of Rocky plateaus in the northern Western Ghats and Konkan
region of Maharashtra, India with recommendations for conservation and management.
Journal of Threatened Taxa 5: 3935–3962.

Weiss MR. 2001. Vision and learning in some neglected pollinators: beetles, flies, moths, and
butterflies. *Cognitive Ecology of Pollination*: 171–190.

Wenzell KE, Skogen KA, Fant J. 2023. Range wide floral trait variation reflects shifts in
pollinator assemblages, consistent with pollinator mediated divergence despite generalized
visitation. *Oikos* e09708.

Wolfe, L. M., & Sellers, S. E. (1997). Polymorphic floral traits in Linaria canadensis

(Scrophulariaceae). *American Midland Naturalist*, 134-139.

Yadav, R., & Gowda, V. (2024). Six new species of Globba L.(Zingiberales, Zingiberaceae)

from the Eastern Himalayas and Northeast India. *PhytoKeys*, *246*, 197.

Yadav R. 2024. Reproductive system in the genus *Globba* L. (Zingiberaceae): Understanding

its origin and maintenance. Unpublished PhD thesis. Indian Institute of Science Education

and Research Bhopal.

Zar, J.H., 1999. *Biostatistical analysis*. Pearson Education India.

Záveská, E., Fér, T., Šída, O., Krak, K., Marhold, K., & Leong-Škorničková, J. (2012).

Phylogeny of Curcuma (Zingiberaceae) based on plastid and nuclear sequences: Proposal of

the new subgenus Ecomata. *Taxon*, *61*(4), 747-763.

Supplementary information (brief legends)

Fig. S1 Plant habit, habitat, and key vegetative features of *Curcuma caulina*.

Fig. S2 Variation in bract color, floral reflectance, and morph frequency of *C. caulina*.

Fig. S3 Hawkmoth pollinators of *C. caulina*.

Fig. S4 Circular histogram and circular statistics values of pollinator visitation rates.

Table S1 List of floral and vegetative morphological characters used for the nMDS analysis.

Table S2 Within-group comparisons for nectar energy, volume, and concentration.

Table S3 Within-group comparisons for peak pollinator visitation rates.

Table S4 Results of the chi-square test for fruiting success rates of hand-pollination.

Table S5 Mean and standard error values for seed count for cross- and self-compatibility

treatments.

Table S6 Between groups comparisons for seed counts for hand pollination treatments.