Type of Article: **ORIGINAL ARTICLE**

A tale of three morphs I: Variations in nectar, reproductive compatibility, and

abundance explain reproductive success of the morphs in a polymorphic ginger from

Western Ghats, India

Authors: Saket Shrotri[1] and Vinita Gowda[1*]

[1]Tropical Ecology and Evolution (TrEE) Lab, Department of Biological Sciences, Indian

Institute of Science Education and Research (IISER) Bhopal, Bhopal, Madhya Pradesh, India,

462066.

*Corresponding author: Vinita Gowda.

Email: gowdav@iiserb.ac.in

ORCiD:

Saket Shrotri: 0000-0002-4596-9451

Vinita Gowda: 0000-0001-8533-0014

Running title:

Shrotri and Gowda; Bract color Polymorphism in *Curcuma caulina*

Abstract (word limit: 300/300)

- **Background and aims:** Floral color polymorphism is common in angiosperms, yet in perennial, clonal plants, proximate trait differences and reproductive differences among sympatric morphs remain poorly understood. This is primarily because perennial individuals can carry signatures of past selection regimes. We studied bract color polymorphism in a nocturnal ginger *Curcuma caulina* to ask: (i) Do nectar rewards, reproductive compatibility and pollinator visitation differ among color morphs? (ii) Does morph abundance correlate with fitness via ecological (pollinator-mediated) or physiological (compatibility) pathways? (iii) Do these traits predict realized reproductive success?

- **Methods:** We measured 22 floral/vegetative traits (n=30 per morph), nectar rewards (106 flowers), pollinator visitation rates (648 hours), and physiological compatibility (471 hand-pollinations treatments) and natural fruit and seed set ($\geq$50 per morph) across three dominant morphs. Path analysis tested direct and indirect effects of morph abundance, nectar traits, pollinator visitation, and compatibility on reproductive fitness.

- **Key results:** Morphs did not differ in their morphological traits but showed differential nectar and reproductive compatibility traits. The rare red-white morph produced the highest nectar energy, received the most visits, had a leaky self-compatibility, and yet showed lowest seed count per fruit, while the most common green-red morph was self-incompatible, showed higher cross-compatibility and had highest seeds per fruit. The second common green-white morph showed intermediate abundance and also showed leaky self-compatibility. Path analysis indicated that morph abundance had both direct and indirect effects on reproductive fitness, mediated by nectar, pollinator visitation and compatibility.

- **Conclusions:** Reproductive success of the polymorphic *C. caulina* is a result of multi-trait interactions including pollinator interaction. That is, nectar traits and mating-system differences shape reproductive fitness of the morphs. We also highlight that in polymorphic perennial plants contemporary selection regimes may be acting alongside genotypic vestiges (historical genotypes present in extant population), thus complicating any frequency-dependent selection regime.

Keywords

Curcuma caulina, Bract color polymorphism, Hawkmoth pollination, Laterite plateaus, Path analysis, Physiological compatibility, Reproductive fitness, Self-incompatibility

Introduction

Floral color polymorphism (henceforth FCP) is the co-occurrence of two or more discrete color forms within a single population. The origins of FCP are usually explained by standard genetic processes such as mutation and recombination (Kellenberger et al., 2019; Stebbins, 2019), and the maintenance of morphs in any population is often explained by ecological selection such as by pollinators. Both vertebrate and non-vertebrate pollinators are well known for their use of floral color as a cue to locate and choose flowers resulting in pollinator-mediated selection shaping the success and decline of morphs within a population (Schiestl and Johnson, 2013; Rausher, 2008; de Jager and Ellis, 2014; Milano et al., 2016; Bigio et al., 2017; Dyer et al., 2021). However, there is mounting evidence that shows that floral color morphs also vary in other traits such as morphological, physiological, reproductive, and ecological traits, which can ultimately affect the morph's fitness. Thus, recognizing the multivariate nature of FCP and characterizing how traits may covary among morphs is an essential precondition in order to identify selective agents that may be operating within a population.

Maintenance of FCP despite weak or absent pollinator preferences has been empirically shown (Schemske and Bierzychudek, 2007; Tang and Huang, 2010; Keasar et al., 2016; Hassa et al., 2020), and in several studies the target of selection within a population polymorphic for its floral and reproductive traits is not their floral color (Warren and Mackenzie, 2001; Subramaniam and Rausher, 2007; Dick et al., 2011; Fry and Rausher, 2017; Vaidya et al., 2018; Dafni et al., 2020; Table 1). Instead, these studies show that traits such as nectar rewards, flower size and number, pollen and ovule attributes, physiological tolerances, or mating-system differences alter reproductive outcomes independently of pollinator color choice and may play a greater role in the maintenance of the floral color morphs (Gomez, 2000; Wolfe, 1993; Carlson and Holsinger, 2013; Mu et al., 2011; Lau and Galloway, 2004; Kosi and Galloway, 2020; Arista et al., 2013; Ortiz et al., 2015). For example, seed viability differences linked to color morphs have been reported in *Boechera stricta* (Vaidya et al, 2018), while abiotic tolerance associated with color has been implicated in species such as

Parrya nudicaulis and *Anemone coronaria* (Dick et al., 2011; Dafni et al., 2020). Some of the

representative cases across plant families and selection regimes are summarized in table 1,

which shows the growing body of examples that associate diverse functional traits with floral

color morphs including life history strategies (annuals vs. perennials).

The consequences of color polymorphism may differ across life-history strategies (Narbona et

al., 2018; Sapir et al., 2021): in annuals, selection acts within a single reproductive cycle

(Vaidya et al., 2018), whereas in long-lived perennials (rhizomatous and non-rhizomatous

plants like trees) polymorphisms may persist or spread across multiple reproductive seasons

(Snell and Aarssen, 2005; Friedman, 2020), allowing potential ecological or physiological

advantages to accumulate over time. Therefore, life history strategy of a polymorphic species

can further influence its morphological and genetic characteristics. Whereas frequency-

dependent selection regimes may not even be detectable in perennials as past selections may

persist in the current population as genotypic vestiges (historical genotypes present in extant

population). This may be because change in morph frequencies often go undetected in

perennial rhizomatous species that also propagate vegetatively (as reported in heterostylous

species: Eckert and Barrett, 1993; Eckert et al., 1999; Barrett, 2015 and in dioecious species:

Irish and Nelson, 1989; Petzold et al., 2013). Thus, clonality may buffer against rapid

frequency shifts resulting in demographic feedback (Barrett, 2015). To the best of our

knowledge there have been no studies that have looked into whether life history strategies

influence the selection of polymorphism in perennial species. This is particularly striking for

FCP, which is frequently reported in perennials but has not yet been empirically investigated.

Thus, plants with floral color polymorphism simplistically reveal at least the following three

important points. First, color is often embedded in a broader phenotypic context; pigments

such as anthocyanins have physiological functions (protection against UV, heat, drought, and

pathogens) in addition to visual signaling (Narbona et al., 2018). Second, trait associations can

alter selection indirectly: a morph that is less preferred by pollinators may nonetheless gain

fitness if it enjoys higher autogamy, greater seed viability, or higher stress tolerance (Turelli et

al., 2001; Schemske and Bierzychudek, 2007; Barazani et al., 2019; Vaidya et al., 2018).

Third, perennial and clonal life histories complicate straightforward pollinator-mediated

selection narratives. Clonal plants with long adult lifespans can accumulate historical

demographic events which can decouple the effect of current selective agents on the extant

morph frequencies.

Given these complex sets of drivers shaping a species that shows floral color polymorphism,

we propose that as a first step it is critical to characterize phenotypic variations in perennial

species that show FCP in order to understand extant selection regimes that may be

maintaining the polymorphism within a population. To establish the variable traits among

morphs, require documenting not just visual features such as color and pollinator visitation

but also reproductive compatibility (autogamy, self-compatibility, cross-compatibility), floral

rewards, other floral and vegetative morphological traits, and performance under abiotic

stress. A multi-dimensional phenotypic characterization provides the empirical scaffolding

needed to (a) identify which traits differ consistently among morphs, (b) evaluate the

demographic and reproductive consequences of those differences, and (c) design targeted

analysis of causal mechanisms.

Here, we explore a polymorphic, nocturnal ginger *Curcuma caulina* from the family

Zingiberaceae. All gingers are perennials due to their rhizomatic habit and they offer ideal

grounds to test the role and maintenance of polymorphism in floral forms. Several genera

show pronounced intraspecific variation in flower and bract colors, and within the Asian

tropics a range of polymorphisms (stylar, bract color, and floral color polymorphisms) has

been reported across genera such as *Alpinia*, *Amomum*, *Curcuma*, *Hedychium* and *Roscoea*

(Poulsen & Lock, 1997; Harris et al., 2000; Leong-Skornicková et al., 2007; Mood et al.,

2013; Yadav, 2024; Yadav and Gowda, 2024). In some taxa, floral polymorphism has been

linked to pollinator shifts and speciation dynamics (Takano et al., 2009; Paudel et al., 2019),

but detailed, multi-trait descriptions remain rare for many species.

In this study we present a comprehensive characterization of bract-color polymorphism in *C.*

caulina, which is endemic to the laterite-plateaus of the Northern Western Ghats (Figure S1).

In our preliminary observations we had noted that morphs showed variable abundances within

a population, with a few dominant, and several rare morphs. Due to the absence of any prior

ecological studies on *C. caulina*, we focused on three complementary trait axes in order to

characterize intraspecific morph variations. They are: morphology (bract and floral display),

pollinator visitation, and reproductive compatibility traits (autogamy, cross-compatibility, and

seed set). These traits form the immediate causal chain linking phenotype to reproductive

output in animal-pollinated plants: bract color and rewards influence pollinator attraction and

visitation; visitation determines pollen transfer; and reproductive compatibility converts

pollen receipt into realized fitness. These axes capture both pre-pollination (pollinator

attraction and pollen receipt) and post-pollination (fertilization and compatibility) processes

and can be quantified in the field. More generally, the questions we ask are ecological. These

traits are analogous to those examined in studies investigating interspecific competition;

however, we utilize them within a polymorphic species that exhibits intraspecific variations

among its morphs.

Using field experiments we predicted that since higher nectar reward is known to increase

visits by pollinators, morphs with better nectar rewards will show higher pollinator visitation

rates, higher pollen transfer, and increased fruit set and seed set. Since we had noted

differences in morph abundances within our study site, we predicted that the most common

morph will show the highest nectar rewards, visitation rates, and fruit set and seed sets. We

also expected that rare morphs may compete for pollinator visits with the other more abundant

morphs, resulting in a trade-off between morph abundance and reproductive compatibility.

Based on these expectations, we addressed the following questions:

(i) Do nectar rewards, reproductive compatibility, and pollinator visitations differ among the

color morphs?

(ii) Do pollinator and plant reproductive traits explain reproductive success and do these

correlate with the observed morph abundance?

Materials and Methods

Study species and study site

Curcuma caulina (Zingiberaceae) is a rhizomatous, mass-flowering, perennial herb that

flowers between July to October (Fig. S1). The plant is 30–50 cm tall with a single shoot

bearing a terminal inflorescence which consists of 10-18 fertile 'lateral bracts' that bear

flowers (Fig. S1). Flowers are born in cincinni (1–3 flowers per cincinnus), are nocturnal

(anthesis at 18:00), and last 12–14 hours only. The stigma is funnel-shaped and projects

beyond the anther, and fruits mature within the bracts. *C. caulina* is endemic to the northern

Western Ghats of India and this study was carried out on Kaas Plateau (कास पठार; 17°43'12.36"

N, 73°49'28.0956" E, elevation 1225m, Fig. S1), Satara, Maharashtra, India between 2019

and 2022. Kaas Plateau is an elevated ferricrete outcrop (red laterite crust) and a UNESCO

Natural World Heritage site recognized for its floral diversity and high endemicity (Shrotri et

al., 2025).

The color of the lateral bracts of *C. caulina* individuals varied within the population from

predominantly green to predominantly red colored inflorescence (Fig. S2a). To determine the

frequency of bract color morphs, we identified and counted all morphs within randomly

placed ten transects, each measuring 10 x 4 m^2. Variations in the color of the lateral bracts

were first identified based on the extent of the reddish-pink coloration (0% to 100%) on the

first or basal lateral bract. This color difference was visually identified and also quantified

using handheld spectrophotometer (Ocean Optics; Fig. S2a, b).

To morphologically characterize the morphs, we measured a total of 22 vegetative and floral

characters using digital vernier caliper and rulers (n=30 per morph; character list in Table S1).

To assess morphological variations among the morphs, we ran a non-metric multidimensional

scaling (nMDS) analysis with Bray–Curtis dissimilarity using the VEGAN package (Dixon,

2003) in R (4.3.0; R core Team, 2021). We used the first three dimensions, to construct three-

dimensional (3D) plots, which were used to visualize both intra-morph and inter-morph

variations. The statistical difference among the overlapping clusters (morphs) was measured

using the ANOSIM test, where the Bray–Curtis dissimilarity matrix with 1000 permutations

was used (VEGAN package; Dixon, 2003).

We measured nectar volume and concentration in 106 bagged flowers (32 to 38 unopened

buds per morph) between 18:00 and 18:30 hrs by destructive sampling. The nectar volume

(μl), concentration (% sucrose), and nectar energy (cal) were compared among the three

morphs. Nectar volume was measured using calibrated glass microcapillary tubes (Drummond

Scientific Company), and the sucrose concentration in nectar was determined using a

handheld refractometer (ERMA BRIX 0~55%). The brix sucrose % value was then converted

into sugar (mg) per 1 ml of nectar using the table given in the CRC Handbook of Chemistry

and Physics (Lide, 2004) and as outlined by Bolten *et al.* (1979). This value was used to

calculate the amount of sugar in a given volume of nectar, and it was converted into calories

(1 mg of sucrose = 4 calories as given in Heinrich, 1975). Differences between the morphs

were tested using the Kruskal-Wallis test followed by Dunn's test as a post-hoc; the

significance was set at $p<0.05$; analysis was carried out in R (4.3.0; R Core Team, 2023).

 Natural pollinator visitations

To determine if pollinators exhibit differential visitation towards a particular morph,

pollinator visits were monitored by night vision CCTV cameras (Ae Zone CCTV IR day/night

vision USB camera) between 18:00 hrs and 06:00 hrs. These observations were carried out in

18 randomly selected observation plots on the plateau, where each plot consisted of three to

six individuals. All morphs were observed for a total of 648 hrs (i.e., 216 hrs per morph).

Cameras were positioned approximately 1 to 1.5 m away from the focal plants such that all

inflorescences were visible in their entirety in the field of view. The recordings were manually

scored, where the total number of visits by each pollinator and the total number of flowers

present in the plot were measured. We calculated the pollinator visitation rate as the total

number of visits per flower per hour for each morph. The visitation rates were summed for

every hour between 18:00 to 06:00 hrs in order to record the temporal shift in pollinator

visitation rate.

Natural pollinator visitation rates (number of visits per flower per hour) were compared

among the morphs on a temporal scale using the circular statistical analysis in ORIANA (v.

4.02 Kovach Computing Services; Kovach, 2011). Temporal differences in visitation rates

between the morphs were tested using the Mardia-Watson Wheeler test (or uniform score test;

Zar, 1999). Since the pollinator observation plots were not always located within the morph-

census transects, the pollinator visitation rates were standardized for morph frequency using

Monte Carlo Simulation (n = 10,000). The morph frequency and pollinator visitation rates for

each morph were utilized to estimate the distribution of their product for the respective

morphs. To test for inter-morph differences in pollinator visitation rates we used the peak

visitation rates from 19:00 hrs to 22:00 hrs, which accounted for ~85% of all pollinator visits.

We used the Kruskal-Wallis test followed by Dunn's test as a post-hoc test at $p<0.05$. All

statistical analyses were carried out in R (4.3.0; R Core Team, 2023).

Physiological compatibility and natural fruit set

To comprehensively evaluate the physiological self-compatibility and cross-compatibility

rates in the three floral color morphs, we carried out a total of 15 hand-pollination treatments.

Since we identified three bract color morphs in this population (refer to results for within

population morph variability), reproductive compatibility was tested for these three morphs,

where one morph was rare and two were common. Each morph was treated separately, and the

pollination treatments included three autogamy, three self-pollination, and nine cross-

pollinations that included three intra-morph cross-pollinations and six inter-morph cross-

pollinations were the identity of a morph as the male or female parent was recorded.

Autogamy treatment included unmanipulated, bagged flowers that were checked for fruit set

after four weeks. For cross-pollination treatments, flowers were emasculated and mixed pollen

grains from 10-15 individuals were manually transferred on marked flowers. Flowers were

observed for fruit set in ~4 weeks and percent success rate in reproduction was measured for

each treatment using the number of fruits per treatment as well as the number of seeds per

fruit (seed count). We statistically tested the reproductive success rates between self- and

cross-pollination treatments for each morph and compared male versus female reproductive

success for the morphs using Fisher's exact tests. Comparative analyses were also carried out

for the seed counts from self- and cross-pollination treatments using the Kruskal-Wallis test

followed by Dunn's post-hoc test with significance set at $p < 0.05$.

We quantified the reproductive success of natural pollination events within the study

population at the end of the flowering season in mid-October by measuring fruit set in ≥ 500

individuals per morph and seed counts in ≥ 50 per morph. For each inflorescence, we

estimated the total number of flowers per inflorescence by multiplying the number of lateral

bracts with the number of flowers estimated in a cincinnus. Fruit-set per inflorescence was

then calculated as the ratio of the number of matured fruits to the total number of flowers

estimated for a particular inflorescence. We used Kruskal-Wallis test to compare both natural

fruit set and natural seed count among morphs. Dunn's post-hoc test was used with a

significance set at $p < 0.05$.

Path analysis to test traits effects on the reproductive fitness of the morphs

We used path analysis to understand the direct and indirect effects of abundance of a morph

on its natural fitness as described by Mitchell (1993). Path analysis was tested for the entire

Kaas population of *C. caulina* as well as for each morph independently (n = 300 individuals;

i.e., 100 per morph) in order to assess the net effect and important selection regimes within the

population and within each morph. Three indirect effects included in the path analyses were-

total nectar reward, pollinator visitation rates, and physiological compatibilities. Values for

the total nectar reward were normalized with morph abundance and of natural fitness with

seed-set using Monte Carlo simulations (n = 10,000). Path analysis was performed in R (4.3.0;

R Core Team, 2023) using the *lavaan* package (Rosseel, 2012) and path estimates were

computed using 10,000 iterations in the bootstrapping procedure. Estimates of model fit were

computed using Chi-squared goodness of fit, Comparative Fit Index (CFI) and Tucker-Lewis

Index (TLI). A nonsignificant chi-squared value indicates that a model is not significantly

different from the observed correlations in the data and is therefore a good fit whereas CFI

and TLI indices above 0.9 are considered to be a good fit (Mitchell, 2001). Finally, we tested

the hypothetical relationship between reproductive fitness and relative abundance using a

regression model which was not part of the main path analysis.

Results

Floral (bract) color polymorphism and morphological variability

We identified six bract color variants with an almost continuous distribution within the

population (Fig. S2c). However, using Ford's criteria for polymorphic forms (Ford, 1945; Fig.

S2), we chose three most common variants (0% red, 60% red, and 100% red; henceforth

referred to as 'morphs') in this study. These three morphs were named based on the coloration

of their lateral and coma bracts as follows: the 0% variant is Green-White morph (GW), the

60% variant is Green-Red morph (GR), and the 100% variant is Red-White morph (RW; Supp

Fig. 1a). The distribution of these morphs was unequal, with GR morphs being the most

common (~ 44% occurrence), followed by GW (~ 29%) and RW (13%). The remaining three

bract color variants (15%, 30%, and 45%) were extremely rare, each constituting less than 5%

of the population (Fig. S2c).

The abundance of the three morphs differed within the population. The GR morph (98.6 ± 7.2

individuals; ~44 %) and GW morph (75.7 ± 7.2 individuals; ~29 %) were the most common

morphs; their frequencies were not significantly different from each other (Kruskal-Wallis $\chi2$

$= -0.692$, df $= 1$, $p = 0.2446$). The rarest of the three morphs was the Red-White morph (RW)

with 24.2 ± 2.0 individuals (~13 %) in the population, and the RW morph had significantly

fewer individuals when compared to both GR and GW (RW vs GR, Kruskal-Wallis $\chi2 = -$

2.427, df $= 1$, $p = 0.0076$ and RW vs GW, Kruskal-Wallis $\chi2 = 1.736$, df $= 1$, $p = 0.0413$

respectively). All other bract color variants totaled to ~14% in the population and thus were

extremely rare. The non-metric multidimensional scaling (nMDS) analysis of the 22 floral and

vegetative morphometric characters (Table S1; Fig. S2) failed to form distinct clusters

separating the morphs by their morphological traits. Further, the Analysis of Similarities

(ANOSIM) test ($p = 0.4835$) and the associated ANOSIM R-value (ANOSIM R = - 0.0038)

also indicated no statistical significance in the differentiation of these clusters for each morph.

Since we detected polymorphism in bract colors and not floral colors, henceforth, the term

color polymorphism refers to only bract color polymorphism in *C. caulina*.

Variability in nectar traits and hawkmoth visitation rates

Nectar is the primary pollinator reward in *C. caulina*, and there was no significant difference

between the total nectar rewards (assessed in calories as mean ± s.e.) of the two common

morphs GR (16.74 ± 1.47) and GW (18.04 ± 2.2). However, the total nectar reward was

highest for the rare RW morph (31.31 ± 2.25) due to both its higher nectar volume and

concentration (Fig. 1b; Table S2; Kruskal-Wallis $\chi2 = 33.53$, df = 2, $p < 0.0001$). Flowers of

C. caulina were visited by two species of hawkmoths with long probosces: *Agrius convolvuli*

(~12.4 cm) and *Hippotion rafflesii* (~4.6 cm, Fig. S3; Sphingidae). During foraging, the moths

insert their proboscis into the floral tube, hover over the flower, and the sticky pollen is

carried on the proboscis of *A. convolvuli* (Fig. S3) or on the head in the case of *H. rafflesii*.

We did not observe the curling of proboscises during foraging, as noted by Smith et al.

(2022). The average time spent by each pollinator per flower was ~ 1.5 seconds for *A.*

convolvuli and ~ 2.5 seconds for *H. rafflesii*, and the rupture in anther theca could be used as a

visual feature of flowers visited by hawkmoths.

The circular statistical analysis of visitation rates across the anthesis time showed no temporal

differences in pollinator visitations among the three morphs (Rayleigh Test; $p < 0.0001$; Fig.

S4). The mean pollinator visitation time, denoted by the circular mean vector, µ was noted to

be between 21:00 to 22:00 for all morphs (Fig. S4). All three morphs showed the highest

hawkmoth activity during the first few hours of anthesis, between 19:00 hrs to 23:00 hrs,

which gradually decreased until there was no activity by 06:00 hrs the following morning

(Fig. S4). Pollinator visitation rates of the three morphs during the peak visitation hours were

statistically different from each other with RW morph showing the highest visitation rate

(mean number of visits per flower per hour ± s.e.; 5.14 ± 0.96), followed by the GW morph

(2.8 ± 0.59), and GR morph (2.19 ± 0.3; Fig. 2c; Kruskal-Wallis chi-square = 6.52; df = 2; $p =$

0.03839; Table S3). However, the visitation rate standardized for morph abundance showed a

significantly lower visitation rate (Kruskal-Wallis chi-square = 3.77; df = 2; $p = 0.01511$) to

the RW morph (mean number of visits per hour ± s.e.; 119.94 ± 10.75), than the common GR

(175.15 ±17.29) and GW morphs (171.39 ± 18.72; Fig. 1d;).

From video recordings of pollinator visitations, we also observed that hawkmoths revisited

the same flowers within an RW morph (~ 10-12 times), once they discovered this rare but

nectar-rich inflorescence. In contrast, the two common morphs GR and GW with low nectar

rewards received lower visitation rates, and pollinators rarely revisited the flowers. Thus

within-inflorescence visits were higher in the rare RW morph compared to the common GR

and GW morphs.

Physiological compatibility and natural fitness vary among the morphs

A total of 471 hand pollinations were carried out (with $n \geq 30$ per treatment) where the

three intra-morph cross-pollination treatments were: GW x GW, GR x GR, and RW x RW

(male x female morph) and will be interchangeably referred to as 'cross-pollination'

treatments. Inter-morph cross-pollination treatments were categorized into three pairs to

investigate the effect of male and female parents in each treatment. These crosses are (male x

female) identified as pair a: GW x GR and GR x GW, pair b: GW x RW and RW x GW, and

pair c: GR x RW and RW x GR. Out of 471 hand-pollination treatments, only 108 treatments

set fruits (Fig. 2a). None of the autogamous treatments set fruits. Selfing treatments failed to

set fruits exclusively in the GR morph, whereas the GW and RW morphs demonstrated leaky

self-compatibility with success rates of 6.67% and 9.68%, respectively. Fruit-set success in

selfed treatments was 5.8%, which was significantly lower ($\chi 2 = 31.299$, $p < 0.0001$; Table

S4) when compared with fruit-set success in the intra-morph cross (48.38%) and inter-morph

cross (58.4%). The GR morph demonstrated the highest fruit set in the intra-morph outcrossed

treatment with a success rate of 46.67%, followed by the RW morph at 27.03% and the GW

morph at 18.75% (Fig. 2a).

In the inter-morph crosses, we noted donor-specific differences in the GW and RW

morphs. The donor specificity was related to whether a morph was used as a male parent or a

female parent in the inter-morph cross-pollination treatments. In crosses where GW morph

was the male parent (i.e., pollen donor), success rates were significantly lower (pair a: GW x

GR, 24.24% and in pair b: GW x RW, 25%) when compared to crosses where it was used as a

female parent, that is pollen recipient (in pair a: GR x GW, 43.33% and in pair b: RW x GW,

44.44%, Fig. 2a, Fisher's exact test $p = 0.027$, odd ratio = 0.419). Whereas when RW acted as

the male parent (pollen donor) in inter-morph crosses, fruit set success was higher (pair b: RW

x GW = 44.44%; pair c: RW x GR = 46.67%) compared to when it served as the female

parent (pair b: GW x RW = 25%; pair c: GR x RW = 37.84%) however, Fisher's exact test did

not yield the significant difference (p = 0.1129, Odd ratio = 1.8).

Among all hand-pollination treatments, self-pollination treatments exhibited the lowest seed

count (mean number of seeds per matured fruit ± s.e.; 8.63 ± 3.15), followed by intra-morph

crosses (16.93 ± 1.8) and then inter-morph crosses (18.63 ± 1.2). In the self-pollination

treatment, the GR morph exhibited a significantly lower seed count (zero) compared to the

GW morph (14.5 ± 6.5) and RW morph (13.33 ± 4.37; Kruskal-Willis $\chi2$ = 5.2499, df = 2, p

= 0.07244; Fig. 2b; Table S5 and S6). In the intra-morph cross-pollination treatment, the seed

counts varied significantly among the morphs. The GW morph exhibited the lowest seed

count (10.72 ± 2.7) compared to the GR morph (20.71 ± 2.8) and the RW morph (15.7 ± 2.7;

Kruskal-Willis $\chi2$ = 5.2499, df = 2, p = 0.07244; Fig. 2b; Table S6). Seed count in inter-

morph cross-pollination treatments was not significantly different among the three pairs (Fig.

2b; Table S6). Finally, the mean total fruit set per plant in open pollination treatments was

recorded to be equal in all three morphs (Fig. 3a). However, the mean seed count per fruit was

significantly different among the morphs (Kruskal-Wallis $\chi2$ = 36.3222, df = 2, p < 0.0001)

with the GR morph being the highest (23.7 ± 1.5), followed by GW (20.3 ± 1.2) and RW

(11.5 ± 1.4; Fig. 3b). We also noted that the seed counts recorded in the open (natural)

pollination and self-pollination treatments of RW were similar (Fig. 2a and 3a).

Associating morph abundance and floral traits

In the path analyses, the five variables resulted in one direct path between abundance and

reproductive fitness (e), seven indirect paths (a to d and f to h), one correlation path between

abundance and type of compatibility (i), and one path on the effect of reproductive fitness on

relative abundance (j; Fig. 4). The Chi-squared goodness of fit test, CFI, and TLI scores

identified a good fit for all four models (*C. caulina*, GR, GW and RW morph; Table 1). In the

case of the GW morph, the model fit as estimated by TLI values was not good (TLI = -0.65),

and we did not find any significant relationships between the traits as well as for the net

effect. We found that both the direct effect (path e in Fig. 4) and the net effect of morph

abundance on reproductive fitness were positive and statistically significant in three (*C.*

caulina, GR morph, and RW morph) of the four path analyses models (Table 1).

In the species-specific model, *C. caulina* in KAS showed a positive and significant effect of

only morph abundance on total nectar reward (a), self-compatibility on reproductive fitness

(d'), and finally, reproductive fitness on relative morph abundance (j). The correlation

between morph abundance and self-compatibility was also noted to be positive (i''). For the

within-morph path analyses, significant effects among indirect effects were observed only in

the GR morph, although positive effects were present across several paths for the other two

morph-specific models (GW and RW). In the GR morph, the following paths were significant

- morph abundance and total nectar reward (a), total nectar reward and reproductive fitness

(g), cross-compatibility and reproductive fitness (d''), and the regression of reproductive

fitness on morph abundance (j). The direct effect between relative morph abundance and

reproductive fitness (e), the regression of the reverse effect, that is, between reproductive

fitness and abundance (j) was significant and positive in RW but negative and not significant

for GW. We failed to detect any significant relationships between the other traits in the RW

morph, although notably, the effect of self-compatibility on reproductive fitness was positive,

whereas the effect of cross-compatibility on reproductive fitness was negative but not

significant (path d; Table 1).

Discussion

Floral color polymorphism, variable morph abundances, and population shifts in morph

frequencies are well-documented across angiosperms. However, only few studies have

explored the breadth of phenotypic landscape of FCP, that is, how floral color covaries with

morphology, physiology, mating systems and ecological interactions, and this is particularly

rare for long-lived, clonal taxa (Elam & Linhart, 1988; Lau & Galloway, 2004; Caruso *et al.*,

2010; Berardi *et al.*, 2016; Koski *et al.*, 2020; Jacquemyn & Brys, 2020). Our study of bract-

color morphs in the nocturnal ginger *Curcuma caulina* illustrates why such breadth matters.

On lateritic plateaus which are known for a strongly seasonal environment, abiotic stressors

are pronounced. We found that morph identity is tied not only to differential nocturnal

visitation by hawkmoths but also to nectar investment and mating-system differences that

together determine realised reproductive outcomes. Interpreting these patterns therefore

requires an integrated view that accounts both for immediate pollinator behaviour and for

longer-term demographic and physiological dynamics characteristic of perennial, clonal

species.

Floral color variation is widespread, and in tropical lineages such as the order Zingiberales

and within the family Zingiberaceae many species exhibit striking floral and bract color

variation (Maas-Van De Kamer et al., 2016; Temeles et al., 2000; Gowda & Kress, 2013;

Leong-Skornicková et al., 2007; Saryan et al., 2020). Previous studies have suggested that specialisation to particular pollinators may drive morph differentiation. At one extreme of this idea lies the 'one pollinator–one morph' scenario (Temeles et al., 2013; Meléndez-Ackerman et al., 2005) that is empirically reported in the Caribbean *Heliconia caribaea* (Temeles et al., 2013) and trans-Himalayan *Roscoea* spp. (Paudel et al., 2019). And the other extreme is, one pollinator-multi morphs and multi pollinator-multi morphs interactions reported in the Caribbean *Heliconia* spp. (Martén-Rodríguez et al. 2011; Gowda et al., 2012; Gowda and Kress, 2013). To understand the role of pollinators in shaping morph frequencies in a polymorphic species we addressed whether pollinator specialisation offers a satisfactory explanation in the case of *C. caulina*. Within the Kaas population all morphs displayed classic moth-pollination syndromes (Johnson *et al.*, 2017), such as nocturnal opening, white colored flowers, and long floral tube with copious nectar. True to its syndrome, we discovered that *C. caulina* is exclusively pollinated by nocturnal hawkmoths and the moths show differential visitation towards the morphs. However, in our study, pollinator identity and visitation patterns alone did not provide a full account of morph-level fitness. Like many classic pollinator-mediated models we expected morphological differences (tube length, flower size, display) to drive pollinator sorting and hence differential fitness (Meléndez Ackerman and Campbell 1998; Gomez 2000; Majetic et al. 2007; Milano et al. 2016); interestingly, in *C. caulina* those morphological traits do not differ among morphs, and instead only nectar investment and mating-system differences were the two most defining axes. Thus we did not see the strict 'one pollinator–one morph' outcome in *C. caulina*. We propose that the trait space through which the hawkmoths discriminate and affect the plant's fitness appears to rely primarily on floral reward rather than due to gross morphological matching.

In *C. caulina*, the first ecological trait that showed direct effect on reproductive fitness is morph abundance (path e, Fig. 4). The rocky lateritic plateaus of Northern Western Ghats are known for their extreme weather conditions suggesting the presence of strong abiotic selection factors (Watve, 2013; Shrotri et al., 2025). Our identification of the tri-modally distributed morphs with distinct physiological features suggests that these morphs may have been under different selection pressures in the past. For example, anthocyanin-rich red bracts of RW morph may confer UV or drought tolerance (Berardi et al., 2016) which may have been advantageous in the rocky laterite habitats like Kaas. Unequal morph abundance has been commonly reported in plants showing FCP. Narbona et al. (2018) noted that more than 80 % of Mediterranean species with FCP exhibit skewed morph distributions that often

fluctuate spatio-temporally due to abiotic pressures (Dormont et al., 2010; Tang et al., 2016).
Thus, we propose that abiotic adaptations, along with vegetative propagation, could decouple
morph frequencies from contemporary selection, allowing relic traits to persist (Barrett, 2015;
Záveská et al., 2012) resulting in a polymorphic population with a few rare but persistent
morphs, as noted in *C. caulina* at Kaas plateau.

Physiological compatibility is the second trait that we identified as affecting the plants
reproductive fitness (path d). While physiological traits are not expected to be different
between morphs, we noted two different reproductive outcomes in two separate morphs of *C.*
caulina - a) improved reproductive fitness via strong self-incompatibility in the most common
GR morph, and b) self-compatibility providing reproductive assurance in the rare RW morph.
The second strategy has been suggested theoretically (Lloyd 1992; Holsinger, 2000) and
empirically in *Ipomea purpurea* (Subramaniam and Rusher, 2000; Fry and Rausher, 2017),
and *Lysimachia arvensis* (Arista et al., 2013; Ortiz et al., 2015). However, our results
demonstrate that contrary to our expectations reproductive strategies can vary significantly
both within a species and within a population.

The third trait that showed a direct effect on reproductive fitness was floral nectar. The
absence of both high nectar reward and higher pollinator visitations translating to higher
reproductive fitness (fruit set and seed set) suggests that the rare morph may have a trade-off
between its nectar investment and observed female reproductive success. This is in contrast to
the pattern noted in *Ipomoea purpurea* where rare, nectar-rich morphs in a population benefit
from higher pollen export (Fry & Rausher, 1997). We speculate that the higher nectar rewards
in the rare RW morph may boost pollen export as noted in *Antirrhinum* sp. (Jones & Reithel
2001) and therefore may be maintained in the population despite the observed trade-off.

So, what factors explain the observed disparity in morph abundances in *C. caulina* at Kaas?
We propose a multi-trait hypothesis involving an interplay between nectar reward, pollinator
attraction, and physiological compatibility (self vs. cross), which collectively influence the
reproductive success of different morphs and subsequently affects their population abundance.
The common morphs show higher reproductive success despite having lower nectar energy
probably because their self-incompatibility ensures that all seeds are outcrossed and with
higher seed-set (fitness). On the other hand, the rare morph may suffer from pollen limitation
due to their lower discovery and high inbreeding by pollinators, they seem to ensure
reproductive success by producing higher quality nectar along with a higher self-compatibility

rate. Since the higher quality nectar resulted in repeat visits by the pollinators most of which

may be self-pollen (Harder *et al.*, 2004; M. Eckhart *et al.*, 2006; Ishii & Harder, 2006),

presence of self-compatibility ensures that there is some seed-set, despite higher selfing. Thus,

the plant's reproductive fitness is correlated with morph abundance via both ecological and

physiological paths suggesting a complex population-scale dynamics that may be responsible

for the observed overall trimodal distribution of the morphs.

Population shifts in morph frequencies are common and have been observed in annual plants.

However, across the last 15 years of our studies at the Kaas plateau (unpublished data,

personal observations by authors) we did not record transitions in morph ratios or elimination

of the rare morphs. This emphasizes the critical role vegetative propagation can play in

maintaining rare morphs within a population. The importance of vegetative propagation has

also been noted in other gingers (Záveská et al. 2012; Skopalíková et al., 2023; Yadav, 2025)

and has been predicted to reduce the intensity of selection otherwise expected in sexually

propagated plants (Jesson and Barrett 2002; Barrett and Harder 2006). Therefore, to

understand factors shaping population structures of rhizomatic plants, genetic contributions

from both sexual and vegetative strategies need to be considered in future studies.

Our study describes morph variability, specifically in reproductive outcomes, and our ongoing

genetic studies are now focused on population genetic structure of the population as well as in

identifying morph-level genotypic differences. Genetic analyses of population structure can

help us partition the contributions of sexual versus vegetative reproduction and will hopefully

aid in distinguishing relic genotypes from recent selection products (Eckert et al., 1999;

Barrett, 2015). We aim to use *C. caulina* as a model species for long-term demographic

studies focused on understanding polymorphism in perennial plants. By carrying out periodic

plant and pollinator census, records of reproductive fitness of the morphs, and climate factors

we hope to identify the transient or stable equilibria of the morph frequencies within and

across populations of these endemic taxa from the lateritic plateaus.

Conclusions

Our findings suggest a pluralistic view of floral polymorphism in perennial plants: pollinator

visitation is a necessary but not sufficient predictor of morph fitness, because nectar

economics, physiological compatibilities, mating differences, and demography jointly shape

reproductive success. By foregrounding a multi-trait, network perspective we provide an

empirical foundation for future examinations (paternity analysis, population genetic studies,

and manipulative experiments) that will be required to separate contemporary selective forces

from the legacies of the past. More broadly, studies of intraspecific polymorphism in long-

lived plants will benefit from routinely measuring both pre- and post-pollination traits and

from explicitly accounting for vegetative reproduction; only then can we robustly assess when

and how color polymorphisms are maintained in nature.

Supplementary Information

Fig. S1 Plant habit, habitat, and key vegetative features of *Curcuma caulina*.

Fig. S2 Variation in bract color, floral reflectance, and morph frequency of *C. caulina*.

Fig. S3 Hawkmoth pollinators of *C. caulina*.

Fig. S4 Circular histogram and circular statistics values of pollinator visitation rates.

Table S1 List of floral and vegetative morphological characters used for the nMDS analysis.

Table S2 Within-group comparisons for nectar energy, volume, and concentration.

Table S3 Within-group comparisons for peak pollinator visitation rates.

Table S4 Results of the chi-square test for fruiting success rates of hand-pollination.

Table S5 Mean and standard error values for seed count for cross- and self-compatibility
treatments.

Table S6 Between groups comparisons for seed counts for hand pollination treatments.

Funding

This work is supported by funds from the Ministry of Human Resource Development

(MHRD) and the National Geographic Society for research funds to VG; Heliconia Society

Internationals-student research grant 2019 to SS and DST for the INSPIRE fellowship

(IF190233) to SS.

Competing interests

The authors have no conflicts of interest to declare.

Author contributions

SS and VG conceived and designed the study. SS conducted the experiments and collected the field data. SS performed the statistical analyses with inputs from VG. SS drafted the manuscript; VG provided conceptual advice and edited the manuscript. All authors gave final approval for publication.

Data availability

The entire dataset and analysis codes used in this study will be made accessible from Dryad or similar online repository after publication of the paper.

Acknowledgements

Authors thank all funding agencies for funds to both authors and IISER Bhopal for infrastructure. We also thank PCCF, Maharashtra Forest Department, DFO of Satara Division (Forest Department), and the Joint Forest Management Committee (JFMC), Kaas, Satara for permitting our fieldwork. We thank Mr. Dilip Shrotri and Azad G for assistance during night work, Amal, Rahul, Viraj, Asawari, Najla, and Sooraj for their assistance during various data collections, including morphometry and Susnata and Sukhraj for assistance during physiological compatibility experiments during the fieldwork. Finally, we thank all fellow TrEE lab members for their valuable advice, discussions, and support.

Literature cited

Arista M, Talavera M, Berjano R, Ortiz PL. 2013. Abiotic factors may explain the geographical distribution of flower color morphs and the maintenance of color polymorphism in the scarlet pimpernel. *The Journal of Ecology* 101: 1613–1622.

Barazani O, Erez T, Ogran A, Hanin N, Barzilai M, Dag A, Shafir S. 2019. Natural variation in flower color and scent in populations of *Eruca sativa* (Brassicaceae) affects pollination behavior of honey bees. *Journal of Insect Science* 19: 6.

Barrett, S. C. (2015). Influences of clonality on plant sexual reproduction. *Proceedings of the*
National Academy of Sciences, *112*(29), 8859-8866.

Benadi, G., & Gegear, R. J. (2018). Adaptive foraging of pollinators can promote pollination
of a rare plant species. *The American Naturalist*, *192*(2), E81-E92.

Berardi AE, Fields PD, Abbate JL, Taylor DR. 2016. Elevational divergence and clinal
variation in floral color and leaf chemistry in *Silene vulgaris*. *American Journal of Botany*
103: 1508–1523.

Berardi AE, Frey F, Denton EM, Wells JH. 2013. Betalain color morphs exhibit differential
growth, defensive ability, and pollen tube growth rates in *Mirabilis jalapa* (Nyctaginaceae).
International Journal of Plant Sciences 174: 1229–1238.

Bigio, L., Lebel, M., & Sapir, Y. (2017). Do different measures of maternal fitness affect
estimation of natural selection on floral traits? A lesson from *Linum pubescens* (Linaceae).
Journal of Plant Ecology, *10*(2), 406-413.

Bolten AB, Feinsinger P, Baker HG, Baker I. 1979. On the calculation of sugar concentration
in flower nectar. *Oecologia* 41: 301–304.

Brunet J, Thairu MW, Henss JM, Link RI, Kluever JA. 2015. The effects of flower, floral
display, and reward sizes on bumblebee foraging behavior when pollen is the reward and
plants are dichogamous. *International Journal of Plant Sciences* 176: 811–819.

Carlson JE, Holsinger KE. 2013. Direct and indirect selection on floral pigmentation by
pollinators and seed predators in a color polymorphic South African shrub. *Oecologia* 171:
905–919.

Caruso C, Scott SL, Wray JC, Walsh CAJ. 2010. Pollinators, herbivores, and the maintenance
of flower color variation: A case study with *Lobelia siphilitica*. *International Journal of Plant*
Sciences 171: 1020–1028.

Dafni, A., Tzohari, H., Ben-Shlomo, R., Vereecken, N. J., & Ne'eman, G. (2020). Flower
colour polymorphism, pollination modes, breeding system and gene flow in *Anemone*
coronaria. *Plants*, *9*(3), 397.

de Jager, M. L., & Ellis, A. G. (2014). Floral polymorphism and the fitness implications of

attracting pollinating and florivorous insects. *Annals of Botany*, *113*(2), 213-222.

Dick, C. A., Buenrostro, J., Butler, T., Carlson, M. L., Kliebenstein, D. J., & Whittall, J. B.

(2011). Arctic mustard flower color polymorphism controlled by petal-specific

downregulation at the threshold of the anthocyanin biosynthetic pathway. *PloS one*, *6*(4),

e18230.

Dixon P. 2003. VEGAN, a package of R functions for community ecology. *Journal of*

vegetation science 14: 927–930.

Dormont L, Delle-Vedove R, Bessière J-M, Hossaert-Mc Key M, Schatz B. 2010. Rare white-

flowered morphs increase the reproductive success of common purple morphs in a food-

deceptive orchid. *The New Phytologist* 185: 300–310.

Dyer, A. G., Jentsch, A., Burd, M., Garcia, J. E., Giejsztowt, J., Camargo, M. G., ... &

Shrestha, M. (2021). Fragmentary blue: resolving the rarity paradox in flower colors.

Frontiers in Plant Science, *11*, 618203.

Eckert, C. G., & Barrett, S. C. (1993). Clonal reproduction and patterns of genotypic diversity

in Decodon verticillatus (Lythraceae). *American Journal of Botany*, *80*(10), 1175-1182.

Eckert, C. G., Dorken, M. E., & Mitchell, S. A. (1999). Loss of sex in clonal populations of a

flowering plant, *Decodon verticillatus* (Lythraceae). *Evolution*, *53*(4), 1079-1092.

Elam DR, Linhart YB. 1988. Pollination and seed production in *Ipomopsis Aggregata*:

Differences among and within flower color morphs. *American Journal of Botany* 75: 1262–

1274.

Ford EB. 1945. Polymorphism. *Biological Reviews of the Cambridge Philosophical Society*

20: 73–88.

Friedman, J. (2020). The evolution of annual and perennial plant life histories: ecological

correlates and genetic mechanisms. *Annual Review of Ecology, Evolution, and Systematics*,

51(1), 461-481.

Fry, J. D., & Rausher, M. D. (1997). Selection on a floral color polymorphism in the tall
morning glory (*Ipomoea purpurea*): transmission success of the alleles through pollen.
Evolution, *51*(1), 66-78.

Fry JD, Rausher MD. 1997. Selection on a floral color polymorphism in the tall morning
glory (*Ipomoea purpurea*): Transmission success of the alleles through pollen. *Evolution* 51:
66–78.

Gegear, R. J., & Laverty, T. M. (2005). Flower constancy in bumblebees: a test of the trait
variability hypothesis. *Animal Behaviour*, *69*(4), 939-949.

Gigord LD, Macnair MR, Smithson A. 2001. Negative frequency-dependent selection
maintains a dramatic flower color polymorphism in the rewardless orchid *Dactylorhiza*
sambucina (L.) Soo. *Proceedings of the National Academy of Sciences, USA* 98: 6253–6255.

Gómez G. 2000. Phenotypic selection and response to selection in *Lobularia maritima*:
importance of direct and correlational components of natural selection. *Journal of*
Evolutionary Biology 13: 689–699.

Gowda, V., Temeles, E. J., & Kress, W. J. (2012). Territorial fidelity to nectar sources by
purple-throated caribs, *Eulampis jugularis*. *The Wilson Journal of Ornithology*, *124*(1), 81-86.

Gowda V, Kress WJ. 2013. A geographic mosaic of plant-pollinator interactions in the
Eastern Caribbean islands. *Biotropica* 45: 224–235.

Goyret J, Pfaff M, Raguso RA, Kelber A. 2008. Why do *Manduca sexta* feed from white
flowers? Innate and learnt color preferences in a hawkmoth. *Die Naturwissenschaften* 95:
569–576.

Hahn M, Brühl CA. 2016. The secret pollinators: an overview of moth pollination with a
focus on Europe and North America. *Arthropod-Plant Interactions* 10: 21–28.

Harder, L. D., & Barrett, S. C. (2006). *Ecology and evolution of flowers*. OUP Oxford.

Harder, L.D., Jordan, C.Y., Gross, W.E. and Routley, M.B., 2004. Beyond floricentrism: the
pollination function of inflorescences. *Plant Species Biology*, *19*(3), pp.137-148.

Harris DJ, Poulsen AD, Frimodt-Møller C, Preston J, Cronk QCB. 2000. Rapid radiation in

Aframomum (Zingiberaceae): Evidence from nuclear ribosomal DNA internal transcribed

spacer (ITS) sequences. *Edinburgh Journal of Botany* 57: 377–395.

Hassa, P., Traiperm, P., & Stewart, A. B. (2020). Pollinator visitation and female reproductive

success in two floral color morphs of *Ipomoea aquatica* (Convolvulaceae). *Plant Systematics*

and Evolution, *306*(6), 88.

Heinrich, B., 1975. Bee flowers: a hypothesis on flower variety and blooming times.

Evolution, 325-334.

Holsinger, K.E., 2000. Reproductive systems and evolution in vascular plants. *Proceedings of*

the National Academy of Sciences, *97*(13), pp.7037-7042.

Imbert E, Wang H, Conchou L, Vincent H, Talavera M, Schatz B. 2014. Positive effect of the

yellow morph on female reproductive success in the flower color polymorphic *Iris lutescens*

(Iridaceae), a deceptive species. *Journal of Evolutionary Biology* 27: 1965–1974.

Irish, E. E., & Nelson, T. (1989). Sex determination in monoecious and dioecious plants. *The*

plant cell, *1*(8), 737.

Ishii H, Harder L. 2006. The size of individual *Delphinium* flowers and the opportunity for

geitonogamous pollination. *Functional Ecology* 20: 1115–1123.

Jacquemyn H, Brys R. 2020. Lack of strong selection pressures maintains wide variation in

floral traits in a food-deceptive orchid. *Annals of Botany* 126: 445–453.

Jesson, L. K., & Barrett, S. C. (2002). Enantiostyly in *Wachendorfia* (Haemodoraceae): the

influence of reproductive systems on the maintenance of the polymorphism. *American*

Journal of Botany, *89*(2), 253-262.

Johnson SD, Moré M, Amorim FW, Haber WA, Frankie GW, Stanley DA, Coccuci AA,

Raguso RA. 2017. The long and the short of it: a global analysis of hawkmoth pollination

niches and interaction networks. *Functional Ecology* 31: 101–115.

Johnson SD, Raguso RA. 2016. The long-tongued hawkmoth pollinator niche for native and

invasive plants in Africa. *Annals of Botany* 117: 25–36.

Jones KN, Reithel JS. 2001. Pollinator-mediated selection on a flower color polymorphism in
experimental populations of *Antirrhinum* (Scrophulariaceae). *American Journal of Botany* 88:
447–454.

Jorgensen, T. H., Petanidou, T., & Andersson, S. (2006). The potential for selection on pollen
colour dimorphisms in *Nigella degenii*: morph-specific differences in pollinator visitation,
fertilisation success and siring ability. *Evolutionary Ecology*, *20*, 291-306.

Keasar, T., Gerchman, Y., & Lev-Yadun, S. (2016). A seven-year study of flower-color
polymorphism in a Mediterranean annual plant. *Basic and Applied Ecology*, *17*(8), 741-750.

Kellenberger RT, Byers KJRP, De Brito Francisco RM, Staedler YM, LaFountain AM,
Schönenberger J, Schiestl FP, Schlüter PM. 2019. Emergence of a floral color polymorphism
by pollinator-mediated overdominance. *Nature Communications* 10: 63.

Koski, M. H., & Galloway, L. F. (2020). Geographic variation in floral color and reflectance
correlates with temperature and colonization history. *Frontiers in plant science*, *11*, 991.

Koski MH, Berardi AE, Galloway LF. 2020. Pollen color morphs take different paths to
fitness. *Journal of Evolutionary Biology* 33: 388–400.

Kovach, W.L., 2011. Oriana–circular statistics for windows, ver. 4. *Kovach Computing*
Services, Pentraeth, Wales, UK.

Lau JA, Galloway LF. 2004. Effects of low-efficiency pollinators on plant fitness and floral
trait evolution in *Campanula americana* (Campanulaceae). *Oecologia* 141: 577–583.

Leong-Skornicková J, Sída O, Jarolímová V, Sabu M, Fér T, Trávnícek P, Suda J. 2007.
Chromosome numbers and genome size variation in Indian species of *Curcuma*
(Zingiberaceae). *Annals of Botany* 100: 505–526.

Lide, D.R. ed., 2004. *CRC handbook of chemistry and physics* (Vol. 85). CRC press.

Liu C-Q, Sun H. 2019. Pollination in *Lilium sargentiae* (Liliaceae) and the first confirmation
of long☐tongued hawkmoths as a pollinator niche in Asia. *Journal of Systematics and*
Evolution 57: 81–88.

Lloyd, D. G. (1992). Self-and cross-fertilization in plants. II. The selection of self-
fertilization. *International journal of plant sciences*, *153*(3, Part 1), 370-380.

M. Eckhart, V., S. Rushing, N., M. Hart, G., & D. Hansen, J. (2006). Frequency–dependent
pollinator foraging in polymorphic *Clarkia xantiana* ssp. *xantiana* populations: implications
for flower colour evolution and pollinator interactions. *Oikos*, *112*(2), 412-421.

M. Eckhart V, S. Rushing N, M. Hart G, D. Hansen J. 2006. Frequency–dependent pollinator
foraging in polymorphic *Clarkia xantiana* ssp. *xantiana* populations: implications for flower
color evolution and pollinator interactions. *Oikos* 112: 412–421.

Maas-Van De Kamer, H., Maas, P. J. M., Wieringa, J. J., & Specht, C. D. (2016). Monograph
of African Costaceae. *Blumea-Biodiversity, Evolution and Biogeography of Plants*, *61*(3),
280-318.

Majetic CJ, Raguso RA, Ashman TL. 2009. The sweet smell of success: floral scent affects
pollinator attraction and seed fitness in *Hesperis matronalis*. *Functional Ecology* 23: 480–487.

Majetic CJ, Raguso RA, Tonsor SJ, Ashman T-L. 2007. Flower color-flower scent
associations in polymorphic *Hesperis matronalis* (Brassicaceae). *Phytochemistry* 68: 865–
874.

Malovrh, K., Bavcon, J., Križman, M., & Ravnjak, B. (2024). Effect of Abiotic Factors on
Nectar Quality and Secretion of Two Early Spring Species, *Galanthus nivalis* L. and
Helleborus niger L. diversity, 16(8), 469.

Martén-Rodríguez, S., John Kress, W., Temeles, E. J., & Meléndez-Ackerman, E. (2011).
Plant–pollinator interactions and floral convergence in two species of *Heliconia* from the
Caribbean Islands. *Oecologia*, *167*(4), 1075-1083.

Meléndez-Ackerman, E. J., Speranza, P., Kress, W. J., Rohena, L., Toledo, E., Cortés, C., ...
& Soltis, D. (2005). Microevolutionary processes inferred from AFLP and morphological
variation in *Heliconia bihai* (Heliconiaceae). *International Journal of Plant Sciences*, *166*(5),
781-794.

Melendez-Ackerman E, Campbell DR. 1998. Adaptive significance of flower color and inter-
trait correlations in an *Ipomopsis* hybrid zone. *Evolution* 52: 1293.

Milano, E. R., Kenney, A. M., & Juenger, T. E. (2016). Adaptive differentiation in floral traits
in the presence of high gene flow in scarlet gilia (*Ipomopsis aggregata*). *Molecular Ecology*,
25(23), 5862-5875.

Mitchell, R. J. (1993). Adaptive significance of *Ipomopsis aggregata* nectar production:
observation and experiment in the field. *Evolution*, *47*(1), 25-35.

Mitchell, R. J. (2001). Path analysis. Design and analysis of ecological experiments.

Mood, J.D., Prince, L.M., Veldkamp, J.F. and Dey, S., 2013. The history and identity of
Boesenbergia longiflora (Zingiberaceae) and descriptions of five related new taxa. *Gardens*
Bulletin Singapore, *65*(1), 47-95.

Mu, J., Li, G., Niklas, K. J., & Sun, S. (2011). Difference in floral traits, pollination, and
reproductive success between white and blue flowers of *Gentiana leucomelaena*
(Gentianaceae) in an alpine meadow. *Arctic, Antarctic, and Alpine Research*, *43*(3), 410-416.

Narbona, E., Wang, H., Ortiz, P.L., Arista, M. and Imbert, E., 2018. Flower color
polymorphism in the Mediterranean Basin: occurrence, maintenance and implications for
speciation. *Plant Biology*, *20*, 8-20.

Ortiz PL, Berjano R, Talavera M, Rodríguez-Zayas L, Arista M. 2015. Flower color
polymorphism in *Lysimachia arvensis*: How is the red morph maintained in Mediterranean
environments? *Perspectives in Plant Ecology, Evolution and Systematics* 17: 142–150.

Paudel, B. R., Kessler, A., Shrestha, M., Zhao, J. L., & Li, Q. J. (2019). Geographic isolation,
pollination syndromes, and pollinator generalization in Himalayan *Roscoea* spp.
(Zingiberaceae). *Ecosphere*, *10*(11), e02943.

Paudel BR, Kessler A, Shrestha M, Zhao JL, Li Q. 2019. Geographic isolation, pollination
syndromes, and pollinator generalization in Himalayan *Roscoea* spp. (Zingiberaceae).
Ecosphere 10: e02943.

Petzold, A., Pfeiffer, T., Jansen, F., Eusemann, P., & Schnittler, M. (2013). Sex ratios and
clonal growth in dioecious *Populus euphratica* Oliv., Xinjiang Prov., Western China. *Trees*,
27, 729-744.

Plos, C., Stelbrink, N., Römermann, C., Knight, T. M., & Hensen, I. (2023). Abiotic

conditions affect nectar properties and flower visitation in four herbaceous plant species.

Flora, 303, 152279.

Poulsen AD, Lock JM. 1997. New species and new records of Zingiberaceae and Costaceae

from Tropical East Africa. *Kew Bulletin* 52: 601–616.

Raju A, Rao S, Ezradanam V, Zafar R, Kalpana P, Kiran P. 2004. The Hawkmoth,

Macroglossum gyrans and its interaction with some plant species at Visakhapatnam. *Zoos'*

print journal 19: 1595–1598.

Rausher, M. D. (2008). Evolutionary transitions in floral color. *International Journal of Plant*

Sciences, *169*(1), 7-21.

R Core Team. (2023). *R: A language and environment for statistical computing* [Software]. R

Foundation for Statistical Computing. https://www.R-project.org/

Ren S, Vitasse Y, Chen X, Peichl M, An S. 2022. Assessing the relative importance of

sunshine, temperature, precipitation, and spring phenology in regulating leaf senescence

timing of herbaceous species in China. *Agricultural and Forest Meteorology* 313: 108770.

Rosseel Y. 2012. Lavaan: An R package for structural equation modeling. *Journal of*

Statistical Software 48: 1–36.

Sapir Y, Gallagher MK, Senden E. 2021. What maintains flower color variation within

populations? *Trends in Ecology & Evolution* 36: 507–519.

Saryan P, Gupta S, Gowda V. 2020. Species complex delimitations in the genus *Hedychium*: a

machine learning approach for cluster discovery. *Applications in Plant Sciences* 8: e11377.

Schemske, D. W., & Bierzychudek, P. (2007). Spatial differentiation for flower color in the

desert annual *Linanthus parryae*: was Wright right? *Evolution*, *61*(11), 2528-2543.

Schiestl, F. P., & Johnson, S. D. (2013). Pollinator-mediated evolution of floral signals.

Trends in ecology & evolution, 28(5), 307-315.

Shrotri, S., Kaur, S., Dhargalkar, R., Najla, P. V., & Gowda, V. (2024). Mass flowering and

flowering asynchrony characterise a seasonal herbaceous community in the Western Ghats.

Biotropica, 57(5) e70080.

Skopalíková, J., Leong☐Škorničková, J., Šída, O., Newman, M., Chumová, Z., Zeisek, V., ...

& Záveská, E. (2023). Ancient hybridization in *Curcuma* (Zingiberaceae)—Accelerator or

brake in lineage diversifications? *The Plant Journal*, *116*(3), 773-785.

Smith GP, Davidowitz G, Raguso RA, Bronstein JL. 2022. Proboscis curling in a pollinator

causes extensive pollen movement and loss. *Ecological Entomology* 47: 234–241.

Smithson, A., & Macnair, M. R. (1997). Negative frequency☐dependent selection by

pollinators on artificial flowers without rewards. *Evolution*, *51*(3), 715-723.

Snell, R., & Aarssen, L. W. (2005). Life history traits in selfing versus outcrossing annuals:

exploring the 'time-limitation' hypothesis for the fitness benefit of self-pollination. *BMC*

ecology, *5*(1), 2.

Stanton, M. L., Snow, A. A., Handel, S. N., & Bereczky, J. (1989). The impact of a

flower☐color polymorphism on mating patterns in experimental populations of wild radish

(*Raphanus raphanistrum* L.). *Evolution*, *43*(2), 335-346.

Stebbins GL. 2019. Chapter VI. Isolation and the Origin of Species. In: Chapter VI. Isolation

and the Origin of Species. *Columbia University Press*, 189–250.

Streinzer M, Roth N, Paulus HF, Spaethe J. 2019. Color preference and spatial distribution of

glaphyrid beetles suggest a key role in the maintenance of the color polymorphism in the

peacock anemone (*Anemone pavonina*, Ranunculaceae) in Northern Greece. *Journal of*

Comparative Physiology A 205: 735–743.

Subramaniam, B., & Rausher, M. D. (2000). Balancing selection on a floral polymorphism.

Evolution, *54*(2), 691-695.

Subramaniam B, Rausher MD. 2000. Balancing selection on a floral polymorphism. *Evolution*

54: 691–695.

Takano, A., Julius, A. and Mohamed, M., 2009. First report of flexistyly in *Plagiostachys*

(Zingiberaceae). *Acta Phytotaxonomica et Geobotanica*, *60*(1), pp.56-59.

Tang X, Liu H, Zhang C, Wang Y, Zhang X, Yi Y. 2016. Flower color polymorphism in

G*eranium nepalense* (Geraniaceae): Adaptation to non-pollinator agents. *Polish Journal of*

Ecology 64: 526–533.

Tang X-X, Huang S-Q. 2010. Fluctuating selection by water level on gynoecium color

polymorphism in an aquatic plant. *Annals of Botany* 106: 843–848.

Temeles, E. J., Pan, I. L., Brennan, J. L., & Horwitt, J. N. (2000). Evidence for ecological

causation of sexual dimorphism in a hummingbird. *Science*, *289*(5478), 441-443.

Temeles EJ, Rah YJ, Andicoechea J, Byanova KL, Giller GSJ, Stolk SB, Kress WJ. 2013.

Pollinator-mediated selection in a specialized hummingbird-*Heliconia* system in the Eastern

Caribbean. *Journal of Evolutionary Biology* 26: 347–356.

Thompson FL, Eckert CG. 2004. Trade-offs between sexual and clonal reproduction in an

aquatic plant: experimental manipulations vs. phenotypic correlations. *Journal of*

Evolutionary Biology 17: 581–592.

Turelli, M., Schemske, D. W., & Bierzychudek, P. (2001). Stable two‐allele polymorphisms

maintained by fluctuating fitnesses and seed banks: protecting the blues in *Linanthus parryae*.

Evolution, *55*(7), 1283-1298.

Twyford AD, Caola AM, Choudhary P, Raina R, Friedman J. 2018. Loss of color

pigmentation is maintained at high frequency in a Monkey flower population. *The American*

Naturalist 191: 135–145.

Vaidya P, McDurmon A, Mattoon E, Keefe M, Carley L, Lee C-R, Bingham R, Anderson JT.

2018. Ecological causes and consequences of flower color polymorphism in a self-pollinating

plant (*Boechera stricta*). *The New Phytologist* 218: 380–392.

Warren J, Mackenzie S. 2001. Why are all color combinations not equally represented as

flower-color polymorphisms? *The New Phytologist* 151: 237–241.

Waser NM, Price MV. 1981. Pollinator choice and stabilizing selection for flower color in

Delphinium nelsonii. *Evolution* 35: 376–390.

Waser NM, Price MV. 1985. The effect of nectar guides on pollinator preference:

experimental studies with a montane herb. *Oecologia* 67: 121–126.

Watve A. 2013. Status review of Rocky plateaus in the northern Western Ghats and Konkan

region of Maharashtra, India with recommendations for conservation and management.

Journal of Threatened Taxa 5: 3935–3962.

Weiss MR. 2001. Vision and learning in some neglected pollinators: beetles, flies, moths, and

butterflies. *Cognitive Ecology of Pollination*: 171–190.

Wenzell KE, Skogen KA, Fant J. 2023. Range□wide floral trait variation reflects shifts in

pollinator assemblages, consistent with pollinator□mediated divergence despite generalized

visitation. *Oikos* e09708.

Wolfe, L. M., & Sellers, S. E. (1997). Polymorphic floral traits in *Linaria canadensis*

(Scrophulariaceae). *American Midland Naturalist*, 134-139.

Wolfe, L. M. (1993). Reproductive consequences of a flower color polymorphism in

Hydrophyllum appendiculatum. *American Midland Naturalist*, 405-408.

Yadav, R., & Gowda, V. (2024). Six new species of *Globba* L.(Zingiberales, Zingiberaceae)

from the Eastern Himalayas and Northeast India. *PhytoKeys*, *246*, 197.

Yadav R. 2025. Reproductive system in the genus *Globba* L. (Zingiberaceae): Understanding

its origin and maintenance. PhD thesis. Indian Institute of Science Education and Research

Bhopal.

Zar, J.H., 1999. *Biostatistical analysis*. Pearson Education India.

Záveská, E., Fér, T., Šída, O., Krak, K., Marhold, K., & Leong-Škorničková, J. (2012).

Phylogeny of *Curcuma* (Zingiberaceae) based on plastid and nuclear sequences: Proposal of

the new subgenus Ecomata. *Taxon*, *61*(4), 747-763.

Tables

Table 1: Results of path analysis for all four models. Bold values represent the statistical

significance. Path IDs are the same as fig. 4.

	For *C.*	GR morph	GW morph	RW morph

		caulina			
Model fit:	Chi-square statistics	0.052	0.037	0.006	0.664
	p-value (Chi-square)	**0.82**	**0.847**	**0.94**	**0.415**
	CFI value	**1**	**1**	1	**1**
	TLI value	**1.332**	**7.138**	-0.65	**1.494**

Regression:	**Path ID**				
Reproductive fitness ~ Relative abundance	e	**2.324**	**8.64**	-3.21	**12.241**
Total nectar reward	g	-0.122	**-1.94**	0.416	-0.63
Pollinator visitation rates	f	0.004	0.048	0.019	0.08
Self-compatibility	d'	**3.077**	NA	1.922	8.672
Cross-compatibility	d''	0.092	**0.428**	0.121	-0.341
Pollinator visitation rates ~ Relative abundance	h	1.058	-8.15	-2.84	-3.509
Total nectar reward	b	0.56	1.221	0.664	-1.327
Self-compatibility ~ Pollinator visitation rates	c'	-0.004	NA	-0.004	< 0.001

Cross-compatibility ~ Pollinator visitation rates	c''	-0.013	-0.038	-0.013	-0.024
Total nectar reward ~ Relative abundance	a	**0.214**	**1.808**	-0.156	-0.112
Relative abundance ~ Reproductive fitness	j	**2.461**	**6.800**	-2.871	**10.942**
Correlation					
Relative abundance ~~ Self-compatibility	i'	**-42.15**	NA	-5.488	-0.303
Relative abundance ~~ Cross-compatibility	i''	214.939	-18.372	35.678	62.978
Net Total Effect		**2.302**	**4.814**	-3.219	**12.045**

Figure legends

Figure 1. Three bract color morphs of *C. caulina*, their nectar rewards energy, and natural

pollinator visitation rates. a. From left, GW, GR, and RW morphs of *C. caulina*, b. The mean

(± s.e.) nectar energy (cal), c. Bar plot represents pollinator visitation rates (mean ± s.e.)

measured for each morph in the observation plots (y-axis) and simulated pollinator visitation

rates (mean ± s.e.) calculated for the population shown by red dots. Different alphabets

indicate significant differences between morphs ($p < 0.05$).

Figure 2. Physiological compatibility between the three floral color morphs of *C. caulina*. a.

Heatmap illustrates the percent success rate (values inside box) from 478 hand-pollination

from 15 treatments. The male morphs (pollen source) are displayed on the x-axis, while the

female morphs (stigma identity) are shown on the y-axis. b. Mean (± s.e.) seed count from the

pollination treatments in the three morphs of *C. caulina*. The male donors (pollen source) are

displayed on the x-axis while the boxes represent female recipients (stigma identity).

Significant differences are shown by different alphabets.

Figure 3. Natural fitness of the three morphs of *C. caulina* measured as A. mean ($\pm$ s.e.)

reproductive success (% success) of open pollination treatments and B. mean ($\pm$ s.e.) seed

count of the successful fruits from those open pollination treatments. Small letters indicate

significant differences between female morphs. Results of the statistical test between morphs

are given in Table S6.

Figure 4. Path diagrams illustrating the effect of morph abundance on reproductive fitness,

analysed a) as species-specific model, b) for the common GR morph and c) for the rare RW

morph. One-headed arrows (paths a—h and j) indicate regression, while the two-headed arrow

(path i) indicates correlation. Thick arrows denote significant effects ($P < 0.05$), with the

strength of regression or correlation specified adjacent to them. Path IDs for all direct and

indirect effects are provided, and their strengths can be referenced in Table 1.

 Figures

Figure 1. Three bract color morphs of *C. caulina*, their nectar rewards energy, and natural
pollinator visitation rates. a. From left, GW, GR, and RW morphs of *C. caulina*, b. The mean
(± s.e.) nectar energy (cal), c. Bar plot represents pollinator visitation rates (mean ± s.e.)
measured for each morph in the observation plots (y-axis) and simulated pollinator visitation
rates (mean ± s.e.) calculated for the population shown by red dots. Different alphabets
indicate significant differences between morphs ($p < 0.05$).

Figure 2. Physiological compatibility between the three floral color morphs of *C. caulina*. a.

Heatmap illustrates the percent success rate (values inside box) from 478 hand-pollination

from 15 treatments. The male morphs (pollen source) are displayed on the x-axis, while the

female morphs (stigma identity) are shown on the y-axis. b. Mean (± s.e.) seed count from the

pollination treatments in the three morphs of *C. caulina*. The male donors (pollen source) are

displayed on the x-axis while the boxes represent female recipients (stigma identity).

Significant differences are shown by different alphabets.

Figure 3. Natural fitness of the three morphs of *C. caulina* measured as A. mean (± s.e.)

reproductive success (% success) of open pollination treatments and B. mean (± s.e.) seed

count of the successful fruits from those open pollination treatments. Small letters indicate

significant differences between female morphs. Results of the statistical test between morphs

are given in Table S6.

Figure 4. Path diagrams illustrating the effect of morph abundance on reproductive fitness,
analysed a) as species-specific model, b) for the common GR morph and c) for the rare RW
morph. One-headed arrows (paths a—h and j) indicate regression, while the two-headed arrow
(path i) indicates correlation. Thick arrows denote significant effects (P < 0.05), with the
strength of regression or correlation specified adjacent to them. Path IDs for all direct and
indirect effects are provided, and their strengths can be referenced in Table 1.