Reconstructing the diversity dynamics of paleo-grasslands using deep learning on superresolution images of fossil Poaceae pollen

Marc-Élie Adaimé[1], Shu Kong[2], Michael A. Urban[1,3], F. Alayne Street-Perrott[4], Dirk Verschuren[5], Surangi W. Punyasena[1]

[1] Department of Plant Biology, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA
[2] Faculty of Science and Technology, University of Macau, Macau 999078, China
[3] Department of Biology, University of New Brunswick, Fredericton, NB, E3B 5A3, Canada
[4] Faculty of Science and Engineering, Swansea University, Singleton Park, Swansea, SA2 8PP, UK
[5] Limnology Unit, Department of Biology, Ghent University, B-9000 Gent, Belgium

Corresponding author: spunya1@illinois.edu

Significance

The pollen of most grass species are morphologically indistinguishable using traditional optical microscopy, but we show that they can be differentiated through deep learning analyses of superresolution images. Abstracted morphological features derived from convolutional neural networks can be used to quantify the biological and physiological diversity of grass pollen assemblages, without *a priori* knowledge of the species present, and used to reconstruct past changes in the taxonomic diversity and relative abundance of C_4 grasses in ancient grasslands. This approach unlocks ecological information that had been previously unattainable from the fossil pollen record and demonstrates that deep learning can solve some of the most intractable identification problems in the reconstruction of past vegetation dynamics.

Abstract

Despite its abundance in the fossil record, grass pollen is largely overlooked as a source of ecological and evolutionary data because most Poaceae species cannot be differentiated using traditional optical microscopy. However, deep learning techniques can quantify the small variations in grass pollen morphology visible under superresolution microscopy. We use the abstracted morphological features output by deep learning to estimate the taxonomic diversity and physiology of fossil grass pollen assemblages. Using a semi-supervised learning strategy, we trained convolutional neural networks (CNNs) on pollen images of 60 widely distributed grass species and unlabeled fossil Poaceae. Semi-supervised learning improved the CNN models' capability to generalize feature recognition in fossil pollen specimens. Our models successfully captured both the taxonomic diversity of an assemblage and morphological differences between C_3 and C_4 species. We applied our trained models to fossil grass pollen assemblages from a 25,000-year lake-sediment record from eastern equatorial Africa and correlated past shifts in grass diversity with atmospheric CO_2 concentration and proxy records of local temperature, precipitation, and fire occurrence. We quantified grass diversity for each time window using morphological variability, calculating both Shannon entropy and morphotype counts from the specimens' CNN features. Reconstructed C_3:C_4 ratios suggest a gradual increase in C_4 grasses with rising temperature and fire activity across the late-glacial to Holocene transition. Our results demonstrate that quantitative machine-learned features of pollen morphology can significantly advance palynological analysis, enabling robust estimation of grass diversity and C_3:C_4 ratio in ancient grassland ecosystems.

Keywords

C_4 photosynthesis, deep learning, diversity estimation, Poaceae, superresolution imaging

Introduction

The evolutionary history of grasses and the ecological history of grasslands are recorded in the fossil pollen record. However, because the pollen of Poaceae species are visually indistinguishable using traditional optical microscopy, this vast paleontological record has been largely limited to estimating changes in past abundance (1, 2). As a result, grass pollen data has contributed little insight into the evolution of the ~11,800 species of this diverse plant lineage (3) or the establishment and paleoecological dynamics of grasslands.

The fossil history of grasses is foundational to understanding the development of modern grasslands and their ecological importance. Grasslands are a geologically recent ecosystem, covering ~40% of Earth's terrestrial surface and contributing to ecosystem services such as soil fertility, carbon storage, and atmospheric cooling (4, 5). Grasslands are characterized by unique ecological properties, including high water-use efficiency and high albedo, and play a critical role in regulating global geochemical cycles, including those of carbon, nitrogen, and silica (4, 6). The productivity and ecological resilience of grasslands is linked to its diversity (7), with high species richness increasing resilience to environmental stress (7, 8).

Grass species employ one of two photosynthetic pathways, C_3 and C_4. C_4 photosynthesis increases water use efficiency and tolerance of low atmospheric CO_2 (9). It evolved independently in multiple grass lineages and allowed these species to eventually dominate the world's grasslands by the late Miocene (10, 11). The global expansion of C_4-dominated grasslands occurred alongside a downward trend in atmospheric CO_2 concentrations during the late Cenozoic and is associated with major environmental changes linked to the growth of ice sheets, such as the expansion of seasonally dry tropical and temperate climates (12, 13).

During the Quaternary (the last 2.5 million years), grassland ecosystems expanded throughout the tropics due to the generally cool and dry climatic conditions that characterized glacial periods in low-latitude regions (14). Pollen records from southern and southeastern Brazil indicate that now forested regions were dominated by grasslands during glacial times (15). Similarly, in Africa, grasslands expanded during the Last Glacial Maximum (LGM) (16). Sensitivity to climate, combined with their

widespread geographic distribution and extensive fossil record, make grasses an

important biological proxy in the reconstruction of past climates and environments.

However, interpreting the fossil record of grasses has been a long-standing

challenge. Poaceae macrofossils can often be difficult to distinguish from other

monocotyledonous plants (12), and grass phytoliths do not directly capture taxonomic

diversity with multiple morphotypes shared between C_3 and C_4 species (11, 17). Pollen

across Poaceae species is morphologically similar, but small variations in the exine

patterning of the pollen grain surface are taxonomically meaningful and can be

quantitatively characterized (2, 18, 19). However, these previous categorizations of

grass morphotypes have been too broad to capture changes in the taxonomic structure

of grassland communities on millennial timescales.

New approaches are therefore needed to effectively reconstruct the diversity and

composition of the grass pollen record. Machine learning has become an important part

of the palynological toolkit over the last decade (e.g., 25–27) but with few exceptions

(23–25), it has been largely used to automate, and not improve upon, pollen

identifications. In this study, we developed deep learning pipelines that use

convolutional neural networks (CNNs) trained on superresolution images (0.04 µm

resolution) of modern and fossil Poaceae pollen (Fig. 1A) with the goal of improving on

expert classifications of grass pollen. The abstracted morphological features derived

from these CNNs capture taxonomic and physiological diversity in grass pollen

assemblages. Our first computational pipeline quantifies taxonomic diversity by

assessing morphological variability within fossil pollen assemblages using Shannon

entropy (Fig. 1B). Our second pipeline uses a random forest classifier (26) to identify

the abstracted morphological features associated with each photosynthetic pathway and

reconstruct changes in the abundance ratio of C_3 and C_4 grass pollen (Fig. 1C).

We applied our trained CNN models to superresolution images of fossil grass

pollen extracted from the 25,000-year sediment record of Lake Rutundu (0.0416° S,

37.4638° E, 3,078 m asl), a subalpine crater lake on Mt. Kenya in eastern equatorial

Africa (27) (Fig. 2). We reconstructed the taxonomic diversity and physiological

composition of the surrounding high-mountain grasslands relative to local environmental

factors such as fire history (28), temperature (29), and precipitation (30), as well as

known atmospheric CO_2 levels (31). The wealth of paleoecological and paleoclimate

data available from previous research on the Lake Rutundu record (27, 28, 32, 33)

allowed direct comparison of our results with prior work on the history of Mt. Kenya's

grasslands. Our findings demonstrate that superresolution imaging and deep learning

enables -- for the first time -- the estimation of grass diversity and C_3:C_4 ratios from fossil

pollen morphology. With these tools, we can interpret the diversity and composition of

grass assemblages even in the absence of species identifications.

Results

CNNs can discriminate among grass species

We trained two separate K-way classification CNNs using superresolution

structured illumination (SR-SIM) (34) images of 1,241 modern and 1,503 fossil pollen

grains (Fig. 1A). Modern pollen was isolated from herbarium specimens of 60 extant

Poaceae species (3 - 37 specimens per species, SI Appendix, Tables S1 and S2).

Fossil material was isolated from a Lake Rutundu sediment core (27) and recently

deposited sediments from five additional Mt Kenya lakes (SI Appendix, Tables S3 and

S4). The first CNN (H-CNN) was trained on maximum intensity projections (MIPs) of the

whole grain. The second CNN (P-CNN) used small cropped segments of the MIP for

training. CNNs are a series of convolutional and pooling layers with a final classification

layer that outputs probabilities of class labels (35). Notably, we employed CNNs to

extract abstracted morphological characters for downstream analyses and not for

classification. Our analysis ignores the final classification layer and instead uses the

penultimate layer containing the highest-level representation of a specimen's

morphology, or "features" (25). However, we assessed the robustness of these CNN

features by calculating the average classification accuracy of our trained models (Fig. 3

and SI Appendix, Fig. S1). The H-CNN and P-CNN achieved an average accuracy of

61.40% and 72.51%, respectively (SI Appendix, Table S5). Fusion of the two CNNs

achieved an average accuracy of 73.95% (SI Appendix, Table S5). Lowest accuracies

were for species with the lowest number of training images (3-4 specimens), but all

accuracies were significantly greater than chance (1.67%).

Morphological variability reflects taxonomic diversity

We used the Shannon entropy (36) of the CNN features of a grass pollen
assemblage as a measure of its taxonomic diversity and tested this assumption by
measuring the morphological entropy of simulated grass communities. We randomly
split our modern Poaceae dataset into 30 species used to train the CNNs and 30
unseen species from which we drew random selections of specimens to form 1000
artificial assemblages of varying species richness and species abundances. For each
artificial assemblage, we calculated Shannon's index (36) based on the number of
species selected and their relative abundances. We then forward-passed the specimens
through our trained CNN models and calculated the assemblages' Shannon entropy
from the distribution of their modified CNN features (see Methods). Linear regression
analysis quantified the strong positive correlation between Shannon entropy and the
Shannon index in our data ($r = 0.972$, $p < 0.0001$) (SI Appendix, Fig. S2). A unit
increase in Shannon entropy corresponded to an equivalent increase in Shannon
diversity, meaning morphological variability was proportional to taxonomic (species)
diversity.

Shifts in grass diversity on Mt Kenya

The Shannon entropy of fossil grass pollen assemblages from the 25,000-year
Lake Rutundu record showed a positive correlation with the evolution of atmospheric
CO_2 (Pearson's $r = 0.548$, $p = 0.006$) and the temperature history of Mt Kenya ($r = $
0.551, $p= 0.008$) over the same period (Fig. 4A and SI Appendix, Fig. S3 and Table S6),
indicating that higher CO_2 concentration and higher local temperatures are associated
with greater morphological variability and taxonomic diversity in these subalpine tropical
grasslands.

Using k-means clustering to translate the entropy values into an estimated
number of distinct morphotypes, we then calculated the Shannon diversity index for
each time window based on the number of morphotypes and their relative abundances
(see Methods). These diversity index values exhibited strong positive correlation with

CO_2 (r = 0.742, p < 0.0001) and temperature (r = 0.783, p < 0.0001) (Fig. 4A and SI

Appendix, Fig. S3 and Table S6). There was no significant relationship, however,

between our time series of either Shannon entropy or Shannon diversity values and

reconstructed changes in local precipitation (Fig. 4A and SI Appendix, Fig. S3 and

Table S6).

Finally, we used the CNN features of our modern pollen dataset and the Rutundu

pollen assemblages as a proxy for changes in taxonomic composition, under the

assumption that differences in the distribution of CNN features reflected differences in

the underlying species composition. We used PCA to reduce the dimensionality of the

dataset and visualize the distribution of CNN features across different time windows.

We found that the morphological diversity of the fossil dataset was a subset of the

diversity present in the modern dataset (SI Appendix, Fig. S4), but the distribution of

features were significantly different, particularly for the first principal component (PC1,

Welch's t-test, p < 0.0001). The morphological space occupied by the fossil

assemblages varied through time, with the space occupied by the four LGM samples

(21,260 - 18,435 cal yr. BP) distinct from that of other time periods along PC1 and

occupying a smaller area in the morphospace defined by PC1 and PC2 (SI Appendix,

Fig. S4).

Pollen morphology captures grass photosynthetic pathway

The phylogenetic PCA (pPCA) of the CNN feature matrix of the 60 modern pollen

samples yielded 59 phylogenetic principal components (pPCs), of which 12 showed a

statistically significant difference between C_3 and C_4 grass taxa (Welch's t-test, p < 0.05)

(SI Appendix, Fig. S5). We identified an optimal subset of five pPCs that discriminated

between the pollen of C_3 and C_4 grasses with an accuracy of 91.25% through recursive

feature elimination (RFE) using a random forest classifier (37, 38) (see Methods). We

then used these five pPCs to train a second random forest classifier to label fossil pollen

grains as C_3 or C_4 and used bootstrapping to calculate confidence intervals for the

estimated C_3:C_4 ratios in fossil samples.

Late Quaternary trends in C$_4$ grass abundance

We projected the standardized CNN features of the fossil specimens onto the

pPCA space defined by the CNN features of the modern grass pollen (see Methods).

This generated a set of five pPCA scores for each fossil pollen grain that was then

forward-passed through the trained random forest classifier for C$_3$/C$_4$ classification. The

percentage of pollen grains within each fossil assemblage that were assigned to C$_4$

grasses ranged between 21.03% and 51.61%, with a slight increase through time over

the past 25,000 years (Fig. 4B). This temporal trend in C$_4$ grass abundance is positively

correlated with past changes in atmospheric CO$_2$ (Pearson's r = 0.437, p = 0.033) and

local biomass burning (Pearson's r = 0.550, p = 0.005 for microcharcoal flux; Pearson's

r = 0.442, p = 0.031 for macrocharcoal flux), suggesting that post-glacial expansion of

C$_4$ grasses in the subalpine zone on Mt Kenya was mediated by increased fire

prevalence.

Our results contrast with a previous reconstruction of C$_4$ grass abundance on Mt

Kenya, inferred from the carbon isotope ratio (δ^{13}C) of fossil grass pollen in the Lake

Rutundu record, that shows a gradual decrease in C$_4$ grass abundance over time (Fig.

4B) (28). However, C$_4$ abundance estimates in this study were highly variable (ranging

between 3% and 100%), with wide confidence intervals indicating a large margin of

error (Fig. 4B). Our results are more consistent with C$_4$ grass estimates based on fossil

grass cuticle preserved in the Lake Rutundu record, which range between 16.32% and

72.67%, and show no consistent trend over the past 25,000 years (Fig. 4B) (27).

Discussion

Deep learning allows us to quantify the sub-micrometer differences in pollen

morphology visible under superresolution microscopy and statistically discriminate

among different morphotypes of grass pollen, opening new opportunities in

paleobotanical research. With these computational tools, we are able to more

consistently discriminate among known species (Fig. 3), capture trends in morphological

diversity that approximate species diversity (Fig. 4A and SI Appendix, Fig. S3), and

capture changes in the relative abundance of C_3 and C_4 grasses through time (Fig. 4B

and SI Appendix, Fig. S3).

Grass diversity increases in East African grasslands

Shannon entropy estimates species diversity without species identifications.

Estimates are most accurate for high-diversity assemblages (SI Appendix, Fig S2).

When applied to the fossil pollen record from Lake Rutundu, we found that grass

diversity in subalpine grasslands on Mt Kenya remained relatively low from 25,000

years ago until the Last Glacial Maximum (LGM) (~20,000 cal. yr BP), after which there

was a gradual increase in taxonomic diversity, as inferred from the variability of CNN

features (Fig. 4A). Changes in grass diversity were significantly correlated with changes

in CO_2 and temperature (SI Appendix, Fig. S3 and Table S6). Converting entropy

measurements into morphotype counts and Shannon diversity index values produced a

more pronounced increase in estimated diversity that significantly correlated with CO_2

levels and temperature (SI Appendix, Fig. S3 and Table S6). These findings suggest

that elevated CO_2 levels, coupled with warmer conditions, favored diversification of

subalpine Poaceae communities around Lake Rutundu following the LGM. This is

consistent with multiple studies that have documented increased biomass productivity

and biodiversity, particularly in grasslands, under conditions of elevated post-glacial

CO_2 levels, as well as in warmer, more fire-disturbed environments (39–43).

Time series of the organic geochemical composition of Lake Rutundu sediments

show that the carbon-nitrogen (C/N) ratio, total organic carbon (TOC), and terrestrial

biomarker fluxes reached some of their lowest recorded levels during the LGM, before

rapidly increasing at ~16,000 cal. yr BP. This suggests terrestrial primary productivity in

the subalpine zone of Mt Kenya was lower during the LGM, when grasses were more

abundant relative to other herbaceous plants such as heather (Ericaceae) (27, 28, 32).

However, our results show that during this period, grass diversity was also at its lowest

point in the last 25,000 years. Notably, as the dominance of Poaceae in the subalpine

zone declined under the warmer conditions of the Holocene, grass diversity increased.

C$_4$ grasses have thinner pollen walls and simpler surface texture

Integrating feature selection with a random forest model allowed us to select the
optimal set of phylogenetic principal components that best predicted photosynthetic
pathway. pPCA allowed us to reduce the influence of relatedness and identify
convergent morphological features. We cannot directly link the 12 principal components
that captured significant differences between C$_3$ and C$_4$ grass pollen (SI Appendix, Fig.
S5) to specific morphological characters, but we found that Shannon entropy and pixel
intensity values were, on average, significantly greater in C$_3$ grass species than C$_4$
grass species (Welch's t-test, $p = 0.029$ and $p = 0.041$ respectively) (Fig. 5). Higher
entropy suggests more complex surface texture and increased pixel brightness
suggests a thicker exine wall. We tested the hypothesis that the pollen grains of C$_3$
grasses have thicker exine walls using direct measurements of exine thickness from
(19). In this alternate set of Poaceae species, exine thickness was also significantly
greater in C$_3$ taxa (Welch's t-test, $p = 0.001$) (Fig. 5).

Morphological differences between pollen from C$_3$ and C$_4$ grasses may be
directly linked to their physiological characteristics. C$_4$ grasses tend to achieve faster
growth by producing less dense leaves (higher leaf area to mass ratio), enabling the
production of more leaves at the same carbon cost (44). C$_4$ grasses also invest more in
their root systems (44). The systematic difference in resource allocation between C$_3$ and
C$_4$ grasses may explain why physiological differences are also encoded in pollen
morphology, independent of phylogenetic relatedness. Grass pollen morphology may
reflect the evolutionary trade-off between investment in roots and shoots. We
hypothesize that C$_4$ grasses may invest less in pollen development and structure,
producing grains with thinner exines and less complex surface patterning.

Changing prevalence of C$_3$ and C$_4$ grasses in East African grasslands

Our analysis of fossil grass pollen from Lake Rutundu showed that the proportion
of C$_4$ grasses changed through time, with a slight increase from ~18,000 to 5,000 cal. yr
BP (Fig. 4B). The increase positively correlated with both local fire prevalence (inferred
from micro- and macrocharcoal fluxes) and atmospheric CO_2 (SI Appendix, Fig. S3 and

Table S6) and is consistent with research that identified increased fire activity as a key

factor in the expansion of C_4-dominated grasslands in Africa during the Late Miocene

(10, 45, 46). Likewise, while full-glacial atmospheric CO_2 concentrations (~180-200

ppm) are limiting for both C_3 and C_4 grasses, the increase in CO_2 concentrations from

glacial to postglacial levels provided favorable conditions for an increase in biomass

production, and may have been more advantageous for C_4 species under certain

environmental conditions, such as water stress (47). Our results reflect this trend,

though C_3 grasses remained dominant overall (Fig. 4B). The increased proportion of C_4

grasses we reconstructed across the glacial-to-Holocene transition may also be linked

to warmer conditions in the Holocene. However, the modern-day occurrence of cold-

tolerant panicoid C_4 grasses in alpine grasslands (48, 49) complicates this

interpretation. Our results did not show significant correlation between the proportion of

C_4 grasses and precipitation (Fig. 4A and SI Appendix Fig. S3 and Table S6). However,

increased biomass burning may have been the result of higher precipitation in eastern

equatorial Africa during the Late-Glacial period and early Holocene (~14,300-12,300

and 11,400-9,600 cal. yr BP), as evidenced by more depleted δD values leaf-wax

alkanes (52) (Fig. 4C), as well as depleted $δ^{13}C$ values of organic matter and water-

level changes in East African lakes and swamps (50).

Our estimates of C_4 grass abundance are consistent with previous estimates

from Lake Rutundu fossil cuticle (27) (Fig. 4B). From ~25,000 to 18,000 cal. yr BP, our

results align with the most conservative estimates of C_4 abundance (based on cuticle

fragments identified with certainty as C_4 grass species) (27). From ~15,000 cal. yr BP to

present, our data align with the median trend, which includes both certain and likely C_4

taxa (27). Our results do not match estimates of C_4 grass abundance based on pollen

carbon isotope measurements before ~10,000 cal. yr BP, but align with estimates after

~10,000 cal. yr BP (28). Our underestimation of C_4 grass abundance compared to these

other proxies, particularly before the Holocene, may stem in part from differences

between the species composition of fossil assemblages from Mt. Kenya and our modern

training dataset. If we use the diversity of CNN features as stand-in for taxonomic

composition, we see that while there is substantial overlap in the morphological space

occupied by our modern and fossil datasets, this range of morphological diversity is

significantly different from that of modern grass pollen (SI Appendix, Fig. S4A). This is

especially true for the range of morphospace where the LGM pollen assemblages fall

(SI Appendix, Fig. S4B). This may have introduced uncertainty in our random forest

estimates and explain some of the mismatch between our time series and those of other

proxies for grass abundance.

Deep-time implications

Deep learning can be a powerful tool for revealing shifts in grass diversity and

composition. It allows us to quantify and use the morphological data captured in fossil

pollen in new and unexpected ways. By analyzing the long-term morphological

variability within pollen assemblages, we can test hypotheses about the evolution and

physiological adaptations of grasses over thousands to millions of years and provide a

deeper understanding of how grasses adapted to changing climates and environments

throughout the Cenozoic.

Grasslands provide crucial habitat for many plant and animal species, driving key

evolutionary adaptations (51). The first putative Poaceae appeared in the Early

Cretaceous (113-101 Mya) (46–48), and grasslands were present on every continent

except Antarctica by the middle Eocene (~38 Mya) (12). Although our study focused on

Late Quaternary grasslands, this same approach can be applied to fossil grass pollen

assemblages further back in the geologic record, providing us deeper insight into the

evolution of this important global biome.

Methods

Pollen samples and paleoclimate data

Pollen samples of 60 extant grass species were isolated from herbarium material

from the Missouri Botanical Gardens (USA), University of Illinois (USA), and Harvard

(USA) or purchased from Sigma-Aldrich (St. Louis, MO, USA) (SI Appendix, Table S1).

These are predominately, but not exclusively, widely distributed species that occur in

East Africa, including the important African cereals sorghum (*Sorghum halepense*),

finger millet (*Eleusine coracana*), and maize (*Zea mays*) (SI Appendix, Table S2).

Thirty-six of these 60 species are C_3 grasses and 24 are C_4. Twenty-four samples of

fossil pollen (SI Appendix, Table S4) were isolated from a 7.55-m sediment sequence

collected from Lake Rutundu in 1996 (27). Rutundu is a 40-ha oligotrophic lake, 3078 m

above sea level (a.s.l.) on the northeast side of Mt. Kenya, just above the current

treeline (Fig. 2) (27). Recently deposited pollen was extracted from the surface

sediments of five additional small lakes on Mt. Kenya distributed along an altitudinal

transect between 1820 and 4585 m a.s.l. (Fig. 2 and SI Appendix, Table S3) (55). All

pollen samples were prepared following (56, 57). (Details provided in the SI Appendix.)

The ages of the fossil samples are derived from the age model for the Rutundu

sediment sequence based on AMS ^{14}C dating of bulk organic carbon (27, 28).

Paleoclimate data for the subalpine zone of Mt. Kenya were taken from published

studies on the Rutundu sediment record. The temperature reconstruction is based on

abundance ratios of preserved glycerol dialkyl glycerol tetraethers (GDGTs) (29) and

the precipitation reconstruction is based on the hydrogen-isotope signature of alkanes in

fossil leaf waxes from terrestrial vegetation (30). For fire history, we used data on

microscopic and macroscopic charcoal influx to Lake Rutundu as a proxy for past

variation in local biomass burning and fire prevalence (28). We used the composite

record of atmospheric CO_2 concentration based on air-bubble measurements in several

Antarctic ice cores (31).

SR-SIM imaging

Pollen specimens were imaged using a Zeiss Elyra S1 superresolution structured

illumination (SR-SIM) microscope. SR-SIM is an optical microscopy method capable of

capturing morphological features <150 nm in length, below the diffraction limit of light

(34). We used a 63x Plan Apochromat (1.4 NA) oil objective and an excitation

wavelength of 561 nm. Image resolution was 0.0397 µm/pixel. Each axial plane of the

Z-stack was constructed with five grid rotations, spaced 0.18 µm apart, with 28-108

planes per grain. Images were cropped to the perimeter of each pollen grain.

Deep learning architectures and dataset setup

We developed two separate K-way classification CNNs based on the ResNeXt-

101 architecture (58): a holistic image CNN (H-CNN) trained on maximum intensity

projections (MIP) of entire image stacks, and a patch CNN (P-CNN) trained on square

crops of the MIP, each covering ~10% of the image (25) (Fig. 1A). Both CNNs were

trained using images of modern reference pollen (labeled data) and images of fossil

pollen (unlabeled data) in a semi-supervised learning (SSL) framework (59). SSL

assigns pseudo-labels to unlabeled images and treats them as labeled in training

classification models. The labeled reference data were divided into training (70%) and

validation (30%) sets for model training and model selection, respectively. Unlabeled

images were assigned pseudo-labels by the model being trained, and images that had

pseudo-label confidence scores greater ≥95% were retained for training. The validation

set of reference images was used for hyperparameter tuning and model selection. SSL

improved the model's ability to generalize across both modern and fossil data. Training

details are included in the SI Appendix.

The H-CNN generated a single *K*-dimensional logit vector for each pollen

specimen, while the P-CNN produced multiple logit vectors corresponding to multiple

input patches per specimen (Fig. 1A). The mean of these vectors by P-CNN was

calculated and normalized using softmax (25). For each specimen, we fused the outputs

of the two CNNs by multiplying the two *K*-dimensional classification probability vectors

and normalizing it to unit length (25).

Feature extraction

Our analysis relied on the features characterized by the two CNNs. We

concatenated the 2048-dimensional feature vectors derived from the penultimate (global

pooling) layer of the H-CNN and P-CNN to produce a 4096-dimensional feature vector

for each specimen (Fig. 1A). This combined feature vector represented a high-level

abstraction of the pollen image that served as input for our morphological analyses of

diversity and physiology. To visualize and compare the morphological features of the

modern grass pollen dataset and the 24 fossil pollen assemblages, we stacked their

classification features into a single standardized dataset and performed a PCA on the

combined feature matrix (Fig S4).

Morphological measures of diversity

We used simulated communities to test our hypothesis that the morphological

diversity within a grass pollen assemblage can serve as a proxy for taxonomic (species)

diversity. From our modern pollen dataset of 60 grass species, we randomly selected 30

species for training our CNNs and used the remaining 30 species (572 specimens) as

an independent test set. From the test set, we randomly selected a random number of

species (between 2 and 30) and a random number of specimens per species (between

1 and the maximum number of available specimens per species). We quantified the

diversity of each simulated community using Shannon's index:

$$H' = -\sum_{i=1}^{S} p_i \log(p_i)$$

where S is the total number of species and p_i is the relative frequency of the i^{th} species.

We next measured morphological diversity of each community using its CNN-

derived features. We standardized the feature matrix and applied PCA. We then

computed Shannon entropy (36) over the probability distribution of the first principal

component (PC1). The distribution was estimated using kernel density estimation (KDE)

(60). The estimated probability density for a given PC1 score, z_i, is calculated as:

$$p(z_i) = \hat{f}(z_i) = \frac{1}{n \times h} \sum_{j=1}^{n} K\left(\frac{z_i - z_j}{h}\right)$$

436   Where $\hat{f}$ is the estimated density calculated using KDE, n is the total number of

437   specimens per sample, h is the bandwidth, and K is the kernel function. We computed

438   entropy H from the density values:

$$H = -\sum_{i=1}^{n} p(z_i) \log(p(z_i))$$

We calculated Shannon entropy and the Shannon diversity index for 1000 simulations

and measured their correlation using linear regression, reporting the Pearson

correlation coefficient (r) and p-value.

Fossil grass diversity estimation and paleoenvironmental correlates

We calculated the Shannon entropy values for the CNN-derived morphological

features of the 24 fossil Rutundu samples. We then used k-means clustering to

translate the entropy values into the number of distinct morphotypes in each time

window. We set the number of morphotypes by selecting the Shannon diversity index

values that best correlated with Shannon entropy values using Pearson's correlation

coefficient (Fig. 1B).

We report both the Shannon entropy and Shannon index values and their

correlation to five paleoenvironmental variables: atmospheric CO_2 level, local

reconstructed temperature and precipitation, and micro- and macrocharcoal influx as

proxy for fire prevalence. Our results include Pearson's correlation coefficient (r) and its

corresponding p-value for each individual regression (SI Appendix, Fig. S3 and Table

S6).

Grass pollen morphology and photosynthetic pathway

We conducted a phylogenetic PCA (pPCA) of the CNN-derived feature matrix

using the phyl.pca function of the package phytools (61, 62) (Fig. 1C). We used a

Bayesian time-calibrated grass phylogeny based on chloroplast DNA sequences

covering approximately 90% of existing grass genera (54) as our input tree. To identify

the pPCs most predictive of the photosynthetic pathway, we implemented a feature

selection process using recursive feature elimination (RFE) with a random forest

classifier, using the functions rfe and randomForest of the 'caret' and 'randomForest'

packages in R (26, 38, 63). Photosynthetic pathway was treated as a categorical

variable. We established a control for RFE using repeated cross-validation with five

folds, repeated 10 times.

Using the selected pPCs, we then trained a final random forest model to classify

the photosynthetic pathway based on the modern data's pPCA scores, using 1000

decision trees to reduce potential overfitting (26). To quantify the uncertainty in our

estimates, we employed a bootstrapping procedure with 1000 resamples. For each

bootstrap sample, we calculated the mean proportion (percent abundance) of C_4

grasses and computed the 95% confidence intervals around the mean estimates for

each time window in the Rutundu record.

Estimates of past C_4 abundance and paleoenvironmental correlates

We standardized the fossil dataset using the mean and standard deviation of the

modern data and projected the standardized fossil data onto the pPCA space computed

using the modern data. We then extracted the optimal pPCs determined using RFE

(Fig. 1C). We used the final random forest model, trained on modern data, to determine

the photosynthetic pathway (C_3 or C_4) for each fossil specimen based on the selected

pPCA components.

We report the correlation between our time series of estimated C_4 abundance

and those of the five paleoenvironmental variables: atmospheric CO_2 concentration,

local temperature, local precipitation, and micro- and macro-charcoal influx. We

calculated Pearson's correlation coefficient (r) and its corresponding p-value for each

individual regression (SI Appendix, Fig. S3 and Table S6).

Quantifying morphological differences between C_3 and C_4 pollen

We used entropy to estimate the complexity of the grass pollen surface texture,

with higher values indicating more random and complex textural patterning. We

calculated Shannon entropy by segmenting the pollen grain image and quantifying the

isolated grain's information content (complexity) based on the distribution of its pixel

intensities (64). Because pollen exine is autofluorescent, pixel intensity allows us to infer

variations in thickness of the pollen wall, with brighter images suggesting a thicker wall

(34). We used direct exine thickness measurements (19) for 35 C_3 and 23 C_4 grass

species to confirm this relationship. We calculated the median of entropy, pixel intensity,

and exine thickness measurements across all specimens of each species in our modern

dataset and used Welch's t-test to assess the statistical significance of observed

differences.

Acknowledgments

504 Imaging costs and postdoctoral support for M.A.U. were provided by NSF-DBI –

505 Advances in Bioinformatics (NSF-DBI-1262561) to S.W.P. S.K. was supported in part

506 by the University of Macau (SRG2023-00044-FST). M.-E.A. was supported in part by

507 the University of Illinois Tom L. Phillips Memorial Fund for Paleobotany. We thank Tim

508 Gallaher for referring us to his recently published grass phylogeny.

Author Contributions

511 M.-E.A., S.K., and S.W.P. designed the research. M.-E.A. conducted the computational

512 analyses and experiments. M.A.U. conducted the imaging, sample preparation, and lab

513 analyses. F.A.S.-P. and D.V. provided samples and collected data. S.W.P. and S.K.

514 supervised the research. M.-E.A., S.W.P., S.K., and M.A.U. wrote the manuscript, with

515 feedback from all authors.

Data Availability

518 Superresolution structured illumination images used in this study were submitted to the

519 Illinois Databank (DOI: 10.13012/B2IDB-1391373_V1) (65). The code base is publicly

520 available on Github (DOI: 10.5281/zenodo.13756262) (66), and can be accessed using

521 the following link: https://github.com/paleopollen/Pollen_Diversity_Dynamics

References

1. M. B. Bush, On the interpretation of fossil Poaceae pollen in the lowland humid neotropics. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **177**, 5–17 (2002).

2. L. Mander, S. W. Punyasena, Grass pollen surface ornamentation: A review of morphotypes and taxonomic utility. *J. Micropalaeontology* **35**, 121–124 (2016).

3. Grass Phylogeny Working Group III, *et al.*, Nuclear phylogenomics of grasses (Poaceae) supports current classification and reveals repeated reticulation. [Preprint] (2024). Available at: http://biorxiv.org/lookup/doi/10.1101/2024.05.28.596153 [Accessed 16 August 2024].

4. J. S. Petermann, O. Y. Buzhdygan, Grassland biodiversity. *Curr. Biol.* **31**, R1195–R1201 (2021).

5. G. J. Retallack, Cenozoic expansion of grasslands and climatic cooling. *J. Geol.* **109**, 407–426 (2001).

6. S. E. Melzer, *et al.*, Fire and grazing impacts on silica production and storage in grass dominated ecosystems. *Biogeochemistry* **97**, 263–278 (2010).

7. D. Tilman, J. A. Downing, Biodiversity and stability in grasslands. **367** (1994).

8. A. Lüscher, *et al.*, Using plant diversity to reduce vulnerability and increase drought resilience of permanent and sown productive grasslands. *Grass Forage Sci.* **77**, 235–246 (2022).

9. R. F. Sage, The evolution of C_4 photosynthesis. *New Phytol.* **161**, 341–370 (2004).

10. S. Hoetzel, L. Dupont, E. Schefuß, F. Rommerskirchen, G. Wefer, The role of fire in Miocene to Pliocene C_4 grassland and ecosystem evolution. *Nat. Geosci.* **6**, 1027–1030 (2013).

11. C. A. E. Strömberg, R. E. Dunn, C. Crifò, E. B. Harris, "Phytoliths in paleoecology: Analytical considerations, current use, and future directions" in *Methods in Paleoecology: Reconstructing Cenozoic Terrestrial Environments and Ecological Communities*, D. A. Croft, D. F. Su, S. W. Simpson, Eds. (Springer International Publishing, 2018), pp. 235–287.

12. C. A. E. Strömberg, Evolution of grasses and grassland ecosystems. *Annu. Rev. Earth Planet. Sci.* **39**, 517–544 (2011).

13. R. Sage, T. Sage, F. Kocacinar, Photorespiration and the evolution of C_4 photosynthesis. *Annu. Rev. Plant Biol.* **63**, 19–47 (2012).

14. L. 't Mannetje, Climate change and grasslands through the ages: An overview. *Grass Forage Sci.* **62**, 113–117 (2007).

15. H. Behling, South and southeast Brazilian grasslands during Late Quaternary times: A synthesis. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **177**, 19–27 (2002).

16. L. Scott, Grassland development under glacial and interglacial conditions in southern
Africa: Review of pollen, phytolith and isotope evidence. *Palaeogeogr. Palaeoclimatol.*
Palaeoecol. **177**, 47–57 (2002).

17. P. C. Twiss, "Predicted world distribution of C_3 and C_4 grass phytoliths" in *Phytolith*
Systematics: Emerging Issues, G. Rapp, S. C. Mulholland, Eds. (Springer US, 1992), pp.
113–128.

18. L. Mander, M. Li, W. Mio, C. C. Fowlkes, S. W. Punyasena, Classification of grass pollen
through the quantitative analysis of surface ornamentation and texture. *Proc. Biol. Sci.* **280**,
20131905 (2013).

19. C. Wei, P. E. Jardine, W. D. Gosling, C. Hoorn, Is Poaceae pollen size a useful proxy in
palaeoecological studies? New insights from a Poaceae pollen morphological study in the
Amazon. *Rev. Palaeobot. Palynol.* **308**, 104790 (2023).

20. E. C. Stillman, J. R. Flenley, The needs and prospects for automation in palynology. *Quat.*
Sci. Rev. **15**, 1–5 (1996).

21. K. Holt, G. Allen, R. Hodgson, S. Marsland, J. Flenley, Progress towards an automated
trainable pollen location and classifier system for use in the palynology laboratory. *Rev.*
Palaeobot. Palynol. **167**, 175–183 (2011).

22. C. M. Barnes, *et al.*, Deductive automated pollen classification in environmental samples
via exploratory deep learning and imaging flow cytometry. *New Phytol.* **240**, 1305–1326
(2023).

23. S. W. Punyasena, D. K. Tcheng, C. Wesseln, P. G. Mueller, Classifying black and white
spruce pollen using layered machine learning. *New Phytol.* **196**, 937–944 (2012).

24. I. C. Romero, *et al.*, Improving the taxonomy of fossil pollen using convolutional neural
networks and superresolution microscopy. *Proc. Natl. Acad. Sci.* **117**, 28496–28505
(2020).

25. M.-É. Adaimé, S. Kong, S. W. Punyasena, Deep learning approaches to the phylogenetic
placement of extinct pollen morphotypes. *PNAS Nexus* **3**, pgad419 (2024).

26. L. Breiman, Random forests. *Mach. Learn.* **45**, 5–32 (2001).

27. K. J. Ficken, M. J. Wooller, D. L. Swain, F. A. Street-Perrott, G. Eglinton, Reconstruction of
a subalpine grass-dominated ecosystem, Lake Rutundu, Mount Kenya: A novel multi-proxy
approach. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **177**, 137–149 (2002).

28. M. A. Urban, D. M. Nelson, F. A. Street-Perrott, D. Verschuren, F. S. Hu, A Late-
Quaternary perspective on atmospheric $p\mathrm{CO_2}$, climate, and fire as drivers of C_4-grass
abundance. *Ecology* **96**, 642–653 (2015).

29. S. E. Loomis, *et al.*, The tropical lapse rate steepened during the Last Glacial Maximum.
Sci. Adv. **3**, e1600815 (2017).

30. S. Garelick, J. M. Russell, S. Dee, D. Verschuren, D. O. Olago, Atmospheric controls on

precipitation isotopes and hydroclimate in high-elevation regions in Eastern Africa since
the Last Glacial Maximum. *Earth Planet. Sci. Lett.* **567**, 116984 (2021).

31.   B. Bereiter, *et al.*, Revision of the EPICA Dome C CO_2 record from 800 to 600 kyr before
present. *Geophys. Res. Lett.* **42**, 542–549 (2015).

32.   M. Wooller, *et al.*, Late Quaternary vegetation changes around Lake Rutundu, Mount
Kenya, East Africa: Evidence from grass cuticles, pollen and stable carbon isotopes. *J.*
Quat. Sci. **18**, 3–15 (2003).

33.   F. A. Street-Perrott, *et al.*, Towards an understanding of late Quaternary variations in the
continental biogeochemical cycle of silicon: Multi-isotope and sediment-flux data for Lake
Rutundu, Mt Kenya, East Africa, since 38 ka BP. *J. Quat. Sci.* **23**, 375–387 (2008).

34.   M. Sivaguru, *et al.*, Comparative performance of airyscan and structured illumination
superresolution microscopy in the study of the surface texture and 3D shape of pollen.
Microsc. Res. Tech. **81**, 101–114 (2018).

35.   Y. Lecun, L. Bottou, Y. Bengio, P. Haffner, Gradient-based learning applied to document
recognition. *Proc. IEEE* **86**, 2278–2324 (1998).

36.   C. E. Shannon, A mathematical theory of communication. *Bell Syst. Tech. J.* **27**, 379–423
(1948).

37.   I. Guyon, J. Weston, S. Barnhill, V. Vapnik, Gene selection for cancer classification using
support vector machines. *Mach. Learn.* **46**, 389–422 (2002).

38.   M. Kuhn, Building predictive models in R using the caret package. *J. Stat. Softw.* **28**, 1–26
(2008).

39.   R. F. Sage, Was low atmospheric CO_2 during the Pleistocene a limiting factor for the origin
of agriculture? *Glob. Change Biol.* **1**, 93–106 (1995).

40.   W. Jiang, Y. Cheng, X. Yang, S. Yang, Chinese Loess Plateau vegetation since the Last
Glacial Maximum and its implications for vegetation restoration. *J. Appl. Ecol.* **50**, 440–448
(2013).

41.   J. Cunniff, M. Charles, G. Jones, C. P. Osborne, Reduced plant water status under sub-
ambient $p$$CO_2$ limits plant productivity in the wild progenitors of C_3 and C_4 cereals. *Ann.*
Bot. **118**, 1163–1173 (2016).

42.   Y.-F. Yao, *et al.*, Pollen-based reconstruction of vegetational and climatic change over the
past ~30 ka at Shudu Lake in the Hengduan Mountains of Yunnan, southwestern China.
PLOS ONE **12**, e0171967 (2017).

43.   T. He, B. B. Lamont, J. G. Pausas, Fire as a key driver of Earth's biodiversity. *Biol. Rev.*
94, 1983–2010 (2019).

44.   R. R. L. Atkinson, *et al.*, C_4 photosynthesis boosts growth by altering physiology, allocation
and size. *Nat. Plants* **2**, 16038 (2016).

45.   J. E. Keeley, P. W. Rundel, Fire and the Miocene expansion of C_4 grasslands. *Ecol. Lett.*
632   **8**, 683–690 (2005).

633   46.   C. P. Osborne, D. J. Beerling, Nature's green revolution: The remarkable evolutionary rise
634   of C_4 plants. *Philos. Trans. R. Soc. B Biol. Sci.* **361**, 173–194 (2006).

635   47.   J. Cunniff, M. Charles, G. Jones, C. P. Osborne, Was low atmospheric CO_2 a limiting
636   factor in the origin of agriculture? *Environ. Archaeol.* **15**, 113–123 (2010).

637   48.   S. P. Long, C_4 photosynthesis at low temperatures. *Plant Cell Environ.* **6**, 345–363 (1983).

638   49.   M. J. Wooller, D. L. Swain, F. A. Street-Perrott, S. Mathai, A. D. Q. Agnew, An altitudinal
639   and stable carbon isotope survey of C_3 and C_4 graminoids on Mount Kenya. *J. East Afr.*
640   *Nat. Hist.* **90**, 69–85 (2001).

641   50.   F. A. Street-Perrott, *et al.*, Late Quaternary changes in ecosystems and carbon cycling on
642   Mt. Kenya, East Africa: A landscape-ecological perspective based on multi-proxy lake-
643   sediment influxes. *Quat. Sci. Rev.* **26**, 1838–1860 (2007).

644   51.   O. Toljagić, K. L. Voje, M. Matschiner, L. H. Liow, T. F. Hansen, Millions of years behind:
645   Slow adaptation of ruminants to grasslands. *Syst. Biol.* **67**, 145–157 (2018).

646   52.   E. A. Kellogg, "Fossil record and dates of diversification" in *Flowering Plants. Monocots:*
647   *Poaceae*, E. A. Kellogg, Ed. (Springer International Publishing, 2015), pp. 103–107.

648   53.   Y. Wu, H.-L. You, X.-Q. Li, Dinosaur-associated Poaceae epidermis and phytoliths from
649   the Early Cretaceous of China. *Natl. Sci. Rev.* **5**, 721–727 (2018).

650   54.   T. J. Gallaher, *et al.*, Grasses through space and time: An overview of the biogeographical
651   and macroevolutionary history of Poaceae. *J. Syst. Evol.* **60**, 522–569 (2022).

652   55.   F. A. Street-Perrott, K. J. Ficken, Y. Huang, G. Eglinton, Late Quaternary changes in
653   carbon cycling on Mt. Kenya, East Africa: An overview of the $\delta^{13}C$ record in lacustrine
654   organic matter. *Quat. Sci. Rev.* **23**, 861–879 (2004).

655   56.   K. Fægri, J. Iversen, P. E. Kaland, K. Krzywinski, *Textbook of Pollen Analysis*, ed. 4th
656   (John Wiley & Sons, 1989).

657   57.   M. A. Urban, I. C. Romero, M. Sivaguru, S. W. Punyasena, Nested cell strainers: An
658   alternative method of preparing palynomorphs and charcoal. *Rev. Palaeobot. Palynol.* **253**,
659   101–109 (2018).

660   58.   S. Xie, R. Girshick, P. Dollar, Z. Tu, K. He, Aggregated residual transformations for deep
661   neural networks in *2017 IEEE Conference on Computer Vision and Pattern Recognition*
662   *(CVPR)*, (IEEE, 2017), pp. 5987–5995.

663   59.   K. Sohn, *et al.*, FixMatch: Simplifying semi-supervised learning with consistency and
664   confidence in *Advances in Neural Information Processing Systems*, (Curran Associates,
665   Inc., 2020), pp. 596–608.

60.   B. W. Silverman, Density Estimation for Statistics and Data Analysis Chapter 1 and 2.

61. L. J. Revell, Size-correction and principal components for interspecific comparative
studies. *Evolution* **63**, 3258–3268 (2009).

62. L. J. Revell, phytools: An R package for phylogenetic comparative biology (and other
things). *Methods Ecol. Evol.* **3**, 217–223 (2012).

63. A. Liaw, M. Wiener, Classification and regression by randomForest. **2** (2002).

64. R. M. Haralick, K. Shanmugam, I. Dinstein, Textural features for image classification. *IEEE*
Trans. Syst. Man Cybern. **SMC-3**, 610–621 (1973).

65. S. W. Punyasena, M. A. Urban, M.-É. Adaimé, I. C. Romero, SR-SIM superresolution
structured illumination microscopy images of fossil and modern pollen of Poaceae.
Champaign, IL, USA: University of Illinois Urbana-Champaign. doi: 10.13012/B2IDB-
1391373_V1.

66. M.-É. Adaimé, paleopollen/Pollen_Diversity_Dynamics: Poaceae Pollen Diversity
Dynamics. (2024). doi: 10.5281/zenodo.13756262. Deposited 13 September 2024.

Figures

Fig. 1. Flowchart illustrating the pipeline of computational analyses performed in this study. Two representations of superresolution micrographs of a pollen specimen (the maximum intensity projection of the image stack and crops of the projection) are passed to convolutional neural networks (CNNs) trained to recognize the holistic morphology (H-CNN) and surface texture of patches (P-CNN) of grass pollen. The feature values extracted by both CNNs are concatenated to form a comprehensive feature vector representing each specimen. These vectors are then stacked across all N specimens in the sample, forming a feature matrix of all the concatenated feature vectors. In our assessments of grass diversity, the PCA scores of the feature matrix are used to

estimate probability distributions for pollen assemblages using kernel density
estimation. Shannon entropy is calculated over these probability distributions to provide
a quantitative diversity estimate. In our assessments of photosynthetic pathway, we
apply a phylogenetic PCA (pPCA) on the feature matrix and pass the most informative
phylogenetic principal components (selected based on the modern data) to a random
forest classifier trained to discriminate among C_3 and C_4 grasses.

Fig. 2. Location of the study area. The left panel illustrates the distribution of major vegetation biomes across Africa, with Mt. Kenya highlighted by a red square. The right panel shows the elevation gradient in the Mt. Kenya region with the locations of our six sources of fossil grass pollen: the 25,000-year sediment record of Lake Rutundu, and the five additional lakes from which we extracted pollen from recently deposited surface sediments (Lake Nkunga, Rumuiku Swamp, Lake Ellis, Small Hall Tarn, and Simba Tarn).

Fig. 3. Confusion matrix depicting the average accuracy of the fused CNN classification
model for modern pollen grains from 60 Poaceae species. Rows represent the true
taxa, with the number of specimens analyzed per species in parentheses. Columns
represent the model's predictions, with the classification accuracy for each species in
parentheses. Individual splits are shown in SI Appendix, Fig. S1.

Fig. 4. Inferred changes in grass diversity and proportion of C_4 grasses in subalpine

grasslands of Mt Kenya over the past 25,000 years, plotted alongside time series of

paleoenvironmental proxies interpolated using moving average with a window size of 4.

(A) Grass diversity as measured by Shannon entropy (dashed green line) and a derived

Shannon index. (B) Proportion of C_4 grasses determined by this study (dashed green

line) compared to results based on the carbon-isotopic signature ($\delta^{13}C$) of individual

grass pollen grains (solid blue line, from (28)) and grass cuticle identification (solid

purple line, from (27)). (C) The ice-core record of atmospheric CO_2 concentration (31)

plotted with reconstructions of temperature and precipitation on Mt. Kenya derived from,

respectively, bacterial membrane lipids and the hydrogen-isotopic signature (δD_{precip}) of

plant leaf-wax alkanes extracted from Lake Rutundu sediments (29, 30). More depleted

δD_{precip} values indicate wetter conditions. (D) History of fire prevalence on Mt. Kenya

reconstructed from estimates of biomass burning based on sedimentary fluxes of micro-

and macrocharcoal (28).

Fig. 5. Box plots showing the median difference in Shannon entropy and pixel intensity

(brightness) between pollen from 35 C_3 and 25 C_4 grass species in our modern

reference dataset and in exine thickness (μm) of 35 C_3 and 23 C_4 grass species (from

(19)). On average, C_3 grass pollen has a significantly thicker exine wall and more

complex surface patterning than C_4 grass pollen.