Deep learning of fossil pollen morphology reveals 25,000 years of

ecological change in East African grasslands

Marc-Élie Adaimé[1], Shu Kong[2], Michael A. Urban[1,3], F. Alayne Street-Perrott[4], Dirk

Verschuren[5], Surangi W. Punyasena[1]

[1] Department of Plant Biology, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

[2] Faculty of Science and Technology, University of Macau, Macau 999078, China

[3] Department of Biology, University of New Brunswick, Fredericton, NB, E3B 5A3, Canada

[4] Faculty of Science and Engineering, Swansea University, Singleton Park, Swansea, SA2 8PP,

UK

[5] Limnology Unit, Department of Biology, Ghent University, B-9000 Gent, Belgium

Corresponding author: spunya1@illinois.edu

Significance

Although the pollen of most grass species are morphologically indistinguishable using

traditional optical microscopy, we show that they can be differentiated through deep

learning analyses of superresolution images. Abstracted morphological features derived

from convolutional neural networks can be used to quantify the biological and

physiological diversity of grass pollen assemblages, without *a priori* knowledge of the

species present, and used to reconstruct past changes in the taxonomic diversity and

relative abundance of C_4 grasses in ancient grasslands. This approach unlocks

ecological information previously unattainable from the fossil pollen record and

demonstrates that deep learning can solve some of the most intractable identification

problems in the reconstruction of past vegetation dynamics.

Abstract

Grass pollen is largely overlooked in investigating grassland evolution because the pollen of most species cannot be differentiated using traditional optical microscopy. However, deep learning can quantify small variations in pollen morphology visible under superresolution microscopy. We use the abstract features output by deep learning to estimate the taxonomic diversity and physiology of fossil grass pollen assemblages. Using a semi-supervised learning strategy, we trained convolutional neural networks (CNNs) on superresolution pollen images of modern grasses and unlabeled fossil Poaceae. Our models captured features that reflected both the taxonomic diversity of grass communities along an elevational gradient and morphological differences between C_3 and C_4 species. We applied our trained models to fossil grass pollen assemblages from a 25,000-year lake-sediment record from eastern equatorial Africa (Mt. Kenya) and correlated past shifts in grass diversity with atmospheric CO_2 concentration and proxy records of local temperature, precipitation, and fire occurrence. We quantified changes in grass diversity using morphological variability of fossil pollen assemblages, approximated by the Shannon entropy of CNN features. Our data show that grassland species diversity was strongly reduced between 21,500 and 16,000 years ago, coincident with most severe regional cooling during the last ice age. C_3:C_4 ratios reconstructed using a gradient-boosted decision tree classifier infer a gradual decrease in C_4 grasses since the late-glacial to Holocene transition, associated with decreasing fire activity and elevated temperatures. Our results demonstrate that CNN features of pollen morphology can advance palynological analysis, enabling robust estimation of grass diversity and C_3:C_4 ratio in ancient grassland ecosystems.

Keywords

C_4 photosynthesis, deep learning, diversity estimation, Poaceae, Quaternary, superresolution imaging

Introduction

The evolutionary and ecological history of grasses and grasslands is recorded in the global fossil pollen record. However, despite its abundance, grass pollen is underutilized in paleoecological analyses because the pollen of Poaceae species tends to be visually indistinguishable using traditional optical microscopy. This vast paleontological record has been largely limited to estimating past changes in grass abundance (1, 2). As a result, the fossil record of grass pollen has contributed little insight into the evolution of the ~11,800 species of this diverse plant lineage (3) or the establishment and community dynamics of grassland ecosystems.

Grasslands are a relatively recent ecosystem in Earth history, first appearing in the Eocene-Oligocene (4). Today they cover ~40% of Earth's terrestrial surface and contribute substantially to ecosystem services such as soil fertility, carbon storage, and biosphere control of atmospheric greenhouse gas concentrations (5, 6). Grasslands are characterized by unique ecological properties, including high water-use efficiency and high albedo, and play a critical role in regulating the global biogeochemical cycles of nitrogen, silica, and carbon (5, 7). The productivity and ecological resilience of grasslands is positively correlated with their diversity (8), with high species richness promoting resilience to environmental stressors (8, 9).

Grass species employ one of two photosynthetic pathways, C_3 and C_4. C_4 photosynthesis uses water more efficiently and is more tolerant of low atmospheric CO_2 than the ancestral C_3 pathway (10). C_4 photosynthesis evolved independently in multiple grass lineages, allowing these species to eventually dominate the world's grasslands by the late Miocene (ca. 8 million years ago) (4, 11, 12). The global expansion of C_4 grasslands during the late Cenozoic occurred alongside a downward trend in atmospheric CO_2 concentration, associated with major environmental changes such as the growth of continental ice sheets and the increased prevalence of seasonally dry tropical and temperate climate regimes (4, 13). During the Quaternary period (the last 2.5 million years), grassland ecosystems expanded throughout the tropics due to the generally cool and dry climatic conditions characterizing Pleistocene glacial periods in low-latitude regions (14). Pollen records from southern and southeastern Brazil indicate

that now forested regions were dominated by grasslands during glacial times (15).

Similarly, in Africa, grasslands expanded during the Last Glacial Maximum (LGM) (16).

Sensitivity to climate, combined with their widespread geographic distribution and

extensive fossil record, make grasses an important paleobiological proxy in the

reconstruction of past climates and environments (17, 18). However, interpreting the

fossil record of grasses has been a long-standing challenge. Fossil Poaceae leaves and

stems are rarely preserved with enough morphological detail to distinguish them from

other monocotyledonous plants (4). Fossil phytoliths have generally provided the most

information about the evolution of open grasslands in contemporary paleoecological

research (4, 12). However, multiple phytolith morphotypes are shared between C_3 and

C_4 species, so phytoliths alone cannot directly capture taxonomic and physiological

diversity (12, 19). Multiproxy approaches combining pollen, starch grains, and phytoliths

have advanced identification of fossil grasses in archeobotanical contexts (20), and

Fourier-transform infrared (FTIR) microspectroscopy can classify modern Poaceae

pollen based on their chemical spectra (21). However, starch grain analysis and FTIR

are not generally applicable to community-level or deep-time (≥2 Ma) paleoecology.

Furthermore, starch and phytolith evidence are context- and tissue-dependent, while

FTIR classifications require specific sample preparation, rely on sporopollenin chemical

spectra that can be altered by sedimentation processes, and have thus far been applied

to only a limited number of taxa (21).

Fossil grass pollen has been largely ignored as a source of paleoecological

information because grass pollen grains are morphologically similar across the family.

All have a single annulated pore and a psilate (visually smooth) outer exine surface.

However, small variations in the patterning of the pollen grain surface visible under

electron microscopy have been shown to be taxonomically meaningful and can be

characterized quantitatively (22, 23, 2, 24–26). While pollen size alone is insufficient to

tell apart grasses, there are also quantifiable differences in size among species (27).

However, all of these previous categorizations of grass pollen morphotypes have been

too broad to capture changes in the taxonomic and physiological composition of

grassland communities on millennial or longer timescales.

New approaches are therefore needed to effectively reconstruct the diversity and
composition of fossil grasslands. Here we propose that morphological differences
among the pollen of different grass species revealed in superresolution microscopic
images, though subtle, can provide the basis for training deep learning models. Machine
learning has become an important component of the palynological toolkit over the last
decade (e.g., 28–30), but with few exceptions (31–34) it has been used to automate,
rather than to improve upon, pollen identifications. In this study, we developed deep
learning pipelines that use convolutional neural networks (CNNs) trained on
superresolution images (0.04 µm resolution) of modern and fossil Poaceae pollen (Fig.
1A), with the goal of improving on expert classifications of grass pollen. The abstract
features derived from these CNNs quantify morphological differences that capture the
taxonomic and physiological diversity of grass pollen assemblages. Our first
computational pipeline quantifies the taxonomic diversity in fossil pollen assemblages
by using Shannon entropy to approximate morphological variability (Fig. 1B). Our
second pipeline uses a gradient-boosted decision tree classifier (35, 36) to identify the
variation within the CNN-derived features of modern Poaceae pollen that are most
strongly associated with either C_3 or C_4 grass species.
We then applied our trained CNN models to superresolution images of fossil
grass pollen assemblages extracted from 24 time windows in the 25,000-year sediment
record of Lake Rutundu (0.0416° S, 37.4638° E), a crater lake located at 3078 m
elevation in the heathland-moorland ('subalpine') zone of Mt. Kenya in eastern
equatorial Africa (37) (Fig. 2). We reconstructed the taxonomic diversity and C_4 fraction
of the high-mountain grasslands surrounding Lake Rutundu relative to changes in local
environmental factors such as fire regime (38), temperature (39), and precipitation (40),
as well as the known evolution of atmospheric CO_2 concentration (41). The wealth of
paleoecological and paleoclimate data available from previous research on the Lake
Rutundu record (37, 38, 42, 43) allowed direct comparison of our results with earlier
reconstructions of the history of grasslands on Mt. Kenya. Our findings demonstrate that
the combination of superresolution imaging and deep learning enables – for the first
time – the estimation of past grass diversity and C_3:C_4 ratios from fossil pollen

morphology. With these tools, we can interpret the diversity and composition of ancient
grass communities even in the absence of species identifications.
146

Results

CNNs can discriminate the pollen of different grass species

We trained two separate K-way classification CNNs in a semi-supervised learning
framework, using superresolution structured illumination (SR-SIM) (44) images of 1,241
modern pollen grains identified to species and 1,503 unlabeled fossil pollen grains.
Modern pollen grains were isolated from herbarium specimens of 60 extant Poaceae
species (3–37 specimens per species; SI Appendix, Tables S1 and S2). Fossil material
was isolated from the principal Lake Rutundu sediment core (37) and from recently
deposited sediments in five other lakes on Mt. Kenya (SI Appendix, Tables S3 and S4).
The first CNN (holistic CNN; H-CNN) was trained on maximum intensity projections
(MIPs) of whole grains. The second CNN (patch CNN; P-CNN) used small, cropped
segments of the MIP for training (Fig. 1A). CNNs are a series of convolutional and
pooling layers with a final classification layer that produces probabilities of class labels
(45). Notably, we employed CNNs to extract abstracted morphological characters for
downstream analyses and not for classification. Our analysis ignores the final
classification layer and instead uses the penultimate layer containing the highest-level
representation of a specimen's morphology, or "features" (33). However, we assessed
the quality of these CNN features by calculating the average classification accuracy of
our trained models (SI Appendix, Fig. S1). The H-CNN and P-CNN achieved an
average accuracy of 61.40% and 72.51%, respectively (SI Appendix, Table S5). Fusion
of the two CNNs achieved an average accuracy of 73.95% (SI Appendix, Table S5). All
but three species had ≥19 specimens for training. *Andropogon chrysostachyus*, with
only 3 specimens for training, was never identified correctly (i.e., 0% accuracy). All other
species had average accuracies >23%, which is significantly greater than chance
(1.67%).

Morphological variation within and among grass pollen is taxonomically

meaningful and reflects species diversity

The CNN-derived morphological features of grass pollen show clear taxonomic

structure. Species belonging to the same genus cluster significantly closer together in

CNN feature space than species from different genera ($p = 0.0001$), indicating that the

learned features capture biologically meaningful variation rather than overfitting to

individual species (SI Appendix, Fig. S2A). We also found that morphological variability

within species is not significantly affected by whether the grains came from a single or

from multiple individuals. The mean within-species variance did not differ between these

two groups ($p = 0.498$), suggesting that intra-individual and inter-individual

morphological variability are broadly similar (SI Appendix, Fig. S2B).

We next used artificial pollen assemblages (i.e., simulated grass communities) to

test whether the Shannon entropy (46) of the CNN features of a fossil grass pollen

assemblage captured its taxonomic diversity. We randomly split our modern Poaceae

dataset into 30 species used to train the CNNs and 30 "unseen" species. We randomly

drew 50 grains from the 30 "unseen" species 100 times to form 100 artificial pollen

assemblages of varying species richness and relative abundance. We forward-passed

these grains through our trained CNN models and calculated the assemblages'

Shannon entropy from the distribution of their CNN features (see Methods). We used

PCA to reduce the dimensionality of our CNN feature matrix, producing three PC

subspaces of Shannon entropy identified as PC1, PC1-PC2, and PC1-PC2-PC3. There

was a strong positive correlation between the Shannon entropy of the CNN features and

the Shannon diversity index (46) of the artificial pollen assemblages ($r = 0.47, 0.58, 0.66$

for each respective PCA subspace; $p < 0.0001$) (SI Appendix, Fig. S3). The Shannon

index is a standard statistical measure of diversity that accounts for both species

richness and evenness (47). These results demonstrate that the Shannon entropy of

CNN features effectively captures variation in species diversity.

Modern lake sediment samples capture modern grass diversity gradients

We used grass pollen isolated from recently deposited sediments of six Mt. Kenya lakes situated between 1820 and 4585 m elevation (equivalent to a 16 °C difference in mean annual temperature [MAT]; ref. 48) to test whether the Shannon entropy of CNN features captures modern elevational gradients in grass diversity. Using the first principal component (PC1) of our CNN feature matrix as the diversity metric, we evaluated variation in diversity across a set of kernel bandwidth multipliers (Fig. S4). Leave-one-out cross-validation (LOO-CV) (49) identified $h^* \approx 3.87$, $h^* \approx 5.99$, and $h^* \approx 6.24$ as the optimal bandwidths for the PC1, PC1-PC2, and PC1-PC2-PC3 subspaces. At smoothing (h) values below the optimal bandwidth, the PC1 subspace recovered a mid-elevational (~3000–3500 m a.s.l.) peak in diversity with lower diversity both at lower and higher elevations (SI Appendix, Fig. S4). This closely matches data from vegetation surveys, which indicate that maximum grass species richness on Mt. Kenya (9–11 species) occurs in the heathland-moorland zone between 3100 and 3400 m altitude (where present-day MAT is *ca.* 8 °C; ref. 38), with lower richness both in low-elevation savanna woodland (1–4 species) and in the upper alpine zone above 4000 m (1–5 species) (50). With greater smoothing (where $h = \alpha h^*, \alpha = \sqrt{2}$), between-site differences in PC1 are reduced, resulting in weak expression of the mid-elevation diversity maximum (Fig. S4).

Changes in grass diversity on Mt. Kenya through time

We applied the same approach (using the PCA scores of the Shannon entropy of the CNN features of a pollen assemblage as a proxy for grass diversity) to fossil grass pollen assemblages isolated from 24 more or less equally spaced time windows in the 25,000-year Lake Rutundu sediment record. We calculated entropy for each PCA subspace at their optimal bandwidths (h^*), as derived from the modern-day elevation gradient in grass species diversity (Fig. S4), and compared the long-term temporal trends in entropy (i.e., the inferred past grass diversity) with time series of environmental variables potentially forcing these trends (Fig. 3C-D).

We find that grass diversity in the subalpine grasslands on Mt. Kenya was
strongly reduced during the LGM and earliest late-glacial period (*ca.* 21,500–16,000
years ago), broadly coeval with the period of most strongly reduced MAT in eastern
equatorial Africa (51) and at high elevations on Mt. Kenya in particular (7–8 °C cooler
than today; ref. 38) (Fig. 3A, C). Grass diversity has been substantially higher from *ca.*
15,000 years ago until the present, and was also high under the climate regime
immediately before the LGM when glacial cooling was less extreme (4–6 °C cooler than
today; ref. 38). Entropy values for all three PCA subspaces fluctuate throughout the
Holocene period but with no sustained trend (Fig. 3A), suggesting no or limited change
in the diversity of Mt. Kenya grasslands at the elevation of Lake Rutundu during this
time period. Over the 25,000-year record as a whole, entropy values for subspaces PC1
and PC1-PC2 consistently and positively correlated with both changes in atmospheric
CO_2 and local temperature, although the strength of the relationships varied. For PC1, r
= 0.67 for CO_2 and 0.68 for temperature ($p < 0.001$); for PC1-PC2, r = 0.47 for CO_2 and
0.48 for temperature ($p < 0.05$) (Fig. 3A and SI Appendix, Fig. S5). Entropy also
correlated significantly with fire regime proxies (micro- and macrocharcoal influx) in the
morphological subspace represented by PC1-PC2 (r = 0.45 for microcharcoal and 0.44
for macrocharcoal, $p < 0.05$), while they were weak and not significant for PC1. No
diversity estimates correlated with reconstructed precipitation and no significant
relationships were detected in the PC1-PC2-PC3 subspace (Fig. 3A and SI Appendix,
Fig. S5).

The CNN features of each Rutundu fossil grass pollen assemblage also capture
the morphological disparity among the assemblages' grass pollen grains. This disparity
reflects the underlying species composition. Therefore, changes in the morphological
space occupied by the CNN features likely represent changes in the taxonomic
composition of grass communities on Mt. Kenya through time. We used PCA to reduce
the dimensionality of the dataset and to visualize the distribution of CNN features in
each time window. We found that the combined morphological diversity of all 24 fossil
samples is contained within the diversity present in our reference dataset of 60 modern
grass species (Fig. 4), suggesting that our reference data capture the majority of the
morphological disparity within our fossil dataset. However, the mean distribution of CNN

features in the fossil dataset is significantly different from that in the modern dataset for
PC1 (Welch's t-test, p < 0.0001). Moreover, the morphological space occupied by the
fossil assemblages varies through time. For example, the four assemblages dated to
the LGM (21,260–18,435 cal yr. BP) are distinct from those of other time windows in
that they occupy a distinctly smaller area in the morphospace defined by PC1 and PC2
(Fig. 4).

Grass pollen morphology captures photosynthetic pathway

To determine whether there are consistent morphological differences between
the pollen of C_3 and C_4 grass species, we averaged the CNN features of our modern
grass pollen dataset by species and used PCA to reduce its dimensionality. The first
two principal components significantly discriminated between the pollen of C_3 and C_4
species (Welch's t-test, p = 0.0075 for PC1 and p < 0.001 for PC2) (Figs. 4, 5D).
However, tests for phylogenetic signal revealed strong evolutionary structure in both
PC1 (λ = 0.854, p < 0.001) and PC2 (λ = 0.738, p < 0.001), indicating that
morphological traits may also be strongly influenced by evolutionary relatedness. When
corrected for phylogenetic relatedness using Phylogenetic Generalized Least Squares
(PGLS) regression (see Methods), the difference in mean PC1 values between C_3 and
C_4 species remains significant (p = 0.014, λ = 0.906), but the difference in mean PC2
values does not (p = 0.470, λ = 0.706).
While we cannot link variation in PC1 to specific pollen morphological differences
between C_3 and C_4 grasses, the median pixel brightness of specimen images was found
to be higher in C_3 grasses (Fig. 5A; Welch's t-test, p = 0.029), which suggests that they
have a thicker exine wall, on average, than the pollen of C_4 grasses. The entropy of the
distribution of pixel values was also higher in C_3 grass pollen (Fig. 5B; Welch's t-test, p
= 0.041), suggesting a more complex surface texture (note that this measure is distinct
from our calculation of Shannon entropy used for diversity estimation). Using actual
measurements of exine thickness in an independent dataset of 35 C_3 and 23 C_4 grass
species assembled by (24), we tested the hypothesis that pollen grains of C_3 grasses

have thicker exine walls and found this to be the case (Fig. 5C; Welch's t-test, p =
0.001).

Late Quaternary changes in C_4 grass fraction on Mt. Kenya

Given the morphological differences between pollen of living C_3 and C_4 grasses,
we finally attempted to reconstruct changes through time in the fraction of C_4 grasses
on Mt. Kenya as registered in the Lake Rutundu record. We first trained a C_3 versus C_4
classifier on the CNN features of all 1,241 individual modern grass pollen grains. We
used PCA to reduce the dimensionality of the feature matrix and applied a gradient-
boosted decision tree classifier to the first 230 principal components, which together
represent 95% of the explained variance in pollen morphology. The classifier achieved
high prediction accuracy in distinguishing C_3 and C_4 grains, confirmed by cross-
validation ($\mu = 91.1\%$, $\sigma = 2.5\%$).

We next projected the CNN features of the 1,204 fossil pollen grains from Lake
Rutundu onto the PCA space defined by the CNN features of the modern grass pollen
(see Methods). We forward-passed the first 230 principal components through the
trained gradient-boosted decision tree classifier to determine the proportion of pollen
derived from C_4 grasses in each of the 24 fossil assemblages of the Rutundu record.
The C_4 fraction ranged between 22–68%, with a general increase from ~40% *ca.* 25,000
years ago to 50–60% during the LGM (with the exception of one outlying value of 28%
ca. 19,500 years ago), followed by an overall decreasing trend from the late-glacial
period *ca.* 15,000–13,000 years ago until the late-Holocene and present (Fig. 3B).
These results are consistent with estimates of C_4 grass fraction inferred from
taxonomically diagnostic grass cuticle fragments preserved in the same Lake Rutundu
record, which range between 16–73% and show a stable fraction of 50–60% C_4 grasses
prior to and during the LGM, followed by a general decline over the past 15,000 years,
albeit punctuated by large fluctuations (Fig. 3B) (37). They are less in agreement with a
reconstruction of C_4 grass abundance based on the carbon isotope ratio ($\delta^{13}C$) of fossil
grass pollen from Lake Rutundu, which also shows a long-term decline in the proportion
of C_4 grasses during the postglacial period, but infers values of 75–100% before and
during the LGM (Fig. 3B) (38). Nevertheless, pairwise correlations between these three

C_4 data series, interpolated using varying degrees of spline smoothing, are all highly

significant ($p < 0.0001$) (SI Appendix, Fig. S6), indicating that they all infer a broadly

similar long-term trend, namely the general decline of C_4 grasses relative to C_3 grasses

during the postglacial period on Mt. Kenya. This reconstructed temporal trend in C_4

grass abundance on Mt. Kenya is negatively related to past changes in atmospheric

CO_2 and temperature (Fig. 3C) and positively related to fire prevalence (Fig. 3D),

although neither of these correlations is significant ($p > 0.05$ in all cases).

Discussion

Deep learning techniques have allowed us to quantify the sub-micrometer scale

differences in pollen morphology made visible by superresolution microscopy, and to

statistically differentiate different morphotypes of grass (Poaceae) pollen. With these

computational tools, we demonstrate the capability to more consistently discriminate

among the pollen of known grass species (SI Appendix, Fig. S1), infer the diversity

(species richness) of grass communities from pollen morphological diversity (Fig. 3A

and SI Appendix, Figs. S3–S5), and capture past changes in the relative abundance of

C_3 and C_4 grasses (Fig. 3B and SI Appendix, Fig. S5). In addition to opening important

new opportunities in paleobotanical research, our results also support the important

assumption in palynology that the morphological disparity among pollen grains from a

single plant specimen is smaller than that between species (52) and provides a

meaningful representation of intraspecific pollen variability. Indeed, the variation in CNN

features of pollen from multiple plants within a single species was found to not differ

significantly from that of a single plant, and pollen from species within the same genus

clustered closer together in feature space than pollen from more distantly related

species. This supports our conclusion that the learned morphological features of grass

pollen captured biologically meaningful, taxonomically structured variation and were not

likely overfit to the class-specific patterns of individual plants.

Peak glacial cooling reduced grass diversity on Mt. Kenya

Shannon entropy, estimated from CNN-derived features, provides a proxy for
grass diversity without relying on taxonomic identifications (SI Appendix, Fig. S3).
Applied to the fossil pollen record from Lake Rutundu, it shows that grass diversity
around 3000 m elevation on Mt. Kenya was substantially lower than today during the
period of most severe cooling during the LGM and earliest late-Glacial between 21,500
and 16,000 years ago (38, 51), and relatively stable at more or less their modern-day
diversity throughout the Holocene (Fig. 3A). Notably, the magnitude of entropy-inferred
diversity loss under peak LGM cooling (MAT 7–8 °C below its modern value) is
quantitatively consistent with the low entropy values (low species richness) of modern-
day grasslands in the upper alpine zone of Mt. Kenya *ca.* 1000 m above Lake Rutundu
(ref. 50; Fig. S4), where MAT is ~6 °C lower (38). This local decrease in grass diversity
is likely the result of the depression of the snowline (the 0° C MAT isotherm) and
movement of underlying vegetation zones to lower altitudes during Quaternary glacial
episodes (45, 46), so that the species-rich heather-moorland grassland around Lake
Rutundu was replaced by a species-poor alpine grassland during cooler periods.
Additionally, the relatively abrupt drop in entropy (inferred grass diversity) at the onset of
peak LGM conditions *ca.* 21,500 years ago, and similarly rapid recovery *ca.* 16,000
years ago (Fig. 3A) suggest a threshold response of Mt. Kenya grasslands involving
reduced local viability of many grasses when glacial cooling exceeded >6 °C relative to
present-day MAT. Across the PC1 and PC1-PC2 subspaces of CNN-inferred
morphological variation, changes in entropy through time are positively correlated with
shifts in atmospheric CO_2 and temperature (Fig. 3C and SI Appendix, Fig. S5),
indicating that elevated CO_2 levels and/or warmer habitat conditions favor more diverse
grass communities on Mt. Kenya. This is consistent with multiple studies that have
documented higher biomass productivity and biodiversity, particularly in grasslands,
under conditions of elevated interglacial CO_2 and temperature compared to glacial
periods (55–58).

The total organic carbon (TOC) concentration, carbon-nitrogen (C/N) ratio and
terrestrial biomarker fluxes of Lake Rutundu sediments were low during the LGM before
rapidly increasing at *ca.* 16,000 cal. yr BP, indicating a reduction of terrestrial primary

productivity in the present-day heathland-moorland zone of Mt. Kenya during the LGM

(37, 42, 38). Thus, despite grasses becoming more abundant relative to other

herbaceous plants such as heather (Ericaceae), grass productivity and biomass

decreased relative to today, because Lake Rutundu was surrounded by the grass-

dominated but low-productivity ecosystem now occurring in the upper alpine zone (37,

38, 42). Finally, the positive relationship between temporal variation in entropy across

the PC1-PC2 subspace (Fig. 3A) and local biomass burning (as represented by micro-

and macrocharcoal influxes; Fig. 3D and SI Appendix, Fig. S5) suggests that climate-

driven changes in fire regime may have played an important role in shaping grassland

diversity on Mt. Kenya. However, considering that this relationship lacks statistical

significance across the other two subspaces of morphological variation, as well as the

complex interplay between temperature and moisture effects on fire dynamics in tropical

Africa (59), we refrain from deriving strong conclusions from this observation at this

time.

C_4 grasses have thinner pollen walls and simpler surface texture

The PC1 of CNN features representing pollen morphological variation among

modern grass species is significantly associated with whether those grasses use the C_3

or C_4 photosynthetic pathway, even after accounting for phylogenetic relatedness (Fig.

5D), indicating that this portion of CNN features captures real morphological differences

independent of shared evolutionary history. We cannot directly link PC1 to specific

morphological differences between C_3 and C_4 grass pollen, but there are statistically

significant differences in median pixel brightness and in the entropy of the distribution of

pixel values, which may indicate a thicker exine and more complex exine patterning in

C_3 versus C_4 species (Fig. 5A-C). Morphological differences between the pollen of C_3

and C_4 grasses may be linked to their distinct physiology. C_4 grasses invest more in

their root systems and tend to achieve faster growth by producing less dense leaves

(higher leaf area to mass ratio), enabling the production of more leaves at the same

carbon cost (60). This systematic difference in resource allocation may explain why

physiological differences are encoded in pollen morphology, independent of

phylogenetic relatedness. We hypothesize that the thinner exine wall and less complex

surface patterning of C_4 grass pollen reflect lower investment in pollen structure

resulting from the evolutionary trade-off between resource allocation to roots and

shoots.

Changing prevalence of C_3 and C_4 grasses in East African grasslands

During the late Miocene, fire activity was a key factor in the establishment and

expansion of C_4-dominated grasslands in Africa (11, 46, 47), and many contemporary

C_4 grasslands are fire-dependent (61, 63). Our results suggest a similar dynamic over

the last 25,000 years on Mt. Kenya. However, we did not find a significant correlation

between changes through time in the proportion of C_4 grasses (Fig. 3B) and

precipitation (Fig. 3C and SI Appendix, Fig. S5). Regional records, however, indicate

that biomass burning increased during wetter but more seasonal climates in eastern

equatorial Africa (~14,300–12,300 and 11,400–9,600 cal. yr BP), as shown by depleted

leaf-wax δD, depleted $δ^{13}C$, and elevated lake levels that mark wetter monsoon phases

(64). These wet-dry oscillations may have promoted greater vegetation growth followed

by burning, leading to greater C_4 abundance.

Full-glacial atmospheric CO_2 concentrations (~180–200 ppm) are limiting for both

C_3 and C_4 grasses, but C_4 species are generally better adapted to lower CO_2 and drier

conditions due to their ability to concentrate CO_2 around the carboxylating enzyme

Rubisco (10) and use water more efficiently (65). However, unlike diversity, the

reconstructed proportion of C_4 grasses correlated only weakly with CO_2 (SI Appendix,

Fig. S5). Likewise, temperature does not appear to have influenced the C_3:C_4 ratio

through time, with a near-zero correlation. Although C_4 grasses may have first evolved

in warm, tropical environments, they subsequently expanded and diversified into cooler

and even cold environments, and today occur across both warm and cold regions (66),

notably including cold-tolerant panicoid C_4 species in modern alpine grasslands (50,

65).

Implications for deep-time paleoenvironmental reconstruction

436   Our results demonstrate that deep learning methods applied to superresolution

microscopic images can be a powerful tool for revealing shifts in grass diversity and

community composition through time, by allowing us to quantify and exploit heretofore

hidden morphological variation in fossil grass pollen. By documenting the long-term

morphological variability contained in grass pollen assemblages, we can test

hypotheses about the evolution and physiological adaptations of grasses over

thousands to millions of years and provide a deeper understanding of how grasses

adapted to changing climates and environments. The first plants attributed to the

Poaceae appeared in the Early Cretaceous (113–101 Mya) (46–48), and by the middle

Eocene (~38 Mya; ref. 4), grasslands were present on every continent except

Antarctica. Our study focused on a fossil record of Late Quaternary grasslands to allow

validation of the method through direct comparison with other indicators of grass

diversity and C_3:C_4 ratio. Applying this same approach to fossil grass pollen

assemblages further back in the geologic record may at last open a true window on the

origin and evolution of this important global biome.

Methods

Pollen samples

454   Pollen samples of 60 extant grass species were isolated from herbarium material

from the Missouri Botanical Gardens (USA), University of Illinois (USA), and Harvard

(USA) or purchased from Sigma-Aldrich (St. Louis, MO, USA) (SI Appendix, Table S1).

These are predominately, but not exclusively, widely distributed species that occur in

East Africa, including the important African cereals sorghum (*Sorghum halepense*),

finger millet (*Eleusine coracana*), and maize (*Zea mays*) (SI Appendix, Table S2). Over

half of our species (56.67%) and over two-thirds of our genera (71.11%) are found on

Mt. Kenya. They represent 26.6% of Mt. Kenya's 124 grass species and 52.6% of its 52

genera (70) (SI Appendix, Table S2). Thirty-six species in our reference pollen dataset

are C_3 grasses and 24 are C_4 (SI Appendix, Table S1). All but three of the extant grass

species were represented by 19 or more pollen grains in our image dataset (SI

Appendix, Table S1).

Twenty-four samples of fossil pollen (SI Appendix, Table S4) were isolated from

a 7.55-m sediment sequence collected from Lake Rutundu in 1996 (37). Rutundu is a

40-ha oligotrophic lake, 3078 m a.s.l. on the northeast side of Mt. Kenya, just above the

current treeline (Fig. 2) (37). Recently deposited pollen was extracted from the surface

sediments of Lake Rutundu and five additional small lakes on Mt. Kenya, distributed

along an altitudinal transect between 1820 and 4585 m a.s.l. (Fig. 2 and SI Appendix,

Table S3), and used as an independent validation dataset (48). All pollen samples were

prepared following (71, 72) (details provided in the SI Appendix). Forty-nine to 50 fossil

grass grains were imaged from the six Mt. Kenya lake surface sediment samples (SI

Appendix, Table S3) and 48–51 fossil grass grains were imaged from each of the 24

Rutundu samples (SI Appendix, Table S4). The ages of the Rutundu samples are

derived from an age model based on AMS ^{14}C dating of bulk organic carbon (37, 38).

Paleoenvironmental data

Paleoenvironmental data for the subalpine zone of Mt. Kenya were taken from

published studies on the Rutundu sediment record. The temperature reconstruction is

based on abundance ratios of preserved glycerol dialkyl glycerol tetraethers (GDGTs)

(39) and the precipitation reconstruction is based on the hydrogen-isotope signature of

alkanes in fossil leaf waxes from terrestrial vegetation (40). For fire history, we used

data on microscopic and macroscopic charcoal influx to Lake Rutundu as a proxy for

past variation in local biomass burning and fire prevalence (38). We used the composite

record of atmospheric CO_2 concentration based on air-bubble measurements in several

Antarctic ice cores (41).

SR-SIM imaging

Pollen specimens were imaged using a Zeiss Elyra S1 superresolution structured

illumination (SR-SIM) microscope. SR-SIM is an optical microscopy method capable of

capturing morphological features <150 nm in length, below the diffraction limit of light

(44). We used a 63× Plan Apochromat (1.4 NA) oil objective and an excitation

wavelength of 561 nm. Image resolution was 0.0397 µm/pixel. Each axial plane of the

Z-stack was constructed with five grid rotations, spaced 0.18 µm apart, with 28–108

planes per grain. Images were cropped to the perimeter of each pollen grain.

Deep learning architectures and dataset setup

We developed two separate K-way classification CNNs based on the ResNeXt-

101 architecture (73): a holistic image CNN (H-CNN) trained on maximum intensity

projections (MIP) of entire image stacks, and a patch CNN (P-CNN) trained on square

crops of the MIP, each covering ~10% of the image (33) (Fig. 1A). Both CNNs were

trained using images of modern reference pollen (labeled data) and images of fossil

pollen (unlabeled data) in a semi-supervised learning (SSL) framework (74). SSL uses

the classification model to assign pseudo-labels to unlabeled images and continues to

train the model over both the pseudo-labeled and the labeled images. From the pseudo-

labeled images, only those that had pseudo-label confidence scores greater ≥95% were

retained for training. In our work, we divided the labeled reference data into training

(70%) and validation (30%) sets and evaluated model performance using 5-fold cross-

validation. The validation set was used for hyperparameter tuning and model selection

(SI Appendix, Fig. S1). SSL improved the model's ability to generalize across both

modern and fossil data. Training details are included in the SI Appendix.

The H-CNN generated a single K-dimensional logit vector for each pollen

specimen, while the P-CNN produced multiple logit vectors corresponding to multiple

input patches per specimen (Fig. 1A). The mean of these vectors by P-CNN was

calculated and normalized using softmax (33). For each specimen, we fused the outputs

of the two CNNs by multiplying the two K-dimensional classification probability vectors

and normalizing it to unit length (33).

Feature extraction

Our analysis relied on the features characterized by the two CNNs. We

concatenated the 2048-dimensional feature vectors derived from the penultimate (global

pooling) layer of the H-CNN and P-CNN to produce a 4096-dimensional feature vector

for each specimen (Fig. 1A). This combined feature vector represented a high-level

abstraction of the pollen image that served as input for our morphological analyses of

diversity and physiology. To visualize and compare morphological variation between the

modern grass pollen dataset and the 24 fossil pollen assemblages, we performed PCA

on the standardized feature matrix of all modern grains and then mapped fossil grains

onto this PCA space (Fig. 4).

Variation within species and genera

A common assumption in palynology is that variation in pollen from a single

individual captures the full range of variation in the morphology of an entire species. To

test whether this assumption holds in our dataset and if CNN features captured

individual or species-level variability, we conducted two analyses. First, we evaluated

whether species from the same genus are closer in feature space than they are to

species of different genera. We grouped all modern specimens by species and

averaged their CNN feature vectors to obtain species-level embeddings. We then

computed pairwise Euclidean distances among all species and used a one-sided

Wilcoxon rank-sum test to compare distances within versus between genera. This

analysis demonstrated whether there was taxonomic structure in our features. Second,

using species in our modern dataset for which pollen samples were available from

multiple individuals, as identified in the pollen metadata as originating from

MilliporeSigma (Burlington, MA, USA) or "other" sources, we assessed whether

variability in feature space differed between pollen samples derived from a single plant

and those derived from multiple individuals of the same species. For each species, we

measured the average feature variance across specimens, then compared the resulting

distributions between the two groups using a Wilcoxon rank-sum test. This analysis

demonstrated whether the variability in pollen from a single plant was representative of

the diversity within the broader species.

Ecological simulations and morphological measures of diversity

We first generated artificial communities to test our hypothesis that the morphological diversity within a grass pollen assemblage can serve as a proxy for taxonomic (species) diversity. To avoid reusing the same training data, we randomly withheld half of the modern dataset. Specifically, 30 of the 60 grass species were used to train a new CNN model, while the remaining 30 species (572 specimens) were reserved as an independent test set. This second model was trained solely for the purpose of the diversity-entropy analysis.

In our simulation, each artificial community contained a fixed number of specimens ($N = 50$), matching the counts available in fossil assemblages. For each replicate, we first drew species richness

$$S \sim \text{Unif}\,\{1, 30\}\,,$$

and then sampled S species without replacement from the 30-species test pool. Relative abundances were generated as

$$\tau \sim \text{LogUnif}\,(0.01, 20)\,, \quad \alpha = (\tau/S)\,1_S\,, \quad p \sim \text{Dirichlet}(\alpha)\,.$$

This parametrization allows assemblages to vary from highly uneven (small τ) to highly even (large τ).

Specimen counts were then sampled from a multinomial distribution,

$$X \sim \text{Multinomial}\,(50, p)\,.$$

If the sampled count for a species exceeded its available specimens, we capped it at the maximum available and redistributed the excess proportionally among the remaining selected species with remaining availability.

We quantified the diversity of each simulated community using Shannon's index from the empirical relative frequencies derived from the sampled counts:

$$H' = -\sum_{i=1}^{S} p_i \log_2(p_i)\,.$$

Where S is the total number of species and p_i is the relative frequency of species i.

578   Morphological disparity was quantified from the CNN feature vectors of each

579   assemblage. Feature matrices were standardized and reduced by PCA; each specimen

580   was represented by its PCA scores

581
$$Z_i = (Z_{i1}, Z_{i2}, Z_{i3}, \dots) \in R^d \,,$$

where d is the number of retained principal components (PCs). The set of all specimens

in a community is: $Z = \{Z_1, Z_2, \dots, Z_n\} \subset R^d$, where n is the number of specimens in an

assemblage.

We estimated the continuous analogue of Shannon entropy, the differential

entropy of the score distribution. For a density function $f(x)$, differential entropy is

defined as

588
$$H(f) = - \int_{R^d} f(x) \, \log_2 f(x) \, dx = - \mathbb{E}_{X \sim f}[\log_2 f(X)] \,.$$

589   This measures uncertainty associated with drawing a specimen at random from the

590   morphological space. Because f is unknown, we estimated it directly from the observed

591   specimens in each assemblage using Gaussian kernel density estimation (KDE) based

592   only on the observed PCA scores:

593
$$\widehat{f_h}(x) = \frac{1}{nh^d} \sum_{j=1}^{n} K\left(\frac{x - Z_j}{h}\right),$$

594
$$K(u) = (2\pi)^{-d/2} \, e^{-\frac{1}{2}\|u\|^2}, \quad u = \frac{x - Z_j}{h}, \quad h > 0 \,.$$

596   Using a leave-one-out (LOO) version to reduce self-influence:

597
$$\hat{f}_{-i,h}(Z_i) = \frac{1}{(n-1)h^d} (2\pi)^{-d/2} \sum_{j \neq i} e^{-\frac{\|Z_i - Z_j\|^2}{2h^2}} \,.$$

598   Since $H(f) = -\mathbb{E}[\log_2 f(X)]$, we estimate entropy by approximating the expectation with

599   the empirical mean:

$$\widehat{H} = -\frac{1}{n}\sum_{i=1}^{n} \log_2 \hat{f}_{-i,h}(Z_i).$$

Bandwidths were selected by maximizing the mean LOO log-likelihood of the pooled

PCA scores, separately for $d \in \{1, 2, 3\}$, evaluated over 500 logarithmically spaced

values between 0.01 and 10. The optimum for each d was denoted as h^*. For the

Rutundu fossil assemblages, entropy was computed only at h^*. For the simulated

assemblages and surface sediment samples, we evaluated sensitivity by scaling h^*

in $\log_2$ space:

607
$$\alpha \in 2^{\{-1, -\frac{1}{2}, 0, \frac{1}{2}\}} = \left\{\frac{1}{2}, \frac{1}{\sqrt{2}}, 1, \sqrt{2}\right\}, \quad h = \alpha h^*.$$

These four values bracket the optimal h^* and allow evaluation of how smoothing

influences diversity reconstruction.

Finally, morphological disparity from the simulated communities was compared

with their Shannon diversity values across 100 replicates using linear regression, with

Pearson's correlation coefficient (r) and its significance reported. For each panel, we

also display uncertainty as 95% confidence bands for the mean response and 95%

prediction bands for new observations (SI Appendix, Fig. S3).

Surface sample analysis

We applied the entropy measure described above to pollen isolated from the

surface sediments of six Mt. Kenya lakes, including Lake Rutundu. For each site,

Shannon entropy (bits) was calculated from PCA scores of the pollen assemblage and

compared with Poaceae species richness measured in vegetation surveys along the

same elevational transect (50). Analyses were performed on three PCA subspaces

(PC1, PC1-PC2, and PC1-PC2-PC3). To test the robustness of the entropy estimates,

we repeated the analysis across the same four bandwidth scaling factors defined

above. To visualize elevational diversity patterns, site-level entropy values were

connected using a piecewise cubic Hermite interpolating polynomial (PCHIP)

implemented in SciPy (v. 1.16.0) (63).

Fossil grass diversity estimation and paleoenvironmental correlates

We next applied the same entropy measure to 24 fossil pollen assemblages

isolated from the Lake Rutundu sediment core, spanning 25,000 years. For each

assemblage, we calculated Shannon entropy across the PC1, PC1-PC2, and PC1-PC2-

PC3 subspaces. These diversity estimates were then compared with five

paleoenvironmental variables: atmospheric CO_2 level, local reconstructed temperature

and precipitation, and micro- and macrocharcoal influx as proxy for fire prevalence.

Correlations were evaluated using Pearson's coefficient (r) and associated p-values (SI

Appendix, Fig. S5). All environmental proxy series were smoothed with a 4-point moving

average prior to visualization and correlation analysis to reduce high-frequency noise.

Grass pollen morphology and photosynthetic pathway

We averaged the CNN feature matrix by species and performed a standard PCA.

We tested whether the principal components discriminated between C_3 and C_4

photosynthetic pathways using Welch's t-tests (Fig. 5).

To determine whether the observed differences reflect genuine functional

differentiation or phylogenetic history, we tested whether the principal components

remained significantly discriminatory when accounting for evolutionary relationships. We

placed all 60 grass species within a recent time-calibrated Poaceae phylogeny of ~90%

of extant grass genera (69) by creating polytomies of multiple congeneric species using

the function bind.tip in the R package ape (76), with branch lengths of 0.001. This

conservative approach inflated phylogenetic signal (77), but allowed us to include all

species in the analysis. We tested for phylogenetic signal in PC scores using Pagel's λ

with the phylosig function in the R package phytools (78), then applied phylogenetic

generalized least squares (PGLS) regression using the phylolm function in the R

package phylolm (79) to determine whether C3–C4 differences persisted after

accounting for shared ancestry. We set all branch lengths to equal values to address

potential biases from artificial polytomies.

Estimates of past C$_4$ abundance and paleoenvironmental correlates

We performed a principal component analysis (PCA) on the CNN features of all individual pollen specimens from our modern Poaceae dataset (766 C$_3$ and 475 C$_4$ grains), retaining the principal components explaining up to 95% of the variance. The fossil pollen CNN feature matrix was standardized using the mean and standard deviation of the modern data and projected onto this PCA space. We then trained a gradient-boosted decision tree classifier (26, 27), implemented in the XGBoost Python library (v. 2.1.4) (80), and evaluated it with stratified five-fold cross-validation. The final trained model was applied to each fossil specimen from the 24 Rutundu samples to infer its photosynthetic pathway (C$_3$ or C$_4$) (Fig. 1C). For comparison, we also report in the same plot the δ^{13}C-based Bayesian mixture model estimates of C$_4$ abundances (38) and the fossil cuticle identifications of C$_4$ taxa (37), each shown as originally reported in their respective age models (Fig. 3B). For the latter, we traced the upper and lower bounding curves from their published figure and linearly interpolated each onto a dense evenly spaced age grid ($\leq$25 Ka). The midpoint (arithmetic mean) of the two bounds was taken as the central trajectory, with the half-difference forming an uncertainty envelope. We show shaded intervals representing 95% Wilson confidence intervals (81) for our estimates, 95% credible intervals for the Bayesian mixture model (38), and the range between "certain only" versus "certain + likely" cuticle identifications (37).

To evaluate temporal consistency between C$_4$ records, all series were smoothed using cubic smoothing splines (82) with varying smoothing factors (α = 0.01–0.4), interpolated onto a common 1000-point age grid spanning the 25 ka BP–present interval, and compared pairwise using Pearson's correlation coefficient (r) (SI Appendix, Fig. S6). We also assessed the correlation between our time series of estimated C$_4$ abundance and those of the five paleoenvironmental variables: atmospheric CO$_2$ concentration, local temperature, local precipitation, and micro- and macro-charcoal influx. We calculated Pearson's r and its corresponding p-value for each individual regression (SI Appendix, Fig. S5).

Quantifying morphological differences between C_3 and C_4 pollen

We used entropy to estimate the complexity of the grass pollen surface texture, with higher values indicating more random and complex textural patterning. We calculated Shannon entropy by segmenting the pollen grain image and quantifying the isolated grain's information content (complexity) based on the distribution of its pixel intensities (83). Because pollen exine is autofluorescent, pixel intensity allows us to infer variations in thickness of the pollen wall, with brighter images suggesting a thicker wall (44). We used direct exine thickness measurements (24) for 35 C_3 and 23 C_4 grass species to confirm this relationship. We calculated species-level medians of pixel intensity, morphological complexity (measured using Shannon entropy), and exine thickness across all specimens of each species and used Welch's t-test to assess the statistical significance of the observed differences between C_3 and C_4 species (Fig. 5).

Acknowledgments

Imaging costs and postdoctoral support for M.A.U. were provided by NSF-DBI – Advances in Bioinformatics (NSF-DBI-1262561) to S.W.P. S.K. was supported in part by the University of Macau (SRG2023-00044-FST), Science and Technology Development Fund of Macau SAR (0067/2024/ITP2), and the Institute of Collaborative Innovation. M.-E.A. was supported in part by the University of Illinois Tom L. Phillips Memorial Fund for Paleobotany. We thank Tim Gallaher for referring us to his recently published grass phylogeny.

Author Contributions

M.-E.A., S.K., and S.W.P. designed the research. M.-E.A. conducted the computational analyses and experiments. M.A.U. conducted the imaging, sample preparation, and lab analyses. F.A.S.-P. and D.V. provided samples and collected data. S.W.P. and S.K. supervised the research. M.-E.A., S.W.P., S.K., and M.A.U. wrote the manuscript, with feedback from all authors.

Data Availability

Superresolution structured illumination images used in this study were submitted to the Illinois Databank (DOI: 10.13012/B2IDB-1391373_V1) (84). The code base developed for this study is publicly available on GitHub (85) and can be accessed using the following link: https://github.com/paleopollen/Pollen_Diversity_Dynamics. In addition, several components of the analytical pipeline were adapted from code published in (86) available at: https://github.com/paleopollen/Novel_Pollen_Phylogenetic_Placement.

References

1. M. B. Bush, On the interpretation of fossil Poaceae pollen in the lowland humid neotropics. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 177, 5–17 (2002).

2. L. Mander, S. W. Punyasena, Grass pollen surface ornamentation: a review of morphotypes and taxonomic utility. *J. Micropalaeontology* 35, 121–124 (2016).

3. Grass Phylogeny Working Group III, *et al.*, Nuclear phylogenomics of grasses (Poaceae) supports current classification and reveals repeated reticulation. [Preprint] (2024). Available at: http://biorxiv.org/lookup/doi/10.1101/2024.05.28.596153 [Accessed 16 August 2024].

4. C. A. E. Strömberg, Evolution of Grasses and Grassland Ecosystems. *Annu. Rev. Earth Planet. Sci.* 39, 517–544 (2011).

5. J. S. Petermann, O. Y. Buzhdygan, Grassland biodiversity. *Curr. Biol.* 31, R1195–R1201 (2021).

6. G. J. Retallack, Cenozoic Expansion of Grasslands and Climatic Cooling. *J. Geol.* 109, 407–426 (2001).

7. S. E. Melzer, *et al.*, Fire and grazing impacts on silica production and storage in grass dominated ecosystems. *Biogeochemistry* 97, 263–278 (2010).

8. D. Tilman, J. A. Downing, Biodiversity and stability in grasslands. 367 (1994).

9. A. Lüscher, *et al.*, Using plant diversity to reduce vulnerability and increase drought resilience of permanent and sown productive grasslands. *Grass Forage Sci.* 77, 235–246 (2022).

10. R. F. Sage, The evolution of C_4 photosynthesis. *New Phytol.* 161, 341–370 (2004).

11. S. Hoetzel, L. Dupont, E. Schefuß, F. Rommerskirchen, G. Wefer, The role of fire in Miocene to Pliocene C_4 grassland and ecosystem evolution. *Nat. Geosci.* 6, 1027–1030 (2013).

12. C. A. E. Strömberg, R. E. Dunn, C. Crifò, E. B. Harris, "Phytoliths in Paleoecology: Analytical Considerations, Current Use, and Future Directions" in *Methods in Paleoecology: Reconstructing Cenozoic Terrestrial Environments and Ecological Communities*, D. A. Croft, D. F. Su, S. W. Simpson, Eds. (Springer International Publishing, 2018), pp. 235–287.

13. R. Sage, T. Sage, F. Kocacinar, Photorespiration and the Evolution of C_4 Photosynthesis. *Annu. Rev. Plant Biol.* 63, 19–47 (2012).

14. L. 't Mannetje, Climate change and grasslands through the ages: an overview.
Grass Forage Sci. 62, 113–117 (2007).

15. H. Behling, South and southeast Brazilian grasslands during Late Quaternary
times: a synthesis. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 177, 19–27 (2002).

16. L. Scott, Grassland development under glacial and interglacial conditions in
southern Africa: review of pollen, phytolith and isotope evidence. *Palaeogeogr.*
Palaeoclimatol. Palaeoecol. 177, 47–57 (2002).

17. M. J. Wooller, K. R. Beuning, Introduction to the reconstruction and modeling of
grass-dominated ecosystems. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 177, 1–3
(2002).

18. C. L. Yost, *et al.*, Phytoliths, pollen, and microcharcoal from the Baringo Basin,
Kenya reveal savanna dynamics during the Plio-Pleistocene transition.
Palaeogeogr. Palaeoclimatol. Palaeoecol. 570, 109779 (2021).

19. P. C. Twiss, "Predicted World Distribution of C_3 and C_4 Grass Phytoliths" in
Phytolith Systematics: Emerging Issues, G. Rapp, S. C. Mulholland, Eds. (Springer
US, 1992), pp. 113–128.

20. I. Holst, J. E. Moreno, D. R. Piperno, Identification of teosinte, maize, and
Tripsacum in Mesoamerica by using pollen, starch grains, and phytoliths. *Proc.*
Natl. Acad. Sci. 104, 17608–17613 (2007).

21. F. Katsi, *et al.*, FTIR spectra from grass pollen: A quest for species-level resolution
of Poaceae and Cerealia-type pollen grains. *Rev. Palaeobot. Palynol.* 321, 105039
(2024).

22. L. Mander, M. Li, W. Mio, C. C. Fowlkes, S. W. Punyasena, Classification of grass
pollen through the quantitative analysis of surface ornamentation and texture. *Proc.*
Biol. Sci. 280, 20131905 (2013).

23. L. Mander, *et al.*, Accuracy and consistency of grass pollen identification by human
analysts using electron micrographs of surface ornamentation. *Appl. Plant Sci.* 2,
1400031 (2014).

24. C. Wei, P. E. Jardine, W. D. Gosling, C. Hoorn, Is Poaceae pollen size a useful
proxy in palaeoecological studies? New insights from a Poaceae pollen
morphological study in the Amazon. *Rev. Palaeobot. Palynol.* 308, 104790 (2023).

25. C.-X. Wei, *et al.*, Grass pollen surface ornamentation is diverse across the
phylogeny: Evidence from northern South America and the global literature. *J.*
Syst. Evol. 62, 687–701 (2024).

26. C. Wei, *et al.*, A 23-million-year record of morphological evolution within
Neotropical grass pollen. *New Phytol.* 246, 365–376 (2025).

27. L. Schüler, H. Behling, Poaceae pollen grain size as a tool to distinguish past
grasslands in South America: a new methodological approach. *Veg. Hist.*
Archaeobotany 20, 83–96 (2011).

28. E. C. Stillman, J. R. Flenley, The needs and prospects for automation in
palynology. *Quat. Sci. Rev.* 15, 1–5 (1996).

29. K. Holt, G. Allen, R. Hodgson, S. Marsland, J. Flenley, Progress towards an
automated trainable pollen location and classifier system for use in the palynology
laboratory. *Rev. Palaeobot. Palynol.* 167, 175–183 (2011).

30. C. M. Barnes, *et al.*, Deductive automated pollen classification in environmental
samples via exploratory deep learning and imaging flow cytometry. *New Phytol.*
240, 1305–1326 (2023).

31. S. W. Punyasena, D. K. Tcheng, C. Wesseln, P. G. Mueller, Classifying black and
white spruce pollen using layered machine learning. *New Phytol.* 196, 937–944
(2012).

32. I. C. Romero, *et al.*, Improving the taxonomy of fossil pollen using convolutional
neural networks and superresolution microscopy. *Proc. Natl. Acad. Sci.* 117,
28496–28505 (2020).

33. M.-É. Adaimé, S. Kong, S. W. Punyasena, Deep learning approaches to the
phylogenetic placement of extinct pollen morphotypes. *PNAS Nexus* 3, pgad419
(2024).

34. M.-É. Adaimé, M. A. Urban, S. Kong, C. Jaramillo, S. W. Punyasena, Pollen
morphology, deep learning, phylogenetics, and the evolution of environmental
adaptations in Podocarpus. *New Phytol.* 247, 1460–1473 (2025).

35. J. H. Friedman, Greedy Function Approximation: A Gradient Boosting Machine.
Ann. Stat. 29, 1189–1232 (2001).

36. T. Chen, C. Guestrin, XGBoost: A Scalable Tree Boosting System in *Proceedings*
of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and
Data Mining, KDD '16., (Association for Computing Machinery, 2016), pp. 785–794.

37. K. J. Ficken, M. J. Wooller, D. L. Swain, F. A. Street-Perrott, G. Eglinton,
Reconstruction of a subalpine grass-dominated ecosystem, Lake Rutundu, Mount
Kenya: a novel multi-proxy approach. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*
177, 137–149 (2002).

38. M. A. Urban, D. M. Nelson, F. A. Street-Perrott, D. Verschuren, F. S. Hu, A late-
Quaternary perspective on atmospheric pCO_2, climate, and fire as drivers of C_4-
grass abundance. *Ecology* 96, 642–653 (2015).

39. S. E. Loomis, *et al.*, The tropical lapse rate steepened during the Last Glacial
Maximum. *Sci. Adv.* 3, e1600815 (2017).

40. S. Garelick, J. M. Russell, S. Dee, D. Verschuren, D. O. Olago, Atmospheric
controls on precipitation isotopes and hydroclimate in high-elevation regions in
Eastern Africa since the Last Glacial Maximum. *Earth Planet. Sci. Lett.* 567,
116984 (2021).

41. B. Bereiter, *et al.*, Revision of the EPICA Dome C CO_2 record from 800 to 600 kyr
before present. *Geophys. Res. Lett.* 42, 542–549 (2015).

42. M. Wooller, *et al.*, Late Quaternary vegetation changes around Lake Rutundu,
Mount Kenya, East Africa: Evidence from grass cuticles, pollen and stable carbon
isotopes. *J. Quat. Sci.* 18, 3–15 (2003).

43. F. A. Street-Perrott, *et al.*, Towards an understanding of late Quaternary variations
in the continental biogeochemical cycle of silicon: multi-isotope and sediment-flux
data for Lake Rutundu, Mt Kenya, East Africa, since 38 ka BP. *J. Quat. Sci.* 23,
375–387 (2008).

44. M. Sivaguru, *et al.*, Comparative performance of airyscan and structured
illumination superresolution microscopy in the study of the surface texture and 3D
shape of pollen. *Microsc. Res. Tech.* 81, 101–114 (2018).

45. Y. Lecun, L. Bottou, Y. Bengio, P. Haffner, Gradient-based learning applied to
document recognition. *Proc. IEEE* 86, 2278–2324 (1998).

46. C. E. Shannon, A mathematical theory of communication. *Bell Syst. Tech. J.* 27,
379–423 (1948).

47. C. E. Shannon, W. Weaver, *The Mathematical Theory of Communication*
(University of Illinois Press, 1998).

48. F. A. Street-Perrott, K. J. Ficken, Y. Huang, G. Eglinton, Late Quaternary changes
in carbon cycling on Mt. Kenya, East Africa: an overview of the $\delta^{13}C$ record in
lacustrine organic matter. *Quat. Sci. Rev.* 23, 861–879 (2004).

49. F. Pedregosa, *et al.*, Scikit-learn: Machine Learning in Python. *J. Mach. Learn.*
Res. 12, 2825–2830 (2011).

50. M. J. Wooller, D. L. Swain, F. A. Street-Perrott, S. Mathai, A. D. Q. Agnew, An
Altitudinal And Stable Carbon Isotope Survey of C_3 and C_4 Graminoids On Mount
Kenya. *J. East Afr. Nat. Hist.* 90, 69–85 (2001).

51. A. J. Baxter, *et al.*, Reversed Holocene temperature–moisture relationship in the
Horn of Africa. *Nature* 620, 336–343 (2023).

52. L. May, T. Lacourse, Morphological differentiation of *Alnus* (alder) pollen from
western North America. *Rev. Palaeobot. Palynol.* 180, 15–24 (2012).

53. I. Farrera, *et al.*, Tropical climates at the Last Glacial Maximum: a new synthesis of
terrestrial palaeoclimate data. I. Vegetation, lake-levels and geochemistry. *Clim.*
Dyn. 15, 823–856 (1999).

54. M. Kageyama, S. P. Harrison, A. Abe-Ouchi, The depression of tropical snowlines
at the last glacial maximum: What can we learn from climate model experiments?
Quat. Int. 138–139, 202–219 (2005).

55. R. F. Sage, Was low atmospheric CO_2 during the Pleistocene a limiting factor for
the origin of agriculture? *Glob. Change Biol.* 1, 93–106 (1995).

56. W. Jiang, Y. Cheng, X. Yang, S. Yang, Chinese Loess Plateau vegetation since the
Last Glacial Maximum and its implications for vegetation restoration. *J. Appl. Ecol.*
50, 440–448 (2013).

57. J. Cunniff, M. Charles, G. Jones, C. P. Osborne, Reduced plant water status under
sub-ambient pCO_2 limits plant productivity in the wild progenitors of C_3 and C_4
cereals. *Ann. Bot.* 118, 1163–1173 (2016).

58. T. He, B. B. Lamont, J. G. Pausas, Fire as a key driver of Earth's biodiversity. *Biol.*
Rev. 94, 1983–2010 (2019).

59. D. M. Nelson, D. Verschuren, M. A. Urban, F. S. Hu, Long-term variability and
rainfall control of savanna fire regimes in equatorial East Africa. *Glob. Change Biol.*
18, 3160–3170 (2012).

60. R. R. L. Atkinson, *et al.*, C_4 photosynthesis boosts growth by altering physiology,
allocation and size. *Nat. Plants* 2, 16038 (2016).

61. J. E. Keeley, P. W. Rundel, Fire and the Miocene expansion of C_4 grasslands.
Ecol. Lett. 8, 683–690 (2005).

62. C. P. Osborne, D. J. Beerling, Nature's green revolution: the remarkable
evolutionary rise of C_4 plants. *Philos. Trans. R. Soc. B Biol. Sci.* 361, 173–194
(2006).

63. J. G. Pausas, S. Paula, Grasses and fire: the importance of hiding buds. *New*
Phytol. 226, 957–959 (2020).

64. F. A. Street-Perrott, *et al.*, Late Quaternary changes in ecosystems and carbon
cycling on Mt. Kenya, East Africa: a landscape-ecological perspective based on
multi-proxy lake-sediment influxes. *Quat. Sci. Rev.* 26, 1838–1860 (2007).

65. S. P. Long, C_4 photosynthesis at low temperatures. *Plant Cell Environ.* 6, 345–363
(1983).

66. T. Watcharamongkol, P.-A. Christin, C. P. Osborne, C_4 photosynthesis evolved in
warm climates but promoted migration to cooler ones. *Ecol. Lett.* 21, 376–383
(2018).

67. E. A. Kellogg, "Fossil Record and Dates of Diversification" in *Flowering Plants.*
Monocots: Poaceae, E. A. Kellogg, Ed. (Springer International Publishing, 2015),
pp. 103–107.

68. Y. Wu, H.-L. You, X.-Q. Li, Dinosaur-associated Poaceae epidermis and phytoliths
from the Early Cretaceous of China. *Natl. Sci. Rev.* 5, 721–727 (2018).

69. T. J. Gallaher, *et al.*, Grasses through space and time: An overview of the
biogeographical and macroevolutionary history of Poaceae. *J. Syst. Evol.* 60, 522–
569 (2022).

70. Y.-D. Zhou, G. Mwachala, G.-W. Hu, Q.-F. Wang, Annotated checklist of the
vascular plants of Mount Kenya, East Africa. *Phytotaxa* 546, 1–108 (2022).

71. K. Fægri, J. Iversen, P. E. Kaland, K. Krzywinski, *Textbook of Pollen Analysis*, ed.
4th (John Wiley & Sons, 1989).

72. M. A. Urban, I. C. Romero, M. Sivaguru, S. W. Punyasena, Nested cell strainers:
An alternative method of preparing palynomorphs and charcoal. *Rev. Palaeobot.*
Palynol. 253, 101–109 (2018).

73. S. Xie, R. Girshick, P. Dollar, Z. Tu, K. He, Aggregated Residual Transformations
for Deep Neural Networks in *2017 IEEE Conference on Computer Vision and*
Pattern Recognition (CVPR), (IEEE, 2017), pp. 5987–5995.

74. K. Sohn, *et al.*, FixMatch: Simplifying Semi-Supervised Learning with Consistency
and Confidence in *Advances in Neural Information Processing Systems*, (Curran
Associates, Inc., 2020), pp. 596–608.

75. P. Virtanen, *et al.*, SciPy 1.0: fundamental algorithms for scientific computing in
Python. *Nat. Methods* 17, 261–272 (2020).

76. E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* 35, 526–528 (2019).

77. R. Molina-Venegas, M. Á. Rodríguez, Revisiting phylogenetic signal; strong or
negligible impacts of polytomies and branch length information? *BMC Evol. Biol.*
17, 53 (2017).

78. L. J. Revell, phytools 2.0: an updated R ecosystem for phylogenetic comparative
methods (and other things). *PeerJ* 12, e16505 (2024).

79. J. Lai, *et al.*, Evaluating the relative importance of phylogeny and predictors in
phylogenetic generalized linear models using the *phylolm.hp* R package. *Plant*
Divers. (2025). https://doi.org/10.1016/j.pld.2025.06.003.

80. T. Chen, C. Guestrin, XGBoost: A Scalable Tree Boosting System in *Proceedings*
of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and
Data Mining, (ACM, 2016), pp. 785–794.

81. E. B. Wilson, Probable Inference, the Law of Succession, and Statistical Inference.
J. Am. Stat. Assoc. 22, 209–212 (1927).

82. D. S. G. Pollock, "Smoothing with Cubic Splines" in *Handbook of Time Series*
Analysis, Signal Processing, and Dynamics, (Elsevier, 1999), pp. 293–322.

83. R. M. Haralick, K. Shanmugam, I. Dinstein, Textural Features for Image
Classification. *IEEE Trans. Syst. Man Cybern.* SMC-3, 610–621 (1973).

84. S. W. Punyasena, M. A. Urban, M.-É. Adaimé, I. C. Romero, SR-SIM
superresolution structured illumination microscope images of fossil and modern
pollen of Poaceae. University of Illinois Urbana-Champaign.
https://doi.org/10.13012/B2IDB-1391373_V1.

85. M.-É. Adaimé, S. Kong, S. W. Punyasena, Computational Tools for Quantifying
Poaceae Pollen Diversity and Photosynthetic Pathway Composition (C_3/C_4) from
Superresolution Images. (2024). https://doi.org/10.5281/zenodo.13756262.
Deposited 13 September 2024.

86. M.-É. Adaimé, S. Kong, S. W. Punyasena, A Computational Framework Integrating
Molecular Phylogenetic Data and CNN-Derived Morphology for the Evolutionary
Placement of Unknown Pollen Types. (2023).
https://doi.org/10.5281/zenodo.8128443. Deposited 9 July 2023.

Figures

Fig. 1. Flowchart illustrating the pipeline of computational analyses performed in this

study. (A) Two superresolution microscopic representations of a grass pollen grain (the

maximum intensity projection of the image stack and overlapping crops of the maximum

intensity projection) are passed to convolutional neural networks (CNNs) trained to

recognize the holistic morphology (H-CNN) and surface texture of patches (P-CNN) of

grass pollen. The morphological features extracted by both CNNs are concatenated to

form a comprehensive feature vector representing each specimen. These vectors are

then stacked across all N grass pollen grains present in an assemblage, forming a

feature matrix of all the concatenated feature vectors. (B) In our assessment of grass

diversity, the PCA scores of the feature matrix are used to estimate probability densities

for pollen assemblages using kernel density estimation. Shannon entropy is calculated

over these estimated densities to provide a quantitative measure of diversity. (C) In our

assessment of $C_3:C_4$ ratios, we apply PCA to the feature matrix and pass the most

informative principal components to a gradient boosted decision tree (XGBoost) trained

to discriminate between C_3 and C_4 grasses.

Fig. 2. Setting of the 25,000-year grass-pollen record used for method validation. (A) Distribution of major vegetation biomes across the African continent, with the location of Mt. Kenya in eastern equatorial Africa highlighted by a red square. (B) Elevation gradient in the Mt. Kenya region with the locations of our six sources of fossil grass pollen: the 25,000-year sediment record from Lake Rutundu, and the five additional lakes (Lake Nkunga, Rumuiku Swamp, Lake Ellis, Small Hall Tarn, and Simba Tarn) located between 1820 m and 4585 m elevation where we extracted subfossil pollen from recently deposited sediment.

Fig. 3. Changes in grass diversity and proportion of C$_4$ grasses in subalpine grasslands

of Mt. Kenya over the past 25,000 years in relation to paleoenvironmental history.

Vertical gray shading delineates major late-glacial climatic intervals: the Last Glacial

Maximum (LGM; 23,000–19,000 cal. yr BP), Heinrich Stadial 1 (H1; 18,000–16,000 cal.

yr BP), and the Younger Dryas (YD; 13,000–11,500 cal. yr BP). Paleoenvironmental

proxies (C, D) were interpolated using moving averages with a window size of 4. (A)

Grass species richness as inferred from the morphological diversity of grass pollen, as

estimated by differential entropy of Gaussian kernel density estimates (KDEs) on PCA

scores (PC1, PC1-PC2, and PC1-PC2-PC3). Bandwidth selection, curve smoothing,

and evaluation are detailed in Methods. Results are normalized for visualization. (B)

Proportion of C_4 grasses of the grassland surrounding Lake Rutundu as estimated by

this study (dashed green line), the carbon-isotopic signature ($\delta^{13}C$) of individual grass

pollen grains (solid blue line; ref. 38), and fossil grass cuticle (solid purple line; ref. 37).

Uncertainty is shown as shaded bands: 95% Wilson confidence intervals for this study,

95% credible intervals from the Bayesian mixture model in (38), and bounds from

"certain only" versus "certain + likely" C_4 cuticle identifications in (37). (C) Antarctic ice-

core record of atmospheric CO_2 concentration (41) and reconstructions of local mean

annual temperature (MAT) and precipitation derived from, respectively, bacterial

membrane lipids and the hydrogen-isotopic signature (δD_{precip}) of plant leaf-wax alkanes

extracted from Lake Rutundu sediments (28, 29). More depleted δD_{precip} values indicate

wetter conditions. (D) History of fire prevalence on Mt. Kenya reconstructed from

estimates of biomass burning based on sedimentary fluxes of micro- and macrocharcoal

(38).

Fig. 4. (A) PC1-PC2 biplots of the morphological variation among 1,241 grass pollen grains in our 60-species modern dataset and the projected placement of 1,204 fossil grains from all 24 analyzed time windows in the Lake Rutundu record. There is partial separation between the pollen of modern C$_3$ (blue) and C$_4$ (yellow) grass species, and

the morphological variation among the fossil Poaceae pollen falls largely within a subset

of the modern range of morphospace. PC1 and PC2 together account for 17.05% of the

variance. (B) PC1-PC2 biplots of the pollen morphological variation in fossil

assemblages from each analyzed time window, with shaded convex hulls delineating

the morphological range present in each assemblage. The colors for each time window

match the boxplots below. (C) Boxplots of the median values and ranges of PC1 and

PC2 scores of the 24 fossil assemblages, highlighting the distinctively low PC1 scores

of the four assemblages dated to between 21,160 and 18,435 BP.

Fig. 5. Measurements by this study and previous morphometric analyses showing that C_3 grass pollen has a thicker exine wall and more complex surface patterning than C_4 grass pollen. Boxplots showing the difference in (A) pixel intensity (brightness) and (B) morphological complexity (measured using Shannon entropy) between pollen from 36 C_3 and 24 C_4 grass species in our modern reference dataset, and (C) in exine thickness (μm) of pollen from 35 C_3 and 23 C_4 grass species determined by (24). (D) PC1-PC2 biplot of mean CNN-derived pollen morphological features across 60 modern Poaceae species illustrating the morphological differentiation between C_3 and C_4 grasses. Each point represents a mean species-level feature vector.