Comparative pan-genomics reveals divergent adaptations in
clinically-relevant members of the *Fusarium solani* species
complex

Phillip J.T. Brassington[1,2], Maria Laura Fabre[1#], Anna Zimmermann[2#], Marion Perrier[1,3], Ronny
Martin[2], Oliver Kurzai[2,4], Amelia E. Barber[1,3]

[1] Junior Research Group Fungal Informatics. Institute of Microbiology. Friedrich Schiller
University. Jena, Germany.
[2] Institute of Hygiene and Microbiology. University of Würzburg. Würzburg, Germany.
[3] Cluster of Excellence Balance of the Microverse. Friedrich Schiller University. Jena, Germany.
[4] National Reference Center for Invasive Fungal Infections (NRZMyk). Leibniz Institute for
Natural Product Research and Infection Biology-Hans Knöll Institute. Jena, Germany.

[#] Maria Laura Fabre and Anna Zimmermann contributed equally. Author order was determined
alphabetically.

Address correspondence to:
Amelia E. Barber
amelia.barber@uni-jena.de

Abstract

The *Fusarium solani* species complex are a group of dual-kingdom fungal pathogens capable of causing devastating disease on a wide range of host plants and life-threatening infections in humans that are difficult to treat. In this study, we generate highly contiguous genomes for three clinical isolates of *Fusarium keratoplasticum* and three clinical isolates of *Fusarium petroliphilum* and compare them with other genomes of the FSSC from plant and animal sources. We find that human pathogenicity is polyphyletic within the FSSC, including in *F. keratoplasticum*. Pan-genome analysis revealed a high degree of gene presence-absence in the complex, with only 41% of genes (11,079/27,068) found in all samples and the presence of accessory chromosomes encoding isolate- and species-specific genes. We also defined conserved long non-coding RNAs (lncRNAs) between *F. keratoplasticum* and *F. petroliphilum,* revealing that they show a similar low degree of presence-absence variation. Secondary metabolite analysis revealed a conserved core set of biosynthetic gene clusters across the FSSC, as well as a unique cluster potentially linked to keratitis. Transcriptomic analysis under stress conditions showed minimal differential gene expression, indicating that both *F. keratoplasticum* and *F. petroliphilum* are well adapted to human infection-relevant conditions. This study provides valuable insights into the evolutionary dynamics, genomic architecture, and potential pathogenicity mechanisms of the FSSC, with implications for understanding multi-kingdom virulence, of increasing relevance as climate change potentially increases the number of fungal species that can grow at human temperatures.

Introduction

The genus *Fusarium* encompasses a highly diverse range of filamentous fungal species, including environmental saprobes, plant pathogens, and organisms capable of causing disease in animals and humans. Well established as plant pathogens, *Fusarium* spp. are causal agents for devastating worldwide food supply disruptions [1–3], causing significant losses in global yields of wheat, maize, rice, potato, and soybean [4], with outbreaks of *Fusarium* head blight causing over 50% wheat losses in China, Europe, and the Americas [5]. Yet beyond this, *Fusarium* spp are largely unique among fungi in that they are multi-kingdom pathogens capable of also infecting higher eukaryotes. In this capacity, *Fusarium* spp. are playing a role in the decline of threatened animal species [6], and are the causative agents with rising numbers of highly lethal human fusariosis infections [7]. In 2022, the World Health Organization identified the genus as one of the highest priority human fungal pathogens, and one with the most urgent need for additional study based on current knowledge gaps [8]. These include lack of diagnostic options, limited

epidemiologic data, and the intrinsic antifungal resistance, which cumulatively contribute to a high
mortality rate of 43-67% [8].

Previous studies have highlighted the exceptional range of genomic diversity and plasticity among
Fusarium spp. Even within a single species, the genome structure is highly flexible, and strains
often possess accessory chromosomes that are variably conserved and, importantly, confer the
ability to infect specific plant species [9]. These accessory chromosomes can even be horizontally
transferred, resulting in a corresponding transfer of host plant specificity [10–12]. Despite the large
number of species in the genus, the majority (reportedly up to 80-90%) of human infections are
due to only two species groups: the *Fusarium oxysporum* species complex and the *Fusarium*
solani species complex [13–17].

The *Fusarium solani* species complex (FSSC) comprises globally ubiquitous problematic
agricultural pathogens and is the most frequent cause of human fusariosis [1,3,13,17,18]. Among
the human pathologies it causes, it is particularly problematic in causing corneal infection and
disseminated disease. More recently, it has been linked to two outbreaks of meningitis in
immunocompetent patients in Mexico due to contaminated anesthetic solutions [19,20]. In the
plant kingdom, the FSSC exhibits a broad host range and is capable of causing root rot in multiple
agriculturally important crops, including peas, cucurbits, and sweet potato [21–24]. Even within
individual species of the complex, there is a broad range of habitats and hosts. For example,
F. keratoplasticum has been found in diverse habitats ranging from plumbing systems, strawberry
roots, Chinese softshell turtle eggs, to human nails, eyes, and endocardium [6,25–29]. This is
also likely a non-exhaustive list, as many studies classify isolates as *forma specialis* (*f. sp*). based
on the host plant, rather than identifying the exact FSSC species member. The diversity of habitats
and pathologies caused by the FSSC in both plants and animals is accompanied by a
correspondingly high level of genomic diversity, including the presence of core and accessory
chromosomes [6]. However, the genomes of FSSC members remain understudied compared to
other *Fusarium* groups like the *Fusarium oxysporum* species complex. Moreover, it is unclear
what, if any, genomic features are specific to particular species or hosts (plant or animal).

In this study, we characterized the genomic diversity of six FSSC member species focusing on
F. keratoplasticum and *F. petroliphilum*, two of the most prominent FSSC species in human
keratitis [28,30], with the latter being particularly understudied and lacking even a reference
genome. To this end, we generated near-chromosomal genomes of three novel clinical isolates

of *F. keratoplasticum*, and provided the first three genomes for *F. petroliphilum*. We also predict

long non-coding RNAs for these species, the first to investigate this class of regulatory genes in

the FSSC. Finally, we integrate these genomes with six previously sequenced, nearly gapless,

genomes from plant and animal isolates of the FSSC [6] to study the genomic diversity of the

complex.

Results

Genome sequencing of novel clinical *Fusarium solani* species complex isolates and

description of the study dataset

In this study we used PacBio long read sequencing combined with Illumina short read sequencing

to assemble the genomes of six novel FSSC isolates that were collected from keratitis cases by

the German National Reference Center for Invasive Fungal Infections (n=3 *F. keratoplasticum*

and n=3 *F. petroliphilum*). To facilitate more in-depth comparisons within the species complex,

and to include isolates of environmental and animal origin, six additional FSSC isolates with

similarly contiguous genomes were incorporated into the study dataset [6]. The published

samples included isolates of *F. falciforme*, *F. keratoplasticum, F. solani* FSSC12, *F. vanettennii*,

and originated from plant (pea plant and orchid flower) and marine animal (including hawksbill

sea turtle, roughtail stingray, and harbor seal) sources (Table 1). The final assemblies of the novel

and published assemblies were highly contiguous, with an average of 58 scaffolds (range 14 to

143) and an average N50 of 3.8 Mb ± 0.4 (Table 1). The FSSC genomes analyzed in this study

ranged in size from 48.0 Mb for one of the newly sequenced *F. keratoplasticum* isolates to 72.9

Mb for *F. vanettenii* Fs6 and encoded between 14,050 and 16,755 protein coding genes.

Parallelling the range of genome sizes, a similar pattern was observed for the fraction of the

genome consisting of repetitive regions, including transposable elements, which varied from 4.5%

for *F. keratoplasticum* NRZ-2015-035 to 14.2% for *F. vanettennii* Fs6. Using BUSCO as a

measure of genome completeness, we found that all genomes were highly complete, recovering

on average >98.5% of the expected single-copy orthologs. Despite originating from different

studies, there were no significant differences in key genome quality metrics such as N50, the

number of protein coding genes, or BUSCO completeness between the isolates sequenced in

this study and the public assemblies (p>0.05 by Mann-Whitney U test).

Human and plant association is polyphyletic in the FSSC.

To determine the phylogenetic relationship of the newly sequenced clinical isolates within the
genus *Fusarium* and the FSSC complex, we constructed a whole genome phylogeny from amino
acid sequence of conserved single copy orthologs. As expected, the new isolates of
F. keratoplasticum and *F. petroliphilum* were located within clade 3 of the FSSC in the phylogeny
(Figure 1). The *F. petroliphilum* isolates formed a monophyletic group, with *Fusarium* FSSC12
being the closest relative in the dataset. The *F. keratoplasticum* isolates belonged to a separate
monophyletic group containing *F. solani sensu stricto*, *F. falciforme*, and the two other
F. keratoplasticum isolates in the dataset. Interestingly, the clinical *F. keratoplasticum* isolates
from Germany did not segregate from the marine host-associated isolates from east Asia,
indicating a lack of geographic population structure or host specificity in the species. This overall
pattern was observed in the entire complex, with virulence in both plant and human hosts being
polyphyletic in the FSSC.
131
41% of genes are common to all FSSC isolates, while pan-gene conservation is higher in
individual species.
We next sought to quantify the genomic diversity of the FSSC by characterizing its pan-genome
and delineating the genes common to all genomes analyzed (core genome) and those which
show presence-absence variation within the species complex (accessory genome). Of the 27,068
orthologous genes identified across the 12 isolates and 6 species, 41% (n=11,079) were core, or
present in all isolates (Figure 2a & b). The FSSC pan-genome exhibited a long tail distribution,
where 6% of the pan-genome (n=1,666) was present in all but one sample and 7% (n=1,946) of
the accessory genome was isolate-specific (7%; n=1,946) (Figure 2b). Despite the low fraction of
core genes, Heaps' law indicates that the FSSC pan-genome is closed (alpha = 1.57, where alpha
values > 1 indicate a closed pan-genome) and the number of pan-genes beginning to plateau
after six genomes (Figure 2c). Protein sequences of core genes were significantly longer than
those of accessory genes, averaging 505 amino acids compared to 377 for the latter (Wilcoxon p
< 0.0001). Within the scope of single species, the fraction of core genes was notably higher. For
F. keratoplasticum, 75% of pan-genes were core, or in all 5 isolates (13,275/17,617), while 88%
of pan-genes were core in *F. petroliphilum* (14,067/15,923 pan-genes; n=3 isolates) (Figure 2d &
e).
149
Diverging patterns in sequence conservation and repetitive elements between core and
accessory chromosomes

In addition to the high degree of gene presence-absence variation, some of the accessory
genome in *Fusarium* species is known to be encoded on discrete accessory chromosomes. These
accessory chromosomes are involved in plant virulence and the ability to infect specific host
plants, but their role in human infection is unclear. Intriguingly, work in the *F. oxysporum* species
complex hints that clinical isolates of this species complex may carry accessory chromosomes
that are distinct from plant-pathogenic isolates [10]. To investigate this in the FSSC, we analyzed
our study dataset for the conservation of both core and accessory contigs among the isolates
from plant and animal origins. We identified 12 core chromosomes, which were present across
the isolates in the dataset with a high degree of nucleotide identity and synteny (Figure 3).
However, the relative proportion of the core chromosomes compared to the total genome varied
considerably within the species complex. Core chromosomes, encompassing a total length of
46.8 Mb in *F. keratoplasticum* Fu6, made up 95% of its total genome length. In comparison, core
chromosomes comprised only 73% of the total genome length of *F. vanettenii* Fs6, which
contained 15 additional large scaffolds or contigs with low identity to any other sequences in the
dataset (Supplementary Figure 1). We also observed that all samples in the clade containing
F. petroliphilum and *F. sp. FSSC12* possessed a substantial chromosomal translocation between
the two largest core chromosomes (Supplementary Figure 3).

In contrast to the high synteny and sequence conservation among the core chromosomes, the
accessory chromosomes were largely species- or isolate-specific or had much lower levels of
sequence identity for the regions that were present (Figure 4). We detected between three and
eight accessory chromosomes or contigs >150kb in the *F. keratoplasticum* isolates, and between
five and nine in the three *F. petroliphilum* isolates. Many accessory chromosomes were
conserved in all isolates of the same species, but not across species. For example, two of the
three accessory chromosomes present in *F. keratoplasticum* Fu6 had high identity with the other
F. keratoplasticum isolates but low synteny with other FSSC isolates (Figure 4). In other cases,
accessory chromosomes were isolate-specific, such as chromosome 13 of *F. keratoplasticum*
LHS11, which displayed <52.3% identity with any other contig in the dataset. However, we did
not identify any accessory chromosomes or contigs in *F. keratoplasticum* that were specific to the
clinical strains. We also observed cases where the distribution of accessory chromosomes or
contigs did not mirror the core genome-generated phylogeny. For example, four
F. keratoplasticum isolates (Fu6, LHS11, 2015-087, and 2016-116) possessed one to three
scaffolds whose nucleotide identity was higher to accessory genomic regions of *F. petroliphilum*
than to the other strains of its own species (Figure 4a). Conversely, F. *petroliphilum* strains 2014-

013 and 2015-035 possessed accessory contigs with higher synteny and nucleotide identity to
F. keratoplasticum than to their own species (Figure 4b).

The accessory chromosomes of all six FSSC species possessed a significantly higher fraction of
repetitive elements, including transposons, compared to the core chromosomes (p <0.05 for all
species by Mann-Whitney U test) (Supplementary Figure 2). For example, the median repeat
content of the 12 core chromosomes among the five *F. keratoplasticum* isolates was 6.7%,
compared to 24.5% for the accessory chromosomes and contigs. The synteny analysis also
revealed that the short regions of lower synteny in the core chromosomes were closely associated
with the presence of repetitive elements. Altogether, we defined 12 core chromosomes with a
high degree of sequence conservation across the FSSC and an assortment of species- and
isolate- specific accessory chromosomes that also had a higher content of repetitive elements.

57% of intergenic lncRNAs in *F. keratoplasticum* are conserved in *F. petroliphilum*
In addition to variation in protein-coding genes, eukaryotic genomes encode myriad long non-
coding transcripts (lncRNAs) that are often involved in gene regulation and evolve faster than
coding genes [31]. Previous work on lncRNAs in *Fusarium graminearum* have highlighted their
roles in regulating sexual reproduction [32] and carotenoid biosynthesis [33], however the lncRNA
landscape of the FSSC has not been explored. To this end, we sequenced the transcriptomes of
F. keratoplasticum NRZ-2014-069 and *F. petroliphilum* NRZ-2014-079 conidia and of mycelia
grown under several environmental conditions to capture the breadth of lncRNAs expressed by
these two species. Using this data, we identified 2,632 and 2,222 putative lncRNAs for the two
species, respectively. Both species shared a comparable ratio of intergenic to antisense lncRNAs,
with the majority of lncRNAs being intergenic and only 16% being antisense in both species
(Supplementary Figure 4). Intergenic lncRNAs were significantly shorter than antisense lncRNAs
and protein-coding genes, with a median length of 795 bp for *F. keratoplasticum* compared to
1,231 bp and 1,317 bp for antisense and for protein coding genes, respectively (p < 0.001 for all
comparisons by Bonferroni-corrected Mann-Whitney U test). Similarly, intergenic lncRNAs in
F. petroliphilum had a median length of 831 bp compared to 1,352 bp for antisense lncRNAs and
1,345 bp for protein coding genes (Supplementary Figure 4).

To identify conserved intergenic lncRNAs between the two species, we used a combination of
reciprocal BLASTn (local alignment), MUMmer (global alignment), and synteny-based
approaches. BLASTn identified 395 conserved lncRNA pairs, representing 18% of the intergenic

lncRNAs in *F. keratoplasticum*. MUMmer alignment identified 385 pairs, with a 95% concurrence

with BLASTn. As lncRNAs are known to have lower levels of sequence conservation compared

to protein-coding genes, we also employed a synteny-based approach as in [34]. Conserved

lncRNAs were identified between the two species if they were flanked by at least one up- and

downstream orthologous genes. This approach identified 1,647 syntenic genomic blocks,

containing lncRNAs (Supplementary Figure 5, Supplementary File 3). Taking the union of the

three approaches revealed that 57% of the intergenic lncRNAs in *F. keratoplasticum* (1264/2224)

were conserved in *F. petroliphilum*, similar to the conservation rate among protein-coding genes

in the two species. Notably, all conserved lncRNAs identified through synteny analysis were

located on core chromosomes. However, we detected 51 species-specific lncRNAs in

F. keratoplasticum and 104 in *F. petroliphilum* that were located on accessory chromosomes.

Altogether, we defined the lncRNAome of two FSSC species and found that the degree of their

conservation was similar to that of protein coding genes.

Fusarium keratoplasticum genomes are enriched in branched-chain amino acid and

metallopeptidases, while all FSSC genomes show features of plant pathogenicity

Given the degree of variation in the pan-genome of the FSSC and the accessory chromosomes

within and between species, we next performed gene set enrichment analysis to identify Gene

Ontology (GO) categories that were over- or underrepresented in species of the FSSC, with a

focus on *F. keratoplasticum* and *F. petroliphilum*. The genomes of *F. keratoplasticum* had an

expansion of genes involved in branched-chain amino acid GO terms relative to other FSSC

genomes, a pathway which is not-conserved in metazoans, but essential to growth and virulence

in several human pathogenic fungi [35,36]. We also observed an enrichment in genes annotated

in protein ubiquitination and metallopeptidase activity in this species, which serve roles in fungal

growth and pathogenicity [37], including tissue invasion [38] and hyphae formation [39]. The

genomes o*f F. petroliphilum* had higher numbers of genes with endonuclease activity and

glutathione metabolic processes relative to the other FSSC species. The former plays a strong

oxidoprotectant role in fungi [40], and the latter of which is involved in defense against reactive

oxygen species and the repair of damaged DNA [41].

Underscoring their evolutionary trajectory as plant pathogens, all FSSC genomes contained an

abundance of secreted effectors and genes with GO annotations related to plant virulence. These

included similar levels of genes with GO terms related to chitosanase activity, pectinesterase

activity, and cellulase activity, all of which are involved in the degradation of plant cell walls

[42,43]. All genomes also had between 477-546 secreted effectors. The fact that these genomic
features were present at similar levels in plant isolates and human and marine animal isolates
suggests that the latter likely retain some degree of plant pathogenicity or, alternatively, that these
factors are also important during interaction with higher eukaryotes. Taken together, the gene
functional analysis revealed that all FSSC genomes share features associated with plant
virulence, while *F. keratoplasticum* and *F. petroliphilum* possessed differential expansion of
genes involved in branched chain amino acid metabolism, oxidative stress protection, and DNA
repair.

Comparison of secondary metabolite biosynthetic gene clusters reveals a core set of
secondary metabolites in the FSSC
Fungi produce an extraordinary array of secondary metabolites, which play key roles in their
interactions with other environmental organisms. They can be used for microbial communication,
nutrient scavenging, and even as antimicrobial weapons. These molecules are typically encoded
in biosynthetic gene clusters (BGCs), or organized units of multiple genes all involved in
metabolite production [44]. *Fusarium* spp, including FSSC, produce a diverse range of
compounds, with 162 novel *Fusarium* metabolites described between the years of 2015-2020
alone [45]. However, the range of BGC produced by FSSC isolates from human and animal origin
has not been studied. Across the 12 genomes, we predicted 42 BGCs and each isolate encoded
a median of 33 BGCs (range 32-37). Of the BGCs identified, 55% (23/42) were present across all
study strains (Figure 6). Among the BGCs present in all genomes that matched a known product
was ochratoxin A, an agricultural contaminant. Among the other core BGCs with known products
were squalestatin S1, fusarubin, metachelin C, tryptoquialanine, and lucilactaene, many of which
have antimicrobial activity [46–50]. The BGC encoding the antimicrobial and immunosuppressant
cyclosporin C was produced by the phylogenetic subclade encompassing *F. keratoplasticum,*
F. falciforme, and *F. sp. Ph1*, but absent in the other FSSC clades examined.

The majority of the detected BGCs (24/42) encoded unknown products. One such example was
BGC19 (Supplementary Figure 6), a secondary metabolite cluster of the isocyanide class, which
are known for their ability to chelate transition metals. This BGC was found in five of the six
keratitis isolates (n=2 *F. petroliphilum* and n=3 *F. keratoplasticum*), but none of the other study
isolates, including the non-keratitis *F. keratoplasticum* isolates, hinting that it might play a role in
virulence in the eye. Also of interest, *F. sp. Ph1* was predicted to produce gibberellin, a BGC
typically restricted to the *Fusarium fujikuroi* species complex. Gibberellin is a plant growth

hormone and its production by *Fusarium fujikuroi* is considered a major factor in the development
of symptoms in rice bakanae disease, which causes up to 50% crop loss [51,52].

Minimal transcriptional changes in *F. keratoplasticum* and *F. petroliphilum* at ocular
infection temperatures

Given the genomic differences observed between *F. keratoplasticum* and *F. petroliphilum* in this
study, we wanted to compare the transcriptomes of these two species under several stressors.
To this end, we performed RNAseq of mycelia from *F. keratoplasticum* NRZ-2014-069 and
F. petroliphilum NRZ-2014-079 grown in complete media at 25ºC, under carbon starvation at this
temperature, and at elevated temperature mimicking the surface temperature of the eye (34ºC).
Additionally, the transcriptome of conidia from each isolate was also assayed. Growth stage
(conidia vs. mycelia) had the largest impact on gene expression with 1,185 and 2,017 differentially
expressed genes (DEG) in *F. keratoplasticum* and *F. petroliphilum* respectively (Supplementary
File 4). Gene set enrichment analysis of GO terms revealed that conidia from *F. keratoplasticum*
had an upregulation of DNA-binding proteins and transporters compared to mycelia (Figure 7,
Supplementary File 5), while *F. petroliphilum* showed an upregulation in genes cell wall
organization and structure.

Relative to complete media, mycelia grown under carbon starvation resulted in 357 DEGs in
F. keratoplasticum and 850 DEGs in *F. petroliphilum* (Supplementary File 4). Both species
showed an upregulation in nucleotide binding and ATP binding. However, only F. petroliphilum
showed an upregulation in genes with metal ion binding activity (Figure 7, Supplementary File 5).
Unexpectedly, elevated temperature (34°C) resulted in very modest change to the transcriptome,
with 95 genes differentially expressed in *F. petroliphilum* and zero in *F. keratoplasticum*
(Supplementary File 4). This suggests that both species are more than capable of growing at
temperatures experienced during mammalian association or infection

Consistent with earlier studies [32], both intergenic and antisense lncRNAs identified in this study
had significantly lower expression relative to protein-coding genes (Mann-Whitney U test with
Bonferroni correction, $p < 0.0001$ for both lncRNA classes in *F. keratoplasticum* and
F. petroliphilum (Supplementary Figure 4b). However, we did identify 169 lncRNAs in
F. keratoplasticum and 190 in *F. petroliphilum* that were uniquely differentially expressed in
conidia relative to mycelia, suggesting they may play a role in asexual development
(Supplementary File 4). Overall, the transcriptional changes of *F. keratoplasticum* and

F. petroliphilum during carbon starvation and elevated temperature were minimal, indicating that

neither infection-relevant condition was a significant stressor to either species.

325   Discussion

In this study we generate the highly contiguous genomes for six isolates of the FSSC from clinical

origin, including the first three genomes available for *F. petroliphilum*. We show that German

keratitis isolates of *F. keratoplasticum* are not phylogenetically distinct from Asian isolates from

marine animal origin. We also reveal a low level of core gene conservation among isolates of the

FSSC, including pronounced variation in lncRNAs, secondary metabolites, and the presence of

accessory chromosomes with isolate- and species- specific genes and a higher repeat content.

Finally, transcriptional profiling of *F. keratoplasticum* and *F. petroliphilum* at temperatures

mimicking the human eye revealed minimal changes, suggesting that neither species experiences

infection temperatures as a significant stress.

In line with previous work, our study highlights the considerable genomic flexibility of *Fusarium*

spp. However, we find that the level of gene conservation within the FSSC (41%) is even lower

than the *Fusarium oxysporum* species complex (FOSC), whose core genome fraction has been

reported at 51-53% [53,54]. The proportion of core genes in the FSSC much also lower than

environmental human fungal pathogens such as *Aspergillus fumigatus* (69%) and *Cryptococcus*

neoformans (80%), but similar to another multi-kingdom pathogen, *Aspergillus flavus* (41-58%)

[55–58]. This may indicate that the FSSC and other multi-kingdom pathogens occupy a broader

niche space and the larger pan-genome has evolved to cope with a more variable set of

environmental conditions, as has been found in bacteria [59]. It is also tempting to hypothesize

that this expanded genomic repertoire and the resulting flexibility underpin the ability of these

organisms to infect multiple kingdoms, with very different metabolic environments and immune

systems; a point that requires further investigation. On the other hand, comparing pan-genomes

across studies is not without its own limitations [60]. It is complicated by differences in the number

of isolates between studies, the pangenome methodology (*e.g.* how orthologous genes are

identified and what fraction of isolates must contain a gene for it to be considered 'core'), and

whether the study defines taxonomic groups using *forma specialis* or species complex definitions.

Previous work comparing a phytopathogenic *F. oxysporum* strain with two *F. oxysporum* isolates

from human disease isolates from humans suggested that the latter may have unique accessory

chromosomes. While all FSSC genomes we examined had accessory chromosomes with higher

repeat content, as has been observed for many *Fusarium* species complexes [6,9,61,62], we did

not find clear evidence of accessory chromosome content differed from that of plant or other non-

clinical sources. We did not find any accessory scaffolds unique to human isolates, with the

exception of accessory chromosomes found only in F. *petroliphilum*. However, since all isolates

of this species in our dataset were of human origin, they lack a phylogenetically matched

comparator. As sequencing becomes even more accessible, we expect that larger datasets of

isolates from plant and human disease will help to further investigate the important question of

whether there are specific FSSC genomic elements associated with human infection.

The genomes of *F. keratoplasticum* were enriched in various genes related to branched-chain

amino-acid metabolism, carbon-nitrogen hydrolysis, and acetolactate activity, the latter of which

catalyzes the first step of the BCAA biosynthetic pathway (BCAA) [63]. BCAA synthesis is

conserved in plants, bacteria, and fungi, but not in higher metazoans [64], and has been shown

to be essential for fungal growth and virulence [35,36], making it a potential target for antifungal

drug development [65]. *F. keratoplasticum* also possessed an enrichment in genes relating to

oxidoreductase activity and base-excision repair, both potentially relevant to the prevention and

repair of DNA damage as a result of host reactive oxygen species [41]. In comparison,

F. petroliphilum was enriched in glutathione metabolism and sulphuric ester hydrolysis.

Glutathione serves as as a crucial oxidative stress protectant in fungi, the biosynthesis of which

has in *F. graminearum* been directly linked to sulphur metabolism [40], while the expansion in

helicase and endonuclease genes could be relevant for the repair of oxidatively damaged DNA.

Although these gene functional enrichments shed light on the different mechanisms by which

these pathogens may cause disease, or resist and repair damage resulting from ROS-mediated

host defenses.

The identification of BGCs for squalestatin, fusarubin, oxyjavanicin, metachelin, and

sansalvamide in all isolates is consistent with previous work exploring the secondary metabolite

diversity of the FSSC using different species [66], suggesting that these are core secondary

metabolites of the complex. However, this study identified the potential siderophores ferricrocin

and malonichrome, as additional core secondary metabolites, while these BGCs were not

predicted for any isolates in our dataset. Conversely, all isolates analyzed in this study encode

ochratoxin A, which was not reported in their study. Also intriguing was the observation that
BGC19, an unknown natural product of isocyanide class, was found in five of the six clinical
isolates but none of the plant or marine animals isolates. Isocyanides have been described in
virulence factors in both bacterial and fungal pathogens of insects and humans [67,68]. These
natural products are involved in copper uptake, a micronutrient that is often limited during
infection, and isocyanide BGCs are upregulated upon copper starvation [69]. We speculate that
this BGC might play a role in copper scavenging during keratitis, but this remains to be
experimentally tested.

lncRNAs are increasingly recognized to play important roles in regulating pathogenicity and
virulence in fungal pathogens of plants and humans [70–72]. However, they have not been
defined in the FSSC or any *Fusarium* species that infects humans. The list of lncRNAs described
in this study will be an important starting point to determine their roles in regulating basic cellular
processes and potentially plant and human virulence. The fact that many lncRNAs are located on
core chromosomes and are conserved between species indeed suggests that they are likely to
have conserved and essential functions. On the other hand, the non-conserved lncRNAs on
accessory chromosomes will be exciting to explore as potential virulence factors in plants and
animals.

In closing, the FSSC is an excellent system to study the molecular mechanisms and potential
evolutionary trade-offs associated with multi-kingdom pathogenicity. This is increasingly important
as climate change is expected to increase the number of environmental fungal species that can
grow at human body temperatures, potentially opening the door for other phytopathogens to
become human pathogens. The genomes and strains provided here are an excellent resource for
unraveling multi-kingdom virulence, such as comparing the transcriptomes of these isolates
during plant and mammalian infection to identify *in planta* virulence strategies that are repurposed
for human pathogenesis.

414 Materials and methods

Strains analyzed in this study

Six isolates originating from cases of keratitis collected by the German National Reference Centre
for Invasive Fungal Infection (NRZMyk) were sequenced in this work. Prior to whole genome
sequencing, species identification was performed by sequencing of the TEF-1α gene as in [28].

The additional FSSC genomes analyzed in this study originate from plant and aquatic animal
hosts and were generated in [6]. The accession numbers of the FSSC genomes analyzed in this
study and those included in the phylogeny are available in Supplementary File 1.
422

Nucleic acid isolation and sequencing
For the isolates sequenced in this study, mycelia were grown shaking at 30°C in complete media.
Mycelia were ground in liquid nitrogen, heat shocked for 1 minute at 95°C, and cellular debris
precipitated using steps and reagents from the MasterPure Yeast DNA Purification Kit (Lucigen).
DNA was finally purified using the Monarch Genomic DNA Purification Kit (New England Biolabs)
for binding and elution.
429

Genome assembly and annotation, including lncRNA annotation
The genomes generated in this study were assembled using the PacBio long reads and Flye
v2.9.1. The draft assemblies then underwent two rounds of polishing using Pilon v1.24 and the
strain-matched Illumina data. Assemblies were then scaffolded using RagTag v2.1.0 with
minimum confidence score set to 0.5, minimum location confidence score set to 0.35, and the
minimum orientation confidence score set to 0.6. The *F. keratoplasticum* genomes were
scaffolded using *F. keratoplasticum* LHS11 as a reference as it was the most contiguous genome
available for the species. As there was no existing chromosome-level genome available for
F. petroliphilum, their genomes were scaffolded using LHS14 because it resulted in the highest
mean location and orientation confidence scores of the FSSC isolates analyzed in this study.
Sequence contamination was identified and removed from genome assemblies using Foreign
Contaminant Screen (FSC) v0.4.0.
442

Before annotation, assemblies were cleaned to remove contigs shorter than 1kb and those with
>95% percent identity and >95% coverage with other contigs in the assembly. A custom FSSC
repeat library was generated using RepeatModeler v2.0.4 and the FSSC genome assemblies
analyzed in this study. RepeatMasker v4.1.2 and the FSSC repeat library was then used to identify
and mask repetitive regions. Genomes were annotated using Funannotate v1.8.9 and all
subsequent software referenced uses the versions packaged in this containerized version. Briefly,
an *ab initio* FSSC gene prediction model was trained using Augustus and the assembled
transcriptome of *Fusarium sp. FSSC11* available from Mycocosm as the RNAseq data generated
in our study was not available yet. Gene prediction was then performed using EvidenceModeler
incorporating predictions from Augustus, GeneMarkES, Snap, and GlimmerHMM. tRNAs were

predicted using tRNAScanSE. When strain-specific RNAseq data became available later, gene
models were corrected using the update function of Funannotate, which also added 5' and 3'
UTRs to the gene predictions. Functional annotations were assigned using PFAM, Eggnog-
mapper, InterProScan, CAZYmes, BUSCO (dikarya_odb9 lineage), SignalP and KEGG.
EffectorP 3.0 was used to predict secreted effectors. To ensure consistent genome annotations
across the study dataset, the genome assemblies from Hoh *et al.* 2022 [6] were downloaded from
NCBI and annotated as above, with the Augustus model trained using the RNAseq data
generated in the same publication. Genome completeness in Table 1 was determined using
BUSCO v5.2.2 and the hypocreales_odb10 lineage set.

To annotate lncRNAs de novo, we used the RNA-seq data from *F. ketatoplasticum* NRZ-2014-
069 and *F. petroliphilum* NRZ-2014-079 generated in this study. Genome-guided transcriptome
assembly was performed using Stringtie v.2.2.1 [73] and the individual transcript assemblies from
each species unified into a single transcript assembly using the Stringtie -merge option. To identify
novel transcripts, assembled transcripts were compared against protein-encoding transcripts
using Gffcompare v0.12.6 and novel transcripts >200 bp selected. Where applicable, the longest
isoform was chosen utilizing gtf2gtf from CGAT apps v0.6.0 [73,74]. Coding potential prediction
was performed using two software tools: CPC2 v3 [75], which was executed against the UniProt
database (https://www.uniprot.org/download), and Feelnc v0.2 [76]. Transcripts were classified
as lncRNAs when identified as non-coding transcripts by both tools. lncRNA transcripts were
categorized as intergenic (class code 'u') or antisense (class code 'x') based on the Gffcompare
output. To evaluate the expression levels of lncRNAs in Supplementary Figure 4, read counts for
lncRNA and protein-coding transcripts were calculated using Featurecounts v2.0.1 [77].
Subsequently, count data was normalized by transcript length and library size, yielding transcripts
per million (TPM) values.

Whole-genome phylogeny
The *Fusarium* spp phylogeny was inferred 1,164 single copy orthologs identified in all species
genomes using OrthoFinder v2.5.4. Amino acid sequences for each orthogroup were aligned
using MUSCLE v3.8, the alignments trimmed using ClipKIT v1.3.0, and then concatenated into
an alignment of length 614,941. The phylogeny was constructed using IQ-TREE v2.2.0.3.
ModelFinder within IQ-TREE was used to identify JTT+F+I+GR as the best fitting substitution
model based on Bayesian information criterion (BIC). Bootstrapping was performed using 1,000
ultrafast bootstraps and all branches were supported by values of >98%.

Pan-genome analysis and identification of core vs. accessory chromosomes and contigs

GENESPACE v1.3.1, incorporating Orthofinder v2.5.4 and MCScanX v01.11.2022, was used for
synteny-guided orthologous gene identification and their presence-absence between strains.
Heap's Law calculation was done using the MicroPan v2.1 R package using 500 permutations.
Chromosomes and contigs were defined as core and accessory based on alignment using the
nucmer tool of MUMmer v4.0.0. For these analyses, *F. keratoplasticum* Fu6, was used as the
reference query. Core chromosomes were identified using a cutoff of >92.5% mean identity
across a minimum alignment length of 1.75 Mb in all strains and were in agreement with those
identified in Hoh et al. 2022 [6]. The remaining chromosomes and contigs, which were not found
in all 12 study strains or showed <92.0% identify, were designated as accessory.
498

To compare the intergenic lncRNAs of *F. keratoplasticum* and *F. petroliphilum*, we used NRZ-
2014-069 and NRZ-2014-079 as representative examples of each species. Cross-species
lncRNA conservation was assessed using three approaches: BLASTn (local alignment), MUMmer
v.4.0.0 (global alignment) and orthology-based synteny. First, we generated two FASTA files
containing the sequences of intergenic lncRNA in *F. keratoplasticum* and *F. petroliphilum*. The
pairwise comparison of reciprocal hit using BLASTn and dnadiff tool from the MUMmer4 suite [78]
were filtered to get those with e-value > 1e-03 and a coverage > 50%. In parallel, a synteny based
orthology approach developed and validated in Pegueroles *et al*. 2019 [34] was performed. We
implemented the custom pipeline with modification to fit the analysis for two species instead of
four, and it was executed using the parameters '3 3 1'. This configuration involved analyzing three
protein-coding genes on either side of a specific lncRNA, requiring at least three common genes
for each pairwise comparison between species, and ensuring a minimum of one common gene
on each side of the lncRNA. To visualize the alignment coordinates, we used Circos v.0.69 to plot
the 1-to-1 coordinates (coordinates available in Supplementary File 3).
513

Functional enrichment annotation and BGC analysis

For determining the functional differences between FSSC genomes, genes were converted into
matching ontology terms and summed for each strain. ANOVA and post-hoc Dunnett test were
then applied to identify annotations over- or under-represented in a single species, using a
Bonferroni-correct p-value cutoff of 0.05. For visualization purposes per-species expression
values were normalized to z-scores, representing the number of standard deviations above/below
the mean.

Biosynthetic gene clusters were predicted for each isolate genome using the fungal version of
antiSMASH v7.0, and unknown metabolites were grouped together across isolates using BiG-
SCAPE v1.1.5. For each predicted BGC cluster, percent identity was calculated via BLASTp.

RNA isolation, sequencing, and transcriptome analysis

Two representative keratitis samples, *F. keratoplasticum* NRZ-2014-069 and *F. petroliphilum*
NRZ-2014-079 were selected for transcriptomic analysis under a set of five growth conditions:
mycelia grown for 12h in complete media (5.0 g/L ammonium sulfate + 10.0 g/L glucose) at 25°C,
carbon-starvation media (0.1 g/L glucose), elevated temperature (34°C), and ungerminated
conidia harvested after 3 days' growth at 28°C. RNAs from two replicates were isolated using a
phenol-chloroform protocol. Briefly, harvested mycelia were pelleted and bead-beaten in a mix of
lysis buffer and acid phenol:chloroform:isoamyl alcohol (PCI). Lysate was centrifuged, upper
phase transferred, then centrifuged again in fresh PCI. Upper phase was incubated at -20 degrees
in sodium acetate and 2-propanol, pelleted, then twice ethanol-washed. The washed pellet was
air-dried and resuspended in RNAse-free dH_2O, then stored at -80 degrees.

Illumina TruSeq stranded library preparation and sequencing was performed by the University of
Würzburg Core Unit Systems Medicine on the NextSeq platform. Raw FastQ data files were
trimmed via TrimGalore v0.6.10, quality threshold ≤ 20. Filtered reads were mapped using STAR
v2.7.11b, counted using RSEM v1.3.3 and differentially expressed genes were identified using
the intersection of DEGs identified by DESeq2 v1.40.2, Limma v3.56.2, and EdgeR v3.42.4, with
DEGs that met the threshold of FDR p-adj < 0.01 for all callers being retained.

For the functional analysis of differentially expressed genes, gene set enrichment analysis was
conducted using the eponymous software GSEA (v.4.3.2) from Broad Institute. GSEA was run in
preranked mode, analyzing a gene list of differentially-expressed genes ranked by expression
log-fold change, against a background of every gene ontology term in the genome. Enriched or
depleted terms were filtered by FDR < 0.25, as recommended by Broad Institute in their
documentation.

Acknowledgements

Funding

This project was supported by The Federal Ministry for Education and Science (Bundesministerium für Bildung und Forschung, BMBF, bmbf.de) within the framework of InfectControl 2020 project FINAR 2.0 (grant no. 03ZZ0834) and by the Interdisciplinary Center for Infection Research (IZKF; ukw.de) at the University Hospital Würzburg (project D-420). AEB and MP are funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, dfg.de) under Germany's Excellence Strategy – EXC 20151 – Project-ID 390813860). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data and regent availability.

The strains sequenced in this study are deposited in and publicly available from the Jena Microbial Resource Collection (JMRC). The PacBio, Illumina WGS, and RNAseq data generated in this study are deposited in the NCBI SRA under BioProject PRJNA1137914. The additional *Fusarium* genomes analyzed in this study are publicly available under the accession numbers listed in Supplementary File 1.

Author Contributions

AEB, RM, and OK conceptualized the study. AZ and RM performed experimental investigations. PJTB, LMF, MP, and AEB performed formal analysis. PJTB, LMF, and AEB were responsible for visualization and writing the original draft. All authors were involved in reviewing and editing the manuscript.

 Tables & Figures

	Scaffolds (n)	N50 (Mb)	Genome Size (Mb)	Protein coding genes	BUSCO completeness	Repeat content	Isolation Source	Reference
Fusarium falciforme Fu3	14	3.88	53.6	15,051	99.1%	10.1%	Hawksbill sea turtle	[6]
Fusarium keratoplasticum Fu6	25	4.21	49.5	14,050	99.0%	7.1%	Hawksbill sea turtle	[6]
Fusarium keratoplasticum LHS11	26	3.50	53.5	15,777	98.9%	7.2%	Harbor seal	[6]
Fusarium keratoplasticum NRZ-2014-069	70	4.16	49.6	14,398	98.9%	4.8%	Keratitis (human eye)	This study
Fusarium keratoplasticum NRZ-2015-087	53	4.02	50.6	14,387	98.9%	7.1%	Keratitis (human eye)	This study
Fusarium keratoplasticum NRZ-2016-116	55	4.24	50.5	14,979	98.9%	6.5%	Keratitis (human eye)	This study
Fusarium petroliphilum NRZ-2014-013	120	3.98	51.2	15,060	99.1%	6.7%	Keratitis (human eye)	This study
Fusarium petroliphilum NRZ-2015-035	143	3.34	48.0	14,212	98.5%	4.5%	Keratitis (human eye)	This study
Fusarium petroliphilum NRZ-2014-079	95	3.33	48.5	14,541	99.0%	4.8%	Keratitis (human eye)	This study
Fusarium sp. FSSC12 LHS14	35	3.82	56.3	14,512	99.1%	11.2%	Roughtail stingray	[6]
Fusarium sp. Ph1	24	3.30	51.5	14,109	99.0%	8.6%	Orchid flower	[6]
Fusarium vanettenii Fs6	39	3.36	72.9	16,755	99.1%	14.2%	Pea plant	[6]

Table 1: Genomic characteristics of FSSC isolates included in this study.
BUSCO completeness quantified as the percent of BUSCOs identified as single copies using the
hypocreales_odb10.2019-11-20 ortholog set.

Figure 1. Phylogenetic reconstruction of *Fusarium* spp, including the FSSC isolates
analyzed in this study

Phylogeny constructed from an amino acid sequence of 1,866 single copy orthologs. Pink box
indicates the *Fusarium solani* species complex. The strains with highly contiguous long-read
genome assemblies included in this study are indicated in bold. Emoji indicates the source for the
FSSC isolates that belong to the same clade as the study dataset members. All branches
supported by ultrafast bootstrap values >98%.

Figure 2. The pan-genome of the FSSC, *Fusarium keratoplasticum*, and *F. petroliphilum*.
(a, d, e). Pan-gene presence-absence matrix of FSSC (a), *F. keratoplasticum* (d), and
F. petroliphilum, where gene presence is indicated with blue and gene absence in black. Colored
bars at the top of the matrices indicate the number of isolates each pangene was present in. (b)
Histogram of pangene frequency in the FSSC, calculated as the number of genomes in which
each pan-gene is present. (c) Pan-gene accumulation plot quantifying the relationship between
the number of genomes analyzed and the number of pan-genes identified (orange) and the
fraction of which were core (grey). Dots represent the individual values from 24 permutations.

599
600 **Figure 3. Sequence conservation of core chromosomes of the FSSC.** Gray-scale fill indicates
601 the nucleotide percent identity between the core chromosome sequence of *F. keratoplasticum*
602 Fu6 and other FSSC members, which are indicated with the colored boxes. Red lines indicate
603 presence of transposable and other repetitive elements.

Figure 4. Conservation of accessory chromosomes and contigs from *Fusarium keratoplasticum* and *Fusarium petroliphilum*. (a-b) Accessory chromosomes and contigs of (a) *F. keratoplasticum* or (b) *F. petroliphilum* strains and their sequence conservation with other FSSC isolates. Shading indicates percent identity across the syntenic block between the indicated isolate and accessory chromosome/contig indicated, versus the best matching chromosome or contig from the other isolates. For simplicity, the subject (reference sequence) for each comparison is omitted from the plot as it would be a solid black band. Red lines indicate the location of transposable elements and other repetitive elements. Only genomic fragments >150 kb are displayed.

615
**Figure 5. Heatmap of over and underrepresented gene ontology (GO) terms in *Fusarium*
keratoplasticum and *petroliphilum* compared to other members of the FSSC.** Significantly
over- or under-represented GO terms for *F. keratoplasticum* and *F. petroliphilum* identified by t-
test and Bonferroni-corrected p < 0.05. Heat map fill represents the Z-scaled number of genes
per genome annotated with each term. GO terms of particular interest or highlighted in the text
are written in bold.

Figure 6. Comparison of secondary metabolite biosynthetic gene clusters reveals a core set of secondary metabolites in the FSSC. BGCs with a known predicted product are indicated in bold and those with unknown products labeled with their natural product class. Percent identity was calculated using amino acid sequence of the backbone enzymes. Red boxes indicate the isolate whose sequence was used as the query when calculating percent identity. The dendrogram on the left represents the phylogenetic relationship of the isolates.

630
Figure 7. Comparison of enriched and depleted Gene Ontology terms in the transcriptomes
of *F. keratoplasticum* and *F. petroliphilum*. Experiments were conducted using NRZ-2014-069
for *F. keratoplasticum* and NRZ-2014-079 for *F. petroliphilum*. Differential expression for the
conditions shown was calculated in comparison to mycelia grown in complete media at 25
degrees. Color indicates enrichment score calculated by gene set enrichment analysis. Stars
indicate strains for which q < 0.25, as recommended by Broad Institute for GSEA.

Accessory chromosomes and contigs

637
Supplementary Figure 1. Conservation of accessory chromosomes and contigs from
Fusarium falciforme, Fusarium sp. FSS12, Fusarium sp. Ph1 and Fusarium vanettenii. (a-
d) Accessory chromosomes and contigs of (a) *F. sp.* Ph1 (b) *F. falciforme* (c) *F. sp FSS12*, and
(d) *F. vanettenii* strains and their sequence conservation with other FSSC isolates. Shading
indicates percent identity across the syntenic block between the indicated isolate and accessory
chromosome/contig indicated, versus the best matching chromosome or contig from the other
isolates. Red lines indicate the location of transposable elements and other repetitive elements.

646
Supplementary Figure 2. Repeat content of core and accessory chromosomes. Repeat
content calculated as the number of repetitive bases masked compared to the total chromosome
length. Only chromosomes and contigs >150kb were included in the analysis. *F. keratoplasticum*
and *F. petrolphilum* boxplots represent aggregated chromosome data from n=3 genomes each;
all other species represent the chromosomes from n=1 genome. Significance determined by
Mann-Whitney U test. * denotes p<0.05, ** p<0.01, *** p<0.001.

653
654 **Supplementary Figure 3. Pairwise synteny of *F. keratoplasticum* Fu6 and the other study**
655 **isolates.** Colors indicate synthetic blocks shared between *F. keratoplasticum* Fu6 and the
656 indicated strain, with the individual chromosome colors representing the syntenic block's
657 chromosome number in Fu6. Only genomic chromosomes and contigs larger than 400 kbp are
658 displayed.

659
660
Supplementary Figure 4. Identification and molecular properties of lncRNAs in *Fusariarium*
keratoplasticum and Fusarium petroliphilum. (a) The number of intergenic and antisense
predicted for *F. keratoplasticum* (upper panel) and *F. petroliphilum* (bottom panel). (b) Density
plots showing the distribution of sequence length (left) and expression (right) between intergenic
lncRNAs, antisense lnRNAs, and protein-coding genes. Dashed lines represent the median.
Statistical significance calculated using the Mann-Whitney U test with Bonferroni correction.
Intergenic lncRNAs are significantly shorter than protein-coding genes (*F. keratoplasticum*: p <
0.0001, *F. petroliphilum*: p < 0.0001). Antisense lncRNAs do not differ significantly in length
compared to protein-coding genes. Both intergenic and antisense lncRNAs show lower
expression levels compared to protein-coding genes (p < 0.0001 for both classes of lncRNAs in
both species compared to protein coding genes.

Supplementary Figure 5. Synteny analysis of conserved intergenic lncRNAs in *Fusariarium*
***keratoplasticum* and *Fusarium petroliphilum*.** Circos plot depicts the syntenic relationships of
intergenic lncRNAs between *F. keratoplasticum* NRZ-2014-069 and *F. petroliphilum* NRZ-2014-
079. For a lncRNA to be considered syntenic, at least three common genes for each pairwise
comparison between species, with a minimum of one common gene on each side of the lncRNA
(parameters 3 3 1). Syntenic regions are visualized as black ribbons connecting segments
between the two species. Only genomic scaffolds >615 kb are displayed.

680
681 **Supplementary Figure 6. Synteny of genes in biosynthetic gene cluster 19 (BCG19).**
682 Alignment performed using BiG-SCAPE. PFAM domains in individual genes are indicated by
683 color.

Supplementary File List

Supplementary File 1. Accession numbers for the genome assemblies used in the phylogeny,
including the publicly available FSSC isolates analyzed.

Supplementary File 2. Gene Ontology terms significantly over- or underrperesneted in the
genomes of *F. keratoplasticum* and *F. petroliphilum*. File include the gene counts and z-scored
values depicted in Figure 5. Description of genome-wide ontology terms.

Supplementary File 3. Conservation of lncRNAs between *F. keratoplasticum* and *F. petroliphilum*
identified through nucleotide alignment of lncRNA sequences and synteny using flanking protein-
coding genes.

Supplementary File 4. Differentially expressed genes in conidia, under carbon starvation, and at
34°C relative to mycelia grown in complete media.Description of DEG file.

Supplementary File 5. Geneset enrichment analysis of differentially expressed genes in
F. keratoplasticum and *F. petroliphilum*.

Supplementary File 6. Newick phylogeny displayed in Figure 1.

Supplementary File 7. Core and accessory genes for the FSSC, *F. keratoplasticum* and
F. petroliphilum

References

1. Zhang N, O'Donnell K, Sutton DA, Nalim FA, Summerbell RC, Padhye AA, et al. Members of the Fusarium solani species complex that cause infections in both humans and plants are common in the environment. J Clin Microbiol. 2006;44: 2186–2190.

2. O'Donnell K, Sutton DA, Wiederhold N, Robert VAR, Crous PW, Geiser DM. Veterinary Fusarioses within the United States. J Clin Microbiol. 2016 [cited 17 May 2024]. doi:10.1128/jcm.01607-16

3. Coleman JJ. The Fusarium solani species complex: ubiquitous pathogens of agricultural importance. Mol Plant Pathol. 2016;17. doi:10.1111/mpp.12289

4. Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N, Nelson A. The global burden of pathogens and pests on major food crops. Nature Ecology & Evolution. 2019;3: 430–439.

5. Singh RP, Singh PK, Rutkoski J, Hodson DP, He X, Jørgensen LN, et al. Disease Impact on Wheat Yield Potential and Prospects of Genetic Control. Annu Rev Phytopathol. 2016;54: 303–322.

6. Hoh DZ, Lee H-H, Wada N, Liu W-A, Lu MR, Lai C-K, et al. Comparative genomic and transcriptomic analyses of trans-kingdom pathogen Fusarium solani species complex reveal degrees of compartmentalization. BMC Biol. 2022;20: 1–18.

7. Batista BG, de Chaves MA, Reginatto P, Saraiva OJ, Fuentefria AM. Human fusariosis: An emerging infection that is difficult to treat. Rev Soc Bras Med Trop. 2020;53. doi:10.1590/0037-8682-0013-2020

8. WHO fungal priority pathogens list to guide research, development and public health action. World Health Organization; 25 Oct 2022 [cited 23 Jul 2024]. Available: https://www.who.int/publications/i/item/9789240060241

9. Ma L-J, van der Does HC, Borkovich KA, Coleman JJ, Daboussi M-J, Di Pietro A, et al. Comparative genomics reveals mobile pathogenicity chromosomes in Fusarium. Nature. 2010;464: 367–373.

10. Zhang Y, Yang H, Turra D, Zhou S, Ayhan DH, DeIulio GA, et al. The genome of opportunistic fungal pathogen Fusarium oxysporum carries a unique set of lineage-specific chromosomes. Communications Biology. 2020;3: 1–12.

11. Genome-wide analysis of Fusarium verticillioides reveals inter-kingdom contribution of horizontal gene transfer to the expansion of metabolism. Fungal Genet Biol. 2019;128: 60–73.

12. Sieber CMK, Lee W, Wong P, Münsterkötter M, Mewes H-W, Schmeitzl C, et al. The Fusarium graminearum Genome Reveals More Secondary Metabolite Gene Clusters and Hints of Horizontal Gene Transfer. PLoS One. 2014;9: e110311.

13. Guarro J. Fusariosis, a complex infection caused by a high diversity of fungal species refractory to treatment. Eur J Clin Microbiol Infect Dis. 2013;32: 1491–1500.

14. van Diepeningen AD, Al-Hatmi AMS, Brankovics B, de Hoog GS. Taxonomy and Clinical
Spectra of Fusarium Species: Where Do We Stand in 2014? Current Clinical Microbiology
Reports. 2014;1: 10–18.

15. van Diepeningen AD, Feng P, Ahmed S, Sudhadham M, Bunyaratavej S, de Hoog GS.
Spectrum of Fusarium infections in tropical dermatology evidenced by multilocus
sequencing typing diagnostics. Mycoses. 2015;58: 48–57.

16. Short DPG, O'Donnell K, Zhang N, Juba JH, Geiser DM. Widespread occurrence of diverse
human pathogenic types of the fungus Fusarium detected in plumbing drains. J Clin
Microbiol. 2011;49: 4264–4272.

17. Garnica M, Nucci M. Epidemiology of Fusariosis. Curr Fungal Infect Rep. 2013;7: 301–305.

18. Hoenigl M, Jenks JD, Egger M, Nucci M, Thompson GR. Treatment of Fusarium Infection
of the Central Nervous System: A Review of Past Cases to Guide Therapy for the Ongoing
2023 Outbreak in the United States and Mexico. Mycopathologia. 2023;188: 973–981.

19. Strong N, Meeks G, Sheth SA, McCullough L, Villalba JA, Tan C, et al. Neurovascular
Complications of Iatrogenic Meningitis. N Engl J Med. 2024;390: 522–529.

20. García-Rodríguez G, Duque-Molina C, Kondo-Padilla I, Zaragoza-Jiménez CA, González-
Cortés VB, Flores-Antonio R, et al. Outbreak of Meningitis in Immunocompetent Persons
Associated With Neuraxial Blockade in Durango, Mexico, 2022-2023. Open Forum Infect
Dis. 2024;11: ofad690.

21. Gibert S, Edel-Hermann V, Gautheron E, Gautheron N, Sol J-M, Capelle G, et al. First
Report of Fusarium avenaceum, Fusarium oxysporum, Fusarium redolens, and Fusarium
solani Causing Root Rot in Pea in France. Plant Dis. 2022 [cited 5 Aug 2024].
doi:10.1094/PDIS-04-21-0833-PDN

22. Pérez-Hernández A, Porcel-Rodríguez E, Gómez-Vázquez J. Survival of Fusarium solani f.
sp. cucurbitae and Fungicide Application, Soil Solarization, and Biosolarization for Control
of Crown and Foot Rot of Zucchini Squash. Plant Dis. 2017 [cited 5 Aug 2024].
doi:10.1094/PDIS-06-16-0883-RE

23. Gómez J, Serrano Y, Pérez A, Porcel E, Gómez R, Aguilar MI. Fusarium solani f. sp.
cucurbitae, affecting melon in Almería Province, Spain. Australas Plant Dis Notes. 2014;9:
1–3.

24. Wang R-Y, Gao B, Li X-H, Ma J, Chen S-L. First Report of Fusarium solani Causing
Fusarium Root Rot and Stem Canker on Storage Roots of Sweet Potato in China. 2013
[cited 5 Aug 2024]. doi:10.1094/PDIS-06-13-0651-PDN

25. Short DPG, O'Donnell K, Geiser DM. Clonality, recombination, and hybridization in the
plumbing-inhabiting human pathogen Fusarium keratoplasticuminferred from multilocus
sequence typing. BMC Evol Biol. 2014;14: 1–14.

26. Vega-Gutiérrez TA, López-Orona CA, Molina-Cárdenas L, López-Urquídez GA, Payán-
Arzapalo MA, Tirado-Ramírez MA. First Report of Fusarium keratoplasticum Causing
Strawberry Root Rot in Sinaloa, Mexico. Plant Dis. 2023 [cited 17 May 2024].
doi:10.1094/PDIS-10-22-2441-PDN

27. Lu L-Y, Ou J-H, Hui RC-Y, Chuang Y-H, Fan Y-C, Sun P-L. High Diversity of Fusarium
Species in Onychomycosis: Clinical Presentations, Molecular Identification, and Antifungal
Susceptibility. Journal of Fungi. 2023;9: 534.

28. Walther G, Stasch S, Kaerger K, Hamprecht A, Roth M, Cornely OA, et al. Fusarium
Keratitis in Germany. J Clin Microbiol. 2017 [cited 25 Jun 2024]. doi:10.1128/jcm.00649-17

29. Peinado-Acevedo JS, Ramírez-Sánchez IC. Endocarditis by Fusarium keratoplasticum.
Mycopathologia. 2020;186: 131–133.

30. Phylogenetic relationships among members of the Fusarium solani species complex in
human infections and the descriptions of F. keratoplasticum sp. nov. and F. petroliphilum
stat. nov. Fungal Genet Biol. 2013;53: 59–70.

31. Mattick JS, Amaral PP, Carninci P, Carpenter S, Chang HY, Chen L-L, et al. Long non-
coding RNAs: definitions, functions, challenges and recommendations. Nat Rev Mol Cell
Biol. 2023;24: 430–447.

32. Kim W, Miguel-Rojas C, Wang J, Townsend JP, Trail F. Developmental Dynamics of Long
Noncoding RNA Expression during Sexual Fruiting Body Formation in Fusarium
graminearum. MBio. 2018;9. doi:10.1128/mBio.01292-18

33. Parra-Rivero O, Pardo-Medina J, Gutiérrez G, Limón MC, Avalos J. A novel lncRNA as a
positive regulator of carotenoid biosynthesis in Fusarium. Sci Rep. 2020;10: 1–14.

34. Pegueroles C, Iraola-Guzmán S, Chorostecki U, Ksiezopolska E, Saus E, Gabaldón T.
Transcriptomic analyses reveal groups of co-expressed, syntenic lncRNAs in four species
of the genus Caenorhabditis. RNA Biol. 2019;16: 320–329.

35. Kingsbury JM, Yang Z, Ganous TM, Cox GM, McCusker JH. Cryptococcus neoformans
Ilv2p confers resistance to sulfometuron methyl and is required for survival at 37 °C and in
vivo. Microbiology. 2004;150: 1547–1558.

36. Kingsbury JM, McCusker JH. Cytocidal amino acid starvation of Saccharomyces cerevisiae
and Candida albicans acetolactate synthase (ilv2Δ) mutants is influenced by the carbon
source and rapamycin. Microbiology. 2010;156: 929–939.

37. Oh Y, Franck WL, Han S-O, Shows A, Gokce E, Muddiman DC, et al. Polyubiquitin Is
Required for Growth, Development and Pathogenicity in the Rice Blast Fungus
Magnaporthe oryzae. PLoS One. 2012;7. doi:10.1371/journal.pone.0042868

38. Rodier M-H, El Moudni B, Kauffmann-Lacroix C, Daniault G, Jacquemin J-L. A Candida
albicans metallopeptidase degrades constitutive proteins of extracellular matrix. FEMS
Microbiol Lett. 1999;177: 205–210.

39. Silva BA, Souza-Gonçalves AL, Pinto MR, Barreto-Bergter E, Santos ALS.
Metallopeptidase inhibitors arrest vital biological processes in the fungal pathogen
Scedosporium apiospermum. Mycoses. 2011;54: 105–112.

40. Park J, Han JW, Lee N, Kim S, Choi S, Lee H-H, et al. Sulfur metabolism-mediated fungal
glutathione biosynthesis is essential for oxidative stress resistance and pathogenicity in the
plant pathogenic fungus Fusarium graminearum. MBio. 2024 [cited 20 Aug 2024].

doi:10.1128/mbio.02401-23

41. Oxidative responses and fungal infection biology. Semin Cell Dev Biol. 2019;89: 34–46.

42. Arana DM, Prieto D, Román E, Nombela C, Alonso-Monge R, Pla J. The role of the cell wall
in fungal pathogenesis. Microb Biotechnol. 2009;2: 308–320.

43. Garcia-Rubio R, de Oliveira HC, Rivera J, Trevijano-Contador N. The Fungal Cell Wall: , ,
and Species. Front Microbiol. 2019;10: 2993.

44. Identification of gene clusters associated with fusaric acid, fusarin, and perithecial pigment
production in Fusarium verticillioides. Fungal Genet Biol. 2012;49: 521–532.

45. Jihua Wei BW. Chemistry and bioactivities of secondary metabolites from the genus
Fusarium. Fitoterapia. 2020;146: 104638.

46. Dawson MJ, Farthing JE, Marshall PS, Middleton RF, O'neill MJ, Shuttleworth A, et al. THE
SQUALESTATINS, NOVEL INHIBITORS OF SQUALENE SYNTHASE PRODUCED BY A
SPECIES OF PHOMA I. TAXONOMY, FERMENTATION, ISOLATION, PHYSICO-
CHEMICAL PROPERTIES AND BIOLOGICAL ACTIVITY. J Antibiot . 1992;45: 639–647.

47. Kurobane I, Vining LC, Gavin Mcinnes A, Gerber NN. METABOLITES OF FUSARIUM
SOLANI RELATED TO DIHYDROFUSARUBIN. J Antibiot . 1980;33: 1376–1379.

48. Kundu A, Mandal A, Saha S, Prabhakaran P, Walia S. Fungicidal activity and molecular
modeling of fusarubin analogues from Fusarium oxysporum. Toxicol Environ Chem. 2020
[cited 31 May 2024]. doi:10.1080/02772248.2020.1770253

49. Costa JH, Wassano CI, Angolini CFF, Scherlach K, Hertweck C, Pacheco Fill T. Antifungal
potential of secondary metabolites involved in the interaction between citrus pathogens. Sci
Rep. 2019;9: 1–11.

50. Kato S, Motoyama T, Futamura Y, Uramoto M, Nogawa T, Hayashi T, et al. Biosynthetic
gene cluster identification and biological activity of lucilactaene from Fusarium sp. RK97-94.
Biosci Biotechnol Biochem. 2020;84: 1303–1307.

51. Suga H, Arai M, Fukasawa E, Motohashi K, Nakagawa H, Tateishi H, et al. Genetic
Differentiation Associated with Fumonisin and Gibberellin Production in Japanese Fusarium
fujikuroi. Appl Environ Microbiol. 2018 [cited 19 Jul 2024]. doi:10.1128/AEM.02414-18

52. Hossain MT, Khan A, Chung EJ, Rashid MH-O, Chung YR. Biological Control of Rice
Bakanae by an Endophytic Bacillus oryzicola YC7007. Plant Pathol J. 2016;32: 228–241.

53. Fayyaz A, Robinson G, Chang PL, Bekele D, Yimer S, Carrasquilla-Garcia N, et al. Hiding
in plain sight: Genome-wide recombination and a dynamic accessory genome drive
diversity in Fusarium oxysporum f.sp. ciceris. Proceedings of the National Academy of
Sciences. 2023;120: e2220570120.

54. van Westerhoven AC, Aguilera-Galvez C, Nakasato-Tagami G, Shi-Kunne X, de la Parte
EM, Chavarro-Carrero E, et al. Segmental duplications drive the evolution of accessory
regions in a major crop pathogen. New Phytol. 2024;242: 610–625.

55. Barber AE, Sae-Ong T, Kang K, Seelbinder B, Li J, Walther G, et al. Aspergillus fumigatus

pan-genome analysis identifies genetic variants associated with human infection. Nature
Microbiology. 2021;6: 1526–1536.

56.　McCarthy CGP, Fitzpatrick DA. Pan-genome analyses of model fungal species. Microbial
Genomics. 2019;5: e000243.

57.　Gangurde SS, Korani W, Bajaj P, Wang H, Fountain JC, Agarwal G, et al. Aspergillus
flavus pangenome (AflaPan) uncovers novel aflatoxin and secondary metabolite associated
gene clusters. BMC Plant Biol. 2024;24: 1–17.

58.　Anne Hatmaker E, Barber AE, Drott MT, Sauters TJC, Alastruey-Izquierdo A, Garcia-
Hermoso D, et al. Pathogenicity is associated with population structure in a fungal
pathogen of humans. bioRxiv. 2024. p. 2024.07.05.602241.
doi:10.1101/2024.07.05.602241

59.　Dewar AE, Hao C, Belcher LJ, Ghoul M, West SA. Bacterial lifestyle shapes pangenomes.
PNAS. 2024;121. doi:10.1073/pnas.2320170121

60.　Perrier M, Barber AE. Unraveling the genomic diversity and virulence of human fungal
pathogens through pangenomics. PLoS Pathog. 2024;20: e1012313.

61.　Möller M, Stukenbrock EH. Evolution and genome architecture in fungal plant pathogens.
Nat Rev Microbiol. 2017;15: 756–771.

62.　Fokkens L, Guo L, Dora S, Wang B, Ye K, Sánchez-Rodríguez C, et al. A Chromosome-
Scale Genome Assembly for the Fusarium oxysporum Strain Fo5176 To Establish a Model
Arabidopsis-Fungal Pathosystem. G3 Genes|Genomes|Genetics. 2020;10: 3549–3555.

63.　McCourt JA, Duggleby RG. Acetohydroxyacid synthase and its role in the biosynthetic
pathway for branched-chain amino acids. Amino Acids. 2006;31: 173–210.

64.　Steyer JT, Todd RB. Branched-chain amino acid biosynthesis in fungi. Essays in
Biochemsitry. 2023;67: 865–876.

65.　Richie DL, Thompson KV, Studer C, Prindle VC, Aust T, Riedl R, et al. Identification and
Evaluation of Novel Acetolactate Synthase Inhibitors as Antifungal Agents. Antimicrob
Agents Chemother. 2013;57: 2272.

66.　Pokhrel A, Coleman JJ. Inventory of the Secondary Metabolite Biosynthetic Potential of
Members within the Terminal Clade of the Species Complex. J Fungi (Basel). 2023;9.
doi:10.3390/jof9080799

67.　Lim FY, Won TH, Raffa N, Baccile JA, Wisecaver J, Rokas A, et al. Fungal Isocyanide
Synthases and Xanthocillin Biosynthesis in Aspergillus fumigatus. MBio. 2018 [cited 12 Aug
2024]. doi:10.1128/mbio.00785-18

68.　Harris NC, Sato M, Herman NA, Twigg F, Cai W, Liu J, et al. Biosynthesis of isonitrile
lipopeptides by conserved nonribosomal peptide synthetase gene clusters in
Actinobacteria. Proceedings of the National Academy of Sciences. 2017;114: 7025–7030.

69.　Won TH, Bok JW, Nadig N, Venkatesh N, Nickles G, Greco C, et al. Copper starvation
induces antimicrobial isocyanide integrated into two distinct biosynthetic pathways in fungi.
Nat Commun. 2022;13: 1–14.

70. Hovhannisyan H, Gabaldón T. The long non-coding RNA landscape of Candida yeast
pathogens. Nat Commun. 2021;12. doi:10.1038/s41467-021-27635-4

71. Kalem MC, Panepinto JC. Long Non-Coding RNAs in Cryptococcus neoformans: Insights
Into Fungal Pathogenesis. Front Cell Infect Microbiol. 2022;12: 858317.

72. Glad HM, Tralamazza SM, Croll D. The expression landscape and pangenome of long non-
coding RNA in the fungal wheat pathogen Zymoseptoria tritici. Microbial Genomics. 2023;9:
001136.

73. Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. StringTie
enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol.
2015;33: 290–295.

74. Sims D, Ilott NE, Sansom SN, Sudbery IM, Johnson JS, Fawcett KA, et al. CGAT:
computational genomics analysis toolkit. Bioinformatics. 2014;30: 1290–1291.

75. Kang Y-J, Yang D-C, Kong L, Hou M, Meng Y-Q, Wei L, et al. CPC2: a fast and accurate
coding potential calculator based on sequence intrinsic features. Nucleic Acids Res.
2017;45: W12–W16.

76. Wucher V, Legeai F, Hédan B, Rizk G, Lagoutte L, Leeb T, et al. FEELnc: a tool for long
non-coding RNA annotation and its application to the dog transcriptome. Nucleic Acids Res.
2017;45: e57.

77. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30: 923–930.

78. Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. MUMmer4: A fast
and versatile genome alignment system. PLoS Comput Biol. 2018;14: e1005944.