An annotated reference library for supporting DNA metabarcoding analysis of aquatic macroinvertebrates
in French freshwater environments

Authors : Paula GAUVIN*, David EME, Isabelle DOMAIZON, Frédéric RIMET

UMR CARRTEL, Université Savoie Mont-Blanc, INRAE, Pole R&D ECLA, 75bis av. de Corzent – CS 50511,
FR-74200, Thonon-les-Bains, France

RiverLY Research Unit, National Research Institute for Agriculture Food and Environment, (INRAE),
Villeurbanne, France

isabelle.domaizon@inrae.fr

frederic.rimet@inrae.fr

david.eme@inrae.fr

*Corresponding author

UMR CARRTEL, Université Savoie Mont-Blanc, INRAE, Pole R&D ECLA, 75bis av. de Corzent – CS 50511,
FR-74200, Thonon-les-Bains, France

Paula.gauvin@inrae.fr

Abstract

Freshwater ecosystems are increasingly threatened by human activities, leading to biodiversity loss and ecosystem degradation. Effective biodiversity monitoring, particularly through the use of aquatic macroinvertebrates as bioindicators, is crucial for assessing ecological health. While traditional morphological methods face limitations, DNA metabarcoding offers higher accuracy and efficiency in species identification using environmental DNA. However, the success of metabarcoding is contingent on the quality of reference libraries, which are often incomplete or biased. This study aims to construct a comprehensive COI-based DNA barcode library for freshwater macroinvertebrates in France, specifically targeting short gene regions amplified with fwhF2/fwhR2N primers, suitable for degraded DNA. Data from the PERLA database, Alpine lake surveys, and Opie-benthos were consolidated and refined using TAXREF to produce a final species checklist for France. The resulting library was analyzed for taxonomic completeness and genetic diversity. The checklist consists of 2,841 species across 10 phyla, with Arthropoda being the most diverse. Despite this, 56% of species lack COI-5P sequences in the Barcode of Life Data System (BOLD), with Mollusca and Nematoda showing particularly low coverage. Alignment challenges were identified for certain taxa, particularly within Coleoptera, Diptera, and Malacostraca. This work highlights the need for expanded sequencing efforts to improve barcode coverage and enhance the accuracy of biodiversity assessments in freshwater ecosystems.

Keywords: freshwater, COI, fwhF2 primers, macroinvertebrates, gap analysis, metabarcoding

Introduction

Freshwater ecosystems face escalating anthropogenic pressures threatening their functionality (Reid et al., 2019) and leading to significant biodiversity loss (Borgwardt et al., 2019; Young et al., 2016). In this context, biodiversity monitoring is crucial to provide relevant ecological diagnoses and support the management and conservation of these ecosystems and the vital services they provide. Aquatic macroinvertebrates are widely used as bioindicators due to their abundant presence, high species diversity and sensitivity to both anthropogenic and natural disturbances (Hering et al., 2006). These organisms primarily inhabit the littoral zone and play pivotal roles in community assembly and food web dynamics, contributing significantly to beta diversity and metacommunity structure across diverse aquatic habitats (May, 2019). Consequently, they serve as key indicators in environmental monitoring programs, facilitating assessments of the ecological status of aquatic ecosystems as mandated by the EU Water Framework Directive (WFD) and national legislation (Mondy et al., 2012).

The standardized methodology for macroinvertebrate biomonitoring in Europe is based on morphological identification through binocular microscopy (ANFOR, 2010; CEN 2019). This methodology encounters significant challenges, including the labor-intensive and expensive nature of collecting and sorting individual benthic invertebrates, which hinders its broader adoption in routine biomonitoring (Blackman et al., 2019; Bonada et al., 2006). Moreover, routine morphological identification poses drawbacks such as high expertise demands and lower taxonomic resolution (Bean et al., 2017; Hering et al., 2018; Leese et al., 2016, 2018). Notably, many taxa can only be identified at limited taxonomic resolution (Caesar et al., 2006), without a timely effort of highly skilled taxonomists. This is the case for Chironomids, a common group in freshwaters, frequently classified at the family level due to challenges in species or genus identification (Beermann et al., 2018). Additionally, immature life stages lacking diagnostic morphological traits hinder accurate species identification and therefore limit the proper quality assessment of freshwater ecosystems (Sweeney et al., 2011). These challenges underscore the need for innovative approaches to enhance the efficiency and accuracy of macroinvertebrate biomonitoring practices.

The recent advent of DNA metabarcoding techniques, integrating amplicon barcoding with high-throughput sequencing, represents a promising advancement in biomonitoring applications (Carraro & Altermatt, 2024; Deiner et al., 2017; Pawlowski et al., 2018), providing a valuable complement to morphology-based approaches for species identifications. Metabarcoding finds application in analyzing environmental DNA (eDNA) samples obtained from water, biofilm or sediment (Sakata et al., 2020; Valentini et al., 2016, Rivera et al. 2021, Gauvin et al. submitted) or DNA extracts derived from a homogenate of the sample's fauna (Taberlet et al., 2012). These molecular innovations enable simultaneous processing of multiple samples, identification of small taxa, immature or larval stages, and offer increased sensitivity and specificity, often revealing hidden diversity, while enhancing time and cost effectiveness (Holman et al., 2019; Pochon et al., 2015; Shokralla et al., 2012). These advantages facilitate direct comparison among sites and studies and enable higher spatial-temporal frequency in monitoring due to increased throughput (Bush et al., 2019).

The efficiency of macroinvertebrates DNA metabarcoding relies heavily on primer sets' effectiveness across a broad taxonomic spectrum. The recovery rate of taxa using metabarcoding is contingent upon various factors, including the taxonomic resolution of the gene marker employed (e.g., COI or ribosomal markers like 16S, e.g, Elbrecht et al., 2016), amplicon length (Meusnier et al., 2008), primer universality, and the number of primer pairs utilized to amplify target taxonomic groups (Elbrecht & Leese, 2015; Gibson et al., 2015). Specifically, a segment of the cytochrome c oxidase subunit I (COI) gene has emerged as the standard DNA barcoding marker for most animal groups (Hebert, Ratnasingham, et al., 2003), with over 95% of species in diverse animal groups exhibiting distinctive COI sequences in test

assemblages (Hebert, Cywinska, et al., 2003; Hebert et al., 2004). Given the considerable phylogenetic
diversity among macroinvertebrates, the high taxonomic resolution and existing reference data for the
COI marker, make it a judicious choice for metabarcoding freshwater macroinvertebrate communities
(Andújar et al., 2018; Ratnasingham & Hebert, 2007). Furthermore, well-established barcoding gaps for
the COI marker in freshwater macroinvertebrates underscore its suitability for such applications (Zhou
et al., 2009). Numerous primers of COI gene of varying base pair sizes are available in the literature for
DNA amplification of macroinvertebrates communities. Environmental DNA released by target taxa can
degrade rapidly (Seymour et al., 2018). Consequently, for amplifying degraded DNA, that is extracted
from water samples for example, targeting a short COI marker region is suggested to enhance
amplification success (Barnes et al., 2014; Jelger Herder et al., 2014).

The effectiveness of DNA metabarcoding programs relies on open, comprehensive and accurate
reference sequence libraries (Briski et al., 2016; Oliveira et al., 2016; Weigand et al., 2019), ensuring
precise taxonomic assignment (Richardson et al., 2018; Rodriguez-Ezpeleta et al., 2021). In that sense,
incomplete DNA barcode libraries, as in poorly represented species, pose a significant challenge, often
leading to false negatives and compromising biodiversity assessments (Ardura, 2019; Duarte et al.,
2021; Leite et al., 2020). Therefore, evaluating these gaps and the quality of sequence data in reference
libraries is imperative for the effective implementation of DNA-based tools in biodiversity assessments
(Duarte et al., 2020). Given the significance of reference libraries completeness, primarily those that are
pertinent in the context of the WFD, numerous studies have been conducted to assess their
representativeness by comparing them with lists of described species (Csabai et al., 2023; Duarte et al.,
2020; Leite et al., 2020; Specchia et al., 2020; Trebitz et al., 2015; Weigand et al., 2019). Despite efforts
to assess their representativeness, biases in taxonomic coverage persist within reference libraries (Li et
al., 2019; Weigand et al., 2019). Indeed, numerous studies have emphasized the construction of tailored
reference libraries to suit the geographic context of the research area (Ficetola et al., 2021; Mugnai et
al., 2023) and emphasized the importance of such databases to be refined with custom sequences
specific to local study areas and free from unexpected taxa. Additionally, research by Abad et al., 2016
and Schenekar et al., 2020 has highlighted the value of possessing DNA barcodes for local species. These
studies have demonstrated that DNA barcodes from local organisms can improve the accuracy of
taxonomic assignments and reveal previously unrecognized biodiversity, leading to adjustments in
taxonomic classifications among species.

Our study is in line with the recommendations proposed by Blackman et al., 2023, which advocate for
compiling a comprehensive list of target species within the study area and assembling accurate
sequences pertinent to those species. Our first objective is to construct a DNA barcode library utilizing
COI primers, particularly focusing on a short region of this gene which is highly effective when targeting
degraded DNA (fwhF2/fwhR2N, 205pb, Vamos et al., 2017), to identify freshwater macroinvertebrate
fauna in France using metabarcoding techniques and to assess its genetic completeness. Our second
objective is to identify the gaps of this new reference library, thereby providing insight into its limitations
and facilitating data interpretation for futures users.

Material and methods

Checklist constitution for French metropolitan freshwater macroinvertebrates species

156   In order to establish a checklist of macroinvertebrates taxa of freshwater aquatic habitats in
157   metropolitan France, three checklists were complied. The first one comes from PERLA, an interactive
158   tool managed by a French regional environmental agency (DREAL Auvergne-Rhône-Alpes) accessible at

http://www.perla.developpement-durable.gouv.fr/index.php. PERLA serves as a national comprehensive checklist for water managers and encompasses larvae, nymphs and adults across various taxonomic groups, including insects, molluscs, crustaceans, and worms, among others, found in rivers and aquatic ecosystems. In the manuscript, we will refer to this checklist as the 'French aquatic ecosystems checklist'. The second checklist is coming from macroinvertebrate surveys conducted in four Alpine lakes (Lake Geneva, Lake Annecy, Lake Bourget and Lake Tignes) from 2015 to 2022. In the manuscript, we will refer to this checklist as the 'French Alpine lakes checklist'. The third checklist we used was from a French NGO, Opie-benthos (https://www.opie-benthos.fr/opie/monde-des-insectes.html), studying freshwater insect's taxonomy and diversity. Currently, Opie-benthos leads comprehensive inventories across France, specifically targeting insect species undergoing a part of their life cycle within diverse aquatic ecosystems. In the manuscript, we will refer to this checklist as the 'French aquatic insect's species list'.

These three checklists were merged into a single one. When the taxonomic resolution of taxa was limited to a level above species (genus, family, class, phylum), we consulted the taxonomic referential of the National Inventory of Natural Heritage (TAXREF v17.0 2024, available at https://inpn.mnhn.fr/telechargement/referentielEspece/taxref/17.0/menu) to select species from those ranks. The phylum Acanthocephala and Rotifera, the classes Copepoda and Ostracoda (Arthropoda, Crustacea) and the families Succineidae (Arthropoda, Insecta, Diptera) and Hydrachnidae (Arthropoda, Arachnida, Trombidiformes), initially listed in French aquatic ecosystems checklist, were omitted from our search from TAXREF v17.0 2024 as they are not considered as freshwater macroinvertebrates according Tachet et al., 2000. Once those macroinvertebrates species were retrieved for each taxonomic levels without species identification, several filters from TAXREF were applied to select only French metropolitan freshwater species from this taxonomic referential. First, we selected only the species occurring in the following habitats: freshwater, marine and freshwater, brackish water, continental (land and freshwater). Then, from this updated list, only species occurring in metropolitan France were kept. Finally, species characterized by one those 8 status over 15 status in total were selected (present (native or undetermined), endemic, sub-endemic, cryptogenic, introduced, invasive introduced, non-established introduced (including cultivated / domestic), occasional).

The checklists described above and the summarized species list resulting from these compilations, inclusive of all species from the mentioned inventories, is referred as the "French metropolitan freshwater macroinvertebrates species list" and all together are available at https://doi.org/10.57745/LMOXEW in the 'Checklists composition data' file for future reference.

Sequences origin and cleaning steps

All the sequences for those species listed in the French metropolitan freshwater macroinvertebrates species list, whatever the gene, were downloaded from October 2023 to May 2024, from the Barcode of Life Data System v4 -Bold- repository (https://boldsystems.org/index.php/databases). We opted to utilize Bold as reference library instead of GenBank due to concerns regarding the latter's unverified submission process, which frequently results in misannotated sequences (Kozlov et al., 2016; Locatelli et al., 2020; Steinegger & Salzberg, 2020).

The absence of sequences for a given species may indicate either that the search in BOLD returned "Unmatched terms" or that while the species was found, no sequences were publicly available, rendering them private sequences (regardless of the reason for their absence, they were all defined as "not available sequences" for this analysis). Before concluding that a species lacked records in BOLD for the species from the French aquatic ecosystems checklist and French Alpine lakes checklist, synonyms

were searched using INPN. The details of taxa resulting from this taxonomic harmonization are available
in Table S1.

From this file containing all the sequences retrieved (whatever the gene) for the queried species of the
French metropolitan freshwater macroinvertebrates species list, the COI-5P sequences belonging to the
COI gene were selected. All the remaining sequences were aligned with the fwhF2/fwhR2N primer pairs
using MAFFT version 7 (Auto) (https://mafft.cbrc.jp/alignment/server/), by genus or species group (as
in all sequences assigned to the same genus or specie were aligned together). Then, sequences
characterized by gaps, insufficiencies in length relative to the COI barcode standard (shorter than 205
pb), or identical (same genetic sequence for one species) from the sequences file, were removed using
Jalview (https://www.jalview.org/). Due to the possibility of identical sequences being shared among
different species, certain species may have been excluded during this stage, as a result of sequence
overlap with other taxa. To address this issue, the list of species obtained from this step was cross-
referenced with the initial species list following the alignment process. This comparison identified any
missing species, whose sequences were then realigned individually to ensure their accurate
representation. This final reference library containing only genetic sequences capable of being aligned
by the fwhF2/fwhR2N primer pair is named 'Aligned DNA library' hereafter.

Graphical analysis
221
We firstly analysed the taxonomic composition of the French metropolitan freshwater
macroinvertebrates species list. Secondly, to highlight any gaps in the availability of sequences in our
list of species, the taxonomic coverage of barcodes at various taxonomic levels was assessed. Thirdly,
we highlighted the taxonomic composition of species with COI-5P sequences that did not align with
fwhF2/fwhR2N primers to reveal which species and taxonomic groups could be misrepresented in
metabarcoding studies using those primer pairs. Then, the distribution of the number of sequences
available for each phylum was examined to explore the species genetic diversity. The categorization of
different capacities to examine genetic variability, ranging from limited (<5 barcodes) to moderate (5-
25 barcodes) to good (>25 barcodes), was adopted from Trebitz et al., 2015, and the species with the
highest number of unique sequences exceeding 100 were identified. Finally, in order to demonstrate
the ability of the primers used in the aligned DNA library to discriminate each species by a unique
sequence, groups of taxa sharing the same genotypes were identified.

Results and discussion
235

Taxonomic composition of the French metropolitan freshwater macroinvertebrates species list and barcoding coverage
236
237

The French metropolitan freshwater macroinvertebrates species list is composed by 10 phyla, 16
classes, 50 orders, 222 families, 670 genera, and 2841 species (Table 1). Figure 1 provides insights into
the taxonomic composition, highlighting that the highest diversity is observed within the phylum
Arthropoda, followed by Mollusca and Nematoda, with 2469, 195, and 99 species, respectively. On the
other hand, taxa such as Entoprocta and Nemerta are represented by only one species (Figure 1, Table
1). Among the insects, Coleoptera dominate, followed by Diptera and Trichoptera, with 684, 630, and
527 species, respectively. The predominant phyla with the highest number of species in our checklist
matches with those identified in a study by Specchia et al., 2020, who conducted a gap analysis of DNA
barcodes available in international repositories using the aquatic macroinvertebrate species checklist of
a south-eastern Apulia region in Italy. Furthermore, our findings align with the prevailing understanding

that Arthropoda, and thus insects, represent the most diverse group of animals, exerting dominance in freshwater ecosystems inventories (Choudhary & Ahi, 2015; Dijkstra et al., 2014; Grosberg et al., 2012; Yeates & Wiegmann, 1999). Moreover, Diptera emerge as the most species-rich group utilized in biomonitoring across Europe, with chironomids recognized for their prevalence and diversity in freshwater habitats (Pinder, 1986). Alongside Diptera, Coleoptera (beetles) stand out for their exceptionally high species numbers across various ecoregions and countries in Europe and serving as the most abundant group of aquatic insects (Jäch & Balke, 2008; Short, 2018). Following these orders, Trichoptera (caddisflies), Plecoptera (stoneflies) and Ephemeroptera (mayflies), collectively known as EPT, emerge as the next three orders in terms of species richness from our checklist. These organisms spend their immature stages in freshwater and are widely employed as biological indicators for freshwater quality assessment (Hering et al., 2004; Sweeney et al., 2011) and ecological investigations, demonstrating robust responses to pollution or climate change (Álvarez-Troncoso et al., 2015). Additionally, Nematoda emerges as a highly diverse group, ranking third in species richness among all phyla listed in our checklist. This outcome may be attributed to the the interest in this phylum is its utility in ecological assessments, as nematodes are used for this purpose since a long time (e.g. Bongers, 1990; Moreno et al., 2011).

From all the sequences retrieved based on the species of the French metropolitan freshwater macroinvertebrates species list, 85.4% were belonging to COI-5P gene (Figure S1). This result was expected as BOLD is the main repository for COI sequences (Ratnasingham & Hebert, 2007). Overall, 56% of species listed in the French metropolitan freshwater macroinvertebrates species list do not have any COI-5P genetic sequences publicly available in BOLD database. There are some exceptions with Nemerta, which have a barcode coverage of 100% and the Entoprocta which have 0% coverage (both phylum with only species referenced). Mollusca and Nematoda, which are the second and third most diverse phylum in terms of species in the French metropolitan freshwater macroinvertebrates species list, have one the worst barcode coverage of all phyla (41% and 5.1%, respectively). Within Arthropoda, most of the class have a barcode coverage above 60%, with the highest being the Malacostraca (73.8%) and the lowest the Branchiura (66.7%). Finally, within insects, Hymenoptera, Lepidoptera, Megaloptera, Odonata and Hemiptera have the highest barcode coverage (>80%). The three most species diverse insects orders (Coleoptera, Diptera, Trichoptera) have a barcode coverage of 73.4%, 55.9% and 67.9%, respectively (Figure 2). Compared to previous gap analyses carried out in other countries, our analysis in freshwater ecosystems in France revealed a slightly lower barcoding coverage for freshwater macroinvertebrates (56%). For instance, investigations in specific regions such as the Apulia Region of Southeast Italy reported DNA barcode availability for 58% of listed aquatic Macroinvertebrate species (Specchia et al., 2020), while a study in Atlantic Iberia documented coverage for 63% of macroinvertebrates (Leite et al., 2020). Similarly, a comprehensive assessment of 4502 freshwater invertebrate species utilized in ecological quality assessments indicated that 60% possessed one or more barcodes (Weigand et al., 2019). Our findings of the lowest barcode coverage at the phylum level align with previous studies, which also reported very low barcode coverage for freshwater Platyhelminthes, with only three species having sequences deposited in examined databases (two in our study). The low taxonomic coverage for Mollusca in our study can be attributed to the high number of DNA barcodes deposited in GenBank rather than BOLD. As our study exclusively utilized BOLD for sequence retrieval, this likely accounts for the observed low barcode coverage of Mollusca, despite it being the second most listed phylum in our checklist. Additionally, although nematodes are taxonomically diverse and ecologically significant, they are generally overlooked in existing surveys (Weigand et al., 2019 and references therein). Insects exhibited the highest number of available COI sequences, consistent with previous studies highlighting the overrepresentation of Arthropoda, particularly insects (Meglécz, 2023). The barcode coverage for Insecta was 69%, aligning with other

findings such as Weigand et al. (2019), who reported that 66% of monitored insect species were barcoded, and Trebitz et al. (2015), who found approximately 70% representation in BOLD for Great Lakes fauna. Within Insecta, Diptera had the lowest coverage on BOLD at 55.9%, while Odonata and Hemiptera were the best covered, with over 80% of species barcoded in each group, similar to Weigand et al. (2019). The large number of DNA barcodes accessible within public databases often reflects the intensity of dedicated studies and associated barcoding projects. Certain taxonomic groups, such as Arthropoda or Mollusca receive disproportionate attention, resulting in a heightened focus and increased deposition of sequences within genetic databases (Ardura, 2019; Briski et al., 2011, 2016). Consequently, the absence of comprehensive reference databases may lead to false-negative outcomes (Klymus et al., 2017), while inaccuracies within these databases can engender false-positive identifications, as evidenced by instances of misreporting species presence (Port et al., 2016). Furthermore, the false assignment of sequences to closely related species may ensue when references for the true species are absent (Couton et al., 2022; Schenekar et al., 2020). Resolving this issue necessitates to sequence new specimens of target taxa and their subsequent integration into reference libraries.

Gap analysis of the aligned DNA library

The figure 3 represents the taxonomic composition of the 27 species from the French metropolitan freshwater macroinvertebrates species list which cannot be aligned with the primer pairs fwhF2/fwhR2N. Arthropoda emerged as the predominant taxonomic groups facing alignment difficulties at the phylum level. Within insects (Arthropoda), Coleoptera and Diptera had the species with the most alignment issues. Specifically, within Malacostraca (Arthropoda), the gammarid (Gammaridae) and crayfish (Astacidea) families had the highest number of species with alignment issues (Figure 3). Several reasons could explain these results. Firstly, although COI-5P sequences were available, some were found to lie outside the primer pairs' intended region during the alignment process, either entirely or partially. In the latter case, where the sequence in question was shorter than the desired amplified barcode, it was subsequently discarded. Another possible explanation is the poor sequence annotation, such as mislabelling (e.g., COI-3P instead of COI5-P) or misidentification of specimens, leading to incorrect species assignments. Misidentification of voucher specimens has been highlighted as a major factor contributing to erroneous records, as morphological identifications of closely related species can be challenging. This issue has been noted by Leite et al., 2020; Paz & Rinkevich, 2021; Pentinsaari et al., 2020, emphasizing its impact on subsequent species identifications using databases like BOLD. This was evident in cases when some species of insects failed to align with the primer pairs, while others belonging to the same genus did. Despite BOLD being a curated database with verification procedures during sequence deposition and a reported error rate of less than 1% for Metazoan sequences at the genus level (Leray et al., 2019), problematic records may still exist, as evidenced in various studies on marine macroinvertebrates (Radulovici et al., 2021) where up to 39% of sequences were considered ambiguous. For the Decapoda species (Malacostraca, Arthropoda) that failed to align with the primers pairs fwhF2/fwhR2N, comprising eight crayfish species and one crab species, their absence in the mock community during primer design could account for this discrepancy (Elbrecht et al., 2017; Elbrecht & Leese, 2015). Although Gammarids (Amphipoda, Arthropoda) were included in primer design, the failure of some species to align with the primers could be due to their vast diversity within aquatic environments (Horton et al., 2023). Molecular studies on Amphipoda have revealed extensive species diversity and the presence of cryptic species complexes (Jażdżewska & Mamos, 2019), suggesting a high genetic variability within species (Lefébure et al., 2006) that may not be compatible with the primer pairs. Furthermore, Vamos et al., 2017 indicated that their analysis of

the efficiency of the primers pairs displayed higher penalty scores for certain taxa of Turbellaria, Mollusca, Trichoptera, and Isopoda, indicating potential underrepresentation due to primer bias. This aligns with findings from other studies suggesting preferential detection of taxonomic groups by different markers and primers (Leduc et al., 2019). Consequently, the incorporation of multiple genetic regions in DNA metabarcoding to ensure the broadest possible taxonomic detection may prove to be a good solution (Duarte et al., 2021). This could be particularly important to monitor the noble crayfish, *Astacus astacus* and the amphipod *Gammarus roeselii* which cannot be detected with fwhF2/fwhR2N primers, (Figure 3), whereas they are the most frequently monitored species of the malacostracans in Europe (Weigand et al., 2019).

Analysis of the number of sequences available for each species within different phyla (Figure 4) provides important information about the potential to detect taxa in natural samples. In our reference library, for most phyla, the majority of species have less than five barcodes, except cnidarians, which have the majority of their species with 5 to 25 barcodes. Only three species presented more than 100 unique sequences: one trichopteran (*Agraylea multipunctata*), one isopod (*Asellus aquaticus*) and one gasteropod (*Physella acuta*). These findings are consistent with the observations of Trebitz et al., 2015, suggesting a noteworthy exception to the prevailing low barcoding rate for invertebrates, with some species being exceptionally well genetically represented. This phenomenon may be attributed to the scientific significance and ecological relevance of these species, such as being acknowledged as a reliable indicator. Indeed, *Asellus aquaticus* is a common species monitored in European countries (Weigand et al., 2019), known as bioindicator for metal pollutants detection (e.g. O'Callaghan et al., 2019). *Physella acuta* is also intensively studied as it is an invasive aquatic Gasteropoda with worldwide distribution (Banha et al., 2014; Vinarski, 2017). The analysis on the ability of the fwhF2/fwhR2N primers to discriminate each species by a unique sequence demonstrated that 57 identical sequences were shared by 116 distinct species (Table S2). Among those sequences, 10 were shared by two or three different genus and 47 were shared by two or three different species. These results could be explained by the fact that some taxa could have been mislabelled or that for this length of barcode (205 pb), no genetic variability is found between two related taxa therefore this barcode is not suitable to decipher those species. Several authors showed the necessity to have multiple sequences for each species to cover correctly their haplotypic diversity (Keck et al., 2023; Leite et al., 2020). This imperative arises from the recognition that the absence of intraspecific variants can pose significant challenges. In instances where a single sequence is available for a species exhibiting high genetic variability, the accurate identification of all haplotypes may be compromised. Moreover, inadequate representation of closely related species within reference libraries can lead to the erroneous assignment of multiple species to a single taxonomic entity (e.g., Jackman et al., 2021), potentially resulting in erroneous assessments of species diversity.

Conclusions

Our study underscores a widespread absence of reference barcodes for numerous extant invertebrate species. Although barcoding offers advantages over morphological identification in biomonitoring, existing gaps in barcode libraries may hinder their effectiveness (Duarte et al., 2020; Feio et al., 2020; Hestetun et al., 2020; Vieira et al., 2021). Substantial efforts are necessary to sequence new individuals for species absent from in the reference barcoding library, and for species with a low number of sequences. This will enable a more efficient and reliable species identification and biodiversity assessment, since the effectiveness and reliability of DNA barcoding identification are linked to the thoroughness of taxonomic curation and completeness of the reference barcoding libraries (Geiger et al., 2021) (Keck et al. 2023). Nevertheless, our work has established a reference barcoding library for

freshwater macroinvertebrates at the French level. This database can be utilized for biodiversity studies employing environmental DNA with short COI primers (fwhF2/fwhR2n) designed for samples containing partially degraded DNA. Furthermore, we have provided insights into the various biases associated with the utilization of this library, that future users will have to take into account to interpret their results. Looking ahead, an important future development for the fwhF2/fwhRNn reference library would be to assign a confidence level to the identification of each taxon (e.g., species). This confidence could be determined based on several factors, including the number of reference sequences available (with higher sequence counts providing greater confidence), geographical coverage (broader coverage being preferable), species delimitation (with monophyletic groups offering more robust identification compared to paraphyletic ones), and the availability of metadata associated with each sequence (e.g., sampling date and location, habitat, sequence quality). These criteria align with recommendations proposed by Fontes et al., 2021.

Acknowledgments

This study was made possible thanks to a funding of the Carnot Eau&Environnement institute, a funding of the INRAE department AQUA and a funding of the pôle INRAE-OFB ECLA (ECosystèmes LAcustres).

References

Abad, D., Albaina, A., Aguirre, M., Laza-Martínez, A., Uriarte, I., Iriarte, A., Villate, F., & Estonba, A. (2016). Is metabarcoding suitable for estuarine plankton monitoring? A comparative study with microscopy. *Marine Biology*, *163*(7), 149. https://doi.org/10.1007/s00227-016-2920-0

Álvarez-Troncoso, R., Benetti, C. J., Sarr, A. B., Pérez-Bilbao, A., & Garrido, J. (2015). Impacts of hydroelectric power stations on Trichoptera assemblages in four rivers in NW Spain. *Limnologica*, *53*, 35–41. https://doi.org/10.1016/j.limno.2015.05.001

Andújar, C., Arribas, P., Yu, D. W., Vogler, A. P., & Emerson, B. C. (2018). Why the COI barcode should be the community DNA metabarcode for the metazoa. *Molecular Ecology*, *27*(20), 3968–3975. https://doi.org/10.1111/mec.14844

Ardura, A. (2019). Species-specific markers for early detection of marine invertebrate invaders through eDNA methods: Gaps and priorities in GenBank as database example. *Journal for Nature Conservation*, *47*, 51–57. https://doi.org/10.1016/j.jnc.2018.11.005

Banha, F., Marques, M., & Anastácio, P. M. (2014). *Dispersal of two freshwater invasive macroinvertebrates* , Procambarus clarkii *and* Physella acuta, *by off-road vehicles*. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *24*(5), 582–591. https://doi.org/10.1002/aqc.2453

Barnes, M. A., Turner, C. R., Jerde, C. L., Renshaw, M. A., Chadderton, W. L., & Lodge, D. M. (2014). Environmental Conditions Influence eDNA Persistence in Aquatic Systems. *Environmental Science & Technology*, *48*(3), 1819–1827. https://doi.org/10.1021/es404734p

Bean, T. P., Greenwood, N., Beckett, R., Biermann, L., Bignell, J. P., Brant, J. L., Copp, G. H., Devlin, M. J., Dye, S., Feist, S. W., Fernand, L., Foden, D., Hyder, K., Jenkins, C. M., Van Der Kooij, J., Kröger, S., Kupschus, S., Leech, C., Leonard, K. S., … Righton, D. (2017). A Review of the Tools Used for Marine Monitoring in the UK: Combining Historic and Contemporary Methods with Modeling and Socioeconomics to Fulfill Legislative Needs and Scientific Ambitions. *Frontiers in Marine Science*, *4*, 263. https://doi.org/10.3389/fmars.2017.00263

Beermann, A. J., Zizka, V. M. A., Elbrecht, V., Baranov, V., & Leese, F. (2018). DNA metabarcoding reveals the complex and hidden responses of chironomids to multiple stressors. *Environmental Sciences Europe*, *30*(1), 26. https://doi.org/10.1186/s12302-018-0157-x

Blackman, R. C., Walser, J., Rüber, L., Brantschen, J., Villalba, S., Brodersen, J., Seehausen, O., & Altermatt, F. (2023). General principles for assignments of communities from ᴇDNA: Open versus closed taxonomic databases. *Environmental DNA*, *5*(2), 326–342. https://doi.org/10.1002/edn3.382

Blackman, R., Mächler, E., Altermatt, F., Arnold, A., Beja, P., Boets, P., Egeter, B., Elbrecht, V., Filipe, A. F., Jones, J., Macher, J., Majaneva, M., Martins, F., Múrria, C., Meissner, K., Pawlowski, J., Schmidt Yáñez, P., Zizka, V., Leese, F., … Deiner, K. (2019). Advancing the use of molecular methods for routine freshwater macroinvertebrate biomonitoring – the need for calibration experiments. *Metabarcoding and Metagenomics*, *3*, e34735. https://doi.org/10.3897/mbmg.3.34735

Bonada, N., Prat, N., Resh, V. H., & Statzner, B. (2006). DEVELOPMENTS IN AQUATIC INSECT BIOMONITORING: A Comparative Analysis of Recent Approaches. *Annual Review of Entomology*, *51*(1), 495–523. https://doi.org/10.1146/annurev.ento.51.110104.151124

Bongers, T. (1990). The maturity index: An ecological measure of environmental disturbance based on nematode species composition. *Oecologia*, *83*(1), 14–19. https://doi.org/10.1007/BF00324627

Borgwardt, F., Robinson, L., Trauner, D., Teixeira, H., Nogueira, A. J. A., Lillebø, A. I., Piet, G., Kuemmerlen, M., O'Higgins, T., McDonald, H., Arevalo-Torres, J., Barbosa, A. L., Iglesias-Campos, A., Hein, T., & Culhane, F. (2019). Exploring variability in environmental impact risk from human activities across aquatic ecosystems. *Science of The Total Environment*, *652*, 1396–1408. https://doi.org/10.1016/j.scitotenv.2018.10.339

Briski, E., Cristescu, M. E., Bailey, S. A., & MacIsaac, H. J. (2011). Use of DNA barcoding to detect invertebrate invasive species from diapausing eggs. *Biological Invasions*, *13*(6), 1325–1340. https://doi.org/10.1007/s10530-010-9892-7

Briski, E., Ghabooli, S., Bailey, S. A., & MacIsaac, H. J. (2016). Are genetic databases sufficiently populated to detect non-indigenous species? *Biological Invasions*, *18*(7), 1911–1922. https://doi.org/10.1007/s10530-016-1134-1

Bush, A., Compson, Z. G., Monk, W. A., Porter, T. M., Steeves, R., Emilson, E., Gagne, N., Hajibabaei, M., Roy, M., & Baird, D. J. (2019). Studying Ecosystems With DNA Metabarcoding: Lessons From Biomonitoring of Aquatic Macroinvertebrates. *Frontiers in Ecology and Evolution*, *7*, 434. https://doi.org/10.3389/fevo.2019.00434

Caesar, R. M., Sörensson, M., & Cognato, A. I. (2006). Integrating DNA data and traditional taxonomy to streamline biodiversity assessment: An example from edaphic beetles in the Klamath ecoregion, California, USA. *Diversity and Distributions*, *12*(5), 483–489. https://doi.org/10.1111/j.1366-9516.2006.00237.x

Carraro, L., & Altermatt, F. (2024). *eDITH: An R-package to spatially project eDNA-based biodiversity across river networks with minimal prior information* [Preprint]. Ecology. https://doi.org/10.1101/2024.01.16.575835

Choudhary, A., & Ahi, J. (2015). *BIODIVERSITY OF FRESHWATER INSECTS: A REVIEW*.

Couton, M., Lévêque, L., Daguin-Thiébaut, C., Comtet, T., & Viard, F. (2022). Water eDNA metabarcoding is effective in detecting non-native species in marinas, but detection errors still

hinder its use for passive monitoring. *Biofouling*, *38*(4), 367–383.

https://doi.org/10.1080/08927014.2022.2075739

Csabai, Z., Čiamporová-Zaťovičová, Z., Boda, P., & Čiampor, F. (2023). 50%, not great, not terrible: Pan-

European gap-analysis shows the real status of the DNA barcode reference libraries in two

aquatic invertebrate groups and points the way ahead. *Science of The Total Environment*, *863*,

160922. https://doi.org/10.1016/j.scitotenv.2022.160922

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer, S., Bista,

I., Lodge, D. M., Vere, N., Pfrender, M. E., & Bernatchez, L. (2017). Environmental DNA

metabarcoding: Transforming how we survey animal and plant communities. *Molecular*

Ecology, *26*(21), 5872–5895. https://doi.org/10.1111/mec.14350

Dijkstra, K.-D. B., Monaghan, M. T., & Pauls, S. U. (2014). Freshwater Biodiversity and Aquatic Insect

Diversification. *Annual Review of Entomology*, *59*(1), 143–163.

https://doi.org/10.1146/annurev-ento-011613-161958

Duarte, S., Leite, B., Feio, M., Costa, F., & Filipe, A. (2021). Integration of DNA-Based Approaches in

Aquatic Ecological Assessment Using Benthic Macroinvertebrates. *Water*, *13*(3), 331.

https://doi.org/10.3390/w13030331

Duarte, S., Vieira, P. E., & Costa, F. O. (2020). Assessment of species gaps in DNA barcode libraries of

non-indigenous species (NIS) occurring in European coastal regions. *Metabarcoding and*

Metagenomics, *4*, e55162. https://doi.org/10.3897/mbmg.4.55162

Elbrecht, V., & Leese, F. (2015). Can DNA-Based Ecosystem Assessments Quantify Species Abundance?

Testing Primer Bias and Biomass—Sequence Relationships with an Innovative Metabarcoding

Protocol. *PLOS ONE*, *10*(7), e0130324. https://doi.org/10.1371/journal.pone.0130324

Elbrecht, V., Taberlet, P., Dejean, T., Valentini, A., Usseglio-Polatera, P., Beisel, J.-N., Coissac, E., Boyer,

F., & Leese, F. (2016). Testing the potential of a ribosomal 16S marker for DNA metabarcoding

of insects. *PeerJ*, *4*, e1966. https://doi.org/10.7717/peerj.1966

Elbrecht, V., Vamos, E. E., Meissner, K., Aroviita, J., & Leese, F. (2017). Assessing strengths and

weaknesses of DNA metabarcoding-based macroinvertebrate identification for routine stream

monitoring. *Methods in Ecology and Evolution*, *8*(10), 1265–1275.

https://doi.org/10.1111/2041-210X.12789

Feio, M. J., Garcia-Reventós, A., Ardura, A., Calapez, A. R., Pujante, A. M., Mortágua, A., Múrria, C.,

Diaz-de-Quijano, D., & Filipe, A. F. (2020). Advances in the use of molecular tools in ecological

and biodiversity assessment of aquatic ecosystems. *Limnetica*, *39*(1), 419–440.

https://doi.org/10.23818/limn.39.27

Ficetola, G. F., Boyer, F., Valentini, A., Bonin, A., Meyer, A., Dejean, T., Gaboriaud, C., Usseglio-

Polatera, P., & Taberlet, P. (2021). Comparison of markers for the monitoring of freshwater

benthic biodiversity through DNA metabarcoding. *Molecular Ecology*, *30*(13), 3189–3202.

https://doi.org/10.1111/mec.15632

Fontes, J. T., Vieira, P. E., Ekrem, T., Soares, P., & Costa, F. O. (2021). BAGS: An automated Barcode,

Audit & Grade System for DNA barcode reference libraries. *Molecular Ecology Resources*,

21(2), 573–583. https://doi.org/10.1111/1755-0998.13262

Geiger, M., Koblmüller, S., Assandri, G., Chovanec, A., Ekrem, T., Fischer, I., Galimberti, A., Grabowski,

M., Haring, E., Hausmann, A., Hendrich, L., Koch, S., Mamos, T., Rothe, U., Rulik, B., Rewicz, T.,

Sittenthaler, M., Stur, E., Tończyk, G., … Moriniere, J. (2021). Coverage and quality of DNA

barcode references for Central and Northern European Odonata. *PeerJ*, *9*, e11192.

https://doi.org/10.7717/peerj.11192

Gibson, J. F., Shokralla, S., Curry, C., Baird, D. J., Monk, W. A., King, I., & Hajibabaei, M. (2015). Large-
520   Scale Biomonitoring of Remote and Threatened Ecosystems via High-Throughput Sequencing.
521   *PLOS ONE*, *10*(10), e0138432. https://doi.org/10.1371/journal.pone.0138432

522   Grosberg, R. K., Vermeij, G. J., & Wainwright, P. C. (2012). Biodiversity in water and on land. *Current*
523   *Biology*, *22*(21), R900–R903. https://doi.org/10.1016/j.cub.2012.09.050

524   Hebert, P. D. N., Cywinska, A., Ball, S. L., & deWaard, J. R. (2003). Biological identifications through
525   DNA barcodes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*,
526   *270*(1512), 313–321. https://doi.org/10.1098/rspb.2002.2218

527   Hebert, P. D. N., Ratnasingham, S., & De Waard, J. R. (2003). Barcoding animal life: Cytochrome *c*
528   oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society*
529   *of London. Series B: Biological Sciences*, *270*(suppl_1). https://doi.org/10.1098/rsbl.2003.0025

530   Hebert, P. D. N., Stoeckle, M. Y., Zemlak, T. S., & Francis, C. M. (2004). Identification of Birds through
531   DNA Barcodes. *PLoS Biology*, *2*(10), e312. https://doi.org/10.1371/journal.pbio.0020312

532   Hering, D., Borja, A., Jones, J. I., Pont, D., Boets, P., Bouchez, A., Bruce, K., Drakare, S., Hänfling, B.,
533   Kahlert, M., Leese, F., Meissner, K., Mergen, P., Reyjol, Y., Segurado, P., Vogler, A., & Kelly, M.
534   (2018). Implementation options for DNA-based identification into ecological status
535   assessment under the European Water Framework Directive. *Water Research*, *138*, 192–205.
536   https://doi.org/10.1016/j.watres.2018.03.003

537   Hering, D., Johnson, R. K., Kramm, S., Schmutz, S., Szoszkiewicz, K., & Verdonschot, P. F. M. (2006).
538   Assessment of European streams with diatoms, macrophytes, macroinvertebrates and fish: A
539   comparative metric-based analysis of organism response to stress. *Freshwater Biology*, *51*(9),
540   1757–1785. https://doi.org/10.1111/j.1365-2427.2006.01610.x

541   Hering, D., Meier, C., Rawer-Jost, C., Feld, C. K., Biss, R., Zenker, A., Sundermann, A., Lohse, S., &
542   Böhmer, J. (2004). Assessing streams in Germany with benthic invertebrates: Selection of
543   candidate metrics. *Limnologica*, *34*(4), 398–415. https://doi.org/10.1016/S0075-
544   9511(04)80009-4

545   Hestetun, J. T., Bye-Ingebrigtsen, E., Nilsson, R. H., Glover, A. G., Johansen, P.-O., & Dahlgren, T. G.
546   (2020). Significant taxon sampling gaps in DNA databases limit the operational use of marine
547   macrofauna metabarcoding. *Marine Biodiversity*, *50*(5), 70. https://doi.org/10.1007/s12526-
548   020-01093-5

549   Holman, L. E., De Bruyn, M., Creer, S., Carvalho, G., Robidart, J., & Rius, M. (2019). Detection of
550   introduced and resident marine species using environmental DNA metabarcoding of sediment
551   and water. *Scientific Reports*, *9*(1), 11559. https://doi.org/10.1038/s41598-019-47899-7

552   Horton, T., De Broyer, C., Bellan-Santini, D., Coleman, C. O., Copilaş-Ciocianu, D., Corbari, L., Daneliya,
553   M. E., Dauvin, J.-C., Decock, W., Fanini, L., Fišer, C., Gasca, R., Grabowski, M., Guerra-García, J.
554   M., Hendrycks, E. A., Hughes, L. E., Jaume, D., Kim, Y.-H., King, R. A., … Zeidler, W. (2023). The
555   World Amphipoda Database: History and progress. *Records of the Australian Museum*, *75*(4),
556   329–342. https://doi.org/10.3853/j.2201-4349.75.2023.1875

557   Jäch, M. A., & Balke, M. (2008). Global diversity of water beetles (Coleoptera) in freshwater.
558   *Hydrobiologia*, *595*(1), 419–442. https://doi.org/10.1007/s10750-007-9117-y

559   Jackman, J. M., Benvenuto, C., Coscia, I., Oliveira Carvalho, C., Ready, J. S., Boubli, J. P., Magnusson, W.
560   E., McDevitt, A. D., & Guimarães Sales, N. (2021). eDNA in a bottleneck: Obstacles to fish
561   metabarcoding studies in megadiverse freshwater systems. *Environmental DNA*, *3*(4), 837–
562   849. https://doi.org/10.1002/edn3.191

Jażdżewska, A. M., & Mamos, T. (2019). High species richness of Northwest Pacific deep-sea
amphipods revealed through DNA barcoding. *Progress in Oceanography*, *178*, 102184.
https://doi.org/10.1016/j.pocean.2019.102184

Jelger Herder, A. Valentini, E. Bellemain, T. Dejean, J.J.C.W. Van Delft, P.F. Thomsen, & P. Taberlet.
(2014). *Environmental DNA - a review of the possible applications for the detection of*
(invasive) species. https://doi.org/10.13140/RG.2.1.4002.1208

Keck, F., Couton, M., & Altermatt, F. (2023). Navigating the seven challenges of taxonomic reference
databases in metabarcoding analyses. *Molecular Ecology Resources*, *23*(4), 742–755.
https://doi.org/10.1111/1755-0998.13746

Klymus, K. E., Marshall, N. T., & Stepien, C. A. (2017). Environmental DNA (eDNA) metabarcoding
assays to detect invasive invertebrate species in the Great Lakes. *PLOS ONE*, *12*(5), e0177643.
https://doi.org/10.1371/journal.pone.0177643

Kozlov, A. M., Zhang, J., Yilmaz, P., Glöckner, F. O., & Stamatakis, A. (2016). Phylogeny-aware
identification and correction of taxonomically mislabeled sequences. *Nucleic Acids Research*,
44(11), 5022–5033. https://doi.org/10.1093/nar/gkw396

Leduc, N., Lacoursière-Roussel, A., Howland, K. L., Archambault, P., Sevellec, M., Normandeau, E.,
Dispas, A., Winkler, G., McKindsey, C. W., Simard, N., & Bernatchez, L. (2019). Comparing
eDNA metabarcoding and species collection for documenting Arctic metazoan biodiversity.
Environmental DNA, *1*(4), 342–358. https://doi.org/10.1002/edn3.35

Leese, F., Altermatt, F., Bouchez, A., Ekrem, T., Hering, D., Meissner, K., Mergen, P., Pawlowski, J.,
Piggott, J., Rimet, F., Steinke, D., Taberlet, P., Weigand, A., Abarenkov, K., Beja, P., Bervoets, L.,
Björnsdóttir, S., Boets, P., Boggero, A., … Zimmermann, J. (2016). DNAqua-Net: Developing
new genetic tools for bioassessment and monitoring of aquatic ecosystems in Europe.
Research Ideas and Outcomes, *2*, e11321. https://doi.org/10.3897/rio.2.e11321

Leese, F., Bouchez, A., Abarenkov, K., Altermatt, F., Borja, Á., Bruce, K., Ekrem, T., Čiampor, F.,
Čiamporová-Zaťovičová, Z., Costa, F. O., Duarte, S., Elbrecht, V., Fontaneto, D., Franc, A.,
Geiger, M. F., Hering, D., Kahlert, M., Kalamujić Stroil, B., Kelly, M., … Weigand, A. M. (2018).
Why We Need Sustainable Networks Bridging Countries, Disciplines, Cultures and Generations
for Aquatic Biomonitoring 2.0: A Perspective Derived From the DNAqua-Net COST Action. In
Advances in Ecological Research (Vol. 58, pp. 63–99). Elsevier.
https://doi.org/10.1016/bs.aecr.2018.01.001

Lefébure, T., Douady, C. J., Gouy, M., & Gibert, J. (2006). Relationship between morphological
taxonomy and molecular divergence within Crustacea: Proposal of a molecular threshold to
help species delimitation. *Molecular Phylogenetics and Evolution*, *40*(2), 435–447.
https://doi.org/10.1016/j.ympev.2006.03.014

Leite, B. R., Vieira, P. E., Teixeira, M. A. L., Lobo-Arteaga, J., Hollatz, C., Borges, L. M. S., Duarte, S.,
Troncoso, J. S., & Costa, F. O. (2020). Gap-analysis and annotated reference library for
supporting macroinvertebrate metabarcoding in Atlantic Iberia. *Regional Studies in Marine*
Science, *36*, 101307. https://doi.org/10.1016/j.rsma.2020.101307

Leray, M., Knowlton, N., Ho, S.-L., Nguyen, B. N., & Machida, R. J. (2019). GenBank is a reliable
resource for 21st century biodiversity research. *Proceedings of the National Academy of*
Sciences, *116*(45), 22651–22656. https://doi.org/10.1073/pnas.1911714116

Li, J., Lawson Handley, L. J., Harper, L. R., Brys, R., Watson, H. V., Di Muri, C., Zhang, X., & Hänfling, B.
(2019). Limited dispersion and quick degradation of environmental DNA in fish ponds inferred
by metabarcoding. *Environmental DNA*, *1*(3), 238–250. https://doi.org/10.1002/edn3.24

Locatelli, N. S., McIntyre, P. B., Therkildsen, N. O., & Baetscher, D. S. (2020). GenBank's reliability is uncertain for biodiversity researchers seeking species-level assignment for eDNA. *Proceedings of the National Academy of Sciences*, *117*(51), 32211–32212. https://doi.org/10.1073/pnas.2007421117

May, M. L. (2019). Dispersal by Aquatic Insects. In K. Del-Claro & R. Guillermo (Eds.), *Aquatic Insects* (pp. 35–73). Springer International Publishing. https://doi.org/10.1007/978-3-030-16327-3_3

Meusnier, I., Singer, G. A., Landry, J.-F., Hickey, D. A., Hebert, P. D., & Hajibabaei, M. (2008). A universal DNA mini-barcode for biodiversity analysis. *BMC Genomics*, *9*(1), 214. https://doi.org/10.1186/1471-2164-9-214

Mondy, C. P., Villeneuve, B., Archaimbault, V., & Usseglio-Polatera, P. (2012). A new macroinvertebrate-based multimetric index (I2M2) to evaluate ecological quality of French wadeable streams fulfilling the WFD demands: A taxonomical and trait approach. *Ecological Indicators*.

Moreno, M., Semprucci, F., Vezzulli, L., Balsamo, M., Fabiano, M., & Albertelli, G. (2011). The use of nematodes in assessing ecological quality status in the Mediterranean coastal ecosystems. *Ecological Indicators*, *11*(2), 328–336. https://doi.org/10.1016/j.ecolind.2010.05.011

Mugnai, F., Costantini, F., Chenuil, A., Leduc, M., Gutiérrez Ortega, J. M., & Meglécz, E. (2023). Be positive: Customized reference databases and new, local barcodes balance false taxonomic assignments in metabarcoding studies. *PeerJ*, *11*, e14616. https://doi.org/10.7717/peerj.14616

O'Callaghan, I., Harrison, S., Fitzpatrick, D., & Sullivan, T. (2019). The freshwater isopod Asellus aquaticus as a model biomonitor of environmental pollution: A review. *Chemosphere*, *235*, 498–509. https://doi.org/10.1016/j.chemosphere.2019.06.217

Oliveira, L. M., Knebelsberger, T., Landi, M., Soares, P., Raupach, M. J., & Costa, F. O. (2016). Assembling and auditing a comprehensive DNA barcode reference library for European marine fishes. *Journal of Fish Biology*, *89*(6), 2741–2754. https://doi.org/10.1111/jfb.13169

Pawlowski, J., Kelly-Quinn, M., Altermatt, F., Apothéloz-Perret-Gentil, L., Beja, P., Boggero, A., Borja, A., Bouchez, A., Cordier, T., Domaizon, I., Feio, M. J., Filipe, A. F., Fornaroli, R., Graf, W., Herder, J., Van Der Hoorn, B., Iwan Jones, J., Sagova-Mareckova, M., Moritz, C., … Kahlert, M. (2018). The future of biotic indices in the ecogenomic era: Integrating (e)DNA metabarcoding in biological assessment of aquatic ecosystems. *Science of The Total Environment*, *637–638*, 1295–1310. https://doi.org/10.1016/j.scitotenv.2018.05.002

Paz, G., & Rinkevich, B. (2021). Gap analysis of DNA barcoding in ERMS reference libraries for ascidians and cnidarians. *Environmental Sciences Europe*, *33*(1), 4. https://doi.org/10.1186/s12302-020-00449-9

Pentinsaari, M., Ratnasingham, S., Miller, S. E., & Hebert, P. D. N. (2020). BOLD and GenBank revisited – Do identification errors arise in the lab or in the sequence libraries? *PLOS ONE*, *15*(4), e0231814. https://doi.org/10.1371/journal.pone.0231814

Pinder, L. C. V. (1986). Biology of Freshwater Chironomidae. *Annual Review of Entomology*, *31*(1), 1–23. https://doi.org/10.1146/annurev.en.31.010186.000245

Pochon, X., Zaiko, A., Hopkins, G. A., Banks, J. C., & Wood, S. A. (2015). Early detection of eukaryotic communities from marine biofilm using high-throughput sequencing: An assessment of different sampling devices. *Biofouling*, *31*(3), 241–251. https://doi.org/10.1080/08927014.2015.1028923

Radulovici, A. E., Vieira, P. E., Duarte, S., Teixeira, M. A. L., Borges, L. M. S., Deagle, B. E., Majaneva, S.,
Redmond, N., Schultz, J. A., & Costa, F. O. (2021). Revision and annotation of DNA barcode
records for marine invertebrates: Report of the 8th iBOL conference hackathon.
Metabarcoding and Metagenomics, *5*, e67862. https://doi.org/10.3897/mbmg.5.67862
Ratnasingham, S., & Hebert, P. D. N. (2007). bold: The Barcode of Life Data System
(http://www.barcodinglife.org). *Molecular Ecology Notes*, *7*(3), 355–364.
https://doi.org/10.1111/j.1471-8286.2007.01678.x
Reid, A. J., Carlson, A. K., Creed, I. F., Eliason, E. J., Gell, P. A., Johnson, P. T. J., Kidd, K. A., MacCormack,
T. J., Olden, J. D., Ormerod, S. J., Smol, J. P., Taylor, W. W., Tockner, K., Vermaire, J. C.,
Dudgeon, D., & Cooke, S. J. (2019). Emerging threats and persistent conservation challenges
for freshwater biodiversity. *Biological Reviews*, *94*(3), 849–873.
https://doi.org/10.1111/brv.12480
Richardson, R. T., Bengtsson-Palme, J., Gardiner, M. M., & Johnson, R. M. (2018). A reference
cytochrome c oxidase subunit I database curated for hierarchical classification of arthropod
metabarcoding data. *PeerJ*, *6*, e5126. https://doi.org/10.7717/peerj.5126
Rodriguez-Ezpeleta, N., Morissette, O., Bean, C. W., Manu, S., Banerjee, P., Lacoursière-Roussel, A.,
Beng, K. C., Alter, S. E., Roger, F., Holman, L. E., Stewart, K. A., Monaghan, M. T., Mauvisseau,
Q., Mirimin, L., Wangensteen, O. S., Antognazza, C. M., Helyar, S. J., Boer, H., Monchamp, M.,
… Deiner, K. (2021). Trade-offs between reducing complex terminology and producing
accurate interpretations from environmental DNA: Comment on "Environmental DNA: What's
behind the term?" by Pawlowski et al., (2020). *Molecular Ecology*, *30*(19), 4601–4605.
https://doi.org/10.1111/mec.15942
Sakata, M. K., Yamamoto, S., Gotoh, R. O., Miya, M., Yamanaka, H., & Minamoto, T. (2020).
Sedimentary eDNA provides different information on timescale and fish species composition
compared with aqueous eDNA. *Environmental DNA*, *2*(4), 505–518.
https://doi.org/10.1002/edn3.75
Schenekar, T., Schletterer, M., Lecaudey, L. A., & Weiss, S. J. (2020). Reference databases, primer
choice, and assay sensitivity for environmental metabarcoding: Lessons learnt from a re-
evaluation of an eDNA fish assessment in the Volga headwaters. *River Research and*
Applications, *36*(7), 1004–1013. https://doi.org/10.1002/rra.3610
Seymour, M., Durance, I., Cosby, B. J., Ransom-Jones, E., Deiner, K., Ormerod, S. J., Colbourne, J. K.,
Wilgar, G., Carvalho, G. R., De Bruyn, M., Edwards, F., Emmett, B. A., Bik, H. M., & Creer, S.
(2018). Acidity promotes degradation of multi-species environmental DNA in lotic mesocosms.
Communications Biology, *1*(1), 4. https://doi.org/10.1038/s42003-017-0005-3
Shokralla, S., Spall, J. L., Gibson, J. F., & Hajibabaei, M. (2012). Next-generation sequencing
technologies for environmental DNA research: NEXT-GENERATION SEQUENCING FOR
ENVIRONMENTAL DNA. *Molecular Ecology*, *21*(8), 1794–1805. https://doi.org/10.1111/j.1365-
294X.2012.05538.x
Short, A. E. Z. (2018). Systematics of aquatic beetles (Coleoptera): Current state and future directions.
Systematic Entomology, *43*(1), 1–18. https://doi.org/10.1111/syen.12270
Specchia, V., Tzafesta, E., Marini, G., Scarcella, S., D'Attis, S., & Pinna, M. (2020). Gap Analysis for DNA
Barcode Reference Libraries for Aquatic Macroinvertebrate Species in the Apulia Region
(Southeast of Italy). *Journal of Marine Science and Engineering*, *8*(7), 538.
https://doi.org/10.3390/jmse8070538

Steinegger, M., & Salzberg, S. L. (2020). Terminating contamination: Large-scale search identifies more
than 2,000,000 contaminated entries in GenBank. *Genome Biology*, *21*(1), 115.
https://doi.org/10.1186/s13059-020-02023-1
Sweeney, B. W., Battle, J. M., Jackson, J. K., & Dapkey, T. (2011). Can DNA barcodes of stream
macroinvertebrates improve descriptions of community structure and water quality? *Journal*
of the North American Benthological Society, *30*(1), 195–216. https://doi.org/10.1899/10-
016.1
Taberlet, P., Coissac, E., Hajibabaei, M., & Rieseberg, L. H. (2012). Environmental DNA:
ENVIRONMENTAL DNA. *Molecular Ecology*, *21*(8), 1789–1793. https://doi.org/10.1111/j.1365-
294X.2012.05542.x
Tachet, Richoux, Usseglio-Polatera, & Bournaud. (2000). *Invertébrés d'Eau Douce. Systématique,*
Biologie, Écologie.
Trebitz, A. S., Hoffman, J. C., Grant, G. W., Billehus, T. M., & Pilgrim, E. M. (2015). Potential for DNA-
based identification of Great Lakes fauna: Match and mismatch between taxa inventories and
DNA barcode libraries. *Scientific Reports*, *5*, 12162. https://doi.org/10.1038/srep12162
Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., Bellemain, E., Besnard, A.,
Coissac, E., Boyer, F., Gaboriaud, C., Jean, P., Poulet, N., Roset, N., Copp, G. H., Geniez, P.,
Pont, D., Argillier, C., Baudoin, J., … Dejean, T. (2016). Next-generation monitoring of aquatic
biodiversity using environmental DNA metabarcoding. *Molecular Ecology*, *25*(4), 929–942.
https://doi.org/10.1111/mec.13428
Vamos, E., Elbrecht, V., & Leese, F. (2017). Short COI markers for freshwater macroinvertebrate
metabarcoding. *Metabarcoding and Metagenomics*, *1*, e14625.
https://doi.org/10.3897/mbmg.1.14625
Vieira, P. E., Lavrador, A. S., Parente, M. I., Parretti, P., Costa, A. C., Costa, F. O., & Duarte, S. (2021).
Gaps in DNA sequence libraries for Macaronesian marine macroinvertebrates imply decades
till completion and robust monitoring. *Diversity and Distributions*, *27*(10), 2003–2015.
https://doi.org/10.1111/ddi.13305
Vinarski, M. V. (2017). The history of an invasion: Phases of the explosive spread of the physid snail
Physella acuta through Europe, Transcaucasia and Central Asia. *Biological Invasions*, *19*(4),
1299–1314. https://doi.org/10.1007/s10530-016-1339-3
Weigand, H., Beermann, A. J., Čiampor, F., Costa, F. O., Csabai, Z., Duarte, S., Geiger, M. F., Grabowski,
M., Rimet, F., Rulik, B., Strand, M., Szucsich, N., Weigand, A. M., Willassen, E., Wyler, S. A.,
Bouchez, A., Borja, A., Čiamporová-Zaťovičová, Z., Ferreira, S., … Ekrem, T. (2019). DNA
barcode reference libraries for the monitoring of aquatic biota in Europe: Gap-analysis and
recommendations for future work. *Science of The Total Environment*, *678*, 499–524.
https://doi.org/10.1016/j.scitotenv.2019.04.247
Yeates, D. K., & Wiegmann, B. M. (1999). CONGRUENCE AND CONTROVERSY: Toward a Higher-Level
Phylogeny of Diptera. *Annual Review of Entomology*, *44*(1), 397–428.
https://doi.org/10.1146/annurev.ento.44.1.397
Young, H. S., McCauley, D. J., Galetti, M., & Dirzo, R. (2016). Patterns, Causes, and Consequences of
Anthropocene Defaunation. *Annual Review of Ecology, Evolution, and Systematics*, *47*(1), 333–
358. https://doi.org/10.1146/annurev-ecolsys-112414-054142
Zhou, X., Adamowicz, S. J., Jacobus, L. M., DeWalt, R. E., & Hebert, P. D. (2009). Towards a
comprehensive barcode library for arctic life—Ephemeroptera, Plecoptera, and Trichoptera of

Churchill, Manitoba, Canada. *Frontiers in Zoology*, 6(1), 30. https://doi.org/10.1186/1742-
9994-6-30

771 **Figure 1**. Taxonomic composition of the French metropolitan freshwater macroinvertebrates species
772 list. Percentage of the number of species according to phylum (a), class within the arthropod phylum
773 (b) and order within the insect class (c). The number above each bar represents the total number of
774 species affiliated to each taxonomic rank.

Phylum	Number of				
	Class	Order	Family	Genus	Species
Annelida	1	2	6	18	40
Arthropoda	4	20	135	508	2469
Bryozoa	2	2	6	7	11
Cnidaria	1	2	3	3	7
Entoprocta	1	1	1	1	1
Mollusca	2	11	24	52	195
Nematoda	2	9	42	68	99
Nemertea	1	1	1	1	1
Platyhelminthes	1	1	3	7	12
Porifera	1	1	1	5	6

781 **Table 1**. Summary of taxonomic composition for each phylum of the the French metropolitan freshwater
782 macroinvertebrates species list.

Figure 2. Barcoding coverage of the French metropolitan freshwater macroinvertebrates species list.
The barcoding coverage gives for each phylum, for each class within Arthropoda and for each order
within Insecta the total percentage of species with no available COI-5P public sequence in BOLD. The
number above each bar represents the total number of species listed in the Freshwater
macroinvertebrate checklist.

Figure 3. Gap analysis of the aligned DNA library. Number of species associated with COI-5P genetic
sequences not alignable with fwh2 primer pairs at the phylum rank (a), within the Arthropoda phylum
(b) and within the Insecta and Malacostraca class (c, c').

Figure 4. Taxonomic composition at the class rank for the species from the aligned DNA library. Each box
at the phylum level represents the distribution of COI-5P sequences relative to the number of species.

Supplementary data Figure 1. Percentage of genetic sequences depending on the type of genes. Only
the genes representing at least 0,1% of the sequences are shown. All the others genes are grouped in
the « Other genes » bar. The numbers above each bar show the total number of sequences retrieved
from BOLD affiliated to each genes.

Original taxa name	Phylum	Class	Order	Family	Genus	Harmonized taxa name	Reasons
Alainites	Arthropoda	Insecta	Ephemeroptera	Baetidae	Baetis	*Baetis*	Synonym
Anaciaeschna isosceles	Arthropoda	Insecta	Odonata	Aeshnidae	Aeshna	*Aeshna isoceles*	Synonym
Anisops sardea	Arthropoda	Insecta	Hemiptera	Notonectidae	Anisops	*Anisops sardeus*	Synonym
Armiger	Mollusca	Gastropoda		Planorbidae	Armiger	*Gyraulus*	Old nomenclature
Bithynella	Mollusca	Gastropoda	Littorinimorpha	Bythinellidae	Bythinella	*Bythinella*	Wrong spelling
Brachycercus harisellus	Arthropoda	Insecta	Ephemeroptera	Caenidae	Brachycercus	*Brachycercus harrisellus*	Synonym
Cambarincola okadai	Annelida	Clitellata	Branchiobdellida	Cambarincolidae	Cambarincola	*Cambarincola montanus*	Synonym
Chimarra marginatus	Arthropoda	Insecta	Trichoptera	Philopotamidae	Chimarra	*Chimarra marginata*	Synonym
Chironomus thumni	Arthropoda	Insecta	Diptera	Chironomidae	Chironomus	*Chironomus thummi*	Synonym
Corophium curvispinum	Arthropoda	Malacostraca	Amphipoda	Corophiidae	Chelicorophium	*Chelicorophium curvispinum*	Synonym
Cyphon	Arthropoda	Insecta	Coleoptera	Scirtidae	Contacyphon	*Contacyphon*	Synonym
Gomphus flavipes	Arthropoda	Insecta	Odonata	Gomphidae	Stylurus	*Stylurus flavipes*	Synonym
Haemisphaera	Arthropoda	Insecta	Coleoptera	Hydrophilidae	Hemisphaera	*Hemisphaera*	Wrong spelling
Helodes	Arthropoda	Insecta	Coleoptera	Scirtidae	Elodes	*Elodes*	Synonym
Hemianax ephippiger	Arthropoda	Insecta	Odonata	Aeshnidae	Anax	*Anax ephippiger*	Synonym
Heptagenia sulfurea	Arthropoda	Insecta	Ephemeroptera	Heptageniidae	Heptagenia	*Heptagenia sulphurea*	Wrong spelling
Hippeutis complanata	Mollusca	Gastropoda		Planorbidae	Hippeutis	*Hippeutis complanatus*	Synonym
Ironiquia dubia	Arthropoda	Insecta	Trichoptera	Limnephilidae	Ironiquia	*Ironoquia dubia*	Wrong spelling
Jaera danubica	Arthropoda	Malacostraca	Isopoda	Janiridae	Jaera	*Jaera sarsi*	Synonym
Lasiocephala	Arthropoda	Insecta	Trichoptera	Lepidostomatidae	Lepidostoma	*Lepidostoma basale*	Synonym
Meladema coriaceum	Arthropoda	Insecta	Coleoptera	Dytiscidae	Meladema	*Meladema coriacea*	Synonym
Metanoea	Arthropoda	Insecta	Trichoptera	Limnephilidae	Drusus	*Drusus*	Synonym
Micrasema difficile	Arthropoda	Insecta	Trichoptera	Brachycentridae	Micrasema	*Micrasema vestitum*	Synonym
Nemurella picteti	Arthropoda	Insecta	Plecoptera	Nemouridae	Nemurella	*Nemurella pictetii*	Synonym
Nymphula stagnata	Arthropoda	Insecta	Lepidoptera	Crambidae	Nymphula	*Nymphula nitidulata*	Synonym
Oligoplectrum maculatum	Arthropoda	Insecta	Trichoptera	Brachycentridae	Brachycentrus	*Brachycentrus maculatus*	Synonym
Oligotrichia striata	Arthropoda	Insecta	Trichoptera	Phryganeidae	Oligotricha	*Oligotricha striata*	Wrong spelling
Parapsectra	Arthropoda	Insecta	Diptera	Chironomidae	Micropsectra	*Micropsectra*	Synonym
Phacopteryx brevipennis	Arthropoda	Insecta	Trichoptera	Limnephilidae	Anabolia	*Anabolia brevipennis*	Synonym
Plea leachi	Arthropoda	Insecta	Hemiptera	Pleidae	Plea	*Plea minutissima*	Synonym
Pothamanthus luteus	Arthropoda	Insecta	Ephemeroptera	Potamanthidae	Potamanthus	*Potamanthus luteus*	Wrong spelling
Somatochlora artica	Arthropoda	Insecta	Odonata	Corduliidae	Somatochlora	*Somatochlora arctica*	Wrong spelling
Suphrodytes dorsalis	Arthropoda	Insecta	Coleoptera	Dytiscidae	Hydroporus	*Hydroporus dorsalis*	Synonym
Thremma gallicium	Arthropoda	Insecta	Trichoptera	Uenoidae	Thremma	*Thremma gallicum*	Wrong spelling
Tricholeiochiton fagesi	Arthropoda	Insecta	Trichoptera	Hydroptilidae	Tricholeiochiton	*Tricholeiochiton fagesii*	Synonym
Ylodes	Arthropoda	Insecta	Trichoptera	Leptoceridae	Triaenodes	*Triaenodes*	Subgenus

Supplementary data Table 1. Taxonomic harmonization of the French metropolitan freshwater macroinvertebrates species list. This table lists the original taxa name given in the French metropolitan freshwater macroinvertebrates species list and their harmonized name. Reasons of this harmonization are specified (synonymy, wrong spelling, subgenus).

Phylum	Class	Order	Family	Genus	Species	Groups
Arthropoda	Insecta	Odonata	Coenagrionidae	Ischnura	*Ischnura elegans*	G1
Arthropoda	Insecta	Odonata	Coenagrionidae	Ischnura	*Ischnura genei*	G1
Arthropoda	Insecta	Odonata	Coenagrionidae	Ischnura	*Ischnura graellsii*	G1
Mollusca	Gastropoda	Littorinimorpha	Bythinellidae	Bythinella	*Bythinella anianensis*	G2
Mollusca	Gastropoda	Littorinimorpha	Bythinellidae	Bythinella	*Bythinella cebennensis*	G2
Arthropoda	Insecta	Hemiptera	Corixidae	Hesperocorixa	*Hesperocorixa castanea*	G3
Arthropoda	Insecta	Hemiptera	Corixidae	Hesperocorixa	*Hesperocorixa moesta*	G3
Arthropoda	Insecta	Coleoptera	Dytiscidae	Graphoderus	*Graphoderus cinereus*	G4
Arthropoda	Insecta	Coleoptera	Dytiscidae	Graphoderus	*Graphoderus zonatus*	G4
Arthropoda	Insecta	Diptera	Limoniidae	Dicranophragma	*Dicranophragma adjunctum*	G5
Arthropoda	Insecta	Diptera	Limoniidae	Dicranophragma	*Dicranophragma separatum*	G5
Arthropoda	Insecta	Diptera	Limoniidae	Dicranophragma	*Dicranophragma separatum*	G6
Arthropoda	Insecta	Diptera	Pediciidae	Tricyphona	*Tricyphona alticola*	G6
Arthropoda	Insecta	Diptera	Limoniidae	Dicranomyia	*Dicranomyia imbecilla*	G7
Arthropoda	Insecta	Diptera	Limoniidae	Dicranomyia	*Dicranomyia quadra*	G7
Arthropoda	Insecta	Trichoptera	Leptoceridae	Athripsodes	*Athripsodes braueri*	G8
Arthropoda	Insecta	Trichoptera	Leptoceridae	Athripsodes	*Athripsodes cinereus*	G8
Arthropoda	Insecta	Trichoptera	Polycentropodidae	Plectrocnemia	*Plectrocnemia laetabilis*	G9
Arthropoda	Insecta	Trichoptera	Polycentropodidae	Polycentropus	*Polycentropus intricatus*	G9
Arthropoda	Insecta	Trichoptera	Limnephilidae	Limnephilus	*Limnephilus fenestratus*	G10
Arthropoda	Insecta	Trichoptera	Limnephilidae	Limnephilus	*Limnephilus rhombicus*	G10
Arthropoda	Insecta	Coleoptera	Heteroceridae	Heterocerus	*Heterocerus flexuosus*	G11
Arthropoda	Insecta	Coleoptera	Heteroceridae	Heterocerus	*Heterocerus fossor*	G11
Arthropoda	Insecta	Coleoptera	Hydraenidae	Hydraena	*Hydraena gracilidelphis*	G12
Arthropoda	Insecta	Coleoptera	Hydraenidae	Hydraena	*Hydraena gracilis*	G12
Arthropoda	Insecta	Coleoptera	Dytiscidae	Graptodytes	*Graptodytes fractus*	G13
Arthropoda	Insecta	Coleoptera	Dytiscidae	Graptodytes	*Graptodytes ignotus*	G13
Arthropoda	Insecta	Coleoptera	Dytiscidae	Hydroporus	*Hydroporus palustris*	G14
Arthropoda	Insecta	Coleoptera	Dytiscidae	Hydroporus	*Hydroporus vagepictus*	G14
Arthropoda	Insecta	Coleoptera	Hydrophilidae	Chaetarthria	*Chaetarthria seminulum*	G15
Arthropoda	Insecta	Coleoptera	Hydrophilidae	Chaetarthria	*Chaetarthria similis*	G15
Arthropoda	Insecta	Coleoptera	Limnichidae	Pelochares	*Pelochares versicolor*	G15
Arthropoda	Insecta	Coleoptera	Hydrophilidae	Chaetarthria	*Chaetarthria similis*	G16
Arthropoda	Insecta	Coleoptera	Hydrophilidae	Chaetarthria	*Chaetarthria simillima*	G16
Arthropoda	Insecta	Trichoptera	Psychomyiidae	Metalype	*Metalype fragilis*	G17
Arthropoda	Insecta	Trichoptera	Psychomyiidae	Tinodes	*Tinodes assimilis*	G17
Arthropoda	Insecta	Diptera	Limoniidae	Limonia	*Limonia dilutior*	G18
Arthropoda	Insecta	Diptera	Limoniidae	Limonia	*Limonia hercegovinae*	G18
Arthropoda	Insecta	Trichoptera	Hydroptilidae	Hydroptila	*Hydroptila dampfi*	G19
Arthropoda	Insecta	Trichoptera	Hydroptilidae	Orthotrichia	*Orthotrichia tragetti*	G19
Arthropoda	Insecta	Coleoptera	Dytiscidae	Rhantus	*Rhantus bistriatus*	G20
Arthropoda	Insecta	Coleoptera	Dytiscidae	Rhantus	*Rhantus consputus*	G20
Arthropoda	Insecta	Trichoptera	Sericostomatidae	Sericostoma	*Sericostoma galeatum*	G21
Arthropoda	Insecta	Trichoptera	Sericostomatidae	Sericostoma	*Sericostoma personatum*	G21
Arthropoda	Insecta	Trichoptera	Hydroptilidae	Hydroptila	*Hydroptila dampfi*	G22
Arthropoda	Insecta	Trichoptera	Hydroptilidae	Orthotrichia	*Orthotrichia tragetti*	G22
Arthropoda	Insecta	Plecoptera	Nemouridae	Protonemura	*Protonemura beatensis*	G23
Arthropoda	Insecta	Plecoptera	Nemouridae	Protonemura	*Protonemura nitida*	G23
Arthropoda	Malacostraca	Amphipoda	Gammaridae	Echinogammarus	*Echinogammarus feminatus*	G24
Arthropoda	Insecta	Trichoptera	Hydropsychidae	Hydropsyche	*Hydropsyche exocellata*	G24
Arthropoda	Insecta	Coleoptera	Scirtidae	Elodes	*Elodes minuta*	G25
Arthropoda	Insecta	Coleoptera	Scirtidae	Elodes	*Elodes pseudominuta*	G25
Arthropoda	Insecta	Trichoptera	Limnephilidae	Anomalopterygella	*Anomalopterygella chauviniana*	G26
Arthropoda	Insecta	Trichoptera	Limnephilidae	Drusus	*Drusus nigrescens*	G26
Arthropoda	Insecta	Diptera	Limoniidae	Tasiocera	*Tasiocera fuscescens*	G27
Arthropoda	Insecta	Diptera	Limoniidae	Tasiocera	*Tasiocera murina*	G27
Arthropoda	Insecta	Trichoptera	Hydropsychidae	Hydropsyche	*Hydropsyche doehleri*	G28
Arthropoda	Insecta	Trichoptera	Limnephilidae	Potamophylax	*Potamophylax nigricornis*	G28
Mollusca	Gastropoda	Littorinimorpha	Bythinellidae	Bythinella	*Bythinella baudoni*	G29
Mollusca	Gastropoda	Littorinimorpha	Bythinellidae	Bythinella	*Bythinella rufescens*	G29

Mollusca	Gastropoda	Littorinimorpha	Bythinellidae	Bythinella	*Bythinella padiraci*	G30
Mollusca	Gastropoda	Littorinimorpha	Bythinellidae	Bythinella	*Bythinella reyniesii*	G30
Arthropoda	Insecta	Coleoptera	Hydraenidae	Hydraena	*Hydraena atrata*	G31
Arthropoda	Insecta	Coleoptera	Hydraenidae	Hydraena	*Hydraena testacea*	G31
Arthropoda	Insecta	Diptera	Limoniidae	Limonia	*Limonia phragmitidis*	G32
Arthropoda	Insecta	Diptera	Limoniidae	Limonia	*Limonia stigma*	G32
Arthropoda	Insecta	Coleoptera	Scirtidae	Contacyphon	*Contacyphon padi*	G33
Arthropoda	Insecta	Coleoptera	Scirtidae	Contacyphon	*Contacyphon variabilis*	G33
Arthropoda	Insecta	Trichoptera	Psychomyiidae	Tinodes	*Tinodes maclachlani*	G34
Arthropoda	Insecta	Trichoptera	Psychomyiidae	Tinodes	*Tinodes maculicornis*	G34
Arthropoda	Insecta	Trichoptera	Limnephilidae	Limnephilus	*Limnephilus binotatus*	G35
Arthropoda	Insecta	Trichoptera	Limnephilidae	Limnephilus	*Limnephilus bipunctatus*	G35
Arthropoda	Insecta	Hemiptera	Corixidae	Sigara	*Sigara limitata*	G36
Arthropoda	Insecta	Hemiptera	Corixidae	Sigara	*Sigara semistriata*	G36
Arthropoda	Insecta	Odonata	Aeshnidae	Aeshna	*Aeshna grandis*	G37
Arthropoda	Insecta	Odonata	Aeshnidae	Aeshna	*Aeshna juncea*	G37
Arthropoda	Insecta	Diptera	Limoniidae	Neolimnomyia	*Neolimnomyia batava*	G38
Arthropoda	Insecta	Diptera	Limoniidae	Pseudolimnophila	*Pseudolimnophila sepium*	G38
Arthropoda	Insecta	Diptera	Culicidae	Culiseta	*Culiseta alaskaensis*	G39
Arthropoda	Insecta	Diptera	Culicidae	Culiseta	*Culiseta morsitans*	G39
Arthropoda	Insecta	Diptera	Culicidae	Aedes	*Aedes caspius*	G40
Arthropoda	Insecta	Diptera	Culicidae	Aedes	*Aedes dorsalis*	G40
Arthropoda	Insecta	Diptera	Culicidae	Aedes	*Aedes dorsalis*	G41
Arthropoda	Insecta	Diptera	Culicidae	Aedes	*Aedes geniculatus*	G41
Arthropoda	Insecta	Diptera	Culicidae	Culiseta	*Culiseta alaskaensis*	G42
Arthropoda	Insecta	Diptera	Culicidae	Culiseta	*Culiseta annulata*	G42
Arthropoda	Insecta	Diptera	Culicidae	Culex	*Culex impudicus*	G43
Arthropoda	Insecta	Diptera	Culicidae	Culex	*Culex territans*	G43
Arthropoda	Insecta	Diptera	Limoniidae	Dicranomyia	*Dicranomyia nigristigma*	G44
Arthropoda	Insecta	Diptera	Limoniidae	Dicranomyia	*Dicranomyia stigmatica*	G44
Porifera	Demospongiae	Spongillida	Spongillidae	Ephydatia	*Ephydatia fluviatilis*	G45
Porifera	Demospongiae	Spongillida	Spongillidae	Ephydatia	*Ephydatia muelleri*	G45
Arthropoda	Malacostraca	Isopoda	Janiridae	Jaera	*Jaera istri*	G46
Arthropoda	Malacostraca	Isopoda	Janiridae	Jaera	*Jaera sarsi*	G46
Arthropoda	Insecta	Trichoptera	Leptoceridae	Ceraclea	*Ceraclea dissimilis*	G47
Arthropoda	Insecta	Trichoptera	Leptoceridae	Ceraclea	*Ceraclea sobradieli*	G47
Arthropoda	Insecta	Diptera	Limoniidae	Gonomyia	*Gonomyia abscondita*	G48
Arthropoda	Insecta	Diptera	Limoniidae	Gonomyia	*Gonomyia conoviensis*	G48
Arthropoda	Insecta	Coleoptera	Helophoridae	Helophorus	*Helophorus grandis*	G49
Arthropoda	Insecta	Coleoptera	Helophoridae	Helophorus	*Helophorus granularis*	G49
Arthropoda	Insecta	Diptera	Syrphidae	Melanogaster	*Melanogaster hirtella*	G50
Arthropoda	Insecta	Diptera	Syrphidae	Melanogaster	*Melanogaster parumplicata*	G50
Arthropoda	Insecta	Diptera	Limoniidae	Dicranomyia	*Dicranomyia autumnalis*	G51
Arthropoda	Insecta	Diptera	Limoniidae	Dicranomyia	*Dicranomyia lutea*	G51
Arthropoda	Insecta	Coleoptera	Scirtidae	Scirtes	*Scirtes hemisphaericus*	G52
Arthropoda	Insecta	Coleoptera	Scirtidae	Scirtes	*Scirtes orbicularis*	G52
Arthropoda	Insecta	Trichoptera	Philopotamidae	Wormaldia	*Wormaldia copiosa*	G53
Arthropoda	Insecta	Trichoptera	Philopotamidae	Wormaldia	*Wormaldia variegata*	G53
Arthropoda	Insecta	Diptera	Syrphidae	Chrysogaster	*Chrysogaster solstitialis*	G54
Arthropoda	Insecta	Diptera	Syrphidae	Chrysogaster	*Chrysogaster virescens*	G54
Arthropoda	Insecta	Diptera	Limoniidae	Dicranomyia	*Dicranomyia chorea*	G55
Arthropoda	Insecta	Diptera	Limoniidae	Dicranomyia	*Dicranomyia quadra*	G55
Arthropoda	Insecta	Trichoptera	Rhyacophilidae	Rhyacophila	*Rhyacophila dorsalis*	G56
Arthropoda	Insecta	Trichoptera	Rhyacophilidae	Rhyacophila	*Rhyacophila meridionalis*	G56
Arthropoda	Insecta	Diptera	Limoniidae	Euphylidorea	*Euphylidorea dispar*	G57
Arthropoda	Insecta	Diptera	Limoniidae	Euphylidorea	*Euphylidorea phaeostigma*	G57

Supplementary data Table 2. Taxonomic composition of species sharing the same COI sequences after alignment with fwhF2/fwhR2N primers. A total of 57 groups were identified, with one group corresponding to one identical genetic sequence identical common to two or three distinct species.