Identifying ancient antibiotic resistance genes in archaeological dental calculus

Francesca J. Standeven[1], Gwyn Dahlquist-Axe[1], Camilla F. Speller[2], Andrew Tedder[1], Conor J. Meehan[3]

[1]School of Chemistry and Biosciences, University of Bradford

[2]Department of Anthropology, University of British Columbia

[3]Department of Biosciences, Nottingham Trent University

Corresponding author: Conor Meehan conor.meehan@ntu.ac.uk

Keywords: Dental calculus, antimicrobial resistance, bacteria, antibiotic resistance genes, metal resistance genes, ancient DNA, bioinformatics, ancient metagenomics, oral microbiome.

Abstract

Research on ancient antimicrobial resistance is limited, and appropriate screening criteria for identifying antibiotic (ARGs) and metal resistance genes (MRGs) in archaeological samples are unclear. We assessed the impact of DNA damage and contamination on ARG and MRG detection in ancient metagenomic sequences. Starting from a set of modern oral metagenomic samples, we simulated diagenetic DNA damage as expected in ancient oral metagenomic samples. Then we estimated the impact of this damage on ARG and MRG prediction at different identity thresholds. We also examined 25 post-industrial (ca. 1850 – 1901) dental calculus samples before and after decontamination to study the rates of false positive (FP) and negative (FN) ARG and MRG predictions introduced by sample contamination. The tests showed that diagenetic damage does not significantly affect resistance gene detection, but contamination does. Furthermore, while high thresholds are advisable when feasible, overall identity thresholds do not significantly affect the rates of FPs and FNs. Additionally, comparing post-industrial and modern dental calculus revealed Tetracycline ARGs as dominant in both contaminated ancient samples and modern samples, and MLS (Macrolide, Lincosamide, and Streptogramins) ARGs as prevalent in historical samples before widespread antibiotic use.

Data summary

The simulated data were generated from 182 human oral biofilm samples, retrieved from the

European Nucleotide Archive (ENA project: PRJNA817430) (Anderson et al., 2023).

Additionally, real ancient (PRJEB1716 and PRJEB12831) and modern (PRJEB1716)

metagenomic sequences were selected from metagenomic datasets published by Standeven et

al. (2024).

Impact statement

Antimicrobial resistance (AMR) is a global health crisis. Studying the adaptability of

microorganisms over centuries allows us to understand key factors that contribute to the

survival and spread of antibiotic-resistant bacteria today. We know that antibiotic abuse is a

key driver of AMR; however, further study into specific environmental niches that promote

the evolution of antibiotic-resistant bacteria is important. For example, the extent to which

the oral microbiome facilitates the increase of certain antibiotic-resistant genes and the

impact of metal pollution on the spread of AMR. To investigate these key areas, it is essential

to examine oral microbiomes across time, providing a complete perspective on the evolution

of AMR. However, ancient metagenomics poses problems for the screening of antibiotic and

metal-resistant genes in ancient bacterial DNA due to nucleotide base damage and short-read

data. Through thorough threshold experimentation to establish optimal screening criteria for

ancient resistance gene identification, and by addressing gaps in knowledge of ancient

resistance genes, this research offers clinical significance to existing research and contributes

to the development of strategies aimed at easing the impact of AMR on public health.

Introduction

58
Antimicrobial resistance (AMR), declared a top global health threat by the World Health

Organisation (Organisation 2020), can be described as the evolution and adaptability of

microorganisms in stressful situations to improve their chances of survival. Stressful

circumstances might include threats from other microorganisms including bacteria, viruses,

phages, and parasites, as well as antibiotic compounds produced by plants or metals.

Antibiotic resistance is often caused by an increase in antibiotic load in the environment due

to antibiotic misuse, overuse, and underuse of antibiotics (Bombaywala et al. 2021) that cause

bacteria to evolve, adapt, and resist drugs created to reduce their spread or destroy them. This

process involves the acquisition of various antibiotic resistance genes (ARGs), or metal

resistance genes (MRGs), that allow bacteria to defend themselves against certain

compounds. Resistance can arise by horizontal gene transfer (HGT), which is a major cause

of AMR, or from mutations that result in the acquisition of resistance-conducting genes (von

Wintersdorff et al. 2016). There are different types of resistance genes, including acquired

(Arzanlou et al. 2017), intrinsic (Blair et al. 2015), proto (Singh et al. 2019), and silent

resistance (Singh et al. 2019).

The use of databases and consistent classification of ARGs is essential (Alcock *et al.* 2019),

and various bioinformatics software, databases, and data-sharing resources are available for

identifying, predicting, cataloguing, and analysing molecular sequences of ARGs (McArthur

and Tsang 2017). Various tools to detect ARGs from metagenomic sequences include

AMR++ (Bonin et al. 2023), CARD RGI (Alcock et al. 2023), ARG-ANNOT (Gupta et al.

2014), ResFinder (Bortolaia et al. 2020), and Meta-MARC (Lakin et al. 2019). There are also

tools made just for identifying metal and biocide-resistant MRGs such as BacMet (Pal et al.

2014); tools such as the AMR++ pipeline can screen for both ARGs and MRGs. These

pipelines all typically identify resistance genes by aligning DNA sequences or whole-

genomes to reference databases (Zankari et al. 2017; Alcock et al. 2019; Feldgarden et al.

2019; Bortolaia et al. 2020; Feldgarden et al. 2021; Florensa et al. 2022; NCBI 2024) to

detect genes and single nucleotide polymorphisms (SNPs). Tools that access a combination of

these databases, like AMR++, can help reduce the impact of database bias.

Microbial DNA recovered from archaeological or historical samples has a high potential to

inform ARG and MRG studies by providing a perspective into resistance gene prevalence

before widespread antibiotic use. However, there is little guidance within bioarcheology on

the right tools and databases to use for ancient ARG and MRG analyses. However, some tools

are known to perform well on DNA with ancient traits and some researchers may prefer

pipelines like AMR++ and ARG-ANNOT, which work effectively with short sequences. To

elaborate, ancient microbial DNA analysis is fraught with challenges, including poor quality,

limited quantity, and high contamination levels of ancient DNA (Gaeta 2021). aDNA is

typically damaged through common forms of chemical processes that degrade DNA over

long periods such as depurination, which causes changes in nucleotide base pairs (Dabney et

al. 2013) and can also cause fragmentation (Kistler et al. 2017). These structural issues of

aDNA have the potential to hinder bioinformatic tools from aligning nucleotides to reference

genomes, but it is not known exactly to what extent. Regarding contamination, it is strongly

advised that environmental and laboratory blanks decontaminate samples before screening for

resistance genes following research on ARGs identified in mummified remains (Santiago-

Rodriguez et al. 2018) that sparked criticism of the lack of decontamination protocols

(Eisenhofer and Weyrich 2018). Environmental decontamination is crucial because

archaeological skeletons can remain *in situ* for thousands of years, accumulating contaminant

DNA from the environment that may accrue similar damage patterns to the endogenous

DNA. Due to the widespread distribution of ARGs across landscapes, the surrounding soil

contains genetic material that was not originally part of the ancient remains.

Bioinformaticians also encounter challenges when determining suitable cut-off thresholds

such as e-values and sequence identity (Ji et al. 2022), and appropriate threshold selection for

ancient ARG identification is especially difficult. This is demonstrated in a study by

Rascovan et al. (2016) that explores ARGs in several previously published ancient

metagenomic datasets. The authors use both stringent (>90% similarity) and non-stringent

(*evalue* $<1E^{-10}$) parameters to screen for ARGs against the ARGANNOT database and

highlighted that three more groups of ARGs were identified using less strict thresholds. The

authors are aware that using less stringent parameters increases the risk of false positives

(FPs), but they argue that stricter thresholds pose an even greater risk of false negatives

(FNs). Except for the study by Rascovan et al. (2016), there is a lack of discussion regarding

the appropriate thresholds for analysing resistance genes in ancient samples.

Ancient resistance genes have been found in arctic soil (Perron et al. 2015), caves (Bhullar et

al. 2012; Pawlowski et al. 2016), and permafrost (D'Costa et al. 2011; Kashuba et al. 2017).

Although the study of ARGs is intriguing in these contexts, the concept of their antiquity is

not surprising (Relman and Lipsitch 2018; Kraemer et al. 2019) because bacteria have always

developed defence mechanisms to survive (Baron et al. 2018). Antibiotics are naturally

occurring (Van Goethem et al. 2018) and most antibiotics today are derived from natural

sources (Hutchings et al. 2019).

Ancient resistance genes in human microbiomes are intriguing because delving into their

ancient presence before the influence of extreme selective pressure allows us to track

evolutionary traits and uncover novel mechanisms of bacteria in clean models. For example,

Rascovan et al. (2016) examined 10[th] and 12[th]-century dental calculus and found a new class

B Beta-lactamase family showing high penicillinase activity. Ancient dental calculus

(fossilized dental plaque) is a promising source of resistance genes because they have been

shown to entrap oral microorganisms and their genes and preserve them over thousands of

years (Warinner et al. 2014; Weyrich et al. 2017; Hendy et al. 2018; Fotakis et al. 2020;

Neukamm et al. 2020).

Examining resistance genes in archaeological contexts helps us understand how significant

lifestyle changes, such as industrialisation, may have influenced the spread and prevalence of

specific resistance genes in human oral microbiomes. For example, smoke pollution in

Manchester, UK, during the Industrial Revolution was particularly severe, and the poorest

inhabitants were compelled to dwell amid the mills and factories in the north-easterly regions,

while some better-paid workers may have avoided the worst of the pollution (Mosley 2008).

Such a significant historical transition as this would likely increase the levels of MRGs in the

environment. It is also noteworthy that heavy metals are shown to operate as co-selecting

agents in the spread of antibiotic resistance in human pathogens across various environments

(Nguyen et al. 2019). Moreover, the coexistence of ARGs, metal resistance genes (MRGs),

and mobile genetic elements in resistomes of modern dental plaque has indicated the

existence of a co-selection process in the oral microbiome (Kang et al. 2021). Given this

evidence, it is worthwhile to explore the role of heavy metals in the evolution of ARGs in

Industrial Age oral microbiomes that were heavily exposed to metal pollution and untouched

by modern prescription antibiotics.

Research on ARGs in ancient human microbiomes is still scarce due to the limited number of

published ancient ARGs identified in dental calculus. Additionally, there are no established

guidelines for their discovery, as the impact of ancient DNA characteristics on ARG

identification has not been comprehensively studied. Therefore, further research is needed to

address these questions and fill other gaps in our understanding of resistance genes, such as

the natural prevalence of resistance genes in oral microbiomes. This study explores how

environmental contamination and damage resulting from diagenesis affects the discovery and

analysis of resistance genes in ancient samples. Moreover, we analyze dental calculus

samples from different historic periods to investigate which resistance genes in the oral

microbiome can be correlated with historical human transitions such as industrialisation and

modern medical innovations.

Materials and Methods

Quantifying the effects of diagenetic damage

To understand the impact of diagenetic damage on ARG identification, we first carried out a

threshold experiment using simulated ancient DNA. First, we downloaded 182 public human

biofilm samples through the European Nucleotide Archive (ENA) from modern oral

microbiomes that were healthy, caries-active, or periodontally diseased, collected from the

University Hospital in Heidelberg, Germany (ENA project title: PRJNA817430) (Anderson et

al. 2023). To simulate 'damaged' versions of these 182 samples, the fastq files were

converted into fasta format and then the 'deamSim' function from gargammel v1.1.4 (Renaud

et al. 2017) was applied with the following default parameters -damage 0.03, 0.4, 0.01, 0.7 to

deaminate nucleotide bases. Deaminated reads were then recombined with their initial per

base quality scores to recreate fastq paired-end read files.

Having a batch of modern DNA and a batch of 'damaged' DNA to simulate aDNA enabled us

to calculate true positive (TP), false positive (FP), true negative (TN), and false negative (FN)

in our ARGs data. To do this, we implemented Nextflow v23.04.1 to run the MEGARes and

AMR++ pipeline v3 (Bonin et al. 2023) first on the modern samples using three sequence

identity percentage parameters: low (25%), medium (50%), and high (75%), and then on the

simulated aDNA with the same thresholds. We used the modern results at a 75% parameter as

the ground truth.

Following AMR++, an analysis of variance (ANOVA) test was performed in R to measure

significant differences of resistance gene counts between ancient and modern threshold

groups. Then, the type of individual resistance genes between groups (based on unique

MEGARes accession numbers) was analysed by counting the number of times each

MEGARes accession code (ARGs or MRGs) was called in each group. This data was used to

count the number of TPs, FPs, TNs, and FNs based on comparison to the modern 75%

sequence identity threshold results. To take a closer look at the small differences between the

types of ARGs being produced between these groups, resistance gene group types (as

opposed to just the single gene identified through a MEGARes accession) were assembled,

counted across the threshold groups, and an ANOVA test was performed to show statistical

differences in gene type across groups.

Contamination experiment

Next, we sought to determine the degree to which microbial contamination from the
environment influenced the recovery of ARGs and MRGs in historical samples, specifically
the level of false positives introduced due to modern contamination of samples. For this
experiment, we used metagenomic data from 25 post-industrial (ca. 1850 – 1901) dental
calculus samples (50 paired-end fastq files) from varying sites across the UK (Table **S1**). Also
included in this study are 10 modern (21st Century) oral metagenomic samples from the
University of Tennessee, USA, (20 paired-end fastq files) which are involved in a separate
experiment outlined below. Detailed information on these metagenomic datasets is reported
in Standeven et al. 2024.

All metagenomic datasets went through a series of pre-processing steps including quality
control, decontamination, and damage authentication, which can be found in more detail in
the data publication by Standeven et al. 2024. To quality check samples, raw fastq read data
were analysed using FastQC v0.11.9 with default parameters (Babraham Bioinformatics
2019). Then, Fastp v0.23.2 (Chen *et al*. 2018) was used with default parameters to filter and
trim poor-quality reads, cut adapters, repair mismatched base pairs, and ensure overall
quality. BBduk (Joint Genome Institute 2023) was utilised with default parameters to
decontaminate samples by filtering out homologous sequences found in paired environmental
and laboratory blanks. To authenticate ancient sequences, we retained human reads from
outputs produced by Centrifuge v1.0.3 (Kim et al. 2016) and used seqtk 'subseq' to convert
them into fastq files which were then mapped to the human genome (hg38) (NCBI 2013)
using BWA mem v0.7.17 (Li and Durbin 2009). SAMtools v1.12 (Danecek et al. 2021) (-
view -rmdup -flagstat -sort -index) was used to format the alignments and sort into BAM files
which were run through mapDamage2 v2.2.2 with default parameters (Jónsson et al. 2013).

The AMR++ pipeline was applied to these 25 post-industrial samples before and after
decontamination, producing two separate resistance gene results. Firstly, we compared the
number of resistance genes before and after decontamination; a list of counts per resistance
gene in each group was also created for comparison. Then, as described above, an ANOVA
test was performed in R to show statistical differences in types of resistance genes between
the contaminated and decontaminated group to calculate TPs, FPs, TNs, and FNs, using the
decontaminated results as the ground truth. We also identified which resistant genes are

responsible for significant differences between the two groups, and R was used to list which

MEGARes numbers were unique to both groups.

Post-industrial versus modern analysis

The discovery and mass manufacture of antibiotics changed the way humans

experience AMR today – it is no longer solely a microbial conflict taking place discreetly in

the environment, but also a large-scale issue affecting human health and lifestyle. Our study

also investigated which genes in particular have driven that transformation and continue to

impact the oral microbiome. Ten modern dental calculus samples (**Table S1**) were put

through the AMR++ pipeline and resistance genes were compared with the post-industrial

results. Since we chose a 50% sequence identity threshold for our ancient data, this parameter

was also applied to our modern data to make the results comparable.

Percentages of ARGs and MRGs were calculated in each group and compared to highlight

resistance genes associated with ancient and modern oral microbiomes, or any increases or

declines in genes across the two periods. To calculate ARGs and MRGs coverage in these

groups, the MEGARes drugs database was downloaded, and an in-house Python script was

made to extract reference genes in fasta format. BWA mem v0.7.17 (Li and Durbin 2009) (-v

3 -t 20) was then utilised to align resistance query genes to reference genes, and then

SAMtools v1.12 (Danecek et al. 2021) (sort --threads 20 coverage) was used to generate

coverage tables.

257
258 Results
259
260
The impact of damage on ARG prediction

ANOVA was conducted to compare the effect of DNA damage on predicted counts of

resistance genes (MEGARes accession numbers). No significant differences were found

between the simulated ancient and modern threshold groups, F value = 0.917, p = 0.469. A

table of counts per resistance gene for each ancient and modern threshold group was

produced (**Table S2**), showing that slightly more than half of individual genes (51%) do not

exhibit differences in counts across the specified modern and ancient threshold groups. From

these data, the TP, FP, TN, and FN percentages were calculated (**Figure 1; Table S3**). The

combination of TP and TN percentages in each group is close to the ground truth group

(modern 75%). FN percentages are low (<15%) and restricted to ancient groups, and the FP

proportion is higher than the ground truth (10%–45%) in all groups.

Although there are no statistically significant differences in the individual resistance

prediction counts, differences in TPs, FPs, TNs, and FNs suggest that there may be

differences per resistance group type. Therefore, the resistance gene group types were

counted per threshold group and are displayed in a heatmap (**Figure 2; Table S4**). ANOVA

was performed to see statistical differences between resistance gene group type counts across

the different threshold groups and no significant differences were found, F value = 0.337, p =

0.89. Many group type counts are consistent across thresholds like Tetracycline,

Sulfonamides, Spiropyrimidinetriones, etc. However, upon closer inspection, these findings

show that certain drug-resistant categories are not reliable under particular thresholds as they

are likely to call more FPs. For example, when looking for Copper, Iron, or Pleuromutilin

resistance, **Figure 2** shows that these categories of drug resistance do not exist in the 75%

ancient and modern threshold groups, the latter being our basis for the ground truth in this

analysis.

The impact of contamination

For real ancient data, the `runsnp` process in the pipeline failed for thresholds of 75% and

50%, likely due to data quality issues in the fastq files, such as low-quality reads. The

pipeline ran without SNP errors at a 25% threshold. Since the main focus of this analysis is

on resistance gene categories rather than an in-depth SNP analysis, the SNP step is not

critical. Achieving results as close to the ground truth as possible by using a higher threshold

is preferred. However, the pipeline at 75% spent more CPU time on failed tasks compared to

50%, thus, a 50% threshold was selected as a compromise.

Resistance genes were counted in contaminated and decontaminated groups and compared

(**Table S5**) to determine significant variation in the resistance gene count between the groups.

A total of 1,286 individual genes were counted in the contaminated group and 111 were

counted in the decontamination group, a significant difference in gene counts between the

groups (p-value < 2e-16). The resistance gene counts were then used to calculate TPs, FPs,

TNs, and FNs, using the decontaminated results as the ground truth (**Table S6**). The

decontaminated group contained all the TPs, however, the exact counts for each gene were

only correct in three instances (MEG_3838, MEG_5405, MEG_5776). The contaminated

group contained 198 FPs.

The decontaminated resistance gene list was removed from the original contaminated list for

each sample and what remained was 177 unique MEGARes accession numbers (different to

counts) unique to the contaminated group (**Table S7**). The percentage of different resistance

gene types identified in the contaminated and decontaminated groups is shown in **Fig 3**

(**Table S8**). Although certain MEGARes accession numbers are unique to the contaminated

group, **Fig 3** shows that some resistance gene categories that these genes belong to are not.

For example, the contaminated group has many unique Betalactam and Aminoglycosides

ARGs that do not exist in the decontaminated group, but this is not to say that all Betalactam

and Aminoglycosides ARGs are contaminants because the decontaminated group also have a

few Betalactam and Aminoglycosides ARGs.

On the other hand, there are entire drug resistance categories in the contaminated group that

are completely absent in the decontamination samples and are likely to be an artefact of

modern contamination: Aminocoumarins; Arsenic; Biocides; Cationic antimicrobial peptides;

Copper; Fluoroquinolones; Fosfomycin; Glycopeptide; Iron; Phenicol; *Mycobacterium*

tuberculosis-specific drugs; Oxazolidinone; Pleuromutilin; Rifampin; Tellurium;

Thiopeptides; Tetracyclines; and Tetracenomycin. These findings show that contamination

has a large impact on the retrieval of ARGs and MRGs in ancient samples that have not been

decontaminated.

325
326

Patterns in modern and ancient resistance genes

The percentage of different resistance gene types identified in ancient (post-industrial) and

modern dental calculus samples is shown in **Figure 4**. The ancient dataset has 15 more

samples than the modern dataset, so direct similarity comparisons cannot be made. However,

it is clear that certain resistance gene types are specific to certain periods. The ancient oral

microbiome revealed the following resistance genes: including MLS (Macrolide,

Lincosamide, and Streptogramins), Beta-lactams, Fusidic acid, Multi-metal, Elfamycins,

Mercury, Multi-drug, Sulfonamides, Spiropyrimidinetriones, and Aminocoumarins. Several

resistance genes are restricted to modern samples: Aminoglycosides, Copper,

Fluoroquinolones, Iron, Lipopeptides, Oxazolidinone, Rifampin, and Tetracyclines.

Resistance to MLS (43.9%) are ARGs heavily linked to the ancient oral microbiome and are

less common in modern oral microbiota (22.55%). Instead, resistance to Beta-lactams

(10.78%) and Tetracyclines (48.04%) are the dominant resistance genes in the modern oral

microbiomes within this study. **Figure 5** and **Table S9** show overall good coverage (%) for

ARGs and MRGs in modern and post-industrial groups.

342
343   Discussion
344

This research investigated factors affecting the retrieval of resistance genes from ancient

samples, focusing on damage and contamination. Through simulated data, it was found that

gene counts and types remained statistically consistent between ancient and modern threshold

groups, indicating minimal impact from damage (deamination). However, false positives and

negatives in ancient samples necessitate caution. Contamination, on the other hand,

significantly affected gene retrieval, with a notable decrease in gene counts and type post-

decontamination. Analysis of post-industrial and modern dental calculus revealed a

prominent peak of Tetracycline resistance genes in modern oral microbiomes, absent in

ancient samples, indicating evidence of widespread use of Tetracycline following the post-

industrial period into the modern era and its clear impact on the oral microbiome.

Minimal effect of damage on resistance gene retrieval

It could be predicted that lowering the sequence identity barrier would increase resistance

gene counts between groups since loose cut-offs allow for the detection of more DNA

sequences. The ANOVA test found no significant difference in resistance gene count between

simulated ancient and modern threshold groups. This finding is informative since it indicates

that the percentage of FPs is not very high when lowering the identity thresholds.

Nonetheless, this is based solely on overall gene count, and it does not into account the

differences in resistance gene types between the groups which could still lead to misleading

results. Instead of counting all of the genes simultaneously, we compared how many times

each gene was counted in each group and compared them. There were no significant

differences in gene type since most gene counts were consistent across each group. This is an

important outcome because it shows that the pipeline detects similar types of genes in ancient

and modern samples, regardless of damage or threshold.

No significant overall differences in gene count and variability between the ancient and

modern threshold groups may come as a surprise as damaged DNA may have the potential to

read differently to modern DNA and produce varying results. The reason for this might be

because, unlike attempting to align longer and more fragmented core bacterial genomes, the

smaller size of ancient variable genes with fewer nucleotides, such as resistance mechanisms

with fewer nucleotides, may be less challenging to identify. Additionally, as a result of the

AMR crisis, it can be argued that there is a strong research focus on ARGs and MRGs, with

extensive sequencing efforts contributing to their detailed characterisation. This is a

promising outcome, as it indicates that researchers can generally confidently conduct

resistance gene analysis without concern that damage will distort the retrieval of resistance

genes from metagenomic sequences. Nonetheless, it is critical to interpret these data with

caution. We added artificial damage to modern bacterial DNA using defined parameters, but

in fact, aDNA can experience several forms of degradation (Uysal Yoca et al. 2021), all of

which could impact results differently. For example, aDNA that has been *in situ* for 5000

years and has undergone considerable deamination (the transformation of cytosine to uracil

and interpreted as thymine) may have a very different structure to DNA that has been buried

for a century and exposed to harsh acidic conditions. This merits further investigation.

Despite that statistical testing showed threshold and damage do cause statistically measurably

impacts, a closer analysis of the TP, FP, TN, and FN count reveals some discrepancies

between groups, indicating that some refinement of the method is needed. There were no FNs

in modern groups at any threshold, whereas there were some FNs (<10%) in the ancient

groups, the highest being in the ancient 75% group. This is expected because tightening the

threshold allows for fewer genes to be identified, which means that some genes may be

overlooked. Given that these genes are also 'ancient', it is expected that sequences with some

damage in key portions of ARG sequences may drop the similarity to modern reference genes

below the 75% threshold. In comparison to the FN proportion, the FP rate is high across all

groups (between 10% and 45%), increases at lower thresholds, and is the highest among the

ancient groups. In general, the relaxing of criteria allows for the identification of more genes,

and unavoidably, some of those genes will not be genuine. Moreover, the high FP rate among

ancient samples may be linked to the program's ability to process and align modified

nucleotides, causing false results, especially at lower thresholds (50% and 25%).

In ancient samples, low thresholds cause more FPs, and high thresholds cause more FNs.

However, the FN proportion is always lower than the FP proportion in ancient samples across

all thresholds. There is a large element of debate in deciding which thresholds are more

appropriate to use regarding the proportion of FPs and FNs and it is down to context to

conclude. Some studies have recommended using stricter sequence identity cutoffs between

80%-95% for detecting modern resistomes (Bengtsson-Palme et al. 2017; Liao et al. 2022).

Other literature argues that using strict cut-offs is reliable in avoiding false positives, but it

can lead to a high false negative rate. Since many ARGs do not match closely with known

databases, strict similarity thresholds can cause many ARGs to be overlooked (Arango-

Argoty et al. 2018). This is an important consideration in the context of aDNA. While

stringent criteria are necessary for accurately identifying very short-read DNA to minimise

false positives, it is essential to recognize that a more rigorous approach may inadvertently

overlook lesser-known or extinct ARGs from ancient metagenomic sequences. For example,

previous research on ancient ARGs has argued that while lenient cut-offs result in FPs to a

certain extent, the frequency of FNs produced by tighter thresholds is more detrimental

(Rascovan et al. 2016). Although there are medical situations where FNs have grave

consequences, our results suggest that in archaeological contexts, the larger proportion of

present untrustworthy resistance genes is more misleading than the fewer resistance genes

that were missed.

The use of low thresholds on aDNA is sometimes unavoidable. This study found that SNP

analysis was unsuccessful at higher thresholds (50% and 75%) in cases of low read quality in

our archaeological aDNA dataset. Since our focus is on resistance categories rather than

specific genetic variations, poor SNP calling did not impact the overall results of the study.

However, researchers aiming to understand the development, evolution and spread of ancient

resistance genes using gene variation may be affected by these threshold settings and might

need to use lower thresholds, which increases the risk of false positives. Studies focusing on

more effective aDNA extraction methods as well as improved sequencing coverage would

help overcome this challenge.

Future research should also consider investigating additional thresholds beyond sequence

identity, such as target coverage cut-offs, which focus on the proportion of the reference gene

that is aligned. For instance, a study by Cooper et al. (2024) demonstrated that antimicrobial

resistance genes (ARGs) can still be detected even when the target organism is present in

lower amounts and the coverage ranges between 40-60%. However, detecting ARGs at these

lower coverage levels may increase the risk of identifying similar but incorrect ARGs

(Cooper et al. 2024). Given this finding, it is important to evaluate how these thresholds

influence the detection of resistance genes in ancient metagenomic sequences.

We investigated if any specific drug categories were driving these minor discrepancies. It was

revealed that Copper, Iron, and Pleuromutilin resistance are unreliable genes in our ancient

groups because they are absent in the 75% ancient and modern threshold groups. Moreover,

some MEG accessions of the same resistance type have different TP and FP rates. For

example, one Aminocoumarin ARG is trustworthy (MEG_5380), while the rest are not

(MEG_3247, MEG_3248, MEG_3249, MEG_5364, MEG_5380, MEG_5389, MEG_5396,

MEG_3247, MEG_3248, and MEG_3249). These small differences are important because

although the statistics suggest that using certain thresholds on ancient data are appropriate,

stricter parameters are recommended when screening for certain drug-resistant categories.

Significant effect of contamination on resistance gene retrieval

We observed that the quantity and types of resistance genes in dental calculus before and after

decontamination are significantly different. This result was attributed to the vast number and

type of genes unique to the contaminated group that did not exist in the decontaminated

group. Seven different Tetracycline ARGs were removed from the dental calculus batch after

filtering due to being false positives. This demonstrates the level of modern contamination

that, if unfiltered, may be found in ancient samples and result in biased conclusions. Although

not surprising given how vulnerable aDNA is to environmental contamination, this is a major

finding because previous publications have expressed significant concerns around research

that does not properly decontaminate aDNA (Eisenhofer and Weyrich 2018). Studies that do

not use environmental (bone, tooth root, dirt, etc.) and laboratory (extraction, library) blanks

to detect and remove putative contaminants may explain why they detect modern ARGs such

as those conferring to Vancomycin and Tetracycline in dental calculus (Santiago-Rodriguez et

al. 2018; Ottoni et al. 2021). Furthermore, Warinner *et al.* 2014 have shown the absence of

Tetracycline ARGs in bacteria from ancient dental calculus, which makes sense considering

the lack of selective pressure from this antibiotic in the past.

On the other hand, it is not completely implausible to suppose that ancient Tetracycline ARGs

could be identified in historical remains. Ancient populations are believed to have created

fermentation mixtures containing bacteria that produce Tetracycline antibiotics (Nelson et al.

2010), and where there are Tetracyclines, there are Tetracycline ARGs. Furthermore, the

presence of certain drug resistance categories in contaminated samples but not in

decontaminated samples does not necessarily imply their absence in ancient resistomes. This

study identified a limited number of ARGs from a restricted sample set. Therefore, additional

research involving a larger number of samples is required to conclusively determine whether

entire categories of drug resistance, such as tetracycline resistance, are truly contaminants.

Instead of filtering out possible ARG contaminants, studies (Ottoni et al. 2021) have used

systems like SourceTracker (Knights et al. 2011) to assess the risk of contamination by

determining the proportion and origin of microorganisms in a sample; however, to properly

reduce the impact of contamination, it should not be used alone and without decontaminating

using both an environment and a laboratory blank. Overall, we argue that based on our

findings, contamination (unlike damage) has a large effect on the type and amount of

resistance genes being retrieved; thus, adequate decontamination protocols are required.

The rise and emergence of specific resistance genes in oral microbiomes

Following decontamination, we were able to reliably analyse resistance genes in our post-

industrial and modern oral microbiomes. Common ARGs found in people's mouths today are

Tetracycline resistance genes (Seville et al. 2009; Rôças and Siqueira 2013; Arredondo et al.

2021; Brooks et al. 2022; Vázquez-Ramos et al. 2022; Morales-Dorantes et al. 2023), and we

can expect to see less of these ARGs in ancient samples before the 1930s because there was

less selective pressure from antibiotics before their discovery and human manufacture. Our

post-industrial versus modern dental calculus analysis has added evidence to this notion, as

48.04% of ARGs in modern samples conferred Tetracycline while none were found in ancient

samples. There was also a rise in Beta-lactam ARG proportions from 4.88% in ancient

samples to 10.78% in modern ones. This shows that certain genes, such as Tetracycline

ARGs, are modern occurrences that have increased alarmingly inside the oral microbiome.

Therefore, this information could be crucial for determining how these resistance genes

spread, whether the oral microbiome is a site of transmission, or if they are spreading in other

areas and the oral microbiome, like other biofilms, (Guo et al. 2018) serves as a source for

their dissemination.

Worthy of discussion in the post-industrial vs modern analysis is how the presence of certain

ARGs and MRGs came to be in ancient samples. Many antibiotics are ancient and produced

naturally by bacteria, fungi, or metals (Tahlan and Jensen 2013; Wolański et al. 2016; Li and

Seiple 2017; McDonald et al. 2017; Yarlagadda et al. 2020; Long et al. 2021; Mori and Abe

2024; Turner 2024), explaining their presence before the mass production of antibiotics

following the 1920s. Other antibiotics, such as Sulfonamides and Spiropyrimidinetriones

ARGs, are synthetic compounds and their existence in the environment today is a result of

modern human pharmaceutical use, making their presence in our post-industrial dental

calculus confusing.

To explore why certain ARGs are found in post-industrial samples, we can re-examine their

occurrence in the previously mentioned contaminated dental calculus. Sulfonamide ARGs

were counted 13 times in the contaminated group, and three times in the decontaminated

group, and Spiropyrimidinetriones ARGs were counted 16 times in the contaminated group

and 10 times in the decontaminated group. Several of these ARGs are FPs, so they are likely

contaminants, but we can offer explanations for the TPs and the reasoning for their presence

in decontaminated post-industrial samples. Some natural antibiotics may have similar

structures to Sulfonamides or Spiropyrimidinetriones, and ancient bacteria may have

therefore unintentionally developed resistance mechanisms against both in the past, which is

why we may see sequences similar to synthetic compounds in ancient samples. For example,

Aminocoumarins (Alt et al. 2011; Rahimi et al. 2016) and Spiropyrimidinetriones (Shi et al.

2019; Govender et al. 2022), although in slightly different ways, both target the DNA gyrase

enzyme in bacteria. Another argument is that the cutoff was too relaxed, and the 50%

sequence identity threshold resulted in FP Spiropyrimidinetriones and Sulfonamide ARGs. In

the ancient and modern threshold analysis using simulated data, the FP rate for these specific

ARGs increased as the threshold was loosened. Therefore, these synthetic ARGs may surface

in ancient ARG analysis employing medium-low sequence identity thresholds. It is also

possible that the decontamination protocol did not effectively remove these specific ARGs.

However, this scenario is unlikely, as any sequences detected in environmental and

laboratory controls would have been excluded from the calculus samples.

The large proportion of MLS ARGs is reasonable in ancient samples because these naturally

manufactured antibiotics are produced by the 380 million-year-old microorganism

Streptomyces (McDonald et al. 2017) a soil-borne bacteria isolated from a range of natural

environments (Laskaris et al. 2010; Augustine et al. 2012; Encheva-Malinova et al. 2014;
Zhao et al. 2016; Arocha-Garza et al. 2017; Sarmiento-Vizcaíno et al. 2018; Núñez-Montero
et al. 2019; Nicault et al. 2020). If there are microorganisms that want to compete with
Streptomyces in the natural world, *Streptomyces* may increase the antibiotic load in that
environment, which would explain the occurrence of macrolide ARGs in ancient ecosystems.
MSL ARGs have been previously observed in ancient dental calculus (Warinner et al. 2014;
Rascovan et al. 2016) and their observation in other ancient samples may provide insight into
how ancient lifestyles and diet influenced oral resistomes.
546
Since handwashing for hygiene was still in its infancy in the mid-1800s (Poczai and
Karvalics 2022), post-industrial populations may have faced greater exposure to
environmental bacteria, such as antibiotic-producing *Streptomyces*, when preparing food for
example, potentially leading to a different form of selective pressure in the mouth compared
to today. People in the 21st century consume a Western diet containing pre-packaged,
processed, high-sugar, high-fat, and fried meals (Clemente-Suárez et al. 2023) and it is
widely accepted that sugary diets lead to oral disease (Echeverria et al. 2022; Guo et al. 2022;
Echeverria et al. 2023; Spatafora et al. 2024), prompting the use of antibiotics that explain the
high levels of resistance genes observed in our modern calculus. Additionally, antibiotics
used to treat other diseases directly or indirectly linked to poor diets, along with antibiotics
present in certain foods like meat, may also contribute to the resistance genes observed in
modern calculus. It is clear that variations in lifestyle factors, such as diet and hygiene
practices, have contributed to the prevalence of resistance to specific antibiotics in different
periods (e.g., MLS in post-industrial samples and Tetracyclines today). This highlights the
significant shifts in bacterial mechanisms within the oral microbiome over the past two
centuries.
563
Moreover, learning how ancient diets impacted the oral resistome can help us gradually
discover what types of foods and lifestyles may contribute to the emergence of AMR, which
can assist in fostering healthier oral microbiomes today. For example, dental plaque is known
to be a reservoir for AMR distribution (Anderson et al. 2023), so it can be argued that
carbohydrate-rich eating habits and poor oral hygiene that help form plaque should be
addressed to combat AMR. In this light, we should consider the oral microbiota a hotbed for
ARGs.
571

Finally, we examine ancient MRGs in our post-industrial and modern dental calculus

analyses. As humans became more industrialised and more metals contaminated the

environment, including people's noses, mouths, and lungs, we would expect more resistance

to metals to emerge. Mercury and multi-metal ARGs were found in both ancient and modern

samples. Before the Industrial Revolution, it would be normal for modest amounts of

mercury to exert selective pressure on bacteria. Mercury is a naturally occurring contaminant

that enters the environment through volcanic eruptions (Bagnato et al. 2009) and forest

ecosystems (Zhou et al. 2021). Furthermore, metals have always been a part of human life,

from cave fires to several metal ages, therefore the development of multi-metal resistant

ARGs is not unexpected. Copper and Iron MRGs, absent in post-industrial dental calculus,

make an appearance in modern calculus. This could indicate that modern industrialisation has

contributed to the spread of AMR. Because ARGs and MRGs are co-selective agents,

addressing AMR should consider the impact of pollution. To effectively quantify the

influence of pollution on the spread of AMR, more data is needed to completely determine

which MRGs rise over pre-industrial, industrial, post-industrial, and modern dental calculus.

Conclusion

Our experiment using simulated ancient metagenomic datasets suggests that DNA damage

has little effect on the quantity and type of ARGs recovered from ancient samples. In

contrast, environmental contamination of ancient metagenomic datasets has a pronounced

effect, artificially inflating the number and type of ARGs recovered. It is also clear that higher

thresholds generate more TPs and TNs than FPs and FNs in comparison to medium-low

thresholds. Therefore, ARGs are highly reliable at a ≥75% identity percentage threshold for

ancient dental calculus when decontamination protocols are followed. High identity

percentage thresholds can cause SNP errors in low-quality aDNA. As a result, medium to low

thresholds may have to be used but certain drug-resistant groups are interpreted with caution.

In conclusion, the principal findings of this study recommend appropriate thresholds for

resistance gene analysis, emphasise the importance of effective environmental and laboratory

decontamination processes and highlight the significance of using ancient and modern oral

microbiomes as sources for ARGs and MRG analysis.

Conflicts of interest

The authors have declared no competing interests.

Author contributions and funding information

610 Conceptualization: F.J.S, C.J.M. Supervision: A.T., C.J.M., and C.S. Funding acquisition:
611 C.S., A.T., C.J.M. Investigation: F.J.S, G D-A., and C.J.M. Formal Analysis: F.J.S, G. D-A.,
612 C.J.M and A.T. Data Curation, Visualization and Writing -Original Draft: F.J.S. Writing -
613 Review & Editing: all authors.

Acknowledgements

615 The authors acknowledge the use of the University of Bradford High-Performance
616 Computing Service in the completion of this work.

Funding information

619 F.J. S. is funded by the University of Bradford Faculty of Life Sciences and G.D.-A. is
620 funded by the University of Bradford School of Chemistry and Biosciences. No other authors
621 have external funding to acknowledge.
622

References
Alcock, B. P., Huynh, W., Chalil, R., Smith, K. W., Amogelang, Wlodarski, M. A.,
    Edalatmand, A., Petkau, A., Syed, S. A., Tsang, K. K., Baker, S. J. C., Dave, M.,
    Madeline, Mukiri, K. M., Nasir, J. A., Golbon, B., Imtiaz, H., Jiang, X., Kaur, K.,
    Kwong, M., Liang, Z. C., Niu, K. C., Shan, P., Yang, J. Y. J., Kristen, Hoad, G. R., Jia,
    B., Bhando, T., Lindsey, Maya, French, S., Gordzevich, R., Rachwalski, K., Megan,
    Bordeleau, E., Dooley, D., Griffiths, E., Zubyk, H. L., Brown, E. D., Maguire, F.,
    Robert, Hsiao, W. W. L., Brinkman, F. S. L., Gary and McArthur, A. G. (2023) CARD
    2023: expanded curation, support for machine learning, and resistome prediction at
    the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Research* 51 (D1),
    D690-D699.
Alcock, B. P., Raphenya, A. R., Lau, T. T. Y., Tsang, K. K., Bouchard, M., Edalatmand, A.,
    Huynh, W., Nguyen, A.-L. V., Cheng, A. A., Liu, S., Min, S. Y., Miroshnichenko, A.,
    Tran, H.-K., Werfalli, R. E., Nasir, J. A., Oloni, M., Speicher, D. J., Florescu, A.,
    Singh, B., Faltyn, M., Hernandez-Koutoucheva, A., Sharma, A. N., Bordeleau, E.,
    Pawlowski, A. C., Zubyk, H. L., Dooley, D., Griffiths, E., Maguire, F., Winsor, G. L.,
    Beiko, R. G., Brinkman, F. S. L., Hsiao, W. W. L., Domselaar, G. V. and McArthur, A.
    G. (2019) CARD 2020: antibiotic resistome surveillance with the comprehensive
    antibiotic resistance database. *Nucleic Acids Research*.
Alt, S., Mitchenall, L. A., Maxwell, A. and Heide, L. (2011) Inhibition of DNA gyrase and
    DNA topoisomerase IV of Staphylococcus aureus and Escherichia coli by

aminocoumarin antibiotics. *Journal of antimicrobial chemotherapy* 66 (9), 2061-2069.

Anderson, A. C., Von Ohle, C., Frese, C., Boutin, S., Bridson, C., Schoilew, K., Peikert, S. A., Hellwig, E., Pelz, K., Wittmer, A., Wolff, D. and Al-Ahmad, A. (2023) The oral microbiota is a reservoir for antimicrobial resistance: resistome and phenotypic resistance characteristics of oral biofilm in health, caries, and periodontitis. *Annals of Clinical Microbiology and Antimicrobials* 22 (1).

Arango-Argoty, G., Garner, E., Pruden, A., Heath, L. S., Vikesland, P. and Zhang, L. (2018) DeepARG: a deep learning approach for predicting antibiotic resistance genes from metagenomic data. *Microbiome* 6 (1), 23-23.

Arocha-Garza, H. F., Canales-Del Castillo, R., Eguiarte, L. E., Souza, V. and De La Torre-Zavala, S. (2017) High diversity and suggested endemicity of culturable Actinobacteria in an extremely oligotrophic desert oasis. *PeerJ* 5, e3247.

Arredondo, A., Blanc, V., Mor, C., Nart, J. and León, R. (2021) Tetracycline and multidrug resistance in the oral microbiota: differences between healthy subjects and patients with periodontitis in Spain. *Journal of Oral Microbiology* 13 (1), 1847431.

Augustine, N., Peter A, W., Kerkar, S. and Thomas, S. (2012) Arctic Actinomycetes as Potential Inhibitors of Vibrio cholerae Biofilm. *Current Microbiology* 64 (4), 338-342.

Bagnato, E., Parello, F., Valenza, M. and Caliro, S. (2009) Mercury content and speciation in the Phlegrean Fields volcanic complex: Evidence from hydrothermal system and fumaroles. *Journal of volcanology and geothermal research* 187 (3), 250-260.

Baron, S. A., Diene, S. M. and Rolain, J.-M. (2018) Human microbiomes and antibiotic resistance. *Human Microbiome Journal* 10, 43-52.

Bengtsson-Palme, J., Larsson, D. G. J. and Kristiansson, E. (2017) Using metagenomics to investigate human and environmental resistomes. *Journal of Antimicrobial Chemotherapy* 72 (10), 2690-2703.

Bhullar, K., Waglechner, N., Pawlowski, A., Koteva, K., Banks, E. D., Johnston, M. D., Barton, H. A. and Wright, G. D. (2012) Antibiotic resistance is prevalent in an isolated cave microbiome. *PloS one* 7 (4), e34953.

Bombaywala, S., Mandpe, A., Paliya, S. and Kumar, S. (2021) Antibiotic resistance in the environment: a critical insight on its occurrence, fate, and eco-toxicity. *Environmental Science and Pollution Research* 28 (20), 24889-24916.

Bonin, N., Doster, E., Worley, H., Pinnell, L. J., Bravo, J. E., Ferm, P., Marini, S., Prosperi, M., Noyes, N., Morley, P. S. and Boucher, C. (2023) MEGARes and AMR++, v3.0: an updated comprehensive database of antimicrobial resistance determinants and an improved software pipeline for classification using high-throughput sequencing. *Nucleic Acids Research* 51 (D1), D744-D752.

Bortolaia, V., Kaas, R. S., Ruppe, E., Roberts, M. C., Schwarz, S., Cattoir, V., Philippon, A., Allesoe, R. L., Rebelo, A. R., Florensa, A. F., Fagelhauer, L., Chakraborty, T., Neumann, B., Werner, G., Bender, J. K., Stingl, K., Nguyen, M., Coppens, J., Xavier, B. B., Malhotra-Kumar, S., Westh, H., Pinholt, M., Anjum, M. F., Duggett, N. A., Kempf, I., Nykäsenoja, S., Olkkola, S., Wieczorek, K., Amaro, A., Clemente, L., Mossong, J., Losch, S., Ragimbeau, C., Lund, O. and Aarestrup, F. M. (2020) ResFinder 4.0 for predictions of phenotypes from genotypes. *Journal of Antimicrobial Chemotherapy* 75 (12), 3491-3500.

Brooks, L., Narvekar, U., McDonald, A. and Mullany, P. (2022) Prevalence of antibiotic resistance genes in the oral cavity and mobile genetic elements that disseminate antimicrobial resistance: A systematic review. *Molecular Oral Microbiology* 37 (4), 133-153.

Clemente-Suárez, V. J., Beltrán-Velasco, A. I., Redondo-Flórez, L., Martín-Rodríguez, A. and
Tornero-Aguilera, J. F. (2023) Global Impacts of Western Diet and Its Effects on
Metabolism and Health: A Narrative Review. *Nutrients* 15 (12), 2749.
Cooper, A. L., Low, A., Wong, A., Tamber, S., Blais, B. W. and Carrillo, C. D. (2024)
Modeling the limits of detection for antimicrobial resistance genes in agri-food
samples: a comparative analysis of bioinformatics tools. *BMC Microbiology* 24 (1),
31-31).
Dabney, J., Meyer, M. and Paabo, S. (2013) Ancient DNA Damage. *Cold Spring Harbor*
Perspectives in Biology 5 (7), a012567-a012567.
D'Costa, V. M., King, C. E., Golding, G. B., Poinar, H. N., Wright, G. D., Kalan, L., Morar,
M., Sung, W. W. L., Schwarz, C., Froese, D., Zazula, G., Calmels, F. and Debruyne,
R. (2011) Antibiotic resistance is ancient. *Nature (London)* 477 (7365), 457-461.
Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A.,
Keane, T., McCarthy, S. A., Davies, R. M. and Li, H. (2021) Twelve years of
SAMtools and BCFtools. *GigaScience* 10 (2).
Echeverria, M. S., Schuch, H. S., Cenci, M. S., Motta, J. V. D. S., Bertoldi, A. D., Britto
Correa, M., Huysmans, M.-C. D. N. J. M. and Demarco, F. F. (2023) Early Sugar
Introduction Associated with Early Childhood Caries Occurrence. *Caries Research* 57
(2), 152-158.
Echeverria, M. S., Schuch, H. S., Cenci, M. S., Motta, J. V. S., Bertoldi, A. D., Hallal, P. C.
and Demarco, F. F. (2022) Trajectories of Sugar Consumption and Dental Caries in
Early Childhood. *Journal of dental research* 101 (6), 724-730.
Eisenhofer, R. and Weyrich, L. (2018) Proper Authentication of Ancient DNA Is Still
Essential. *Genes* 9 (3), 122.
Encheva-Malinova, M., Stoyanova, M., Avramova, H., Pavlova, Y., Gocheva, B., Ivanova, I.
and Moncheva, P. (2014) Antibacterial potential of streptomycete strains from
Antarctic soils. *Biotechnology & Biotechnological Equipment* 28 (4), 721-727.
Feldgarden, M., Brover, V., Gonzalez-Escalona, N., Frye, J. G., Haendiges, J., Haft, D. H.,
Hoffmann, M., Pettengill, J. B., Prasad, A. B., Tillman, G. E., Tyson, G. H. and
Klimke, W. (2021) AMRFinderPlus and the Reference Gene Catalog facilitate
examination of the genomic links among antimicrobial resistance, stress response, and
virulence. *Scientific Reports* 11 (1).
Feldgarden, M., Brover, V., Haft, D. H., Prasad, A. B., Slotta, D. J., Tolstoy, I., Tyson, G. H.,
Zhao, S., Hsu, C.-H., McDermott, P. F., Tadesse, D. A., Morales, C., Simmons, M.,
Tillman, G., Wasilenko, J., Folster, J. P. and Klimke, W. (2019) Using the NCBI
AMRFinder Tool to Determine Antimicrobial Resistance Genotype-Phenotype
Correlations Within a Collection of NARMS Isolates. Cold Spring Harbor Laboratory.
Florensa, A. F., Kaas, R. S., Clausen, P. T. L. C., Aytan-Aktug, D. and Aarestrup, F. M. (2022)
ResFinder – an open online resource for identification of antimicrobial resistance
genes in next-generation sequencing data and prediction of phenotypes from
genotypes. *Microbial Genomics* 8 (1).
Fotakis, A. K., Denham, S. D., Mackie, M., Orbegozo, M. I., Mylopotamitaki, D.,
Gopalakrishnan, S., Sicheritz-Pontén, T., Olsen, J. V., Cappellini, E., Zhang, G.,
Christophersen, A., Gilbert, M. T. P. and Vågene, Å. J. (2020) Multi-omic detection of
Mycobacterium leprae in archaeological human dental calculus. *Philos. Trans. R. Soc.*
Lond. B Biol. Sci. 375 (1812), 20190584.
Gaeta, R. (2021) Ancient DNA and paleogenetics: risks and potentiality. *Pathologica* 113 (2),
141-146.
Govender, P., Müller, R., Singh, K., Reddy, V., Eyermann, C. J., Fienberg, S., Ghorpade, S.
R., Koekemoer, L., Myrick, A., Schnappinger, D., Engelhart, C., Meshanni, J., Byl, J.

A. W., Osheroff, N., Singh, V., Chibale, K. and Basarab, G. S. (2022) Spiropyrimidinetrione DNA Gyrase Inhibitors with Potent and Selective Antituberculosis Activity. *Journal of Medicinal Chemistry* 65 (9), 6903-6925.

Guo, A., Wide, U., Arvidsson, L., Eiben, G. and Hakeberg, M. (2022) Dietary intake and meal patterns among young adults with high caries activity: a cross-sectional study. *BMC oral health* 22 (1), 190-190.

Guo, X.-p., Yang, Y., Lu, D.-p., Niu, Z.-s., Feng, J.-n., Chen, Y.-r., Tou, F.-y., Garner, E., Xu, J., Liu, M., Hochella, M. F. and Pacific Northwest National Lab, R. W. A. (2018) Biofilms as a sink for antibiotic resistance genes (ARGs) in the Yangtze Estuary. *Water research (Oxford)* 129 (C), 277-286.

Gupta, S. K., Padmanabhan, B. R., Diene, S. M., Lopez-Rojas, R., Kempf, M., Landraud, L. and Rolain, J.-M. (2014) ARG-ANNOT, a New Bioinformatic Tool To Discover Antibiotic Resistance Genes in Bacterial Genomes. *Antimicrobial Agents and Chemotherapy* 58 (1), 212-220.

Hendy, J., Warinner, C., Bouwman, A., Collins, M. J., Fiddyment, S., Fischer, R., Hagan, R., Hofman, C. A., Holst, M., Chaves, E., Klaus, L., Larson, G., Mackie, M., McGrath, K., Mundorff, A. Z., Radini, A., Rao, H., Trachsel, C., Velsko, I. M. and Speller, C. F. (2018) Proteomic evidence of dietary sources in ancient dental calculus. *Proceedings of the Royal Society B: Biological Sciences* 285 (1883), 20180977.

Hutchings, M. I., Truman, A. W. and Wilkinson, B. (2019) Antibiotics: past, present and future. *Current Opinion in Microbiology* 51, 72-80.

Ji, B., Pi, W., Zhang, X. and Peng, S. (2022) HyperVR–A hybrid prediction framework for virulence factors and antibiotic resistance genes in microbial data. Cold Spring Harbor Laboratory.

Kang, Y., Sun, B., Chen, Y., Lou, Y. and Li, Z. (2021) Dental Plaque Microbial Resistomes of Periodontal Health and Disease and Their Changes after Scaling and Root Planing Therapy. *mSphere* 6 (4), 1;e0016221;-20.

Kashuba, E., Dmitriev, A. A., Kamal, S. M., Melefors, O., Griva, G., Römling, U., Ernberg, I., Kashuba, V. and Brouchkov, A. (2017) Ancient permafrost staphylococci carry antibiotic resistance genes. *Microbial Ecology in Health and Disease* 28 (1), 1345574.

Kim, D., Song, L., Breitwieser, F. P. and Salzberg, S. L. (2016) Centrifuge: rapid and sensitive classification of metagenomic sequences. *Genome Research* 26 (12), 1721-1729.

Kistler, L., Ware, R., Smith, O., Collins, M. and Allaby, R. G. (2017) A new model for ancient DNA decay based on paleogenomic meta-analysis. *Nucleic acids research* 45 (11), 6310-6320.

Knights, D., Kuczynski, J., Charlson, E. S., Zaneveld, J., Mozer, M. C., Collman, R. G., Bushman, F. D., Knight, R. and Kelley, S. T. (2011) Bayesian community-wide culture-independent microbial source tracking. *Nature Methods* 8 (9), 761-763.

Kraemer, S. A., Ramachandran, A. and Perron, G. G. (2019) Antibiotic Pollution in the Environment: From Microbial Ecology to Public Policy. *Microorganisms* 7 (6), 180.

Lakin, S. M., Kuhnle, A., Alipanahi, B., Noyes, N. R., Dean, C., Muggli, M., Raymond, R., Abdo, Z., Prosperi, M., Belk, K. E., Morley, P. S. and Boucher, C. (2019) Hierarchical Hidden Markov models enable accurate and diverse detection of antimicrobial resistance sequences. *Communications Biology* 2 (1).

Laskaris, P., Tolba, S., Calvo-Bado, L. and Wellington, L. (2010) Coevolution of antibiotic production and counter-resistance in soil bacteria. *Environmental microbiology* 12 (7), 2048-2048.

Li, H. and Durbin, R. (2009) Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* 25 (14), 1754-1760.

Li, Q. and Seiple, I. B. (2017) Modular, Scalable Synthesis of Group A Streptogramin Antibiotics. *Journal of the American Chemical Society* 139 (38), 13304-13307.

Liao, H., Li, H., Duan, C.-S., Zhou, X.-Y., An, X.-L., Zhu, Y.-G. and Su, J.-Q. (2022) Metagenomic and viromic analysis reveal the anthropogenic impacts on the plasmid and phage borne transferable resistome in soil. *Environment international* 170, 107595.

Long, J., Ji, W., Zhang, D., Zhu, Y. and Bi, Y. (2021) Bioactivities and Structure–Activity Relationships of Fusidic Acid Derivatives: A Review. *Frontiers in pharmacology* 12, 759220-759220.

McDonald, B. R., Currie, C. R. and Univ. of Wisconsin, M. W. I. (2017) Lateral Gene Transfer Dynamics in the Ancient Bacterial Genus Streptomyces. *mBio* 8 (3).

Morales-Dorantes, V., Domínguez-Pérez, R. A., Pérez-Serrano, R. M., Solís-Sainz, J. C., García-Solís, P., Espinosa-Cristóbal, L. F., Cabeza-Cabrera, C. V. and Ayala-Herrera, J. L. (2023) The Distribution of Eight Antimicrobial Resistance Genes in Streptococcus oralis, Streptococcus sanguinis, and Streptococcus gordonii Strains Isolated from Dental Plaque as Oral Commensals. *Tropical Medicine and Infectious Disease* 8 (11), 499.

Mori, T. and Abe, I. (2024) Lincosamide Antibiotics: Structure, Activity, and Biosynthesis. *Chembiochem : a European journal of chemical biology* 25 (6), e202300840-n/a.

Mosley, S. (2008) ***The chimney of the world : a history of smoke pollution in Victorian and Edwardian Manchester***. London, Routledge, Taylor & Francis Group.

NCBI (2013) *Genome assembly GRCh38.* https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_000001405.26/ Accessed 31 July.

NCBI (2024) *National Center for Biotechnology Information*.

Nelson, M. L., Dinardo, A., Hochberg, J. and Armelagos, G. J. (2010) Brief communication: Mass spectroscopic characterization of tetracycline in the skeletal remains of an ancient population from Sudanese Nubia 350-550 CE. *American journal of physical anthropology* 143 (1), 151-154.

Neukamm, J., Pfrengle, S., Molak, M., Seitz, A., Francken, M., Eppenberger, P., Avanzi, C., Reiter, E., Urban, C., Welte, B., Stockhammer, P. W., Teßmann, B., Herbig, A., Harvati, K., Nieselt, K., Krause, J. and Schuenemann, V. J. (2020) 2000-year-old pathogen genomes reconstructed from metagenomic analysis of Egyptian mummified individuals. *BMC Biol.* 18 (1), 108.

Nguyen, C. C., Hugie, C. N., Kile, M. L. and Navab-Daneshmand, T. (2019) Association between heavy metals and antibiotic-resistant human pathogens in environmental reservoirs: A review. *Frontiers of environmental science & engineering* 13 (3), 1-17.

Nicault, M., Tidjani, A.-R., Gauthier, A., Dumarcay, S., Gelhaye, E., Bontemps, C. and Leblond, P. (2020) Mining the Biosynthetic Potential for Specialized Metabolism of a Streptomyces Soil Community. *Antibiotics* 9 (5), 271.

Núñez-Montero, K., Lamilla, C., Abanto, M., Maruyama, F., Jorquera, M. A., Santos, A., Martinez-Urtaza, J. and Barrientos, L. (2019) Antarctic Streptomyces fildesensis So13.3 strain as a promising source for antimicrobials discovery. *Scientific Reports* 9 (1).

Organisation, W. H. (2020) *Antibiotic resistance.* https://www.who.int/news-room/fact-sheets/detail/antibiotic-resistance

Ottoni, C., Borić, D., Cheronet, O., Sparacello, V., Dori, I., Coppa, A., Antonović, D., Vujević, D., Price, T. D., Pinhasi, R. and Cristiani, E. (2021) Tracking the transition to agriculture in Southern Europe through ancient DNA analysis of dental calculus. *Proceedings of the National Academy of Sciences* 118 (32), e2102116118.

Pal, C., Bengtsson-Palme, J., Rensing, C., Kristiansson, E., Larsson, D. G. J., Department of
Mathematical Sciences, M. S., Naturvetenskapliga, f., Institutionen för biomedicin, a.
f. i., Faculty of, S., Göteborgs, u., Gothenburg, U., Sahlgrenska, A., Institute of
Biomedicine, D. o. I. M., Sahlgrenska, a. and Institutionen för matematiska
vetenskaper, m. s. (2014) BacMet: antibacterial biocide and metal resistance genes
database. *Nucleic acids research* 42 (Database issue), D737-D743.
Pawlowski, A. C., Wang, W., Koteva, K., Barton, H. A., McArthur, A. G. and Wright, G. D.
(2016) A diverse intrinsic antibiotic resistome from a cave bacterium. *Nature*
Communications 7 (1), 13803.
Perron, G. G., Whyte, L., Turnbaugh, P. J., Goordial, J., Hanage, W. P., Dantas, G. and Desai,
M. M. (2015) Functional Characterization of Bacteria Isolated from Ancient Arctic
Soil Exposes Diverse Resistance Mechanisms to Modern Antibiotics. *PLOS ONE* 10
(3), e0069533.
Poczai, P. and Karvalics, L. Z. (2022) The little-known history of cleanliness and the
forgotten pioneers of handwashing. *Frontiers in Public Health* 10.
Rahimi, H., Najafi, A., Eslami, H., Negahdari, B. and Moghaddam, M. (2016) Identification
of novel bacterial DNA gyrase inhibitors: An in silico study. *Research in*
pharmaceutical sciences 11 (3), 250-258.
Rascovan, N., Telke, A., Raoult, D., Rolain, J. M. and Desnues, C. (2016) Exploring
divergent antibiotic resistance genes in ancient metagenomes and discovery of a novel
beta-lactamase family. *Environmental Microbiology Reports* 8 (5), 886-895.
Relman, D. A. and Lipsitch, M. (2018) Microbiome as a tool and a target in the effort to
address antimicrobial resistance. *Proceedings of the National Academy of Sciences*
115 (51), 12902-12910.
Rôças, I. N. and Siqueira, J. F. (2013) Detection of antibiotic resistance genes in samples
from acute and chronic endodontic infections and after treatment. *Archives of oral*
biology 58 (9), 1123-1128.
Santiago-Rodriguez, T. M., Fornaciari, G., Luciani, S., Toranzos, G. A., Marota, I., Giuffra,
V., Sangwan, N. and Cano, R. J. (2018) Tetracycline-like resistome of ancient human
guts. *Human microbiome journal* 10, 21-26.
Sarmiento-Vizcaíno, A., Espadas, J., Martín, J., Braña, A. F., Reyes, F., García, L. A. and
Blanco, G. (2018) Atmospheric Precipitations, Hailstone and Rainwater, as a Novel
Source of Streptomyces Producing Bioactive Natural Products. *Frontiers in*
Microbiology 9.
Seville, L. A., Patterson, A. J., Scott, K. P., Mullany, P., Quail, M. A., Parkhill, J., Ready, D.,
Wilson, M., Spratt, D. and Roberts, A. P. (2009) Distribution of Tetracycline and
Erythromycin Resistance Genes Among Human Oral and Fecal Metagenomic DNA.
Microbial Drug Resistance 15 (3), 159-166.
Shi, C., Zhang, Y., Wang, T., Lu, W., Zhang, S., Guo, B., Chen, Q., Luo, C., Zhou, X. and
Yang, Y. (2019) Design, Synthesis, and Biological Evaluation of Novel DNA Gyrase-
Inhibiting Spiropyrimidinetriones as Potent Antibiotics for Treatment of Infections
Caused by Multidrug-Resistant Gram-Positive Bacteria. *Journal of medicinal*
chemistry 62 (6), 2950-2973.
Singh, S., Verma, N. and Taneja, N. (2019) The human gut resistome: Current concepts &
future prospects. *Indian journal of medical research (New Delhi, India : 1994)* 150
(4), 345-358.
Spatafora, G., Li, Y., He, X., Cowan, A. and Tanner, A. C. R. (2024) The Evolving
Microbiome of Dental Caries. *Microorganisms* 12 (1), 121.
Standeven, F. J., Dahlquist-Axe, G., Hendy, J., Fiddyment, S., Holst, M., McGrath, K.,
Collins, M., Mundorff, A., Meehan, C. J., Tedder, A. and Speller, C. F. (2024) An

extensive archaeological dental calculus dataset spanning 5000 years for ancient human oral microbiome research. *bioRxiv*, 2024.09.17.613443.

Tahlan, K. and Jensen, S. E. (2013) Origins of the β-lactam rings in natural products. *The Journal of Antibiotics* 66 (7), 401-410.

Turner, R. J. (2024) The good, the bad, and the ugly of metals as antimicrobials. *BioMetals* 37 (3), 545-559.

Uysal Yoca, Ö., Efe, H. and Yüce, Z. (2021) Ancient DNA Research: Ongoing Challenges and Contribution to Medical Sciences. *Journal of Basic and Clinical Health Sciences* 5 (2), 182-189.

Van Goethem, M. W., Pierneef, R., Bezuidt, O. K. I., Van De Peer, Y., Cowan, D. A. and Makhalanyane, T. P. (2018) A reservoir of 'historical' antibiotic resistance genes in remote pristine Antarctic soils. *Microbiome* 6 (1).

von Wintersdorff, C., Penders, J., van Niekerk, J., Mills, N. D., Majumder, S., van Alphen, L., Savelkoul, P. H. M. and Wolffs, P. F. G. (2016) Dissemination of Antimicrobial Resistance in Microbial Ecosystems through Horizontal Gene Transfer. *Frontiers in microbiology* 7, 173-173.

Vázquez-Ramos, V. R., Pérez-Serrano, R. M., García-Solís, P., Solís-Sainz, J. C., Espinosa-Cristóbal, L. F., Castro-Ruíz, J. E. and Domínguez-Pérez, R. A. (2022) Root canal microbiota as an augmented reservoir of antimicrobial resistance genes in type 2 diabetes mellitus patients. *Journal of applied oral science* 30, e20220362-e20220362.

Warinner, C., Rodrigues, J. F. M., Vyas, R., Trachsel, C., Shved, N., Grossmann, J., Radini, A., Hancock, Y., Tito, R. Y., Fiddyment, S., Speller, C., Hendy, J., Charlton, S., Luder, H. U., Salazar-García, D. C., Eppler, E., Seiler, R., Hansen, L. H., Castruita, J. A. S., Barkow-Oesterreicher, S., Teoh, K. Y., Kelstrup, C. D., Olsen, J. V., Nanni, P., Kawai, T., Willerslev, E., Von Mering, C., Lewis, C. M., Collins, M. J., Gilbert, M. T. P., Rühli, F. and Cappellini, E. (2014) Pathogens and host immunity in the ancient human oral cavity. *Nature Genetics* 46 (4), 336-344.

Weyrich, L. S., Duchene, S., Soubrier, J., Arriola, L., Llamas, B., Breen, J., Morris, A. G., Alt, K. W., Caramelli, D., Dresely, V., Farrell, M., Farrer, A. G., Francken, M., Gully, N., Haak, W., Hardy, K., Harvati, K., Held, P., Holmes, E. C., Kaidonis, J., Lalueza-Fox, C., de la Rasilla, M., Rosas, A., Semal, P., Soltysiak, A., Townsend, G., Usai, D., Wahl, J., Huson, D. H., Dobney, K. and Cooper, A. (2017) Neanderthal behaviour, diet, and disease inferred from ancient DNA in dental calculus. *Nature* 544 (7650), 357-361.

Wolański, M., Łebkowski, T., Kois-Ostrowska, A., Zettler, J., Apel, A. K., Jakimowicz, D. and Zakrzewska-Czerwińska, J. (2016) Two transcription factors, CabA and CabR, are independently involved in multilevel regulation of the biosynthetic gene cluster encoding the novel aminocoumarin, cacibiocin. *Applied Microbiology and Biotechnology* 100 (7), 3147-3164.

Yarlagadda, V., Medina, R., Johnson, T. A., Koteva, K. P., Cox, G., Thaker, M. N. and Wright, G. D. (2020) Resistance-Guided Discovery of Elfamycin Antibiotic Producers with Antigonococcal Activity. *ACS infectious diseases* 6 (12), 3163-3173.

Zankari, E., Allesøe, R., Joensen, K. G., Cavaco, L. M., Lund, O. and Aarestrup, F. M. (2017) PointFinder: a novel web tool for WGS-based detection of antimicrobial resistance associated with chromosomal point mutations in bacterial pathogens. *Journal of Antimicrobial Chemotherapy* 72 (10), 2764-2768.

Zhao, F., Qin, Y.-H., Zheng, X., Zhao, H.-W., Chai, D.-Y., Li, W., Pu, M.-X., Zuo, X.-S., Qian, W., Ni, P., Zhang, Y., Mei, H. and He, S.-T. (2016) Biogeography and Adaptive evolution of Streptomyces Strains from saline environments. *Scientific Reports* 6 (1), 32718.

944 Zhou, J., Obrist, D., Dastoor, A., Jiskra, M. and Ryjkov, A. (2021) Vegetation uptake of
945   mercury and impacts on global cycling. *Nature Reviews Earth & Environment* 2 (4),
946   269-284.

947
948
949

Figure legends

Figure 1. (a) False positive (FP), (b) false negative (FN), (c) true positive (TP), and (d) true
negative (TN) count and percentage per modern and ancient threshold group.

Figure 2. The colour gradient key on the heatmap represents the count of reads that are
mapped to a specific gene in a given threshold group, ranging from white for 0 counts to light
purple for counts up to 500. As the count increases, the colour transitions to dark purple at
1000 counts and shifts to dark blue beyond 1500 counts. The gradient shows the increasing
abundance of genes in across groups, with lighter colours corresponding to lower counts and
darker reflecting higher counts.

Figure 3. Percentage of putative resistance genes identified in contaminated and
decontaminated subgroups.

Figure 4. Percentage of putative resistance genes identified in post-industrial and modern
oral resistomes.

Figure 5. Coverage (%) for resistance genes in ancient and modern groups.

Type of drug resistance

■ Ancient ■ Modern