CHAMP1 complex directs heterochromatin assembly and promotes homology-directed DNA repair

Feng Li[1,*], Tianpeng Zhang[3], Aleem Syed[1], Amira Elbakry[1], Noella Holmer[1], Huy Nguyen[1,2], Sirisha Mukkavalli[1], Roger A. Greenberg[3], Alan D. D'Andrea[1,2,4,*]

1 Division of Radiation and Genome Stability, Department of Radiation Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA 02215, USA.

2 Center for DNA Damage and Repair, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA 02215, USA.

3 Department of Cancer Biology, Penn Center for Genome Integrity, Basser Center for BRCA, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

4 Lead contact

*Corresponding authors: feng_li@dfci.harvard.edu, alan_dandrea@dfci.harvard.edu

ABSTRACT

The CHAMP1 complex, a little-known but highly conserved protein complex consisting of CHAMP1, POGZ, and HP1α, is enriched in heterochromatin though its cellular function in these regions of the genome remain unknown. Here we show that the CHAMP complex promotes heterochromatin assembly at multiple chromosomal sites, including centromeres and telomeres, and promotes homology-directed repair (HDR) of DNA double strand breaks (DSBs) in these regions. The CHAMP1 complex is also required for heterochromatin assembly and DSB repair in highly-specialized chromosomal regions, such as the highly-compacted telomeres of ALT (Alternative Lengthening of Telomeres) positive tumor cells. Moreover, the CHAMP1 complex binds and recruits the writer methyltransferase SETDB1 to heterochromatin regions of the genome and is required for efficient DSB repair at these sites. Importantly, peripheral blood lymphocytes from individuals with CHAMP1 syndrome, an inherited neurologic disorder resulting from heterozygous mutations in *CHAMP1*, also exhibit defective heterochromatin clustering and defective repair of local DSBs, suggesting that a defect in DNA repair underlies this syndrome. Taken together, the CHAMP1 complex has a novel role in heterochromatin assembly and the enhancement of HDR in heterochromatin.

INTRODUCTION

DNA double strand breaks (DSBs), resulting from intrinsic DNA replication errors or from external exposure to DNA damaging agents, can occur at any site throughout the genome. Homology-directed repair (HDR), including classical homologous recombination (HR) and break-induced replication (BIR), functions to repair these DSBs. Some regions of chromosomes may require specific mechanisms of DSB repair. For example, heterochromatic regions of chromosomes, such as centromeres and telomeres, are highly-compacted structures and pose a significant barrier to HDR-mediated DSB repair. Accordingly, cells may require distinct mechanisms to mediate HDR in these chromosomal regions[1-4].

Heterochromatin regions of the genome are characterized by increased trimethylation of histone H3 on lysine 9 (H3K9me3), a key modification that interacts with histone reader proteins[5] and is critical for heterochromatin clustering and function[6]. Heterochromatin assembly is regulated by the coordinated activity of writer H3K9 methyltransferases and eraser H3K9 demethylases. These enzymes regulate the level of H3K9me3 in heterochromatin[6,7], and these mechanisms are coupled to the HDR repair process[8-12].

Some heterochromatin structures have a unique nucleosome organization, and these structures may also require specialized mechanisms for heterochromatin assembly and local HDR repair. For example, approximately 10-15% of cancers employ a telomerase-independent mechanism known as the alternative lengthening of telomeres (ALT pathway). This process relies on HDR for template driven telomere lengthening by a break induced replication-like mechanism[13,14]. Moreover, ALT tumor cells have a high level of H3K9me3 and a specialized heterochromatic microenvironment that facilitates DNA synthesis and telomere expansion[15,16]. Whether the CHAMP1 complex plays a specific role in ALT telomere maintenance or ALT telomere DSB repair remains unknown.

A previous report[17] identified a distinct heterochromatin complex- namely, the CHAMP1/POGZ/HP1α complex- with H3K9me3-binding activity. CHAMP1 (CHromosome Alignment Maintaining Phosphoprotein 1), a highly-conserved zinc finger protein, functions as a regulator of chromosome segregation in mitosis and is required for correct alignment of chromosomes on the metaphase plate[18]. The CHAMP1 binding partner POGZ (POGO transposable element with ZNF domain), also a zinc-finger protein, functions in mitotic cell cycle progression and kinetochore assembly[19]. HP1α, a member of the HP1 family of "hub

proteins", interacts with several chromosomal proteins and binds to H3K9me3 through its

chromodomain (CD). Whether this CHAMP1/POGZ/HP1α complex regulates heterochromatin

assembly or mediates HDR in condensed regions of chromosomes remains unknown.

Humans with *de novo* heterozygous mutations in *CHAMP1*, known as CHAMP1

syndrome, present with a neurodevelopmental disorder characterized by intellectual disability,

behavioral symptoms, and distinct dysmorphic features [20-24]. Interestingly, *de novo* heterozygous

mutations in the *POGZ* gene also result in a rare but highly-related neurodevelopment syndrome

called the White Sutton Syndrome (WSS) [25-33]. Whether these inherited mutations result from

defects in heterochromatin assembly or HDR in heterochromatin remains unknown.

Recent studies have elucidated at least one mechanism by which CHAMP1 and POGZ

promote HR [34-36]. CHAMP1 has a unique, high affinity site for the DNA repair HORMA

protein, REV7.   By binding to REV7, the CHAMP1/POGZ complex steals REV7 from the

Shieldin (REV7/SHLD) complex[36]. The Shieldin complex normally blocks DNA end resection,

promotes NHEJ repair, and inhibits HR repair [37-41].   By binding to REV7, the CHAMP1/POGZ

complex increases DSB end resection and promotes HR repair. Whether this mechanism of

enhanced HR repair occurs in heterochromatin, or whether its use is confined to euchromatic

regions, remains unknown.

In the current study we show that the CHAMP1/POGZ/HP1α heterochromatin complex

performs a direct role in heterochromatin clustering and HDR repair. The complex promotes the

deposition of H3K9me3, the clustering of heterochromatin, and the recruitment of HDR proteins

(both HR proteins and BIR proteins) to DSBs at heterochromatic sites. While the complex

subunits promote heterochromatin clustering at multiple sites, including centromeres and

telomeres, it has a distinct and obligate role in maintaining ALT telomeres. Depletion of the

CHAMP1/POGZ/HP1α complex results in reduced H3K9me3 and loss of heterochromatin

clustering. Interestingly, the POGZ subunit also binds and recruits the methyltransferase

SETDB1 to sites of heterochromatin.   Finally, peripheral blood lymphocytes from individuals

with CHAMP1 syndrome also exhibit defective heterochromatin clustering and HR, suggesting

that a defect in DNA repair underlies this syndrome. Taken together, the CHAMP1/POGZ/HP1α

is a novel molecular machine which regulates heterochromatin clustering and contributes to local

HDR repair in heterochromatin.

RESULTS

Structural modelling of the heterochromatic CHAMP1-POGZ-HP1α complex

Quantitative proteomic analyses of histone epigenetic marks and readers identified

CHAMP1-POGZ-HP1α as a heterochromatin complex that binds H3K9me3[17]. Recent studies

have shown that the CHAMP1[34,36] and POGZ subunits[35] promote HR-mediated DSB repair

through interaction with REV7.　Nevertheless, the relationship between the heterochromatin

localization of the complex and HR is unclear.

To investigate the organization of the multi-subunit CHAMP1 complex, we performed *in*

silico structural modeling of binary interactions among the subunits with AlphaFold2-Multimer

(AF2)[42,43]. This allowed us to predict a high-confidence interaction between the CHAMP1 N-

terminal ZnF domain (1-87 aa) and the structured C-terminus of POGZ (1021-1410 aa) (**Fig.**

1a,b). The interaction between CHAMP1 and POGZ is formed primarily by hydrophobic

residues at the interface (**Fig. 1a**). Conversely, our modelling showed that the CHAMP1 C-

terminal ZNF domain forms an extended β-sheet-type interaction with residues (16-22 aa) from

the N-terminus of HP1α (**Extended Data Fig. 1a**). Furthermore, HP1α homodimerizes through

its C-terminal chromodomains (**Extended Data Fig. 1b**), while the HPZ domain of POGZ

interacts with the same C-terminal chromodomain of HP1α from the opposite side of the

dimerization interface (**Extended Data Fig. 1c**).　Since CHAMP1 binds to the same site on the

N-terminus of HP1α as H3K9me3, HP1α homodimerization through its C-terminal

chromodomain may allow a HP1α dimer to bind CHAMP1 and H3K9me3 simultaneously.

Through these AF2 predictions and previously published studies on the interactions

between POGZ and HP1α[19], HP1α and H3K9me3 [5], and HP1α dimerization [44,45], we built a

composite structural model of the CHAMP1 complex containing CHAMP1, POGZ, HP1α and

H3K9me3 (**Fig. 1c**). In this model, the N-terminus of CHAMP1 interacts with POGZ, while the

C-terminus of CHAMP1 directly engages with an HP1α dimer[5]. The CHAMP1-POGZ-HP1α

complex appears to function as a bridge, connecting two H3K9me3. We next validated these

predictions using a series of mutations in CHAMP1 and HP1α (homodimerization mutant) (**Fig.**

1c-f, and Extended Data Fig. 1d). As expected, N-terminal mutants of CHAMP1 (ΔN, ΔNΔC

and QLT/RRR) failed to bind to POGZ (**Fig. 1d, lanes 7,8, and Extended Data Fig. 1d, lane 6**),

and C-terminal mutants of CHAMP1 (ΔNΔC and ΔC) failed to bind to HP1α (**Fig. 1d, lane 8,**

and Fig. 1e, lane 10). Furthermore, the HP1α dimerization mutant I165E[19,44] failed to bind to

CHAMP1, while POGZ could independently bind to HP1α and H3K9me3 (**Fig. 1f, lane 6**).

Interestingly, the GFP-CHAMP1-2A mutant, which failed to bind to REV7, still pulled

down POGZ and HP1α (**Fig. 1e, lane 8**), suggesting that REV7 is not required for the formation

of the CHAMP1/POGZ/HP1α heterochromatin complex. GFP-HP1α successfully pulled down

CHAMP1, POGZ, and H3K9me3, but did not pull down REV7 (**Fig. 1g, lanes 5,6**). Moreover,

GFP-REV7 pulled down CHAMP1 and POGZ but did not pull down HP1α or H3K9me3 (**Fig.**

1h, lanes 5,6). Taken together, our data suggest that the CHAMP1 complex may function in a

REV7-independent pathway while interacting with heterochromatin.

CHAMP1 complex promotes heterochromatin clustering at multiple genomic regions

To investigate the potential function of the CHAMP1 complex in the formation of

heterochromatin clusters, we initially assessed the localization of CHAMP1 and POGZ within

heterochromatin foci by immunofluorescence. Remarkably, we observed a strong colocalization

of CHAMP1 and POGZ with H3K9me3, which we used as a marker of heterochromatin, in

human U2OS and mouse NIH-3T3 cells. Colocalization was further enhanced by ionizing

radiation (IR) induced DSBs (**Fig. 2a,b and Extended Data Fig. 2a,b**). Consistently, the

association between CHAMP1 and HP1α was stronger after cellular exposure to IR (**Extended**

Data Fig. 2c). This interaction was not affected by the loss of REV7 interaction (i.e., the

CHAMP1-2A mutant), supporting the idea that the CHAMP1 heterochromatin complex

functions independently of REV7. Interestingly, radiation also induced the interaction of

H3K9me3 with HP1α, but not with HP1β or HP1γ (**Extended Data Fig. 2d**), suggesting a

specific role of HP1α in the heterochromatin enrichment at DSBs.

Notably, the CRISPR-knockout (KO) of CHAMP1 or POGZ in U2OS cells resulted in a

significant reduction in the formation of H3K9me3 and HP1α foci (**Fig. 2c-f**). However, the

knockout of CHAMP1 or POGZ did not affect the overall expression level or chromatin binding

of H3K9me3 and HP1α (**Extended Data Fig. 2e,f, lane 9-12**). To further determine whether the

CHAMP1 heterochromatin complex promotes H3K9me3 accumulation at centromeres and

telomeres, well-known heterochromatin sites, we next performed immunofluorescence with a

centromere marker (Cent FISH) or a telomere marker (anti-TRF2 antibody)[46]. As expected, we

observed a significant loss of H3K9me3 foci at both centromeres and telomeres following

deletion of the CHAMP1 or POGZ in U2OS cells (**Fig. 2g-j).** Consistently, ChIP-qPCR also

demonstrated a decreased level of H3K9me3 at telomeres in CHAMP1 or POGZ KO U2OS cells

(**Fig. 2k**). Taken together, the CHAMP1 complex plays a role in assembling the H3K9me3

heterochromatin mark at multiple sites in the genome, including sites with damage-induced

DSBs as well as at centromeres and telomeres.

Recent studies on telomeres in ALT tumor cells suggest a distinctive heterochromatin

structure characterized by high levels of H3K9me3, low levels of ATRX, and elevated HDR

processes at telomeres[47-49]. We therefore decided to investigate the possible role of the CHAMP1

complex in the maintenance of these ALT telomere structures and HDR repair. We first

examined the possible localization of the CHAMP1 complex at telomeres in ALT U2OS cells.

Notably, we observed a strong colocalization of CHAMP1 and POGZ with telomere foci,

particularly with large telomere foci, in U2OS cells (**Extended Data Fig.3a,b**). This suggests

that the CHAMP1 complex may play a role in promoting the formation of telomere clusters.

DSBs at telomeres are known to initiate long-range movements and clustering of telomeres, and

these processes are critical for homology-directed telomere synthesis[50]. Accordingly, we

evaluated the response of U2OS cells to engineered DSBs at telomeres, using the TRF1-

FokI[50,51]. TRF1-FokI induced telomeric DSBs (**Extended Data Fig.3c,d**), resulting in telomere

clustering (**Fig. 3a,b**), as measured by an increase in the average size of telomere foci in ALT

cells and a decrease in the total number [50] (**Fig. 3c,d**). Interestingly, this DSB-induced telomere

clustering was significantly diminished in CHAMP1 or POGZ knockout U2OS cells (**Fig. 3c,d**).

Depletion of CHAMP1 or POGZ also led to a decrease in the hallmarks of ALT recombination,

including a reduction in ALT-associated PML bodies (APBs)[14,52] (**Fig. 3e,f**), which are sites of

telomeric clustering and recombination[15,53], and a decrease of non-S phase telomeric DNA

synthesis[16] (**Fig. 3g-i**). Taken together, our results demonstrate that the CHAMP1 complex plays

a critical role in facilitating damage-induced telomere clustering and in maintaining ALT activity

(**Fig. 3j**).

CHAMP1 complex promotes HDR at ALT telomeres

We next used the PICh (<u>P</u>roteomics of <u>I</u>solated <u>Ch</u>romatin Segments) method to identify

the recruitment of telomere-associated proteins following DSBs, in the presence or absence of

the CHAMP1 heterochromatin complex (**Fig. 4a and Extended Data Fig. 4a**). Consistent with

previous observations[51], multiple DNA repair- and telomere maintenance-related biological

processes were significantly upregulated at telomeres following TRF1-FokI induction (**Extended**

Data Fig. 4b). CRISPR knockout of CHAMP1 resulted in a decrease of other CHAMP1

complex proteins (i.e., POGZ and HP1α) at telomeres with DSBs, as well as a decrease of many

proteins associated with HR (e.g., RAD51, BRCA1, and BARD1), as measured by mass

spectrometry (**Fig. 4b**). These results are consistent with the known role of CHAMP1 and POGZ

in HR repair[34-36]. Additionally, CHAMP1 depletion also resulted in a reduction in proteins

involved in break-induced replication (BIR) or break-induced telomere synthesis (BITS) [51,54]

(i.e., POLD3, POLD4, RFC and PCNA) (**Fig. 4b**). Western blot analysis of PICh-purified

telomeres revealed an enrichment of the CHAMP1 complex (i.e., CHAMP1 and HP1α) and

BITS proteins (i.e., POLD3, RAD18 and PCNA) at telomeres in U2OS cells following TRF1-

FokI induction, but a reduction in CHAMP1-depleted U2OS cells (**Fig. 4c, lanes 9-12**).

Immunofluorescence further confirmed the reduction of CHAMP1 complex (i.e., POGZ and

HP1α) and H3K9me3 at telomeres in CHAMP1 knockout U2OS cells (**Fig. 4d-g and Extended**

Data Fig. 4c,d). Consistently, POGZ depletion also resulted in a reduction of proteins associated

with HR and BITS at telomeres with DSBs (**Extended Data Fig. 4e**). Consistent with the local

reduction in HR proteins, knockout of CHAMP1 or POGZ in U2OS cells resulted in a reduction

in HR activity at telomeres, as measured by RAD51 and pRPA2(S33) foci at damaged telomeres

(**Fig. 4h,i and Extended Data Fig. 4f,g**). Taken together, these results support a model in which

the CHAMP1 complex promotes HDR, potentially through the clustering of heterochromatin at

telomeres.

CHAMP1 complex binds and recruits SETDB1 to telomeric heterochromatin clusters

221   To further evaluate the regulation of telomere heterochromatin deposition, we examined

H3K9 methyltransferase levels and activity. Interestingly, SETDB1, but not other H3K9

methyltransferases, was recruited to telomeres after TRF1-FokI induction in U2OS cells,

whereas this recruitment was not observed in CHAMP1- and POGZ-depleted U2OS cells (**Fig.**

5a). Immunofluorescence confirmed the reduction in SETDB1 at telomeres following TRF1-

FokI induction in CHAMP1 or POGZ knockout U2OS cells (**Fig. 5b,c**). Based on previously

published data, indicating a POGZ and SETDB1 interaction[55], we reasoned that the CHAMP1

complex might directly recruit SETDB1 to telomeres. A strong interaction between SETDB1

and the Zn-fingers (type 2-7) of POGZ was predicted (iPTM =0.85) (**Fig. 5d**). Furthermore, an

N-terminal deletion of CHAMP1, causing loss of the interaction with POGZ, also exhibited

reduced interaction with SETDB1 (**Fig. 5e**). These results suggest that the CHAMP1 complex

may recruit SETDB1 to telomeres through POGZ (**Extended Data Fig. 5a**).

Recently, it has been shown that ALT activity relies on a high level of H3K9

trimethylation, facilitated by SETDB1, and on the recruitment of recombination factors [48].

Consistently, knockout of SETDB1 resulted in a decrease of H3K9me3 foci at telomeres

(**Extended Data Fig. 5b,c**). Importantly, SETDB1 depletion decreased CHAMP1 recruitment to

telomeres (**Fig. 5f,g**). Furthermore, SETDB1 depletion in U2OS cells led to reduced damage-

induced telomere clustering and APBs (**Fig. 5h,i**). Collectively, these findings suggest that the

CHAMP1 heterochromatin complex recruits SETDB1, thereby promoting a positive feedback

loop that enhances the formation of telomere heterochromatin clusters (**Fig. 5j**).

CHAMP1 and HP1α are epistatic in the regulation of heterochromatin clustering

Since HP1α is a critical subunit of the CHAMP1 heterochromatin complex, we next

examined its role in the regulation of telomeric HR and clustering. Consistent with previous

findings, HP1α is required for APBs formation[56]. HP1α knockdown resulted in a significant

decrease of damage-induced telomere clustering and APBs formation (**Fig. 6a,b and Extended**

Data Fig. 6a). HP1α knockdown did not result in a further decrease in damage-induced

telomere clustering and APBs in CHAMP1 knockout U2OS cells (**Fig. 6a,b and Extended Data**

Fig. 6a), indicating that CHAMP1 and HP1α are epistatic for APB formation and telomere

clustering. Consistent with the requirement of HP1α dimerization in the function of the

CHAMP1 complex, the wild-type HP1α protein, but not the HP1α-I165E dimerization mutant,

restored telomere clustering and APBs in cells with an HP1α knockdown (**Fig. 6c,d and**

Extended Data Fig. 6b). Furthermore, the C-terminal deletion mutant of CHAMP1, which lacks

HP1α binding, and the N-terminus deletion mutant of CHAMP1, which lacks POGZ binding, did

not restore RAD51 foci at telomeres or telomere clustering after CHAMP1 depletion (**Fig. 6e,f**

and Extended Data Fig. 6c). These findings highlight the critical importance of the multi-

subunit CHAMP1 heterochromatin complex in telomeric HR and clustering.

REV7 binding is not required for heterochromatin clustering or functional HR repair

The interaction of CHAMP1 and REV7 appears to be dispensable for the telomere HR and clustering (**Fig. 6e,f and Extended Data Fig. 6c**). REV7 does not co-immunoprecipitate with the CHAMP1 complex (**Fig. 1e,g,h**). Moreover, the CHAMP1-2A mutant protein, which has a disruption of the CHAMP1 binding site for REV7, can still assemble into the CHAMP1 complex, promote heterochromatin cluster formation, and promote HR activity (ie, RAD51 foci at sties of DSBs) (**Fig. 6e,f and Extended Data Fig. 6c**). Taken together, these results demonstrate that the activity of the REV7-independent CHAMP1 heterochromatin complex is essential for heterochromatic clustering and HR repair in heterochromatin.

Patients with CHAMP1 syndrome display impaired heterochromatin clustering and defects in HR repair

We next examined EBV-immortalized lymphoblast cell lines derived from patients with CHAMP1 syndrome, resulting from an inherited mutation in one allele of the *CHAMP1* gene. The cells were obtained via the Coriell mutant cell line repository (**Fig. 7**). These cell lines express the mutant (truncated) CHAMP1 proteins (**Fig. 7a, and Extended Data Fig. 7**), as predicted by their genomic sequencing. These truncated proteins, lacking the C-terminal HP1α binding domain, may act as dominant negative proteins or lead to haploinsufficiency. Interestingly, these CHAMP1 patient-derived cell lines exhibit a defect in heterochromatin clustering, as indicated by a decrease of H3K9me3 foci (**Fig. 7b,c**) compared to wild-type control cells. The CHAMP1 patient-derived cell lines also have a defect in RAD51 foci assembly following IR (**Fig. 7d,e**), consistent with a deficiency in HR, and exhibit heightened sensitivity to both IR and PARP inhibitors (**Fig. 7f,g**). In summary, lymphocytes from individuals carrying CHAMP1 mutations exhibit a defect in heterochromatin clustering and deficiencies in homologous recombination, providing a possible functional diagnostic assay for the CHAMP1 syndrome.

DISCUSSION

Taken together, our studies show that the CHAMP1 heterochromatin complex, consisting of CHAMP1, POGZ, and HP1α, has a direct role in heterochromatin assembly and Homology-Directed DNA Repair. The complex binds to H3K9me3 in heterochromatin sites, including

centromeres and telomeres, and contributes to the H3K9 trimethylation at these sites. The

complex is also recruited to heterochromatic sites with DSBs generated by ionizing radiation.

Moreover, the CHAMP1 heterochromatin complex plays a role in HDR at these heterochromatic

sites.

Our results support the presence of two distinct CHAMP1/POGZ complexes. One is

REV7-dependent, and the other is REV7-independent. We previously showed that the

CHAMP1/POGZ complex promotes HR by binding to REV7 and limiting the binding of REV7

to proteins in the Shieldin complex (i.e., the REV7, SHLD1, SLD2, SHLD3 complex). The

Shieldin complex is known to block the resection of DSB double strand breaks and thereby limit

the level of HR repair. While this HR mechanism primarily occur in euchromatic regions, it may

also occur to some extent in heterochromatic regions. However, in the current study, we identify

another CHAMP1 complex, the CHAMP1 heterochromatin complex, whose function was

previously unknown. This latter complex is enriched in heterochromatin and excludes the

CHAMP1 binding partner REV7. Here, we show that this complex interacts directly with

HP1α and promotes the H3K9me3 marks in heterochromatin. Moreover, this complex includes

the SETDB1 methyltransferase. The complex is required for the recruitment of HDR proteins to

heterochromatic regions of the chromosome. Loss of the CHAMP1 heterochromatin complex

results in a decrease in repair of TRF1-FokI mediated DSBs at telomeres. Knockdown of

CHAMP1 complex subunits, or mutation of individual subunits preventing complex assembly,

reduced pRPA and RAD51 foci at these sites.

Evaluation of the CHAMP1-2A mutant protein, which fails to bind to REV7[36] is

especially informative. CHAMP1-2A protein does not enable HDR repair by preventing the

binding of REV7 with the Shieldin complex[36]. However, the CHAMP1-2A mutant protein is still

functional for HDR repair through its interaction with the CHAMP1 heterochromatin complex.

As a subunit of the CHAMP1 heterochromatin complex, the CHAMP1-2A mutant protein still

allows heterochromatin clustering, recruitment of HDR proteins to heterochromatin, and repair

of DSBs in heterochromatin (i.e., formation of RAD51 foci).

Our study also demonstrates that the CHAMP1 heterochromatin complex is especially

important to heterochromatin clustering and HDR repair at specific heterochromatic sites. For

instance, the telomeres of ALT tumor cells are known to have high levels of H3K9me3, high

HDR activity, and high levels of replication stress. ALT telomeres appear to have a unique

heterochromatin environment [48,57]. The CHAMP1 heterochromatin complex is required for

assembly and maintenance of ALT telomeres, and knockdown or knockout of the CHAMP1

complex results in decreased DSB-induced telomere cluster formation and reduced HDR activity,

coincident with a loss of telomere-associated histone H3 lysine 9 trimethylation (H3K9me3).

The CHAMP1 heterochromatin complex may play additional roles in heterochromatin structures.

For instance, the complex may be regulated by the ATM kinase[2] or may function in HDR repair

at the periphery of the cell nuclease[9,18]. The complex may play a role in specific signaling

events required for assembling heterochromatin at stressed replication forks[58,59]. Future studies

will be required to assess the role of the CHAMP1 heterochromatin complex at these other

heterochromatic sites.

A reduction in H3K9me3 is known to result from a mislocalization or inactivation of the

SETDB1, a methyltransferase known to trimethylate H3K9[60]. Consistent with these findings,

our AF2 analysis, immunoprecipitation studies, and PICh results revealed an interaction of

SETDB1 with the POGZ subunit of the complex, required for the recruitment of SETDB1 to

sites of heterochromatin. Taken together, the CHAMP1 complex appears to be a critical

determinant of heterochromatin clustering. Genetic ablation of a subunit of the complex, or a

mutation in a subunit which disrupts complex assembly, results in failure to recruit SETDB1 and

in loss of heterochromatin clustering.

Previous studies have suggested that tightly-packed heterochromatin regions have an

overall decrease in HDR activity[3]. Paradoxically, in our studies and in other recent reports[48,61,62],

the increase in H3K9 trimethylation or other modifications in the heterochromatin composition at

telomeres, may enhance HDR activity. Consistent with these findings, the CHAMP1 complex

may promote HDR by functioning as a bridge between two heterochromatin nucleosomes, either

intra- or interchromosomally, through its interaction with H3K9me3 in heterochromatin. This

bridge may facilitate the formation of heterochromatin clusters, thereby creating an environment

that promotes the recruitment and stabilization of HDR proteins.

Finally, individuals with *de novo* heterozygous mutations in *CHAMP1*, known as

CHAMP1 syndrome, present with a neurodevelopmental disorder characterized by intellectual

disability, behavioral symptoms, and distinct dysmorphic features. Interestingly, *de novo*

heterozygous mutations in the *POGZ* gene also result in another rare but highly-related

neurodevelopment syndrome called the White Sutton Syndrome (WSS) [25,27,28,30,31,63]. The

mutant proteins encoded by pathogenic alleles of CHAMP1 have C-terminus truncations, and

perhaps dominant negative activity, and cannot assemble into functional CHAMP1 complexes.

Accordingly, lymphoblast cultures derived from individuals with CHAMP1 syndrome and

expressing these mutant proteins have defects in heterochromatin and homologous

recombination. These functional defects may also underlie the neurodevelopmental defects in

these individuals.   Lymphoblasts from these individuals have phenotypic characteristics similar

to engineered human cells lacking the CHAMP1 complex. Taken together, these cellular

characteristics could serve as a functional diagnostic screening tool for identifying individuals

with CHAMP1 syndrome among the much larger number of individuals with autism spectrum

disorders. In conclusion, we provide evidence that the CHAMP1 heterochromatin complex is

critical for the heterochromatin HDR repair, not only in tumor cells but also in normal tissue

cells. The clinical significance of the CHAMP1 complex will require further studies.

Fig. 1. CHAMP1-POGZ-HP1α form a REV7 independent heterochromatin complex. a.
AlphaFold2-Multimer (AF2)-predicted structural model of POGZ_Cter (peach)-
CHAMP1_NZnF (cyan) complex. The key residues (shown in sticks) contributing to the protein-
protein interactions are highlighted in the inset including CHAMP1 residues (Q56, L61 and T70)
that are mutated to disrupt the interaction between POGZ and CHAMP1. **b**. The AF2-predicted
model of POGZ_Cter-CHAMP1_NZnF is colored based on the confidence in the model
prediction (100-high (blue) and 50-low (red)) and corresponding predicted aligned error plot
(PAE) matrix showing the confidence in the predicted interaction between CHAMP1_NZnF and
POGZ_Cter domains along with interface-predicted template scores (iPTM) for the AF2
prediction. **c.** (Top) Schematic of CHAMP1 wild-type (WT) and mutants. N-ZNFs (N-terminus
C2H2-Zn finger domains), SPE (PxxSPExxK motifs), WK (SPxxWKxxP motifs), FPE (FPExxK
motifs), C-ZNFs (C-terminus C2H2-Zn finger domains), and SBM (REV7 seatbelt-binding
motif). 2A indicates W334A/K335A mutant. (Bottom) Schematic of CHAMP1 complex
connecting two H3K9me3 sites. The N-terminus of CHAMP1 binds to the C-terminus of POGZ.
Additionally, both the C-terminus of CHAMP and the HPZ motif of POGZ can independently
bind to HP1α. CHAMP1 shows a preference for binding to the HP1α dimer, while POGZ is
capable of binding to either the HP1α dimer or monomer. N-terminus of HP1α directly interacts
with H3K9me3. **d**. Western blot showing GFP-immunoprecipitation of GFP-empty vector, GFP-
CHAMP1 WT or N- or C- terminal deletion mutants, and the co-immunoprecipitation of
endogenous POGZ, HP1α and REV7. **e.** Western blot showing GFP-immunoprecipitation of
GFP-empty vector, GFP-CHAMP1 WT and GFP-CHAMP1 mutants in 293T cells, and the co-
immunoprecipitation of endogenous POGZ, HP1α and REV7. **f**. Western blot showing GFP-
immunoprecipitation of GFP-empty vector, GFP-HP1α WT or I165E mutant, and the co-
immunoprecipitation of endogenous POGZ, CHAMP1 and H3K9me3. GAPDH acts as a
negative control. **g.** Western blot showing GFP-immunoprecipitation of GFP-empty vector, GFP-
HP1α in 293T WT and sgCHAMP1 cells, and the co-immunoprecipitation of endogenous POGZ,
CHAMP1, REV7 and H3K9me3. GAPDH acts as negative control for IP. **h.** Western blot
s
h
o
397/399 w
398/400 i

Fig. 2. The CHAMP1 complex is required for the nuclear distribution of H3K9me3. a.
Representative immunofluorescence (IF) confocal images of the colocalized CHAMP1 and
H3K9me3 foci with/without IR (5Gy) treatment in U2OS cells. DAPI was used to stain the
nuclei. Scale bar, 5 μm. **b**. Square analysis of the intensity and colocalization of H3K9me3 and
CHAMP1 foci. **c**. Representative IF images of H3K9me3 foci in WT, sgCHAMP1 and sgPOGZ
U2OS cells. DAPI was used to stain the nuclei. Scale bar, 5 μm. **d**. Quantification of H3K9me3
foci in (c). Error bars indicate SEM. More than 100 cells were counted. ****P<0.0001.
Statistical analysis was performed using two-tailed Student's t test. **e**. Representative IF images
of HP1α foci in WT, sgCHAMP1 and sgPOGZ U2OS cells. DAPI was used to stain the nuclei.
Scale bar, 5 μm. **f**. Quantification of HP1α foci in (e). Error bars indicate SEM. More than 100
cells were counted. ****P<0.0001. Statistical analysis was performed using two-tailed Student's
(410) t test. **g**. Representative IF-FISH confocal images of H3K9me3 colocalization with centromere
(411) (Cent) in WT, sgCHAMP1 and sgPOGZ U2OS cells. DAPI was used to stain the nuclei. Scale
p
r
e
c
i
p

bar, 5 µm. **h**. Quantification of H3K9me3-Centromere colocalizations in (g). Error bars indicate
SEM. More than 100 cells were counted. ****P<0.0001. Statistical analysis was performed
using two-tailed Student's t test. **i**. Representative IF confocal images of H3K9me3-TRF2
colocalization in WT, sgCHAMP1 and sgPOGZ U2OS cells. TRF2 was used as a telomere
maker. DAPI was used to stain the nuclei. Scale bar, 5 µm. **j**. Quantification of H3K9me3
colocalization with TRF2 in (i). Error bars indicate SEM. More than 100 cells were counted.
****P<0.0001. Statistical analysis was performed using two-tailed Student's t test. **k**. ChIP-
qPCR analyses showing relative enrichment levels of H3 (Top) and H3K9me3 (Bottom) at
telomeres. Level of H3K9me3 was normalized to total H3 levels (H3K9me3/H3). Error bars
indicate SD. P-values calculated using Student's T-test (***P<0.001; ns, non-significant).
484
485
Fig. 3. The CHAMP1 Complex promotes break-induced telomere clustering and synthesis.
a. Schematic of U2OS-TRF1-FokI cell and time course of IF/FISH following Dox/4-OHT
treatment. **b**. Representative FISH confocal images with telomere probe (green) treated
with/without TRF1-FokI in U2OS cells. DAPI was used to stain the nuclei. Scale bar, 5 µm. **c-d**.
CHAMP1 and POGZ KO U2OS cells were examined by telomere FISH after TRF1-FokI
treatment (a). Average telomere (Telo) foci size (c) and telomere number (d) per nucleus was
calculated using ImageJ. Error bars indicate SEM. More than 100 cells were counted.
****P<0.0001. Statistical analysis was performed using two-tailed Student's t test. **e**. PML
immunostaining combined with telomere FISH in wild-type (WT), sgCHAMP1 and sgPOGZ
U2OS cells after TRF1-FokI treatment (a). Colocalization of PML foci and telomere signals
(APBs) are shown as indicated. DAPI was used to stain the nuclei. Scale bar, 5 µm. **f**.
Quantification of APBs shown in (e). Error bars indicate SEM. More than 100 cells were
counted. ****P<0.0001. Statistical analysis was performed using two-tailed Student's t test. **g**.
Schematic of U2OS-TRF1-FokI cell and time course of IF assay following Dox/4-OHT and EdU
treatment. **h**. Representative confocal images of immunostaining of TRF2 and EdU foci in S
phase and non-S phase of U2OS cells following the treatment in (g). DAPI was used to stain the
nuclei. Scale bar, 5 µm. **i**. Quantification of the colocalized TRF2 and EdU foci in WT,
sgCHAMP1 and sgPOGZ U2OS cells. Error bars indicate SEM. More than 100 cells were
counted. ****P<0.0001. Statistical analysis was performed using two-tailed Student's t test. **j.** A
cartoon showing that CHAMP1 complex promotes telomere clustering, APBs formation and
facilitates Break-induced new telomeric DNA synthesis.
507
508
Fig. 4. The CHAMP1 complex establishes telomeric heterochromatin and promotes the
recruitment of HR and BIR factors. a. Schematic of the PICh approach used to define the
telomere specific proteome in WT and sgCHAMP1 U2OS cells after TRF1-FokI (-/+Dox/4-
OHT) treatment. Fractionated and precleared chromatin was hybridized to a biotinylated
telomeric probe and subsequently captured on magnetic beads. Telomere-associated proteins
were analyzed by silver staining, western blot analysis and mass spectrometry. PFA,
paraformaldehyde. **b.** Scatterplot of the telomere specific DSB response proteome profile
enriched by PICh in wild-type (WT) and sgCHAMP1 U2OS cells induced with TRF1-FokI.
Mass Spectrometry data was shown in Table S1. Scatterplot of log2 FC[(+Dox+1)/(-Dox+1) of
total peptide number] from two independent experiments. FC, fold change. Homologous
recombination (HR) pathway proteins with Log2 FC > 1 in WT were highlighted in red. Break-

induced replication (BIR) pathway proteins with Log2 FC > 1 in WT were highlighted in green.
CBX5 (HP1α) was highlighted in blue. The red dotted line indicates the same value of log2
FC(+Dox/-Dox) in both WT and sgCHAMP1 cells. **c.** Western blot from PICh experiments in
U2OS with/without Dox-induced TRF1-FokI. Telomere-associated proteins are blotted with the
indicated antibodies. **d**. Representative IF confocal images depicting the colocalization of HP1α
(Green) with TRF1-FokI (T1-F1, mCherry tagged, Red) in WT and sgCHAMP1 U2OS cells
induced with/without TRF1-FokI for 2 hours. DAPI (blue) was used to stain the nuclei. Scale
bar, 5 μm. **e**. Quantification of HP1α-TRF1-FokI colocalization events in (d). Error bars indicate
SEM. More than 100 cells were counted. ****P<0.0001. Statistical analysis was performed
using two-tailed Student's t test. **f**. Representative IF confocal images depicting the
colocalization of H3K9me3 (Green) with TRF1-FokI (T1-F1, mCherry tagged, Red) in WT and
sgCHAMP1 U2OS cells induced with/without TRF1-FokI for 2 hours. DAPI (blue) was used to
stain the nuclei. Scale bar, 5 μm. **g**. Quantification of H3K9me3-TRF1-FokI colocalization
events in (f). Error bars indicate SEM. More than 100 cells were counted. ****P<0.0001.
Statistical analysis was performed using two-tailed Student's t test. **h**. Representative IF confocal
images depicting the colocalization of RAD51 (Green) with TRF1-FokI (T1-F1, mCherry
tagged, Red) in WT, sgCHAMP1 and sgPOGZ U2OS cells induced with/without TRF1-FokI for
2 hours. DAPI (blue) was used to stain the nuclei. Scale bar, 5 μm. **i**. Quantification of RAD51-
TRF1-FokI colocalization events in (h). Error bars indicate SEM. More than 100 cells were
counted. ****P<0.0001. Statistical analysis was performed using two-tailed Student's t test.
540
541
Fig. 5. The CHAMP1 complex promotes the recruitment of SETDB1 to telomeres. a.
Scatterplot of Log2 (+Dox+1)/(-Dox+1) of total peptide number from the telomere specific DSB
response proteome profile enriched by PICh in wild-type (WT), sgCHAMP1 and sgPOGZ U2OS
cells induced with TRF1-FokI. Mass Spectrometry data was shown in Table S2. SETDB1 is
highlighted. **b**. Representative IF confocal images of SETDB1 (Green) colocalization with
telomeres with double-strand breaks (T1-F1, mCherry, Red) in wild-type (WT), sgCHAMP1 and
sgPOGZ U2OS cells induced with TRF1-FokI for 2 h. Arrow indicates SETDB1 and TRF1-FokI
colocalization. DAPI was used to stain the nuclei. Scale bar, 5 μm. **c**. Quantification of (a) for
number of SETDB1-mCherry colocalization events. Error bars indicate SEM. More than 100
cells were counted. ****P<0.0001. Statistical analysis was performed using two-tailed Student's
t test. **d.** (Top) AlphaFold2-Multimer (AF2)-predicted structural model of POGZ_ZnFcore
(peach)-SETDB1_Cter (cyan) complex. The key residues (shown in sticks) contributing to the
protein-protein interactions are highlighted in the inset including residues (C532, C535, H548
and H534) from POGZ_ZnFcore potentially forming a C2H2-type Zn finger that could be
critical in maintaining interaction between POGZ and SETDB1. (Bottom) The AF2-predicted
model of POGZ_ZnFcore-SETDB1_Cter complex is colored based on the confidence in the
model prediction (100-high (blue) and 50-low (red)) and corresponding PAE matrix showing the
confidence in the predicted interaction between POGZ_ZnFcore and SETDB1_Cter domains
along with iPTM socres for the AF2 prediction. **e.** Western blot showing GFP-
immunoprecipitation of GFP-vector control, GFP-CHAMP1 WT or deletion mutants, and the co-
immunoprecipitation of endogenous SETDB1 and POGZ. GAPDH acts as a negative control for
IP. **f.** Representative IF-FISH confocal images of CHAMP1 colocalization with telomeres in
wild-type (WT) and sgSETDB1 U2OS cells. Arrow indicates CHAMP1-telomere colocalization.
DAPI (blue) was used to stain the nuclei. Scale bar, 5 μm. **g**. Quantification of (f) for number of

CHAMP1-telomere colocalization events. Error bars indicate SEM. More than 100 cells were
counted. ****P<0.0001. Statistical analysis was performed using two-tailed Student's t test. **h**.
SETDB1 KO TRF1-FokI U2OS cells were examined by FISH using a FITC-labeled telomere
probe after DOX/4-OHT treatment. Average telomere foci size per nucleus was calculated using
ImageJ. Error bars indicate SEM. More than 100 cells were counted. ****P<0.0001. Statistical
analysis was performed using two-tailed Student's t test. **i.** PML immunostaining combined with
telomere FISH in SETDB1 KO U2OS cells after DOX/4-OHT treatment. Colocalization of PML
foci and telomere signals (APBs) are shown. Error bars indicate SEM. More than 100 cells were
counted. ****P<0.0001. Statistical analysis was performed using two-tailed Student's t test. **j.** A
cartoon showing that the recruitment of SETDB1 by CHAMP1 complex could potentially
facilitates a positive feedback loop that boosts the formation of telomere heterochromatin
clusters.
578
579
Fig. 6. HP1α and CHAMP1 are epistatic in the regulation of telomeric HR and clustering.
a-b. Wild-type (WT) and sgCHAMP1 U2OS-TRF1-FokI cells were treated with siControl or
siHP1α, followed by examination of FISH using a telomere probe, or PML combined with
telomere IF-FISH staining after DOX/4-OHT treatment. (a) Average telomere foci size per
nucleus was calculated using ImageJ. Error bars indicate SEM. More than 100 cells were
counted. ****P<0.0001, **P<0.01. Statistical analysis was performed using two-tailed Student's
t test. (b) Quantification of colocalization of PML foci and telomere signals (APBs). Error bars
indicate SEM. More than 100 cells were counted. ****P<0.0001, **P<0.01. Statistical analysis
was performed using two-tailed Student's t test. **c-d.** WT, sgHP1α, and sgHP1α U2OS cells with
ectopically expressed HP1α WT or mutant I165E were examined with FISH using a telomere
probe, or PML combined with telomere IF-FISH staining using a FITC-labeled telomere probe
after DOX/4-OHT treatment. (c) Average telomere foci size per nucleus was calculated using
ImageJ. Error bars indicate SEM. More than 100 cells were counted. ****P<0.0001. Statistical
analysis was performed using two-tailed Student's t test. (d) Quantification of colocalization of
PML foci and telomere signals (APBs). Error bars indicate SEM. More than 100 cells were
counted. ****P<0.0001. Statistical analysis was performed using two-tailed Student's t test. **e-f.**
WT, sgCHAMP1, and sgCHAMP1 U2OS cells with ectopically expressed CHAMP1 WT or
mutants (ΔN, 2A and ΔC) were examined with IF using anti-RAD51 or FISH using a telomere
probe. (e) Quantification of RAD51-TRF1-FokI colocalization events. Error bars indicate SEM.
More than 100 cells were counted. ****P<0.0001. Statistical analysis was performed using two-
tailed Student's t test. (f) Average telomere foci size per nucleus was calculated using ImageJ.
Error bars indicate SEM. More than 100 cells were counted. ****P<0.0001, **P<0.01.
Statistical analysis was performed using two-tailed Student's t test.
603
604
Fig. 7. Peripheral blood lymphocytes from Patients with CHAMP1 syndrome display
impaired heterochromatin clustering and defects in HR repair. a. Schematic of CHAMP1
mutants from patients with CHAMP1 syndrome. **b**. EBV-immortalized lymphoblast cell lines
derived from several CHAMP1 patients (Coriell) were examined by immunostaining with
H3K9me3 antibody. Representative images show the H3K9me3 foci. DAPI was used to stain the
nuclei. Scale bar, 5 μm. **c**. Quantification of H3K9me3 foci of the CHAMP1 patient lymphoblast
cells in (b). More than 100 cells were counted. ****P<0.0001. Statistical analysis was performed

using one-way ANOVA. **d**. Representative images of RAD51 foci formation in normal and
CHAMP1 patient lymphoblast cells following with/without 5Gy IR treatment. DAPI was used to
stain the nuclei. Scale bar, 5 μm. **e**. Quantification of RAD51 foci of the CHAMP1 patient
lymphoblast cells following with 5Gy IR treatment. More than 100 cells were counted.
****$P<0.0001$. Statistical analysis was performed using one-way ANOVA. **f**. 3-day cytotoxicity
analysis of the normal and CHAMP1 patient lymphoblast cells treated with various dose of
radiation. Cell viability was detected by CellTiter Glo (Promega). Error bars indicate SD, n=3
independent experiments. **g**. 3-day cytotoxicity analysis of the normal and CHAMP1 patient
lymphoblast cells treated with various dose of Olaparib. Cell viability was detected by CellTiter
Glo (Promega). Error bars indicate SD, n=3 independent experiments.
622

References

Dabin, J., Giacomini, G., Petit, E. & Polo, S. E. New facets in the chromatin-based regulation of genome maintenance. *DNA Repair (Amst)* **140**, 103702 (2024). https://doi.org/10.1016/j.dnarep.2024.103702

Goodarzi, A. A. *et al.* ATM signaling facilitates repair of DNA double-strand breaks associated with heterochromatin. *Mol Cell* **31**, 167-177 (2008). https://doi.org/10.1016/j.molcel.2008.05.017

Price, B. D. & D'Andrea, A. D. Chromatin remodeling at DNA double-strand breaks. *Cell* **152**, 1344-1354 (2013). https://doi.org/10.1016/j.cell.2013.02.011

Watts, F. Z. Repair of DNA Double-Strand Breaks in Heterochromatin. *Biomolecules* **6** (2016). https://doi.org/10.3390/biom6040047

Bannister, A. J. *et al.* Selective recognition of methylated lysine 9 on histone H3 by the HP1 chromo domain. *Nature* **410**, 120-124 (2001). https://doi.org/10.1038/35065138

Padeken, J., Methot, S. P. & Gasser, S. M. Establishment of H3K9-methylated heterochromatin and its functions in tissue differentiation and maintenance. *Nat Rev Mol Cell Biol* **23**, 623-640 (2022). https://doi.org/10.1038/s41580-022-00483-w

Hyun, K., Jeon, J., Park, K. & Kim, J. Writing, erasing and reading histone lysine methylations. *Exp Mol Med* **49**, e324 (2017). https://doi.org/10.1038/emm.2017.11

Chiolo, I. *et al.* Double-strand breaks in heterochromatin move outside of a dynamic HP1a domain to complete recombinational repair. *Cell* **144**, 732-744 (2011). https://doi.org/10.1016/j.cell.2011.02.012

Jakob, B. *et al.* DNA double-strand breaks in heterochromatin elicit fast repair protein recruitment, histone H2AX phosphorylation and relocation to euchromatin. *Nucleic Acids Res* **39**, 6489-6499 (2011). https://doi.org/10.1093/nar/gkr230

Noon, A. T. *et al.* 53BP1-dependent robust localized KAP-1 phosphorylation is essential for heterochromatic DNA double-strand break repair. *Nat Cell Biol* **12**, 177-184 (2010). https://doi.org/10.1038/ncb2017

Ryu, T. *et al.* Heterochromatic breaks move to the nuclear periphery to continue recombinational repair. *Nat Cell Biol* **17**, 1401-1411 (2015). https://doi.org/10.1038/ncb3258

Tsouroula, K. *et al.* Temporal and Spatial Uncoupling of DNA Double Strand Break Repair Pathways within Mammalian Heterochromatin. *Mol Cell* **63**, 293-305 (2016). https://doi.org/10.1016/j.molcel.2016.06.002

Dilley, R. L. & Greenberg, R. A. ALTernative Telomere Maintenance and Cancer. *Trends Cancer* **1**, 145-156 (2015). https://doi.org/10.1016/j.trecan.2015.07.007

Henson, J. D., Neumann, A. A., Yeager, T. R. & Reddel, R. R. Alternative lengthening of telomeres in mammalian cells. *Oncogene* **21**, 598-610 (2002). https://doi.org/10.1038/sj.onc.1205058

Draskovic, I. *et al.* Probing PML body function in ALT cells reveals spatiotemporal requirements for telomere recombination. *Proc Natl Acad Sci U S A* **106**, 15726-15731 (2009). https://doi.org/10.1073/pnas.0907689106

Zhang, J. M., Yadav, T., Ouyang, J., Lan, L. & Zou, L. Alternative Lengthening of Telomeres through Two Distinct Break-Induced Replication Pathways. *Cell Rep* **26**, 955-968 e953 (2019). https://doi.org/10.1016/j.celrep.2018.12.102

17 Vermeulen, M. *et al.* Quantitative interaction proteomics and genome-wide profiling of
epigenetic histone marks and their readers. *Cell* **142**, 967-980 (2010).
https://doi.org/10.1016/j.cell.2010.08.020
18 Itoh, G. *et al.* CAMP (C13orf8, ZNF828) is a novel regulator of kinetochore-microtubule
attachment. *EMBO J* **30**, 130-144 (2011). https://doi.org/10.1038/emboj.2010.276
19 Nozawa, R. S. *et al.* Human POGZ modulates dissociation of HP1alpha from mitotic
chromosome arms through Aurora B activation. *Nat Cell Biol* **12**, 719-727 (2010).
https://doi.org/10.1038/ncb2075
20 Garrity, M. *et al.* Neurodevelopmental phenotypes in individuals with pathogenic
variants in CHAMP1. *Cold Spring Harb Mol Case Stud* **7** (2021).
https://doi.org/10.1101/mcs.a006092
21 Hempel, M. *et al.* De Novo Mutations in CHAMP1 Cause Intellectual Disability with
Severe Speech Impairment. *Am J Hum Genet* **97**, 493-500 (2015).
https://doi.org/10.1016/j.ajhg.2015.08.003
22 Isidor, B. *et al.* De Novo Truncating Mutations in the Kinetochore-Microtubules
Attachment Gene CHAMP1 Cause Syndromic Intellectual Disability. *Hum Mutat* **37**, 354-
358 (2016). https://doi.org/10.1002/humu.22952
23 Levy, T. *et al.* CHAMP1 disorder is associated with a complex neurobehavioral phenotype
including autism, ADHD, repetitive behaviors and sensory symptoms. *Hum Mol Genet* **31**,
2582-2594 (2022). https://doi.org/10.1093/hmg/ddac018
24 Tanaka, A. J. *et al.* De novo pathogenic variants in CHAMP1 are associated with global
developmental delay, intellectual disability, and dysmorphic facial features. *Cold Spring
Harb Mol Case Stud* **2**, a000661 (2016). https://doi.org/10.1101/mcs.a000661
25 Dentici, M. L. *et al.* Expanding the phenotypic spectrum of truncating POGZ mutations:
Association with CNS malformations, skeletal abnormalities, and distinctive facial
dysmorphism. *Am J Med Genet A* **173**, 1965-1969 (2017).
https://doi.org/10.1002/ajmg.a.38255
26 Duan, J. *et al.* White-Sutton syndrome and congenital heart disease: case report and
literature review. *BMC Pediatr* **23**, 158 (2023). https://doi.org/10.1186/s12887-023-
03972-9
27 Ferretti, A. *et al.* POGZ-related epilepsy: Case report and review of the literature. *Am J
Med Genet A* **179**, 1631-1636 (2019). https://doi.org/10.1002/ajmg.a.61206
28 Fukai, R. *et al.* A case of autism spectrum disorder arising from a de novo missense
mutation in POGZ. *J Hum Genet* **60**, 277-279 (2015).
https://doi.org/10.1038/jhg.2015.13
29 Matsumura, K. *et al.* De novo POGZ mutations in sporadic autism disrupt the DNA-
binding activity of POGZ. *J Mol Psychiatry* **4**, 1 (2016). https://doi.org/10.1186/s40303-
016-0016-x
30 Tan, B. *et al.* A novel de novo POGZ mutation in a patient with intellectual disability. *J
Hum Genet* **61**, 357-359 (2016). https://doi.org/10.1038/jhg.2015.156
31 Ye, Y. *et al.* De novo POGZ mutations are associated with neurodevelopmental disorders
and microcephaly. *Cold Spring Harb Mol Case Stud* **1**, a000455 (2015).
https://doi.org/10.1101/mcs.a000455

32 Zhao, W. *et al.* POGZ de novo missense variants in neuropsychiatric disorders. *Mol Genet*
Genomic Med **7**, e900 (2019). https://doi.org/10.1002/mgg3.900

33 Zhao, W. *et al.* Rare inherited missense variants of POGZ associate with autism risk and
disrupt neuronal development. *J Genet Genomics* **46**, 247-257 (2019).
https://doi.org/10.1016/j.jgg.2019.04.002

34 Fujita, H. *et al.* CHAMP1-POGZ counteracts the inhibitory effect of 53BP1 on homologous
recombination and affects PARP inhibitor resistance. *Oncogene* **41**, 2706-2718 (2022).
https://doi.org/10.1038/s41388-022-02299-6

35 Heath, J. *et al.* POGZ promotes homology-directed DNA repair in an HP1-dependent
manner. *EMBO Rep* **23**, e51041 (2022). https://doi.org/10.15252/embr.202051041

36 Li, F. *et al.* CHAMP1 binds to REV7/FANCV and promotes homologous recombination
repair. *Cell Rep* **40**, 111297 (2022). https://doi.org/10.1016/j.celrep.2022.111297

37 Dev, H. *et al.* Shieldin complex promotes DNA end-joining and counters homologous
recombination in BRCA1-null cells. *Nat Cell Biol* **20**, 954-965 (2018).
https://doi.org/10.1038/s41556-018-0140-1

38 Ghezraoui, H. *et al.* 53BP1 cooperation with the REV7-shieldin complex underpins DNA
structure-specific NHEJ. *Nature* **560**, 122-127 (2018). https://doi.org/10.1038/s41586-
018-0362-1

39 Gupta, R. *et al.* DNA Repair Network Analysis Reveals Shieldin as a Key Regulator of NHEJ
and PARP Inhibitor Sensitivity. *Cell* **173**, 972-988 e923 (2018).
https://doi.org/10.1016/j.cell.2018.03.050

40 Mirman, Z. *et al.* 53BP1-RIF1-shieldin counteracts DSB resection through CST- and
Polalpha-dependent fill-in. *Nature* **560**, 112-116 (2018).
https://doi.org/10.1038/s41586-018-0324-7

41 Noordermeer, S. M. *et al.* The shieldin complex mediates 53BP1-dependent DNA repair.
Nature **560**, 117-121 (2018). https://doi.org/10.1038/s41586-018-0340-7

42 Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021). https://doi.org/10.1038/s41586-021-03819-2

43 Liu, J. *et al.* Enhancing alphafold-multimer-based protein complex structure prediction
with MULTICOM in CASP15. *Commun Biol* **6**, 1140 (2023).
https://doi.org/10.1038/s42003-023-05525-3

44 Brasher, S. V. *et al.* The structure of mouse HP1 suggests a unique mode of single peptide
recognition by the shadow chromo domain dimer. *EMBO J* **19**, 1587-1597 (2000).
https://doi.org/10.1093/emboj/19.7.1587

45 Machida, S. *et al.* Structural Basis of Heterochromatin Formation by Human HP1. *Mol*
Cell **69**, 385-397 e388 (2018). https://doi.org/10.1016/j.molcel.2017.12.011

46 Palm, W. & de Lange, T. How shelterin protects mammalian telomeres. *Annu Rev Genet*
42, 301-334 (2008). https://doi.org/10.1146/annurev.genet.41.110306.130350

47 Cubiles, M. D. *et al.* Epigenetic features of human telomeres. *Nucleic Acids Res* **46**, 2347-
2355 (2018). https://doi.org/10.1093/nar/gky006

48 Gauchier, M. *et al.* SETDB1-dependent heterochromatin stimulates alternative
lengthening of telomeres. *Sci Adv* **5**, eaav3673 (2019).
https://doi.org/10.1126/sciadv.aav3673

49   Udugama, M. *et al.* Mutations inhibiting KDM4B drive ALT activation in ATRX-mutated
754   glioblastomas. *Nat Commun* **12**, 2584 (2021). https://doi.org/10.1038/s41467-021-
755   22543-z

756   50   Cho, N. W., Dilley, R. L., Lampson, M. A. & Greenberg, R. A. Interchromosomal homology
757   searches drive directional ALT telomere movement and synapsis. *Cell* **159**, 108-121
758   (2014). https://doi.org/10.1016/j.cell.2014.08.030

759   51   Zhang, T. *et al.* Break-induced replication orchestrates resection-dependent template
760   switching. *Nature* **619**, 201-208 (2023). https://doi.org/10.1038/s41586-023-06177-3

761   52   Yeager, T. R. *et al.* Telomerase-negative immortalized human cells contain a novel type of
762   promyelocytic leukemia (PML) body. *Cancer Res* **59**, 4175-4179 (1999).

763   53   Zhang, H. *et al.* Nuclear body phase separation drives telomere clustering in ALT cancer
764   cells. *Mol Biol Cell* **31**, 2048-2056 (2020). https://doi.org/10.1091/mbc.E19-10-0589

765   54   Dilley, R. L. *et al.* Break-induced telomere synthesis underlies alternative telomere
766   maintenance. *Nature* **539**, 54-58 (2016). https://doi.org/10.1038/nature20099

767   55   Sun, X. *et al.* POGZ suppresses 2C transcriptional program and retrotransposable
768   elements. *Cell Rep* **42**, 112867 (2023). https://doi.org/10.1016/j.celrep.2023.112867

769   56   Jiang, W. Q. *et al.* Induction of alternative lengthening of telomeres-associated PML
770   bodies by p53/p21 requires HP1 proteins. *J Cell Biol* **185**, 797-810 (2009).
771   https://doi.org/10.1083/jcb.200810084

772   57   Hartlieb, S. A. *et al.* Alternative lengthening of telomeres in childhood neuroblastoma
773   from genome to proteome. *Nat Commun* **12**, 1269 (2021).
774   https://doi.org/10.1038/s41467-021-21247-8

775   58   Alabert, C. & Groth, A. Chromatin replication and epigenome maintenance. *Nat Rev Mol*
776   *Cell Biol* **13**, 153-167 (2012). https://doi.org/10.1038/nrm3288

777   59   Gaggioli, V. *et al.* Dynamic de novo heterochromatin assembly and disassembly at
778   replication forks ensures fork stability. *Nat Cell Biol* **25**, 1017-1032 (2023).
779   https://doi.org/10.1038/s41556-023-01167-z

780   60   Wang, H. *et al.* mAM facilitates conversion by ESET of dimethyl to trimethyl lysine 9 of
781   histone H3 to cause transcriptional repression. *Mol Cell* **12**, 475-487 (2003).
782   https://doi.org/10.1016/j.molcel.2003.08.007

783   61   Shi, G. *et al.* A critical role of telomere chromatin compaction in ALT tumor cell growth.
784   *Nucleic Acids Res* **48**, 6019-6031 (2020). https://doi.org/10.1093/nar/gkaa224

785   62   Voon, H. P. J., Collas, P. & Wong, L. H. Compromised Telomeric Heterochromatin
786   Promotes ALTernative Lengthening of Telomeres. *Trends Cancer* **2**, 114-116 (2016).
787   https://doi.org/10.1016/j.trecan.2016.02.003

788   63   White, J. *et al.* POGZ truncating alleles cause syndromic intellectual disability. *Genome*
789   *Med* **8**, 3 (2016). https://doi.org/10.1186/s13073-015-0253-0

Break-induced telomere synthesis