The Neural Underpinnings of Aphantasia: A Case Study of Identical Twins

---

Emma Megla[1*], Deepasri Prasad[2*], & Wilma A. Bainbridge[1,3]

1—Department of Psychology, University of Chicago, Chicago, IL

2—Department of Psychological and Brain Sciences, Dartmouth College, Hanover, NH

3—Neuroscience Institute, University of Chicago, Chicago, IL

* These authors contributed equally

Correspondence to: Emma Megla

emegla@uchicago.edu

Institute for Mind and Biology

940 E. 57th St.

Chicago, IL 60637

Summary

Aphantasia is a condition characterized by reduced voluntary mental imagery. As this lack of

mental imagery disrupts visual memory, understanding the nature of this condition can provide

important insight into memory, perception, and imagery. Here, we leveraged the power of case

studies to better characterize this condition by running a pair of identical twins, one with

aphantasia and one without, through mental imagery tasks in an fMRI scanner. We identified

objective, neural measures of aphantasia, finding less visual information in their memories which

may be due to lower connectivity between frontoparietal and occipitotemporal lobes of the brain.

However, despite this difference, we surprisingly found more visual information in the

aphantasic twin's memory than anticipated, suggesting that aphantasia is a spectrum rather than a

discrete condition.

Keywords

Visual imagery, long-term memory, perception, fMRI, encoding-recall similarity, functional

connectivity, SVM searchlight, representational similarity analysis

Introduction

What does your bedroom look like? For many of us, we can form a vivid mental image of

this place, filling our "mind's eye" with its visual details. However, the nature of these mental

representations—and their relationship to perception—is debated in the field. Are these

representations a recapitulation of what we viewed during perception, or have they been altered

in memory? The key to this fundamental question may be those with *aphantasia*, a condition

characterized by the lack of voluntary mental imagery.[1,2] Since aphantasia may serve as a natural

"knock-out" model of visual imagery and recall, it could highlight potential differences in our

perceptual and mnemonic representations. Here, we leveraged the identical genetics and shared

experiences of a unique case study: a pair of identical twins—one with aphantasia and one with

normal imagery—using neuroimaging to pinpoint differences in their memories stemming from

their different imagery experiences.

It is currently debated how visual perception relates to visual long-term memory. On one

hand, a collection of research has identified similarities between perception and memory, finding

that the same voxels activated during perception are reactivated during memory[3–6]. On the other

hand, more recent studies have identified meaningful differences between perception and

memory, uncovering that the voxels activated during memory are anterior to—rather than the

same as—those activated during perception.[7] Moreover, entirely different networks may even be

involved in perception than in memory.[8–10] In fact, the existence of aphantasia suggests

differences between perception and memory: if memory is a reinstatement of perception, then

how do aphantasics have intact perception, but disrupted memory?[11]

Although aphantasia could help define the relationship between perception and memory,

we first need to understand the nature of this condition. As aphantasia has largely been identified

through the Vividness of Visual Imagery Questionnaire (VVIQ),[12,13] resulting in estimates that

roughly 4% of the population has aphantasia,[2,14] this subjective measure has led to claims that it

could instead be a metacognitive or psychometric condition.[15] Indeed, there is little objective

evidence of aphantasia, although measures have been quantified in recent years. For aphantasics,

forming a mental image does not bias perception during subsequent binocular rivalry,[1] unlike for

those with normal imagery.[16,17] Additionally, those with aphantasia have a reduced skin-

conductance response when reading a frightening story compared to controls,[18] as they cannot

"see" these events in their minds. Therefore, is lack of imagery a *subjective* or *objective*

experience? Neuroimaging could help reveal the underlying nature of this condition, and

potentially identify additional objective measures.

To date, there has only been four published neuroimaging studies of aphantasia,[19–22] with

these studies largely taking a network approach. These neuroimaging studies suggest that

aphantasics may have reduced connectivity between their visual-occipital regions and other

regions of the brain, such as the prefrontal cortex[20] or temporal lobe regions,[19] but increased

connectivity among non-visual areas.[19,20] Similarly, a recent electroencephalography (EEG)

study found that mental imagery may be evoked starting in the left temporal lobe for aphantasics

compared to frontal areas in normal imagers.[21] Therefore, it seems that aphantasics may have

different networks dedicated to memory than their control counterparts. However, better

understanding aphantasia and memory's relationship to perception will also depend on

understanding the neural *representations* during memory. Ongoing work has started to tackle this

topic, such as comparing representations for low-level features of different items (e.g., faces and

shapes).[23]

In the present study, we use functional magnetic resonance imaging (fMRI) to identify

some of the first neural underpinnings of aphantasia, and examine the nature of memory, through

a pair of identical twins—one with aphantasia and one with normal imagery. The identical

genetics and shared experiences of these twins ensures that meaningful differences in memory

are likely due to their differing imagery experiences, making this an ideal sample to pinpoint

neural markers of aphantasia. By having the twins view and mentally imagine the same items,

we found that although the aphantasic twin does have lower memory quality, their memories still

contained an unexpected amount of visual information. Additionally, we also observed reduced

connectivity between occipitotemporal and fronto-parietal areas in the aphantasic twin. These

results not only identify some of the first objective, neural measures of aphantasia, but also

suggest that memory is more than a recapitulation of perception.

Results

We examined the mental imagery abilities of a pair of identical twins (31 years old, female)

raised in the same household, where one has aphantasia ("aphantasic twin") and one does not

("imager twin"). The twins engaged in two mental imagery tasks while in the fMRI scanner (**Fig**

1a). In the *Novel Imagery* task,[7] the twins encoded and subsequently mentally imagined the same

set of novel scene and object images. After mentally imagining each image, they rated the

vividness of that mental image using a three-point scale: 1—high vividness, 2—low vividness, or

3—no memory. In the *Familiar Imagery* task,[10] the twins were shown the text label of a familiar

person or place (e.g., 'Childhood Bedroom')—generated before the experiment—and were

subsequently asked to mentally imagine that person or place. They then rated the vividness of

that mental image using the same three-point scale.

In the following sections, we examine behavioral and neural differences between the

twins due to their differing mental imagery experiences. First, we verified differences in the

strength of their mental imagery using behavioral measures, including questionnaire results,

drawings, and vividness reports. Then, we used univariate and multivariate approaches to

identify neural correlates of aphantasia, where we unexpectedly found evidence for perceptual

information present in the aphantasic's imagery, though to a lesser degree than the imager.

Lastly, we found that the lesser degree of perceptual information in the aphantasic's imagery

may be due to different underlying strength of their functional connectivity patterns.

Figure 1. Methods and behavioral results. (a) Methods for the two imagery tasks. In the Novel Imagery task, participants first encoded a novel scene or object image for 6 sec. Then, there was a 4 sec distractor period in which the participants indicated an intact image amongst a stream of scrambled images. After a 1-4 sec randomized jitter, participants then recalled the original image using mental imagery for 6 sec. Lastly, they rated the vividness of their mental image using a three-point scale. There was a total of 96 trials. In the Familiar Imagery task, participants were first given a prompt which consisted of the text label of a personally familiar person or place. After a 1 sec mask of scrambled alphanumeric characters, the participants then mentally imagined the corresponding text prompt for 10 sec before rating the vividness of their mental imagery using a three-point scale. There was a 5-7 sec randomized jittered fixation between trials and 144 trials total. (b) Behavioral results. Whereas both twins drew many objects in detail from a scene during perception, the aphantasic twin drew starkly less from memory compared to the imager twin. The aphantasic twin additionally reported significantly lower vividness during mental imagery for both the novel imagery and familiar imagery tasks.

Lower subjective imagery strength for aphantasic twin

The strength of mental imagery is often measured through self-report surveys, such as the VVIQ,

which assesses the overall strength of mental imagery, and the Object-Spatial Inventory

Questionnaire (OSIQ), which probes object and spatial imagery abilities separately. The imager

twin reported scores within the standard imagery range (VVIQ=47, Object-OSIQ=56, Spatial-

OSIQ=38). However, the aphantasic twin had scores indicative of overall diminished imagery

and object imagery levels (VVIQ=24, Object-OSIQ=22), but intact spatial imagery (Spatial-

OSIQ=49), as is typical for aphantasic individuals.[24]

Drawings made from memory have been shown to be a more objective measure of

imagery experience.[24] When the twins drew three scene images from memory and perception

(see *Drawing Experiment*), we observed the same trends as the reported survey results. Whereas

both twins were able to accurately draw the scenes in detail during perception, the aphantasic

twin used starkly less detail—including no color—when drawing the scenes from memory (**Fig.**

1b).

Additionally, we analyzed the vividness reports collected after each in-scanner imagery

trial. A 2-way ANOVA of successfully remembered trials with participant (imager/aphantasic)

and task (novel imagery/familiar imagery) as factors revealed a significant effect of both

participant ($F_{(1,330)}=246.68$, $p<0.001$) and task ($F_{(1,330)}=8.44$, $p<0.004$), as well as a

significant interaction ($F_{(1,330)}=7.77$, $p=0.006$). For novel images, the imager twin reported

significantly higher imagery vividness for novel scenes and objects (*M*=1.91, *SD*=0.29) than the

aphantasic twin (*M*=1.38, *SD*=0.49; $t(190)=9.16$, $p<0.001$). The imager twin also reported

significantly higher vividness in their imagery for familiar people and places (*M*=1.90, *SD*=0.30)

than the aphantasic twin *(M*=1.14, *SD*=0.35; $t(140)=13.89$, $p<0.001$). However, although the

imager reported high vividness for both tasks—with no significant difference between tasks

(t(148.91)=0.10, p=0.92)—the aphantasic twin actually reported significantly *lower* vividness

during the familiar imagery than the novel imagery task (t(164.90)=3.61, p=0.003). This is

opposite to the trend typically found in control participants,[25] suggesting that despite the twins

having increased perceptual experience with familiar people and places than novel images, this

experience does not benefit imagery in aphantasics like it does those with normal imagery.

Similar univariate activation during imagery

Given the diminished mental imagery reported by the aphantasic twin, is there any information

contained in their mental images? If there is category information during aphantasic imagery,

then we should see differential univariate activation during imagery for different categories of

items during the Novel Imagery task. Indeed, we observed higher activation for scenes than

objects in scene-selective perceptual areas, such as the parahippocampal place area (PPA; **Fig.**

2a).

155   We additionally looked at the location of these areas, with a focus on the PPA (**Fig. 2a**),

as an anterior shift in peak voxel activity from perception to imagery (or memory) is thought to

reflect the more conceptual nature of mnemonic compared to perceptual representations.[26,27] The

peak voxel within the imager twin's right PPA was anteriorly shifted from perception (y=29) to

memory (y=34), although not for their left PPA (perception: y=30; imagery: y=30). However, we

found a similar magnitude—or even smaller—of an anterior shift in the aphantasic's left PPA

between perception (y=26) and memory (y=28), suggesting items do not get more semanticized

Figure 2. Univariate brain activity during the imagery tasks for both twins. (a) The location of PPA during perception and memory of the Novel Imagery task. The vertical green line indicates the location of the peak voxel activity in each condition. We observed an anterior shift in the peak voxel activity of PPA between perception and memory in both twins, with an equal (or even smaller) shift in the aphantasic compared to the imager. (b) A people>places contrast during the Familiar Imagery task. Using this contrast, we identified the recently discovered "familiar memory regions" in the medial parietal cortex in both twins, with their characteristic alternating pattern between familiar people and place selectivity. Each image is shown at a threshold of *p<0.001* unless otherwise noted, and all images are from the sagittal view. See also *Fig. S3* and *Table S2*.

in aphantasic memory. We also found the anterior shift to be unilateral in the aphantasic twin,

with no evidence of an anterior shift in their right PPA (perception: $y=29$; imagery: $y=29$).

Are regions sensitive to the recall of familiar concepts also active during memory? Areas

in the medial parietal cortex have been identified that alternate in their selectivity for familiar

people and familiar places during imagery.[9] To see if we could identify these areas in the

aphantasic twin, we tested a univariate contrast of people>places in each twin (**Fig. 2b**). We

found the characteristic alternating pattern of these familiar people and place memory regions in

both twins, suggesting that information specific to familiar people and places are also present

during aphantasic imagery. The locations of these areas aligned with where they have been found

previously.[9,28]

Different, though similar, brain patterns during imagery

As univariate activity revealed similarities—rather than differences—between the twins, what is

causing their phenomenological differences? We hypothesized that these differences may be

reflected in different *multivariate patterns* of activation during mental imagery, indicating

different information stored in their mental images. To test this, we ran a whole-brain support

vector machine (SVM) searchlight analysis on the neuroimaging data from the Novel Imagery

task (**Fig. 3a**). To determine the similarity in representations across participants and across task

phases, we trained an SVM to decode between objects and scenes in one condition (e.g., imager

perception), and tested the decoding accuracy between objects and scenes in the other condition

(e.g., aphantasic perception) within each searchlight region. To determine whether decoding was

above chance, we ran a permutation test within each searchlight region.

Figure 3. SVM searchlight methods and results. (a) Methods for cross decoding between conditions. Using the brain patterns within each searchlight region, we trained an SVM to distinguish between objects and scenes in one condition and tested on the other condition. Conditions were either between-participants (e.g., training on imager perception, testing on aphantasic perception) or within-participants (e.g., training on imager perception, testing on imager recall). To determine whether the voxels within a searchlight region were able to cross-decode above chance, we randomly swapped the image class labels for half of the training and test trials. We did this 100 times to build a null distribution to compare to the true decoding accuracy. (b) Voxels with significant decoding accuracy. Between-participants, there were many significant voxels able to cross-decode between the twins' representations during perception, whereas there were far fewer during their recall. The decoding accuracy between the twins' perceptual representations was also significantly higher than between their recall representations. Within-participants, there was a significantly higher decoding accuracy within the imager twin. However, the aphantasic twin had a surprisingly similar number of voxels as well as decoding accuracy as the imager twin. (c) Voxels with significantly higher decoding accuracy in one condition versus another**.** Whereas visual areas, including PHC, were significantly more similar between the twins' perception than their recall, few areas emerged with higher similarity between their recall. Surprisingly, visual areas, including the PHC, shared significantly more similarity in their perceptual and mnemonic representations for the aphantasic than the imager. Each image is shown at a threshold of *p<0.001*, but all key regions reported survive cluster threshold correction (see *Supplemental Results 1* and *Fig. S1*). See also *Supplemental Results 2* and *Fig. S2* for an ROI-based approach.

First, given that aphantasics are thought to have intact perception but disrupted imagery,

we hypothesized that there would be similarity between the twins' perceptual representations,

but not their recall representations (**Fig. 3b**). When we trained on the imager's perceptual

representations and tested on the aphantasic's, we found a large number of voxels that were able

to decode above chance (4511 voxels), with an average above-chance decoding accuracy of

66.8% (*SD*=5.5%). We also surprisingly found decodability between the twins' recall

representations, suggesting at least some shared information during imagery. However, this

average decoding accuracy (*M*=60.7%, *SD*=2.0%) was significantly lower than between their

perceptual representations (t(1192.95)=-48.64, p<0.001), and far fewer voxels were able to

decode above chance (423 voxels).

Further, we hypothesized that if there is less perceptual information in aphantasic

imagery, then there should be higher similarity between the imager twin's perceptual and

mnemonic representations than between the aphantasic's (**Fig. 3b**). Within the imager, we found

similarity between their perceptual and mnemonic representations, with 311 voxels able to

decode above chance with 68.78% (*SD*=2.24) accuracy. However, we found a *surprisingly*

similar degree of successful decoding between the aphantasic's perceptual and mnemonic

representations, with a similar number of significant voxels (263 voxels) and average decoding

accuracy (67.89%, *SD*=2.50%). Although the imager twin's decoding accuracy was significantly

higher than the aphantasic's (t(530.57)=4.49, p<0.001), the numerical difference of only ~1%

suggests that there might be more visual information present in memory for the aphantasic twin

than we originally predicted. We replicated similar cross-decodability of perception and memory

in both twins when targeting mental imagery areas as a region of interest (see *Supplemental*

Results 2 and *Fig. S2*).

What is the content of the information shared between conditions? To answer this, we

first determined areas that had significantly higher decoding accuracy between the twins'

perceptual than between their recall representations using permutation testing (**Fig. 3c**). Many

visual areas, including those extending along the parahippocampal cortex (PHC), had

significantly higher decoding accuracy between the twins' perceptual than between their recall

representations. However, only a few areas—and none visual—had significantly higher decoding

accuracy between their recall than between their perceptual representations. These results are in-

line with what we would expect, with more visual information shared between the twins during

perception than recall. Within-participants, given the lack of visual information in aphantasic

memory, we predicted that visual memory areas would share significantly more information

between perception and recall in the imager than the aphantasic twin. However, we surprisingly

found evidence contrary to our prediction, with PHC and the hippocampus sharing significantly

more similar representations during perception and memory in the aphantasic twin. As this

posterior PHC region aligns with where the PPA is typically found,[29] this surprisingly suggests

the presence of visual information in aphantasic memory. Within the imager, we found that the

inferior parietal lobule (IPL) had significantly more similar representations during perception

and memory in the imager than the aphantasic twin, which could suggest some immediate

consolidation of visual information in the imager twin.[30]

Different brain patterns during familiar imagery

Although we found evidence of visual information in memory for newly-learned images for the

aphantasic twin, do we find this same evidence for more consolidated, highly familiar items? We

tested this using the Familiar Imagery task—as this task required mentally imagining a

Figure 4. Representational similarity during familiar imagery. To determine whether there is coarse level (person vs. place) visual information in aphantasic memory during familiar imagery, we correlated brain activity from the PHC region between every pair of stimuli. We quantified the amount of coarse level information by calculating a discrimination index (D) for each twin, which subtracts the degree of neural similarity within category – between category. Although we found evidence of coarse level visual information in the imager twin, we found nearly next to no discrimination between people and places in the aphantasic twin. Indeed, D was significantly higher in the imager than the aphantasic twin.

personally familiar person or place *without* any perceptual information, the twins had to conjure

visual detail from longer-term memory stores to accomplish this task.

We constructed a representational similarity matrix[31] by correlating brain activation

between pairs of trials in the same PHC region that contained higher similarity between

perception and memory in the aphantasic twin than the imager twin (see **Fig. 4**). We quantified

coarse level information by calculating discrimination indices (*D*), which subtracts the

correlation *between* conditions (e.g., people and places) from the correlation *within* conditions

(e.g., people and people). Therefore, if there is visual information in the aphantasic twin's mental

imagery when pulling from longer-term stores, then there should be positive discriminability for

people versus places. However, we found a *D* close to 0 in the aphantasic twin, which was

significantly lower than the positive *D* (0.098) in the imager twin (permutation testing: p<0.001).

In other words, although this region contained category-level visual information for newly-

learned images for the aphantasic twin, this visual information seemed to dissipate when drawing

from longer-term memory stores.

Lower and less visually-based connectivity between key regions in aphantasic twin

What could be the cause for the different amount of visual information in aphantasic memory

between the two imagery tasks? To explore this, we quantified the strength of the twins'

functional connectivity during rest (**Fig. 5**). To compare the strength of their connections, we

subtracted their correlation values between each pair of nodes (imager – aphantasic) and

averaged across all the connections within lobes.

We overall found trends that replicate prior work, finding that the aphantasic twin had

lower connectivity between their occipital lobe and both the prefrontal lobe[20,23] as well as the

parietal lobe.[23] However, we did not find lower connectivity between the occipital and temporal

lobe as found previously.[19] We additionally found that the aphantasic twin had reduced

connectivity between their temporal lobe and both their prefrontal and parietal lobes (though

only in the right hemisphere). Overall, the disconnect between occipitotemporal and fronto-

parietal lobes in the aphantasic twin could interestingly hint at visual information initially

making it into aphantasic memory, but unsuccessfully being consolidated into longer-term stores.

All correlation values between lobes are reported in *Table S1*.

Different language lateralization

Although identical twins raised in the same household are as similar as possible for two

individuals, there is one notable physiological difference between these twins. Namely, whereas

the imager twin self-reported as right-handed, the aphantasic twin self-reported as left-handed

Figure 5. Differences in resting state functional connectivity between the lobes of the brain. Red means a higher correlation between two lobes in the imager, whereas blue means higher correlation in the aphantasic. Interestingly, we generally found lower connectivity between lobes housing immediate memory processes (temporal and occipital) and lobes housing consolidated memory processes (parietal and prefrontal) in the aphantasic twin, which could account for the differences we found between imagery tasks. These connections of interest are outlined in black. See *Table S1* for all correlation values.

(called "mirror twins"). This opposite handedness was verified using the Edinburgh Handedness

Inventory (EHI; imager twin EHI=1, aphantasic twin EHI=-1). As handedness can correlate with

brain lateralization,[33] this meant that the laterality between the twins could also be opposite. We

determined lateralization of the brain through a language localizer, in which the twins read words

versus nonwords, to calculate a laterality index (LI). We found that the imager twin has left-side

language localization (LI=0.225), but the aphantasic twin has bilateral dominance (LI=0.177),

with a trend towards left-side language localization.

Discussion

In this work, we leveraged a rare sample of participants—identical twins, one with

aphantasia and one with normal imagery—which allowed us to identify some of the first neural

underpinnings of aphantasia. First, we found similarity between the aphantasic and imager using

univariate methods, with areas such as PPA and "familiar memory regions" active during

aphantasic memory. Second, when examining differences in multivariate patterns during

memory between the twins, although we found significantly more similarity between the

imager's perceptual and mnemonic representations, we also found unexpected similarity between

these representations for the aphantasic. In fact, visual areas in the PHC contained significantly

higher similarity between the aphantasic's perceptual and mnemonic representations than the

imager's. Although these findings suggest visual information in aphantasic memory, we did not

find evidence for this during familiar imagery. Lastly, we found that the lack of visual

information in aphantasic memory may be attributed to lower functional connectivity between

occipitotemporal and frontoparietal areas.

As there have only been a handful of published neuroimaging studies on aphantasia—and

none looking at the content of aphantasic memory—the current results help build a foundation

for our current understanding of the condition. Here, we found evidence that there is indeed a

difference between aphantasic memory content compared to controls, with our results suggesting

that there is significantly less visual information in memory for both newly-learned images and

familiar people and places. These neural results suggest that the lack of imagery is an objective

experience, supporting other objective behavioral findings,[1,18] and that there may be ways to

identify aphantasia on the neural level. The overall finding of less visual information in

aphantasic memory also aligns with previous, more subjective measures of aphantasic memory

content, such as recalling fewer objects and using less color when drawing scenes from

memory.[24]

However, we also found evidence that memory content for the aphantasic may contain

more visual information than we originally predicted. Although the difference in decoding

accuracy between perception and memory in the imager twin was significantly higher than in the

aphantasic twin, the accuracy was unexpectedly similar (only ~1% difference). In fact, visual

areas in the PHC had significantly *higher* decoding between perception and memory in the

aphantasic twin, suggesting that there is still a surprising degree of perceptual information in

aphantasic memory for newly-learned images. Univariate approaches also revealed intact

category-level visual information for the aphantasic, with activation of PPA, OPA, and LO

during memory for newly-learned images. We even found activation of regions selective to recall

of familiar people and places ("familiar memory regions"), suggesting at least some intact

memory content during familiar imagery as well.

The finding of less visual information in aphantasic memory during familiar imagery

compared to novel imagery also suggests that the amount of visual information may depend on

when the information was learned. Whereas novel imagery involved mentally imagining an

image that was shown shortly before, familiar imagery required mentally imagining a familiar

person or scene without any previous visual information shown. Therefore, it is possible that

aphantasics can maintain some visual information shortly after encoding, but that this visual

information dissipates more in aphantasics than imagers when the memory becomes

consolidated. Indeed, upcoming work may support this idea, which reports aphantasics

maintaining visual information in early visual cortex during working memory.[34]

Additionally, our results may suggest that there is a transformation between perception

and memory representations, even for those with normal imagery. As the aphantasic twin's

memory lacks perceptual information, it likely undergoes a transformation in representation from

perception (e.g., becomes more semanticized). As a result, their memory serves as a powerful

comparison to determine whether such a transformation occurs even for those who report more

visual information in memory. Since we found that there was comparable cross-decoding

between perception and memory between the twins, this suggests the removal of some perceptual

information—and thereby a transformation—in the imager's memory as well. Similarly, we also

found a similar degree of an anterior shift in peak voxel activity for areas like PPA between the

imager and aphantasic. These results therefore align with previous studies that have found

differences between perception and memory.[7–10]

In addition to the novel insight our findings provide on aphantasia, our work also

coincides with previous network-based studies. Specifically, we found both lower connectivity

between the aphantasic twin's occipital lobe and their prefrontal[20,23] and parietal lobes.[23]

However, we did not find lower connectivity between the aphantasic's occipital and temporal

lobe as reported previously,[19] and we found a new pattern of lower connectivity between the

aphantasic's temporal lobe with their prefrontal and parietal lobes. In total, these trends in

connectivity in the aphantasic suggest that there could be a lack of access to visual information

for consolidated memories. Indeed, the occipital lobe is widely known to house visual areas like

OPA[35] and the temporal lobe to house the hippocampus and other visual areas like PPA.[36] In

contrast, the medial prefrontal cortex and posterior parietal cortex are thought to be two areas

that house longer-term stores in the greater neocortex after memory consolidation,[37,38] which

involves migration of memories from the hippocampus.[30] Therefore, it is possible that more

visual information is lost during consolidation for aphantasics than imagers.

Lastly, the present study raises important avenues of research for future work. Although

the twins are identical, we did find that the imager twin processed language in the left

hemisphere, whereas the aphantasic twin processed language bilaterality (though left-hemisphere

leaning). As visual processing is more right lateralized,[39,40] it is possible that bilateral language

processing inhibited some of this visual processing. Therefore, future work could investigate

whether there is a connection between language lateralization and imagery ability. In addition,

the presence of visual information in aphantasic memory in the present study suggests that

aphantasia may be more of a spectrum than a discrete condition. Indeed, although the aphantasic

twin reported imagery within the aphantasic range, they did report some visual information in

imagery. Therefore, it may be valuable for future studies investigating aphantasia to recruit

participants with the lowest VVIQ score, indicating the complete absence of visual imagery.

In conclusion, this case study of identical twins was able to characterize aphantasia in

new and valuable ways, quantifying their memories as lacking visual information even on the

most objective, neural level. However, this study also revealed that there can still be a surprising

level of visual information for someone with aphantasia, at least for newly-learned images.

Methods

Participants

Two identical twins (31 years old, female) raised in the same household participated in this

experiment. Their overall mental imagery ability was assessed using the Vividness of Visual

Imagery Questionnaire (VVIQ),[12] and separable object and spatial imagery abilities using the

Object-Spatial Imagery Questionnaire (OSIQ).[41] Handedness was also assessed using the

Edinburgh Handedness Inventory.[42] The subjects had corrected vision and wore MRI-compatible

lenses during the scan. Subjects consented to participation, following the guidelines approved by

the University of Chicago Institutional Review Board (IRB20-0233), and were compensated for

their time.

Tasks

Perceptual localizer

A perceptual localizer was run to identify scene-, face-, and object-selective regions, but was not

used in analysis for this paper. In this localizer, participants viewed four 16 sec blocks of images.

Each block contained a single category of images: faces, objects, scenes, or scrambled images.

Participants indicated consecutive repeats of images by pressing the response button. Participants

completed one run of this task. All tasks performed in the scanner were displayed using

Psychtoolbox.[43]

Novel imagery

The item-based imagery task was adapted from Bainbridge, Hall, et al.[7] In each trial, participants

were presented with an image to view for 6 sec. After a 1 sec fixation, they completed a 4 sec

distractor where they viewed a series of scrambled images with one intact image and responded

when they saw the intact image. After a 1-4 sec jittered fixation, they were instructed to recall

the previously shown image for 6 sec as vividly as possible. At the end of each trial, they rated

the vividness of their memory as either high vividness, low vividness, or no memory.

Participants viewed 48 images of scenes and 48 images of objects presented against a white

background, for a total of 96 images. Subjects completed 4 runs of this task with 24 trials each,

equally balanced across the stimulus hierarchy. The order of image presentation was the same

between participants.

Familiar imagery

The familiar imagery task was modified from Steel et al.[10] Before the scan, the twins generated a

list of 36 personally familiar places and 36 personally familiar people together. In each trial of

the experiment, participants were prompted with the name of one of the people (e.g., mom) or

places (e.g., childhood bedroom) for 1 sec. After a 1 sec dynamic alphanumeric mask, they were

asked to recall the person or place associated with the prompt as vividly as possible for 10 sec.

After recall, they were asked to rate the vividness of their imagery as either high vividness, low

vividness, or no memory. Participants completed 4 runs of this task with 18 trials in each run.

Language localizer

The language localizer was adapted from Fedorenko et al.[44] In each trial, participants were

presented with a 12-unit sequence of either words that formed a sentence or nonwords. Each unit

was presented individually for 450 msec. Subjects completed 1 run of this task, with 16 blocks of

3 trials of either words or nonwords.

Drawing experiment

The drawing experiment was performed to obtain visual representations of the participants'

perceptual and mnemonic content. During the experiment, participants first encoded three scene

images sequentially (a bedroom, living room, and kitchen) for 10 sec each before recalling them

using drawing. The canvas used to create the drawings matched the size of the encoded images

(500 × 500 pixels). While drawing, participants had access to a range of colors, an erasure tool,

and an undo tool. Participants next performed a short old/new recognition task with the three

target images and three foil images from the same scene categories. Lastly, the participants were

sequentially shown the original three scene images alongside the drawing canvas. The

participants were instructed to copy each image using drawing.

MRI data collection and analysis

Neuroimaging data was collected at the University of Chicago using a 3T Philips Achieva MRI

scanner with a 32-channel phased-array head coil. Anatomical scans used a T1 MPRAGE

structural scan with a resolution of 1×1×1 mm voxels. Functional scans used a gradient echo-

planar T2* sequence (39 axial slices parallel to the anterior commissure-posterior commissure

line; 64×64 matrix; FoV=192×192 mm; TR=2000 msec; TE=28; 0.5 mm gap; flip angle=77

degrees; 3×3×3 mm voxels). We preprocessed the functional scans using the same protocol as

prior studies,[7] which included slice time correction and motion correction using the Analysis of

Functional NeuroImages (AFNI) software.[45] No spatial smoothing was applied. Functional data

were aligned to Montreal Neurological Institute (MNI) space.

Whole-brain univariate analyses

We ran general linear models (GLM) to perform whole-brain univariate analyses on the Novel

Imagery and Familiar Imagery tasks. For Novel Imagery, all trials were modeled separately (e.g.,

recalling farmhouse #4). Whole-brain *t*-contrasts were then calculated by grouping trials along

the dimensions of object/scene and encoding/recall. Distractor periods were modeled separately.

For Familiar Imagery, all trials were also modeled separately (e.g., Student Art Gallery), with

trials grouped along the dimensions of recalling people/places for whole-brain *t*-contrasts. Both

GLMs additionally included six regressors for head motion. The individual trial beta values were

used for multivariate analyses for both imagery tasks.

Defining regions of interest

We identified regions of interest (ROIs) using functional and anatomical criteria using data from

the Novel Imagery and Familiar Imagery tasks. From the Novel Imagery task, we identified and

focused analyses on scene-selective area parahippocampal place area (PPA) using a

scenes>objects contrast. Additionally, we also used a scenes>objects contrast to identify

occipital place area (OPA) and medial place area (MPA) as well as an objects>scenes contrast to

identify object-selective area lateral occipital (LO; see *Table S2* for coordinates of these

additional regions). These regions were localized during perception and memory separately.

From the Familiar Imagery task, we identified "familiar memory regions" that have been found

in the medial parietal area using a people>places contrast, and compared the coordinates in MNI

space to where they have been localized previously.[9,28]

When determining the location of the ROIs during perception and memory of the Novel

Imagery task, we located their peak voxel activation in MNI space. We first used the threshold of

p<0.001 to identify these regions, but iteratively lowered the threshold until we were able to

identify them. Although PPA was largely identified using our most conservative threshold, the

aphantasic's left and right PPAs were notably not identified during memory until we used a more

liberal threshold (left: p<0.02; right: p<0.01). However, we also found similar evidence for an

anterior shift when using a more conservative threshold and expanding to the greater medial

temporal lobe region (see *Fig. S3*).

<u>Whole-brain SVM searchlight analyses</u>

We performed four whole-brain SVM searchlight analyses to determine representational

differences between participants and tasks during the Novel Imagery task. Between-participants,

we used the voxels within each searchlight (sphere radius=3 voxels) to train an SVM to

differentiate between objects and scenes in the imager and tested the model to differentiate

objects and scenes in the aphantasic. We did this twice: first for their representations during

perception, and second for their representations during memory. If there are similar

representations between the imager and aphantasic in a searchlight area, then there will be higher

(and above chance) decoding accuracy. Significance at each searchlight area was determined

through permutation testing, in which we performed 100 iterations of randomly swapping half of

the scene and object labels during training, and swapping those same labels during test to build a

null distribution. We used the same logic within-participants (but between tasks), where we

trained an SVM on a participant's perception and tested on their recall. We performed this for

both the aphantasic twin and the imager twin.

We additionally performed a second set of permutation tests to determine searchlight

areas that had significantly different decoding accuracy for either (1) encoding or recall or (2) for

the imager or the aphantasic twin. We ran this permutation test by randomly swapping half of the

condition labels (e.g., encoding or recall) between training and test. We performed this random

swap 100 times to build a null distribution, with significance set at $p<0.05$ for all permutation

tests. For all SVM searchlight results, statistics and brain visualizations are shown using the

uncorrected threshold of $p<0.001$, but we find the same trends and key brain regions when using

cluster threshold correction (*see Supplemental Results 1 and Fig. S1*). For cluster threshold

correction, we performed 1-sided thresholding for the initial SVM searchlight analyses and bi-

sided thresholding when comparing between the SVM searchlight conditions (i.e., when results

could be positive and negative).

<u>Representational similarity analyses and discrimination indices</u>

We conducted representational similarity analyses[31] on the Familiar Imagery task data to

determine the amount of visual information present in memory when recalling familiar people

and places. We performed these analyses in the PHC region identified from the whole-brain

searchlight SVM analyses as having unexpectedly higher similarity between perception and

memory for the aphantasic than the imager twin. We built a representational similarity matrix

(RSM) for each twin by correlating (Pearson's correlation) the activation of each voxel of the

PHC region between each pair of trials. Therefore, trials with a higher Pearson's correlation have

more similar representations.

If there is visual information present during memory of familiar people and places, then

we would expect more similarity for within-category trials (e.g., within-people) than between-

category trials (i.e., between people and places) as there would be more shared visual

information within-categories. Therefore, we calculated discrimination indices (*D*) by

subtracting the average of all between-category correlations from the average of all within-

category correlations.[46] To determine whether there was any significant difference in the

discriminability of people vs. places between participants, we performed a permutation test by

calculating the difference in D between participants when half of the trials were randomly

swapped between participants. We performed 1000 iterations of this random swapping to build a

null distribution to compare to the true difference in discriminability between participants.

<u>Functional connectivity analysis</u>

To determine participants' functional connections in the brain at rest, we had the twins watch a

10-min video titled *Inscapes*[47] that contained no semantic or social information while in the

scanner. The functional data were preprocessed and analyzed using a separate pipeline from the

other collected MRI data to more closely follow recent studies of functional connectivity.[48,49]

Preprocessing involved using *afni_proc* to remove outliers, perform time-slice correction, align

to the anatomical scan, register volumes to the TR with the least amount of motion, and align to

MNI space. Additionally, white matter and CSF masks were created for each participant using

FMRIB Software Library (FSL) and were regressed out of the data. To control for motion,

volumes in which 5% of the voxels contained motion outliers were removed as well as volumes

following a change of at least 0.2 mm of motion from the volume. As this last step left too few

volumes for analysis for the aphantasic twin, we used a slightly more liberal threshold of

censoring out volumes with at least 0.3 mm of motion for the aphantasic twin.

For the functional connectivity analysis, we parcellated the pre-processed data into the

268 nodes of the Shen Brain Atlas.[50] Since analysis depended on directly comparing the strength

of functional connections between the twins, we removed three nodes (2 prefrontal nodes, 1

temporal node) that were filtered out during preprocessing in at least one participant. After

calculating the mean time series for each node, we used Pearson's correlation to correlate the

mean times series between each pair of nodes and create a functional connectivity matrix. Lastly,

to compare the strength of connections between participants on the lobe level, we averaged

across the correlation values within each lobe (i.e., averaged across the nodes) and subtracted

these averages between participants (imager twin – aphantasic twin).

Language localizer analysis

We ran a language localizer analysis to test whether the twins' language areas were lateralized to

different hemispheres. We determined language lateralization using the previously established

method of calculating a laterality index (LI),[51,52] which involved comparing the number of voxels

active in the right (RH) versus the left hemisphere (LH) for words>nonwords at a threshold of

$p \leq 0.001$. Specifically, we followed the formula $LI=(LH–RH)/(LH+RH)$ and did not include

voxels within the cerebellum, as the cerebellum can show opposite trends to the rest of the

brain.[53] Therefore, a positive LI indicates laterality towards the left hemisphere, whereas a

negative LI indicates laterality towards the right hemisphere. However, we set a laterality

threshold of 0.2 following the most common convention,[54,55] which meant that laterality was

considered bilateral until the LI was > +/- 0.2.

Mental imagery ROI SVM analysis

To determine whether the SVM searchlight results replicate using a region of interest (ROI)-

based approach, we used the tool *Neurosynth*[56] to localize a mental imagery ROI. When given a

term, Neurosynth performs a meta-analysis on all published fMRI studies in its database that

includes that term in the abstract, and then determines the voxels that are preferentially active for

that term. These voxel maps are then corrected using a false discovery rate (FDR) of 0.01.

Therefore, this is a powerful method of deriving ROIs based on data from many studies.

For the present ROI, we used the term "mental imagery", which created a brain map

based on 84 published studies. We focused on replicating the SVM searchlight results with a

mental imagery ROI because this area should theoretically (1) be more active during the

imager's than the aphantasic's recall, given that the aphantasic has impaired mental imagery and

(2) share some similarity between perception and recall, given that imagery is thought to involve

some reactivation of perception. These were the same key hypotheses tested with the SVM

searchlight. Therefore, to see whether we found the same trend of results, we averaged across the

decoding accuracies for each voxel within the mental imagery ROI for each SVM searchlight

condition. We only included voxels that were in the mask and inside the participants' brains.

Results for this analysis are reported in *Supplemental Results 2* and *Fig. S2*.

Acknowledgements

We would like to thank Hayoung Song and Ziwei Zhang for their help with the functional

connectivity analysis. The authors would like to thank the MRI Research Center, the University

of Chicago (MRIRC, RRID:SCR_024723) for their assistance in MRI data acquisition of this

study. The present work was supported by the National Eye Institute (R01-EY034432) to

W.A.B.

References

1. Keogh, R. & Pearson, J. The blind mind: No sensory visual imagery in aphantasia. *Cortex*

105, 53–60 (2018).

2. Zeman, A., Dewar, M. & Della Sala, S. Lives without imagery – Congenital aphantasia.

Cortex **73**, 378–380 (2015).

3. Cichy, R. M., Heinzle, J. & Haynes, J.-D. Imagery and Perception Share Cortical

Representations of Content and Location. *Cereb. Cortex* **22**, 372–380 (2012).

4. Ishai, A., Ungerleider, L. G. & Haxby, J. V. Distributed Neural Systems for the Generation

of Visual Images. *Neuron* **28**, 979–990 (2000).

5. O'Craven, K. M. & Kanwisher, N. Mental Imagery of Faces and Places Activates

Corresponding Stimulus-Specific Brain Regions. *J. Cogn. Neurosci.* **12**, 1013–1023 (2000).

6. Reddy, L., Tsuchiya, N. & Serre, T. Reading the mind's eye: Decoding category information

during mental imagery. *NeuroImage* **50**, 818–825 (2010).

7. Bainbridge, W. A., Hall, E. H. & Baker, C. I. Distinct Representational Structure and

Localization for Visual Encoding and Recall during Visual Imagery. *Cereb. Cortex* **31**,

1898–1913 (2021).

8. Baldassano, C., Esteva, A., Fei-Fei, L. & Beck, D. M. Two Distinct Scene-Processing

Networks Connecting Vision and Memory. *eNeuro* **3**, ENEURO.0178-16.2016 (2016).

9. Silson, E. H., Steel, A., Kidder, A., Gilmore, A. W. & Baker, C. I. Distinct subdivisions of

human medial parietal cortex support recollection of people and places. *eLife* **8**, (2019).

10. Steel, A., Billings, M. M., Silson, E. H. & Robertson, C. E. A network linking scene

perception and spatial memory systems in posterior cerebral cortex. *Nat. Commun.* **12**, 2632

(2021).

11. Dijkstra, N., Bosch, S. E. & van Gerven, M. A. J. Shared Neural Mechanisms of Visual

Perception and Imagery. *Trends Cogn. Sci.* **23**, 423–434 (2019).

12. Marks, D. F. Visual Imagery Differences in the Recall of Pictures. *Br. J. Psychol.* **64**, 17–24

(1973).

13. Zeman, A. *et al.* Phantasia–The psychological significance of lifelong visual imagery

vividness extremes. *Cortex* **130**, 426–440 (2020).

14. Dance, C. J., Ipser, A. & Simner, J. The prevalence of aphantasia (imagery weakness) in the

general population. *Conscious. Cogn.* **97**, 103243 (2022).

15. de Vito, S. & Bartolomeo, P. Refusing to imagine? On the possibility of psychogenic

aphantasia. A commentary on Zeman et al. (2015). *Cortex J. Devoted Study Nerv. Syst.*

Behav. **74**, 334–335 (2016).

16. Pearson, J., Clifford, C. W. G. & Tong, F. The Functional Impact of Mental Imagery on

Conscious Perception. *Curr. Biol.* **18**, 982–986 (2008).

17. Pearson, J., Naselaris, T., Holmes, E. A. & Kosslyn, S. M. Mental Imagery: Functional

Mechanisms and Clinical Applications. *Trends Cogn. Sci.* **19**, 590–602 (2015).

18. Wicken, M., Keogh, R. & Pearson, J. The critical role of mental imagery in human emotion:

insights from fear-based imagery and aphantasia. *Proc. R. Soc. B Biol. Sci.* **288**, 20210267

(2021).

19. Zeman, A. Z. J. *et al.* Loss of imagery phenomenology with intact visuo-spatial task

performance: A case of 'blind imagination'. *Neuropsychologia* **48**, 145–155 (2010).

20. Milton, F. *et al.* Behavioral and Neural Signatures of Visual Imagery Vividness Extremes:

Aphantasia versus Hyperphantasia. *Cereb. Cortex Commun.* **2**, tgab035 (2021).

21. Furman, M. *et al.* Cortical activity involved in perception and imagery of visual stimuli in a

subject with aphantasia. An EEG case report. *Neurocase* **28**, 344–355 (2022).

22. Zhao, B., Della Sala, S., Zeman, A. & Gherri, E. Spatial transformation in mental rotation

tasks in aphantasia. *Psychon. Bull. Rev.* **29**, 2096–2107 (2022).

23. Liu, J. *et al.* Ultra-high field fMRI of visual mental imagery in typical imagers and

aphantasic individuals. 2023.06.14.544909 Preprint at

https://doi.org/10.1101/2023.06.14.544909 (2023).

24. Bainbridge, W. A., Pounder, Z., Eardley, A. F. & Baker, C. I. Quantifying aphantasia

through drawing: Those without visual imagery show deficits in object but not spatial

memory. *Cortex* **135**, 159–172 (2021).

25. Ragni, F., Lingnau, A. & Turella, L. Decoding category and familiarity information during

visual imagery. *NeuroImage* **241**, 118428 (2021).

26. Favila, S. E., Lee, H. & Kuhl, B. A. Transforming the Concept of Memory Reactivation.

Trends Neurosci. **43**, 939–950 (2020).

27. Srokova, S., Hill, P. F. & Rugg, M. D. The Retrieval-Related Anterior Shift Is Moderated by

Age and Correlates with Memory Performance. *J. Neurosci.* **42**, 1765–1776 (2022).

28. Bainbridge, W. A. & Baker, C. I. Multidimensional memory topography in the medial

parietal cortex identified from neuroimaging of thousands of daily memory videos. *Nat.*

Commun. **13**, 6508 (2022).

29. Epstein, R. A. & Baker, C. I. Scene Perception in the Human Brain. *Annu. Rev. Vis. Sci.* **5**,

373–397 (2019).

30. Himmer, L., Schönauer, M., Heib, D. P. J., Schabus, M. & Gais, S. Rehearsal initiates

systems memory consolidation, sleep makes it last. *Sci. Adv.* **5**, eaav1695 (2019).

31. Kriegeskorte, N., Mur, M. & Bandettini, P. Representational similarity analysis - connecting

the branches of systems neuroscience. *Front. Syst. Neurosci.* **2**, (2008).

32. Ke, J. *et al.* The neural signatures of ongoing thoughts during rest. (in prep).

33. Szaflarski, J. P., Holland, S. K., Schmithorst, V. J. & Byars, A. W. fMRI study of language

lateralization in children and adults. *Hum. Brain Mapp.* **27**, 202–212 (2006).

34. Weber, S., Christophel, T., Görgen, K., Soch, J. & Haynes, J.-D. Working memory signals in

early visual cortex do not depend on visual imagery. 2023.02.13.528298 Preprint at

https://doi.org/10.1101/2023.02.13.528298 (2023).

35. Dilks, D. D., Julian, J. B., Paunov, A. M. & Kanwisher, N. The Occipital Place Area Is

Causally and Selectively Involved in Scene Perception. *J. Neurosci.* **33**, 1331–1336 (2013).

36. Epstein, R. & Kanwisher, N. A cortical representation of the local visual environment.

Nature **392**, 598–601 (1998).

37. Sekeres, M. J., Winocur, G. & Moscovitch, M. The hippocampus and related neocortical

structures in memory transformation. *Neurosci. Lett.* **680**, 39–53 (2018).

38. Tompary, A. & Davachi, L. Consolidation Promotes the Emergence of Representational

Overlap in the Hippocampus and Medial Prefrontal Cortex. *Neuron* **96**, 228-241.e5 (2017).

39. Hugdahl, K. Hemispheric asymmetry: contributions from brain imaging. *WIREs Cogn. Sci.*

2, 461–478 (2011).

40. *The Asymmetrical Brain*. (MIT Press, Cambridge, Mass, 2003).

41. Blajenkova, O., Kozhevnikov, M. & Motes, M. A. Object-spatial imagery: a new self-report

imagery questionnaire. *Appl. Cogn. Psychol.* **20**, 239–263 (2006).

42. Veale, J. F. Edinburgh Handedness Inventory – Short Form: A revised version based on

confirmatory factor analysis. *Laterality* **19**, 164–177 (2014).

43. Brainard, D. H. The Psychophysics Toolbox. (1997) doi:10.1163/156856897X00357.

44. Fedorenko, E., Hsieh, P.-J., Nieto-Castañón, A., Whitfield-Gabrieli, S. & Kanwisher, N.

New Method for fMRI Investigations of Language: Defining ROIs Functionally in

Individual Subjects. *J. Neurophysiol.* **104**, 1177–1194 (2010).

45. Cox, R. W. AFNI: Software for Analysis and Visualization of Functional Magnetic

Resonance Neuroimages. *Comput. Biomed. Res.* **29**, 162–173 (1996).

46. Kravitz, D. J., Peng, C. S. & Baker, C. I. Real-World Scene Representations in High-Level

Visual Cortex: It's the Spaces More Than the Places. *J. Neurosci.* **31**, 7322–7333 (2011).

47. Vanderwal, T., Kelly, C., Eilbott, J., Mayes, L. C. & Castellanos, F. X. *Inscapes*: A movie

paradigm to improve compliance in functional magnetic resonance imaging. *NeuroImage*

122, 222–232 (2015).

48. Chamberlain, T. A. *et al.* High performers demonstrate greater neural synchrony than low

performers across behavioral domains. *Imaging Neurosci.* **2**, 1–17 (2024).

49. Zhang, Z. & Rosenberg, M. D. Brain network dynamics predict moments of surprise across

contexts. 2023.12.01.569271 Preprint at https://doi.org/10.1101/2023.12.01.569271 (2024).

50. Shen, X., Tokoglu, F., Papademetris, X. & Constable, R. T. Groupwise whole-brain

parcellation from resting-state fMRI data for network node identification. *NeuroImage* **82**,

403–415 (2013).

51. Desmond, J. E. *et al.* Functional MRI measurement of language Lateralization in Wada-

tested patients. *Brain* **118**, 1411–1419 (1995).

52. Binder, J. R. *et al.* Determination of language dominance using functional MRI. *Neurology*

46, 978–984 (1996).

53. Hubrich-Ungureanu, P., Kaemmerer, N., Henn, F. A. & Braus, D. F. Lateralized organization

of the cerebellum in a silent verbal fluency task: a functional magnetic resonance imaging

study in healthy volunteers. *Neurosci. Lett.* **319**, 91–94 (2002).

54. Springer, J. A. *et al.* Language dominance in neurologically normal and epilepsy subjects: A

functional MRI study. *Brain* **122**, 2033–2046 (1999).

55. Deblaere, K. *et al.* MRI language dominance assessment in epilepsy patients at 1.0 T: region

of interest analysis and comparison with intracarotid amytal testing. *Neuroradiology* **46**,

413–420 (2004).

56. Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. Large-scale

automated synthesis of human functional neuroimaging data. *Nat. Methods* **8**, 665–670

(2011).

689