1   Absent, but not glucocorticoid-modulated, corticotropin-releasing hormone (Crh)

2   regulates anxiety-like behaviors in mice

Hannah R. Friedman[1], Lindsey S. Gaston[1], Li F. Chan[2], Joseph A. Majzoub[1, 3]

[1]Division of Endocrinology, Department of Pediatrics, Boston Children's Hospital,

Harvard Medical School, Boston, MA; [2]Centre for Endocrinology, William Harvey

Research Institute, Faculty of Medicine and Dentistry, Queen Mary University of

London, London, United Kingdom

[3]Address all correspondence to: Joseph A. Majzoub, MD, Division of Endocrinology,

Boston Children's Hospital, 300 Longwood Avenue, Boston, MA 02115,

joseph.majzoub@childrens.harvard.edu, ORCID 0000-0002-4983-5143

All authors declare they have no conflicts of interest to disclose

Abstract (248 words)

The hypothalamic-pituitary-adrenal (HPA) axis is a well characterized endocrine

response system. Hypothalamic Crh in the paraventricular nucleus of the hypothalamus

(PVH) initiates HPA axis signaling to cause the release of cortisol (or corticosterone in

rodents) from the adrenal gland. PVH-specific deletion of *Crh* reduces anxiety-like

behaviors in mice. Here we report that manipulation of PVH *Crh* expression in primary

adrenal insufficiency or by dexamethasone (DEX) treatment do not alter mouse anxiety

behaviors. In Experiment 1, we compared wildtype (WT) mice to those with primary

adrenal insufficiency (*Mrap*KO) or global deletion of Crh (*Crh*KO). We analyzed

behaviors using open field (OF) and elevated plus maze (EPM), PVH Crh mRNA

expression by spatial transcriptomics, and plasma ACTH and corticosterone after a 15-

minute restraint test with ELISAs. EPM analysis showed *Crh*KO mice were less anxious

than WT and *Mrap*KO mice, and *Mrap*KO mice had no distinguishing behavioral

phenotype. In Experiment 2, we evaluated HPA axis habituation to chronically elevated

Crh expression by comparing mice treated with 5-8 weeks of DEX with those similarly

treated followed by DEX withdrawal for 1 week. All mice regardless of genotype and

treatment showed no significant behavioral differences. Our findings suggest that

reduced anxiety associated with low *Crh* expression requires extreme deficiency,

perhaps outside of those PVH Crh neurons negatively regulated by glucocorticoids. If

these findings extend to humans, they suggest that increases in Crh expression with

primary adrenal insufficiency, or decreases with exogenous glucocorticoid therapy, may

not alter anxiety behaviors via modulation of Crh expression.

Introduction

The World Health Organization estimates that approximately 300 million people have some form of an anxiety disorder, with this number increasing by 26% during the COVID-19 pandemic.[1] In general, patients living with anxiety disorders have excessive worry and/or fear. Typically, patients with anxiety disorders have significant disturbances in their thinking which often result in dysregulation of their emotions and/or behavior that may consequently impair daily functions.[1] There is a wide spectrum in the severity of symptoms patients may experience, and many factors may contribute to the source of their anxiety.

Stressors are defined as conditions in which there is a threat to homeostasis. This includes perceived or anticipated threats established from prior experience and perceived threats within a current situation.[2] When the body is subjected to these stressors, the stress response incudes autonomic, endocrine, and behavioral components. In the presence of an acute stressor, the autonomic nervous system utilizes both the sympatho-adrenomedullary limb, which secretes epinephrine from the adrenal medulla into the systemic circulation, and the sympatho-neuronal limb, which secretes norepinephrine from sympathetic neurons distributed throughout the body's organs, to elicit a rapid-onset 'fight-or-flight' response.[2-4] These catecholamines work to restore homeostasis by increasing heart rate, blood pressure, plasma glucose, and oxygen consumption within seconds.[2,5]

Endocrine responses to stressors have a longer time course and are mediated by the

hypothalamus, anterior pituitary gland, and adrenal gland, which together constitute the

hypothalamic-pituitary-adrenal (HPA) axis.[2-7] In response to a stressor, corticotropin-

releasing hormone (CRH) is released from parvocellular neurons within the medial

paraventricular nucleus of the hypothalamus (PVH). Acting through its G-protein

coupled CRH receptor-1 (CRHR1), located in corticotrophs of the anterior pituitary

gland, CRH stimulates these cells to increase the synthesis and secretion of

adrenocorticotropin hormone (ACTH). ACTH stimulates zona fasciculata cells of the

adrenal cortex through its G protein-coupled melanocortin-2 receptor (MC2R) to

stimulate the synthesis and release of cortisol (or corticosterone in rodents). In addition

to their effects to maintain vascular, energy and metabolic homeostasis by promoting

vasoconstriction, hepatic gluconeogenesis, proteolysis in muscle, and lipolysis in

adipocytes, glucocorticoids cause negative feedback inhibition on CRH in the PVH and

ACTH in anterior pituitary corticotrophs via glucocorticoid receptors (GR) to restore the

axis to its baseline state.[2,4,6]

The behavioral stress-response has yet to be fully characterized. While CRH plays a

role in mouse anxiogenesis, the biological conditions and mechanisms remain unclear.[8]

[3,9-14] This study aims to expand upon our and others' prior work by characterizing the

relationship between the body's endocrine and behavioral stress responses. We have

previously shown that deletion of hypothalamic Crh using a Cre recombinase[15] driven

by the expression of transgenic *Sim1* (a transcription factor that regulates PVH

development[16,17]), results in both suppression of the HPA stress response and an

anxiolytic behavioral phenotype in mice.[18] This phenotype persisted in mice with *Sim1*

deletion of Crh even when they were administered corticosterone-supplemented

drinking water, suggesting that the reduction in anxiety was due to loss of hypothalamic

Crh rather than the reduction in blood corticosterone concentration.[18] Another group

demonstrated that PVH Crh neurons coordinate rodent behavior in response to stress,

as their metaplastic[19] capability to anticipate[20] and interpret environmental cues[21]

promotes adaptable behavioral decision making.[20,22] Through

photoactivation/inactivation of PVH Crh neurons in mice, this group found a relationship

between PVH Crh activity and the onset of anxiety-like behaviors.[23] They found that

optogenetic stimulation of PVH Crh neurons caused a rapid onset of intense grooming

behavior that rapidly ceased with termination of photostimulation[21]. These studies

demonstrate that modulation of Crh in PVH neurons controls not only the endocrine

response but also aspects of the behavioral response to stressors. This could provide

an explanation for the tight coupling between these two aspects of the stress response,

which are invariably linked following exposure to a stressor. [21]

Our present study examines the impact of more subtle, physiological changes in PVH

Crh on behavioral responses to stressors by modulating PVH Crh via the negative

feedback of glucocorticoids upon PVH Crh expression and comparing these responses

to those in *Crh*KO mice. We have used *Mrap*KO mice as a model of high endogenous

Crh. These animals have a genetic form of severe, primary adrenal insufficiency due to

deletion of *Mrap*, a single-pass transmembrane accessory protein required for signaling

of ACTH via Mc2r, causing markedly elevated PVH *Crh* expression.[24] We have used

mice treated with dexamethasone (DEX) as a model of low endogenous Crh. DEX is a

potent, synthetic glucocorticoid that suppresses HPA axis activity, as it inhibits both PVH

Crh and anterior pituitary ACTH secretion through GR mediation.[25] [26] We administered a

supraphysiological amount of DEX in mouse drinking water to WT, *Crh*KO and *Mrap*KO

mice, and examined the endocrine and behavioral stress responses in these mice

before, during, and after withdrawal from DEX treatment. Surprisingly, we found that

while *Crh*KO mice with global deletion of *Crh* have markedly decreased anxiety-like

behaviors, modulation of endogenous Crh in mice via alteration in glucocorticoid

negative feedback has no effect on behavioral stress responses.

Methods

<u>Animal Husbandry</u>

Adult male mice ages 12-16 weeks were used for all experiments. Adrenal function [27]

including the effect of Crh on adrenal activity [28] is sexually dimorphic in mice, such that

future studies are warranted in female mice. All mice were bred on a C57BL/6J

background and housed in Boston Children's Hospital's animal facility following the

regulations of the Institutional Animal Care and Use Committee, which approved all

experiments. Mice were kept on a 12-hour light/dark cycle (lights on 7 am) in a

temperature and humidity-controlled room (21°C with relative humidity 35-64% set point

+/- 5%). Animals were kept on a standard chow diet *ad libitium* (<u>Prolab IsoPro</u>

<u>RMH3000</u>). *Mrap*KO mice have global deletion of *Mrap* [24] and were bred from an

*Mrap*KO father paired with a heterozygous (glucocorticoid-sufficient) *Mrap*KO mother.

*Crh*KO mice were generated as previously described from *Crh*KO x *Crh*KO pairings. [18]

To assure viability of postnatal mice, when either *Crh*KO or heterozygous *Mrap*KO mice

were visibly pregnant (around E10.5), mothers were switched from standard drinking

water to corticosterone-supplemented drinking water (see below). Mothers and pups

were kept on corticosterone-supplemented water until pups were weaned (P21). Mice

treated with DEX (see below) were immediately weaned onto DEX-supplemented water

(3.6 mcg/mL) and maintained on this treatment for 5-8 weeks.

Corticosterone Treatment

A 15 mg/mL stock solution was prepared by adding 500mg >92% pure corticosterone

(Sigma-Aldrich, C2505-500 mg) to 33 mL pure, non-denatured 200 proof ethanol

(Sigma-Aldrich, 493546) and vortexing until the material was completely dissolved. The

final stock solution was aliquoted and stored at -20°C until ready for use. Before use,

the aliquots were thawed and vortexed until all corticosterone was solubilized and then

added into mouse drinking water at a final concentration of 30 mcg/mL, as previously

described.[29]

Dexamethasone Treatment

A 10 mg/mL DEX stock solution was prepared by mixing 100 mg water-soluble DEX

(Sigma-Aldrich, D2915-100mg) with 10 mL of tap water. The material was vortexed until

completely dissolved. The stock solution was aliquoted and stored at -20°C until ready

for use. Before use, the aliquots were thawed and mixed by pipetting up and down and

then added to mouse drinking water at a final concentration of 3.6 mcg/mL.

<u>Behavioral Experiments</u>

Open Field

Mice were transferred to Boston Children's Hospital's Neurobehavioral Core's testing

facilities (Day 0) and single-housed one week prior to the first testing date to allow the

mice to acclimate to their new environment. All experiments were performed starting at

9 am. Open field (OF) experiments were performed (Day 7) in a 42 cm x 42 cm square

box and scored using Kinder Scientific Motor Monitor II tracking software. Mice were

placed at a 6 o'clock position facing towards the wall at the start time. Each run lasted

for 15 minutes. Between each run, each OF box was wiped down with

disinfecting/deodorizing wipes. Each mouse was returned to their home cage after

testing.

Elevated Plus Maze

Elevated plus maze (EPM) experiments were performed (Day 8) on a platform raised 74

cm from the floor. Each arm of the platform was 77 cm in total length. Mice were placed

at the center opening so that they were facing the open arms at the start time. Each trial

ran for 6 minutes, and the data were automatically scored using Ethovision XT software

(version 17.0). EPM heatmaps represent the average amount of time spent by location

of each mouse. The average time and location data were generated by the same

Ethovision XT software. In Fig. 2, the horizontal arms, denoted with "O", represent the

open arms of the platform, and the vertical arms, denoted by "C", represent the closed

arms on the platform. The center area of the EPM is not enclosed by walls, so it is

considered an open area of the maze. Between each run, the maze was wiped down

with disinfecting/deodorizing wipes. Each mouse was returned to its home cage after

testing. For both OF and EPM experiments, a lamp with an 8.5 W (800 Lumens) light

bulb was pointed up at the ceiling for low level ambient lighting.

Restraint Test and retro-orbital phlebotomy

On Day 9 of Experiments 1 and 2, one day after completing the EPM experiment, mice

were restrained in 50cc tubes that previously had the end of the tube cut off for

ventilation. After 15 minutes of restraint, mice were subjected to retro-orbital phlebotomy

using EDTA-coated capillary tubes and returned to their home cage. Blood samples

were immediately placed on ice and then centrifuged at 4°C for 10 minutes at 1500 x g.

The plasma was immediately aspirated and stored at -80°C until hormone analysis. A

separate group of WT, *Crh*KO heterozygotes, and *Crh*KO homozygotes was restrained

for 2 h to verify *Crh* mRNA expression in the *Crh*KO model used in this study that was

created by recombineering[18] and different from our initial *Crh*KO model.[28]

In Experiment 2, due to unavoidable experiment interference, 1 WT untreated animal

was restrained for 28 min instead of 15 min. Therefore, data from this animal were

removed from both plasma ACTH and corticosterone concentration evaluations. No

interference occurred during the prior behavioral experiments, so those data from this

animal were retained in OF and EPM analyses.

<u>Hormone Assays</u>

Plasma was slowly thawed on ice. Corticosterone concentration was measured using a

mouse/rat corticosterone ELISA kit (ALPCO; 55-CORMS-E01) as directed by the

manufacturer. The lower limit of detection is 0.061 mcg/dL. Animals which had

undetectable plasma corticosterone levels were recorded as having a value halfway

between 0 mcg/dL and the assay's lower limit of detection. ACTH levels were measured

using ACTH ELISA (ALPCO; 21-ACTH-E01) following the manufacturer's protocol. The

lower limit of detection is 0.22 pg/mL. Additional ACTH 0 calibrator (ALPCO; 11-C-270-

10-G25) was used to dilute plasma as necessary. Animals that had undetectable

plasma ACTH levels were recorded as having a value halfway between 0 pg/mL and the

assay's lower limit of detection.

In situ Hybridization

Experimental animals were euthanized (Day 10) by rapid decapitation without

anesthesia. Brains were removed and fixed in 10% formalin overnight and then

transferred to 70% ethanol for histology. Paraffin embedding was performed by either

the Histology Core Facility at Beth Israel Deaconess Medical Center or at Boston

Children's Hospital's Histology Department. Paraffin blocks were sectioned and H&E

stained by Boston Children's Hospital's Pathology Department. Three 5μM-thick coronal

brain sections were sectioned onto each slide, and every other slide was H&E stained.

We surveyed H&E slides under a Nikon E800 microscope to identify adjacent unstained

slides that contained PVH. Four unstained slides that were flanked by H&E-identified,

PVH-containing brain sections were selected for *Crh* mRNA hybridization using

RNAscope (ACDBio). The middle brain section on each of the 4 slides was stained for

Crh mRNA using a mouse *Crh* probe (ACD #316091) and 2.5 HD RED detection kit

(ACD, #316091) following manufacturer's instructions. Similar brain sections from

*Crh*KO mice of all treatment conditions for were analyzed for *Crh* mRNA as a negative

control.

For an individual biological sample, the brain section with the highest *Crh* mRNA

expression among the 4 stained sections was analyzed densitometrically (see below).

These selected slides were then compared across their respective biological replicates.

If there was variation in *Crh* expression among biological replicates, the selected slides

were ranked from least to greatest *Crh* expression, and the median image was chosen

as the final representative image. Semi-quantification of RNAscope images was

performed following the RNAscope manufacturer's protocol for FIJI (Fig. S1,

Supplemental Materials[30]). The area of the PVH was determined by first drawing two

triangles that were large enough to encompass the *Crh* signal on all slides (Fig. S1A,

Supplemental Materials[30]). These same triangles were then transposed onto each

image. The areas of the two transposed triangles were measured, and the average area

of these triangles was calculated. *Crh* expression was quantified by dividing the area of

the *Crh* densitometric signal by the average area of the PVH triangles.

<u>Genotyping</u> (see Supplemental Materials[30])

<u>Statistical Analyses</u>

Statistical analyses of behavioral experiments were performed using Prism GraphPad

10. Means were evaluated using One-way ANOVA multiple comparison testing with the

Bonferroni method and compared to untreated WT values. In Experiment 2, Figure 3G

used the same WT untreated values from Experiment 1 that were shown in Figure 1D.

Asterisks on graphs denote significance (*≤0.05, **≤0.001, and ***≤0.0001). Error bars

on graphs represent mean + SEM. Due to multiple undetectable plasma ACTH

concentrations in DEX treated animals, Gaussian distribution cannot be assumed.

Therefore, one-way ANOVA analysis is unable to test for significance compared to WT

untreated animals.

Results

<u>*Crh*KO mice lack hypothalamic *Crh* mRNA and have low basal and stress-induced</u>

<u>CORT responses</u>

The *Crh*KO model used in this study was created by recombineering[18], whereas our

previously described *CrhKO* model was developed with restriction fragment ligation and

retained a *Pgk*neo cassete in the deleted *Crh* locus[28]. We thus first sought to phenotype

this novel, recombineered model by assaying hypothalamic *Crh* mRNA expression as

well as plasma corticosterone and ACTH levels both in the basal and stressed state in

Crh WT, *Crh*KO heterozygous, and *Crh*KO homozygous animals. Stress was induced

with a 2h physical restraint. Overall, *Crh* mRNA expression significantly differed by both

genotype and treatment condition using 2-way ANOVA (Fig S1; p=0.007). *Crh* mRNA

increased following 2h restraint in both WT and *Crh*Het animals, and expression was

higher in WT vs. *Crh*HET animals under both conditions. As expected, *Crh* mRNA was

absent in both basal and stressed *Crh*KO mice (Fig S1, Supplemental Materials[30]).

Corticosterone levels were similarly higher after 2h restraint compared to the basal state

in WT and *Crh*HET but not *CrhKO* mice ($P<0.0001$), though there was no significant

effect of genotype nor treatment condition on plasma ACTH levels ($P=0.44$, Fig S2,

Supplemental Materials[30]).

<u>Influence of endogenous *Crh* and glucocorticoids on behavioral and endocrine</u>

<u>responses to stress in WT, *Mrap*KO and *Crh*KO mice. (Experiment 1)</u>

In Experiment 1, we tested whether *Crh* expression links the endocrine and behavioral

stress responses. First, we exposed mice to a 15-minute open field (OF) trial followed

by a 6-minute elevated plus maze (EPM) trial 24 hours after the OF test. OF total

locomotor activity was not different among WT, *Mrap*KO and *Crh*KO mice (WT vs.

*Mrap*KO $P>0.999$; WT vs. *Crh*KO $P>0.999$), indicating there were no baseline locomotor

abnormalities in the mutant genotypes (Fig. S3A, Supplemental Materials[30]). Further,

*Mrap*KO and *Crh*KO mice were not different from WT mice in total distance traveled in

the OF center (WT vs. *Mrap*KO $P=0.286$; WT vs. *Crh*KO $P>0.999$), time spent in the OF

center (WT vs. *Mrap*KO $P=0.0719$; WT vs. *Crh*KO $P=0.836$), and number of entries into

the OF center (WT vs. *Mrap*KO $P=0.462$; WT vs. *Crh*KO $P>0.999$) (Figs. S3B-D,

Supplemental Materials[30]). These three measures are indicators of anxiety, suggesting

that neither absent (*Crh*KO) nor elevated (*Mrap*KO) *Crh* expression affects these

aspects of mouse anxiety-like behavior.

The morning following the OF trial, we exposed mice to a 6-minute EPM trial. Increased

time spent and number of entries into the EPM open areas indicate decreased anxiety-

like behavior in mice. An arial view heatmap, displaying genotype-averaged position x

time during 6-minute trials, showed that *Crh*KO mice (Fig. 1C, D), compared to WT mice

(Fig. 1A, D), spent more time in the open areas of the maze (*Crh*KO 145.2±10.8 s vs.

WT 69.5±7.6 s; P<0.0001). *Crh*KO mice also entered the open areas more frequently

than WT mice during the 6-minute test (*Crh*KO 49.4±4.7 entries vs. WT 29.7±2.5

entries; P=0.0015) (Fig. 1E). Interestingly, *Mrap*KO mice, with elevated PVH *Crh*

expression, scored comparably to WT mice in time spent in open areas (*Mrap*KO

99.4±9.6 s vs. WT 69.5±7.6 s; P=0.0595), with a nonsignificant trend towards more time

in these areas (Fig. 1B, D). Further, the number of entries into the open areas was not

different between *Mrap*KO and WT mice (*Mrap*KO 35.1±3.2 entries vs. WT 29.7±2.5

entries; P=0.541) (Fig. 1E). An increase in latency to enter the open areas of the EPM is

associated with increased anxiety-like behavior. Both *Mrap*KO mice (23.6±10.0 s;

P=0.911) and *Crh*KO mice (11.9±4.7 s; P>0.999) scored comparably to WT mice

(15.0±6.1 s) (Fig. S3E, Supplemental Materials[30] in this metric.

24 h after completing the EPM trial in Experiment 1, we characterized the HPA axis

status in these same WT, *Mrap*KO and *Crh*KO mice after 15 minutes of restraint stress

by measuring PVH *Crh* mRNA expression by RNAscope and plasma corticosterone

concentrations. Following this stressor, *Mrap*KO mice tended towards increased

expression of PVH *Crh* mRNA compared to WT (WT 5.8±0.2 vs. *Mrap*KO 16.4±2.6

arbitrary units; P=0.052) while no *Crh* transcript was detected in *Crh*KO mice (Figs. 2A-

D). After 15 minutes of restraint, WT mice plasma corticosterone was 11.6±1.5 mcg/dL,

whereas it was undetectable in *Mrap*KO mice (P<0.0001) (Fig. 2E). As we have

previously found[18], *Crh*KO had low plasma corticosterone concentrations following

restraint (0.59±0.24 mcg/dL compared to WT, P<0.0001) (Fig. 2E).

<u>Dexamethasone reduces PVH *Crh* expression and the HPA axis response to stress,</u>

<u>and its subsequent withdrawal restores PVH Crh expression and HPA axis</u>

<u>responsiveness, but neither treatment affects anxiety-like behaviors. (Experiment 2)</u>

In Experiment 2, we asked if the lack of increased anxiety-like behaviors in *Mrap*KO

mice in Experiment 1 was due to habituation to chronically elevated *Crh* expression. To

test this, we disrupted HPA axis signaling by administration of dexamethasone (DEX)-

supplemented drinking water. We compared two treatment groups to the WT-untreated

animals studied in Experiment 1, (Fig. 2A, D). The first group consisted of WT, *Mrap*KO

and *Crh*KO mice treated with DEX for 5-8 weeks. The second treatment group

consisted of WT, *Mrap*KO and *Crh*KO mice treated with DEX for 5-8 weeks and then

withdrawn from treatment for 1 week.

325
We measured mouse behavior using the same 15-minute OF and 6-minute EPM tests

as in Experiment 1. Unlike in Experiment 1, where untreated *Crh*KO mice compared to

WT mice had reduced anxiety-like behaviors in OF and EPM tests, following either DEX

treatment or its subsequent withdrawal, *Crh*KO mice behaved no differently than

untreated WT mice in time spent in EPM open areas (*Crh*KO-DEX 73.8± 16.9 s vs, WT-

untreated 100.8± 9.1 s, P=0.616; *Crh*KO-DEX withdrawn 95.8± 13.0 s vs. WT-untreated

100.8± 9.1 s, P>0.999) (Fig. 3A) or entry frequency in EPM open areas (*Crh*KO-DEX

39.7±8.4 entries vs. WT-untreated 40.9±2.4 entries, P>0.999; *Crh*KO DEX-withdrawn

41.8±6.8 entries vs, WT-untreated 40.9±2.4 entries, P>0.999) (Fig. 3B). Similarly,

behavior of *Mrap*KO mice in both treatment groups (DEX-treated, and DEX-withdrawn)

was not different from that of untreated WT (Fig. 3). All genotypes in both treatment

groups (DEX-treated or DEX-withdrawn) had no difference in OF total locomotor activity

compared to WT untreated mice, indicating DEX treatment does not interfere with

overall mouse locomotion (Fig. S4A, Supplemental Materials[30]). Distance traveled, time

spent, and number of entries in the OF center were also not different among all

genotypes and treatment groups compared to WT untreated animals (Figs. S4B-D,

Supplemental Materials[30]). EPM analysis of latency to open areas also found no

significant differences in animals on DEX treatment, nor in animals who were DEX-

withdrawn compared to WT-untreated animals (Fig. S4E, Supplemental Materials[30]).

Twenty-four hours after completion of behavioral testing, we analyzed the impact of

DEX treatment and its withdrawal on PVH *Crh* mRNA expression and HPA axis

hormones after a 15-minute restraint. Compared with *Crh* mRNA expression in

untreated WT mice (5.8±0.2 arbitrary units, Figs. 2A, D), DEX-treated WT (Fig. 4A, G,

1.6±0.1 arbitrary units) and *Mrap*KO (Figs. 4B, G, 1.6±1.4 arbitrary units) were lower.

Following 1 week of DEX withdrawal PVH *Crh* was higher in both WT (Fig. 4D, G,

6.6±0.4 arbitrary units) and *Mrap*KO (Figs. 4E, G, 7.7±4.3 arbitrary units) mice. As

expected, *Crh*KO mice had undetectable *Crh* mRNA irrespective of DEX treatment

status (Figs. 4C, F, G). None of the differences in PVH *Crh* were significant between WT

and MrapKO genotypes or DEX treatment status, possibly because of the small number

of animals studied (Fig 4G; P=0.4948).

After 15 minutes of restraint stress in WT and *Mrap*KO DEX-treated animals, plasma

ACTH was undetectable compared to WT mice not given DEX (410.1±33.9 pg/mL) (Fig.

5A). Plasma ACTH was also very low in DEX-treated *Crh*KO mice (31.8±26.9 pg/mL).

One week after DEX withdrawal, WT mice had plasma ACTH concentrations not

different from those of WT-untreated mice (WT-withdrawn 833.6±230.6 pg/mL vs. WT-

untreated 410.1±33.9 pg/mL; P>0.999) (Fig. 5A), whereas *Mrap*KO mice had

significantly elevated plasma ACTH (2399±841.3 pg/mL; P<0.0001) (Fig. 5A). Notably,

two DEX-withdrawn *Mrap*KO animals had low (357.9 pcg/mL) or undetectable plasma

ACTH after the restraint test. When analyzing the behavior of these two animals, they

were not identified as outliers when using the Grubbs' outlier test (alpha=0.05). Plasma

ACTH remained low in DEX-withdrawn *Crh*KO mice (14.6±10.6 pg/mL) (Fig. 5A).

After 15 minutes of restraint stress, plasma corticosterone in WT-untreated mice was

10.9±0.7 mcg/dL (Fig. 5B), similar to that observed in Experiment 1 (Fig. 2E) and was

markedly lower in DEX-treated WT mice (0.2±0.1 mcg/dL; P<0.0001) (Fig. 5B). Plasma

corticosterone was also very low in DEX-treated *Mrap*KO mice (0.8±0.3 mcg/dL;

P<0.0001) and *Crh*KO mice (1.0±0.7 mcg/dL; P<0.0001) (Fig. 4B), with levels similar to

those found following restraint without DEX treatment (Fig.1E). Although WT mice

withdrawn from DEX treatment had increased plasma ACTH compared to restrained WT

mice that had not received DEX (Fig. 4A), corticosterone concentrations after restraint

in DEX-withdrawn compared to DEX-untreated restrained WT mice were markedly

lower (WT-withdrawn 2.3±0.6 mcg/dL vs. WT-untreated 10.9±0.7 mcg/dL; P<0.0001)

(Fig. 4B).

Discussion

Many studies have explored the role of Crh and its receptor, Crhr1, in anxiety behavior

regulation in rodents, [9,31-35], including some that have found administration of Crhr1

antagonists results in a reduction of anxiety-like behavioral phenotypes,[10,36], and

another which demonstrated that a Crhr1 antagonist reversed an anxiogenic phenotype

in mice with Crh-neuron specific deletion of GABA.[37] Based on these preclinical data,

several pharmaceutical companies developed small molecule CRH receptor 1

antagonists, which unfortunately all failed to reduce anxiety in multiple human clinical

trials. [38] We previously found that selective deletion of murine *Crh* in the brain regions of

Sim1 expression, including in the PVH, reduces both stressor-induced anxiety-like

behaviors and endocrine responses of the HPA axis.[18] Our findings are consistent with

the fact that behavioral and endocrine responses to stressors are usually temporally

associated[21], and suggest that CRH, at least in part, may mediate this linkage. [23]

In our prior studies of selective deletion of PVH *Crh, Sim1Crh*KO mice were deficient in

both hypothalamic Crh as well as in systemic corticosterone, the major mouse

glucocorticoid. Thus, their reduced anxiety could have been due to the reduction in

either hormone. To evaluate this further we supplemented those mice with short-term

systemic glucocorticoid via their drinking water and found that they still exhibited

anxiolytic behavior.[18] We therefore concluded that this behavior was due to

hypothalamic Crh, rather than glucocorticoid, deficiency. We undertook the current

studies to further examine the impact of manipulation of endogenous Crh via changes in

negative feedback by glucocorticoids in mediating stressor-related behaviors. To that

end, we have compared wildtype mice to those with global deletion of either *Crh*

(*Crh*KO mice) or *Mrap* (*Mrap*KO mice). Mrap is necessary for ACTH function via

signaling through the adrenal ACTH/Mc2r receptor. [24] Humans with mutations in *MRAP*

have a severe form of primary adrenal insufficiency, Familial Glucocorticoid Deficiency

Type 2 (FGD2).[39] Mice with global deletion of *Mrap* have a similar phenotype, with

severe glucocorticoid deficiency causing secondary elevation in hypothalamic Crh and

pituitary ACTH.[24] We reasoned that since *Mrap*KO mice have both very low systemic

glucocorticoid concentrations and very high hypothalamic PVH *Crh* expression (Fig. 2),

they would be useful in understanding whether modulation of one or both of these

hormones is involved in stressor-related anxiety behaviors.

In this study, *Crh*KO mice in Experiment 1 exhibited a reduction in anxiety-like

behaviors, as they spent more time in the EPM open areas (Fig. 2) and entered the

open areas more frequently compared to WT mice (Fig. 2). Their anxiolytic behavioral

phenotype was not due to differences in locomotor activity between null mutants and

WT mice as determined by both groups' comparable total distance traveled during the

15-minute OF test (Fig. S3, Supplemental Materials[30]). Thus, the total absence of Crh

expression reduces anxiety-like behaviors in mice. We had previously found that a

different *Crh*KO mouse line had no anxiolytic phenotype[9], which was confirmed by

another research group using this same mouse. [40] However, the mouse model used for

those studies[28] was different from the *Crh*KO mice we used in the present study. The

cloned genomic DNA in the original mouse model was obtained from a Balb/c DNA

library and resulted in a *Crh*KO mouse on a mixed 129SV/C57BL6 background. The

gene-targeting vector replaced the coding region of *Crh* with a neomycin resistance

gene that was continuously expressed from a *Pgk* promoter. The *Crh*KO mouse used in

the current study was created by breeding a mouse with loxP sites flanking *Crh* exon 2

with an *EIIA*Cre mouse, resulting in germline deletion of the entire *Crh* exon 2.[18]

Further, we have studied this mouse on a pure C57BL6 background instead of on a

mixed 129SV/C57BL6 background.

To reverify our new *Crh*KO mouse strain, we used *in situ* transcriptomics to

demonstrate our ability to manipulate *Crh* expression in WT and Crh-deficient mice, and

affirm our ability to accurately locate and measure PVH *Crh* mRNA. We compared PVH

Crh expression in WT, *Crh*HET, and *Crh*KO animals in their basal state (Fig. S1,

Supplemental Materials[30]) and after a 2 h restraint test (Fig. S2, Supplemental

Materials[30]). WT animals had the highest basal PVH *Crh* expression which increased

after restraint. *Crh*HET animals had decreased PVH *Crh* expression compared to their

respective WT treatment controls and as expected, *Crh*KO animals had no PVH *Crh*

expression detected in either. Though WT mice trended towards higher plasma ACTH

than *Crh*HET, there was no significant difference among genotypes and treatment

groups after 2 h of restraint, possibly because the rise in corticosterone during the 2 h

restraint period (Fig. S2, Supplemental Materials[30]) inhibited ACTH production and/or

secretion.

The hypothalamus is the central regulator of the body's endocrine response to an

external stressor, as it initiates the HPA axis signaling cascade with Crh. Within the

hypothalamus, the highest density of Crh-expressing neurons is located in the PVH,[10 41]

and PVH Crh is activated in the presence of multiple external stressors. In one study,

mice were exposed to physical (ether), physiological (high salt), psychological

(restraint), immune (lipopolysaccharide), and innate (predator odor) stressors. Out of

the 95 brain regions that were examined, the PVH was 1 of 12 areas of Crh expression

that had significant increase in Fos expression compared to controls when exposed to

all 5 stressors.[41] The crucial endocrine role of PVH Crh, the dense population of Crh-

expressing cells within the PVH, the selective activation of PVH in response to a variety

of stressors neurons, and numerous studies that demonstrate the behavioral

consequences when PVH Crh is activated or inactivated[18 22], indicate that PVH Crh has

significant importance in rodent stress-response regulation.

Since *Crh*KO mice, with global deletion of Crh, have reduced anxiety-like behaviors, we

hypothesized that mice with increased Crh would display an increase in anxiety-like

behaviors. However, *Mrap*KO mice, with high PVH Crh expression due to primary

adrenal insufficiency (Fig. 2), have normal behavioral responses to stressors (Fig. 1). In

fact, analysis of time spent in EPM open areas showed *Mrap*KO mice had a non-

significant decrease in anxiety-like behaviors compared to WT mice, spending more

time in the open areas. That this might be due to the glucocorticoid deficiency in

*Mrap*KO mice is supported by a study that found that WT mice given corticosterone-

supplemented drinking water for 4 weeks developed an anxiogenic phenotype.[42] If an

increase in circulating glucocorticoids is an anxiogenic factor, it is reasonable to

question if the converse is true. *Mrap*KO mice have undetectable plasma corticosterone

which raises the possibility that despite high hypothalamic Crh, their concomitant low

glucocorticoid state could result in reduced anxiogenic behaviors, such that the two

opposing influences of high Crh and low glucocorticoid in the *Mrap*KO mouse model

counterbalance each other, leading to a normal behavioral phenotype.

The amygdala communicates with the hypothalamus to interpret perceived

threats.[13,14,37,43,44] Activation of Crh in central nucleus of the amygdala (CeA) neurons

using an AAV-DREADD system increased anxiety-like behavior in mice.[14] Rats with CeA

Crh over expression via stereotaxic viral injection have increased anxiety-like behaviors

when evaluated by EPM but not when analyzed with OF. [43] Unlike PVH Crh, Crh in the

central nucleus of the amygdala (CeA) is stimulated by GC secretion.[44] The low GC

state in *Mrap*KO mice may cause reduced Crh expression in the CeA, leading to a

reduction in anxiety that may counterbalance the anxiogenic effect of elevated PVH Crh.

Consistent with our behavioral results in *Mrap*KO mice, another mouse model with high

PVH Crh but also high systemic corticosterone did not have a distinct behavioral

phenotype.[45] Those investigators deleted exon 3 of *Nr3c1*, which encodes the

glucocorticoid receptor (GR) within the hypothalamus using *Sim1*Cre mice.

Homozygous *Nr3c1*FLOX::*Sim1*Cre+ mice had elevated PVH and CeA Crh expression

and elevated systemic corticosterone as a consequence of the PVH failing to

appropriately respond to GR-mediated glucocorticoid negative feedback. CeA GR

remained intact, as the *Sim1* field does not extend to the CeA, resulting in high systemic

corticosterone stimulating CeA Crh.[45].The absence of increased anxiety-like behaviors

in *Mrap*KO (low corticosterone) and in *Nr3c1*FLOX::*Sim1*Cre+ (high corticosterone)

mouse models suggests that glucocorticoids may make minimal contributions to the

manifestation of anxiety-like behaviors in mice.

It is also possible that the HPA axis in *Mrap*KO mice has somehow undergone

physiologic and/or behavioral adaptations to chronically elevated PVH Crh and/or

undetectable corticosterone production. A properly functioning stress response system

should adapt to learned, recurring internal and external stimuli. [46] Some researchers

propose neural network habituation as a mechanism for this adaptation, where repeat

exposure to a stimulus results in a reduction of stimulus-response pathway

transmission.[47] Anxiety disorders may arise due to perceived threats or disruption in

these brain signaling pathways. If a recurring threat or chronic neuronal dysregulation,

such as elevated PVH Crh, is then perceived as non-threatening, it could lead to a

decrease or cessation of the stress response and therefore any subsequent changes in

behavior.[13] Here we have shown how differing levels of PVH Crh or circulating

glucocorticoids do not have a strong relation to anxiety-like behaviors in mice. Though

inconclusive, these results together may have broader implications on how investigators

should define 'stress' in rodents, as many studies use plasma corticosterone

concentrations as the sole indicator of the stress response.

In the current study, we tested the hypothesis that the lack of increased anxiety-like

behaviors in *Mrap*KO mice was the result of HPA axis adaptations. We studied *Mrap*KO

mice that had PVH Crh suppressed by several weeks of DEX treatment, as well as

*Mrap*KO mice that had DEX treatment discontinued 1 week prior to behavioral testing.

Under these conditions, the endocrine axes of all three genotypes (WT, *Crh*KO, and

*Mrap*KO) behaved as anticipated (Figs. 4, 5): *Crh* mRNA expression was suppressed

by DEX treatment in WT and *Mrap*KO mice, and recovered 1 week after discontinuation

of DEX. Although we did not see total supp ression of PVH *Crh* in mice on DEX

treatment, we had previously found that a 70-80% reduction in PVH *Crh* mRNA (by *in*

situ hybridization) was associated with an anxiolytic behavioral phenotype in mice.[18]

Restraint-induced plasma ACTH was suppressed in all three genotypes by DEX

treatment and recovered above baseline one week after DEX was stopped. Plasma

corticosterone was also suppressed during DEX treatment in WT and *Crh*KO and was

undetectable in *Mrap*KO, as expected. [24] One week after DEX withdrawal, WT

corticosterone had not recovered to normal, untreated levels, despite simultaneous

ACTH concentrations being above normal.

Both during DEX treatment and 1 week following its withdrawal, none of the three

genotypes exhibited any changes in anxiety-like behaviors compared to untreated WT

mice (Figs. 3, S4). By testing mice one week after termination of DEX treatment, we had

hoped to study them during the acute phase of the increase in PVH Crh. However, it is

possible that by the end of the 1-week withdrawal period, these mice had already

adapted to their prior, pre-treatment neuroendocrine status, and thus negated our

attempt to avoid habituation. It is also possible that global genetic deletion of *Crh* and

glucocorticoid-mediated suppression of PVH Crh (and stimulation of CeA Crh) that

results in different behavioral outcomes Thus, we have found a reduction in anxiety-like

behavior only in mice with genetic manipulation of *Crh* using *Sim1*-Cre in our prior

study[18] and global *Crh* deletion in the present study. In contrast, glucocorticoid-

dependent manipulation of Crh expression did not result in any observable behavioral

phenotype whether Crh expression was increased or decreased. Our finding that

glucocorticoid-induced suppression of PVH Crh is not anxiolytic makes physiological

sense—otherwise feedback inhibition of Crh by prior activation of glucocorticoid

secretion by a stressor might terminate anxiety behaviors prematurely in response to

that stressor. The lack of anxiogenesis we have found in *Mrap*KO mice with elevated

PVH Crh requires further exploration.

In our prior studies of anxiolytic behavior in *Sim1CreCrhKO* mice, *Crh* was deleted in

~70% of neurons within the *Sim1* field. [18] Although the strongest activity of *Sim1*Cre is in

the PVH, other brain structures that express Crh also express *Sim1* including the

anterior/lateral hypothalamic areas and the bed nucleus of the stria terminalis (BNST)

[15], but not the CeA.[18,45] Glucocorticoid modulates only a subset of brain neurons that

express Crh[48], so it is possible that glucocorticoid insufficiency in *Mrap*KO mice or

excess in DEX-treated mice does not affect all Crh-expressing neurons involved in

behavior.

In conclusion, we have found that complete deficiency of Crh is associated with

decreased anxiety-like behaviors in mice. These results are consistent with other

groups' work[22], which establishes Crh as a link between the endocrine and behavioral

responses to stress in rodents. The studies which document significant behavioral

impact of endogenous Crh, including our own, utilize profound technical manipulations

of *Crh* expression (e.g. global deletion, PVH-specific deletion [18], optogenetics) that may

have forced extreme behavioral outcomes that would otherwise not occur in nature. In

our present study, we manipulated *Crh* expression in a way that mimics what could

occur in disease (elevation of PVH Crh in primary adrenal insufficiency) or during clinical

treatment (PVH Crh suppression during and after glucocorticoid treatment). Although we

observed no changes in anxiety-like behaviors following modulation of *Crh* expression

by glucocorticoids, it may still be possible that PVH Crh links the endocrine and

behavioral responses to stress but in a more complex pathway than we initially

hypothesized. PVH Crh along with other limbic regions of the brain that express Crh,

may work together to interpret a perceived threat and then produce a behavioral

response—explaining why we see a reduction in anxiety when Crh is absent from all

brain regions in *Crh*KO mice. Humans with primary adrenal insufficiency[49] or secondary

adrenal insufficiency following exposure to long-term glucocorticoid [50] experience

increased anxiety. Further characterization of this system could aid in the development

of anxiety treatments and improve the quality of life for patients such as these who

experience anxiety as a consequence of disorders of their adrenal axis.

Acknowledgements

We thank Rong Zhang for helpful discussions. Support was provided by NIDDK

F32DK131795-02, a Pediatric Endocrine Society Clinical Scholar Award, and a Boston

Children's Hospital Office of Faculty Development/Basic & Clinical Translational

Research Executive Committees Faculty Career Development Fellowship (LSG); the

Thomas Morgan Rotch Harvard Medical School endowment (JAM); and the Medical

Research Council UK and Academy of Medical Sciences fellowship grant (G0802796),

a Wellcome Trust grant (217543/Z/19/Z), European Union's Horizon 2020 research and

innovation programme under grant agreement 848077, Barts Charity grant (G-002162)

and BBSRC (BB/W018276/1) (LFC). This work was also supported by NICHD IDDRC

grant U54 HD090255 at Boston Children's Hospital.

Supplemental Materials

Supplemental materials (Figures S1-S4 and genotyping methods[30]) are available at:

References

1.   World Health Organization. Mental Disorders. Updated June 8, 2022. Accessed

October 2, 2023. https://www.who.int/news-room/fact-sheets/detail/mental-disorders

2.   Deussing JM, Chen A. The Corticotropin-Releasing Factor Family: Physiology of

the Stress Response. *Physiol Rev*. Oct 1 2018;98(4):2225-2286.

doi:10.1152/physrev.00042.2017

3.   Naughton M, Dinan TG, Scott LV. Chapter 5 - Corticotropin-releasing hormone

and the hypothalamic–pituitary–adrenal axis in psychiatric disease. In: Fliers E,

Korbonits M, Romijn JA, eds. *Handbook of Clinical Neurology*. Elsevier; 2014:69-91.

4.   Leistner C, Menke A. Hypothalamic-pituitary-adrenal axis and stress. *Handb Clin*

Neurol. 2020;175:55-64. doi:10.1016/B978-0-444-64123-6.00004-7

5.   Ulrich-Lai YM, Herman JP. Neural regulation of endocrine and autonomic stress

responses. *Nat Rev Neurosci*. Jun 2009;10(6):397-409. doi:10.1038/nrn2647

6.   Jacobson L. Hypothalamic-pituitary-adrenocortical axis regulation. *Endocrinol*

Metab Clin North Am. Jun 2005;34(2):271-92, vii. doi:10.1016/j.ecl.2005.01.003

7.   Sperling M, Majzoub JA, Menon RK, Stratakis CA, Elsevier. *Sperling pediatric*

endocrinology. Fifth edition. ed. Elsevier; 2021.

8.   Sutton RE, Koob GF, Le Moal M, Rivier J, Vale W. Corticotropin releasing factor

produces behavioural activation in rats. *Nature*. May 27 1982;297(5864):331-3.

doi:10.1038/297331a0

9.   Weninger SC, Dunn AJ, Muglia LJ, Dikkes P, Miczek KA, Swiergiel AH, Berridge

CW, Majzoub JA. Stress-induced behaviors require the corticotropin-releasing hormone

(CRH) receptor, but not CRH. *Proc Natl Acad Sci U S A*. Jul 6 1999;96(14):8283-8.

doi:10.1073/pnas.96.14.8283

10.   Smagin GN, Heinrichs SC, Dunn AJ. The role of CRH in behavioral responses to

stress. *Peptides*. May 2001;22(5):713-24. doi:10.1016/s0196-9781(01)00384-9

11.   Bale TL, Vale WW. CRF and CRF receptors: role in stress responsivity and other

behaviors. *Annu Rev Pharmacol Toxicol*. 2004;44:525-57.

doi:10.1146/annurev.pharmtox.44.101802.121410

12.   Mathew SJ, Price RB, Charney DS. Recent advances in the neurobiology of

anxiety disorders: implications for novel therapeutics. *Am J Med Genet C Semin Med*

Genet. May 15 2008;148C(2):89-98. doi:10.1002/ajmg.c.30172

13.   Calhoon GG, Tye KM. Resolving the neural circuits of anxiety. *Nat Neurosci*. Oct

2015;18(10):1394-404. doi:10.1038/nn.4101

14.   Paretkar T, Dimitrov E. The Central Amygdala Corticotropin-releasing hormone

(CRH) Neurons Modulation of Anxiety-like Behavior and Hippocampus-dependent

Memory in Mice. *Neuroscience*. Oct 15 2018;390:187-197.

doi:10.1016/j.neuroscience.2018.08.019

15.   Balthasar N, Dalgaard LT, Lee CE, Yu J, Funahashi H, Williams T, Ferreira M,

Tang V, McGovern RA, Kenny CD, Christiansen LM, Edelstein E, Choi B, Boss O,

Aschkenasi C, Zhang CY, Mountjoy K, Kishi T, Elmquist JK, Lowell BB. Divergence of

melanocortin pathways in the control of food intake and energy expenditure. *Cell*. Nov 4

2005;123(3):493-505. doi:10.1016/j.cell.2005.08.035

16.   Fan CM, Kuwana E, Bulfone A, Fletcher CF, Copeland NG, Jenkins NA, Crews

S, Martinez S, Puelles L, Rubenstein JL, Tessier-Lavigne M. Expression patterns of two

murine homologs of Drosophila single-minded suggest possible roles in embryonic

patterning and in the pathogenesis of Down syndrome. *Mol Cell Neurosci*. Jan

1996;7(1):1-16. doi:10.1006/mcne.1996.0001

17.   Michaud JL, Rosenquist T, May NR, Fan CM. Development of neuroendocrine

lineages requires the bHLH-PAS transcription factor SIM1. *Genes Dev*. Oct 15

1998;12(20):3264-75. doi:10.1101/gad.12.20.3264

18.   Zhang R, Asai M, Mahoney CE, Joachim M, Shen Y, Gunner G, Majzoub JA.

Loss of hypothalamic corticotropin-releasing hormone markedly reduces anxiety

behaviors in mice. *Mol Psychiatry*. May 2017;22(5):733-744. doi:10.1038/mp.2016.136

19.   Sterley TL, Baimoukhametova D, Fuzesi T, Zurek AA, Daviu N, Rasiah NP,

Rosenegger D, Bains JS. Social transmission and buffering of synaptic changes after

stress. *Nat Neurosci*. Mar 2018;21(3):393-403. doi:10.1038/s41593-017-0044-6

20.   Daviu N, Fuzesi T, Rosenegger DG, Rasiah NP, Sterley TL, Peringod G, Bains

JS. Paraventricular nucleus CRH neurons encode stress controllability and regulate

defensive behavior selection. *Nat Neurosci*. Mar 2020;23(3):398-410.

doi:10.1038/s41593-020-0591-0

21.   Fuzesi T, Daviu N, Wamsteeker Cusulin JI, Bonin RP, Bains JS. Hypothalamic

CRH neurons orchestrate complex behaviours after stress. *Nat Commun*. Jun 16

2016;7:11937. doi:10.1038/ncomms11937

22.   Rasiah NP, Loewen SP, Bains JS. Windows into stress: a glimpse at emerging

roles for CRH(PVN) neurons. *Physiol Rev*. Apr 1 2023;103(2):1667-1691.

doi:10.1152/physrev.00056.2021

23.   Daviu N, Bains JS. Should I Stay or Should I Go? CRHPVN Neurons Gate State

Transitions in Stress-Related Behaviors. *Endocrinology*. Jun 1

2021;162(6)doi:10.1210/endocr/bqab061

24.   Novoselova TV, Hussain M, King PJ, Guasti L, Metherell LA, Charalambous M,

Clark AJL, Chan LF. MRAP deficiency impairs adrenal progenitor cell differentiation and

gland zonation. *FASEB J*. Jun 7 2018;32(11):fj201701274RR.

doi:10.1096/fj.201701274RR

25.   Hindmarsh PC, Brook CG. Single dose dexamethasone suppression test in

children: dose relationship to body size. *Clin Endocrinol (Oxf)*. Jul 1985;23(1):67-70.

doi:10.1111/j.1365-2265.1985.tb00184.x

26.   Dogra P, Vijayashankar NP. Dexamethasone Suppression Test. *StatPearls*. 2024.

27.   Toews JNC, Philippe TJ, Hill LA, Dordevic M, Miguelez-Crespo A, Homer NZM,

Nixon M, Hammond GL, Viau V. Corticosteroid-binding Globulin (SERPINA6)

Establishes Postpubertal Sex Differences in Rat Adrenal Development. *Endocrinology*.

Oct 11 2022;163(11)doi:10.1210/endocr/bqac152

28.   Muglia L, Jacobson L, Dikkes P, Majzoub JA. Corticotropin-releasing hormone

deficiency reveals major fetal but not adult glucocorticoid need. *Nature*. Feb 2

1995;373(6513):427-32. doi:10.1038/373427a0

29.   Venihaki M, Carrigan A, Dikkes P, Majzoub JA. Circadian rise in maternal

glucocorticoid prevents pulmonary dysplasia in fetal mice with adrenal insufficiency.

Proc Natl Acad Sci U S A. Jun 20 2000;97(13):7336-41. doi:10.1073/pnas.97.13.7336

30.   Repository Supplemental Materials.

31.   Smith GW, Aubry JM, Dellu F, Contarino A, Bilezikjian LM, Gold LH, Chen R,

Marchuk Y, Hauser C, Bentley CA, Sawchenko PE, Koob GF, Vale W, Lee KF.

Corticotropin releasing factor receptor 1-deficient mice display decreased anxiety,

impaired stress response, and aberrant neuroendocrine development. *Neuron*. Jun

1998;20(6):1093-102. doi:10.1016/s0896-6273(00)80491-2

32.   Bale TL, Picetti R, Contarino A, Koob GF, Vale WW, Lee KF. Mice deficient for

both corticotropin-releasing factor receptor 1 (CRFR1) and CRFR2 have an impaired

stress response and display sexually dichotomous anxiety-like behavior. *J Neurosci*.

Jan 1 2002;22(1):193-9. doi:10.1523/jneurosci.22-01-00193.2002

33.   Bale TL, Vale WW. Increased depression-like behaviors in corticotropin-releasing

factor receptor-2-deficient mice: sexually dichotomous responses. *J Neurosci*. Jun 15

2003;23(12):5295-301. doi:10.1523/jneurosci.23-12-05295.2003

34.   Zorrilla EP, Heilig M, de Wit H, Shaham Y. Behavioral, biological, and chemical

perspectives on targeting CRF(1) receptor antagonists to treat alcoholism. *Drug Alcohol*

Depend. Mar 1 2013;128(3):175-86. doi:10.1016/j.drugalcdep.2012.12.017

35.   Uba AI, Scorese N, Dean E, Liu H, Wu C. Activation Mechanism of Corticotrophin

Releasing Factor Receptor Type 1 Elucidated Using Molecular Dynamics Simulations.

ACS Chem Neurosci. May 5 2021;12(9):1674-1687.

doi:10.1021/acschemneuro.1c00126

36.   Philbert J, Belzung C, Griebel G. The CRF_1 receptor antagonist SSR125543

prevents stress-induced cognitive deficit associated with hippocampal dysfunction:

comparison with paroxetine and D-cycloserine. *Psychopharmacology (Berl)*. Jul

2013;228(1):97-107. doi:10.1007/s00213-013-3020-1

37.   Gafford GM, Guo JD, Flandreau EI, Hazra R, Rainnie DG, Ressler KJ. Cell-type

specific deletion of GABA(A)alpha1 in corticotropin-releasing factor-containing neurons

enhances anxiety and disrupts fear extinction. *Proc Natl Acad Sci U S A*. Oct 2

2012;109(40):16330-5. doi:10.1073/pnas.1119261109

38.   Spierling SR, Zorrilla EP. Don't stress about CRF: assessing the translational

failures of CRF(1)antagonists. *Psychopharmacology (Berl)*. May 2017;234(9-10):1467-

1481. doi:10.1007/s00213-017-4556-2

39.   Chan LF, Campbell DC, Novoselova TV, Clark AJ, Metherell LA. Whole-Exome

Sequencing in the Differential Diagnosis of Primary Adrenal Insufficiency in Children.

Front Endocrinol (Lausanne). 2015;6:113. doi:10.3389/fendo.2015.00113

40.   Dunn AJ, Swiergiel AH. Behavioral responses to stress are intact in CRF-

deficient mice. *Brain Res*. Oct 16 1999;845(1):14-20. doi:10.1016/s0006-

8993(99)01912-5

41.   Horvath K, Juhasz B, Kuti D, Ferenczi S, Kovacs KJ. Recruitment of

Corticotropin-Releasing Hormone (CRH) Neurons in Categorically Distinct Stress

Reactions in the Mouse Brain. *Int J Mol Sci*. Jul 21

2023;24(14)doi:10.3390/ijms241411736

42.   David DJ, Samuels BA, Rainer Q, Wang JW, Marsteller D, Mendez I, Drew M,

Craig DA, Guiard BP, Guilloux JP, Artymyshyn RP, Gardier AM, Gerald C, Antonijevic IA,

Leonardo ED, Hen R. Neurogenesis-dependent and -independent effects of fluoxetine

in an animal model of anxiety/depression. *Neuron*. May 28 2009;62(4):479-93.

doi:10.1016/j.neuron.2009.04.017

43.   Flandreau EI, Ressler KJ, Owens MJ, Nemeroff CB. Chronic overexpression of

corticotropin-releasing factor from the central amygdala produces HPA axis hyperactivity

and behavioral anxiety associated with gene-expression changes in the hippocampus

and paraventricular nucleus of the hypothalamus. *Psychoneuroendocrinology*. Jan

2012;37(1):27-38. doi:10.1016/j.psyneuen.2011.04.014

44.   Makino S, Gold PW, Schulkin J. Effects of corticosterone on CRH mRNA and

content in the bed nucleus of the stria terminalis; comparison with the effects in the

central nucleus of the amygdala and the paraventricular nucleus of the hypothalamus.

Brain Res. Sep 19 1994;657(1-2):141-9. doi:10.1016/0006-8993(94)90961-x

45.   Laryea G, Schutz G, Muglia LJ. Disrupting hypothalamic glucocorticoid receptors

causes HPA axis hyperactivity and excess adiposity. *Mol Endocrinol*. Oct

2013;27(10):1655-65. doi:10.1210/me.2013-1187

46.   Peters A, McEwen BS. Stress habituation, body shape and cardiovascular

mortality. *Neurosci Biobehav Rev*. Sep 2015;56:139-50.

doi:10.1016/j.neubiorev.2015.07.001

47.   Anthony TE, Dee N, Bernard A, Lerchner W, Heintz N, Anderson DJ. Control of

stress-induced persistent anxiety by an extra-amygdala septohypothalamic circuit. *Cell*.

Jan 30 2014;156(3):522-36. doi:10.1016/j.cell.2013.12.040

48.   Frim DM, Robinson BG, Pasieka KB, Majzoub JA. Differential regulation of

corticotropin-releasing hormone mRNA in rat brain. *Am J Physiol*. Apr 1990;258(4 Pt

1):E686-92. doi:10.1152/ajpendo.1990.258.4.E686

49.   Hahner S, Loeffler M, Fassnacht M, Weismann D, Koschker AC, Quinkler M,

Decker O, Arlt W, Allolio B. Impaired subjective health status in 256 patients with

adrenal insufficiency on standard therapy based on cross-sectional analysis. *J Clin

Endocrinol Metab*. Oct 2007;92(10):3912-22. doi:10.1210/jc.2007-0685

50.   Hurtado MD, Cortes T, Natt N, Young WF, Jr., Bancos I. Extensive clinical

experience: Hypothalamic-pituitary-adrenal axis recovery after adrenalectomy for

corticotropin-independent cortisol excess. *Clin Endocrinol (Oxf)*. Dec 2018;89(6):721-

733. doi:10.1111/cen.13803

Figure Legends

Figure 1. Analysis of anxiety-like behaviors in WT, *Mrap*KO and *Crh*KO mice. (**A-C**) Arial-view heat map of EPM displaying genotype-averaged position x time during 6 min EPM test. In the pictogram, 'O' denotes the open arms of the maze while 'C' denotes the closed arms. Color gradient indicates the relative time spent within a specified region (Red=maximum, Black = minimum). (**D**) *Crh*KO mice, compared with WT mice, spend the most time in the open areas of the EPM (P<0.0001), whereas *Mrap*KO show no significant difference compared to WT mice (P=0.0595). (**E**) *Crh*KO mice, compared with WT mice, enter the open areas of the EPM more frequently during the 6 min test period (P=0.0015), while *Mrap*KO show no difference compared to WT mice (P=0.5407).

Figure 2. HPA axis status in WT, *Mrap*KO and *Crh*KO mice following 15 minutes exposure to restraint. Representative images of PVH *Crh* mRNA expression in (**A**) WT, (**B**) *Mrap*KO, and (**C**) *Crh*KO mice. Compared to WT, *Mrap*KO have increased, and *Crh*KO have absent *Crh* mRNA expression. (**D**) PVH *Crh* mRNA semi-quantification analysis shows *Mrap*KO mice have a non-significant trend towards increased PVH *Crh* mRNA compared to WT mice (P=0.052). (**E**) Plasma corticosterone concentrations in WT, *Mrap*KO, and *Crh*KO mice immediately after 15 min of restraint stress. Asterisks denote P≤ 0.0001 compared to WT using One-way ANOVA with Bonferroni correction. Compared to WT mice, *Mrap*KO have undetectable, and *Crh*KO have minimal response to restraint stress. Scale bars = 100 μM.

Figure 3. Analysis of anxiety-like behaviors in untreated WT mice compared to WT,

*Mrap*KO and *Crh*KO mice either treated with dexamethasone or withdrawn from

dexamethasone treatment for 1 week. (**A**) During a 6 min EPM trial, compared to

untreated WT mice, all 3 genotypes on dexamethasone treatment or withdrawn from

treatment spend comparable amounts of time in the open areas. (**B**) During a 6 min

EPM trial, compared to untreated WT mice, all 3 genotypes on dexamethasone

treatment or withdrawn from treatment enter the open areas of the maze at the same

frequency.

Figure 4. Comparison of paraventricular hypothalamic *Crh* mRNA expression in mice

treated with 5-8 weeks of dexamethasone. Representative images of (**A**) WT and (**B**)

*Mrap*KO show low *Crh* mRNA expression during dexamethasone treatment. When

dexamethasone treatment was discontinued for 1 week, (**D**) WT and (**E**) *Mrap*KO *Crh*

mRNA expression increased compared to dexamethasone-treated mice. (**C, F**) *Crh*KO

mice have no detectable *Crh* mRNA expression. (**G**) Semi-quantification of PVH *Crh*

mRNA shows no significant difference in *Crh* expression when compared to the WT

untreated animal values in Figure 2D (P=0.427). Scale bar = 100 µM.

Figure 5. Evaluation of the endocrine response in mice after a 15 min restraint stress

test. (**A**) Mice treated with dexamethasone have suppressed plasma ACTH

concentrations compared with untreated WT mice. ACTH secretion recovers after 1

week of withdrawing glucocorticoid treatment, with *Mrap*KO mice having significantly

elevated plasma ACTH concentrations (P<0.0001). (**B**) Plasma corticosterone

concentrations in mice either treated with dexamethasone or withdrawn from

dexamethasone. Compared with untreated WT mice, all dexamethasone-treated mice

have attenuated corticosterone responses to the restraint stress test, even after

withdrawal of dexamethasone 1 week previously. Asterisks denote $P<0.0001$.

Fig. 1

Fig. 2