A membrane-associated conveyor belt controls the rotational direction of the bacterial type 9 secretion system.

Authors. Abhishek Trivedi[1-3], Jacob A. Miratsky[2,4], Emma C. Henderson[1-3], Abhishek Singharoy[2,4], Abhishek Shrivastava[1-3].

Affiliations. [1]School of Life Sciences, Arizona State University, Tempe, AZ, USA; [2]Biodesign Institute, Arizona State University, Tempe, AZ, USA; [3]Center for Biological Physics, Arizona State University, Tempe, AZ, USA; [4]School of Molecular Sciences, Center for Applied Structural Discovery, Arizona State University, Tempe, AZ, USA.

Corresponding authors. Abhishek Shrivastava. Email: ashrivastava@asu.edu; Abhishek Singharoy. Email: asinghar@asu.edu

Keywords. Bacterial protein secretion, Type 9 Secretion System (T9SS), molecular motors, bacterial motility, gliding motility, Flavobacterium, chemotaxis.

ABSTRACT

Many bacteria utilize the type 9 secretion system (T9SS) for gliding motility, surface

colonization, and pathogenesis. This dual-function motor supports both gliding motility

and protein secretion, where rotation of the T9SS plays a central role. Fueled by the

energy of the stored proton motive force and transmitted through the torque of membrane-

anchored stator units, the rotary T9SS propels an adhesin-coated conveyor belt along the

bacterial outer membrane like a molecular snowmobile, thereby enabling gliding motion.

However, the mechanisms controlling the rotational direction and gliding motility of T9SS

remain elusive. Shedding light on this mechanism, we find that in the gliding bacterium

Flavobacterium johnsoniae, deletion of the C-terminus of a conveyor belt protein controls,

and in fact, reverses the rotational direction of T9SS from counterclockwise (CCW) to

clockwise (CW). Largescale conformational changes at the interface of the T9SS ring with

the C-terminus of the conveyer belt, as well as those of the ring protein themselves, in

concert with a CW bias of the stators general rotation brings forth a 'tri-component

gearset' model: the conveyor belt controls the conformation of the T9SS ring, and thereby

its rotational direction. Consequently, the CW rotating stator either push the outer edge

of the T9SS rings, causing its CCW rotation or press against the inner surface of the rings,

resulting in CW rotation. This regulatory mechanism exemplifies how an outer membrane

associated conveyor belt adjusts the rotational direction of its driver, the T9SS, thus

providing adaptive sensory feedback to control the motility of a molecular snowmobile.

INTRODUCTION

To date, three classes of ion-driven biological rotary motors have been identified: the type 9 secretion system (T9SS), ATP synthase, and the bacterial flagellar motor. The T9SS, discovered recently, is utilized by the Bacteroidetes-chlorobi-fibrobacteres superphylum bacteria. Over 200,000 distinct proteins secreted by T9SS have been cataloged on the InterPro[1], showcasing a remarkable diversity. These proteins span a wide range of functions and include motility adhesins[2], virulent proteases[3,4], and enzymes that break down complex polysaccharides[5,6]. Changes in Bacteroidetes abundances in human microbiota correlate with several diseases. The T9SS of a gut Bacteroidetes, *Paraprevotella clara*, plays a crucial role in defending against viral infections[7]. In contrast, in the case of oral microbiota, the inflammatory response amplification induced by enzymes secreted by T9SS of *Porphyromonas gingivalis* is associated with the onset of periodontitis and Alzheimer's disease[8,9]. Furthermore, the T9SS is required for the development of virulent biofilms[10–12], which are harmful to a variety of hosts including humans[9], birds[13], and fish[14,15].

T9SS also enables the gliding motility of Bacteroidetes over human and animal tissues, fish scales, sediments, and plant roots[16–18]. *Flavobacterium johnsoniae,* found in the soybean rhizosphere[19], is a model organism for investigation of both the macromolecular mechanics of T9SS and the gliding motility of Bacteroidetes. T9SS rotates in place around a stationary axis and its rotational motion is driven by a proton motive force[3]. Following measurements of fluorescently labeled T9SS, tracking of motility adhesins on the conveyor belt, and tethered cell analysis, it has been suggested that

similar to a macromolecular rack and pinion assembly, T9SS propels a polymeric cell-surface conveyor belt over the cell body, thus mobilizing a molecular snowmobile that enables the gliding motility of Bacteroidetes[20] (Fig. 1A). Generally, T9SS stands out as a multifunctional apparatus - it is known to provide competitive advantage to a cell by converting the energy of ionic flow into rotational motion, with the generated rotational energy being utilized to enable either secretion processes, or direct mechanical coupling with a conveyor belt, thus facilitating cellular locomotion. Although gliding motility represents a distinctive example of cellular evolution, essentially nothing is known about molecular mechanisms regulating rotational direction or the mechanical linkage between T9SS and the conveyor belt. In *F. johnsoniae*, known for its robust gliding, the rotation of the GldKN ring propels the linear motion of a cell-surface adhesin, SprB, along a conveyor belt made of the protein GldJ[21–23]. The direct surface interaction between SprB and an external substratum enables gliding motility[21]. Intriguingly, GldJ is missing from T9SS in non-motile bacteria which utilize T9SS exclusively for protein secretion and pathogenesis. For instance, *P. gingivalis* is known to possess a large GldKN (PorKN) ring secretes virulent gingipain proteases via T9SS[8].

T9SS consists of around twenty unique proteins working together to enable both gliding motility and protein secretion[24]. These building-block proteins do not exhibit sequence homology with the extensively studied bacterial flagellar motor that enables swimming motility of other bacteria. The mechanical motion in T9SS is enabled by proton conducting GldLM stator units exhibiting 5:2 structural stoichiometry. Predominantly, GldL is cytoplasmic with transmembrane helices forming a pentameric cage which encloses a

periplasmic GldM dimer. GldL localizes on the rotational axis of tethered cells, and in

gliding cells it remains stationary[25,26]. The GldM (PorM) periplasmic arm has 18-fold

symmetry, and it interacts with the outer-membrane embedded GldKN (PorKN) ring[27].

Macromolecular structure and 5:2 stoichiometry of GldLM are both reminiscent of the

flagellar MotAB proteins, suggesting somewhat similar principles in their evolution. T9SS

and bacterial flagellar motor both use ion channels to spin a ring but the location of the

ring varies. Specifically, in bacterial flagellar motor proton-conducting MotAB stator units

drive rotational motion of a cytoplasmic ring (C-ring), and the rotational bias is controlled

by a cytoplasmic protein sensory transduction protein, CheY-P, to the C-ring[28,29]. In

contrast, in T9SS, the GldLM-driven rotation of GldKN ring occurs in the outer membrane.

Extending the logic that allosteric changes in its C-ring determine the rotational direction

in bacterial flagellar motor, it could be hypothesized that some protein binding to the

GldKN ring might regulate the T9SS rotational bias. This suggests the presence of an

outer membrane-associated sensory transducer, but any evidence is currently absent.

Here, we find that an unlikely candidate, namely, the cell-surface conveyor belt

controls the rotational bias in T9SS. Deletion of 8 amino acids from the C-terminus of the

conveyor belt protein GldJ (GldJ ΔC8) switches T9SS from CCW to CW direction. Further

experimental evolution revealed mutations in the T9SS ring protein GldK which partially

complemented the loss of GldJ's C-terminal region, suggesting conservation of

interaction energy between the T9SS ring and the conveyor belt. Molecular Dynamics

(MD) simulations reveal a substantial decrease in the interactions between GldK and the

truncated GldJ monomer. The simulation further brought to light, movement of the

functional GldK residues into the empty region left at the GldJ-GldK interface from removal of the C-terminus, offering insights into the potential recovery of some lost interface interactions. Additional MD simulations show that in the absence of C-terminal region of GldJ, the ring protein GldK itself undergoes deformation. The T9SS motor is driven by direct interaction of individual GldM proteins within GldLM stator units with the GldKN ring[25]. Unraveling the dynamics of the stator, our steered and Bayesian MD simulations suggest a CW rotational bias for GldM. Therefore, when a CW-rotating stator interacts with the outer periphery of a ring, it induces a CCW rotation of the ring itself. In contrast, when the ring undergoes conformational changes, e.g., due to its exposure to the mutant conveyor belt with lower interaction energy, the CW-rotating GldLM stator units may exert force on the inner periphery of the ring, resulting in CW directionality of T9SS. Hence, we suggest a 'tri-component' gearset model where changes in the conformation of the T9SS rotor arise due to a change in its interaction with the conveyor belt protein GldJ, thereby regulating the rotational direction of T9SS.

The gliding motility of Bacteroidetes showcases seamless coordination of multiple macromolecular units that drive directed cellular motion, yet its actual macromolecular mechanisms have long remained enigmatic[30,31]. Combined, our results here suggest that GldLM stator units, GldKN ring, and GldJ conveyor belt constitute a tri-component gearset driving the gliding motility in Bacteroidetes. Namely, the stator complex harnesses proton motive force to propel the GldKN ring which, in its turn, facilitates and controls the mechanical motion of a closed GldJ conveyor belt. There exists a unique molecular mechanism which alters the directionality of cellular motility, whereby a component of the

cell-surface conveyor belt (GldJ) enables a regulatory feedback loop influencing the

rotational bias of its associated gear (GldKN ring) and thereby altering the directionality

of the propulsive motion of the conveyor belt.

RESULTS

Conveyer-belt protein GldJ controls directionality of T9SS motor. It has been

reported that, in wild-type *F. johnsoniae*, 90% of T9SS units rotate CCW, whereas the

remaining 10% rotate CW[3]. We gathered additional data and quantified rotational

preferences of tethered T9SS in sheared wild-type cells. In agreement with earlier

findings, our data confirms that T9SS is capable of rotating bi-directionally with a bias

toward CCW direction (Fig. 1B, Supplementary video 1). T9SS motors appear to drive

the cell-surface GldJ protein of the conveyor belt[32] (Fig. 1A). Deletion of GldJ destabilizes

GldK and cellular levels of GldK are significantly reduced[33]. Yet, when 8 specific amino

acids on the C-terminus of GldJ are deleted (GldJ ΔC8), GldK levels are maintained.

However, despite near-wild-type levels of GldK, GldJ ΔC8 cells do not exhibit swarming

motility on agar[33]. These observations suggest that GldJ might somehow influence the

rotation or assembly of T9SS. To test this conjecture, GldJ ΔC8 cells were sheared and

tethered to a glass surface. Surprisingly, our experimental findings revealed a distinct

change in T9SS rotational directionality as tethered T9SS motors rotated exclusively CW

in GldJ ΔC8 cells (Fig. 1B, Supplementary video 2). In wild-type cells, T9SS exhibited an

average rotational speed of 1.3 Hz (CCW) whereas in GldJ ΔC8 cells the rotational speed

was about 0.6 Hz (CW, Fig. 1B). The above results indicate that the 8 C-terminal amino

acids of GldJ play a crucial role in regulating both the speed and directionality of the T9SS

motor. In the absence of those amino acids of the conveyor belt protein GldJ, T9SS

rotates exclusively CW.

Experimental evolution revealed that point mutations in GldK in GldJ ΔC8 cells

restored swarming. Wild-type *F. johnsoniae* are known to robustly swarm on PY2 agar,

whereas GldJ ΔC8 cells do not swarm on PY2 agar[33]. Experimental evolution of GldJ ΔC8

cells yielded suppressor strains that were capable of swarming on agar (Fig. 2A, Fig. S1).

Three individual suppressor strains were isolated and their whole genome was

sequenced. All the three turned out to be point mutations in the T9SS ring protein GldK,

with the mutations occurring at the R73S, N307S, and M77L sites of the strains,

respectively.

Single-cell motility over a glass surface of the wild-type, GldJ ΔC8 strain, and the

GldK suppressor strains was recorded, and individual trajectories of cellular motion were

plotted. Wild-type cells exhibited long runs followed by relatively quick turning events

characterized by either back-and-forth cellular motion or rare head-on flips, whereas GldJ

ΔC8 cells mostly exhibited minimal back-and-forth cellular motion. Cells of wild-type and

GldJ ΔC8 strains exhibited average locomotion velocities of about 2 μm/s and 0.4 μm/s,

respectively (Fig. 2B, Supplementary video 3). The above results indicate that in the GldJ

ΔC8 cells the conveyor belt may either be shortened, or even depolymerized. In contrast,

GldK R73S, M77L and N307S suppressor mutants of GldJ ΔC8 background did exhibit

runs of reduced length, with the average locomotion velocities being about 0.89, 0.64,

and 0.88 μm/s. Notably, the oscillatory back-and-forth motion was present along with the runs. The longest runs among the three evolved GldJ ΔC8 strains occurred within the N307S population (Fig. 2C, Supplementary video 4). In summary, the experimentally evolved strains exhibited an improvement in swarming and single-cell motility as compared with the parent GldJ ΔC8 strain. This implies that point mutations in the T9SS rotor protein GldK, which has around 30% sequence similarity to GldJ (Fig. S1B), functionally restores, to a certain degree, the mostly non-operational conveyor belt of the parent GldJ ΔC8 strain.

To gather more information on the integrity of the conveyor belt, a mobile cell-surface adhesin, SprB, which is attached to the conveyor belt, was fluorescently labeled and its movement was tracked. Trajectory of a single SprB signal provided information about the structural integrity of the conveyor belt. As anticipated, wild-type cells exhibited long trajectories with SprB moving at about 2 μm/s, and they appeared to have fully formed, closed conveyor belts looping around cell-poles. In contrast, GldJ ΔC8 strain exhibited drastically short SprB trajectories characteristic of the oscillatory motion at about 0.5 μm/s, suggesting either immobilization or depolymerization of the conveyor belt, but N307S strain of GldJ ΔC8 background exhibited longer trajectories akin to the wild type. This indicates that the functionality of the GldJ conveyor belt is restored, at least, partially, because of the above-mentioned point mutations in the GldK ring of T9SS. In the latter case, the velocity of SprB was about 1 μm/s, half of that characteristic of wild-type cells, suggesting that the T9SS rotor to conveyor belt interaction was partially restored (Fig. 3, Supplementary video 5). To summarize, the detailed analysis of swarming motility and

single-cell motility, facilitated by SprB trajectory tracking, provided evidence that the

interaction between C-terminal region of GldJ and the three experimentally identified

mutations of GldK is crucial for maintaining proper operation of the conveyor belt, which

might enable the cell to search a larger area for nutrients. Additionally, the three GldK

suppressor strains isolated from our screen also had a N178K mutation in a putative

LuxR-like protein (*Fjoh_4220*) and a G223V mutation in an Acyl Carrier Protein (ACP)

Synthase (*Fjoh_3810*). A detailed genetic, single-cell motility, and swarming analysis

showed that *fjoh_4220* and *fjoh_3810* do not impact gliding motility (Fig. S2, Fig. S3,

Supplementary text, and Supplementary videos 6-9).

Experimental evolution unlocked some of the CW-only T9SS motors. As evident by

the data presented in Fig. 1, T9SS of the parent GldJ ΔC8 strain are "locked" in the CW-

only rotational state. Because the evolved GldK point mutations partially restored cellular

motility lost due to ΔC8 in GldJ (Fig. 2, Fig. 3), we tested if some of the T9SS motors in

the evolved strains could rotate CCW. Indeed, the GldK point mutations influenced the

rotational bias as well, with some T9SS motors of the evolved strains rotating CCW, albeit

about 5-10 times slower than those having wild-type GldK. At such low rotational

frequencies, it is tough to differentiate between rotational motion and Brownian

fluctuations. Therefore, only the faster population of T9SS motors that exhibited at least

one full revolution were further analyzed. The above criterion increased the statistical

significance of our results, but, simultaneously, it reduced the overall population of T9SS

motors that were subjected to the analysis. We found that in the R73S strain of GldJ ΔC8

background one out of six T9SS motors taken into consideration rotated CCW (Fig. S4).

In the N307S strain the above was true for one out of sixteen T9SS motors (Supplementary video 10). Contrary to the wild-type units that exhibit CCW bias, T9SS of the evolved strains retained the CW bias of the parent strain. However, in both wild-type and experimentally evolved strains, cellular motility was observed irrespective of the direction of the rotational bias (Fig. 2). In both cases, a rotational bias (whether CCW or CW) of about 85 % to 95 % and about 5 % to 15 % noise contribution was associated with long runs that are characteristic of typical gliding trajectories. In the absence of noise, *i.e*, 100 % CW bias characteristic of the GldJ ΔC8 strain, the cellular motility is diminished (Fig. 1B, 2C).

   In the ideal case, a gliding cell would be equipped with about 8 to 12 T9SS units rotating in unison[34,26]. If one of the units were to rotate in the opposite direction, it would contribute about 8 % to 12.5 % to the noise, which is in agreement with the CCW:CW bias observed in our tethered-cell data (Fig. 1B). Given that multiple T9SS motors control a single conveyor belt, a random change in the rotational direction of any motor, due to inherent noise, could lead to slipping of the conveyor belt. This slippage might then cause disorientation and potentially change the gliding direction of the cell. Within seconds, the slipped conveyor belt might diffuse and re-engage with T9SS motors rotating concertedly, leading to restoration of smooth gliding behavior, albeit in a new direction. The observation that gliding cells follow straight trajectories interspersed with sudden turning events (Fig. 2C) further supports the mechanism suggested above. In their natural environments, gliding cells release enzymes to decompose complex polysaccharides

such as chitin and cellulose found in insoluble food sources[35,36]. Their long gliding

movements, interspersed with sudden turns, may enhance their ability to locate these

food sources. This behavior suggests evolutionary pressure for a strong CCW rotational

bias, with significant CW noise. In the lab, we have observed this pattern in wild-type

strains (Fig. 1B, 2C) and its reversal in evolved strains, where a prominent CW bias is

coupled with noticeable CCW noise (Fig. S4). The observed "inverse state," characterized

by slower cell movement, suggests that the interaction energy between T9SS units and

the conveyor belt differs between wild-type and evolved strains.

A predicted tertiary structure of GldK shows that the residues R73, M77, and N307

are packed close together in an arrangement which resembles a cleft, with the R73 to

N307, R73 to M77, and M77 to N307 inter-residue distances being 14 Å, 12 Å, and 13 Å,

respectively (Fig. 3E). Molecular docking of GldJ and GldK was performed using

HDOCK[37]. One of the models, which has an 86% accuracy score, indicates that the C-

terminal region of GldJ fits into the cleft formed by GldK and aligns with the genetic data

(Fig. 3E). Therefore, it was selected for further analysis. MD simulations performed on

the docked GldJ-GldK model revealed that deleting the C8 region of GldJ causes the cleft

region of GldK, which contains residues R73, M77, and N307, to become more dynamic

and expand towards the area previously occupied by C8 of GldJ (Fig. S5). This explains

why mutations in the ring protein GldK can compensate for the loss of the C-terminal

region of the conveyor belt protein GldJ. This provides further evidence that the

phenotypes described above might be due to changes in the interaction energy between

the T9SS ring and the conveyor belt. Interestingly, the combination of GldK mutations

R73S, M77L, and N307S in a GldJ ΔC8 background does not restore either swarming on agar or single-cell motility over a glass surface (Fig. S6 and supplementary text). This suggests that these three GldK point mutations together might completely hinder the interaction of the rotor with the conveyor belt. Any one of the three GldK mutations is sufficient to alter the function of the cleft region of GldK and consequently its interaction energy with GldJ ΔC8.

The conformation of the T9SS ring protein GldK is determined by its interaction with the conveyor belt protein GldJ. The experimental evidence presented suggests that the interaction mechanism between GldJ and GldK determines the directionality of rotation of the T9SS. To further explore this mechanism, we estimated the interaction energy between monomers of GldJ and GldK using MD. The resulting trajectories were analyzed to assess van der Waals and electrostatic interactions between GldJ and GldK at every timestep over a 100 ns simulation period. We observed that, for the monomers, the total non-bonded GldJ-GldK interaction energy decreased by about 25% upon removal of the 8 C-terminal amino acids of GldJ (Fig. 4A).

We further observed a significant conformational change in the tertiary structure of monomeric GldK upon interaction with GldJΔC8 compared to wild-type GldJ (Fig. 4B and Fig. 4C). Additionally, RMSD values for GldK reformations increased by 10 Å after the interaction interface with GldJ was altered by the C8 deletion (Fig. 4D). In the docked model, the distance between the centers of mass of GldK and both forms of GldJ increased by 7.5 Å during the MD simulation (Fig. S7), reinforcing our inference that the

tertiary structure of GldK undergoes conformational changes after the deletion of the C8

region of GldJ. In stark contrast, the structural changes in GldJ lacking the eight C-

terminal amino acids are similar to those in wild-type GldJ (Fig. S8), thus providing an

internal control for the MD simulations. An overlay of GldK monomer from the GldJΔC8

model with the wild-type GldK shows several regions in which the conformation of GldK

changes > 1 nm (Fig. 4E).

A tri-component gearset engages the conveyor belt, T9SS rotor, and GldLM stator

units to drive bacterial gliding motility. The array of evidence described above from

genetics, cell tethering assays, and MD simulations coupled with the established

knowledge that GldLM stator units power the T9SS[25,34], suggests that the interactions

among the T9SS GldK ring, the GldJ conveyor belt, and the GldLM stator units resemble

the meshing of three distinct mechanical gears. We refer to this complex interaction as a

tri-component gearset. Here, the GldLM stator unit is the primary gear and driver, as it

harnesses proton motive force. The stator unit spins the middle gear, the GldK ring, which,

in turn, propels the outer part, the GldJ conveyor belt. The conveyor belt forms a closed

loop and within a single cell, multiple GldJ loops have been observed by Cryo-ET[23]. When

examined in the 2D plane, the outer loops of the tri-component gearset appear as linear

conveyor belts or multirail tracks (Fig. 5a).

In a gliding cell, the GldKN ring interacts with the GldLM stator units, which harness

pmf to power the rotation of the T9SS[27]. Recent research has demonstrated that in

flagellated bacteria, the MotAB components of the bacterial flagellar motor rotate CW,

driving the rotation of C-rings of the bacterial flagellar motor[38]. The proton-conducting

GldLM stator units have an asymmetric 5:2 structure similar to MotAB. However, despite

this conceptual similarity, the actual rotational dynamics of GldLM stator units at the

atomistic level have not yet been explored. Given that MotAB components rotate

unidirectionally CW[38], analyzing whether GldLM stator units are bidirectional or

unidirectional may help elucidate if GldLM play any role directional switching of T9SS. To

investigate this, we employed MD simulations to model the rotational dynamics of GldLM.

The probability of CW vs CCW GldLM rotation was estimated via MD. Notably,

GldLM has a characteristic structural asymmetry with two GldM chains distributed among

five GldL subunits. The latter feature precludes uniformity of GldL-GldM interactions, and

it has been suggested that a single GldM chain can only be engaged into a channel-

specific ion pairing involving protonated ARG 9 of GldM and deprotonated GLU 49 of

GldL at a time[25]. This ion pair is implicated in transporting protons and generating a proton

motive force. With the PDB all-atom macromolecular structure featuring GldLM symmetry

mismatch as an input (see PDB: 6YS8), three different MD simulations were performed

on the GldLM stator units.

We began with an all-atom equilibrium MD simulation where the non-bonded

interaction energies were monitored across the simulation trajectory to explore the

contacts between ARG 9 of GldM and GLU 49 of GldL, where GldL is either in CCW or

CW orientation. We obtained the mean values of -57.5 and -94.6 kcal/mol for the CCW-

orient GLU 49 and CW-oriented GLU 49 on GldL, respectively (Fig. 5B). The rotation of

transmembrane helices of GldM within GldL is inaccessible using traditional MD

techniques, hence 20 replicas of 10 ns-long steered MD (SMD) simulations were

explored. The results are indicative of a significant relationship between average work

and rotational directionality preference, suggesting a bias towards CW (Fig. 5C).

Prompted by the apparent differences in work profiles, we sought to obtain a more

reliable description of the CCW vs. CW occupancies in GldM through Modeling Employing

Limited Data or MELD-guided MD simulations[39]. We began with a configuration where

the ARG 9 residue was equidistant from the CCW and CW deprotonated GLU 49 of GldL

to avoid any rotational bias in our model. Upon convergence, our MELD simulation

revealed that around 82% of ARG tend to be spontaneously located in the immediate

vicinity of the CW-oriented GLU, in the absence of any directive steering force throughout

the simulation. Thus, as opposed to the (CCW biased) GldK ring of T9SS, the proton

conducting GldM-GldL transmembrane helix complex of T9SS have a CW bias (Fig. 5D).

In our MD simulations, GldLM ion channels rotated mostly CW, and we observed no

transitioning to CCW rotation (Supplementary video 11). Consequently, the MD

simulations revealed that the directional bias change in the T9SS cannot be ascribed to

a change in rotational directionality of the GldLM ion channels that drive this system.

The mechanism for directional switching of T9SS. Our findings suggest that the GldJ

conveyor belt adjusts the rotational directionality of the GldK-containing T9SS ring, while

the GldLM stator units, responsible for generating force, rotate CW unidirectionally,

propelling the middle gear (GldK ring) of the tri-component gearset. We find that the

interaction energy between GldJ and GldK influences the size of the GldK ring, with the

ring undergoing a conformational change when the C8 of GldJ is removed (Fig. 4).

According to our theoretical framework, shifts in rotational directionality in T9SS might

occur due to this conformational change of the GldK ring. Tethered cell assays indicate

that the wild-type T9SS predominantly rotates CCW (Fig. 1B). Therefore, the CW rotation

of the GldLM stator units in wild-type cells likely propels near the outer edge of the GldK

ring. This setup is comparable to a mechanical system where a CW rotating driving gear

moves a second gear. In this case, the second gear (GldK ring) rotates CCW which is

opposite of the CW rotation of the GldLM driving gear (Fig. 6). Interestingly, Song *et al*.

reported various conformational states of the GldKN (PorKN) ring. Using cryo-ET, they

demonstrated that the T9SS GldKN (PorKN) ring of *P. gingivalis* adopts three distinct

conformational classes, forming angles of 90°, 70°, and 50° with the outer membrane[27]

(Fig. 6B). Hence cryo-ET supports our model which explains the two experimentally

observed rotational states of T9SS (Fig. 6).

GldK forms a 32/34-meric ring of T9SS[27] and as demonstrated by tethering assays,

the T9SS ring of *F. johnsoniae* completes one full rotation, on average, every second

(Fig. 1B). For a 32-mer ring, if the conveyor belt is positioned on one side, a GldJ

monomer will interact with a rotating GldK monomer of the ring every 1/32 of a second

(about 31 millisecond). Consequently, each GldK monomer receives an allosteric

stimulus from GldJ approximately every 31 milliseconds, ensuring that the conveyor belt

allosterically controls the conformation of the GldK ring. The short axis cross-section of

the GldM dimer, responsible for propelling the GldK ring, measures around 1.7 nm (Fig.

6C). MD simulations suggest that, in the absence of the C8 region of GldJ, around 1

nanometer sized conformational changes occur within GldK (Fig. 4). This alters the

position of the GldK ring in such a way that the GldM stator protein pushes closer to the

inner region of the ring. This altered positioning of the GldK ring relative to the CW rotating

GldM stator protein results in both the ring and stator units rotating in the same CW

direction, as observed in our tethered cell assay with GldJ ΔC8 cells (Fig. 6D,

Supplementary video 12). The rotation of the GldK ring, either CW or CCW, depends on

where the CW rotating GldLM stator units interact with the GldK ring in a manner

reminiscent to that occurring between the MotAB stators and the flagellar C-ring[38]. At a

cursory glance, the mechanism behind directional switching of T9SS presented here

seems similar to how the rotation direction of the bacterial flagellar motor is regulated[40].

However, the proteins composing the T9SS, and gliding machinery bear no sequence

resemblance to those forming the bacterial flagellar motor, indicating an instance of

convergent evolution. Unlike the bacterial flagellar motor, which is regulated by soluble

cytoplasmic signaling proteins, the outer-membrane-associated conveyor belt protein

GldJ uniquely controls the conformation of the T9SS ring and, consequently, the rotational

directionality of T9SS. This presents a novel mechanism for controlling bacterial cellular

motility, resembling a microbial snowmobile moving over external surfaces, with the

added aspect of adaptive feedback from the cell-surface conveyor belt.

DISCUSSION

The data presented here introduces a novel mechanism that governs the rotational

direction of T9SS, essential for both gliding motility and protein secretion[24,33]. T9SS in

various bacterial species secretes over 300,000 distinct proteins, crucial for processes

such as pathogenicity, host colonization, immunomodulation, polysaccharide

degradation, and interbacterial interactions[13,35,41]. To date, little is known about the

regulation of rotation of T9SS. We report that an outer-membrane conveyor belt protein

GldJ regulates the conformation of the T9SS GldK ring, which, in its turn, changes the

rotational directionality of T9SS. From a cellular perspective, a shift in the rotational

direction of T9SS can prompt the cell to reorient or reverse its movement. In their natural

habitat, bacteria glide on surfaces coated with viscous polysaccharides. It is possible that

if a cell gets stuck in highly viscous environment, the tension on the conveyor belt might

alter, leading to disengagement from the T9SS rotor. Since the interaction energy

between the conveyor belt and the T9SS rotor dictates the directionality of T9SS,

disengagement could prompt the rotor to switch, enabling the cell to maneuver backward

or sideways to escape highly viscous surfaces. This potential scenario has opened new

avenues for research on the implications of feedback control of T9SS by its conveyor belt.

Earlier research posits that GldJ servs as a pathway for the cell-surface adhesin,

SprB, to traverse the length of the cell[21,22]. Our findings, however, unveil yet another role

of the GldJ conveyor belt, specifically of its C-terminal moiety, which operates as a

molecular switch. When full-length GldJ is present, the majority of T9SS units rotate

CCW. On the contrary, in the absence of the C-terminal moiety, T9SS predominantly

rotates CW. Furthermore, experimental evolution of GldK reveal the flexibility of T9SS

rotational motor machinery via a partial restoration of bacterial motility even in the

absence of the GldJ C-terminal. It appears that differing evolutionary pressures

characteristic of a variety of environmental niches have likely shaped the distinct

components and control mechanisms of the rotor and conveyor belt that are specific to a

given niche. The evolved rotor features GldK point mutations in moderately conserved

regions and the GldK sequences of certain species also display varying amino acids at

these sites (Fig. S9). This variability indicates interspecies differences in the

conformations of the GldK ring and suggests the presence of a control mechanism that

enables diverse bacterial species to navigate surfaces with varying chemical properties,

stiffness, and fluidity.

The T9SS machinery stands out from any other known type of proton motive force-

driven biological rotary motor, such as the ATP synthase or bacterial flagellar motor. In

both the ATP synthase and flagellar motor, switching the rotation direction of the rotary

motor significantly impacts crucial functions. For ATP synthase, a change in rotation

direction affects whether ADP is catalyzed into ATP or vice versa. In the case of the

flagellar motor, chemotaxis is controlled by the switching of the rotation direction of the

FliG ring. Bacteria propelled by flagella use a cytoplasmic sensory transduction protein

CheY to alter the rotation direction of their flagellar motor. When CheY is phosphorylated,

it binds to the C-ring of the flagellar motor, causing it to switch rotation from CCW to

CW[28,29] . This action leads to the separation of the flagellar bundle in peritrichous bacteria,

causing the cell to transition from straight swimming (a run) to random swimming in a

small area (a tumble). Reorientation enables the cell to change its swimming direction,

facilitating movement towards chemical attractants[28]. Despite significant differences in

mechanical structure, protein sequences, and biological functions between the T9SS and

the bacterial flagellar motor, their regulation appears to adhere to similar principles, indicative of convergent evolution. In both instances, changes in ring conformation—resulting from structural adjustments within the flagellar FliG ring[40] or from conveyor belt-driven conformational remodeling of the T9SS GldK ring (Fig. 6), affect the positioning of the MotAB or GldLM stator units, respectively. This shift in positioning, in turn, causes the direction of rotation of the FliG or GldK ring to reverse.

Powered by proton motive force, the T9SS stator units play a role akin to that of the driving motor in a mechanical conveyor system. The GldK ring, on the other hand, is akin to the gear that translates rotational energy of the motor into linear motion of the conveyor belt. Meanwhile, the GldJ loop represents the conveyor belt itself, but with an added layer of complexity - unlike a passive mechanical conveyer belt that we commonly encounter in our daily life, GldJ plays a dual role; in addition to be passively driven by forces exerted on it by the GldK gear, it also feeds back into the system to influence and control the rotational directionality of its driving gear itself. In traditional human-engineered mechanical conveyor systems, a series of motors and gears propel the belt along a predetermined path. However, more advanced systems, known as flexible or cognitive mechanical conveyors, are equipped with routing algorithms that adjust the direction of motion of the conveyor belt to adapt to varying requirements[42]. Shaped by evolutionary pressures, the bacterial gliding machinery described here represents a unique example of a conveyor belt equipped with sensory feedback. This biological system could inspire the design of future bio-inspired robots requiring dynamic adaptability. The data presented here demonstrate that bacteria have evolved a smart

snowmobile with a conveyor belt system that is capable of reciprocally altering the

rotational bias of its driver. Thereby, it appears to function as a dynamic and adaptive

sensor or controller. Unveiling such intricate bi-directional feedback deepens our

understanding of fundamental controllers of a bacterial cell whose inner workings

resemble a controllable biological snowmobile.

METHODS

Strains, media, and culture conditions. For standard culture maintenance, genetics,

and tethered cell analysis, wild-type *F. johnsoniae* CJ 1827[43] cells were grown in Casitone

Yeast Extract (CYE) broth (10 g casitone per liter, 5 g yeast extract per liter, 1 M Tris-HCl

pH 7.5) at 30°C with 1.5% agar. *Escherichia coli* cells used for preparation of genetic

constructs were grown in Luria Bertani medium (10g Tryptone per liter, 10g Nacl per liter,

5g yeast extract) at 37°C. Unless otherwise noted, antibiotics were used at the following

concentrations: ampicillin (100 μg/ml), erythromycin (100 μg/ml), streptomycin (100

μg/ml), and cephalexin (75 μM or 26 μg/ml).

Measurement of swarming motility on agar. *F. johnsoniae* cells were grown in PY2

broth (2 g peptone per liter, 0.5 g yeast extract per liter, pH 7.3) overnight at 25°C with

shaking at 50 rpm. Cells were then washed once with PY2 medium and adjusted to an

OD$_{600}$ of 0.1. A 2 μl sample was spotted onto the center of a 1% PY2 agar petri plate,

which was then incubated at 25°C for 48 hours. After incubation, the plates were imaged

using an Amscope LED Trinocular Zoom Stereo microscope (SM-2TZZ-LED-18M3) with

a camera (AmScope 18MP Color CMOS, C-Mount Microscope Camera, MU1803).

Analysis of gliding motility on a glass surface. *F. johnsoniae* cells were grown overnight in CYE broth. Then, 50 µl of cells were inoculated in 5 ml of MM and grown for 6 hours at 25°C with shaking at 50 rpm until the OD_{600} reached 0.4. After that, 45 µl of the culture was introduced into a tunnel slide, incubated for 5 minutes, washed with 40 µl of MM, and imaged with a Nikon Optiphot phase contrast microscope fitted with Thor lab CMOS camera (DCC1545M). Videos were recorded at 15 fps. Custom Python scripts were used to analyze the trajectory and speed of the cells.

Suppressor mutagenesis. *F. johnsoniae* strains lacking 8 amino acids from C-terminal of GldJ (GldJ553/CJ2443) were grown in MM at 30°C for 24 hr. A volume of 20 µl was taken and spotted at the center of twenty different PY2 plates, dried for 20 min, and kept for incubation (lid up) in a 100% relative humidity chamber at 25°C for 3-4 days. Spreading colonies flared out on PY2 plates (Fig. S1). These flares were streaked for isolation and single spreading colonies were selected. Genomic DNA was isolated from the cell using GeneJET Genomic DNA Purification Kit (K0722) and the whole genome was sequenced (Supplementary data S1). As a result, three individual strains, which had the following mutations - GldK R73S (strain name: FJASU1), GldK N307S (strain name: FJASU3), GldK M77L (strain name: FJASU4) flared out as suppressor mutants from (GldJΔC8/CJ2443). All spreading colonies also had the G223V mutation in ACPsynthase (*fjoh_3810*) and N178K mutation in putative LuxR (*fjoh_4220*). Separate mutagenesis of each gene revealed that *fjoh_3810* and *fjoh_4220* did not influence this phenotype (Fig. S2, S3).

Construction of point mutations. To obtain the 2 kb fragment of *gldK* from the suppressor mutants FJASU1, FJASU3, and FJASU4, Phusion high-fidelity PCR master mix (Thermo Scientific) was used with primers PASU52 and PASU53. The resulting fragment was then digested with BamHI and SalI restriction enzymes and cloned into the pRR51 plasmid previously digested with the same enzymes. This plasmid was then introduced into *F. johnsoniae* through triparental conjugation and mutants were isolated as described previously[43]. Briefly, a mixture of PRR51 containing *E. coli* strain, helper *E. coli* strain, and CJ1827 was spotted on a CYE agar plate containing CaCl$_2$. Next day, the mixture was plated with dilutions on a CYE plate containing erythromycin to select for cells that had undergone the first recombination step. The erythromycin resistant colonies were cultured overnight in CYE broth and then plated with dilutions on a PY2 agar plate containing streptomycin to select for cells that had undergone the second recombination step. Point mutations in ACP and LuxR were generated via a similar method with the following exceptions – primers PASU26 and PASU27 were used to amplify ACP synthase region while primers PASU50 and PASU51 were used to amplify putative LuxR region. All primers and plasmids used in this study are in Tables S1 and S2. All mutations were confirmed by Sanger sequencing (Azenta life sciences) All sequencing data is available on dropbox (link).

Immunofluorescence labelling of SprB and imaging via TIRF microscopy. Single colonies of the *F. johnsoniae* GldK N307S in GldJ ΔC8 strain (FJASU19) were grown overnight in MM at 25°C with shaking at 50 RPM. From the overnight saturated culture, 100 μL was used as an initial inoculum to subculture 5 mL of MM containing cephalexin

and incubated at 25°C with shaking at 50 RPM, resulting in the generation of elongated cells. After 6 hours, 100 microliter of culture was washed by centrifugation at 5000 g for 1 min. The resulting pellet was resuspended in 45 µl of MM and 5 microliters of 1:10 diluted purified anti-sprB antibody was added. Anti-SprB antibody[44] was purified using Melon Gel IgG Spin Purification Kit (Thermo Scientific, 45206) as described previously[1]. After incubation at room temperature for 10 min, cells were washed again via centrifugation at 5000 g for 1 min and resuspend in 45 µl MM. The cells were treated with 2 μL of secondary antibody, Alexa Fluor 555-labeled anti-rabbit IgG (Abcam, Cambridge,UK), and incubated at 25°C for 10 minutes. The cells were washed thrice at 5000 g and were pipetted into a tunnel slide. Cells were incubated for 5 min at RT and washed gently with MM in the tunnel slide. Imaging was performed on ONI Nanoimager microscope (Oxford Nanoimaging) equipped with 405 nm diode pump solid state lasers. Optical magnification was provided by a Å~100 oil-immersion objective (Olympus, 1.4 numerical aperture) and images were acquired using an ORCA-Flash4.0 V3 CMOS camera (Hamamatsu). TIRF images were acquired using a 20-ms exposure time, with the 561-nm photoactivation laser at 2% power.

Tethered cell assay. The cells were grown overnight in MM at 25°C with shaking at 50 RPM. From the overnight culture, 100 μL was sub-cultured into fresh 5 mL of MM media and incubated at 25°C with shaking at 50 RPM. When the culture reached an OD_{600} of 0.4, cell tethering was performed as previously described[3]. Briefly, a polyethylene tubing with an inner diameter of 0.58 mm was used to shear cells by passing 500 μL of the culture 50 times through the tubing using 1 mL syringes with 23 G stub adapters. (Instech

PE-50 tubing, Instech Luer stub LS23S, Fisherbrand sterile plastic syringe). The sheared

cells were washed with MM by centrifugation at 5000 g for 1 min. A volume of 40 μL of

this sample was incubated with anti-sprB antibody diluted (1:10) for 20 minutes at room

temperature. After another wash at 5000 g for 1 minute in MM, the cells were pipetted

into the tunnel slide. After 5 minutes, 200 microliters of MM were gently flowed through

the tunnel slides multiple times until residual antibody was washed away and freely

rotating cells were observed.

Preparation of the MD model. The initial model of the membrane protein system was

prepared using CHARMM-GUI membrane builder. The protein structure, consisting of five

GldL subunits and two GldM transmembrane helices[25], was uploaded in PDB format, and

the system was solvated in a preequilibrated phosphatidylcholine (POPC) lipid bilayer.

TIP3P water molecules were added to the system, along with 0.15 M NaCl, to reproduce

cellular conditions.

Equilibrium MD. All-atom molecular dynamics simulations were performed using

Nanoscale Molecular Dynamics (NAMD) 3.0 using the CHARMM36m force field for

protein, lipids, and ions. Periodic boundary conditions were implemented in all

dimensions. The system was energy-minimized followed by an equilibration phase under

NPT conditions at 300 K and 1 atm for 1μs.

Steered MD. Non-equilibrium molecular dynamics simulations were carried out using a

spin angle CV; one of a two-part decomposition of a full rotation CV effectively measuring

the angle of rotation of the GldM TMH around the z-axis. A total of 20 simulations were

executed, evenly bifurcated into two directional categories. The rotation parameter was

established at ± 36°. Negative magnitudes denote a counterclockwise torsional

movement, whereas positive magnitudes signify a clockwise torsion. Subsequently, after

ten iterations, the GldM TMH underwent a complete rotation of ±360°. All simulations

were performed using NAMD3 software using the CHARMM36m force field for protein,

lipids, and ions. Periodic boundary conditions were implemented in all dimensions. The

system was energy-minimized followed by a non-equilibration phase under NPT

conditions at 300 K and 1 atm. Each simulation was run for 10 ns.

Modeling Employing Limited Data (MELD). MELD is a physics-based, Bayesian

computational technique leveraging Open Molecular Mechanics (OpenMM) MD software.

The simulation schematic involved using a 16-replica parallelization assigning one GPU

per replica. The individual simulations spanned a duration of 250 nanoseconds each, with

exchange attempts initiated at regular intervals of 10 picoseconds respectively. There

was an implemented linear temperature range from 300 to 380 K across the replica ladder

i.e., 300 K at replica 1, 380 K at replica 16 and uniform temperature interpolation for the

remaining replicas. To effectively model solvation effects in our system, we utilized the

Generalized Born implicit solvent model (GB-Neck2).

We chose the Boltzmann distribution as the prior, as it provides an unbiased view

of the system's conformational landscape. The likelihood distribution includes all external

information, manifested as flat-bottom harmonic potentials, and comprised of non-

negative constraints on geometric degrees of freedom. The external information was

incorporated from generated contact maps between selected residues whose specificities

are outlined below: We applied distance constraints to the heavy atoms within GldL with

an absolute confidence of 100%, ensuring the inclusion of every computed contact; we

imposed identical constraints within the GldM TMH. These heavy atom constraints

maintained the structural integrity of the model especially at high-temperature replicas.

There were no restraints imposed between GldM and GldL therefore allowing flexibility

and rotation. The prior and likelihood constituted the posterior distribution which was

ultimately sampled during MELD.

FIGURES

Figure 1. The C-terminal region of a conveyor belt protein, GldJ controls the rotational direction of T9SS. (A) *A cartoon showing the current rack and pinion model where the GldK ring of rotary T9SS (pinion) drives the GldJ conveyor belt (rack).* ***(B)*** *Frequency distribution of rotational speed shows that T9SS of wild-type cells (n= 33) rotate primarily in the counterclockwise (CCW) direction whereas T9SS of cells lacking the C-terminal region of GldJ (GldJ ΔC8, n= 21) rotate in the clockwise (CW) direction.*

Figure 2. Experimental evolution restored gliding motility in cells that lack the C-terminal region of GldJ. (A) *Cells lacking the C-terminal region of GldJ (GldJ ΔC8) do not swarm on agar. Experimental evolution of the GldJ ΔC8 strain resulted in partial restoration of swarming. All spontaneous mutations in the evolved strains were in the T9SS ring protein GldK. Scale bar = 6 mm.* ***(B)*** *Frequency distribution of the speed with which wild-type and mutant cells glide over a glass surface.* ***(C)*** *Trajectories of wild-type and mutant cells gliding over a glass surface (n=30).*

Figure 3. **Motion of the cell-surface adhesin SprB is partially restored in GldK N307S carrying GldJ ΔC-terminal cells. (A-C)** *A representative trajectory of fluorescently labeled SprB moving along the conveyor belt of wild-type and mutant cells. (D) Box plot showing speed of at least six SprB molecules for each strain. Statistical significance was analyzed via student's t test. P value: (*P < 0.05, 10^{-3} < **P > 10^{-2}, ****P <= 10^{-4}). (E) Predicted structure of GldK shows that the amino acids R73, M77, and N307 identified via the suppressor screen are in close vicinity. Docking of the conveyor belt protein GldJ (cyan) with the T9SS rotor protein GldK (gray) shows that the GldJ C8 region fits into a GldK cleft region that contains R73 and N307 with M77 in vicinity.*

Figure 4. The size of the T9SS GldK ring is determined by its interaction with the conveyor belt protein GldJ. (A) *Nonbonded interaction energy analysis illustrating a stronger interaction between GldJ and GldK in the wild-type GldJ and GldK system (-776.4 Kcal/mol average) compared to a weaker interaction in the GldJΔC8 and GldK system (-537.3 Kcal/mol average).* **(B)** *An overlay of frames collected for the final 50 ns*

*of the MD simulation for the wild-type GldJ (red) and GldK (blue) system with the GldJ C8 region highlighted in green. **(C)** Upon deletion of the C8 region of GldJ, an appreciable expansion of GldK is observed in contrast to the wild-type system shown in panel B. **(D)** RMSD analysis of GldJ and GldK in both the WT and mutant systems. GldK deviates significantly from its initial structure in the GldJ ΔC8 model reaching a final RMSD of ~20 Å, which is around 10 Å larger than the wild-type model. **(E)** An overlay of the GldK from the wild-type and GldJΔC8 systems depicts conformational changes that are > 1nm.*

Figure 5. The driver gear, GldM, in the tri-component gearset rotates unidirectionally in a CW direction. (A) *A cartoon showing the tri-component gearset with GldM stator unit as the driver of the gliding machinery.* **(B)** *Equilibrium MD simulations and non-bonded energy calculations show that the central arginine residue of GldM interacts more favorably with the glutamate residue of GldL in the CW direction.*

*The Kernel Density Estimation (KDE) method was used to estimate the probability density function of the non-bonded energy from the simulation trajectory. **(C)** Central: Unaltered equilibrium configuration of the initial structure, excluding the membrane for the sake of clarity. Left: Structure after a CCW GldM trans-membrane helix (TMH) rotation of -36°. Right: Result after +36° CW rotation. The plot illustrates the non-equilibrium energetics from ten consecutive SMD simulations (360° total in each direction), which reveal a lower work value for CW GldM TMH rotation. **(D)** Three-dimensional representation of the central structure shown in panel C. A top-down view highlights the plane used for the statistical analysis of post-MELD simulations. The spatial coordinates of the unbound arginine residue are mapped in a contour plot, with lighter shades indicating higher density, which suggests an unbiased preference for CW rotation. The arrows indicate the trajectory of the positional displacements of the unbound arginine. A green circle superimposed on the plot serves as the boundary where the probability of a datum falling within the enclosed region was computed.*

Figure 6. The mechanism for directional switching of T9SS. *(A) A cross-sectional view of the gliding machinery shows the proton motive force (pmf) powered driver gear, GldM, rotating unidirectionally clockwise (CW) and pushing the T9SS ring. This action results in the movement of the GldJ conveyor belt and SprB adhesin either towards the viewer or away from the current image plane. Images are not to scale. (B) A cartoon summarizing the different conformational states of T9SS ring[27]. (C) A partial structure of GldM showing that the cross-section of the region of GldM engaging with the T9SS ring is approximately 1.7 nm. (D) A model demonstrating that when the wild-type conveyor belt protein GldJ interacts with the middle gear, the GldK ring, GldK maintains a conformation where GldM pushes the outer periphery of the GldK ring, causing the T9SS*

to rotate counterclockwise (CCW). Conversely, when the interaction energy between GldJ and GldK is reduced, as seen with GldJΔC8, the GldK ring changes conformation. Consequently, GldM pushes near the inner periphery of the GldK ring, resulting in CW rotation of the T9SS. Images are not to scale.

Acknowledgements. A. Shrivastava was supported by an NIH-NIGMS MIRA award R35GM147131. A. Singharoy was supported by an NSF CAREER award MCB-1942763. The MD simulation used an Oak Ridge Leadership Computing Facility, supported an award from the Office of Science, Department of Energy DE-AC05-00OR22725. We thank Mohammed Kaplan, Grant J. Jensen, and Stefano Maggi for helpful comments on the manuscript. We thank Pushkar P. Lele for helpful suggestions and Rizal F. Hariadi for help with TIRF microscopy. The images were created using biorender.com.

Author contributions. AT generated the mutants and performed the cell tethering and TIRF assays. AT and ECH collected and analyzed the cell-motility data. A. Shrivastava generated the initial suppressor strains. JAM and A. Singharoy performed the MD simulations. A. Shrivastava and AT wrote the manuscript.

Data Availability. Custom Python codes for image analysis and example data sets are freely available on our GitHub. The molecular dynamics models, and additional images are available upon request. All other study data are included in the article and/or supporting information.

References.

1. Paysan-Lafosse, T. *et al.* InterPro in 2022. *Nucleic Acids Research* **51**, D418–D427 (2023).

2. Shrivastava, A., Johnston, J. J., van Baaren, J. M. & McBride, M. J. Flavobacterium johnsoniae GldK, GldL, GldM, and SprA are required for secretion of the cell surface gliding motility adhesins SprB and RemA. *J Bacteriol* **195**, 3201–3212 (2013).

3. Shrivastava, A., Lele, P. P. & Berg, H. C. A rotary motor drives Flavobacterium gliding. *Curr Biol* **25**, 338–341 (2015).

4. Saiki, K. & Konishi, K. Identification of a Porphyromonas gingivalis novel protein sov required for the secretion of gingipains. *Microbiol Immunol* **51**, 483–491 (2007).

5. Kharade, S. S. & McBride, M. J. Flavobacterium johnsoniae PorV is required for secretion of a subset of proteins targeted to the type IX secretion system. *J Bacteriol* **197**, 147–158 (2015).

6. Zhu, Y. & McBride, M. J. The unusual cellulose utilization system of the aerobic soil bacterium Cytophaga hutchinsonii. *Appl Microbiol Biotechnol* **101**, 7113–7127 (2017).

7. Li, Y. *et al.* Identification of trypsin-degrading commensals in the large intestine. *Nature* **609**, 582–589 (2022).

8. Sato, K. *et al.* A protein secretion system linked to bacteroidete gliding motility and pathogenesis. *Proc Natl Acad Sci U S A* **107**, 276–281 (2010).

9. Dominy, S. S. *et al.* Porphyromonas gingivalis in Alzheimer's disease brains: Evidence for disease causation and treatment with small-molecule inhibitors. *Sci Adv* **5**, eaau3333 (2019).

10.   Kita, D. *et al.* Involvement of the Type IX Secretion System in Capnocytophaga ochracea Gliding Motility and Biofilm Formation. *Appl Environ Microbiol* **82**, 1756–1766 (2016).

11.   Shrivastava, A. *et al.* Cargo transport shapes the spatial organization of a microbial community. *Proc Natl Acad Sci U S A* **115**, 8633–8638 (2018).

12.   Ratheesh, N. K., Calderon, C. A., Zdimal, A. M. & Shrivastava, A. *Bacterial Swarm-Mediated Phage Transportation Disrupts a Biofilm Inherently Protected from Phage Penetration*. http://biorxiv.org/lookup/doi/10.1101/2021.06.25.449910 (2021) doi:10.1101/2021.06.25.449910.

13.   Guo, Y. *et al.* Riemerella anatipestifer Type IX Secretion System Is Required for Virulence and Gelatinase Secretion. *Front Microbiol* **8**, 2553 (2017).

14.   Pisarenko, O., Studneva, I. & Khlopkov, V. Metabolism of the tricarboxylic acid cycle intermediates and related amino acids in ischemic guinea pig heart. *Biomed Biochim Acta* **46**, S568-571 (1987).

15.   Barbier, P. *et al.* The Type IX Secretion System Is Required for Virulence of the Fish Pathogen Flavobacterium psychrophilum. *Appl Environ Microbiol* **86**, e00799-20 (2020).

16.   How, K. Y., Song, K. P. & Chan, K. G. Porphyromonas gingivalis: An Overview of Periodontopathic Pathogen below the Gum Line. *Front Microbiol* **7**, 53 (2016).

17.   LaFrentz, B. R. *et al.* The fish pathogen Flavobacterium columnare represents four distinct species: Flavobacterium columnare, Flavobacterium covae sp. nov., Flavobacterium davisii sp. nov. and Flavobacterium oreochromis sp. nov., and

emended description of Flavobacterium columnare. *Systematic and Applied Microbiology* **45**, 126293 (2022).

18.   Yin, C. *et al.* Rhizosphere community selection reveals bacteria associated with reduced root disease. *Microbiome* **9**, 86 (2021).

19.   Niraula, S., Rose, M. & Chang, W.-S. Microbial co-occurrence network in the rhizosphere microbiome: its association with physicochemical properties and soybean yield at a regional scale. *J Microbiol* **60**, 986–997 (2022).

20.   McBride, M. J. *Bacteroidetes* Gliding Motility and the Type IX Secretion System. *Microbiol Spectr* **7**, 7.1.15 (2019).

21.   Nakane, D., Sato, K., Wada, H., McBride, M. J. & Nakayama, K. Helical flow of surface protein required for bacterial gliding motility. *Proceedings of the National Academy of Sciences* **110**, 11145–11150 (2013).

22.   Shrivastava, A., Roland, T. & Berg, H. C. The Screw-Like Movement of a Gliding Bacterium Is Powered by Spiral Motion of Cell-Surface Adhesins. *Biophysical Journal* **111**, 1008–1013 (2016).

23.   Shibata, S. *et al.* Filamentous structures in the cell envelope are associated with bacteroidetes gliding machinery. *Commun Biol* **6**, 94 (2023).

24.   Trivedi, A., Gosai, J., Nakane, D. & Shrivastava, A. Design Principles of the Rotary Type 9 Secretion System. *Front Microbiol* **13**, 845563 (2022).

25.   Hennell James, R. *et al.* Structure and mechanism of the proton-driven motor that powers type 9 secretion and gliding motility. *Nat Microbiol* **6**, 221–233 (2021).

26.	Vincent, M. S. *et al. Dynamic Proton-Dependent Motors Power Type IX Secretion and Gliding Adhesin Movement in* Flavobacterium. http://biorxiv.org/lookup/doi/10.1101/2021.10.19.464928 (2021) doi:10.1101/2021.10.19.464928.

27.	Song, L. *et al.* A unique bacterial secretion machinery with multiple secretion centers. *Proc Natl Acad Sci U S A* **119**, e2119907119 (2022).

28.	Sarkar, M. K., Paul, K. & Blair, D. Chemotaxis signaling protein CheY binds to the rotor protein FliN to control the direction of flagellar rotation in Escherichia coli. *Proc Natl Acad Sci U S A* **107**, 9370–9375 (2010).

29.	Chang, Y. *et al.* Molecular mechanism for rotational switching of the bacterial flagellar motor. *Nat Struct Mol Biol* **27**, 1041–1047 (2020).

30.	Glaser, J. & Pate, J. L. Isolation and characterization of gliding motility mutants of Cytophaga columnaris. *Archiv. Mikrobiol.* **93**, 295–309 (1973).

31.	Lapidus, I. R. & Berg, H. C. Gliding motility of Cytophaga sp. strain U67. *J Bacteriol* **151**, 384–398 (1982).

32.	Braun, T. F. & McBride, M. J. Flavobacterium johnsoniae GldJ is a lipoprotein that is required for gliding motility. *J Bacteriol* **187**, 2628–2637 (2005).

33.	Johnston, J. J., Shrivastava, A. & McBride, M. J. Untangling Flavobacterium johnsoniae Gliding Motility and Protein Secretion. *J Bacteriol* **200**, e00362-17 (2018).

34.	Shrivastava, A. & Berg, H. C. A molecular rack and pinion actuates a cell-surface adhesin and enables bacterial gliding motility. *Sci Adv* **6**, eaay6616 (2020).

35.   Kharade, S. S. & McBride, M. J. Flavobacterium johnsoniae chitinase ChiA is required for chitin utilization and is secreted by the type IX secretion system. *J Bacteriol* **196**, 961–970 (2014).

36.   Zhu, Y. & McBride, M. J. Deletion of the Cytophaga hutchinsonii type IX secretion system gene sprP results in defects in gliding motility and cellulose utilization. *Appl Microbiol Biotechnol* **98**, 763–775 (2014).

37.   Yan, Y., Tao, H., He, J. & Huang, S.-Y. The HDOCK server for integrated protein-protein docking. *Nat Protoc* **15**, 1829–1852 (2020).

38.   Rieu, M., Krutyholowa, R., Taylor, N. M. I. & Berry, R. M. A new class of biological ion-driven rotary molecular motors with 5:2 symmetry. *Front Microbiol* **13**, 948383 (2022).

39.   Shekhar, M. *et al.* CryoFold: determining protein structures and data-guided ensembles from cryo-EM density maps. *Matter* **4**, 3195–3216 (2021).

40.   Chang, Y., Carroll, B. L. & Liu, J. Structural basis of bacterial flagellar motor rotation and switching. *Trends in Microbiology* **29**, 1024–1033 (2021).

41.   Wang, S., Zhao, D., Bai, X., Zhang, W. & Lu, X. Identification and Characterization of a Large Protein Essential for Degradation of the Crystalline Region of Cellulose by Cytophaga hutchinsonii. *Appl Environ Microbiol* **83**, e02270-16 (2017).

42.   El-sayed, M. E., Youssef, A. W., Shehata, O. M., Shihata, L. A. & Azab, E. Computer vision for package tracking on omnidirectional wheeled conveyor: Case study. *Engineering Applications of Artificial Intelligence* **116**, 105438 (2022).

43.   Rhodes, R. G., Pucker, H. G. & McBride, M. J. Development and use of a gene

deletion strategy for Flavobacterium johnsoniae to identify the redundant gliding

motility genes remF, remG, remH, and remI. *J Bacteriol* **193**, 2418–2428 (2011).

44.   Nelson, S. S., Bollampalli, S. & McBride, M. J. SprB is a cell surface component

of the Flavobacterium johnsoniae gliding motility machinery. *J Bacteriol* **190**, 2851–

2857 (2008).

Supplementary Information

A membrane-associated conveyor belt controls the rotational direction of the bacterial type 9 secretion system.

Authors. Abhishek Trivedi[1-3], Jacob A. Miratsky[2,4], Emma C. Henderson[1-3], Abhishek Singharoy[2,4], Abhishek Shrivastava[1-3].

Affiliations. [1]School of Life Sciences, Arizona State University, Tempe, AZ, USA; [2]Biodesign Institute, Arizona State University, Tempe, AZ, USA; [3]Center for Biological Physics, Arizona State University, Tempe, AZ, USA; [4]School of Molecular Sciences, Center for Applied Structural Discovery, Arizona State University, Tempe, AZ, USA.

Corresponding authors. Abhishek Shrivastava. Email: ashrivastava@asu.edu; Abhishek Singharoy. Email: asinghar@asu.edu

SUPPLEMENTARY FIGURES

Figure S1. (A) Images from the experimental evolution screen of GldJ ΔC8 which lead to the following suppressors that had the ability to swarm on agar (i) GldK R73S (FJASU1) (ii) GldK N307S (FJASU3) and (iii) GldK M77L (FJASU4). Scale bar, 3 mm. (B) Multiple sequence alignments of *F. johnsoniae* GldJ and Gld

Figure S2. The evolved strains containing either GldK R73S, N307S or M77L restore the swarming motility of the parent GldJ ΔC8 (GldJ ΔC-term background) on PY2 agar plates. Scale bar, 6 mm. Whole genome sequencing revealed that evolved strains also had secondary point mutations in genes encoding an ACP synthase (G223V) and LuxR-like protein (N178K). Individual and combinatorial mutagenesis of ACP synthase and LuxR showed that these two genes do not play a role in restoration of swarming motility.

Figure S3. Microscopic imaging and single cell tracking over a glass surface provided further evidence that G223V mutation in ACP synthase and N178K mutation in a LuxR-like protein do not play a role in the restoration of motility.

Figure S4. The rotational direction of T9SS motors of GldJ ∆C8 cells with GldK N307S mutation (evolved strain) is predominantly CW but unlike the parent GldJ ∆C8 cells where all T9SS motors are CW, few CCW motors (red) are also observed in the evolved strain.

Figure S5. MD stimulations show that in the GldJ ΔC8 strain, the functional GldK residues identified by the suppressor screen (blue, white, and green stick diagrams) can dynamically replace the region previously occupied C8 of wild-type GldJ (shown as a green surface).

Figure S6. Combination of all three GldK point mutations (GldK R73S, M77L, and N307S) in a GldJ ΔC8 background does not restore (A) swarming on agar and (B) Single cell-motility over a glass surface. Scale bar for panel A is 6 mm.

Figure S7. The inter center of mass distances, including all heavy atoms, between GldK and GldJ in the wild-type and GldJ ΔC8 systems recorded periodically throughout the MD simulation.

Figure S8. The RMSD of GldJ in the GldJ-GldK system, compared to GldJΔC8 in the

GldJΔC8-GldK system, indicates that GldJ does not expand in a manner similar to GldK

when C8 of GldJ is deleted.

Figure S9. Multiple sequence alignment (MSA) of GldK protein, highlighting point mutations in moderately conserved regions across various species. Each mutation is indicated by a specific color, Red GldKR73S, Yellow GldK M77L and Green GldKN307S reflecting its evolutionary conservation and potential impact on protein function.

SUPPLEMENTARY TEXT

Single amino acid substitutions of LuxR and Acyl Carrier Protein (ACP) Synthase in a GldJ ΔC8 background does not improve motility. The three GldK suppressor strains isolated from our screen also had a N178K mutation in a putative LuxR-like protein (*Fjoh_4220*) and a G223V mutation in an Acyl Carrier Protein (ACP) Synthase (*Fjoh_3810*). No reports suggest that these two proteins influence Bacteroidetes gliding motility, and it's plausible that these mutations were already present in the parent strain prior to the suppressor screen. To gain a deeper understand of their role, individual strains that carry LuxR N178K, ACP G223V, and LuxR N178K plus ACP G223V mutation in a GldJ ΔC-terminal background were generated. When comparing the new strains with mutations in LuxR and ACP to the original GldJ ΔC-terminal strain, there were no noticeable improvements in swarming and single-cell motility (Supplementary video 6, Supplementary video 7) The strains that had both GldK R73S, M77L, and N307S mutations, along with the LuxR and ACP mutations, showed similar swarming and single-cell motility characteristics as the strains with only GldK R73S, M77L, and N307S mutations in a GldJ ΔC-terminal background. Similarly, for LuxR N178K and ACP G223V individually GldJ ΔC-terminal background exhibited no motility. Furthermore, when both LuxR N178K and ACP G223V were present in individual GldK point mutants no improvement in speed was observed (Supplementary video 8, 9). These results indicate that the suppressor phenotype is not influenced by the LuxR N178K and ACP G223V mutations (Figs. S3, S4).

Combinatorial mutagenesis of GldK point mutations in a GldJ ΔC-terminal

background does not restore motility. A triple GldK mutant strain, which includes the

mutations GldK R73S, M77L, and N307S in the GldJ ΔC8 background, was created.

However, this mutant strain did not exhibit swarming on agar, and its single cells displayed

sluggish back and forth motion (Fig. S6). These characteristics were comparable to the

behavior observed in cells of the original GldJ ΔC8 strain.

SUPPLEMENTARY TABLES.

Table S1. Strain List

Strain	Genotype	Reference
CJ1827	rpsL2; (Sm^r). Wild-type strain, derived from ATCC 17061T (UW101), utilized in the construction of the deletion mutants described below.	(Rhodes et al., 2017)
CJ2443	*GldJ ΔC8*	(Joseph et al., 2017)
FJASU_1	*GldJ ΔC8 fjoh_4220* (N178K) *fjoh_3810* (G223V) *GldK* R73S.	This study
FJASU_3	*GldJ ΔC8 fjoh_4220 fjoh_3810 GldK* N307S.	This study
FJASU_4	*GldJ ΔC8 fjoh_4220 fjoh_3810 GldK* M77L.	This study
FJASU_11	*GldJ ΔC8 fjoh_3810* (G223V)	This study
FJASU_12	*GldJ ΔC8 fjoh_4220* (N178K)	This study
FJASU_13	*GldJ ΔC8 fjoh_*4220 (N178K) *GldK* R7S	This study
FJASU_14	*GldJ ΔC8 fjoh_4220* (N178K) *GldK* N307S	This study
FJASU_15	*GldJ ΔC8 fjoh_4220* (N178K) *GldK* M77L	This study
FJASU_16	*GldJ ΔC8 fjoh_3810* (G223V) *GldK* R73S	This study

FJASU_17	*GldJ* ΔC8 *fjoh_3810* (G223V) *GldK* N307S	This study
FJASU_18	*GldJ* ΔC8 *fjoh_3810* (G223V) *GldK* M77L	This study
FJASU_19	*GldJ* ΔC8 *GldK* N307S	This study
FJASU_20	*GldJ* ΔC8 *GldK* M77L	This study
FJASU_21	*GldJ* ΔC8 *GldK* R73S	This study
FJASU_54	*GldJ* ΔC8 *fjoh_3810* (G223V) *fjoh_4220* (N178K)	This study
FJASU_75	*GldJ* ΔC8 with *GldK* R7SS, M77L, N307S	This study

Table S2. **List of plasmids used in this study**.

Plasmid	Description	Reference
pRR51	rpsL-containing suicide vector; Ap^r (Em^r);	(Rhodes et al., 2017)
ES27	Suicide plasmid used to introduce point mutation in *fjoh_4220*; *fjoh_4220* (N178K) using pRR51.	This study
ES28	Suicide plasmid used to introduce point mutation in *fjoh_3810*; *fjoh_3810* (G223V) using pRR51.	This study
ES29	Suicide plasmid used to introduce point mutation in *GldK*; *GldK* (R73S) using pRR51.	This study
ES30	Suicide plasmid used to introduce point mutation in *GldK*; *GldK* (N307S) using pRR51.	This study
ES31	Suicide plasmid used to introduce point mutation in *GldK*; *GldK* (M77L) using pRR51.	This study
ES35	Suicide plasmid used to introduce wild type copy of *fjoh 4220*. using pRR51	This study
ES36	Suicide plasmid used to introduce wild type copy of *fjoh_3810* using pRR51.	This study
ES37	Suicide plasmid used to introduce all three point mutations in *GldK*; *GldK* (R73S,N307S,M77L) using pRR51.	This study

The antibiotic resistance phenotypes are as follows: ampicillin (Ap[r]), erythromycin (Em[r]), and streptomycin (Smr). The resistance phenotypes indicated in parentheses are those expressed in *F. johnsoniae*, while those without parentheses are expressed in *E. coli*

Table S3. List of primers used in this study.

Primer	Sequence 5'-3'	Used in construction of
PASU26	GCTAGG<u>GATCC</u>GGCACAAAGAGTTTTATTGC	ES28, BamH1 site is underlined. (GCTAG additional sequence added to all primers to improve cutting efficiency)
PASU27	GCTAG<u>GTCGAC</u>CATGCTCCAAAGTATCCAAA	ES28, SalI site is underlined.
PASU50	GCTAGG<u>GATCC</u>AGATATTTTATGGCCGATGT	ES27, BamH1 site is underlined.
PASU51	GCTAG<u>GTCGAC</u>CAGGACCAGAAAACTTAAT	ES27, SalI site is underlined.
PASU52	GCTAGG<u>GATCC</u>AATACATCGTTCCGCTGGAA	ES29, ES30 and ES31 BamH1 site is underlined.
PASU53	GCTAG<u>GTCGAC</u>ATTGCTTTTGCAGCTCCTTC	ES29, ES30 and ES31 SalI site is underlined.

SUPPLEMENTARY VIDEOS

Supplementary video 1. Timelapse images of a tethered wild-type cell with a counterclockwise rotating T9SS.

Supplementary video 2. Timelapse images of a tethered GldJ ΔC8 cell with a clockwise rotating T9SS.

Supplementary video 3. Timelapse images showing gliding of wild-type cells and GldJ ΔC8 cells over a glass surface.

Supplementary video 4. Timelapse images of cells containing GldK R73S, GldK M77L and GldK N307S individual point mutations in GldJ ΔC8 background over a glass surface.

Supplementary video 5. Timelapse TIRF images of immunofluorescent SprB motion on the surface of a wild-type cell, GldJ ΔC8 cell, GldK N307S in GldJ ΔC8 background.

Supplementary video 6. Timelapse images, over a glass surface, of GldJ ΔC8 cells containing GldK R73S ACP G223V, GldK M77L ACP G223V, and GldKN307S ACP G223V.

Supplementary video 7. Timelapse images, over a glass surface, of GldJ ΔC8 cells containing GldK R73S LuxR N178K, GldK M77L LuxR N178K, and GldKN307S LuxR N178K.

Supplementary video 8. Timelapse images, over a glass surface, of cells containing ACP_G223V and LuxR_N178K mutation individually a GldJ ΔC8 background and ACP_G223V and LuxR_N178K together in a GldJ ΔC8 background.

Supplementary video 9. Timelapse images, over a glass surface, of cells containing (i) GldK R73S, ACPG223V, and LuxR N178K mutations in a GldJ ΔC8 background. (ii) GldK R73S, ACP G223V, and LuxR N178K mutations in a GldJ ΔC8 background. (iii) GldK N307S, ACP G223V, and LuxR N178K mutations in a GldJ ΔC8 background.

Supplementary video 10. Time-lapse images from tethered cell assays of the evolved GldK mutant strains in a GldJ ΔC8 background, illustrating examples with both CW and CCW rotating T9SS.

Supplementary video 11. Timelapse MD simulation images showing CW rotation of the GldLM stator units.

Supplementary video 12. A model for directional switching of T9SS.