Evolutionary unique *N*-glycan-dependent protein quality control system plays pivotal

roles in cellular fitness and extracellular vesicle transport in *Cryptococcus neoformans*

Catia Mota[1], Kiseung Kim[1], Ye Ji Son[1], Eun Jung Thak[1], Su-Bin Lee[1], Ju-El Kim[2], Jeong-

Kee Yoon[2], Min-Ho Kang[3,4], Heeyoun Hwang[5], Yong-Sun Bahn[6], J. Andrew Alspaugh[7], and

Hyun Ah Kang[1*]

[1] Department of Life Science, Chung-Ang University, Seoul, 06974, Republic of Korea

[2] Department of Systems Biotechnology, Chung-Ang University, Anseong-Si, Gyeonggi-Do,

17546, Republic of Korea

[3] Department of Biomedical-Chemical Engineering, The Catholic University of Korea,

Bucheon-Si, 14662, Republic of Korea

[4] Department of Biotechnology, The Catholic University of Korea, Bucheon-Si, 14662,

Republic of Korea

[5] Digital OMICs Research Center, Korea Basic Science Institute, Cheongju-si, 28119,

Republic of Korea

[6] Department of Biotechnology, Yonsei University, Seoul, 03722, Republic of Korea

[7] Department of Medicine, Duke University, Durham, North Carolina, USA

*For correspondence (H. A. K.) E-mail: hyunkang@cau.ac.kr; Tel.: +82-2-820-5863; Fax:

+82-2-825-5206

Abstract

A conserved *N*-glycan-dependent endoplasmic reticulum protein quality control (ERQC) system has evolved in eukaryotes to ensure accuracy during glycoprotein folding. The human pathogen *Cryptococcus neoformans* possesses a unique *N*-glycosylation pathway that affects microbial physiology and interactions with the infected host. To investigate the molecular features and functions of the ERQC system in *C. neoformans,* we characterized a set of mutants with deletion of genes coding for the ERQC sensor UDP-glucose:glycoprotein glucosyltransferase (*UGG1*) and putative α1,2-mannose-trimming enzymes (*MNS1*, *MNS101*, *MNL1*, and *MNL2*). The *ugg1Δ*, *mns1Δ*, *mns101Δ*, and *mns1Δ101Δ* mutants showed alterations in *N*-glycan profiles, defective cell surface organization, decreased survival in host cells, and varying degrees of reduced *in vivo* virulence. The *ugg1Δ* strain exhibited severely impaired extracellular secretion of capsular polysaccharides and virulence-related enzymes. Comparative transcriptome analysis showed the upregulation of protein folding, proteolysis, and cell wall remodeling genes, indicative of induced ER stress. However, no apparent changes were observed in the expression of genes involved in protein secretion or capsule biosynthesis. Additionally, extracellular vesicle (EV) analysis combined with proteomic analysis showed significant alterations in the number, size distribution, and cargo composition of EVs in *ugg1Δ*. These findings highlight the essential role of the functional ERQC system for cellular fitness under adverse conditions and proper EV-mediated transport of virulence factors, which are crucial for the full fungal pathogenicity of *C. neoformans*.

Keywords: *Cryptococcus neoformans*, glycoprotein quality control, UDP-glucose: glycoprotein glucosyltransferase, extracellular vesicles, pathogenicity

Introduction

Glycoproteins, destined for the secretory pathways, enter the endoplasmic reticulum (ER) lumen where protein folding occurs (Rapoport, 2007). The accumulation of misfolded proteins affects cell viability and homeostasis; therefore, eukaryotes evolved a conserved ER quality control (ERQC) system that recognizes folding defects, repairs them, or ensures the translocation of irreparable misfolded proteins into the cytosol for proteasome-mediated degradation via the ER-associated degradation (ERAD) system (Thibault and Ng, 2012; Xu and Ng, 2015; Balchin et al., 2016), a process that heavily relies on N-glycosylationto determine protein folding conformation (Aebi, 2013; Varki, 2017).

Many aspects of N-glycosylation are highly conserved, and most eukaryotes initiate this process by synthesizing a Dol-PP-linked $Glc_3Man_9GlcNAc_2$ oligosaccharide as a common core N-glycan, which attaches to nascent polypeptides in the ER via an asparagine residue. Glucosidases I and II (Gls1 and Gls2) remove two glucose residues from the core oligosaccharide. Proteins containing monoglucosylated N-glycans ($Glc_1Man_9GlcNAc_2$) enter a calnexin and/or calreticulin (CNX/CRT in mammals; Cne in yeast) chaperone-mediated folding cycle. Finally, Gls2 cleaves the remaining glucose residue. If proteins are misfolded, they are recognized by the ERQC checkpoint enzyme UDP-glucose:glycoprotein glucosyltransferase (UGGT), which reglucosylates them for re-entry into the folding cycle (Fig. 1A). Then, the N-glycans of the accurately folded proteins are further processed by Gls2 and α1,2-mannosidase I (Mns1) and moved to the Golgi apparatus. However, irreparably misfolded glycoproteins are targeted for ERAD, where they undergo demannosylation and retro-translocation for proteasomal degradation in the cytosol. Recently, N-glycan precursors in the ER of some protists and fungi were found to be shorter than the typical 14-sugar N-glycan precursors in most eukaryote organisms. The length of these N-glycan precursors significantly impacts N-glycan-dependent QC of glycoprotein folding and ERAD (Banerjee

et al., 2007; Samuelson et al., 2015).

The basidiomycetous fungus *Cryptococcus neoformans* is an opportunistic

encapsulated human pathogen that primarily affects immunocompromised individuals,

causing fatal meningoencephalitis (Gottfredsson and Perfect, 2000; Kwon-Chung et al.,

2000). The *N*-glycosylation pathway of *C. neoformans* is evolutionarily conserved;

nevertheless, the structure and biosynthesis of its *N*-glycans includes several unique features

(Park et al., 2012). *C. neoformans* contains serotype-specific high-mannose-type *N*-glycans

with or without a β-1,2-xylose residue attached to the trimannosyl core. Additionally, acidic

N-glycans of *C. neoformans* contain xylose phosphates attached to the mannose residues both

within the *N*-glycan core and outer mannose chains. The intact core *N*-glycan structure is

crucial for *C. neoformans* pathogenicity (Thak et al., 2020); hence, alterations in the *N*-glycan

structure modulates the interaction between the cell surface mannoproteins and host cells

(Lee et al., 2023). Additionally, *C. neoformans* lacks homologous genes to the Asn-Linked

Glycosylation (ALG) genes *ALG6*, *ALG8*, and *ALG10*, which are evolutionary conserved

across most eukaryotic organisms and encode the glucosyltransferases that add the glucose

residues to the core *N*-glycan before its attachment to proteins (Park et al., 2012).

$Man_7GlcNAc_2$ and $Man_8GlcNAc_2$ without glucose residues are primarily detected in Dol-PP-

linked glycans of *C. neoformans* (Samuelson et al., 2005). Moreover, the mature core *N*-

glycan structures assembled on the cell surface mannoproteins of *C. neoformans* are

primarily $Man_{6-7}GlcNAc_2$, which are shorter than the expected $Man_8GlcNac_2$ (Park et al.,

2012). This observation led to the speculation that the terminal α-1,2-mannose residues of *C.*

neoformans N-linked glycans may be more susceptible to trimming by ER α-1,2

mannosidases due to the lack of glucose residues compared to most eukaryotes. Alternatively,

the presence of multiple α-1,2 mannosidases may generate more extensively trimmed core *N*-

glycans in *C. neoformans*.

C. neoformans employs several virulence factors, including an extensive polysaccharide capsule composed of glucuronoxylomannan (GXM) and galactoxylomannan (GXMGal) (Doering et al., 2009), melanin (Qiu et al., 2012), and various extracellular enzymes such as phosphatase and urease (Singh et al., 2013), all of which contribute to immune evasion and enhance fungal pathogenicity. To facilitate the export of these virulence-associated molecules, C. neoformans employs both conventional (Yoneda and Doering, 2006; Panepinto et al., 2009) and unconventional secretion pathways (Rodrigues et al., 2007; Casadevall et al., 2019; Rizzo et al., 2021). Classical secretion relies on the ER-Golgi network, while non-classical mechanisms, such as extracellular vesicle (EV)-mediated transport, provide an alternative route for cargo delivery beyond the conventional secretory pathway. C. neoformans EVs have been described as a heterogeneous population of "virulence bags" containing numerous fungal survival and pathogenicity-associated cargo. Proteomic analyses of EV cargo have identified several cell surface glycoproteins, including members of the CDA family, well-known immunomodulators (Spetch et al., 2017; Rizzo et al., 2021). The ERQC system is essential for maintaining protein homeostasis by ensuring proper folding, glycosylation, and directed trafficking of secreted proteins to downstream compartments within the conventional secretory pathway. Defects in ERQC affect the fidelity of protein secretion not only by accumulation or degradation of misfolded proteins, leading to the impaired secretion of functional proteins, but also by the escape of misfolded forms, thus resulting in increased secretion of non-properly processed proteins (Marcus and Perlmutter, 2000; Chen et al., 2024). However, the potential of ERQC to modulate EV-mediated protein trafficking is not yet systematically investigated.

This study presents the first systematic analysis of N-glycan-dependent ERQC in C. neoformans using mutant strains lacking ERQC gene homologs. Our findings highlight the

121   critical roles of ERQC in maintaining cellular fitness and facilitating EV-mediated transport

122   of virulence factors.

Results

Evolutionary unique features of *C. neoformans* ERQC components

We performed BLAST analysis of the *C. neoformans* H99 genome to identify the ERQC pathway-associated homologous genes of *C. neoformans*, followed by comparison with the ERQC components in other eukaryotic organisms (Supplementary Fig. S1, A). Unlike most eukaryotes, several yeast species within the Ascomycota phylum, such as *Saccharomyces cerevisiae* and *Candida albicans,* do not possess a functional UGGT. In contrast, UGGT homologs were identified in most Basidiomycota fungal species, including *C. neoformans* (Supplementary Fig. S1, A and B). The *C. neoformans* UGGT (CNAG_03648), which has been named as Ugg1, consists of 1,582 amino acids (aa) and features a signal peptide (1–20 aa) along with four tandem-like thioredoxin-like (TRXL) domains: TRXL12 (33–320 aa), TRXL13 (28–416 aa), TRXL14 (432–618 aa), and TRXL15 (712–950 aa). Ugg1 also contains a glucosyltransferase (GT) 24 domain with a DXD motif (1,369–1,371 aa), and a KDEL-like ER retention signal (1579–1582 aa), which facilitates its retrieval from the Golgi apparatus (Fig. 1B, Supplementary Fig. S2, A).

Additionally, we identified two *C. neoformans* ORFs, Mns1 (CNAG_02081) and Mns101 (CNAG_03240), as homologs of the eukaryote α1,2-mannosidase I, which processes *N*-glycans before exporting them to the Golgi. The *C. neoformans* Mns1 and Mns101 show 42.4% and 28.8% amino acid identities to *S. cerevisiae* ER α1,2-mannosidase I, respectively, and share 30.9% identity between them. Furthermore, *C. neoformans* Mnl1 (CNAG_01987) and Mnl2 (CNAG_04498) were identified as putative components of ERQC in *C. neoformans* (Supplementary Fig. S1, A and C). *C. neoformans* Mnl1 is a homolog of the yeast α1,2-mannosidase-like protein Htm1, which processes *N*-glycans targeted for ERAD. In contrast, *C. neoformans* Mnl2 encodes a mannosidase that does not share significant

similarity with mannosidases from other eukaryotes. Mns1, Mns101, and Mnl1 possess

glucosyl hydrolase (GH) 47 domains, essential for mannosidase activity, whereas Mnl2

contains a GH92 domain, also associated with mannosidase activity (Fig. 1B). The Mns1 and

Mnl1 families are characterized by conserved cysteine (Cys) and alanine (Ala) residues

within their activity domains, as previously described (Jakob et al., 2001). These conserved

residues are present in *C. neoformans* Mns1, Mns101, and Mnl1 but are absent in Mnl2

(Supplementary Fig. S2, B). Notably, Mns101 and Mnl2 are Basidiomycota-specific proteins

(Supplementary Fig. S1, A) and appear to have diverged early from other fungal mannosidase

clades based on phylogenetic analysis (Supplementary Fig. S1, C). Interestingly, neither

Mnl1 nor Mnl2 contains canonical KDEL/HDEL-like ER retention signals. In *S. cerevisiae*,

the ER retention of Mnl1/Htm1 is mediated through its interaction with protein disulfide

isomerase Pdi1, which carries an HDEL sequence (Gauss et al., 2011). Similarly, *C.*

neoformans Mnl1 and Mnl2 may employ a non-canonical retention mechanism, likely

facilitated by interactions with other ER-resident proteins, to achieve ER localization.

Loss of *UGG1, MNS1*, and *MNS101* causes alteration of *N*-glycan profiles in *C.*

neoformans

We performed high-performance liquid chromatography (HPLC) analysis of the cell

wall mannoproteins (cwMPs) from both the wild-type (WT) and ERQC mutant strains to

investigate the ERQC malfunction-induced structural differences in *N*-glycans (Fig. 2). The

HPLC profiles of cwMPs from the WT strain showed an M8 peak as the major species. This

peak corresponded to *N*-glycans with 8 mannose residues (Fig. 2A, top). The glycan structure

at the M8 peak ($Man_8GlcNAc_2$) in the WT primarily corresponded to the $Man_7GlcNAc_2$ core

N-glycan with an additional mannose residue that is linked via an α1,6-linkage and added in

the Golgi apparatus (Park et al., 2012). In the *ugg1*Δ mutant, M8 was also the main *N*-glycan

species, but the pools of hypermannosylated *N*-glycans (larger than the M11 peak) were

markedly reduced (Fig. 2A, middle). The altered *N*-glycan profile of the *ugg1*Δ mutant was

restored to that of the WT after complementation with the WT *UGG1* gene (Fig. 2A, bottom).

The lectin blotting analysis using *Galanthus nivalis* agglutinin (GNA), which specifically

binds to terminal α1,2-, α1,3-, and α1,6-linked mannose residues, showed a distinctive

increase of glycoproteins with lower molecular weight in the secretory proteins in *ugg1*Δ

compared to those in WT (Fig. 2B). This observation aligns with the decrease of

hypermannosylated *N*-glycans in the *ugg1*Δ mutant. Overall, these results suggest that Ugg1

is involved in mediating the hypermannosylation of *N*-glycans in the Golgi apparatus.

We also assessed putative α1,2-mannosidase I genes for roles in protein

mannosylation. The *mns1*Δ mutant *N*-glycan profile showed a peak shift from an M8 to an

M9 form, which strongly indicates that *C. neoformans* Mns1 could be the primary ER α1,2-

mannosidase I (Fig. 2C, *mns1*Δ). Notably, the loss of Mns101, which is present only in

Basidiomycota, increased the fractions containing hypermannosylated glycans (> M10) while

maintaining M8 as a primary core *N*-glycan form. This suggests that the basidiomycete-

specific Mns101 may potentially be a novel α1,2-mannosidase that functions to remove

mannose residues from hypermannosylated *N*-glycans in the Golgi apparatus or further trims

the M8 glycan in the ER before the glycoproteins are transported to the Golgi (Fig. 2C,

*mns101*Δ). The matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) mass

spectrometer analysis of neutral *N*-glycans (Fig. 2C), further confirmed the shift of the major

M8 peak in the WT strain to M9 in the *mns1*Δ mutant and the increase in the

hypermannosylation profile in the *mns101*Δ mutant (Fig. 2D). The *N*-glycan profiles of the

*mns1*Δ*101*Δ double mutant showed the combined effect of each null mutation, and it showed

both increased hypermannosylated glycans and a shift from the M8 to the M9 peak in both

the HPLC (Fig. 2C, *mns1Δ101Δ*) and MALDI-TOF analyses. This indicates that Mns1 and

Mns101 serve as mannosidases and play independent roles at different stages of *N*-glycan

processing in *C. neoformans*.

In contrast, the loss of both *MNL1* and *MNL2* did not show notable differences in the

HPLC profile of *N*-glycans from cell surface mannoproteins (Supplementary Fig. S3, A).

Considering the expected function of Mnl1 and Mnl2 in ERAD, their substrates are likely

misfolded proteins that have not been transported to the Golgi apparatus. Thus, we speculate

that only the normally processed *N*-glycan profiles of cell surface mannoproteins in the

mnl1Δ mnl2Δ mutant strain were observed.

207   Loss of ERQC components results in defective growth fitness and increased stress

208   sensitivity

To investigate the changes in ERQC-related gene expression under stress conditions,

we conducted quantitative reverse transcription PCR (qRT-PCR) analysis of *C. neoformans*

following treatment with tunicamycin (TM, 5 µg/ml), a well-established inhibitor of *N*-

glycosylation, or dithiothreitol (DTT, 20 mM), which disrupts disulfide bond formation and

induces the accumulation of misfolded proteins. Additionally, we examined gene expression

after growth at 37 °C (Fig. 3A). When misfolded proteins accumulate in the ER, the unfolded

protein response (UPR) system is activated to restore homeostasis (Chakraborty et al., 2016).

As expected, *KAR2*, encoding a molecular chaperone associated with the UPR system, was

upregulated in response to TM treatment, DTT treatment, and heat stress at 37 °C. The ERQC

components *UGG1*, *MNS1*, *MNS101*, *MNL1*, and *MNL2* were upregulated by DTT treatment;

however, only *UGG1* was slightly induced following TM treatment. The high-temperature

conditions also induced the expression of *KAR2*, *UGG1* and *MNL2*. Notably, DTT treatment

induced a 10-fold higher expression of *MNS1* and *MNL1*. These results strongly suggest that

the *C. neoformans* genes *UGG1*, *MNS1*, *MNS101*, *MNL1*, and *MNL2* are crucial components

of the ERQC and ERAD systems, although each respond uniquely to various cell/ER stress

conditions.

We next investigated the growth of the *ugg1*Δ mutant under various stress-inducing

conditions to elucidate the roles of the ERQC components in ER stress response and

adaptation, which are closely associated with *C. neoformans* virulence. The *ugg1*Δ mutant

showed impaired growth even under normal growth conditions (Fig. 3B) and increased

sensitivity to ER stress-inducing agents such as DTT, TM and cell wall stressors, including as

calcofluor white (CFW), congo red (CR), sodium dodecyl sulfate (SDS), and caffeine.

Additionally, *ugg1*Δ showed hindered growth in the presence of antifungal drugs such as

azole agents (fluconazole and ketoconazole) and the glucose transport inhibitor fludioxonil.

These findings suggest that Ugg1 is crucial for the robust growth and survival of *C.*

neoformans under various stress conditions. In contrast, the single deletion (*mns1*Δ and

*mns101*Δ) and double deletion *(mns1*Δ*101*Δ) strains did not exhibit noticeable phenotypic

changes except for a slight increase in sensitivity of *mns101*Δ and *mns1*Δ*101*Δ to higher

fludioxonil concentrations (Supplementary Fig. S3, B). Similarly, neither the single nor

double disruption of *MNL1* and *MNL2* produced detectable changes in the tested growth

conditions (Supplementary Fig. S3, C).

Amino acid analogues have been used for evaluating the functionality of protein

folding, as its incorporation into newly synthesized proteins interrupts proper protein folding

(Trotter et al., 2002; Cowie et al., 1959). We hypothesized that, in the context of a defective

ERQC pathway, misfolded proteins cannot be adequately repaired and accumulate, thus

triggering ER stress, which may ultimately inhibit cell growth in the presence of amino acid

analogues. In the presence of the leucine analogue 5',5',5'-trifluoroleucine (TFL), *ugg1*Δ

showed noticeably inhibited growth, whereas the WT strain showed no growth inhibition (Fig.

3C). The *mns1Δ101Δ* strain also exhibited slightly increased sensitivity to TFL compared

with that of the WT strain.

As a defense mechanism against misfolded protein accumulation-mediated ER stress,

UPR is induced by the unconventional splicing of the *HXL1* transcription factor in *C.*

neoformans (Cheon et al., 2011). RT-PCR analysis revealed a significantly elevated level of

spliced *HXL1* in the *ugg1Δ* strain compared to the WT, even under normal culture conditions

(Fig. 3D), strongly indicating that the loss of *UGG1* leads to the accumulation of misfolded

proteins, triggering ER stress under standard growth conditions. Furthermore, green

fluorescence protein (GFP)-tagged Ugg1, Mns1, and Mns101 proteins colocalized with the

ER marker, supporting their role as confirming them to be functional ERQC components

based on their subcellular localization in the ER (Supplementary Fig. S4).

ERQC defects lead to virulence attenuation in *C. neoformans*

We investigated the effects of a defective ERQC on *in vitro* virulence phenotypes of *C.*

neoformans by analyzing capsule and melanin production, as these are two of the major

virulence factors of *C. neoformans*. Culturing on L-DOPA-containing plates showed that

melanin production was reduced in the *ugg1Δ* strain compared with that in the WT (Fig. 4A,

top), whereas the *mns1Δ101Δ* mutant cells did not show detectable defects at either 30 or

37 °C (Fig. 4A, bottom). Considering the decreased growth of the *ugg1Δ* strain, the melanin

production of *C. neoformans* cells cultivated in liquid L-DOPA was measured and normalized

by cell density, which further confirmed the decreased melanin production activity of the

ugg1Δ strain (Fig. 4B). India ink staining of the capsule showed significantly reduced capsule

thickness in the *ugg1Δ* strain and a moderate defect in the *mns1Δ101Δ* strain (Fig. 4C).

Defects in the ERQC system led to a more apparent decrease in pathogenicity in a

murine model of systemic cryptococcosis. Notably, the *ugg1Δ* strain was almost avirulent

with no detectable signs of illness in the infected animals (Fig 4D). Although *MNS1* and

MNS101 single disruptions did not cause a detectable decrease in *C. neoformans*

pathogenicity, their double disruption resulted in decreased virulence. Histopathological

analysis of infected mice lungs suggested significantly reduced lung colonization by both

ugg1Δ and *mns1Δ101Δ* cells (Supplementary Fig. S5, A). Additionally, the fungal burden of

ugg1Δ-infected animals showed a notable decrease in organ colonization at 60 days post-

infection (dpi) (Supplementary Fig. S5, B). Analysis of the fungal burden at 7 dpi showed

significantly reduced organ colonization of the *mns1Δ101Δ* mutant compared with that of the

WT (Supplementary Fig. S5, C). Consistent with the *in vivo* survival data, the number of

surviving *ugg1Δ* and *mns1Δ101Δ* cells within the macrophage-like cell line J447A.1 was

significantly lower compared with that of the WT strain (Fig. 4E). These data suggest that a

functional ERQC system is critical for maintaining full *in vivo* pathogenicity and ensuring

robust survival within host immune cells.

Loss of Ugg1 function manifests defects in extracellular transport of GXM

Cryptococcal capsule materials are synthesized intracellularly and secreted to the cell

surface. Subsequently, they are assembled and bound to the cell wall (Yoneda and Doering,

2006). The observed capsule defects in the *ugg1Δ* and *mns1Δ101Δ* mutant strains may be

attributed to issues in the capsule polysaccharide synthesis or transport steps. Alternatively,

defective capsule formation could arise from an increased shedding of capsule polymers,

preventing their proper attachment to the cell surface. To determine the cause of the capsule

defects, we examined the amounts of capsular polysaccharides synthesized intracellularly and

those shed into the culture supernatant by a capsule blotting assay, using the 18B7 antibody

directed against GXM, the major capsule component (Casadevall et al., 1998). As controls,

we also tested capsule production and secretion in the *cap59Δ* strain with a defect in GXM

synthesis (Grijpstra et al., 2009) and the *rim101Δ* strain with a capsule attachment defect

despite normal polysaccharide synthesis and secretion (O'Meara et al., 2010). Intracellular

GXM quantities did not notably differ between the WT, *ugg1Δ*, *mns1Δ101Δ*, and *rim101Δ*

strains, although *ugg1Δ* showed slightly reduced GXM level (Fig. 5A, left). In contrast, no

intracellular GXM was detected in *cap59Δ*, which was consistent with its defect in GXM

synthesis. Moreover, the *ugg1Δ* strain was defective in capsule polymer secretion/shedding,

whereas the *mns1Δ101Δ* strain displayed enhanced shedding of GXM-containing polymers

that appeared to be shorter in length than the WT-type polymers, suggestive of poor

polysaccharide polymerization (Fig. 5A, right). Therefore, these results suggest that *ugg1Δ*

contains a defect in polysaccharide trafficking to the extracellular space, whereas *mns1Δ101Δ*

has a defect in polysaccharide polymerization, resulting in defective capsule elaboration,

albeit with differential degrees of severity in both ERQC mutants.

Furthermore, to determine whether the ERQC mutants are also impaired in capsule

attachment, we performed a capsule transfer assay, to assess whether exogenously shed

capsule polysaccharides can bind to acapsular mutants (Reese and Doering, 2003). The

hypocapsular *rim101Δ* mutant did not show reassociation of capsular polysaccharides;

however, the acapsular *cap59Δ* mutant showed recovery of capsule formation when co-

incubated with exogenous GXM (Fig. 5B). Similarly, *ugg1Δ* reverted its acapsular phenotype

on incubation with the WT-shed polysaccharides, strongly suggesting that the defective

capsule phenotype of *ugg1Δ* is primarily due to defective GXM secretion rather than

impaired attachment of capsule polymers. We further conducted a capsule transfer assay

using capsular polysaccharides shed from *mns1Δ101Δ*. The *cap59Δ* mutant reverted to the

capsular phenotype by attaching to the *mns1Δ101Δ*-shed polymers. However, the

fluorescence intensity of the capsule generated was significantly lower than that of the cells

attached with polysaccharides obtained from the WT. Although the decreased length of

capsule polysaccharides should be validated by techniques specifically measuring GXM size

(De Jesus et al., 2010), this result suggests that the *mns1Δ101Δ* double mutant secretes

incomplete capsule polysaccharides (Fig. 5C), leading to a hypocapsular phenotype. These

results collectively indicate that disrupting the ERQC results in impaired capsule formation

by defective GXM trafficking to the extracellular space.

327   Transmission electron microscopy (TEM) imaging of the WT and mutant cells

distinctly showed a diminished capsule structure and loss of capsule shedding in the *ugg1Δ*

strain compared with that of the WT strain (Fig. 5D). Additionally, we observed considerable

thinning of the cell walls in both ERQC mutant cells. Notable changes in intracellular

structures along with an increase in the number of pigmented vesicles were observed in the

mutant cells even under YPD culture conditions. Under capsule-inducing conditions, a

significant accumulation of vesicular structures (electron-lucent structures) was observed

intracellularly, particularly in *ugg1Δ*. We performed lipid droplet (LD) staining to determine

whether these vesicular structures might be LDs. We observed a notable increase in LDs in

ugg1Δ under capsule-inducing culture conditions (Supplementary Fig. S6, A). LDs impact

proteostasis by sequestering misfolded proteins intended for degradation and providing an

"escape hatch" when the ERQC is overloaded (Ploegh, 2007; Vevea et al., 2015).

Additionally, FM4-64 dye staining showed increased number of vacuoles in the *ugg1Δ* cells

compared with that in the WT (Supplementary Fig. S6, B). Moreover, all the observed

abnormal phenotypes were less pronounced in the *mns1Δ101Δ* mutant than in the *ugg1Δ*

strain. Collectively, the altered vesicular structures observed in the *ugg1Δ* and *mns1Δ101Δ*

strains highlight the critical role of ERQC in maintaining proper vesicular architecture and

cell surface organization. These functions are tightly linked to extracellular trafficking of

virulence factors and cell wall remodeling.

ugg1Δ transcriptomic profiling shows induced ER and cell wall integrity stress

responses

To elucidate the mechanisms underlying *UGG1* deletion-induced physiological changes

in *C. neoformans*, we performed RNA sequencing (RNAseq)-based transcriptome analysis of

the WT and *ugg1Δ* strains under standard growth conditions (YPD medium, 30 °C).

Comparative transcriptome analysis showed statistically significant alterations exceeding 2-

fold in the expression patterns of 146 genes, of which 85 were upregulated and 61 were

downregulated compared with that of the WT strain (Fig. 6A, B, Supplementary Tables S2, A

and B).

Notably, the *ugg1Δ* mutant showed upregulation of genes such as *SKN1* and *KRE6*

(transmembrane glucosidases involved in the sphingolipid biosynthesis and β-glucan

biosynthesis), *CAT2* (a putative peroxisomal catalase), *CHS7* (chitin synthase export

chaperone), and CNAG_05458 (a putative endo-1,3(4)-β-glucanase). In contrast, genes

encoding *ERG3* and *ERG25* (associated with the ergosterol biosynthesis pathway) and *FHB1*

(flavohemoglobin denitrosylase associated with counteracting nitric oxide stress) were

significantly downregulated. Gene Ontology (GO) analysis in *ugg1Δ* showed pronounced

induction of genes implicated in various cellular processes, including the hydrolysis of *O*-

glycosyl compounds for cell wall remodeling, DNA replication, proteolysis, ribosome

biogenesis, protein folding, and serine/threonine kinase activity. In contrast, genes associated

with carbohydrate and lipid metabolism, chaperone binding, iron homeostasis, and

mitochondrial intermembrane space were considerably downregulated (Fig. 6C). The

transcriptome profile strongly suggests that loss of Ugg1 function induces the expression of

several genes involved in maintaining cell wall integrity to compensate for cell wall defects,

particularly those associated with chitin (*CDA2*, *CHS7*, *QRI1*, and CNAG_06898) and glucan

biosynthesis (*SKN1*, *KRE6*, *EBG1*, *LPI9*, CNAG_05458, and *BLG2*). Furthermore, genes

coding for ER chaperones that aid in protein folding (*ERO1*, *KAR2*, *LHS1*, and *PDI1*) and

chaperone regulator (*SCJ1*), and genes involved in proteolysis in the ER (CNAG_04635,

CNAG_06658) were induced, likely as part of the ER stress response induced by the

accumulation of misfolded proteins in the presence of a defective ERQC.

Further investigation of the effects of *UGG1* disruption in the expression of genes

involved in capsule biosynthesis, cell wall remodeling, and the secretion pathways of protein

virulence factors using qRT-PCR analysis (Fig. 6D) showed no noticeable changes in the

expression of genes related to capsule biosynthesis and conventional or unconventional

secretion pathways at mRNA level, despite the aberrantly displayed defective phenotypes.

EV-mediated protein trafficking and non-conventional secretion is impaired in the

***ugg1Δ* mutant strain**

Many enzymes contribute to the composite cryptococcal virulence phenotype. Some of

these virulence-associated enzymes are secreted through traditional secretion pathways,

whereas others are packaged into EVs and released into the extracellular milieu via non-

conventional secretion mechanisms (Amelida et al., 2015). The defects observed in the

ERQC mutant strains, particularly in the production of melanin and polysaccharide capsules,

both of which are transported by EVs, suggest a possible impairment in the EV-mediated

extracellular transport of virulence-associated enzymes in these mutants. Therefore, we

examined the *in vitro* activity of urease, and laccase and acid phosphatase in both intracellular

and extracellular fractions.

Comparative analysis of the WT and *ugg1Δ* strains showed minimal changes in the

intracellular activities of urease, laccase, and acid phosphatase. However, their activities were

significantly reduced in the extracellular fractions of the *ugg1Δ* cells (Fig. 7A, 7B, 7C, and

7D). Importantly, urease activity was performed in both solid and liquid media,

demonstrating a notable loss of secreted urease activity in *ugg1Δ* cells, with a slight decrease

in the *mns1Δ101Δ* double mutant cells (Fig. 7A and 7B). As urease lacks a signal peptide

required for conventional protein secretion, reduced urease secretion strongly indicates a

compromised EV-mediated non-conventional secretory pathway in the *ugg1Δ* mutant strain.

The decrease in extracellular laccase activity (Fig. 7C) corroborates our melanization assay

results, in which *ugg1Δ* showed defective melanin production (Fig. 4A).

We next examined cellulase and α-amylase activities to evaluate the impact of ERQC

disruption on the conventional secretion pathway of non-virulence-related enzymes (Fig. 7E

and 7F). The extracellular activities of these carbohydrate polymer-degrading enzymes in the

ugg1Δ mutant were slightly lower than those observed in the WT, likely reflecting reduced

intracellular enzyme production rather than a substantial impairment of secretion efficiency.

Additionally, we analyzed the localization of chitin deacetylase I (Cda1), a

glycosylphosphatidylinositol (GPI)-anchored protein involved in converting chitin to chitosan,

essential for maintaining cell wall integrity (Baker et al., 2007; Baker et al., 2011; Upadhya et

al., 2018; Upadhya et al., 2021). Cda1 was primarily detected in the insoluble cellular protein

fraction, encompassing the cell wall and membrane compartments (Fig. 7G, left), consistent

with its cell surface localization via a GPI anchor. Cda1 is secreted extracellularly upon

cleavage of the GPI anchor. Notably, the *ugg1Δ* mutant exhibited significantly higher

intracellular and extracellular Cda1 levels compared to the WT and *mns1Δ101Δ* strains (Fig.

7G, right), aligning with a 1.87-fold increase in *CDA1* mRNA expression observed in the

RNA-seq data. These findings indicate that the surface localization and secretion of Cda1

remain efficient, suggesting that the conventional secretion pathway is largely unperturbed in

ugg1Δ. Collectively, these results suggest that while Ugg1-mediated ERQC defects have a

pronounced negative effect on EV-mediated protein transport, their impact on the

conventional secretion pathway is minimal.

Extracellular vesicle biogenesis and cargo loading are defective in *ugg1Δ* strain

We thus analyzed the number and size distribution of EVs to determine possible

abnormal EV-mediated trafficking of virulence factors in the *ugg1Δ* mutant. Nanoparticle

tracking analysis (NTA) showed a major peak in size distribution at approximately 150

(134±28) nm in the WT strain, which was similar in size to mammalian exosomes and typical

microbial EVs. Additionally, a minor peak ranging from 300–500 nm was observed and

corresponded to microvesicles (Fig. 8A). Notably, the size distribution of *ugg1Δ* EVs was

more heterogeneous with smaller EVs ranging from 50–150 nm (82±16; 124±14) compared

with that of the WT. The *cap59Δ* EVs displayed a major distribution of approximately 150

nm (120±21). Microvesicles were barely detected in either *cap59Δ* or *ugg1Δ* strains.

We observed a significant reduction in the total number of secreted EVs in the *ugg1Δ*

mutant (approximately 40%), suggesting defective EV biogenesis and/or stability in the

absence of functional Ugg1 (Fig. 8B). Cryo-TEM analysis of EV morphology confirmed that

the sizes of EVs released by *ugg1Δ* were significantly smaller and more diverse than those of

the WT or the acapsular mutant (Fig. 8C), further supporting the NTA results. Examining EV

size distribution by measuring their diameter (Fig. 8D) further confirmed that although the

EVs in the WT strain ranged from approximately 50–550 nm in size with distribution

primarily concentrated at approximately 150 nm, the EVs in the *ugg1*Δ mutant were smaller

(< 100 nm) with heterogeneous sizes. Notably, *cap59*Δ EVs were uniformly distributed at

approximately 150 nm and were present at a higher concentration than those of EVs from the

WT. An increase in EV release and virulence factors has been also observed in acapsular

mutant strains of *C. neoformans* and *C. gattii* (Rodrigues et al., 2007; Reis et al., 2019),

supporting the notion that the capsule serves as a barrier in EV release in capsule forming

species.

To characterize the protein composition of EVs from WT and *ugg1*Δ strains, EV-

associated and whole-cell lysate (WCL) proteins were extracted and subjected to proteomic

analysis (Fig. 8E). In the WCL, 4,678 proteins were identified in the WT strain, with 333

exhibiting differential expression (>2-fold) in *ugg1*Δ versus the WT strain, including 123

upregulated and 210 downregulated proteins (Supplementary Fig. S7, A, left). In EVs, 2,075

proteins were detected in the WT strain, of which 693 were differentially expressed in EVs

from *ugg1*Δ relative to EVs from the WT strain (>2-fold), comprising 273 upregulated and

420 downregulated proteins (Supplementary Fig. S7, A, right; Supplementary Table S3).

Additionally, comparative proteomic analysis revealed 180 proteins uniquely enriched in EVs

(Supplementary Fig. S7, B). The GO enrichment analysis of differentially expressed proteins

(> 2-fold) showed distinctive patterns in biological process (BP) and cellular components

(CC) between EVs and the WCL (Supplementary Fig. S7, C). Three proteomic analyses of *C.*

neoformans EVs have been reported previously (Rodrigues et al., 2008; Wolf et al., 2014;

Rizzo et al., 2021), identifying 76, 202, and 1,847 proteins associated with EVs, respectively.

We compared the EV protein content between WT and *ugg1*Δ identified in our study,

focusing on overlapped proteins from the three EV proteomic data sets reported. Among 88

commonly reported EV-associated proteins, 29 were downregulated while 16 were

overexpressed in *ugg1Δ* EVs compared to WT EVs (Fig. 8E; Supplementary Table S4). The

differential expression pattern of the EV-associated proteins is quite distinctive from that of

WCL, revealing significantly detectable changes in EVs. Thus, the absence of functional

Ugg1 not only caused morphological alterations and reduced EV numbers but also altered EV

protein abundance in *C. neoformans*.

To further examine possible cargo loading defects, we investigated the presence of

GXM in the *ugg1Δ* EVs, comparing GXM quantities in the total cell extracts from WT,

ugg1Δ, and *cap59Δ* strains (Fig. 8F). Cda1 was used as the representative EV-associated

protein as its presence was reported within the *C. neoformans* EV membrane, despite a

substantial portion of Cda1 being secreted through the conventional secretion pathway (Rizzo

et al., 2021). GXM polysaccharides were not detected in the EVs released from *ugg1Δ*,

although it was present in the whole-cell extract. Taken together with the reduced quantity of

EV-associated proteins, the absence of GXM polysaccharides in *ugg1Δ* EVs strongly

indicates that EV cargo loading is defective when the ERQC is dysfunctional in *C.*

neoformans.

Discussion

As glycoproteins pass through the ER-Golgi secretory pathway, their *N*-glycans

undergo extensive modifications in association with protein quality control. The ERQC

system is highly conserved among eukaryotes but also diverged with distinct and species-

specific features. The ERQC composition of *C. neoformans* is unique in that it possesses

UGGT but lacks the three glucosyltransferases necessary for the addition of glucose residues

to the core precursor *N*-glycans (Park et al., 2012). Additionally, it lacks CRT, and carries

multiple α1,2-mannosidases. In this study, we investigated the molecular features and

functions of the *N*-glycan-dependent ERQC in *C. neoformans*, which generates unique *N*-

glycan precursors without glucose addition, and shorter in length than those of most

eukaryotes. Our data strongly suggest that despite the incomplete composition of the ERQC

components, the UGGT-centered ERQC plays pivotal roles in cellular fitness, and

particularly in the EV-mediated extracellular transport, which is crucial for pathogenicity.

The virulence of the *C. neoformans ugg1Δ* mutant was almost abolished in mice (Fig.

4D), a phenotype consistent with the observed defects in key virulence determinants, such as

the capsule and melanin, as well as poor growth at 37 °C (Fig. 3B; Fig. 4A and 4B). The

mns1Δ101Δ mutant displayed intermediate phenotypes between those of the WT and *ugg1Δ*.

Our data from the UPR induction analysis and comparative transcriptomic analysis strongly

indicate that the ERQC mutation generates ER stress; this accounts for significant similarity

of the defective phenotypes of the *C. neoformans* ERQC mutants to those reported in other

mutants that exhibit defective protein folding in the ER, particularly in terms of increased

stress sensitivity and decreased virulence. Connections between ER stress and

thermotolerance have previously been established in *C. neoformans*, as growth at 37 °C

requires key ER protein chaperones and protein processing machinery, including components

of the UPR signaling pathway Ire1 and the ER stress-responsive transcription factor Hxl1

(Cheon et al., 2011; Havel et al., 2011; Jung et al., 2013). Other mutations causing defects in

ER function in *C. neoformans*, such as mutants lacking *DNJ1* (an ER J-domain containing

co-chaperone) and *CNE1* (an ER chaperone), resulted in growth inhibition. The *dnj1Δ* mutant

strain displayed impaired elaboration of virulence factors, such as the exopolysaccharide

capsule and extracellular urease activity, when cultured at human body temperature

(Horianopoulos et al., 2021). Altogether with our data from the ERQC mutants, these

findings strongly support the notion that maintaining ER homeostasis is crucial for survival

and virulence factor production at elevated temperatures.

It is quite notable that the *C. neoformans ugg1Δ* EVs exhibited defective loading of

GXM and many protein cargo, alongside a decrease in the number and changes in the size

distribution (Fig. 8). Combined with compromised cell fitness, these defects in EV biogenesis

and cargo loading, even under normal growth conditions, likely contribute to the complete

loss of virulence owing to the defective export of virulence factors in the *ugg1Δ* mutant.

Interestingly, our comparative proteome analysis of culture supernatants (secretome) revealed

that the *ugg1Δ* strain exhibited more pronounced defects in the secretion of proteins lacking

signal peptides, which are often linked to unconventional secretion mechanisms, compared to

those with canonical secretion signals (Supplementary Fig. S8; Supplementary Table S5).

These findings further support the role of ERQC in modulating non-conventional secretion

pathways.

It is intriguing how the absence of Ugg1, leading to ERQC defects, results in

defective EV biogenesis and cargo loading in *C. neoformans*. In fungi, EV production and

release occur through the maturation of endosomes into multivesicular bodies (MVBs), which

are directed to the cell surface. Upon fusion with the plasma membrane, MVBs release

vesicles into the extracellular environment, functioning as an unconventional secretion

pathway (Oliveira et al., 2010). MVBs formation relies on the functionality of the endosomal

sorting complex required for transport (ESCRT), a highly intricate pathway involving a series

of finely regulated events (Henne et al., 2011). Deletion of several ESCRT complex-related

genes involved in EV transport directly impacts the cryptococcal capsule, notably resulting in

reduced capsule size. Specifically, the *C. neoformans vps27Δ* strain showed altered EV size

distribution, reduced capsule dimensions, defects in laccase export to the cell wall, and poor

extracellular export of urease (Park et al., 2020). Other regulators of unconventional secretion

are also linked to EV biogenesis. For example, the Golgi reassembly and stacking proteins

(GRASPs) regulate EV cargo and dimensions in *C. neoformans* (Peres et al., 2018). A *graspΔ*

mutant strain produced EVs with dimensions that significantly differed from those produced

by WT cells, along with attenuated virulence and abnormal RNA composition. Moreover, the

GRASP protein Grasp homology 1 (Grh1) serves as a chaperone that directly influences EV

cargo (Malhotra, 2013; Peres da Silva et al., 2018). Autophagy regulators, which participate

in EV formation in other eukaryotes, also play a role in cryptococcal EV formation. An *atg7Δ*

strain manifests hypovirulence, and EVs produced by this strain show slightly different RNA

composition compared with that of the WT cells (Oliveira et al., 2016). Notably, our data of

the transcriptome and proteome analysis did not show significant changes in the ESCRT

complex and GRASPs.

In addition to protein folding and secretion, the ER is crucial for regulating lipid

metabolism (Moncan et al., 2021). Consequently, ER stress significantly impacts lipid and

sterol synthesis, although some of these mechanisms are yet to be clarified. LDs aid the UPR

and ERAD in degrading misfolded proteins during ER stress in *S. cerevisiae* (Garcia et al.,

2021). In the present study, we observed a drastic increase in LDs in the *ugg1Δ* mutant

(Supplementary Fig. S6, A), consistent with the previous hypothesis that LDs help

maintaining ER homeostasis. Furthermore, we observed increased number of vacuoles in

ugg1Δ (Supplementary Fig. S6, B). This may thus indicate abnormal lipid homeostasis

caused by the ERQC defects, which could, in turn, affect EV biogenesis. The importance of

lipids and membrane regulators in proper EV formation and GXM export has been suggested

in a previous study on the Apt1 flippase in *C. neoformans* (Rizzo et al., 2018). Additionally,

ergosterol is essential for membrane fluidity, permeability, and protein transport (Ermakova

and Zuev, 2017). Erg6 is involved in the ergosterol biosynthesis pathway and was identified

as essential for the trans-Golgi network transport of proteins (Proszynski et al., 2005; Nes et

al., 2009). In *C. neoformans*, the *erg6Δ* mutant released EVs with a significantly larger

diameter than those of the WT, carrying increased levels of proteins and sterols, highlighting

the role of ergosterol in cryptococcal EV biogenesis (Oliveira et al., 2020). Notably, the

ugg1Δ mutant exhibited loss of microvesicles, which are derived from the plasma membrane

(Fig. 8A). Our preliminary analysis data on the surface lipid fraction of the *ugg1Δ* mutant

indicated detectable alteration of sphingolipids and sterol profiles. Altogether, these findings

suggest that ER stress caused by misfolded glycoprotein accumulation in ERQC-defective

mutants may alter lipid composition, which could affect EV biogenesis.

A recent study using kidney cells presented strong evidence for a key role of ER

stress in modulating EV biogenesis, demonstrating that ER stress decreases exosome

production in mice (Fukuoka et al., 2023). They showed that T-cadherin is downregulated by

ER stress through IRE1α activation at mRNA and protein levels and that ER stress decreases

EV production through adiponectin/T-cadherin-independent way, which may involve

interferon pathway activation in mice. We similarly observed induced activity of the ER

stress sensor Ire1 in *ugg1Δ*, even under normal growth conditions, suggesting a possible

association between the Ire1-mediated UPR pathway and EV biogenesis (Fig. 9). Additionally,

glycosylation regulates the biogenesis of small EVs and affects protein cargo loading

efficiency in melanoma cells (Harada et al., 2020). This suggests that the altered *N*-

glycosylation observed in *C. neoformans ugg1Δ* may influence cargo loading of certain EV-

targeting glycoproteins. Despite their findings not being conducted in fungi, they provide

valuable guides on the investigation of conserved mechanisms shared with *C. neoformans*.

Further studies to investigate the mechanisms underlying the ERQC-mediated modulation of

EV biogenesis and cargo loading in *C. neoformans* will provide insights into understanding

the regulation, production, composition, and diversity of fungal EVs, enabling a better

understanding of their biological function. Expanding our knowledge on pathogenic fungal

EVs would pave the way for utilizing native or engineered EVs as promising candidates for

therapeutic applications, including fungal infection diagnosis and vaccine development.

Materials and Methods

Strains, culture conditions, plasmids, and primers

The *C. neoformans* strains that were constructed and used in this study are listed in

Supplementary Table S1A. The plasmids and primers used are listed in Supplementary Tables

S1B and S1C, respectively. The construction of the deletion mutants is described in the

Supplementary Information. The strains were typically cultured in YPD medium (1% yeast

extract, 2% bacto peptone, and 2% glucose) at 30 °C with shaking (220 rpm). *C. neoformans*

transformants were selected by culturing on YPD solid medium containing 100 µg/ml

nouseothricin acetyltransferase (Jena Bioscience, Germany; indicated as YPD_{NAT}), YPD solid

medium with 100 µg/ml hygromycin B (Sigma-Aldrich, USA; indicated as YPD_{HyB}), or YPD

solid medium with 100 µg/ml G418 (Duchefa, Netherlands; indicated as YPD_{NEO}). For

capsule induction, *C. neoformans* cells were cultured in liquid Sabouraud dextrose medium

(Difco) at 30 °C for 16 h and incubated in 10% Sabouraud dextrose medium (pH 7.3)

supplemented with 50 mM morpholinepropanesulfonic acid (MOPS) at 30 °C for 2 days. For

melanin production analysis, the cells were spotted onto an L-DOPA agar plate (7.6 mM L-

asparagine monohydrate, 5.6 mM glucose, 22 mM KH_2PO_4, 1 mM $MgSO_4 \cdot 7H_2O$, 0.5 mM L-

DOPA, 0.3 mM thiamine-HCl, and 20 nM biotin) and incubated at 30 °C or 37 °C for 2 days.

Transmission electron microscopy

C. neoformans cells were cultured either at an initial optical density of 600 nm

(OD_{600}) of 0.2 at 30 °C in YPD medium until OD_{600} reached 0.8, or for 2 days in 10%

Sabouraud media at 30 °C. The cell pellets were washed twice in PBS and fixed for 12 h in 2%

glutaraldehyde-2% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4). The cells were

washed in 0.1 M phosphate buffer and post-fixed with 1% OsO_4 in 0.1 M phosphate buffer

for 2 h. The cells were dehydrated using an ascending ethanol series (50, 60, 70, 80, 90, 95,

100%) for 10 min each and infiltrated with propylene oxide for 10 min. The specimens were

embedded with a Poly/Bed 812 kit (Polysciences Inc., USA) and polymerized in an electron

microscope oven (TD-700, DOSAKA, Japan) at 65 °C for 12 h. The block was cut into 200-

nm semi-thin sections with a diamond knife in the ultramicrotome and stained with toluidine

blue for observation using optical microscopy. The region of interest was further cut into 80

nm thin sections using the ultramicrotome, placed on copper grids, double stained with 3%

uranyl acetate for 30 min and 3% lead citrate for 7 min, and imaged using a transmission

electron microscope (JEM-1011, JEOL, Tokyo, Japan) equipped with a Megaview III CCD

camera (Soft imaging system-Germany) at the acceleration voltage of 80 kV.

Capsule transfer and shedding analysis

Capsule transfer assays were performed as described previously (Reese and Doering,

2003). Briefly, conditioned medium (CM) was prepared as a source of GXM by culturing the

respective strains for 5 days in YPD medium, followed by filtering and storing the culture

supernatant at 4 °C. The acceptor strains were cultured overnight at 30 °C in YPD medium.

In total, 2×10^6 acceptor cells were incubated with 1 µl of CM for 1 h at room temperature

under rotation (18 rpm) and washed twice with PBS. Capsule acquisition was visualized by

incubating the cells with an anti-GXM (18B7) antibody conjugated with AlexaFluor 488

(ThermoFisher Scientific, USA) for 1 h at 37 °C, and observing under an Eclipse Ti-E

fluorescence microscope (Nikon, Japan), equipped with a Nikon DS-Qi2 camera. The images

were processed using the NIS-elements microscope imaging software (Nikon, Japan).

Capsule shedding analysis was performed using a modified previously described protocol

(Yoneda and Doering, 2008). The respective strains were cultured in 10% Sabouraud media

for 2 days, and the culture supernatant was sterile filtered. Enzyme denaturation was

performed by subjecting the filtrate to heating at 70 °C for 15 min, followed by centrifugation

at 13,000 rpm for 3 min. Intracellular GXM analysis was performed by resuspending the

pellets in TNE buffer (50 mM Tris-HCl [pH 7.5], 150 mM NaCl, 5 mM EDTA) with the same

volume of glass beads (425–600 μm in diameter, Sigma-Aldrich, USA). The cells were

disrupted four times for 15 s at 5,000 rpm using a Precellys 24 Tissue Homogenizer (Bertin

Technologies, France), followed by centrifugation at 16,000 rpm for 5 min at 4 °C. Next, 10

μl of the supernatant of either secreted or intracellular polysaccharide was mixed with 6X

loading dye and run on a 0.6% certified megabase agarose (Bio-rad, USA) gel in 0.5X TBE

(44.5 mM Trisma base, 44.5 mM boric acid, 1 mM EDTA [pH 8.0]) at 25 V for 16 h. The

polysaccharides were transferred onto a nylon membrane using the southern blotting

technique. The membrane was air-dried, blocked using 5% skim milk, and treated overnight

with 2 μg/ml 18B7 antibody. After washing, the membrane was incubated with an anti-mouse

peroxidase-conjugated   secondary   antibody   and   subjected   to   detection   using

chemiluminescence.

Detection of enzymatic activities in intracellular and secretary fractions

Biochemical enzymatic activities of acid phosphatase, urease, and laccase in the

intracellular and secreted fractions were assayed spectrophotometrically, using previously

described methods (Lev et al., 2014; Fu et al., 2018; de Sousa et al., 2022). Acid phosphatase

activity was assayed by culturing cells in MM-KCL medium (0.5% KCl, 15 mM glucose, 10

mM MgSO$_4$·7H$_2$O, 13 mM glycine, and 3 μM thiamine) for 3 h at 30 °C. The culture

supernatant and soluble cell lysate were allowed to react with 2.5 mM p-nitrophenyl

phosphate (pNPP) for 30 min at 37 °C. Urease activity was determined after culturing the

cells in Rapid Urea broth (RUH) for 24 h at 30 °C and subjecting the culture supernatant and

soluble cell lysate to incubation with phenol red. Laccase activity was assayed by culturing

the cells in asparagine salts media (7.6 mM L-asparagine, 0.1% glucose, 22 mM KH_2PO_4, 1

mM $MgSO_4.7H_2O$, 0.3 mM thiamine-HCl, and 20 nM biotin) for 48 h at 30 °C and allowing

the supernatant and soluble cell lysate to react overnight with 10 mM L-DOPA. The reactions

were quantified by measuring OD_{420} (acid phosphatase), OD_{570} (urease), or OD_{480} (laccase).

Cellulase activity was determined by culturing the cells in cellulose media (0.1% $NaNO_3$, 0.1%

K_2HPO_4, 0.1% KCl, 0.5% $MgSO_4$, 0.5% yeast extract, 0.1% glucose, and 0.5% low viscosity

carboxymethyl cellulose) for 24 h at 30 °C. The culture supernatants were concentrated using

an Amicon tube (30 kDa cutoff, Sigma-Aldrich, USA). The concentrated supernatants and

soluble cell lysates were allowed to react at 40 °C for 10 min with the substrate 4,6-O-(3-

ketobutylidene)-4-nitrophenyl-β-D-cellopentaoside (BPNPG5), which was provided in the

cellulase assay kit (CellG5 Method, Megazyme, Ireland). The reaction was terminated by

adding 2% [w/v] Tris buffer (pH 10), and the absorbance of 4-nitrophenol was measured at

400 nm. Cellulase activity (CellG5 Units/ml) was calculated as indicated in the cellulase

assay kit. α-amylase activity was measured using the α-Amylase Activity Colorimetric Assay

Kit (Biovision Technologies, USA) according to the manufacturer's instructions. Yeast cells

were cultured overnight in YPD medium at 30 °C, and the culture supernatants were

concentrated using an Amicon tube (30 kDa cutoff, Sigma-Aldrich, USA). The concentrated

supernatants and soluble cell lysates were incubated for 1 h at 25 °C in assay buffer with the

substrate ethylidene-pNP-G7, which was provided in the kit. OD_{405} was measured. All the

activity analysis results were normalized according to cell density (OD_{600}).

Purification, nanoparticle tracking analysis (NTA), and Cryo-TEM imaging of

extracellular vesicles

EV purification was performed according to a previously published protocol (Reis

et al., 2019) with modifications. One loop of cells was inoculated into 10 ml of liquid YPD

and incubated at 30 °C for 24 h with shaking (220 rpm). The cells were washed twice with

10 ml of PBS, counted, and diluted in PBS to a density of 3.5×10^7 cells/ml. Aliquots of cell

suspension (300 µl) were spread onto synthetic dextrose (SD) solid medium plates and

incubated for 24 h at 30 °C. The cells were carefully recovered from each plate using an

inoculation loop, gently resuspended in 30 ml PBS, and pelleted through centrifugation at

4,000 rpm for 10 min at 4 °C. The supernatant was collected and centrifuged again at 15,000

$\times g$ for 15 min at 4 °C. Then, the supernatant was filtered through 0.45-µm syringe filters and

ultracentrifuged at 100,000 $\times g$ for 1 h at 4 °C (MLA-50 fixed angle rotor, Beckman Coulter,

Germany). The supernatant was discarded, and the EV pellets were collected and

resuspended in 1 ml of PBS for immediate use or stored at -80 °C for further experiments.

The samples were diluted 100-fold, and EV sizes were measured using an NTA instrument

(NFEC-2024-03-295455, Nanosight Pro, Malvern Panalytical, Netherlands) coupled to a 532-

nm laser (Malvern Panalytical, Netherlands), SCMOS camera (Hamamatsu Photonics. Japan),

and syringe pump (Malvern Panalytical, Netherlands). The data were analyzed using the NS

Xplorer software (v1.1.0.6, Malvern Panalytical, Netherlands).

Cryo-TEM imaging was performed by loading purified EVs (3 µl) onto a lacey

carbon grid (Lacey Carbon, 300mesh Cu, Ted Pella Inc., USA), which was glow discharged

at 15 mA for 60 s. The sample-loaded grid was blotted for 3 s at 15 °C and 100% humidity

and immediately plunge-frozen in liquid ethane. The process was performed by Vitrobot

Maek IV (Thermofisher Scientific, USA, SNU, CMCI). The frozen grids were imaged using

TEM (JEM-2100F, JEOL, Japan); the temperature of the grid was maintained at

approximately -180 °C at an acceleration voltage of 200 keV. The images were recorded

709 using an ultrascan 1000 electron detector.

Proteomic analysis based on mass spectrometry and data processing

712   The samples for proteomic analysis were prepared as described in Supplementary

713 Information. The labeled peptides (total, 100 μg proteins) were combined prior to offline

714 basic reverse-phase liquid chromatographic (bRPLC) fractionation. Linear gradient was

715 performed using buffer A (10 mM TEAB in water) and buffer B (10 mM TEAB in 90%

716 acetonitrile), and 10 fractions were analyzed in total using am LC-MS/MS system. The

717 samples were dissolved in 0.1% formic acid using an UltiMate 3000 RSLCnano system and

718 analyzed using an Orbitrap Eclipse Tribrid mass spectrometer (Thermo Fisher Scientific,

719 USA). All MS raw files were converted into mzML and ms2 file formats using the

720 MSConvert (version 3.0.20033) software. The *C. neoformans* proteome was determined by

721 downloading a protein FASTA file from Uniprot (http://uniprot.org), which included 7,492

722 reviewed (Swiss-Prot) and unreviewed (TrEMBL) proteins entries. A proteome search

723 database with reversed sequence and contaminants in the Integrated Proteomics Pipeline (IP2,

724 version 5.1.2, Integrated Proteomics Applications Inc., San Diego, CA) was generated. The

725 proteome search from 20 ms2 files was performed with IP2 and its following parameters,

726 followed by evaluation for a false discovery rate (FDR) using IP2 and Proteininferencer

727 (version 1.0, Integrated Proteomics Applications Inc., USA). Protein quantification and

728 statistical analysis for discovery of differentially expressed proteins (DEPs) was performed

729 using the ms2 files with tandem mass tag (TMT) reporter ions using an in-house program

730 coded using Python 3.8, where t-test and Pearson's correlation analysis between comparison

731 samples was performed using the scikit-learn (version 0.23.2), Scipy (version 1.6.0), and

732 statsmodels (0.12.1) Python libraries.

Acknowledgments

735   We thank Arturo Casadevall for providing the 18B7 antibody, Jennifer K. Lodge for

736   providing the anti-Cda1 antibody, and Ji-Yeon Kang for technical assistance with MALDI-

737   TOF analysis.

Competing Interests

740   The authors declare no conflicts of interest.

Additional information

Funding

745   This study was supported by the National Research Foundation of Korea (Grant nos. NRF-

746   2022R1A2C1012699, NRF2018R1A5A1025077, and RS-2023-00212663) and by the Korea

747   Institute of Marine Science & Technology Promotion (Grant no. RS-2024-00405273).

Author Contributions

750   Catia Mota: Investigation, Construction of mutants, Performing all virulence analysis, Data

751   curation, and Writing—original draft; Kiseung Kim: Construction of mutants and Performing

752   phenotype analysis; Ye Ji Son: Transcriptome analysis and Validation; Eun Jung Thak and

753   Su-Bin Lee: Glycan analysis; Ju-El Kim and Jeong-Kee Yoon: NTA analysis; Min-Ho Kang:

754   Cryo-TEM analysis; Heeyoun Hwang: Proteome analysis; Yong-Sun Bahn and J. Andrew

Alspaugh: Supervision, Data curation, Writing—review, and editing; Hyun Ah Kang: Conceptualization, Project administration, Supervision, Data curation, Writing—original draft, review, and editing.

Author ORCIDs

Catia Mota: orcid.org/0009-0001-6129-0000

Ye Ji Son: orcid.org/0009-0006-4324-5800

Eun Jung Thak: orcid.org/0000-0003-4523-142X

Ju-El Kim: orcid.org/0009-0002-3028-1125

Jeong-Kee Yoon: orcid.org/0000-0003-0111-9911

Heeyoun Hwang: orcid.org/0000-0002-0124-2533

Yong-Sun Bahn: orcid.org/0000-0001-9573-5752

J. Andrew Alspaugh: orcid.org/0000-0003-3009-627X

Hyun Ah Kang: orcid.org/0000-0002-3722-525X

Additional files

Supplementary files

Supplementary file 1 – Supplementary material & methods. (a) Construction of *C. neoformans ugg1Δ* and *UGG1* complementation strains. (b) Construction of *C. neoformans mns1Δ*, *mns101Δ*, *mns1Δ101Δ*, and respective complementation strains. (c) Construction of *C. neoformans mnl1Δ*, *mnl2Δ* and, *mnl1Δmnl2Δ* strains. (d) Construction and localization analysis of *C. neoformans* GFP-Ugg1, Mns1-GFP, and Mns101-GFP fusion proteins. (e) HPLC and MALDI-TOF-based *N*-glycan structure analysis. (f) Animal study and *in vitro* survival analysis. (g) Subcellular fractionation and western blot analysis of Cda1. (h) RNA

preparation, qRT-PCR, and RNA-sequencing. (i) Sample preparation for proteomic analysis.

Supplementary file 2 - Supplementary figures. (a) Figure S1. Evolutionary conservation and

divergence of endoplasmic reticulum quality control (ERQC)-related gene homologues

predicted in different eukaryote organisms. (b) Figure S2. Domain analysis of putative *C.*

neoformans ERQC-associated proteins. (c) Figure S3. Phenotype characterization of *C.*

neoformans α1,2-mannosidases. (d) Figure S4. Subcellular localization of GFP-tagged Ugg1,

Mns1, and Mns101. (e) Figure S5. Fungal burden of mice infected with *C. neoformans*

ERQC-related mutant cells. (f) Figure S6. Lipid and vacuole staining of WT and *ugg1Δ*

strains. (g) Figure S7. Comparative analysis of whole-cell lysate (WCL) and EV proteomes

between WT and *ugg1Δ* strains. (h) Figure S8. Comparative analysis of whole-cell lysate

(WCL) proteome and secretome between WT and *ugg1Δ* strains.

Supplementary file 3 - Table S1. A. List of strains used in this study; B. List of plasmids

used in this study; C. List of primers used in this study.

Supplementary file 4 - Table S2. A. Up-regulated (> 2-fold) *Cryptococcus neoformans* genes

in *ugg1Δ* than that of wild type (WT) cultivated in yeast extract peptone dextrose (YPD) at

30 °C, B. Down-regulated (> 2-fold) *Cryptococcus neoformans* genes in *ugg1Δ* than that of

wild type (WT) cultivated in YPD at 30 °C.

Supplementary file 5 - Table S3. Excel file containing the list of proteins identified in the

proteomic analysis of the wild type (WT) and *ugg1Δ* extracellular vesicles (EVs) and whole-

cell lysates (WCL).

Supplementary file 6 - Table S4. Representative extracellular vesicle (EV)-associated

proteins commonly detected in this study and previously reported *Cryptococcus neoformans*

EV proteome data sets.

Supplementary file 7 - Table S5. Excel file containing the list of proteins identified in the

whole-cell lysate (WLC) proteome and secretome analysis in the wild type (WT) and *ugg1Δ*

strains.

Data availability

The raw RNA sequencing data have been submitted to the NCBI GEO database under

accession no. GSE254772.

References

Aebi M. 2013. *N*-linked protein glycosylation in the ER. *Biochim Biophys Acta*. **1833**: 2430-2437.

Balchin D, Hayer-Hartl M, Hartl FU. 2016. In vivo aspects of protein folding and quality control. *Science* **353**: aac4354.

Baker LG, Specht CA, Donlin MJ, Lodge JK. 2007. Chitosan, the deacetylated form of chitin, is necessary for cell wall integrity in *Cryptococcus neoformans*. *Eukaryot Cell* **6**: 855-867.

Baker LG, Specht CA, Lodge JK. 2011. Cell wall chitosan is necessary for virulence in the opportunistic pathogen *Cryptococcus neoformans*. *Eukaryot Cell* **10**: 1264-1268.

Banerjee S, Vishwanath P, Cui J, Kelleher DJ, Gilmore R, Robbins PW, Samuelson J**.** 2007. The evolution of *N*-glycan-dependent endoplasmic reticulum quality control factors for glycoprotein folding and degradation. *Proc Natl Acad Sci USA* **104**: 11676-11681.

Casadevall A, Coelho C, Cordero RJB, Dragotakes Q, Jung E, Vij R, Wear MP**.** 2019. The capsule of *Cryptococcus neoformans*. *Virulence* **10**: 822-831.

Casadevall A, Cleare W, Feldmesser M, Glatman-Freedman A, Goldman DL, Kozel TR, Lendvai N, Mukherjee J, Pirofski LA, Rivera J, Rosas AL, Scharff MD, Valadon P, Westin K, Zhong Z. 1998. Characterization of a murine monoclonal antibody to *Cryptococcus neoformans* polysaccharide that is a candidate for human therapeutic studies. *Antimicrob Agents Chemother*. **42**:1437-1446.

Chakraborty R, Baek JH, Bae EY, Kim WY, Lee SY, Kim MG. 2016. Comparison and contrast of plant, yeast, and mammalian ER stress and UPR. *Appl Biol Chem.* **59**: 337-347

Chen J, Wen P, Tang YH, Li H, Wang Z, Wang X, Zhou X, Gao XD, Fujita M, Yang G. 2024. Proteome and glycoproteome analyses reveal regulation of protein glycosylation site-specific occupancy and lysosomal hydrolase maturation by *N*-glycan-dependent ER-quality control. *J Proteome Res*. **23**: 4409-4421.

Cheon SA, Jung KW, Chen YL, Heitman J, Bahn YS, Kang HA**.** 2011. Unique evolution of the UPR pathway with a novel bZIP transcription factor, Hxl1, for controlling pathogenicity of *Cryptococcus neoformans*. *PLoS Pathog* **7**: e1002177.

Cowie DB, Cohen GN, Bolton ET, Robichon-Szulmajster H. 1959. Amino acid analog incorporation into bacterial proteins. *Biochim Biophys Acta*. **34**: 39-46.

De Jesus M, Chow SK, Cordero RJ, Frases S, Casadevall A. 2010. Galactoxylomannans from
Cryptococcus neoformans varieties *neoformans* and *grubii* are structurally and antigenically
variable. *Eukaryot Cell*. **9**: 1018-1028.

de Sousa HR, de Oliveira GP Jr, Frazão SO, Gorgonha KCM, Rosa CP, Garcez EM, Lucas J
Jr, Correia AF, de Freitas WF, Borges HM, Brito Alves LG, Paes HC, Trilles L, Lazera MDS,
Teixeira MM, Pinto VL Jr, Felipe MSS, Casadevall A, Silva-Pereira I, Albuquerque P, Nicola
AM. 2022. Faster *Cryptococcus* melanization increases virulence in experimental and human
cryptococcosis. *J Fungi* **8**: 393.
859
Doering TL. 2009. How sweet it is! Cell wall biogenesis and polysaccharide capsule
formation in *Cryptococcus neoformans*. *Annu. Rev. Microbiol* **63**: 223-247.

Fu MS, Coelho C, De Leon-Rodriguez CM, Rossi DCP, Camacho E, Jung EH, Kulkarni M,
Casadevall A. 2018. *Cryptococcus neoformans* urease affects the outcome of intracellular
pathogenesis by modulating phagolysosomal pH. *PLoS Pathog* **14**: e1007144.

Fukuoka K, Mineo R, Kita S, Fukuda S, Okita T, Kawada-Horitani E, Iioka M, Fujii K,
Kawada K, Fukushima Y, Nishizawa H, Maeda N, Shimomura L. 2023. ER stress decreases
exosome production through adiponectin/T-cadherin-dependent and -independent pathways. *J
Biol Chem* **299**: 105-114.

Garcia EJ, Liao PC, Tan G, Vevea JD, Sing CN, Tsang CA, McCaffery JM, Boldogh IR, Pon
LA. 2021. Membrane dynamics and protein targets of lipid droplet microautophagy during
ER stress-induced proteostasis in the budding yeast, *Saccharomyces cerevisiae*. *Autophagy* **17**:
2363-2383.

Gauss R, Kanehara K, Carvalho P, Ng DT, Aebi M. 2011. A complex of Pdi1p and the
mannosidase Htm1p initiates clearance of unfolded glycoproteins from the endoplasmic
reticulum. *Mol Cell*. **42**: 782-793.

Grijpstra J, Tefsen B, van Die I, de Cock H. 2009. The *Cryptococcus neoformans* cap10 and
cap59 mutant strains, affected in glucuronoxylomannan synthesis, differentially activate
human dendritic cells. *FEMS Immunol Med Microbiol* **57**: 142-150.

Gottfredsson M, Perfect JR. 2000. Fungal Meningitis. *Semin. Neurol* **20**: 307-322.

Harada Y, Nakajima K, Suzuki T, Fukushige T, Kondo K, Seino J, Ohkawa Y, Suzuki T, Inoue
H, Kanekuta T, Dohmae N, Taniguchi N, Maruyama I. 2020. Glycometabolic regulation of
the biogenesis of small extracellular vesicles. *Cell Rep* **33**: 108261.

Havel VE, Wool NK, Ayad D, Downey KM, Wilson CF, Larsen P, Djordjevic JT, Panepinto
JC. 2011. Ccr4 promotes resolution of the endoplasmic reticulum stress response during host
temperature adaptation in *Cryptococcus neoformans*. *Eukaryot Cell* **10**: 895–901.

Henne WM, Buchkovich NJ, Emr SD. 2011. The ESCRT pathway. *Dev Cell* **21**: 77-91.

Horianopoulos LC, Lee CWJ, Hu G, Caza M, Kronstad JW. 2021. Dnj1 promotes virulence
in *Cryptococcus neoformans* by maintaining robust endoplasmic reticulum homeostasis under
temperature stress. *Front Microbiol* **12**: 727039.

Jakob CA, Bodmer D, Spirig U, Battig P, Marcil A, Dignard D, Bergeron JJ, Thomas DY,
Aebi M. 2001. Htm1p, a mannosidase-like protein, is involved in glycoprotein degradation in
yeast. *EMBO Rep.* **2**: 423-430

Jung KW, Kang HA, Bahn YS. 2013. Essential roles of the Kar2/BiP molecular chaperone
downstream of the UPR pathway in *Cryptococcus neoformans*. *PLoS One* **8**: e58956.

Kwon-Chung KJ, Sorrel TC, Dromer F, Fung E, Levitz SM. 2000. Cryptococcosis: clinical
and biological aspects. *Med. Mycol* **38**: 205-213

Lee SB, Mota C, Thak EJ, Kim J, Son YJ, Oh DB, Kang HA. 2023. Effects of altered *N*-
glycan structures of *Cryptococcus neoformans* mannoproteins, MP98 (Cda2) and MP84
(Cda3), on interaction with host cells. *Sci Rep* **13**: 1175.

Lev S, Crossett B, Cha SY, Desmarini D, Li C, Chayakulkeeree M, Wilson CF, Williamson
PR, Sorrel TC, Djordjevic JT. 2014. Identification of Aph1, a phosphate-regulated, secreted,
and vacuolar acid phosphatase in *Cryptococcus neoformans*. *mBio* **5**: e01649-14.

Malhotra V. 2013. Unconventional protein secretion: an evolving mechanism. *EMBO J* **32**:
1660-1664.

Marcus NY, Perlmutter DH. 2000. Glucosidase and mannosidase inhibitors mediate increased
secretion of mutant alpha1 antitrypsin Z. *J Biol Chem.* **275**:1987-92.

Moncan M, Mnich K, Blomme A, Almanza A, Samali A, Gorman AM. 2021. Regulation of
lipid metabolism by the unfolded protein response. *J Cell Mol Med.* **25**: 1359-1370.

Nes WD, Zhou W, Ganapathy K, Liu JL, Vatsyayan R, Chamala S, Hernandez K, Miranda M.
2009. Sterol 24-C-methyltransferase: An enzymatic target for the disruption of ergosterol
biosynthesis and homeostasis in *Cryptococcus neoformans*. *Arch. Biochem. Biophys* **481**:
210-218.

Oliveira DL, Nakayasu ES, Joffe LS, Guimarães AJ, Sobreira TJP, Nosanchuk JD, Cordero
RJB, Frases S, Casadevall A, Almeida IC, Nimrichter L, Rodrigues ML. 2010.
Characterization of yeast extracellular vesicles: evidence for the participation of different
pathways of cellular traffic in vesicle biogenesis. *PLoS ONE* **5**: e11113.

Oliveira DL, Fonseca FL, Zamith-Miranda D, Nimrichter L, Rodrigues J, Pereira MD,
Reuwsaar JC, Schrank A, Staats C, Kmetzsch L, Vainstein MH, Rodrigues ML. 2016. The
putative autophagy regulator Atg7 affects the physiology and pathogenic mechanisms of
Cryptococcus neoformans. *Future Microbiol* **11**: 1405-1419.

Oliveira FFM, Paes HC, Peconick LDF, Fonseca FL, Marina CLF, Bocca AL, Homem-de-
Mello M, Rodrigues ML, Albuquerque P, Nicola AM, Alspaugh JA, Felipe MSS, Fernandes L.
2020. Erg6 affects membrane composition and virulence of the human fungal pathogen
Cryptococcus neoformans. Fungal Genet. Biol **140**: e103368.

O'Meara TR, Norton D, Price MS, Hay C, Clements MF, Nichols CB, Alspaugh JA. 2010.
Interaction of *Cryptococcus neoformans* Rim101 and protein kinase A regulates capsule.
PLoS Pathogens **6**: e1000776.

Park JN, Lee DJ, Kwon O, Oh DB, Bahn YS, Kang HA. 2012. Unraveling unique structure
and biosynthesis pathway of *N*-linked glycans in human fungal pathogen *Cryptococcus*
neoformans by glycomics analysis. *J Biol Chem* **287**: 19501–19515.

Park YD, Chen SH, Camacho E, Casadevall A, Williamson PR. 2020. Role of the ESCRT
pathway in laccase trafficking and virulence of *Cryptococcus neoformans*. *Infect Immun* **88**:
e00954-19.

Peres da Silva R, Martins SDT, Rizzo J, Dos Reis FCG, Joffe LS, Vainstein M, Kmetzsch L,
Oliveira DL, Puccia R, Goldenberg S, Rodrigues ML, Alves LR. 2018. Golgi reassembly and
stacking protein (GRASP) participates in vesicle-mediated RNA export in *Cryptococcus*
neoformans. *Genes* **9**: 400.

Ploegh HL. 2007. A lipid-based model for the creation of an escape hatch from the
endoplasmic reticulum. *Nature* **448**: 435–438.

Proszynski TJ, Klemm RW, Gravert M, Hsu PP, Gloor Y, Wagner J, Kozak K, Grabner H,
Walzer K, Bagnat M, Simons K, Walch-Solimena C. 2005. A genome-wide visual screen
reveals a role for sphingolipids and ergosterol in cell surface delivery in yeast. *Proc. Natl.*
Acad. Scie. U.S.A **102**: 32201128.

Qiu Y, Davis M.J, Dayrit JK, Hadd Z, Meister DL, Osterholzer JJ, Williamson PR, Olszewski
MA. 2012. Immune modulation mediated by cryptococcal laccase promotes pulmonary
growth and brain dissemination of virulent *Cryptococcus neoformans* in mice. *PLoS ONE* **7**:
e47853.

Rapoport TA. 2007. Protein translocation across the eukaryotic endoplasmic reticulum and
bacterial plasma membranes. *Nature* **450**: 663-669.

Reis FCG, Borges BS, Jozefowicz LJ, Sena BAG, Garcia AWA, Medeiros LC, Martins ST,
Honorato L, Schrank A, Vainstein MH, Kmetzsch L, Nimrichter L, Alves LR, Staats CC,
Rodrigues ML. 2019. A novel protocol for the isolation of fungal extracellular vesicles
reveals the participation of a putative scramblase in polysaccharide export and capsule
construction in *Cryptococcus gattii*. *mSphere* **4**: e00080-19.

Rizzo J, Wong SSW, Gazi AD, Moyrand F, Chaze T, Commere PH, Novault S, Matondo M,
Péhau-Arnaudet G, Reis FCG, Vos M, Alves LR, May RC, Nimrichter L, Rodrigues ML,

Aimanianda V, Janbon G. 2021. *Cryptococcus* extracellular vesicles properties and their use
as vaccine platforms. *J Extracell Vesicles* **10**: e12129.

Rizzo J, Colombo AC, Zamith-Miranda D, Silva VKA, Allegood JC, Casadevall A, Del Poeta
M, Nosanchuk JD, Kronstad JW, Rodrigues ML. 2018. The putative flippase Apt1 is required
for intracellular membrane architecture and biosynthesis of polysaccharide and lipids in
Cryptococcus neoformans. Biochim Biophys Acta Mol Cell Res **1865**: 532-541.
966
967 Reese AJ, Doering TL. 2003. Cell wall alpha-1,3-glucan is required to anchor
968 the *Cryptococcus neoformans* capsule. *Mol. Microbiol.* **50**: 1401–1409.
969
970 Rodrigues ML, Nimrichter L, Oliveira DL, Frases S, Miranda K, Zaragoza O, Alvarez M,
971 Nakouzi A, Feldmesser M, Casadevall A. 2007. Vesicular polysaccharide export
972 in *Cryptococcus neoformans* is a eukaryotic solution to the problem of fungal trans□cell wall
973 transport. *Eukaryot Cell* **6**: 48–59.
974
975 Rodrigues ML, Nakayasu ES, Oliveira DL, Nimrichter L, Nosanchuk JD, Almeida IC,
976 Casadevall A. 2008. Extracellular vesicles produced by *Cryptococcus neoformans* contain
977 protein components associated with virulence. *Eukaryot Cell* **7**: 58–67.
978
979 Samuelson J, Banerjee S, Magnelli P, Cui J, Kelleher DJ, Gilmore R, Robbins PW. 2005. The
980 diversity of dolichol-linked precursors to Asn-linked glycans likely results from secondary
981 loss of sets of glycosyltransferases. *Proc Natl Acad Sci USA* **102**: 1548-1553.
982
983 Samuelson J, Robbins PW. 2015. Effects of *N*-glycan precursor length diversity on quality
984 control of protein folding and on protein glycosylation. *Semin Cell Dev Biol.* **41**: 121-128.
985
986 Singh A, Panting RJ, Varma A, Saijo T, Waldron KJ, Jong A, Ngamskulrungroj P, Chang YC,
987 Rutherford JC, Kwon-Chung KJ. 2013. Factors required for activation of urease as a
988 virulence determinant in *Cryptococcus neoformans. mBio* **4**: e00220-13.
989
990 Thak EJ, Kim J, Lee DJ, Kim JY, Kang HA. 2018. Structural analysis of *N*-/*O*-glycans
991 assembled on proteins in yeasts. *J Microbiol* **56**: 11–23.
992
993 Thak EJ, Lee SB, Xu-Vanpala S, Lee DJ, Chung SY, Bahn YS, Oh DB, Shinohara ML, Kang
994 HA. 2020. Core *N*-glycan structures are critical for the pathogenicity of *Cryptococcus*
995 *neoformans* by modulating host cell death. *mBio* **11**: e00711-00720.
996
997 Thak EJ, Son YJ, Lee DJ, Kim H, Kim JH, Lee SB, Jang YB, Bahn YS, Nichols CB,
998 Alspaugh JA, Kang HA. 2022. Extension of *O*-linked mannosylation in the Golgi apparatus is
999 critical for cell wall integrity signaling and interaction with host cells in *Cryptococcus*
1000 *neoformans* pathogenesis. *mBio* **13**: e0211222.
1001

1002 Thibault G, Ng DT. 2012. The endoplasmic reticulum-associated degradation pathways of
1003 budding yeast. *Cold Spring Harb Perspect Biol.* **4**: a013193.

1004

1005 Trotter EW, Kao CM, Berenfeld L, Botstein D, Petsko GA, Gray JV. 2002. Misfolded
1006 proteins are competent to mediate a subset of the responses to heat shock in *Saccharomyces*
1007 *cerevisiae*. *J Biol Chem*. **277**: 44817-44825.

1008

1009 Upadhya R, Baker LG, Lam WC, Specht CA, Donlin MJ, Lodge JK. 2018. *Cryptococcus*
1010 *neoformans* Cda1 and its chitin deacetylase activity are required for fungal pathogenesis.
1011 *mBio*. **9**: e02087-18.

1012

Upadhya R, Lam WC, Hole CR, Parchment D, Lee CK, Specht CA, Levitz SM, Lodge JK.
2021. *Cryptococcus neoformans* Cda1 and Cda2 coordinate deacetylation of chitin during
infection to control fungal virulence. *Cell Surf* **7**: 100066.

Varki A. (2017) Biological roles of glycans. *Glycobiology*. **27**:3-49.

Vevea JD, Garcia EJ, Chan RB, Zhou B, Schultz M, Di Paolo G, McCaffery JM, Pon LA.
2015. Role for lipid droplet biogenesis and microlipophagy in adaptation to lipid imbalance
in yeast. *Dev Cell* **35**: 584–599.

Wolf JM, Espadas-Moreno J, Luque-Garcia JL, Casadevall A. 2014. Interaction of
Cryptococcus neoformans extracellular vesicles with the cell wall. *Eukaryot Cell* **13**: 1484-
1493.

1023

Yoneda A, Doering TL. 2006. A eukaryotic capsular polysaccharide is synthesized
intracellularly and secreted via exocytosis. *Mol Biol Cell* **17**: 5131–5140.

Yoneda A, Doering TL. 2008. Regulation of *Cryptococcus neoformans* capsule size is
mediated at the polymer level. *Eukaryot Cell* **7**: 546–549.

Xu C, Ng DT. 2015. Glycosylation-directed quality control of protein folding. *Nat Rev Mol*
Cell Biol **16**: 742-752.

 Figure legends

 Figure 1. **Presence of UDP-glucose:glycoprotein glucosyltransferase (UGGT) and α1,2-**

 mannosidases as endoplasmic reticulum protein quality control (ERQC) components in

 Cryptococcus neoformans. (A) Schematic representation of the ERQC pathway in *Homo*

 sapiens, Saccharomyces cerevisiae, and *Cryptococcus neoformans*. In mammals, the

 oligosaccharyltransferase (OST) complex attaches $Glc_3Man_9GlcNAc_2$ to nascent

 polypeptides, followed by glucose trimming by glucosidases I and II (GLS1/GLS2). This

 generates $Glc_1Man_9GlcNAc_2$, which binds to calnexin (CNX) or calreticulin (CRT) for

 folding. UGGT reglucosylates misfolded proteins, allowing refolding, while properly folded

 proteins undergo mannose trimming by ERMan I before Golgi transport. In contrast, fungal

 ERQC systems differ in key components. *S. cerevisiae* lacks UGGT, relying instead on

 Gls1/Gls2 and calnexin (Cne1). *C. neoformans* possesses UGGT but lacks ER

 glucosyltransferases (Alg6, Alg8 and Alg19) and CRT, resulting in a distinct ERQC system.

 (B) Domain structures of proteins encoded by *C. neoformans UGG1* (CNAG_03648), *MNS1*

 (CNAG_02081), *MNS101* (CNAG_03240), *MNL1* (CNAG_01987), and *MNL2*

 (CNAG_04498).

 Figure 2. *N*-glycan profile analysis of *C. neoformans* **ERQC mutant strains.** The HPLC

 and MALDI-TOF-based *N*-glycan structure analysis were carried out as described in

 Supplementary information. (A) HPLC-based analysis of *N*-glycan profiles of the *ugg1*Δ

 mutant. (B) Lectin blotting of sodium dodecyl sulphate (SDS)-polyacrylamide gels

 containing intracellular or secreted proteins into the culture supernatants of the wild type

 (WT) and *ugg1*Δ strains. Yeast cells were cultivated in YPD medium for 24 h, harvested, and

 subjected to sample preparation of soluble intracellular proteins and secreted proteins. The

proteins (30 μg) were loaded on 15% SDS-polyacrylamide gel and analyzed using silver

staining (left) or blotting (right) with *Galanthus nivalis* agglutinin (Roche), which recognizes

terminal α1,2-, α1,3-, and α1,6-linked mannose residues. (C) HPLC analysis of total *N*-glycan

profiles of *mns1Δ*, *mns101Δ*, and *mns1Δ101Δ* mutants. (D) MALDI-TOF profiles of neutral

N-glycans of *mns1Δ*, *mns101Δ*, and *mns1Δ101Δ* mutants. The *N*-glycans of cell wall

mannoproteins from *C. neoformans* cells were AA-labelled and analyzed using HPLC. For

MALDI-TOF analysis, neutral *N*-glycan fractions were obtained from the HPLC fractionation

of total *N*-glycans.

1062

Figure 3. Growth phenotype of *C. neoformans* ERQC mutant strains. (A) Expression

analysis of ERQC genes in *C. neoformans*. Yeast cells were cultured in YPD medium to a

mid-logarithmic phase and exposed to dithiothreitol (DTT; 20 mM), tunicamycin (TM; 5

μg/ml) or cultured at 37 °C for 1 h. The relative transcript levels of *C. neoformans* genes

were analyzed using qRT-PCR and normalized with that of *ACT1*. Error bars represent

standard deviation of duplicated assays. All statistical data were determined based on one-

way ANOVA and Dunnett's post-hoc test. *** $P < 0.0005$, ** $P < 0.003$, *** $P < 0.005$, * $P <$

0.05. (B) Spotting analysis of *C. neoformans* *ugg1Δ* mutant strains under various stress

conditions such as heat stress (37 °C and 39 °C), ER stress (DTT and TM), cell-wall stress

(CFW: calcofluor white, CR: Congo red, SDS: Sodium dodecyl sulfate, caffeine), osmotic

stress (NaCl, KCl, sorbitol) and treatment with antifungal drugs (fluconazole, ketoconazole,

fludioxonil). (C) Growth analysis in the presence of 5',5',5'-trifluoroleucine (TFL).

Respective strains were spotted on SC-Leucine media with or without TFL supplementation.

Plates were incubated for 3 days at 30 °C. (D) RT-PCR analysis of *IRE1*-dependent splicing

of *HXL1*. Strains were cultured in YPD supplemented with 5 μg/ml TM.

Figure 4. *In vitro* and *in vivo* virulence-associated phenotypes of *C. neoformans UGG1,*

***MNS1*, and *MNS101* mutant strains.** (A) Melanin synthesis analysis on L-DOPA plates. WT,

ugg1Δ, ugg1Δ::UGG1, mns1Δ, mns101Δ, mns1Δ101Δ, mns1Δ101Δ::MNS1,

mns1Δ101Δ::MNS101, and *cac1Δ* (negative control) strains were serially diluted, plated on

L-DOPA plates, and incubated at 30 °C and 37 °C. (B) Melanin synthesis activity per cell of

the *ugg1Δ* mutant. WT, *ugg1Δ, ugg1Δ::UGG1 w*ere cultured in liquid L-DOPA medium. The

amount of melanin in the culture supernatant was measured and normalized by cell density.

(C) Capsule formation. Cells were cultured for 2 days in 10% Sabouraud media at 30 °C and

observed under the microscope. Statistical significance: ****, $P < 0.0001$, ns, not significant).

(D) *In vivo* virulence analysis. A/Jcr mice (n=8) were infected with 10^5 cells of WT, *ugg1Δ,*

and *ugg1Δ::UGG1, mns1Δ, mns101Δ, mns1Δ101Δ, mns1Δ101Δ::MNS1*, and

mns1Δ101Δ::MNS101 strains, and survival was monitored for 2 months, as described in the

Supplementary Information. (E) Survival of *C. neoformans* in macrophages. Survival of *C.*

neoformans cells within the J774A.1 macrophage-like cell line was determined by counting

colony formation unit (CFU) obtained from lysed macrophages from two biologically

independent experiment sets. **** $P < 0.0001$, *** $P < 0.0005$, * $P < 0.05$, ns, not significant.

All statistical data were determined based on one-way ANOVA and Dunnett's post-hoc test.

1095

Figure 5. Capsule shedding and transfer analysis of *C. neoformans UGG1, MNS1*, and

***MNS101* mutant strains.** (A) Capsule shedding analysis. The presence of intracellular (left)

and shed (right) glucuronoxylomannan (GXM) was assessed by blotting a cell culture filtrate

using the monoclonal antibody 18B7. The arrow indicates the direction of electrophoresis. (B)

Capsule transfer analysis using exogenous capsule material from WT. The capsule transfer

assay was performed using the indicated strains as acceptors. Surface capsules were probed

using the anti-GXM antibody 18B7 conjugated with AlexaFluor 488. Quantitative

measurement of fluorescence intensity was calculated based on independent triplicate

experiments with standard deviations presented as error bars. (C) Capsule transfer analysis

using exogenous capsule material from WT or *mns1Δ101Δ*. Statistical significance: **** *P <*

0.0001. All statistical data were determined based on one-way ANOVA and Dunnett's post-

hoc test. (D) Transmission electron microscopy (TEM) of *C. neoformans* WT, *ugg1Δ*, and

mns1Δ101Δ strains. Yeast cells were grown overnight at 30 °C in YPD medium and fixed in 2%

glutaraldehyde and 2% paraformaldehyde. A Zeiss Axioscope (A1) equipped with an

AxioCan MRm digital camera was used to visualize India ink-stained *C. neoformans* cells.

Specimens were prepared using critical point drying prior to TEM microscopy. Capsule and

yeast cell body diameters were measured using ImageJ (National Institute of Health).

1113

Figure 6. Transcriptome analysis of *C. neoformans* WT and *ugg1Δ* cells. (A) Volcano plot

comparing a 2-fold differential gene expression between *ugg1Δ* and WT strains under normal

growth conditions. (B) Number of genes upregulated and downregulated by ≥ 2-fold in

ugg1Δ compared with that of the WT. (C) Gene ontology (GO) analysis of differentially

expressed genes between WT and *ugg1Δ* strains. Significantly upregulated genes in *ugg1Δ*

are shown in red, whereas significantly downregulated genes in *ugg1Δ* are shown in blue.

Total RNA was extracted and subjected to RNAseq analysis as described in the

Supplementary Information. (D) qRT-PCR analysis of mRNA expression levels of a set of

genes responsible for capsule biosynthesis, cell wall remodeling, and both conventional and

non-conventional secretion in *ugg1Δ* vs WT under normal growth conditions from three

biologically independent experiment sets.

1125

Figure 7. Analysis of protein secretion in *C. neoformans UGG1*, *MNS1*, and *MNS101*

mutant strains. (A) Spot assay for urease analysis on Christensen's urea agar. Absence of

pink coloration indicates loss of urease activity. (B, C, D) Analysis of secretion for virulence-

related enzymes such as urease, laccase, and acid phosphatase from three biologically

independent experiment sets. Statistical significance: **** $P < 0.0001$, ** $P < 0.003$, *** $P <$

0.005, * $P < 0.05$. All statistical data were determined based on one-way ANOVA and

Dunnett's post-hoc test. (E, F) Analysis of secretion for non-virulence-related enzymes such

as cellulase and α-amylase. Statistical significance: *** $P<0.0005$, ** $P<0.003$ * $P < 0.05$, ns,

not significant. All statistical data were determined based on one-way ANOVA and Dunnett's

post-hoc test. (G) Analysis of the conventional secretion of Cda1 in *C. neoformans*. Presence

of Cda1 was analyzed in total (T), soluble (S), and insoluble (I) fractions of intracellular

extracts (left), along with the secreted fraction (right). Subcellular fractionations were

performed as previously described (Thak et al., 2022), and the fractions were subjected to

western blotting analysis using an anti-Cda1 antibody.

1140

Figure 8. Analysis of extracellular vesicles purified from WT, *ugg1*Δ, and *cap59*Δ cells.

(A, B) Nanoparticle tracking analysis (NTA) of EVs extracted from WT, *ugg1*Δ, and *cap59*Δ

strains and quantification of total extracellular vesicle (EV) concentration per cell density.

Quantitative measurements were derived from three independent experiments with standard

deviations presented as error bars. Statistical significance: *** $P<0.0005$, * $P< 0.05$. All

statistical data were determined based on one-way ANOVA and Dunnett's post-hoc test. (C, D)

Cryo-TEM imaging of purified EVs and comparative analysis of EV size in WT, *ugg1*Δ, and

*cap59*Δ strains. Scale bar, 100 nm. The outer EV diameter of a total number of 100 EVs per

strain, captured using cryo-TEM, were measured. (E) Heatmap representation of fold change

between WT and *ugg1Δ* EV-associated proteins, commonly detected in this study and in

previously reported EV proteome datasets (*ugg1Δ*/WT). Upregulated proteins in *ugg1Δ* are

shown in red, whereas downregulated proteins are shown in blue. The proteome data of

whole-cell lysates (WCL), generated from the cell pellets obtained after EV separation, were

included for comparison. (F) Blotting analysis of GXM in the *C. neoformans* cells and EVs

of WT, *ugg1Δ*, and *cap59Δ* strains. The 8 M urea extracts were obtained from EVs and cell

pellets, from which EVs are generated. The urea extracts (5 µg total proteins) were loaded on

8% SDS-polyacrylamide gel and subjected to silver staining or blotting analysis using the

anti-GXM 18B7 (α-GXM) and anti-Cda1 (α-Cda1) antibodies, respectively. Left: total cell

extract. Right: total EV extract.

1160

Figure 9. Impact of ERQC disruption on glycoprotein folding and EV-mediated

transport of virulence factors in *C. neoformans*. (A) In WT strain, *C. neoformans* UGGT

homolog, Ugg1, functions as a sensor for misfolded glycoproteins within the ER, playing a

crucial role in protein quality control. Functional ERQC is essential not only for ensuring the

proper folding of glycoproteins, which is critical for maintaining cellular fitness, but also for

facilitating EV-mediated secretion of capsule polysaccharides and virulence-related enzymes

necessary for pathogenicity. (B) In the UGGT-deficient strain (*ugg1Δ*), ER stress is increased

because of misfolded protein accumulation within the ER lumen. This heightened stress leads

to decreased cellular fitness, which negatively impacts EV biogenesis and cargo loading.

Consequently, significant defects occur in EV-mediated transport, which ultimately leads to a

complete loss of virulence. Nc: nucleus, Vc: vacuoles. This figure was partially created using

BioRender.

A

H. sapiens

ALG6 ALG8 ALG10 OST — GLS1 GLS2 — GLS2 / CNX/CRT cycle / UGGT — UDP UDP — ERManI EDEM2 — Golgi — EDEM3 EDEM1 — Os9 — Proteasomal degradation

S. cerevisiae

Alg6 Alg8 Alg10 OST — Gls1 Gls2 — Gls2 Cne1 — Mns1 — Golgi — Htm1 — Yos9 — Mnl2 — Yos9 — Proteasomal degradation

C. neoformans

OST — UGGT (Ugg1)? — Gls2 / CNE cycle / Ugg1 — UDP UDP — Mns1? — Mns101? Golgi — Mnl1? — Os9 — Mnl2? — Os9 — Proteasomal degradation

● Glucose ● Mannose ■ N-acetylglucosamine 🗲 Misfolded protein ☆ Folded protein

B

C. neoformans Ugg1 — TRXL12 TRXL13 TRXL14 TRXL15 UDP-g GGTase GT24 — 1,582 a.a

C. neoformans Mns1 — GH47 — 603 a.a

C. neoformans Mns101 — GH47 — 865 a.a

C. neoformans Mnl1 — GH47 — 940 a.a

C. neoformans Mnl2 — GH92N GH92 — 913 a.a

■ Transmembrane domain
■ Signal peptide
■ ER retention signal
Ψ N-glycosylation site

A

Intracellular GXM

Shed GXM

B

Acceptors of cell surface capsule

Capsule transfer using WT shed GXM

C

Acceptors of cell surface capsule

Capsule transfer using WT shed GXM

Capsule transfer using *mns1Δ101Δ* shed GXM

D

YPD

10% Sabouraud

A WT

Exosomes

Microvesicles

MVB

Golgi

Vc

ER

Nc

Acid phosphatase · Urease · Laccase · Sterols · GXM · Secreted proteins · CDA1

ER

Golgi

Mns101

Gls2

CNE
cycle

Ugg1

UDP   UDP

Ugg1

Mns1

ERAD

Proteasome
degradation

Ire1

UPR
activation

B *ugg1*Δ

MVB

Golgi

Vc

ER

Nc   **ER
stress**

Vc

ER

Golgi

Mns101

Gls2

CNE
cycle

Ugg1

UDP   UDP

Ugg1

Mns1

ERAD

Proteasome
degradation

Ire1

UPR
activation

Acid phosphatase   Urease   Laccase   Sterols   GXM   Secreted proteins   CDA1

→ Conventional secretion   → Non-conventional secretion   → Transport to vacuole