The electrogenicity of the Na$^+$/K$^+$-ATPase poses challenges for computation in highly active spiking cells

Liz Weerdmeester[1,2]   Jan-Hendrik Schleimer[1,2]   Susanne Schreiber[1,2,*]

[1] Institute for Theoretical Biology, Department of Biology, Humboldt-Universität zu Berlin, Philippstraße 13, Haus 4, 10115 Berlin, Germany

[2] Bernstein Center for Computational Neuroscience, 10115 Berlin, Germany

[*] Correspondence: Susanne Schreiber <s.schreiber@hu-berlin.de>

1   Summary

The evolution of the Na$^+$/K$^+$-ATPase laid the foundation for ion homeostasis and electrical signalling. While not required for restoration of ionic gradients, the electrogenicity of the pump (resulting from its 3:2 stoichiometry) is useful to prevent runaway activity. As we show here, electrogenicity also comes with disadvantageous side effects: (1) an activity-dependent shift in a cell's baseline firing and (2) interference with computation, disturbing network entrainment when inputs change strongly. We exemplify these generic effects in a mathematical model of the weakly electric fish electrocyte, which spikes at hundreds of Hz and is exposed to abrupt rate changes when producing behaviourally-relevant communication signals. We discuss biophysical strategies allowing cells to mitigate the consequences of electrogenicity at additional metabolic cost and postulate an interesting role for a voltage-dependence of the Na$^+$/K$^+$-ATPase. Our work shows that the pump's electrogenicity opens an additional axis of vulnerability that is likely to play a role in brain disease.

2   Keywords

sodium-potassium pump, energetics of excitable cells, network synchronization, cellular biophysics, homeostasis, jury-rigged evolution, electric-organ discharge, mathematical modelling

3   Introduction

The evolution of P-type ATPases in ancestral methanogenic archaea (1,2) also lay the foundation for the energetics of energy-intensive signaling tissues like the metazoan nervous systems billions of years later (3). In particular, one of the ATPases, the Na$^+$/K$^+$-pump, plays a significant role in charging the batteries required to operate the nervous systems that the motile life of all multicellular animals is

so reliant on. The Na^+/K^+-pump exchanges intracellular sodium for extracellular potassium ions in a 3:2 ratio, and thereby generates a net outward current (Fig. 1 A). This electrogenic property of the pump appears not only as a useful exaptation for osmoregulation in eukaryotes (4), but also invokes activity-dependent changes in nerve cell excitability (5), mediating hyperpolarisation after repetitive stimulation and firing rate adaptation (6–12). These mechanisms are in turn exploited in specific neuronal encoding paradigms (13–18), cell-intrinsic bursting dynamics (16,19–21), and accelerated ion homeostasis (22–24) and thus pose an example of jury-rigging in evolution. In the study at hand, we show that for nerve and muscle cells that need to be tonically active for long stretches of time (on the order of minutes to hours), the electrogenicity of the pump has further, less explored consequences requiring special adaptations of the biophysical components that underlie sustained electrical activity.

Generally, it is assumed that the Na^+/K^+-ATPase instantaneously restores ionic gradients, ensuring the robustness of electrical signaling. Furthermore, it is assumed that the net current that is produced by pump activity has limited effects on spiking activity. Under these assumptions, generally, pump currents are not explicitly modeled and reversal potentials are kept constant at all times. At first glance, this seems a reasonable pragmatic approach both for the interpretation of experimental data as well as for computational models of neural dynamics. Accordingly, relatively little attention has been given to the interference between electrogenic pumping and neuronal voltage dynamics, a trend reinforced by the success of models capturing neural dynamics without pump currents, solely based on fixed ion reversal potentials, such as the classical Hodgkin-Huxley model (25). Here, we demonstrate that, contrary to this notion, electrogenic Na^+/K^+-ATPases can exert a significant direct impact on computational properties of highly active excitable cells, and that Na^+/K^+-ATPase electrogenicity poses a challenge for robust spike-based signaling. For strongly active cells operating at high firing rates, additional mechanisms balancing out the pump's effect on computation are required, imposing extra costs on cell signaling.

While, due to their generic nature, the described mechanisms pose challenges for any excitable cell that has to rely on electrogenic pumps, we here showcase them in the electrocyte of the weakly electric fish (26), chosen because of its persistently high firing rates permanently exceeding hundreds of Hz and its, consequently, significant energetic demand. Electrocytes are the cells that make up the electric organ (EO), creating the electric field in the animal's environment that is vital for communication (27). Electrocyte firing rates, and thus the frequency of the electric organ discharge (EODf), are key for the animal's survival: the frequency of the resulting oscillating weakly electric field can be sensed by other individuals through electroreceptor afferents (28) and spans a range of 400 Hz across individuals (29). It constitutes the primary signal transmitting information about sex and hierarchy and is also used in interspecific communication. Due the high-frequency spiking activity and thus high ATP requirement of electroyctes (30,31), the EOD cannot only be expected to have been under a severe evolutionary pressure for energetic efficiency, it also exposes the cells to relatively strong electrogenic pump currents that alter cell excitability. The activity-dependent pump currents thus directly influence electrocyte firing rates, as we argue here, complicating the precise regulation of the excitable cells' activity.

We identify two major effects of electrogenic pumps on neural computation: (1) a significant shift in a cell's baseline activity requiring compensation and (2) strong computational side effects of electrogenic

pumping in the presence of functionally relevant input changes. While the first effect is intuitive – electrogenic pumping permanently contributes a hyperpolarising current that requires a compensation to keep firing rate set points – the second effect is less so. In particular, it can result in unexpected rate changes up to a complete silencing of spiking activity in cases where a drastic firing increase was required and induce spontaneous activity where silencing was required. Even when effects are milder, causing only graded firing rate changes but not silencing, the pump's electrogenicity interferes with cellular computation when spike timing is to be synchronised across excitable cells. As we show, entrainment of electrocytes by their pacemaker neurons can be disrupted in these cases, weakening the EOD and presumably impairing the fitness of weakly electric fish. Based on the identified mechanisms, we suggest a diverse set of biophysical options that permanently mitigate the side effects of pump electrogenicity in both cases.

To do so, we first isolate the effects of pump currents on cell excitability with constant stimulation. We then demonstrate the impact of pump currents in a simple network context, when an electrocyte is driven by a pacemaker neuron. We outline the consequences of electrogenic pumping for two behaviourally-relevant signals in weakly electric fish: so-called chirps (i.e., short interruptions in the EOD) and frequency rises (transient frequency sweeps). Finally, we discuss how a voltage dependence of the Na^+/K^+-ATPase – compared to a pump that adapts its rate exclusively as a function of bulk ionic concentrations and hence only on long timescales – can be useful in alleviating some of the detrimental effects of electrogenicity on cellular dynamics.

4 Results

The electrogenic property of the Na^+/K^+-ATPase plays a role in the osmoregulation of single eukaryotic cells (4) as well as, on the organismal level, in marine osmoregulatory organs (32). In metazoan nerve cells, the Na^+/K^+-pump restores the cross-membrane ion gradients that generate resting and action potentials (AP), whilst its electrogenicity induces a hyperpolarising membrane current proportional to its pumping activity. The pump rate's concentration dependence creates a negative feedback loop, leading to an increase in the hyperpolarising current if nerve cells discharge at higher frequencies where more pumping is required (6–12). This feedback loop could thus reduce, terminate or prevent overactivation. On the other hand, for nerve and muscle cells that need to be tonically active for long stretches of time (on the order of minutes to hours), the electrogenicity poses substantial challenges and its consequences need to be balanced by additional mechanisms, the latter of which further restrict energetic efficiency, as we demonstrate in the following.

4.1 Na^+/K^+-ATPase affects tonic spiking

Most computational models of excitable cells follow the principles of conductance-based models for action potential (AP) generation derived by Hodgkin and Huxley (33). Accordingly, they assume that the compensation of the transmembrane ion currents is carried out perfectly without interfering with voltage dynamics, that the hyperpolarising current introduced by the pump is negligible and that the reversal potentials remain constant (25). An explicit inclusion of the pump into mathematical models

of such cells is hence not required. To challenge this view for highly active cells, we first consider the following scenario: Let's assume that an electrogenic Na^+/K^+-ATPase is added to the Hodgkin-Huxley type spiking mechanism to compensate ion flow across the membrane. Specifically, we illustrate the effects in an experimentally well constrained model of the tonically active electrocyte of weakly electric fish (26). For simplicity, the Na^+/K^+-ATPase in this first case is assumed to only depend on intra- and extracellular ion concentrations as in (19,34), and ATP levels are assumed to be sufficiently high not to impair Na^+/K^+-ATPase activity. Specifically, a voltage-dependence of the Na^+/K^+-ATPase (described in some previous studies, see for example (35–39)), is omitted here, following the approach taken in most previous concentration-dependent models. The layer of complexity that would be added by a voltage dependence of the pump, however, is discussed in section 4.4.

Due to the relatively slow dynamics of ion concentrations, on the timescale of the action-potential generation the pump current is approximately constant. Therefore, under the assumption of a constant ion channel composition, a strong hyperpolarising pump current will decrease the input-induced firing rate of the cell, potentially up to the extreme point of silencing it. This effect can be seen in Fig. 1 B, there reflected in a right-shift of the frequency-input curve (tuning curve) of a model with electrogenic pump relative to the model without. In other words, for a constant input that elicits high-frequency tonic spiking without this hyperpolarising pump current, the addition of a pump current decreases a cell's firing rate; for very strong pump currents, firing is eliminated alltogether (Fig. 1 C).

By measuring sodium currents over time, one can estimate the pump rates necessary to facilitate high frequency firing ((26), Methods 6.1.2). To sustain physiological electrocyte firing rates, howewer, 'appropriate' pump currents (Fig. 1 D) would hyperpolarise the cell to an extent that very high current inputs are needed to elicit tonic high frequency firing (Fig. 1 B,C). These high inputs would come at a significant metabolic cost of synaptic transmission (specifically the cost related to production and packaging of AChR molecules (26,40)).

Alternatively, the pump-induced raise in current rheobase could be compensated for by adequate adjustments of the cell's ion channel composition. Specifically, for an excitable cell to operate in a regime of tonic firing, the constant net outward pump current could be balanced by an additional constant inward current, which can be achieved via co-expression of Na^+/K^+-ATPases and, for example, sodium leak channels (Fig. 1 E, F, G, Methods 6.1.1). Although we are not aware of quantitative data on the regulation of ATPase expression in electrocytes, it seems reasonable to assume that the number of pumps expressed in electrocytes scales with the average energetic demand of its spiking activity. An electrocyte that generally fires at higher rates thus requires more pumps to maintain ionic homeostasis. The discharge of the electrical organ (EOD) of E. virescens (chosen as a typical representative and whose physiology has been well quantified) approximates to the summed activity of electrocytes and individual fish have different baseline EODfs (29), and thus different electrocyte firing rates. Energetic requirements and, consequently, pump expression levels will therefore also vary among individuals. In turn, pump current and the resulting shift in rheobase (that requires compensation) are likely to be unique for each organism. As can be seen in Fig. 1 E, an appropriately chosen co-expression factor between Na^+/K^+-ATPases and sodium leak channels (Methods 6.1.1) suffices to stabilise the rheobase for a wide range of pump currents (Fig. 1 F, G) which are produced within the regime of physiological

firing rates (Fig. 1 D). Therefore, this postulation of a co-expression mechanism (similar to those that have been described for homeostatic regulation of intrinsic excitability, see (41)) provides an elegant solution that allows for reliable tonic high-frequency firing with strong pump activity despite cell-to-cell differences in firing rates and pump expression levels.

As the overlap of the relatively constant opposing outward pump current with the compensatory inward current, which comprises one-third of the sum of all inward currents (see Methods 6.1.2, Eq. 17), results in a largely electroneutral exchange of positive ions (Fig. 1 H), a surprisingly high fraction of the pump's energy is spent on pumping out sodium ions that do not directly contribute to the action potential of the cell but only compensate for the additional pump current (Fig. 1 I). This current redundancy is particularly severe for systems operating at high average firing rates which require high pump densities to maintain ionic homeostasis (Fig. 1 D). Therefore, the energetic efficiency of action potentials is reduced (42) because of the electrogenicity of the Na^+/K^+-ATPase by one third compared to the hypothetical scenario of electroneutral pumping, where no additional inward currents would be needed to enable tonic firing (Fig. 1 J). In a tonically active cell, the negative feedback loop that the electrogenic Na^+/K^+-ATPase provides to enhance ionic homeostasis for action-potential firing thus is likely to come at the cost of a more constrained ion channel composition and sub-optimal energetic efficiency.

Figure 1: Homeostatically relevant Na$^+$/K$^+$-ATPase electrogenicity comes at the cost of a more constrained ion channel composition and sub-optimal energetic efficiency. *(A) The Na$^+$/K$^+$-ATPase (grey) exchanges 3 intracellular sodium ions for 2 extracellular potassium ions against their gradients, which generates a net outward current. (B) Increased pump activity, and thus an increased hyperpolarising pump current, reduces cell excitability as a bigger inward input current is needed to activate voltage gated channels. (C) For a fixed input current that generates tonic firing in an excitable cell, an increase in pump current 'silences' the cell. (D) High firing rates require significant pump activity, which generates a significant hyperpolarising pump current. (E, F, G) Sodium leak channels facilitate a depolarising current that balances out the hyperpolarising pump current. If Na$^+$/K$^+$-ATPase and sodium leak channels are co-expressed (E), the impact of increased pump activity on cell excitability is minimised (F, G). (H) The additional inward sodium current (dark red) that is required to balance the outward pump current (grey) creates a redundancy in current flows which hampers energetic*

efficiency. Currents are plotted as contributions to the total in- and outward currents (top) and as separate currents (bottom). (I) Action potentials are similar in size when generated with a (balanced out) pump current and without a pump current. (J) For the former however, the redundancy in current flows requires more sodium ions to be pumped against the gradient per action potential.

4.2   Na$^+$/K$^+$-ATPase affects the tuning curve

As outlined above, quantitatively, the compensation required because of the pump's electrogenicity depends on a cell's spiking activity. Consequently, even if ion channels and Na$^+$/K$^+$-ATPases were co-expressed, and both were optimised to facilitate tonic firing in an excitable cell (Methods 6.1.2), the electrogenicity of the Na$^+$/K$^+$-ATPase still would interfere with neural processing when firing rates change drastically due to changes in the input, as outlined in the following.

Most excitable cells do not operate at a fixed firing rate. Often, a flexible, transient modulation of firing rates is required either to track and encode stimuli (43) or to control and adapt motor programs (44). Such a modulation is evoked by alterations in synaptic inputs that differ from baseline activity (Fig. 2 A, D). If the pump rate remains constant despite such changes in firing rates, ion accumulation or drain is to be expected. This can lead to critical transitions in cell excitability and function (as intrinsic cell dynamics depend on ion concentrations (5,21,45,46)), and, in the extreme case, diminish the ion concentration gradient to the extent that firing is completely impaired (47).

At first glance, the pump's sensitivity to ionic concentrations (Methods 6.1.3) seems an adequate solution that can alleviate such effects of drastically changing firing rates. The dependence of the pump on ionic concentrations contributes to an activity-dependent restoration of ion gradients, i.e., an appropriately calibrated concentration dependence of Na$^+$/K$^+$-ATPase activity can help to match the energetic demand of the cell's recent activity (48) (Fig. 2 C, F). The adapted pump activity, however, is accompanied by a change in hyperpolarising current; the cell is pushed to a regime that the system was not originally tuned to (Fig. 2 B, E). Therefore, even in a perfectly controlled environment, an excitable cell can assume different firing rates in response to the same input, where the immediate output of the cell depends on previous activity (16). If a fixed input-output mapping is key to the function of an excitable cell, the electrogenicity of the Na$^+$/K$^+$-ATPase induces yet another trade-off between ionic homeostasis and cell function.

Figure 2: Homeostatic feedback loops based on Na⁺/K⁺-ATPase activity affect firing responses through altered pump currents. *(A, B, C) Synaptic input suppression (A) initially silences the cell (B). The Na⁺/K⁺-ATPase adjusts to the reduction in energetic demand through reduced activity. This reduces the pump current (C), which increases cell excitability and results in spontaneous firing without synaptic inputs (B, black). Without a pump current (blue), spontaneous firing is not induced. (D, E, F) Increased synaptic inputs (D) initially increase firing rates (E). The Na⁺/K⁺-ATPase adjusts to the increase in energetic demand through increased activity. This increases the pump current (F), which decreases cell excitability and results in reduced firing rates (E, black).*

4.3   Na⁺/K⁺-ATPase affects entrainment

As we illustrate next, a pump-induced alteration of neuronal response properties is especially problematic if excitable cells are to be entrained in a network or by a pacemaking system. Weakly electric fish electrocytes, like most excitable cells, do not operate in isolation. In order to create an oscillating weakly electric field (Fig. 3 A), electrocyte firing needs to be coordinated across their population, which is enabled by a common drive from an upstream pacemaker (Fig. 3 B, C, Methods 6.2). In order to serve a variety of communication paradigms with largely different EOD patterns and hence electrocyte firing rates, an accurate manipulation of the electrocyte firing rates by the pacemaker nucleus is crucial for electric fish.

Each electrocyte in the electric organ is innervated on the posterior side by the spinal motor nerve, which transmits signals from the pacemaker nucleus to the electrocyte (Fig. 3 B) (49). Electrocytes are not synaptically connected among each other; they receive a unidirectional synaptic input from the pacemaker nucleus and firing patterns are only driven by the pacemaker. Therefore, we can model the effects of upstream cells on the electrocytes by simulating the periodic input currents that originate from neurotransmitter release in the synaptic cleft, modulated by pacemaker nucleus activity (Fig. 3 C, top) (26). The pacemaker entrains the electrocyte on a spike-by-spike basis, and the electrocyte

firing rates should faithfully follow the firing rates of the pacemaker (Fig. 3 C, bottom) to give rise to a strong, high-amplitude EOD signal.

The electrocyte operates in a mean-driven regime, i.e., the mean of the time-varying input it receives from the pacemaker suffices to invoke tonic spiking in the electrocyte (Fig. 3 D,E,F). Whether the electrocyte is entrained by the pacemaker depends on characteristics of the electrocyte's voltage dynamics (like the susceptibility to perturbations, reflected in the so-called phase-response curve) as well as the frequency mismatch between the pacemaker and the firing rate of the electrocyte in response to the stimulus mean. If the frequency mismatch is too large, entrainment fails (Fig. 3 G, Appendix 11.1, Fig. A1). As the pump current affects the mean-driven firing rates of the electrocyte, it can significantly impact entrainment in this simple network, in particular, when the pacemaker frequency and hence mean input to the electrocyte changes.

Figure 3: *The weakly electric fish electrocyte: an example system with strong Na⁺/K⁺-ATPase activity and strict entrainment requirements.* *(A) and (B) are modified from (49). (A) The Electric Organ Discharge (EOD) creates a weakly electric field that can be use to detect objects and communicate with conspecifics. (B) The weakly electric organ consists of a line of excitable cells called electrocytes. Every electrocyte is innervated by the spinal motor neuron that delivers signals from the pacemaker nucleus (PN). (C) The input current to the electrocyte stemming from the PN (top) sets the high frequency firing rate of the electrocyte (bottom). (D) Constant input currents (top) also elicit tonic high frequency firing in the electrocyte (bottom). (E) Tonic electrocyte firing is realised for constant input currents that lie within the mean of the input currents that are generated in the behaviourally relevant regime (200-600 Hz) (F). (G) There is a frequency mismatch between the pacemaker firing rate and the mean driven electrocyte firing rates, which influences signal entrainment.*

For an individual *E. virescens*, pacemaker firing rates can remain constant over long periods of time (50,51). If the electrocyte repeatedly receives the same input and thus produces action potentials that displace a fixed amount of ions per unit of time, pump rates and co-expressed inward leak channels can be tuned to maintain tonic firing and ionic homeostasis. When searching for food, hiding from predators, and courting, however, substantial deviations from baseline occur. In such cases of drastic change in pacemaker firing rate, the pump rate and thus the pump current adapts via its concentration dependence to the new firing statistics of the electrocyte which, consequently, alters the tuning curve (Fig. 1 B) and hence also the mean-driven firing rates (Fig. 2). In electrocytes of *E. virescens*, behaviourally-relevant deviations from baseline firing come in several forms which include chirps and frequency rises (both used for inter-individual communication), and have different consequences for cell entrainment, as we illustrate in more detail in the following.

4.3.1 The Na$^+$/K$^+$-ATPase affects the ability to generate EOD chirps

Chirps consist in short cessations (type A) or period doublings (type B) of the EOD, meant to send a submissive signal in dominance fights (52–54) and to avoid attacks from predators or conspecifics (55). Type A chirps, which essentially correspond to short 'pauses' in EOD generation, are generated in electrocytes through short interruptions of PN firing (Fig. 4 A) (56). As electrocytes are only innervated by excitatory synapses, successful chirp generation thus relies on an electrocyte that is 'silent' when devoid of input.

From experimental observations it is known that the length of type A chirps in *E. virescens* can extend beyond twenty times the length of one EOD (29,57). Repetitive emission of such long type A chirps (Fig. 4 B, top) decreases mean firing rates of electrocytes and thereby the action-potential-induced ion displacement, ultimately resulting in a lowered pump current (Fig. 4 B, bottom). We find that the effect of an individual chirp on pump currents is small and does not alter electrocyte excitability to an extent that firing rates are severely affected (Fig. 4 C, left). In case of consecutive chirping, however, the hyperpolarising pump current progressively weakens with time (Fig. 4 B, bottom), eventually leading to spontaneous electrocyte firing in absence of pacemaker input (Fig. 4 C, right). This effect limits the number and duration of chirps that can be induced by the pacemaker (Fig. 4 B, top). The observed dynamics suggest that mechanisms increasing the timescale of the pump feedback loop, such as extracellular potassium buffering (Methods 6.2.4), are suited to diminish the effects of chirps on the pump current (Fig. 4 D) and thereby the effect of variable input signals on chirp generation (Fig. 4 E). We find that extracellular potassium buffering is particularly efficient in dampening Na$^+$/K$^+$-ATPase effects on cell excitability, because Na$^+$/K$^+$-ATPase rates of excitable cells are especially sensitive to extracellular potassium concentrations (58,59) (Methods 6.1.3, equation 18) and potassium buffering reduces the variability of potassium concentrations in extracellular space. Potential metabolic costs of potassium buffering, however, constitute an additional expense in the total energy budget of the organism.

Figure 4: Homeostatic feedback loops on Na⁺/K⁺-ATPase activity impede chirp generation in electrocytes and can be mitigated through potassium buffering. *(A) Schematic illustration of the chirp setting. Left: the electrocyte (black) is coupled to the pacemaker nucleus (PN, blue) with an excitatory synapse. Right: PN spikes (blue, top) induce chirps in the electrocytes through cessation of inputs and thereby temporarily shut off the Electric Organ Discharge (EOD, middle). When chirps are properly generated, instantaneous firing rates (bottom) of the electrocyte (black) equal those of the PN (blue). (B) The generation of consecutive long chirps by the pacemaker (indicated by black arrows and instantaneous PN firing rates $\frac{1}{ISI_{PN}}$) lowers the mean firing rate of the electrocyte (black, top) and thereby the energetic demand of the electrocyte, which is fed back into a decreased pump current (bottom). This decreased pump current increases cell excitability, which over time (in this paradigm after 400 ms) leads to a mismatch between PN and electrocyte firing rates (top). (C) Electrocyte (black) and PN (blue) spikes (top) and electrocyte membrane voltage (bottom) during chirps before (left) and after (right) a significant decrease in excitability-altering pump current. After a significant deviation in pump current, electrocyte firing occurs during chirps (right). (D,E) Same as (B, C) with extracellular potassium buffering. Extracellular potassium buffering reduces the timescale of the*

homeostatic feedback loop of Na$^+$/K$^+$-ATPase activity on energetic demand which reduces the effect of transient firing rate deviations on pump current (D, bottom). This reduces the deviation in pump current to an extend that chirps can be reliably generated (D (top), E).

4.3.2 The Na$^+$/K$^+$-ATPase affects generation of frequency rises

During courtship behaviour, sudden frequency rises of the EOD followed by an exponential frequency decay back to the original baseline in the course of 2-40 seconds constitute an important signal (60). Frequency rises are produced by the pacemaker and the increase in PN firing rate is meant to entrain the electrocyte accordingly (Fig. 5 A). To this end, the mean-driven firing rates of the electrocyte should be sufficiently similar to the transiently elevated PN firing rates, because entrainment fails if frequency mismatches are too large (61,62). Accordingly, electrocytes with very slow mean-driven dynamics cannot be entrained to very fast PN inputs (see Appendix 11.1, Fig. A1).

Short frequency rises in *E. virescens* of around 2 seconds have been measured to encompass frequency elevations of up to 40 Hz (29). Repetitive emission of such frequency rises (Fig. 5 B, top) increases mean firing rates and thereby the action-potential-induced ion displacement, resulting in an increased pump current (Fig. 5 B, bottom). Comparable to the observation for chirps, we find that the effect of a single frequency rise on pump currents is small and does not alter electrocyte excitability to an extent that impedes entrainment (Fig. 5 C). With repetition of these communication signals, however, the hyperpolarising pump current significantly increases over time (Fig. 5 B, bottom), eventually decreasing the electrocyte's mean-driven firing rate to the point where a precise 1:1 locking between PN and electrocyte breaks down (Fig. 5 D, entrainment index is statistically smaller than in C (63)). Again, this effect of the pump imposes a limit on the number and duration of such frequency rises that can be induced without impairment of electrocyte entrainment and hence the EOD strength (Fig. 5 B, top). Mechanistically, the ability of the electrocyte to entrain to the pacemaker does not only depend on their frequency mismatch, but also on the strength of their synaptic coupling (Appendix 11.1, equation 28). An increase in synaptic coupling strength (Methods 6.2.4) extends the maximum frequency mismatch that still allows for synchronization (Appendix 11.1, equations 33, 32). Both effects of frequency mismatch and synaptic coupling can be illustrated by the so-called Arnold tongue (64). We therefore hypothesise that a strong synapse facilitates electrocyte entrainment and can prolong phases without pump-induced entrainment breakdown (Fig. 5 E-G). Yet it comes at the energetic cost of the increase in neurotransmitter release (including the production and packaging of AChR molecules) (26,40).

Figure 5: Homeostatic feedback loops on Na^+/K^+-ATPase activity impede the generation of frequency rises and can be mitigated through strong synaptic coupling. *(A) Schematic illustration of the generation of frequency rises. Left: the electrocyte (black) is coupled to the pacemaker nucleus (PN, blue) with an excitatory synapse. Right: Frequency rises are generated through a rapid increase in PN firing rates which exponentially decay back to baseline rates (blue, top). As the electrocytes are entrained by the PN (bottom), their firing rates mimick that of the PN and also show a frequency rise (black, top). (B) The generation of consecutive frequency rises by the pacemaker (blue) increases the mean firing rate of the electrocyte (black, top) and thereby the energetic demand of the electrocyte, which is fed back into a increased pump current (bottom). This increased pump current decreases cell excitability, which over time (in this paradigm after 15 seconds) leads to a mismatch between PN and electrocyte firing rates (top). (C, D) Electrocyte (black) and PN (blue) spikes (top) and electrocyte membrane voltage (bottom) during frequency rises before (D) and after (E) a significant increase in excitability-altering pump current. After a significant deviation in pump current, not all PN spikes are*

reproduced in the electrocyte which leads to 'missing' spikes (D). This is reflected in the synchronization index $\overline{R}$, which decreases with increasing pump current deviation. (E-G) Same as (B-D) with strong synaptic coupling. Strong synaptic coupling attenuates the effect of altered pump currents on electrocyte entrainment and enables reliable production of frequency rises (E (top), F, G) but require more ATP-demanding pumping (E, bottom) compared to weak coupling (B, bottom).

The analysis of both types of signals, chirps and frequency rises, shows that the electrogenic Na^+/K^+-ATPase can have significant effects on the computational properties of highly active excitable cells, potentially requiring energetically costly countermeasures for normal operation, especially if cells are to be entrained by a pacemaker or in a network. This suggests that even though Na^+/K^+-ATPase was jury-rigged to support the generation of action potentials in excitable cells, the fact that their original function required them to be electrogenic, inevitably calls for countermeasures that lower the energy-efficiency of signaling.

4.4   The role of Na^+/K^+-ATPase voltage-dependence

A pump current that only varies on the longer time scale of changes in ion concentrations acts like a constant current on the time scale of spike-generation (Fig. 1 H) and will horizontally shift the tuning curves as described above (Fig. 1 B). The consequence of this shift (if uncompensated as described above) is a drastic change in firing rates (Fig. 1 C, Fig. 2, Methods 6.1.1). We next explore whether changes of pump rates on the shorter timescale of action potentials can alleviate the pump-current induced firing rate adaptation. In particular, we illustrate in the following that a voltage-dependence of the pump may constitute an interesting means to limit pump-induced firing-rate modifications and at the same time save on the energetic cost of action potentials.

A common supposition is that (to restore the ionic gradients that get depleted during the generation of action potentials) the pump only depends on ion concentrations and hence, to first approximation, displays constant activity during spiking. In this case, the hyperpolarising pump current counteracts the sodium-currents at the depolarisation phase of the AP upstroke and, in fact, assists the potassium-currents in repolarising the cell during the action-potential downstroke (Fig. 6 A, left). In the following, we contrast this constant pump to a voltage-dependent pump that activates selectively only during the action-potential downstroke (Fig. 6 A, right). The pump's voltage dependence could benefit a neuron in two ways: First, by not affecting sodium-based depolarisation it may reduce the shift that the pump current induces in the tuning curve. Second, by aiding potassium in repolarisation it could provide some energetic benefits for spiking cells.

The exact kinetics and voltage-dependence of Na^+/K^+-ATPases differ per cell type and organism (38,39,65) and have likely evolved differently in distinct cells to support their unique function and energetic demand. To highlight the positive effects of a voltage-dependence of the Na^+/K^+-ATPase in electrocytes, whose dependence on voltage to our knowledge currently is unknown, we use a thought experiment based on an idealised voltage-dependent pump that optimally compensates tuning curve shifts and reduces energy demand (Fig. 6 A, right).

Specifically, the dynamics of this idealised voltage-dependent pump is assumed to exactly mimic the

hyperpolarising potassium current in the following way: Start with a classical Hodgkin and Huxley equation without pump. Then reduce the repolarising potassium current to $\frac{2}{3}$ of its strength. The missing $\frac{1}{3}$ is now substituted by a pump current with exactly the same time course $I_p = \frac{1}{3}I_K$, such that,

$$C_\mathrm{m}\dot{v} = -I_\mathrm{Na} - I_\mathrm{K} = -I_\mathrm{Na} - \frac{2}{3}I_\mathrm{K} - \frac{1}{3}I_\mathrm{K} = -I_\mathrm{Na} - \frac{2}{3}I_\mathrm{K} - I_\mathrm{p}.$$

In the model without pump, the cumulative sodium and potassium currents have to add up to zero after one period (see Appendix 11.2). This, together with the equation above, implies that the chosen I_p meets the requirements of the pump stoichiometry $\oint I_\mathrm{p}dt = \frac{1}{3}\oint I_\mathrm{Na}dt$ and $\oint I_\mathrm{p}dt = \frac{1}{2}\oint I_\mathrm{K}dt$ and thereby perfectly counteracts currents flowing during a complete action-potential cycle to maintain ion homeostasis. The equation also shows that replacing one-third of the potassium channels with the idealised voltage-dependent Na$^+$/K$^+$-ATPase would leave the action potential shape unchanged compared to the model without pump. Importantly, as the pump substitutes for potassium channels, it reduces the flux through these channels by a factor of $\frac{1}{3}$. In addition, compared to the constant pump scenario, there is no need for sodium leak channels to cancel out the hyperpolarising pump current. This additionally reduces the cumulative flow of sodium ions by approximately $\frac{1}{3}$ with respect to an excitable cell with relatively constant pumping. Taken together, the reduction in flow of sodium and potassium ions reduces the pump load by approximately $\frac{1}{3}$ (Fig. 6 B).

Besides lowering the energetic demand, voltage-sensitivity can also reduce the effect of the Na$^+$/K$^+$-ATPases on the neuron's tuning curve and, consequently, firing rate adaptation. Two main factors help reducing the adaptation: First, the pump current with a potassium-like voltage-dependence is almost completely inactive at spiking onset (approximately -80 mV), in contrast to the fully active constant pump current. Alterations in the former pump current hence induces almost no shift in the tuning curve (Fig. 6 C). Second, as the pump optimally scales with membrane currents at any firing rate in the voltage-dependent case, it instantaneously leads to near-perfect homeostasis. Therefore, action-potential induced concentration changes are minimal. In contrast to the case of a constant pump (Fig. 2), pacemaker-induced jumps in electrocyte drive hence do not elicit a substantial firing rate adaptation because concentrations remain largely unchanged (Fig. 6 D). We note that even if they would, their effect would be small due to the first argument (Fig. 6 C).

Consequently, suppression of synaptic inputs (like during a chirp) does not result in spontaneous firing for a pump with the described voltage dependence and a cell can be reliably 'silenced' for extended periods of time (Fig. 6 D, left). Furthermore, there is a fixed input-output mapping from synaptic input to firing rates in this case (Fig. 6 D, right), which benefits the robustness of entrainment between pacemaker and electrocyte.

While it remains to be established experimentally whether Na$^+$/K$^+$-ATPases in electrocytes do exhibit a voltage dependence resembling the one postulated here, our thought experiment demonstrates the generic potential of a voltage dependence of pumps to mitigate negative side effects of ion pumping in highly active cells and to lower the need for a costly investment into alternative compensatory measures such as the co-expression of additional ion channels, increased synaptic weights, or extracellular

potassium buffering.

Figure 6: Ideal voltage dependence of the Na⁺/K⁺-ATPase for energy-efficient action potentials and minimal firing rate adaptation *(A) Current contributions to the total in- and outward currents (top) and absolute current flows (bottom) for one electrocyte action potential with a Na⁺/K⁺-ATPase without voltage dependence (left) and with an optimal voltage dependence that mimics potassium channels (right). The voltage-dependent pump current takes on the role of a potassium channel and contributes significantly to the net current at the AP downstroke. (B) Total amount of sodium (red) and potassium (blue) ions, and net ion tranfer of the pump (grey) that are relocated per AP for a cell with a voltage-insensitive pump (left) and a voltage-dependent pump (right). (C) The effect of pump density on the tuning-curve is minimal for ideal voltage-dependent pumps (bottom) compared to a non-voltage dependent pump (top). (D) Signal generation in a cell with Na⁺/K⁺-ATPases with optimal voltage dependence. Synaptic input suppression (top left) silences the cell (center left) and reduces the pump current (bottom left). Firing rates are however not affected, and the cell remains silent. Synaptic input doubling (top right) increases firing rates (center right) and increases time-averaged pump currents (bottom right). Firing rates are however not affected. Note that the instantaneous pump current (bottom, grey) varies on the timescale of action potentials, which is highly compressed in this 100 second time-window.*

5 Discussion

We investigated the rarely acknowledged "side effects" of the electrogenic Na^+/K^+-ATPase on the computational properties of a highly active spiking cell: the weakly-electric fish electrocyte. Our findings highlight that the electrogenicity of the Na^+/K^+-ATPase poses challenges for robust signal processing in highly active cells; a pump that would have evolved for the sole purpose of maintaining ionic gradients would be more efficient if it was electroneutral. We dissect the mechanisms involved and show that energy-intensive countermeasures are required to ensure robust performance in the presence of the pump when a tonically active cell is driven by inputs of alternating mean – a perspective that is underrepresented in the current literature. While analysed in electrocytes, the model is and identified mechanisms are sufficiently generic to translate to other excitable cells across the animal kingdom operating at high firing rates, such as Purkinje neurons (66), vestibular nuclear neurons (67), and fast-spiking interneurons (68,69).

When Hodgkin and Huxley established their pioneering conductance-based model of action potential generation in 1952 (25), it was generally assumed that the sodium-potassium pump, which maintains the ionic gradient across the cell membrane, was electroneutral (6). Their computational model, which was adapted over the following 70 years to model numerous types of excitable cells in diverse tissues and species, therefore did not include a pump current. Even later, after the 3:2 stoichiometry of the sodium-potassium pump and thus its electrogenicity were proven, the pump current was often not included in simple point-neuron models, presumably because of its relatively small amplitude and effect size (70,71). Although this may be a suitable argument for excitable cells that are only moderately active, it is less likely to hold for cells that need to be tonically active over long periods of time. In fact, tonically active cells are expected to operate closer to the strong-activity-inducing, post-tetanic stimulation protocols used in the 1960s to render to pump current more visible by artificially increasing their effect size (72–74). Additionally, the functional consequences of other, more subtle pump properties, such as its voltage-dependence, have not been explored. Here, we identify a range of possible compensation mechanisms for strong pump currents and speculate that, at least theoretically, the pump itself may hold a key to rendering its effects less computationally invasive.

5.1 Regulatory mechanisms

We discussed four mechanisms that improve firing rate control under strong electrogenic Na^+/K^+-ATPase currents: co-expression of Na^+/K^+-ATPases and sodium leak channels, extracellular potassium buffering, stronger synaptic coupling and pump voltage-dependence. All of these mechanisms treat the 'symptoms' of electrogenic Na^+/K^+-ATPase, and could be replaced by any other mechanism that achieves the same effect, i.e., providing an opposing current, diminishing the deviations from baseline pump currents, increasing the entrainment range of a cell, and limiting the pump activity to specific periods of the spike-generation cycle.

Opposing currents do not necessarily have to stem from a sodium leak current, but could also be achieved by other depolarising currents, like h-currents (68) or through the (relatively small) current that results from co-transport of H^+ ions by the Na^+/K^+-ATPase itself (75). The former has previously

been shown to counterbalance pump currents in the leech central pattern generator neurons (19). Accordingly, the sodium leak also does not have to result from voltage-agnostic channels, but could, for example, be facilitated by a decreased mean half-activation voltage of voltage-dependent sodium channels which could be achieved by transcriptional regulation of channels with different splice forms (76). We lastly note that even though additional supposedly 'wasteful' sodium currents might serve a secondary purpose of balancing out fluctuating currents produced via sodium-coupled transport of metabolites (77).

As we showed, deviations in pump currents, resulting from the susceptibility of the Na^+/K^+-ATPase to changes in firing rate, can be diminished by prolonging the timescale of the pump's feedback on cell activity, for example via buffering of extracellular potassium. Similar effects can be obtained from volume increases in intra- and extracellular space or a weaker concentration dependence of the pump. As we showed, for communication with chirps, an uncompensated pump could result in (undesired) spontaneous spiking of electrocytes. In these cases, an increase in the input current required to reach threshold (mediated by additional leak channels) can suppress such activity. Interestingly, compensation could also be provided on the behavioural level: First, an appropriate timing of communication signals, specifically an alternation between frequency rises and chirps, can alleviate undesired changes in baseline pump current due to their opposite impact on pump activity (i.e., increase versus decrease in pump rate). Second, a limitation of individual and cumulative signal duration and amplitude of chirps or frequency rises, respectively, similarly constrains effects on the pump current. This suggests that not only the evolution of ion channels is relevant for shaping communication signals (78) but that also the Na^+/K^+-ATPas may have played a significant role.

Finally, we argue that effects of the Na^+/K^+-ATPase on neuronal dynamics can be partially avoided if the pump was equipped with an appropriately calibrated voltage dependence. A voltage dependence of the pump has been reported experimentally (35,65). Specifically, the pumping process depends on many individual steps, each involving a different time scale and voltage dependence (36,37). While it remains unclear whether a pump could exhibit a voltage dependence as ideal as in our thought experiment in section 4.4 (in particular with respect to the restriction of its activity to periods of potassium channel activation), at least partial benefits from a modulation of pump activity along some of the qualitative principles described in the thought experiment could be expected. Future experimental investigations on the pump's voltage dependence, in particular in highly active cells, will therefore be of interest to support or reject the hypothesis predicted from our modelling approach. Interestingly, properties of the pump's kinetics and voltage dependence have been reported to be highly adaptable in evolution, showing a large heterogeneity in different tissues (38,39), including different splice variants (79) as well as a regulation via RNA editing (65).

A voltage dependence of the hypothesised ideal pump, however, requires a higher pump density. As the idealised pump's activity is constrained to specific time windows, its peak pump rate during these periods needs to be higher. In our model, the effective pump rate during these (limited) times is elevated by a factor of four (compare Fig. 6 A) in comparison to a voltage-agnostic yet constantly active pump. This brings other constraints, such as available membrane space and the cost of pump synthetization and transport to the table.

5.2 Implications for disease

Several neurological diseases (80–83) have been linked to mutations in the α subunit of the Na^+/K^+-ATPase. The origin of many symptoms observed in these diseases, such as epileptic seizures, lies in pathological neuronal network activity including hyperexcitability and altered oscillatory activity (84). Our modeling work elucidates one mechanism by which altered pump physiology has detrimental effects on cellular computation and can induce pathological network activity. For example, a mutation associated with the rare neurological disease Aternating Hemiplegia of Childhood (AHC) prevents the Na^+/K^+-ATPase co-transport of H^+ ions, the latter of which normally mitigates some of the pump's negative effects due to its depolarising contribution (75,85). From our work, we can conclude that the negative side effects of the pump on network computation should be exacerbated by this mutation. We also hypothesise that not only direct impairment of the Na^+/K^+-ATPase may contribute to pathological electrical activity, but also deficits in the postulated compensatory mechanisms, thus opening up additional points of physiological vulnerability to pathology.

5.3 Other factors of relevance

5.3.1 Reversal potentials

An increase in cellular activity reduces reversal potentials, lowering the 'driving force' in action potential generation, and hence affects firing rates (5); vice versa for decreases in activity. The effects of activity-induced concentration changes on neuronal activity therefore stem from a combination of the change in reversal potentials and in pump currents (16). Such effects of the reversal potentials were included in the model and we note that they only contributed mildly (< 5%) to the firing rate adaptation described.

5.3.2 ATP availability and hypoxia

In addition to ion concentrations, pump rates depend on ATP (86). Limitations in its availability – a phenomenon not uncommon in the weakly electric fish due to its foraging in hypoxic environments (87) – can alter pump rates more drastically than firing-induced changes in ionic concentrations. In theory, a drastic reduction in pump rate (and thus in pump current) results in elevated cell excitability followed by a depolarisation block (Fig. 1 B, (21)). In reality, in such cases the EOD of the *E. virescens* only reduces in amplitude but does not significantly change in frequency. The EOD only terminates after very long exposure to annoxia (88), suggesting the existence of compensatory mechanisms diminishing the effects of altered pump currents on cellular activity.

5.3.3 Temperature fluctuations

Weakly electric fish are poikilothermic. Inhabiting affluent streams of the Amazon river (87), with temperature variations of four degrees during a day-night cycle (89), the effects of temperature on spike generation by voltage-dependent channels (90–92) and Na^+/K^+-ATPases in *E. virescens* need to be well balanced to keep cell firing in the physiological range.

5.3.4 Spatial effects in electrocytes

Electrocytes are approximately 1 mm long and only excitable on the posterior site (49). The subcellular localization of Na$^+$/K$^+$-ATPases is hence likely to modulate the impact of the pump's electrogenicity on cell firing (26). Pumps located on the posterior side are likely to exhibit a more drastic effect on the firing rate of the electrocyte than those on the anterior side. We also note that potential effects of locally-constrained ion concentration changes have, in absence of data about such distributions, been neglected in our model. Constraining changes in ion concentration to specific subcellular locations could influence our results in both directions, either by local amplification or dampening of ion concentration changes (for example via efficient local buffering).

Taken together, our study demonstrates substantial effects of the Na$^+$/K$^+$-ATPase's electrogenicity on voltage dynamics in highly active excitable cells. While this property of the most common pump in the nervous system is assumed to serve as a mechanism preventing overexcitability, we show that it significantly interferes with cellular voltage dynamics via the immediate effects of its highly variable, hyperpolarising current – posing a particular challenge for highly active cells like electrocytes but also any other fast-spiking cell in nervous systems. This ultimately calls for strict regulatory mechanisms and may provide an additional evolutionary explanation for the abundance of differential ion channels and the diversity of pump isoforms expressed in excitable cell membranes to not only serve action potential generation, but also the stabilization of firing in biologically realistic environments.

6 Methods

6.1 Modelling an excitable cell with Na$^+$/K$^+$ pump

All simulations were done in `brian2` (93) with a time-step of 0.001ms. We simulated the weakly electric fish electrocyte model from (26), which we incrementally expanded by adding components relevant for modeling the Na$^+$/K$^+$ pump and corresponding ion concentration dynamics. All parameters were kept the same as in (26), except for $[\mathrm{Na}^+]_{\mathrm{in}}$, which was set to 13.5 mM, and $g_{\mathrm{Na}_{\max}}$ which was set to 900 μS.

First, the Na$^+$/K$^+$ pump current was added and a suitable co-expression factor between pump density and sodium leak channels was determined in order to counteract the depolarising effects of the pump current. Then, dynamical equations for the ion concentrations were added and the energetic demand at the baseline firing regime was estimated to tune pump densities to maintain steady state ion concentrations. Lastly, the feedback loop of pump density on ion concentrations was modeled to maintain ion homeostasis in firing regimes that deviate from baseline. Each of these steps is explained below in more detail.

6.1.1 Modeling the pump current and sodium leak channel co-expression

In previous work on the computational model of the weakly electric fish electrocyte, it was mentioned that there was an intention to include a Na$^+$/K$^+$ pump. This was however abstained from as the writer noted that adding an additional current I_{pump} would leave the model inexcitable. We identified the

effects of this pump current on electrocyte excitability, and propose additional currents to counteract these effects.

The Na$^+$/K$^+$ pump uses one ATP molecule to exchange three intracellular Na$^+$ ions for two extracellular K$^+$ ions (94). This leads to a net outflux of one positive ion every time the Na$^+$/K$^+$ pump performs an ion exchange. This net outflux is modeled as an additional current term in the membrane potential evolution equation

$$C\frac{dv}{dt} = -I_{\mathrm{NaT}} - I_{\mathrm{NaP}} - I_K - I_L - I_{\mathrm{pump}} + I_{\mathrm{stim}}(t).$$
(1)

In the present model, the Na$^+$/K$^+$ pump strength and thus I_{pump} does not depend on the membrane voltage, but on the intra- and extracellular ion concentrations (59). As ion concentration dynamics evolve on much longer timescales than the membrane potential, I_{pump} can in this case be assumed to be approximately constant on the timescale of the membrane potential.

When creating the firing rate (ω) *vs.* input ($\overline{I_{\mathrm{stim}}}$) curve ($f$-$I$ curve) of the electrocyte, where $\omega = f(\overline{I_{\mathrm{stim}}})$, an additional outward pump current creates a horizontal translation of ω, as $\omega = f(\overline{I_{\mathrm{stim}}} - I_{\mathrm{pump}})$. In other words, more inward $\overline{I_{\mathrm{stim}}}$ is needed to balance out the outward pump current and push the electrocyte to a firing regime.

To counterbalance the hyperpolarising outward current of the electrogenic Na$^+$/K$^+$ pump, we introduce an additional inward current. As the first approximation of the Na$^+$/K$^+$-ATPase current is constant, we modeled this balancing inward current as a relatively constant leak current

$$I_{\mathrm{NaL}} = g_{\mathrm{NaL}}(v - E_{\mathrm{Na}}).$$
(2)

To avoid confusion, we renamed the leak current already present in the model, which has a reversal potential equal to E_K, to I_{KL}

$$I_{\mathrm{KL}} = g_{\mathrm{KL}}(v - E_K).$$
(3)

For I_{NaL} to balance out I_{pump}, $I_{\mathrm{NaL}} = -I_{\mathrm{pump}}$ should hold for all t. In contrast to I_{pump} however, I_{NaL} is highly varying over time as it is dependent on v. One condition we can satisfy is for I_{NaL} to cancel out I_{pump} close to the rheobase of the f-I curve. This should render the firing onset of the electrocyte unchanged. We furthermore assume that g_{NaL} is adjusted at a larger timescale than I_{pump}, and that g_{NaL} is expressed to counteract I_{pump} at a baseline level I_{pump_0}. Setting $I_{\mathrm{NaL}_{\mathrm{rheobase}}} = I_{\mathrm{pump}_0}$ and rearranging to get g_{NaL} as a function of I_{pump_0} gives

$$g_{\mathrm{NaL}} = \frac{I_{\mathrm{pump}_0}}{v_{\mathrm{onset}} - E_{\mathrm{Na}}}.$$
(4)

Here, v_{onset} is the membrane voltage of the electrocyte just before firing onset. Upon injecting a

stimulus of 47nA, we find that v_{onset}=-81mV. After tuning g_{NaL} to counteract the effect of I_{pump_0} using equation 4, we find that the f-I curve is least affected by I_{pump_0} when slightly increasing v_{onset} to -76mV.

6.1.2 Modeling dynamic ion concentrations and deriving required pump densities for steady state ion concentrations

The function of the Na$^+$/K$^+$ pump is to maintain intra- and extracellular ion concentrations at fixed levels. If there were no Na$^+$/K$^+$ pump in an excitable cell, sodium ions would accumulate inside the cells and potassium ions would accumulate in extracellular space. This would reduce the concentration differences between ions in intra- and extracellular space, which impedes the firing of action potentials. The goal of the Na$^+$/K$^+$ pump is therefore to retain fixed sodium and potassium reversal potentials by maintaining ion homeostasis. This is achieved when the energetic supply, which would be the rearranging of ions by the Na$^+$/K$^+$ pump exactly equals the energetic demand which is the ion displacement caused by the action potentials.

In order to fully understand the influence of the Na$^+$/K$^+$ pump on cell excitability, we modeled the ion displacements of action potentials and the pump explicitly. To this end, we added ion concentration dynamics of intra- and extracellular sodium and potassium to the model equations (eq. (5)-eq. (8)) similarly to (59), and introduced a dependency of the reversal potentials on these ion concentrations via the Nernst equation (eqs 9, 10):

$$\frac{d[\mathrm{Na}^+]_{\mathrm{in}}}{dt} = \frac{-I_{\mathrm{NaT}} - I_{\mathrm{NaP}} - I_{\mathrm{NaL}} - \mathrm{syn}(t)I_{\mathrm{AChRNa}} - 3I_{\mathrm{pump}}}{FV_{\mathrm{in}}}, \tag{5}$$

$$\frac{d[\mathrm{K}^+]_{\mathrm{in}}}{dt} = \frac{-I_{\mathrm{K}} - I_{\mathrm{KL}} - \mathrm{syn}(t)I_{\mathrm{AChRK}} + 2I_{\mathrm{pump}}}{FV_{\mathrm{in}}}, \tag{6}$$

$$[\mathrm{Na}^+]_{\mathrm{out}} = [\mathrm{Na}^+]_{\mathrm{out}_0} - V_r([\mathrm{Na}^+]_{\mathrm{in}_0} - [\mathrm{Na}^+]_{\mathrm{in}}), \tag{7}$$

$$[\mathrm{K}^+]_{\mathrm{out}} = [\mathrm{K}^+]_{\mathrm{out}_0} - V_r([\mathrm{K}^+]_{\mathrm{in}_0} - [\mathrm{K}^+]_{\mathrm{in}}), \tag{8}$$

$$E_{\mathrm{Na}} = \frac{RT}{zF} \ln \frac{[\mathrm{Na}^+]_{in}}{[\mathrm{Na}^+]_{out}}, \tag{9}$$

$$E_{\mathrm{K}} = \frac{RT}{zF} \ln \frac{[\mathrm{K}^+]_{in}}{[\mathrm{K}^+]_{out}}. \tag{10}$$

Here, F is the Faraday constant and V_{in} is the intracellular volume which, as measured by (49) is roughly $0.424\,\mathrm{mm}^3$. V_r is the ratio between the volumes of the intra- and extracellular space. As electrocytes

are relatively large compared to their environment, we assumed V_r to be 2. Note that $I_{\text{stim}}(t)$ has been decomposed into $\text{syn}(t)I_{\text{AChRNa}}$ and $\text{syn}(t)I_{\text{AChRK}}$. This has been done to separately track the sodium and potassium displacement caused by the input stimulus.

To simplify the analysis, the model equations were rewritten to cancel out the state variable $[\text{K}^+]_{\text{in}}$, and have the model depend only on $[\text{Na}^+]_{\text{in}}$. First, we equate eq. (5) and eq. (6) to our rewritten version of the membrane potential equation

$$C\frac{dv}{dt} = -I_{\text{NaT}} - I_{\text{NaP}} - I_{\text{NaL}} - \text{syn}(t)I_{\text{AChRNa}} - I_K - I_{\text{KL}} - \text{syn}(t)I_{\text{AChRK}} - I_{\text{pump}}, \quad (11)$$

where all currents are separated into Na$^+$ and K$^+$ currents. This gives us

$$C\frac{dv}{dt} = FV_{\text{in}}\left(\frac{d[\text{Na}^+]_{\text{in}}}{dt} + \frac{d[\text{K}^+]_{\text{in}}}{dt}\right). \quad (12)$$

Integrating on both sides gives us

$$\frac{C\Delta v}{FV_{\text{in}}} = \Delta[\text{Na}^+]_{\text{in}} + \Delta[\text{K}^+]_{\text{in}}. \quad (13)$$

As the membrane conductance C is small, the Faraday constant F is very big, and the intracellular volume V_{in} is also relatively big, we can approximate $\frac{C\Delta v}{FV_{\text{in}}} \approx 0$. This proves that in our model, the macroscopic changes in intracellular ion concentrations can always be related by

$$\Delta[\text{Na}^+]_{\text{in}} \approx -\Delta[\text{K}^+]_{\text{in}}. \quad (14)$$

This allows us to rewrite equations 6 and 8 to depend only on state variable $[\text{Na}^+]_{\text{in}}$

$$[\text{K}^+]_{\text{in}} = [\text{K}^+]_{\text{in}_0} - ([\text{Na}^+]_{\text{in}_0} - [\text{Na}^+]_{\text{in}}), \quad (15)$$

$$[\text{K}^+]_{\text{out}} = [\text{K}^+]_{\text{out}_0} + V_r([\text{Na}^+]_{\text{in}_0} - [\text{Na}^+]_{\text{in}}). \quad (16)$$

With a perfectly working Na$^+$/K$^+$ pump, the macroscopic change in intracellular sodium $\Delta[\text{Na}^+]_{\text{in}}$ is zero, which signifies that the energetic supply of the pump exactly equals the energetic demand of the action potentials. From eqs 7, 15 and 16 we can conclude that if there is no macroscopic change in intracellular sodium, extracellular sodium concentrations and intra- and extracellular potassium concentrations also remain constant.

To facilitate long-term ion homeostasis in a high frequency firing electrocyte, we tune I_{pump} so that $\Delta[\text{Na}^+]_{\text{in}}$ (eq. 5) is zero i.e.

$$I_{\text{pump}} = \frac{1}{3} \sum \overline{I_{\text{Na}}}, \tag{17}$$

where $\sum \overline{I_{\text{Na}}}$ is the sum of the time average of all sodium currents. As $\sum \overline{I_{\text{Na}}}$ depends on I_{pump} due to the co-expression of pumps and sodium leak channels, we iteratively recompute the baseline I_{pump} according to equation 17 until the condition is satisfied with an error margin of 1 nA.

6.1.3   Modeling the feedback loop of ion concentrations on pump density

Assuming that pump densities are tuned to sustain a fixed baseline firing rate, deviations from this baseline firing will lead to a mismatch between the ion displacement caused by action potentials and the ion restoration of the Na$^+$/K$^+$ pump. This will lead to a shift in intra- and extracellular ion concentrations. As the pump rate and thus I_{pump} is a function of intra- and extracellular ion concentrations (59), the pump rate will adjust accordingly. We model the dependency of I_{pump} on intracellular sodium concentrations $[\text{Na}^+]_{\text{in}}$ and extracellular potassium concentrations $[\text{K}^+]_{\text{out}}$ similarly to (59)

$$I_{\text{pump}} = \frac{4 I_{\text{pump}_0}}{\left(1 + e^{\frac{[\text{Na}^+]_{\text{in}_0} - [\text{Na}^+]_{\text{in}}}{3}}\right)\left(1 + e^{[\text{K}^+]_{\text{out}_0} - [\text{K}^+]_{\text{out}}}\right)}. \tag{18}$$

For simplicity, we adjusted the terms within the exponents so that $I_{\text{pump}} = I_{\text{pump}_0}$ when $\Delta[\text{Na}^+]_{\text{in}} = 0$. Here, I_{pump_0} is the pump current that is tuned to facilitate ion homeostasis at the baseline firing rate. As the pump current saturates at $4 I_{\text{pump}_0}$, which is proportional to the number of Na$^+$/K$^+$-ATPases that are expressed, the baseline pump current $4 I_{\text{pump}_0}$ is also proportional to the pump density. A shift in I_{pump}, which can now happen as a consequence of deviations from baseline firing, without co-expression of g_{NaL}, which is unlikely on small timescales, will lead to a shift in cell excitability.

6.1.4   Modeling an optimal voltage-dependence of the pump

Dynamics of action potential firing would be unaffected by the presence of voltage-dependent electrogenic pumps if the membrane voltage would modulate their activity in a way that the pump current mimics hyperpolaring voltage-gated and leaky potassium current. We substantiate this idea by modeling a voltage-dependence of the pump that copies the dynamics of potassium currents. To achieve this, we rewrite the baseline pump current I_{pump_0} as a function of the membrane voltage, and a transformation of the membrane voltage that takes into account the history of the membrane voltage (n)

$$I_{\text{pump}_0} = \frac{1}{3}(g_{\text{K}_{\text{L}}} + g_{\text{K}_{\text{max}}} n^4)(v - E_{\text{K}}), \tag{19}$$

which is essentially a scaled version of a combination of the equations that describe the voltage-gated and leaky potassium currents. As the pump current now behaves like $\frac{1}{3}$ of the potassium currents, we can reduce the potassium conductances by $\frac{1}{3}$ and still get qualitatively the same APs, through

$$I_{\mathrm{K}} = \frac{2}{3} g_{\mathrm{K_{max}}} n^4 (v - E_{\mathrm{K}}), \tag{20}$$

and

$$I_{\mathrm{K_L}} = \frac{2}{3} g_{\mathrm{K_L}} (v - E_{\mathrm{K}}). \tag{21}$$

By restructuring the current and pump equations as such, the pump current should always equal $\frac{1}{2}$ of the potassium currents, which exactly satisfies the energetic demand of the cell. There is however a third potassium current, I_{AChRK}, that is activated by neurotransmitter release. As we cannot expect the pumps to also be sensitive to neurotransmitter release, the voltage-dependent pump described in equation 19 will pump slightly less than necessary to maintain ionic homeostasis. We therefore let the model run until steady state ion concentrations were reached, which happens in close proximity to the baseline concentration of $[\mathrm{Na}^+]_{\mathrm{in}} = 13.500$ mM, which is $[\mathrm{Na}^+]_{\mathrm{in}} = 13.517$ mM.

6.2 Effects of pump-mediated cell excitability in a (very simple) network

To showcase the effects of altered cell excitability through a dynamic pump current in a real-world scenario, we modeled its effects in the simplest network possible, which is an excitable cell that is coupled to a pacemaker. In this scenario, there is only one excitatory connection from the pacemaker to the excitable cell. We did not explicitly model the pacemaker, but only the time dependent currents that would result from the excitatory synapse, as implemented by (26).

6.2.1 Generation of communication signals

Synaptic inputs to the electrocyte were modeled as in (26), where the parameter that reflects synaptic conductance $syn_{\mathrm{clamp}}(t_p)$ is modeled by a piecewise function that resets $t_p \to 0$ for every spike that arrives from the PN;

$$syn_{\mathrm{clamp}}(t_p) = \begin{cases} \frac{t_p}{0.05}, & t_p \leq 0.05 \\ 1, & 0.05 < t_p \leq 0.25 \\ e^{\frac{-(t_p - 0.25)}{0.1}}, & t_p > 0.25. \end{cases} \tag{22}$$

As we aimed to show scenarios in which communication signals have maximal effect on the pump current, we modeled relatively long chirps in a fish with high baseline firing rates, and relatively high frequency rises in a fish with low baseline firing rates. Both baseline firing rates and chirp duration and frequency rise amplitude and timescale are representative of EOD signals found in experimental settings (29).

To model chirps, the reset frequency of t_p was set to 600 Hz, and after 100 ms, chirps were generated where t_p was only reset after a period of 20 times the baseline ISI. After this period, t_p was again reset with a frequency 600 Hz for 100 ms. This was repeated 10 times to simulate 10 consecutive chirps.

To model frequency rises, the reset frequency of t_p was set to 260 Hz, and after 6 seconds, frequency rises were generated where t_p was reset with a frequency set by following formula

$$\omega_{t_p} = \omega_r e^{\frac{-t}{\tau}} + \omega_0, \tag{23}$$

where ω_r is the amplitude of the frequency rise, which was set to 40 Hz, t is the passed time since the onset of the frequency rise, τ is the timescale of the frequency rise which was set to 1 second, and ω_0 is the baseline frequency which was set to 260 Hz. Frequency rises were initiated every 2 seconds 10 times in a row.

6.2.2 Parameter tuning

When simulating the communication paradigms, it was assumed that pump densities and voltage-gated sodium channel densities were optimised to support baseline firing rates. Therefore, pump densities were tuned to perfectly maintain ion homeostasis for baseline firing, and voltage-gated sodium channels were tuned to allow for sufficient spike amplitudes ($\approx$ 13mV) at baseline firing.

6.2.3 Spike entrainment measure

To quantify the accuracy of entrainment between spikes emmited by PN at times, $t_1^{\mathrm{p}} < ... < t_m^{\mathrm{p}}$ and spikes emmited by electrocyte at times $t_1^{\mathrm{e}} < ... < t_n^{\mathrm{e}}$, where $t_n^{\mathrm{e}} < t_m^{\mathrm{p}}$.

Let $j(t) = j : t_j^{\mathrm{p}} < t < t_{j+1}^{\mathrm{p}}$ denote the index of the PN spike interval in which a given electrocyte spike time resides. Then define the phases of electrocyte spikes relative to pacemaker inter spike intervals as

$$\theta_k = \frac{t_k^{\mathrm{e}} - t_{j(t_k)}^{p}}{t_{j(t_k)+1}^{p} - t_{j(t_k)}^{p}}.$$

The rigidity of the entrainment phase is then quantified by the circular variance as the mean resultant length

$$\bar{R} = \frac{1}{n} \left| \sum_{k=1}^{n} \exp i 2\pi \theta_k \right|.$$

This score, which is referred to as the entrainment index, is shown on top of spike trains in figures 5 (C,D,F,G).

6.2.4 Mechanisms that improve entrainment

Extracellular potassium buffering was implemented to decrease the variation in ΔI_{pump} and thereby the variation in mean-driven electrocyte properties, which is beneficial for entrainment. As we were

mainly interested in the effects of extracellular potassium buffering on the feedback loop of the pump current (eq 18), and did at this stage not put importance on the transient effects of the potassium buffer, we added a very simple 'instant' potassium buffer by setting

$$[\mathrm{K}^+]_{\mathrm{out}} = [\mathrm{K}^+]_{\mathrm{out}_0}. \tag{24}$$

Synaptic coupling, which also affects entrainment, was modeled by adding a pre-factor to $\mathrm{syn}(t)$, where weak coupling (Fig. 5 B-D) was modeled by setting

$$\mathrm{syn}(t) = \frac{1}{2}\mathrm{syn}(t),$$

and strong coupling was modeled by the original synaptic strength

$$\mathrm{syn}(t).$$

7   Data availability statement

All results presented in this article can be reproduced via the code published at https://itbgit.biologie. hu-berlin.de/compneurophys_pub/electrocyte_nakatpase.

8   Author contributions

Conceptualization, L.W., J-H.S and S.S.; Methodology, L.W. and J-H.S.; Software, L.W.; Writing – Original Draft, L.W., J-H.S., S.S.; Writing, – Review & Editing, L.W., J-H.S. and S.S.; Supervision, J-H.S. and S.S.; Funding Acquisition, S.S.

9   Declaration of interests

The authors declare no competing interests.

10   Acknowledgments

We thank Dr. Louisiane Lemaire and Mahraz Behbood for fruitful discussions.

This project has received funding from the Einstein Foundation Berlin (grant number EP-2021-621).

11 Appendix

11.1 Phase oscillator theory to quantify entrainment

The observation that each PN spike typically causes one electrocyte spike suggests that electrocytes are excitable cells, kicked above threshold by synaptic PN inputs and then return to rest. However, taking into account the sustained high frequency PN firing rates, ω_{pn}, and the kinetics of the electrocyte acetylcholine receptor shows that the model is better understood as a mean driven entrained oscillator. To see this, the membrane voltage evolution equation of the electrocyte is rewritten to separate the stimulus current into a time averaged DC component, eq. (25), and a time-dependent zero mean component, eq. (26), by averaging over one PN period

$$\overline{I_{\mathrm{stim}}} = \omega_{\mathrm{pn}} \int_0^{\frac{1}{\omega_{\mathrm{pn}}}} I_{\mathrm{stim}}(t)\, dt. \tag{25}$$

As will be shown below, in many cases the mean drive $\overline{I_{\mathrm{stim}}}$ is sufficient to induce spiking the the electrocyte. The model then reads

$$C\frac{dv}{dt} = \underbrace{-I_{\mathrm{NaT}} - I_{\mathrm{NaP}} - I_K - I_L + \overline{I_{\mathrm{stim}}}}_{\text{mean driven electrocyte}} + \underbrace{(I_{\mathrm{stim}}(t) - \overline{I_{\mathrm{stim}}})}_{\text{time dependent PN input}}. \tag{26}$$

We can now characterise the mean driven electrocyte by the relation of its firing rate to mean input in terms of its f-I curve (Fig. 3 E). We can furthermore determine the time averaged stimulus for various PN driving frequencies (Fig. 3 F). When comparing the mean driven frequency of the electrocyte, ω_e, to the PN driving frequency, ω_{pn}, we find mismatches in frequencies that can go up to 180 Hz (Fig. 3 G). As proven by (26), the electrocyte model can be entrained by ω_{pn} ranging from 200 to 600 Hz. We can therefore assume that the input stimulus is strong enough to overcome large frequency mismatches between ω_e and ω_{pn}.

To quantify the allowed frequency mismatches between ω_e and ω_{pn} for synchronization, we treat the mean-driven electrocyte as a periodic oscillator that is to be entrained by an external force, which is the PN. The evolution of the phase of the electrocyte can now be expressed as

$$\frac{d\phi}{dt} = \omega_e + Z(\phi)x(t). \tag{27}$$

Here, ω_e is the frequency of the mean driven electrocyte, $Z(\phi)$ is the change in phase of the electrocyte evoked by perturbations of the membrane potential, and $x(t)$ is the time-dependent perturbation caused by the zero mean pacemaker input stimulus

$$x(t) = \frac{1}{C}(I_{\mathrm{stim}}(t) - \overline{I_{\mathrm{stim}}}). \tag{28}$$

One can now define a variable that describes the phase difference between the electrocyte and the PN as

$$\psi = \phi - \omega_{\mathrm{pn}}t. \tag{29}$$

If there exists a phase ψ for which ψ does not change over time i.e. $\frac{d\psi}{dt} = 0$, the electrocyte and the PN will be phase-locked, or entrained. The equation for evolution of the phase difference is obtained by plugging eq 29 in eq 27

$$\frac{d\psi}{dt} = \omega_e - \omega_{\mathrm{pn}} + Z(\psi + \omega_{\mathrm{pn}}t)x(t). \tag{30}$$

As $\psi \ll \omega_{\mathrm{pn}}t$ is a slow variable, one period of ψ will 'see' a lot of PN periods. Hence, the method of averaging yields

$$\frac{d\psi}{dt} = \omega_e - \omega_{\mathrm{pn}} + \omega_{\mathrm{pn}} \int_0^{\frac{1}{\omega_{\mathrm{pn}}}} Z(\psi + \omega_{\mathrm{pn}}t)x(t)\, dt. \tag{31}$$

From this equations the minimum and maximum ω_{pn} for which the electrocyte and the PN can be phase-locked is defined as

$$\omega_{\mathrm{pn_{max}}} = \omega_e + \omega_{\mathrm{pn}} \max_{0 \leq \psi \leq 1} \int_0^{\frac{1}{\omega_{\mathrm{pn}}}} Z(\psi + \omega_{\mathrm{pn}}t)x(t)\, dt, \tag{32}$$

and

$$\omega_{\mathrm{pn_{min}}} = \omega_e + \omega_{\mathrm{pn}} \min_{0 \leq \psi \leq 1} \int_0^{\frac{1}{\omega_{\mathrm{pn}}}} Z(\psi + \omega_{\mathrm{pn}}t)x(t)\, dt. \tag{33}$$

The range $[\omega_{\mathrm{pn_{min}}}, \omega_{\mathrm{pn_{max}}}]$ is referred to as the entrainment range.

The mean-driven features ω_e and $Z(\phi)$ are altered upon a baseline deviation of the pump current ΔI_{pump} (Fig. A1 A, B), which is in essence the same as a deviation in $\overline{I_{\mathrm{stim}}}$. ΔI_{pump} thus affects $[\omega_{\mathrm{pn_{min}}}, \omega_{\mathrm{pn_{max}}}]$ and thereby the entrainment of the pacemaker-driven electrocyte (Fig. A1 C).

Figure A1: A deviation in pump current ΔI_{pump} alters mean-driven electrocyte properties and thereby its entrainment region. *(A) Mean-driven electrocyte frequency ω_e as a function of ΔI_{pump}. (B) Phase Response Curves (PRCs, $Z(\phi)$) as a function of ΔI_{pump}. (C) The entrainment range $[\omega_{pn_{min}}, \omega_{pn_{max}}]$, which is a function of mean-driven electrocyte properties (A, B, eqs 33, 32), changes upon deviations in pump current. For very strong deviations in I_{pump}, the pacemaker frequency ω_{pn} falls out of the entrainment range which means that the electrocyte will not lock to the pacemaker in this regime.*

The influence of the synaptic weight on entrainment (Fig. 5 B-G) can also be explained by equations 33, 32, as it increases x(t) and therefore $[\omega_{pn_{min}}, \omega_{pn_{max}}]$.

11.1.1 Phase Response Curves

To solve eqs 32 and 33, PRCs need to be computed. PRCs were obtained for various $\overline{I_{stim}}$ by first injecting constant input current $\overline{I_{stim}}$ to the electrocyte. Then, the phase of the electrocyte was defined as the peak of one spike in the tonically firing electrocyte to the next peak. The membrane voltage was then perturbed by 1mV at 20 different phases linearly interpolated between 0 and 1, and the resulting time delays or advances of the next spike was recorded as the phase response.

11.2 Relation between integrated sodium and potassium currents over one ISI

Let's assume all voltage dynamics are described by the sum of sodium and potassium currents:

$$C\frac{dv}{dt} = -I_{\mathrm{Na}} - I_{\mathrm{K}}. \tag{34}$$

If v is in a limit cycle with length T,

$$v(t) = v(t+T)\forall t,$$

and

$$v(t) - v(t+T) = 0\forall t. \tag{35}$$

If we integrate the voltage equation (Eq. 34) on both sides from t to $t+T$ we get

$$C[v(t+T) - v(t)] = -\int_t^{t+T} I_{\mathrm{Na}}(t)dt - \int_t^{t+T} I_{\mathrm{K}}(t)dt. \tag{36}$$

If we plug in eq 35, we get

$$\int_t^{t+T} I_{\mathrm{Na}}(t)dt + \int_t^{t+T} I_{\mathrm{K}}(t)dt = 0, \tag{37}$$

which means that the integrated sodium and potassium currents over the time course of one ISI should always be equal to each other.

References

1. Lane N, Martin WF. The origin of membrane bioenergetics. Cell [Internet]. 2012 Dec;151:1406–16. Available from: http://dx.doi.org/10.1016/j.cell.2012.11.050

2. Dibrova DV, Galperin MY, Koonin EV, Mulkidjanian AY. Ancient systems of sodium/potassium homeostasis as predecessors of membrane bioenergetics. Biochemistry (Moscow) [Internet]. 2015 May;80:495–516. Available from: http://dx.doi.org/10.1134/s0006297915050016

3. Palmgren M. Evolution of the sodium pump. Biochimica et Biophysica Acta (BBA)-Molecular Cell Research. 2023;119511.

4. Stein WD. The sodium pump in the evolution of animal cells. Philosophical Transactions of the Royal Society of London Series B: Biological Sciences. 1995;349(1329):263–9.

5. Contreras SA, Schleimer JH, Gulledge AT, Schreiber S. Activity-mediated accumulation of potassium induces a switch in firing pattern and neuronal excitability type. PLoS Computational Biology. 2021;17(5):e1008510.

6.   Thomas R. Electrogenic sodium pump in nerve and muscle cells. Physiological reviews. 1972;52(3):563–94.

7.   Ritchie J, Straub R. The hyperpolarization which follows activity in mammalian non-medullated fibres. The Journal of Physiology. 1957;136(1):80.

8.   Connelly C. Recovery processes and metabolism of nerve. Reviews of modern physics. 1959;31(2):475.

9.   Nakajima S, Onodera K. Membrane properties of the stretch receptor neurones of crayfish with particular reference to mechanisms of sensory adaptation. The Journal of Physiology. 1969;200(1):161–85.

10.   Sokolove P, Cooke I. Inhibition of impulse activity in a sensory neuron by an electrogenic pump. The Journal of General Physiology. 1971;57(2):125–63.

11.   Sawczuk A, Powers RK, Binder MD. Contribution of outward currents to spike-frequency adaptation in hypoglossal motoneurons of the rat. Journal of Neurophysiology. 1997;78(5):2246–53.

12.   Gustafsson B, Wigström H. Hyperpolarization following long-lasting tetanic activation of hippocampal pyramidal cells. Brain research. 1983;275(1):159–63.

13.   Arganda S, Guantes R, De Polavieja GG. Sodium pumps adapt spike bursting to stimulus statistics. Nature neuroscience. 2007;10(11):1467–73.

14.   Forrest MD. The sodium-potassium pump is an information processing element in brain computation. Vol. 5, Frontiers in physiology. Frontiers Media SA; 2014. p. 472.

15.   Zylbertal A, Yarom Y, Wagner S. The slow dynamics of intracellular sodium concentration increase the time window of neuronal integration: A simulation study. Frontiers in computational neuroscience. 2017;11:85.

16.   Megwa OF, Pascual LM, Günay C, Pulver SR, Prinz AA. Temporal dynamics of na/k pump mediated memory traces: Insights from conductance-based models of drosophila neurons. Frontiers in Neuroscience. 2023;17:1154549.

17.   Benda J, Longtin A, Maler L. Spike-frequency adaptation separates transient communication signals from background oscillations. Journal of Neuroscience. 2005;25(9):2312–21.

18.   Benda J. Neural adaptation. Current Biology. 2021;31(3):R110–6.

19.   Kueh D, Barnett WH, Cymbalyuk GS, Calabrese RL. Na+/k+ pump interacts with the h-current to control bursting activity in central pattern generator neurons of leeches. Elife. 2016;5:e19322.

20.   Zylbertal A, Yarom Y, Wagner S. Synchronous infra-slow bursting in the mouse accessory olfactory bulb emerge from interplay between intrinsic neuronal dynamics and network connectivity. Journal of Neuroscience. 2017;37(10):2656–72.

21.   Behbood M, Lemaire L, Schleimer JH, Schreiber S. The na+/k+-ATPase generically enables deterministic bursting in class i neurons by shearing the spike-onset bifurcation structure. PLOS Computational Biology. 2024;20(8):e1011751.

22.   Gulledge AT, Dasari S, Onoue K, Stephens EK, Hasse JM, Avesar D. A sodium-pump-mediated afterhyperpolarization in pyramidal neurons. Journal of Neuroscience. 2013;33(32):13025–41.

23.   Morotti S, Clair JRS, Proenza C, Grandi E. Dynamic regulation of pacemaker activity by the na+-k+ pump. In: 2016 computing in cardiology conference (CinC). IEEE; 2016. p. 765–8.

24.   Niemeyer N, Schleimer JH, Schreiber S. Biophysical models of intrinsic homeostasis: Firing rates and beyond. Current Opinion in Neurobiology. 2021;70:81–8.

25.   Hodgkin AL, Huxley AF, Katz B. Measurement of current-voltage relations in the membrane of the giant axon of loligo. The Journal of physiology. 1952;116(4):424–48.

26.   Joos B, Markham MR, Lewis JE, Morris CE. A model for studying the energetics of sustained high frequency firing. PLOS ONE [Internet]. 2018 Apr;13:e0196508. Available from: http://dx.doi.org/10.1371/journal.pone.0196508

27.   Dunlap KD, Silva AC, Smith GT, Zakon HH. Weakly electric fish: Behavior, neurobiology, and neuroendocrinology. In: Hormones, Brain and Behavior [Internet]. Elsevier; 2017. p. 69–98. Available from: http://dx.doi.org/10.1016/b978-0-12-803592-4.00019-5

28.   Bullock TH. Electroreception. Annual review of neuroscience. 1982;

29.   Hopkins CD. Electric communication: Functions in the social behavior of eigenmannia virescens. Behaviour. 1974;50(3-4):270–304.

30.   Attwell D, B. Laughlin S. An energy budget for signaling in the grey matter of the brain: Journal of Cerebral Blood Flow & Metabolism [Internet]. 2016 Aug 31; Available from: https://journals.sagepub.com/doi/10.1097/00004647-200110000-00001

31.   Salazar VL, Krahe R, Lewis JE. The energetics of electric organ discharge generation in gymnotiform weakly electric fish. Journal of Experimental Biology. 2013;216(13):2459–68.

32.   Chung KF, Lin HC. Osmoregulation and na,k-ATPase expression in osmoregulatory organs of scylla paramamosain. Comparative Biochemistry and Physiology Part A: Molecular \& Integrative Physiology [Internet]. 2006 May;144:48–57. Available from: http://dx.doi.org/10.1016/j.cbpa.2006.02.003

33.   Meunier C, Segev I. Playing the devil's advocate: Is the hodgkin–huxley model useful? TRENDS in Neurosciences. 2002;25(11):558–63.

34.   Hübel N, Schöll E, Dahlem MA. Bistable dynamics underlying excitability of ion homeostasis in neuron models. PLoS computational biology. 2014;10(5):e1003551.

35.   Weer PD, Gadsby DC, Rakowski R. Voltage dependence of the na-k pump. Annual Review of Physiology. 1988;50(1):225–41.

36.   Wuddel I, Apell HJ. Electrogenicity of the sodium transport pathway in the na, k-ATPase probed by charge-pulse experiments. Biophysical journal. 1995;69(3):909–21.

37.   Gadsby DC, Bezanilla F, Rakowski RF, De Weer P, Holmgren M. The dynamic relationships between the three events that release individual na+ ions from the na+/k+-ATPase. Nature communications. 2012;3(1):669.

38. Garcia A, Rasmussen HH, Apell HJ, Clarke RJ. Kinetic comparisons of heart and kidney na+, k+-ATPases. Biophysical journal. 2012;103(4):677–88.

39. Stanley CM, Gagnon DG, Bernal A, Meyer DJ, Rosenthal JJ, Artigas P. Importance of the voltage dependence of cardiac na/k ATPase isozymes. Biophysical Journal. 2015;109(9):1852–62.

40. Squire L, Berg D, Bloom FE, Du Lac S, Ghosh A, Spitzer NC. Fundamental neuroscience. Academic press; 2012.

41. O'Leary T, Williams AH, Caplan JS, Marder E. Correlations in ion channel expression emerge from homeostatic tuning rules. Proceedings of the National Academy of Sciences. 2013;110(28):E2645–54.

42. Yu Y, Hill AP, McCormick DA. Warm body temperature facilitates energy efficient cortical action potentials. PLoS computational biology. 2012;8(4):e1002456.

43. Hildebrandt KJ, Benda J, Hennig RM. Multiple arithmetic operations in a single neuron: The recruitment of adaptation processes in the cricket auditory pathway depends on sensory context. Journal of Neuroscience. 2011;31(40):14142–50.

44. Hürkey S, Niemeyer N, Schleimer JH, Ryglewski S, Schreiber S, Duch C. Gap junctions desynchronize a neural circuit to stabilize insect flight. Nature [Internet]. 2023 Jun;618:118–25. Available from: http://dx.doi.org/10.1038/s41586-023-06099-0

45. Barreto E, Cressman JR. Ion concentration dynamics as a mechanism for neuronal bursting. Journal of biological physics. 2011;37:361–73.

46. Kager H, Wadman WJ, Somjen GG. Simulated seizures and spreading depression in a neuron model incorporating interstitial space and ion concentrations. Journal of neurophysiology. 2000;84(1):495–512.

47. Donnan FG. The theory of membrane equilibria. Chemical reviews. 1924;1(1):73–90.

48. Crambert G, Hasler U, Beggah AT, Yu C, Modyanov NN, Horisberger JD, et al. Transport and pharmacological properties of nine different human na, k-ATPase isozymes. Journal of Biological Chemistry. 2000;275(3):1976–86.

49. Ban Y, Smith BE, Markham MR. A highly polarized excitable cell separates sodium channels from sodium-activated potassium channels by more than a millimeter. Journal of neurophysiology. 2015;114(1):520–30.

50. Bullock TH. The reliability of neurons. The Journal of general physiology. 1970;55(5):565–84.

51. Moortgat KT, Bullock TH, Sejnowski TJ. Precision of the pacemaker nucleus in a weakly electric fish: Network versus cellular influences. Journal of Neurophysiology [Internet]. 2000 Feb;83:971–83. Available from: http://dx.doi.org/10.1152/jn.2000.83.2.971

52. Black-Cleworth P. The role of electrical discharges in the non-reproductive social behaviour of gymnotus carapo (gymnotidae, pisces). Animal Behaviour Monographs. 1970;3:1–IN1.

53. Westby GM. Further analysis of the individual discharge characteristics predicting social dominance in the electric fish, gymnotus carapo. Animal Behaviour. 1975;23:249–60.

54.   Westby GM, Box HO. Prediction of dominance in social groups of the electric fish, gymnotus carapo. Psychonomic Science. 1970;21(3):181–3.

55.   Max Westby G. Assessment of the signal value of certain discharge patterns in the electric fish, gymnotus carapo, by means of playback. Journal of comparative physiology. 1974;92:327–41.

56.   Szabo T, Enger PS. Pacemaker activity of the medullary nucleus controlling electric organs in high-frequency gymnotid fish. Zeitschrift für vergleichende Physiologie. 1964;49:285–300.

57.   Stöckl A, Sinz F, Benda J, Grewe J. Encoding of social signals in all three electrosensory pathways of eigenmannia virescens. Journal of neurophysiology. 2014;112(9):2076–91.

58.   Erecińska M, Dagani F. Relationships between the neuronal sodium/potassium pump and energy metabolism. Effects of k+, na+, and adenosine triphosphate in isolated brain synaptosomes. The Journal of general physiology. 1990;95(4):591–616.

59.   Hübel N, Dahlem MA. Dynamics from seconds to hours in hodgkin-huxley model with time-dependent ion concentrations and buffer reservoirs. PLoS computational biology. 2014;10(12):e1003941.

60.   Hagedorn M, Heiligenberg W. Court and spark: Electric signals in the courtship and mating of gymnotoid fish. Animal Behaviour. 1985;33(1):254–65.

61.   Pikovsky A, Rosenblum M, Kurths J. Synchronization: A universal concept in nonlinear science. American Association of Physics Teachers; 2002.

62.   Schreiber S, Fellous JM, Tiesinga P, Sejnowski TJ. Influence of ionic conductances on spike timing reliability of cortical neurons for suprathreshold rhythmic inputs. Journal of neurophysiology. 2004;91(1):194–205.

63.   Fisher NI. Statistical analysis of circular data. Cambridge University Press; 1993.

64.   Pikovsky A, Rosenblum M, Kurths J, Synchronization A. A universal concept in nonlinear sciences. Self. 2001;2:3.

65.   Colina C, Palavicini JP, Srikumar D, Holmgren M, Rosenthal JJ. Regulation of na+/k+ ATPase transport velocity by RNA editing. PLoS biology. 2010;8(11):e1000540.

66.   Person AL, Raman IM. Purkinje neuron synchrony elicits time-locked spiking in the cerebellar nuclei. Nature. 2012;481(7382):502–5.

67.   Gittis AH, Moghadam SH, Du Lac S. Mechanisms of sustained high firing rates in two classes of vestibular nucleus neurons: Differential contributions of resurgent na, Kv3, and BK currents. Journal of neurophysiology. 2010;104(3):1625–34.

68.   Roth FC, Hu H. An axon-specific expression of HCN channels catalyzes fast action potential signaling in GABAergic interneurons. Nature Communications. 2020;11(1):2248.

69.   Anderson TR, Huguenard JR, Prince DA. Differential effects of na+–k+ ATPase blockade on cortical layer v neurons. The Journal of physiology. 2010;588(22):4401–14.

70.   Hodgkin AL, Keynes R. Active transport of cations in giant axons from sepia and loligo. The Journal of physiology. 1955;128(1):28.

71. Hodgkin A, Keynes R. Experiments on the injection of substances into squid giant axons by means of a microsyringe. The Journal of Physiology. 1956;131(3):592.

72. Straub R. MECHANISM OF POST-TETANIC HYPERPOLARIZATION IN MYELINATED NERVE FIBRES FROM FROG. In: JOURNAL OF PHYSIOLOGY-LONDON. CAMBRIDGE UNIV PRESS 40 WEST 20TH STREET, NEW YORK, NY 10011-4211; 1961. p. P19.

73. Nakajima S, Takahashi K. Post-tetanic hyperpolarization and electrogenic na pump in stretch receptor neurone of crayfish. The Journal of Physiology. 1966;187(1):105.

74. Gage P, Hubbard J. The origin of the post-tetanic hyperpolarization of mammalian motor nerve terminals. The Journal of physiology. 1966;184(2):335–52.

75. Vedovato N, Gadsby DC. Route, mechanism, and implications of proton import during na+/k+ exchange by native na+/k+-ATPase pumps. The Journal of general physiology. 2014;143(4):449.

76. Liu H, Wu M-m, Zakon HH. A novel na+ channel splice form contributes to the regulation of an androgen-dependent social signal. Journal of Neuroscience [Internet]. 2008 Sep;28:9173–82. Available from: http://dx.doi.org/10.1523/jneurosci.2783-08.2008

77. Berndt N, Holzhütter HG. The high energy demand of neuronal cells caused by passive leak currents is not a waste of energy. Cell biochemistry and biophysics. 2013;67:527–35.

78. Zakon HH, Lu Y, Zwickl DJ, Hillis DM. Sodium channel genes and the evolution of diversity in communication signals of electric fishes: Convergent molecular evolution. Proceedings of the National Academy of Sciences [Internet]. 2006 Mar;103:3675–80. Available from: http://dx.doi.org/10.1073/pnas.0600160103

79. Blanco G, Mercer RW. Isozymes of the na-k-ATPase: Heterogeneity in structure, diversity in function. American Journal of Physiology-Renal Physiology. 1998;275(5):F633–50.

80. Carvalho Aguiar P de, Sweadner KJ, Penniston JT, Zaremba J, Liu L, Caton M, et al. Mutations in the na+/k+-ATPase α3 gene ATP1A3 are associated with rapid-onset dystonia parkinsonism. Neuron. 2004;43(2):169–75.

81. Heinzen EL, Swoboda KJ, Hitomi Y, Gurrieri F, Nicole S, Vries B de, et al. De novo mutations in ATP1A3 cause alternating hemiplegia of childhood. Nature genetics. 2012;44(9):1030–4.

82. Rosewich H, Thiele H, Ohlenbusch A, Maschke U, Altmüller J, Frommolt P, et al. Heterozygous de-novo mutations in ATP1A3 in patients with alternating hemiplegia of childhood: A whole-exome sequencing gene-identification study. The Lancet Neurology. 2012;11(9):764–73.

83. Demos MK, Karnebeek CD van, Ross CJ, Adam S, Shen Y, Zhan SH, et al. A novel recurrent mutation in ATP1A3 causes CAPOS syndrome. Orphanet journal of rare diseases. 2014;9:1–9.

84. Yu FH, Mantegazza M, Westenbroek RE, Robbins CA, Kalume F, Burton KA, et al. Reduced sodium current in GABAergic interneurons in a mouse model of severe myoclonic epilepsy in infancy. Nature neuroscience. 2006;9(9):1142–9.

85. Sasaki M, Ishii A, Saito Y, Morisada N, Iijima K, Takada S, et al. Genotype–phenotype correlations in alternating hemiplegia of childhood. Neurology. 2014;82(6):482–90.

86. Soltoff SP, Mandel LJ. Active ion transport in the renal proximal tubule. III. The ATP dependence of the na pump. The Journal of general physiology. 1984;84(4):643–62.

87.   Crampton WG. Electric signal design and habitat preferences in a species rich assemblage of gymnotiform fishes from the upper amazon basin. Anais-Academia Brasileira de Ciencias. 1998;70:805–48.

88.   Reardon EE, Parisi A, Krahe R, Chapman LJ. Energetic constraints on electric signalling in wave-type weakly electric fishes. Journal of Experimental Biology [Internet]. 2011 Dec;214:4141–50. Available from: http://dx.doi.org/10.1242/jeb.059444

89.   Rueda-Delgado G, Wantzen KM, Tolosa MB. Leaf-litter decomposition in an amazonian flood-plain stream: Effects of seasonal hydrological changes. Journal of the North American Benthological Society. 2006;25(1):233–49.

90.   Alonso LM, Marder E. Temperature compensation in a small rhythmic circuit. eLife [Internet]. 2020 Jun;9. Available from: http://dx.doi.org/10.7554/elife.55470

91.   Yu Y, P. Hill A, A. McCormick D. Warm body temperature facilitates energy efficient cortical action potentials. PLOS Computational Biology [Internet]. 8. Available from: https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1002456

92.   Roemschied FA, Eberhard MJ, Schleimer JH, Ronacher B, Schreiber S. Cell-intrinsic mechanisms of temperature compensation in a grasshopper sensory receptor neuron. Elife. 2014;3:e02078.

93.   Stimberg M, Brette R, Goodman DF. Brian 2, an intuitive and efficient neural simulator. elife. 2019;8:e47314.

94.   Guyton AC, Hall JE, et al. Textbook of medical physiology. Vol. 548. Saunders Philadelphia; 1986.