Chemical signatures of social information in Barbary macaques

Brigitte M. Weiß[1,2*], Claudia Birkemeyer[3], Marlen Kücklich[1,2°] & Anja Widdig[1,2°]

[1] Behavioural Ecology Group, Leipzig University, Leipzig, Germany

[2] Department of Human Behaviour, Ecology and Culture, Max-Planck-Institute for Evolutionary

Anthropology, Leipzig, Germany

[3] now Claudia Wiesner, Mass Spectrometry Group, Institute of Analytical Chemistry, Leipzig

University, Leipzig, Germany

°contributed equally

*Corresponding author: Brigitte M. Weiß, Institute of Biology, Talstraße 33, 04103 Leipzig, Germany,

phone: ++49-341-9736-754, email: brigitte.schloegl@uni-leipzig.de

Word count: 4505

<u>Abstract</u>

Primates are well-known for their complex social lives and intricate social relationships, which

requires them to obtain and update social knowledge about conspecifics. The sense of smell may

provide access to social information that is unavailable in other sensory domains or enhance the

precision and reliability of other sensory cues. However, the cognition of social information in

catarrhine primates has been studied primarily in the visual and auditory domain. We assessed the

social information content of body odor in a catarrhine primate, the Barbary macaque (*Macaca*

sylvanus), in three semi-free ranging groups at Affenberg Salem, Germany. In particular, we related

socially relevant attributes (identity, genetic relatedness, rank, sex, age, group membership) to

chemical profiles of body odor. We applied non-invasive active sampling via thermal desorption

tubes and analyzed samples by gas chromatography–mass spectrometry. We found robust evidence

for individual odor signatures and limited support for kin signatures. Chemical profiles were also

related to group membership, but little to rank, sex or age. The detected associations between

chemical profiles and socially relevant attributes form the theoretical foundations for olfactory

information transfer, highlighting the potential of body odor to provide valuable social information in

this highly visually oriented primate.

Keywords: individual signature, catarrhine primate, relatedness, social information, body odor

Introduction

The development and maintenance of social relationships requires animals to acquire and update social information about conspecifics. For instance, animals that form stable relationships need to be able to recognize the unique features of other individuals, and indeed, individual recognition is widespread in the animal kingdom [e.g. 1–3]. Additionally, social animals may interact differently with each other depending on attributes such as sex, rank, group membership or genetic relatedness [4]. Primates are particularly well-known for forming intricate social relationships and frequently show strong and long-lasting relationships among kin [5], but also with unrelated individuals [6,7]. Representations of these social relationships are based on the recognition of traits such as identity, rank and kinship [8]. For instance, rhesus macaques (*Macaca mulatta)* discriminate between individuals, and between familiar as well as unfamiliar kin [1,9,10]. Various macaque species also appear to classify their group members' social relationships with others, suggesting that they attend to interactions in which they themselves are not involved (i.e. third-party relationships [see 11]). Similarly, baboons discriminate between individuals according to their identity and rank, and they also appear to recognize third-party relationships [8,11,12].

The social information underlying differentiated interactions can be emitted and received via different sensory modalities [13,14]. Olfaction is among the evolutionarily oldest of the animal senses [15], and social communication is considered as one of its major functions [16]. Indeed, social interactions are mediated via olfaction in numerous mammals such as African elephants (*Loxodonta africana*) [17], black rhinoceros (*Diceros bicornis*) [18] and various primates. For instance, many strepsirrhine (lemurs, lorises and galagos) and platyrrhine (South American) primates show pronounced scent-marking behavior [e.g. 19–21]. Observational, experimental and chemical studies suggest that primate scent marks as well as odors emanating directly from the body may convey a wealth of information about a sender such as identity, sex, age, rank or reproductive state [e.g. 20,22–24], and that scent marks primarily serve in intra-sexual competition and mediating reproductive behavior [13,22,25].

Notably, vision has gained more and more importance during primate evolution [see 26], and catarrhine

(African and Asian) primates, in particular, are highly visually oriented animals [27]. Scent-marking, on

the other hand, is uncommon in catarrhines [28], and in contrast to evolutionarily older primates,

catarrhines lack a functional accessory olfactory system [29]. Accordingly, catarrhine primates have

long been regarded as little predisposed for olfactory communication [30]. However, the main olfactory

system too has an important part in the processing of cues relevant to social recognition, rendering

the lack of a functional accessory olfactory system of little significance for the communicative

functions of olfaction [see 16,31]. Furthermore, catarrhine olfactory sensitivity may be on par or even

exceed those of other mammals [32,33]. Research on catarrhine olfactory communication has primarily

focused on humans [34]. Similar to other primates, human body odor has been associated with socially

relevant information such as identity, sex, age, kinship or reproductive status (reviewed in [35]),

although the latter is debated [see 36]. Also in mandrills (*Mandrillus sphinx*), one of the few catarrhine

species known to have scent glands, body odor has been related to sex, age, dominance status,

group membership and, possibly, individual identity [37,38]. Results of recent studies further show that

the body odor of other catarrhine species too contains information on socially relevant attributes

such as sex, rank, female reproductive status, genetic relatedness or group membership may be

recognized and acted upon by conspecifics (rhesus macaques, *M. mulatta* [39,40] chimpanzees, *Pan*

troglodytes [41]*,* olive baboons, *Papio anubis* [42]*,* Japanese macaques, *M. fuscata* [43]). Nonetheless, with

the exception of a few species such as humans or mandrills, olfactory communication remains

severely understudied in catarrhine primates [30]. This is particularly true for chemical profiles of body

odor (i.e. the presence/absence and relative abundance of chemical compounds), which restricts our

understanding of which olfactory cues may be available to conspecifics within catarrhines at all. To

reduce this gap, we assessed the social information content of (i.e. potential olfactory cues in) body

odor in a catarrhine primate by relating chemical profiles of body odor to socially relevant attributes

of Barbary macaques living in natural social structures under semi-free conditions at Affenberg

Salem, Germany.

Like other macaque species, Barbary macaques live in multi-male, multi-female groups with female

philopatry and male dispersal [44,45]. Matrilineal kin (especially mother-offspring and maternal half-

sibling dyads) form strong bonds throughout their lives and are therefore highly familiar with each

other [5,46]. Patrilineal kin (e.g. father-offspring or paternal half-sibling dyads), on the other hand,

typically are less closely associated and may even reside in different groups so that they can be

expected to be less familiar with each other [see 47,48]. Females show matrilineal rank acquisition in

which female relatives take similar ranks [49,50]. Adult males form dominance hierarchies in which older

males tend to rank higher, but males beyond their prime (i.e. older than 14-16 years) are subordinate

to prime age males due to being in inferior physical condition [51]. Male Barbary macaques also form

strong, enduring affiliative relationships often referred to as "friendships" [52], whereby triadic

interactions between two males and an infant represent an important form of male affiliative

interaction [53].

Various studies suggest that Barbary macaques can discriminate between individuals, sexes, age

classes and group members using visual or acoustic cues [54–56]. They further distinguish familiar

maternal kin from non-kin using acoustic cues [57]. However, Barbary macaques, which are considered

to be mating particularly promiscuously [58], do not seem to differentiate between unfamiliar kin and

non-kin either in a mating context [47] or in infant care [53,59]. It remains unclear whether this is due to i)

a lack of kin-related cues, ii) Barbary macaques not being able to recognize available cues or iii)

Barbary macaques recognizing kin-related cues but not acting upon them. Kin-related cues in body

odor composition have been shown in a closely related species (rhesus macaques) [40], and Barbary

macaques also use olfaction in social contexts [60], but systematic studies on olfactory kin signatures or

other socially relevant information in Barbary macaque body odor are, to date, absent.

In the present study we aimed to i) identify variation in body odor composition (i.e. presence and

abundance of chemicals) that is systematically associated with socially relevant information about

individuals, and ii) to pinpoint potential semiochemicals (i.e. compounds and compound mixtures

involved in chemical interactions between organisms[15]) associated with the respective traits. Our

particular focus was on traits that form the basis for differentiated social relationships, i.e. individual identity, genetic relatedness between and social rank of individuals. We further assessed effects of sex, age and group membership on chemical profiles of body odors while statistically controlling for potential seasonal and methodological variation (such as sampling distance and volume, or analytical batch) in the data. We hypothesized that the odorants emanating from the skin and fur of Barbary macaques contain socially relevant information. In particular, we expected the similarity between chemical profiles to be greater within than between individuals, and higher among related than among unrelated individuals. Because maternal kin are in frequent physical contact and may exchange odors from the environment or share more of the odor-producing microbiome, we expected maternal kin to exhibit stronger kin signatures than paternal kin, for whom kin signatures should primarily result from similarities in their genetic architecture. We also expected individuals of similar rank to have more similar chemical profiles and/or to show rank-related differences in the relative abundance of particular compounds, as also suggested in [61], because individuals of similar rank may have similar hormonal profiles (e.g. androgens or glucocorticoids), access to food resources and/or space use [see 62–65]. Finally, given the comprehensive evidence for systematic differences in body odor composition with respect to sex, age and group membership in other primates, we also expected similarities between chemical profiles and/or the relative abundance of compounds to be associated with these traits.

Methods

Study animals

The study was conducted at Affenberg Salem, Germany, home to ~ 200 Barbary macaques in three social groups roaming year-round within a 20 ha forested enclosure. The monkeys are well-habituated to humans, as the park is open to visitors from spring to autumn, and to researchers year-round (see [66] for details on the park). The monkeys' diet consists of plants and insects found naturally

within the park as well as fruit, vegetables and grain that is provided daily by park staff. Ponds and

water troughs are distributed throughout the park and grant *ad libitum* access to water. Individuals

differ in facial pigmentation patterns, coat color and stature and have an alphanumeric code

tattooed on the inner thigh, which makes them well distinguishable for experienced observers. Park

staff monitor the groups on a near-daily basis and record major life history events such as births,

deaths or group changes. About two thirds of the adult females receive hormonal implants

(Implanon NXT) to control population size.

We collected odor samples from a total of 72 focal males and females from all three social groups

("C", "F" & "H") and across all ages (2 - 27 years, mean ± SD = 10.5 ± 5.4 years) that were most

suitable to address the main study questions. Focal animals included 13 females without and 18

females with hormonal contraception throughout the study period; two females received

contraception for part of the study period (see Supplement 1 Tab. S2 and statistical analyses section

below). To assess individual odor signatures, we included 24 particularly well-habituated

reproductively mature individuals (> 4 years) that could be sampled frequently (10 samples per ID,

Pollard et al. 2010). To address kinship, we included mothers/fathers and their offspring (> 2 years,

because younger individuals could not be reliably habituated to sampling yet) from a total of 12

maternal and 12 paternal kin units. For rank of males and females, we selected individuals from the

highest- and lowest-ranking adults in their respective groups (17 individuals per rank category, see

details in statistical analysis section).

Odor sampling

One experimenter (B.M.W.) collected 476 odor samples from the focal animals (mean ± SD = 6.6 ±

2.6, median = 5, range 1 – 11 samples per ID) in five sampling periods between October 2020 and

April 2022. Sampling took place in spring (N = 213 samples from 51 individuals), summer (N = 113

samples from 39 individuals) and autumn (N = 150 samples from 56 individuals), but, for logistic

reasons, not during the peak of the mating season in the months of November, December and

January. As Barbary macaques, like other macaque species, do not appear to have specialized scent

glands, we non-invasively collected the airborne volatile and semi-volatile compounds emanating

from the skin or fur of the animals across different parts of the body (see below). We employed

active sampling with thermal desorption (TD) tubes (stainless steel, Supelco), which contained two

adsorbent polymers (Tenax TA and XAD-2, Sigma Aldrich) that trap a wide range of chemical

compounds [67]. The tubes were connected to a silently operating air pump (BiVOC2, Holbach) via a

silicone hose [68]. To be able to trap the air in the immediate vicinity of the animals (1 to 15 cm, mean

± SD = 3.5 ± 2.6 cm), monkeys were habituated to retrieve pieces of peanuts from a pipe collar (7.5

cm diameter) strapped to a horizontal branch or rail while the experimenter stood next to the setup

and pointed a wooden stick with the TD tube mounted at its end towards the monkey. Once a

monkey calmly started to retrieve the peanuts, the experimenter opened the TD tube, positioned it

close to the monkey and started the air flow through the tube with the flow rate set to 1 L/min.

Sampling was resumed until an air volume of 0.5 – 1 L was collected. If the animal moved away

before the sample was completed, the pump was immediately stopped, the tube closed, the volume

already sampled noted and sampling resumed as soon as possible onto the same TD tube (mean ± SD

= 2.3 ± 1.5 subsamples). A sample was typically completed with < 5 min between subsamples, except

for 10 of the 476 samples in which the sample was completed on different days (see Supplement

GCMSdata). Samples were collected from body parts that could be reliably sampled without

disturbing the animals, which primarily constituted the shoulders, flanks or outer thighs (54 % of

samples being collected exclusively from these regions, and 86 % of samples included these regions).

Whenever possible we avoided sampling the hands, feet or facial area, which were the body parts

most likely contaminated with food, feces or other items from the environment. To be able to

statistically control for potential variation arising from the sampled body parts we grouped them into

4 categories: "upper" constituted samples collected from the upper part of the body (i.e. arms,

shoulders and/or the back of the head), "central" the middle part of the body (i.e. flanks, central back

or chest), and "lower" the lower part of the body (i.e. hips, thighs and legs). Finally, we grouped all

samples collected from multiple regions of the body (i.e. upper, central and/or lower) into the

category "mixed" (see Supplement GCMS data).

We avoided sampling when conspecifics were within 3 m of the target animal or higher-ranking

individuals were within sight, when animals had body contact with group members less than 5

minutes before [68], when the monkeys' fur was moist or wet, or when it was very windy. To avoid

disturbances during sampling and contamination with human odors, we did not collect any samples

when visitors or park staff were within 15 m. All equipment was handled wearing disposable gloves.

The experimenter wore a medical mouth-nose cover during sampling and avoided the use of

perfumed products or strongly smelling foods before and on sampling days.

To control for chemical compounds originating from the sampling material and sampling

environment *per se*, we further collected samples of air in the absence of animals (N = 17), TD tubes

mounted for sampling in the absence of animals but without drawing air through the tube (N = 6)

and TD tubes taken to the field but remaining closed at all times (N = 9) as blanks. Before each use,

we cleaned the TD tubes in a thermal conditioner (TD Clean-Cube, Scientific Instruments

Manufacturer) under a constant stream of nitrogen for 120 minutes. When not in use, the tubes

were closed tightly with Swagelok brass caps, wrapped in aluminum foil and stored in airtight bags at

ambient temperature.

Chemical analysis & profiling

Samples were analyzed with gas chromatography – mass spectrometry (GC-MS) on a Shimadzu

TQ8040 GC-MS coupled to a thermal desorption unit (TD-20) with a scan range from 30 to 500 *m/z*

following established GC-MS protocols for the analysis of TD tube samples [67,68]. In the process, the

trapped organic compounds were entirely desorbed from the tube, which then was conditioned and

re-used for sampling. To remove any potential residues from previous measurements from the

column, we ran three blank samples at the start and end of each batch as well as after every tenth

sample.

GC-MS data were processed following an established, semi-automated procedure [67,68], in which a

preliminary compound library was created through automated peak detection and alignment of

peaks by retention time using AMDIS (version 2.71 [69]). We manually corrected this library by taking

into account characteristic mass-to-charge ratios of peaks in addition to retention times. We

subsequently characterized all unique compounds by their retention time and mass spectral patterns

but did not attempt to chemically identify them unless their abundance was later determined to be

statistically associated with particular traits (see below). For the 264 compounds in this library, we

extracted the peak areas for the highest ion not present in adjacent peaks using the Shimadzu GCMS

Browser software. In a subsequent data cleaning step we excluded 81 compounds consistently found

in blanks with peak areas higher than in the animal samples from further analysis, as they are unlikely

to have derived from the target animal [detailed in 68], and performed statistical analyses using the

remaining 182 compounds (see Supplement 2 for full list of retention times and characteristic mass-

to-charge ratios). Furthermore, we assigned undetected peaks, peaks with a signal to noise ratio < 1

and with a peak area below the noise threshold (visually determined to be at 1000) a peak area of 0,

because these peaks were not reliably separable from background noise. All these steps were done

based on uninformative sample numbers and thus blind to the social information about the sampled

individuals. We tentatively identified compounds statistically related to particular traits (see below)

by comparing their mass spectra with the best matches of the NIST Mass Spectral Library (NIST14;

National Institute of Standards and Technologies, Gaithersburg, MD, USA).

Parentage analysis

Pregnancies and births have been monitored consistently by Affenberg staff since the 1970s, so that

partial pedigrees could be built tracing back maternal relatedness over 5-6 generations for most

individuals based on these long-term observational records.

For genetic parentage analysis we used DNA extracted from blood, tissue, hair or feces of 144

individuals (including 68 of the 72 odor-sampled animals, see Supplement 1 for details on genetic

samples and paternity analysis). We genotyped all individuals at an average of 16.7 ± 2.8

microsatellite loci using a marker panel previously established for the study population [70] (see

Supplement 1, Tab. S1).

By applying a combination of exclusion and likelihood methods we were able to assign fathers to 23

of the 28 offspring from cohorts 2017-2019 and to 18 individuals initially sampled as mothers or

potential sires in the present study and born between 2011 and 2016 (see Supplement 1 for details).

For the 68 odor-sampled and genotyped individuals, we were able to genetically confirm 25 fathers

and 41 mothers.

Based on the genetic analysis and maternal pedigree data from long-term records, we calculated

pedigree relatedness (r) per kin line (maternal or paternal/mixed) for each possible dyad in our study

using the software TRACE (version 0.1.0) [71]. Maternal r reflected genetic relatedness between two

individuals of either sex exclusively via female ancestors, while paternal/mixed r (hereafter just called

"paternal") reflected any connections involving at least one male ancestor linking two given

individuals. Because no genetic samples could be obtained for potential sires deceased before the

onset of the study, paternities could only be reliably determined for the younger cohorts, and

therefore our values of pedigree relatedness for the paternal line are likely underestimated in this

analysis.

Dominance and rank

At the onset of the study, we asked experienced park staff to classify individuals from low to high-

ranking, and to identify the alpha male and female in each group based on observed agonistic

interactions and priority of access to resources. This initial judgement was used to pre-select focal

individuals for habituation to the sampling setup. We further conducted dynamic Elo-rating [72] using

ad libitum observations of agonistic interactions (as defined in [73]) throughout all sampling periods.

Dynamic Elo-rating allows for incorporating intensities of interactions as well as prior knowledge and

therefore requires fewer observed interactions and interacting dyads than unmodified Elo-rating [74]

or matrix methods to obtain robust rank estimates [72]. We used the initial classification by park staff,

supplemented by a classification by B.M.W., to define starting values for Elo-rating, and updated this

information with 462 female-female and 240 male-male interactions of varying intensities observed

between October 2020 and April 2022. We computed ranks separately per sex and group, using the

observations during the first two sampling periods as a "burn-in" phase, and extracted the ranks of

the focal individuals at the ends of the second to fifth sampling period for further analysis.

Ethical note

All experimental protocols were approved by Affenberg Salem and complied with the ARRIVE

guidelines. All samples collected specifically for this study (i.e. odor and fecal samples) were collected

non-invasively and the animals participated in the odor sampling procedure voluntarily. In line with

the animal protection law in Germany, we therefore required no further animal ethics approval. All

methods were performed in accordance with the relevant guidelines and regulations.

Statistical analysis

General procedures

We conducted all analyses in R (version 4.3.3) [75].

To investigate chemical similarity between samples, we performed permutational multivariate

analyses of variance (perMANOVAs) using the function "adonis2" from the package "vegan" (version

2.6-4)[76]. These analyses relate pairwise chemical (dis)similarities between all samples entered into a

given analysis to (dis)similarities between traits using distance matrices automatically assembled

from the sample-specific information provided. As measure of distance (or dissimilarity) we used

Bray-Curtis indices calculated from standardized, $\log(x+1)$-transformed peak areas [68,77]. Depending on

the investigated trait(s) we used different data subsets and predictor variables as described for the

different traits below. Unless noted otherwise, we set ID as stratum to account for repeated

measures per individual. For perMANOVAs with multiple test predictors we first conducted a full-null

model comparison by setting the argument "by" to "NULL", and only if this was significant ($p < 0.05$)

we investigated the effects of the single predictor variables by setting the argument "by" to

"margin". In case of non-significant interactions, we re-ran the marginal model without the

interaction to be able to assess the effect of the respective main terms. We ran all perMANOVAs with

the default of 999 iterations.

When the analysis of chemical (dis)similarities involved predictor variables describing pairs of

samples (e.g. pedigree relatedness) and/or required the selection of specific pairs of samples (which

is not possible with the perMANOVA), we further conducted multi-membership models. Multi-

membership models extend multilevel models by allowing to fit data with multiple membership

structures that are not purely hierarchical [78], such as terms whose levels are present in more than

one variable (e.g. ID in either sample 1 or sample 2 of a given pair-wise comparison). For these

models we computed Bray-Curtis dissimilarities from standardized, log(x+1)-transformed peak areas

using the package "vegan" and, for easier interpretation, transformed them into similarities by

subtracting the dissimilarity scores from 1. We fitted these models with chemical similarity as

response variable and various test and control predictors as described below. As similarity scores

follow a beta distribution, we computed the models using a Bayesian approach in the package

"brms" (version 2.21.0) [79], which is the only package we are aware of that implements multi-

membership models with beta distribution. We used the default settings for priors, i.e. uninformative

priors.

We further conducted compositional mixed model analyses, suitable for data whose components are

constrained to sum to a constant value, for a detailed analysis of chemical profiles that considers the

peak areas of all compounds per sample rather than overall (dis)similarities between samples, and

may therefore aid in detecting (suites of) compounds associated with particular traits. For this

purpose, we vectorized the data matrix and included the matrix rows and columns (i.e. the samples

and compounds) as random intercepts to avoid pseudo-replication and heteroscedastic variance [68,80].

As response variable we used standardized peak areas. Because standardized areas are bound

between 0 and 1 and sum up to 1 per sample, we applied a centered log-ratio (clr) transformation [81]

and modelled the transformed values with a Gaussian error distribution. As effects of interest in this

analysis we modelled the interactions between the predictors and compound, i.e. the random slopes

& random interactions of fixed effects test predictors within compound [see 68]. To determine which

compounds were associated with particular traits, we subsequently extracted the compound-specific

random slopes and random interaction estimates for each trait. We considered a given compound to

be robustly associated with a trait if its credible intervals did not overlap zero (in case of continuous

predictors) or did not overlap between trait levels (in case of categorical predictors). To follow a

consistent modelling approach, we also performed this model in the package brms.

In all mixed models we z-transformed continuous predictor variables to a mean of 0 and an SD of 1

prior to fitting the model to facilitate convergence and interpretation of model estimates [82]. We

assessed collinearity between predictor variables using the function "vif" from the R package "car"

version 3.1-2 [83], whereby we considered values < 4 as unproblematic. To assess model convergence we

inspected rhat values and effective sample sizes (ESS), with rhat values deviating from 1 by less than

0.01and ESS values > 400 indicating satisfactory convergence [84].

We present model results most relevant for the interpretation of the test predictors in the main text,

and full model results, including rhat and ESS values, in Supplement 1 (Tables S3-S6).

We could not account for contraception statistically in any of the analyses given that contraceptive

state applied only to a minority of samples and individuals.

Individual identity

To investigate individual signatures, we analyzed distance matrices using a perMANOVA with ID as

test predictor while controlling for sex, age (in years), group and season of the year (spring, summer

or autumn). Because ID was the test predictor, we did not set ID as stratum. For this analysis we used

data from 24 individuals (12 males, 12 females) sampled 10 times each. As individual signatures

commonly result from the unique combination and abundance of compounds (i.e. the overall

similarity between chemical profiles) rather than certain compounds that vary consistently between

all individuals [85,but see 43], we did not conduct a more detailed analysis of chemical profiles

(compositional models) for individual identity and consequently also did not attempt to identify

specific compounds associated with individual odor signatures.

Genetic relatedness

Similar to ID, kin signatures should result from similarities in the overall composition of compounds

rather than specific compounds. We therefore assessed whether the chemical (dis)similarities

between samples are explained by genetic relatedness. In the first of two sets of analyses, we

computed perMANOVAs using only samples of genetically confirmed units of close kin. Each kin unit

comprised 2 – 4 individuals in up to 3 generations and as such, parent-offspring, grandparent-

grandchild and/or half-sibling dyads, reflecting close kin with relatedness coefficients ranging from

0.25 to 0.5. We ran the perMANOVAs separately per kin line, using the maternal or paternal kin unit

(i.e. belonging to the same or a different kin unit) as test predictor, while controlling for sex, age and

season of the year (spring, summer or autumn). All individuals used in these analyses were sampled

for odor at least 5 times. For maternal kin we used 220 samples from 31 individuals and for paternal

kin 179 samples from 29 individuals in 12 kin units per kin line (see Supplement 1, Tab. S2).

Maternal/paternal relatedness between individuals from different units ranged from 0 to 0.125,

reflecting distant or non-kin.

While belonging to a certain kin unit was genetically confirmed in this first set of analyses, it treated

relatedness categorically as belonging to a given kin unit or not, but lacked the possibility to

differentiate the pair-wise degree of relatedness. In a second analytical approach we therefore

conducted a multi-membership model to relate pair-wise chemical similarities between samples to

kinship while accounting for a variety of fixed and random predictor variables. For the multi-

membership model we used the data from all genotyped and odor-sampled individuals (N = 68),

irrespective of whether their maternities or paternities were genetically solved. We fitted dyadic

Bray-Curtis similarities as beta-distributed response variable and relatedness between members of a

dyad as two separate test predictors: one reflecting maternal kinship (i.e. individuals linked

exclusively via the maternal line) and one reflecting paternal & mixed kinship (i.e. containing at least

one male linking two individuals, which breaks the familiarity between maternal kin typical for

matrilineally structured primate groups). To control for other sources of biological variation we also

fitted the age and rank difference between the two individuals in a given dyad, the number of days

between the two samples and the difference in the time of day the sample was collected. We further

included whether or not the two individuals were of the same sex, belonged to the same group, were

sampled in the same season of the year and from the same body region. To control for variation in

sampling and measurement we included the difference in sampling distance, sampling volume, the

number of subsamples needed to complete a sample and whether or not samples were analyzed in

the same GC-MS batch. To account for repeated measures we fitted the sample, the individual and

the dyad (i.e. the unique combination of individuals) and random intercepts, whereby sample and

individual were fitted as multi-membership terms because a given sample or individual could occur in

either of the two columns describing the elements of a dyad. For a more accurate estimation of fixed

effects, we further included all technically possible random slopes of fixed effects within the random

intercepts of individual and dyad. Using 9000 post-warmup draws, the model converged with all rhat

values deviating from 1 by < 0.01 and all ESS > 400 (see Table 1).

Similar to individual identity we did not conduct compositional models for genetic relatedness and

did not attempt to identify specific compounds associated with kin odor signatures.

Dominance and rank

To address potential dominance and rank-related signatures, we first conducted a perMANOVA with

rank as a categorical test predictor (high/low) in interaction with sex while controlling for age, group

and season of the year. For this analysis we used 260 samples from 34 individuals sampled at least 5

times each and classified as particularly high- or low-ranking in their respective groups based on

judgements of staff and confirmed with behavioral data (high-ranking: 9 females & 8 males, low-

ranking: 8 females & 9 males). To keep confounds between dominance class and kinship arising from

matrilineal rank acquisition to a minimum, we chose no more than two females from the same

matriline, so that the 17 females used for this analysis represented 12 different matrilines.

We further conducted a multi-membership model using chemical data and rank as a continuous

measure (i.e. Elo scores) using 476 samples from all 72 sampled individuals. We only considered

same-sex dyads for this analysis, because Barbary macaque males outrank females and Elo scores

were thus computed separately per sex. We fitted dyadic Bray-Curtis similarities as beta-distributed

response variable and the difference in Elo scores and its interaction with sex as test predictors.

Similar to the model on genetic relatedness we also controlled for potential sources of biological and

technical variance by fitting dyad-specific overall relatedness and differences in age, sampling day,

time of day, group, season of the year, body region, sampling distance, sampling volume, number of

subsamples and GC-MS batch. As above we fitted sample and individual as multi-membership terms

as well as dyad as random intercepts, and all technically possible random slopes within individual and

dyad. We subsequently removed the interaction between the difference in Elo scores and sex

because it explained no variance (estimate 0.0004 +- 0.0028, lower/upper CI -0.005 – 0.006) and

fitted both as main terms instead. Using 12000 post-warmup draws, the model without interaction

converged with all rhat values deviating from 1 by < 0.01 and all ESS > 400 (see Table 2).

To consider the possibility that rank information could be encoded by a few specific compounds

(rather than or additional to overall similarities), we further included rank as predictor in a

compositional mixed model as described in the analyses of sex, age and group below.

Sex, age and group

To address potential signatures of sex, age or group membership we conducted a perMANOVA with

these terms as test predictors while controlling for season of the year. For this analysis we used all

476 samples from 72 individuals.

To assess to what extent specific (suites of) compounds may encode information about individual

traits, we further computed a compositional mixed model. As fixed effects predictors we fitted sex,

age, group and rank (Elo scores) along with season of the year, the sampled body part, time of day,

sampling distance, sampling volume and the number of subsamples. As random terms we fitted the

intercepts of sample, compound, individual, date, GC-MS batch and TD tube used for sampling, as

well as all technically possible random slopes, but no random correlations to avoid an overly complex

model. Using 8000 post-warmup draws, the model converged with all rhat values deviating from 1 by

< 0.01 and all ESS > 400 (see Table 3).

Results

Individual identity

Odor samples from the same individuals were significantly more similar than samples from different

individuals, with ID explaining ~ 20 % of the variance in Bray-Curtis indices (perMANOVA, r^2 = 0.203,

df = 21, F = 3.838, p = 0.001, example chromatograms in Supplement 1, Fig. S1). An individual

chemical signature was further supported by the posterior mean estimates for ID in the multi-

membership models on genetic relatedness and rank (see Tables 1 and 2).

Genetic relatedness

Odor samples from close maternal kin were significantly more similar than samples from different kin

units (perMANOVA, r^2 = 0.099, df = 11, F = 2.537 , p = 0.001). Furthermore, individuals from the

same paternal kin unit tended to have more similar chemical profiles than individuals from different

kin units (perMANOVA , r^2 = 0.132, df = 11, F = 2.772 , p = 0.067).

When relatedness was treated as a continuous variable (i.e. using pedigree r) across all genotyped

individuals in the data set, neither maternal nor paternal relatedness robustly affected similarities in

chemical profiles (Table 1).

Table 1: Estimates for fixed effects and random intercepts from the multi-membership model on
genetic relatedness. Terms of primary interest are indicated in italics. CrI = Credible Interval. ESS =
effective sample size.

fixed terms	Estimate	Est.Error	2.5 % CrI	97.5 % CrI	Rhat	bulk ESS	tail ESS
Intercept	0.1419	0.0264	0.0891	0.1935	1.0017	1592	2884
maternal r	*0.0001*	*0.0010*	*-0.0018*	*0.0020*	*1.0000*	*9718*	*7456*
paternal/mixed r	*-0.0016*	*0.0015*	*-0.0048*	*0.0012*	*1.0003*	*6365*	*5522*
d_elo score	-0.0002	0.0010	-0.0022	0.0018	1.0001	10551	7505
d_sex (same)	-0.0018	0.0018	-0.0053	0.0017	1.0001	9026	7614
d_age	-0.0010	0.0016	-0.0042	0.0022	1.0003	6827	6459
d_group (same)	0.0120	0.0055	0.0009	0.0229	1.0014	2163	3956
d_season (same)	-0.0244	0.0119	-0.0480	-0.0010	1.0032	562	1399
d_date	-0.3204	0.0095	-0.3391	-0.3013	1.0075	528	944
d_time	0.0005	0.0013	-0.0019	0.0030	0.9999	4987	6121
d_bodyregion (same)	0.0026	0.0028	-0.0028	0.0082	1.0005	4418	5675
d_s.distance	0.0043	0.0023	-0.0002	0.0088	1.0006	3783	5261
d_s.volume	0.0065	0.0051	-0.0035	0.0164	1.0032	984	1902
d_subsamples	-0.0072	0.0020	-0.0112	-0.0034	1.0007	4902	6101
d_batch (same)	0.1994	0.0154	0.1690	0.2302	1.0011	1051	1941
random terms							
sample (Intercept)	0.3994	0.0144	0.3724	0.4293	1.0031	1074	2166
ID (Intercept)	0.1490	0.0277	0.0960	0.2037	1.0033	471	973
dyad (Intercept)	0.0189	0.0013	0.0164	0.0214	1.0010	1690	3502

Dominance and rank

The interaction between dominance classes (high vs. low) and sex (male vs. female) had no

significant effect on the similarity between chemical profiles (perMANOVA, r^2 = 0.003, df= 1, F =

0.932 , p = 0.982), but individuals from the same dominance class had slightly more similar chemical

profiles than individuals from different dominance classes (perMANOVA, r^2 = 0.002, df = 1, F = 0.879,

p = 0.008). When treated as a continuous variable across all individuals in the data set, rank (Elo

score) had no robust effect on similarities between chemical profiles (Table 2).

Table 2: Estimates for fixed effects and random intercepts from the multi-membership model on
rank. Terms of primary interest are indicated in italics. CrI = Credible Interval. ESS = effective sample
size.

fixed term	Estimate	Est.Error	2.5 % CrI	97.5% CrI	Rhat	bulk ESS	tail ESS
Intercept	0.1544	0.0394	0.0756	0.2315	1.0010	2680	5074
d_Elo score	*-0.0001*	*0.0014*	*-0.0027*	*0.0026*	*1.0002*	*19102*	*9841*
total r	-0.0021	0.0014	-0.0049	0.0007	1.0002	18584	8977
sex (male)	-0.0228	0.0526	-0.1265	0.0798	1.0014	2879	5192
d_age	0.0013	0.0021	-0.0030	0.0054	1.0001	14446	9510
d_group (same)	0.0159	0.0070	0.0022	0.0297	1.0001	5497	7723
d_season (same)	-0.0267	0.0108	-0.0484	-0.0059	1.0015	2736	4694
d_date	-0.3250	0.0104	-0.3456	-0.3053	1.0023	1088	2508
d_time	0.0005	0.0012	-0.0019	0.0030	1.0002	15821	9039
d_bodyregion (same)	0.0041	0.0033	-0.0024	0.0107	1.0002	11941	8932
d_s.distance	0.0054	0.0025	0.0006	0.0103	1.0007	11074	9460
d_s.volume	0.0081	0.0054	-0.0026	0.0187	1.0014	2684	3564
d_subsamples	-0.0070	0.0021	-0.0113	-0.0029	0.9999	15094	10231
d_batch (same)	0.1838	0.0182	0.1477	0.2188	1.0011	2652	5513
random term							
sample (Intercept)	0.3952	0.0141	0.3684	0.4239	1.0016	2096	4665
ID (Intercept)	0.1509	0.0282	0.0988	0.2076	1.0058	891	1598
dyad (Intercept)	0.0189	0.0019	0.0152	0.0226	1.0016	2254	4976

Also the specific composition of chemical profiles showed no pronounced variation in relation to rank

(see random slope of Elo score within compound in Table 4), and none of the compounds was

robustly associated with rank (see random slopes estimates in Supplement 3).

Sex, age and group

When testing sex, age and group membership, the set of predictor variables significantly affected

chemical similarities (perMANOVA, $r^2 = 0.213$, df = 6, F = 21.145, p = 0.001). In particular, sex, age as

well as group explained significant, albeit small proportions of the variance in Bray-Curtis indices

(3.5 – 5 %, Table 3). In comparison, season of the year (included as a control predictor) explained

more variance than sex, age and group together (14.5 %, Table 3).

Table 3: Results of the perMANOVA investigating effects of sex, age and group on chemical

similarities.

Term	Df	SumOfSqs	R^2	F	P
sex	1	0.466	0.004	2.332	0.001
age	1	0.425	0.004	2.124	0.001
group	2	6.106	0.051	15.273	0.001
season	2	17.225	0.145	43.083	0.001
Residual	469	93.756	0.787		
Total	475	119.117	1		

This pattern was similar when considering chemical profiles in detail. Slope estimates from the

compositional model suggest that the effects of sex, age and group on relative peak areas differed

between chemical compounds, whereby the variance was close to 0 for age, only slightly higher for

sex and more pronounced for group. Again, variance related to control predictors such as season of

the year was considerably higher than for sex, age and group (see Table 4).

Table 4: Estimates for random slope terms within the random intercept of compound for the

compositional model. Terms of primary interest are indicated in italics. CrI = Credible Interval. ESS =

effective sample size.

slope in compound	Estimate	Est.Error	2.5 % CrI	97.5% CrI	Rhat	bulk ESS	tail ESS
Elo score	*0.0166*	*0.0118*	*0.0007*	*0.0435*	*1.0020*	*3304*	*3855*
sex	*0.1023*	*0.0191*	*0.0626*	*0.1380*	*1.0006*	*2819*	*3082*
age	*0.0524*	*0.0175*	*0.0121*	*0.0831*	*1.0017*	*1709*	*1218*
group	*0.2901*	*0.0157*	*0.2606*	*0.3214*	*1.0012*	*3772*	*4889*
season	1.0605	0.0417	0.9822	1.1461	1.0019	2698	4371
time	0.0397	0.0182	0.0041	0.0724	1.0032	1657	2624
bodyregion	0.1333	0.0176	0.0987	0.1672	1.0009	2690	3865
s.distance	0.2624	0.0177	0.2290	0.2983	1.0003	2989	5028
s.volume	0.8104	0.0420	0.7341	0.8990	1.0015	1140	2388
subsamples	0.1280	0.0147	0.0993	0.1569	0.9999	3441	5449

When investigating the compound-specific estimates per test predictor, the credible intervals for all

compound-specific age estimates overlapped 0, although one amide, one aromatic hydrocarbone

and an unidentified compound overlapped by < 20 %. Similarly, the credible intervals for all

compound-specific sex estimates overlapped between the sexes, although one aldehyde, one

monoterpene and 4 unidentified compounds showed only a minor overlap (i.e. central estimate for

one sex outside the credible interval of the other sex, Supplement 3). Putative chemical

identifications for these compounds are reported in Table S7 of Supplement 1.

The group estimates for 8 compounds did not overlap between at least 2 of the 3 groups, suggesting

that the peak areas of these compounds differ distinctly between groups (Fig. 1, Supplement 3). We

tentatively identified 5 of these compounds as a ketone, a carboxylic acid, an alcohol, an aromatic

hydrocarbon and a catechol derivate (Table 5).

Table 5: Compounds associated with group differences in chemical profiles of Barbary macaque
samples. RT = retention time (in min.), base peak describes the largest mass fragment of the peak. SI
= similarity index (from 0 - 100) for the match between mass spectra and their respective best hit in
the NIST library.

RT	base peak	best NIST library hit	SI	substance class	found in
2.64	43	2-Pentanone, 3-methyl-	91	ketone	humans [106,113], Japanese macaques [43]
2.68	43	Acetic acid	94	carboxylic acid	various mammals [38,103,106,114–117]
5.16	149	unidentified	< 80		
9.08	73	3-Pentanol, 2,4-dimethyl-	81	alcohol	*human excretions [106,113]
10.38	79	unidentified	< 80		
13.23	115	1H-Indene, 3-methyl-	92	aromatic hydrocarbon	unknown
19.50	179	4-tert-Butylcatechol, dimethyl ether	87	catechol derivate	** beaver castoreum [116]
28.15	91	unidentified	< 80		

* these studies identified a different derivate (3-Pentanol, 3-methyl)

** this study identified two different catechol derivates

Figure 1: Group differences (C, F and H) in peak areas (centered log-ratio transformed) of 8

compounds with retention times in [min] as indicated in the upper left corner of each graph.

Compounds are sorted by order of importance as derived from their compound-specific slope

estimates. Vertical lines show mean peak areas per group, points the estimates per group (with all

other predictors at their average) and whiskers the 95 % Credible Intervals.

<u>Discussion</u>

In the present study, we investigated to what extent the chemical composition of Barbary macaque

body odor is related to socially relevant traits and might thus have the potential to convey social

information. Our analyses support the existence of individual odor signatures, while support for kin

odor signatures differed between our analytical approaches. Chemical profiles were also related to

group membership, but we found little to no support that rank, sex or age affect chemical similarity

or detailed composition of odors emanating from the skin and fur of Barbary macaques.

Individual identity showed the strongest and most robust signature related to social information in

Barbary macaque body odor. Our results are in line with the extensive evidence for individually

distinct body odor across a range of mammals [e.g. 86–88], including primates [e.g. 89,90,43,91,92,reviewed in 85].

While the majority of evidence thus far has come from scent-marking species, our results, along with

studies in Japanese macaques [43] and on human axillary odor [93], indicate that individually distinct

odors may also be present in catarrhine primates that do not show apparent scent-marking.

Individual identity explained only slightly less variance (r^2 of 0.2) in the chemical profiles of Barbary

macaques than in Japanese macaque urine (r^2 of 0.25) or vaginal secretions (r^2 of 0.29). Although the

comparison between Japanese and Barbary macaques certainly needs to be viewed in light of

different methods and a small sample size in the Japanese macaque study[43], it suggests that odor

emitted from the body surface may carry identity information of similar magnitude as potentially

more intense odor sources. Furthermore, the present study shows that individually distinct

information can be conveyed by the chemicals emitted from the body surface, not just by odors

deposited in the environment in the form of scent marks or excretions.

The comparison of close vs. distant or non-kin based on overall chemical similarities suggested

olfactory signatures of both maternal and paternal relatedness, although paternal signatures were

statistically less robust. Units of close kin also showed a more similar chemical composition in owl

monkeys (*Aotus nancymaae*) [94,but see 91] and rhesus macaques [40]. Similar to our results in Barbary

macaques, kin signatures in rhesus macaques were only statistically robust among maternal, but not

paternal half-siblings [40]. These patterns suggest that similarities in body odor between maternal

relatives may to some extent be mediated by frequent physical contact and/or a shared

environment, which could result in an exchange of odor-producing microbes, as e.g. suggested for

group odor signatures in hyenas [95]. However, when treating relatedness as a continuous variable as

well as accounting for more other potential sources of variation in body odor, we found no support

whatsoever for relatedness predicting chemical similarity between individual odors. Possibly,

chemical profiles of surface body odor allow only a coarse, but no fine-scaled differentiation of

relatedness. Alternatively, what appeared to be kin signatures in the perMANOVAs may actually have

been variation related to other predictors that could not be as accurately controlled for in these

analyses. Furthermore, kin odor signatures may only be robustly expressed in certain contexts or

seasons, when kin discrimination is particularly important. For instance, kin signatures in glandular

secretions of ring-tailed lemurs (*Lemur catta*) were robustly expressed only in the breeding season,

when they may aid in inbreeding avoidance [96,97], but we did not collect odor samples during the

mating season for logistic reasons.

The lack of robust chemical information about kinship involving the paternal line (at least outside the

mating season) is especially interesting with respect to the triadic interactions and care that Barbary

macaque males frequently engage in with infants. Particularly, male-infant interactions have been

suggested to be independent of paternity to the respective infant [53], which may be due to low

phenotypic similarity between father and offspring when the offspring is still in its infancy. Studies in

rhesus macaques testing exclusively adult subjects suggest that unfamiliar, paternal kin discriminate

relatives based on visual or acoustic phenotypic cues [9,10]. Moreover, humans rating the facial

similarity of rhesus macaque parents and offspring performed above chance level [98] and better if the

offspring were older [99]. The paternal kin units sampled in the present study indeed comprised cases

in which the offspring was still rather young (2 years old), but in half of the cases all members of a

paternal kin unit were reproductively mature (i.e. 4 years or older). Hence, the chemical data

presented here do not point towards olfactory signatures alone being robust enough to allow

Barbary macaque males to reliably distinguish between own and unrelated offspring.

Regardless of the analytical approach, we detected no robust support for chemical profiles being

related to dominance classes or rank. Although we found a significant effect of dominance class on

the similarity between chemical profiles, the association was very weak, and ranks across the entire

dominance hierarchy were associated with neither chemical similarity nor the abundance of specific

compounds. Dominance-related odors have been described in various mammals [e.g. 61,reviewed in 92],

including male mandrills [37,38], but were not detected in ring-tailed lemurs [92] or female mandrills [37]. In

female rhesus macaques, individual rank had no effect on overall chemical similarities, but affected

the relative abundance of certain compounds [40]. As pointed out by Smith [85], olfactory communication

of dominance requires either a specific dominance-related odor or olfactory individual signatures

that can be associated with other dominance-related information. Evidence for the latter comes from

a study in ring-tailed lemurs, in which males discriminated between the odors of dominant and

subordinate individuals, but only if they were familiar with the odor donors [21]. Hence, while our

results provide no robust evidence for dominance-related body odor in either male or female

Barbary macaques, the detected individual odor signatures certainly open the possibility for a more

indirect olfactory communication of dominance among familiar individuals. Furthermore, our results

imply that Barbary macaques of similar rank either shared spaces less than we had assumed, or that

sharing similar spaces had no detectably effect on chemical profiles. The latter seems contradictory

to our suggestion that a shared environment may have contributed to the more robust odor

signatures among close maternal than paternal kin. However, a higher degree of spatial proximity

and physical contact among close maternal kin than among unrelated individuals of similar rank

could lead to kin, but not rank signatures to be affected by the environment.

Based on behavioral and/or chemical evidence, the odors of numerous primates and other mammals

convey sex-specific [18,40,91,100,101] and age-specific information [24,86,94,100,102,103], although this need not

always be the case [102,also see 30]. Age, for instance, was related to chemical profiles in one subset of

individuals in rhesus macaques, but not in another [40]. In the present study, sex and age, although

significantly associated with chemical similarities, explained very little variation in chemical profiles,

and none of the chemical compounds distinctly differed in abundance between sexes or showed

consistent changes with age. It is possible that sex and age information may actually be more

pronounced in a different context (e.g. mating), in other odor sources (e.g. urine) or other body parts

(e.g. the ano-genital region), all of which, along with differences in sampling and analytical methods,

may affect the expression of chemical signatures or our ability to detect them [30,43,see 97,104].

Furthermore, sex differences may have been attenuated in the study population due to the use of

contraceptives in 60 % of the adult females (see below).

Group membership explained about 5 % of the variance in chemical similarities and affected the

relative abundance of 8 compounds. For most of these compounds, abundances were consistently

higher or lower in group F, but similar in the other two groups. Group odor signatures have been

described in a range of group-living mammals, including rhesus macaques [39,40], mandrills [38] and

chimpanzees [41], with explanations for the origins of these differences including the environment, diet

and microbial differences between groups [91,95,101]. In species living in family groups, group odor

signatures may further reflect relatedness among individuals (e.g. in owl monkeys [94]). Relatedness is

an unlikely explanation for the group differences observed in the present study, because all groups

comprised multiple matrilines and average relatedness among genotyped focal animals was low both

within (r = 0.04) and between groups (r = 0.02, data in Supplement dyadic relatedness data). The

three groups also received the same supplemental food and had spatially (but not temporally)

overlapping home ranges, but there are subtle differences in the vegetation in different areas of the

park that could potentially affect natural food sources as well as background volatiles or substances

the animals get into contact with. Several of the tentatively identified compounds could fit such an

explanation. 3-methyl-2-pentanone and 2,4-dimethyl-3-pentanol, for instance, could be microbial or

plant products metabolized by the mammalian body [see 105] and similar compounds have been found

in human feces before [106]. Furthermore, indenes have been described in plants [e.g. 107], although we

could not find a source specifically for 3-methyl-1H-indene, which was the best mass spectral library

match for the compound at RT 13.23 min. Alternatively, group odor differences could be related to

demographical and social differences between groups. The focal animals of group F comprised a

slightly higher proportion of females than groups C and H (57 % vs. 43 % and 32 %) and were

marginally younger (average age in group F 9.2 years vs. 11.7 and 11.3 yrs in groups C and H, see

Supplement 1 Tab. S2). However, because sex and age effects on body odor were very small (see

above) and statistically accounted for, they are unlikely to contribute much to the observed group

differences. Group F also is the largest and highest-ranking of the three groups, which could affect

stress levels or other physiological parameters [see 108] and thereby, body odor (reviewed in [109]), but we

have no physiological data available to further assess this possibility.

Main assets of our study certainly are the large numbers of samples and focal individuals, the semi-

natural environment and natural social conditions. We also were able to collect all samples non-

invasively, thereby reducing the chance that odors were affected by sampling-related stress. At the

same time, the study faced several limitations, some of which we were able to account for during

data analysis, and others not. For instance, we collected samples from different body parts that we

could approach most easily, i.e. primarily the shoulders, flanks and thighs, and controlled for the

variation in sampled body regions during statistical analysis. However, it is difficult to assess how

relevant these body parts actually are for day-to-day information exchange. Recent observational

data in the study population suggest that sniffs directed at conspecifics frequently target the ano-

genital region of adult females in a mating context [60]. The ano-genital region might thus be a more

appropriate sampling region at least for certain types of individuals and contexts, but would have

been feasible to sample only in a subset of very well-habituated females at frequent intervals. As the

social information detectable in animal odors may differ between odor sources, even within species

[see 30,43], it is possible that results would have differed if we had sampled other body regions or odor

sources such as urine.

Our data collection periods spanned much of the year, but for logistic reasons we did not collect data

during most of the park's winter break, which corresponds to the peak of the mating season in the

study population. Indeed, the season of the year did affect chemical profiles, even more so than the

socially relevant variables investigated. In particular, seasonal variation may have affected the

volatile background in the environment, which we could at least in part account for by using

environmental control samples. Furthermore, aspects such as reproductive physiology, diet, ambient

temperature or state of the fur (winter vs. summer coat) may additionally affect the chemical

compounds emitted by the body and how readily they become available to recipients (or sampling

tube). Such aspects are likely to be at the base of the detected seasonal changes in chemical profiles,

and we statistically accounted for them in our analyses of socially related olfactory signatures by

using season of the year as a control variable. What remains open is whether any of the investigated

effects might have been more pronounced during the mating season, as shown, e.g., for kin

signatures in ring-tailed lemurs [96,97].

Another aspect to consider is that the size of the study population is managed via hormonal

contraception, and 60 % of the sampled females received contraception during part or all of the

study period. Studies on the effects of hormonal contraception on primate body odor are scarce and

contradicting: hormonal contraception using medroxyprogesterone acetate has previously been

shown to alter the chemical composition of female odors in ring-tailed lemurs [110] but not in owl

monkeys [91]. Male mandrills showed more flehmen responses towards odor samples from cycling

females than from females contracepted with etonogestrel [111], which is the active ingredient of the

implants used in the studied Barbary macaques. However, while contraception also tended to

modulate age-specific differences in ano-genital sniffs by male Barbary macaques, it did not affect

subsequent sexual behavior in our studied population [60]. As we did not assess individual attributes

immediately related to mating and female fertility, contraception might be considered as a source of

noise in the data. Unlike several other sources of noise, however, we could not account for

contraception statistically given that it applied only to a minority of samples and individuals. Overall,

it is thus possible that certain signatures, in particular those of sex, might have been more

pronounced in a sample without females receiving contraception.

Our results also suggest some variation in chemical profiles related to sampling and GC-MS

measurement, whereby different variables affected different aspects of the composition: chemical

similarities between samples were greater if samples were collected closer in time, and to a lesser

extent also when analyzed within the same GC-MS batch. In contrast, variables such as the sampling

distance, sampling volume or the sampled body part had little effect on chemical similarities of

samples, but appeared to have some effect on the relative abundance of certain compounds.

However, while we were unable to eliminate these sources of variation in the study design, we

statistically accounted for them in all mixed model analyses.

Finally, results need to be put into perspective of what we can measure vs. what animals can

perceive. Statistically detectable associations between chemical profiles and socially relevant traits

show the potential for olfactory information transfer, but we do not know if or how animals use this

information. Animal sensory systems have evolved for the detection of relevant information and may

thus be superior to even the most modern of our analytical techniques, at least for certain

compounds (reviewed in [112]). Accordingly, it may be possible that animals are able to pick up upon

even very subtle differences in odors, or differences that our analyses are unable to pick up at all,

especially as we are unlikely to sample and detect all chemical substances emitted by an animal's

body across the possible ranges of volatility, polarity etc. [104] despite the constant improvements in

sampling and analytical methods. Hence, future studies should combine chemical data with

(experimental) assessments of behavioral and physiological reactions towards conspecific odors

across a range of contexts to obtain a detailed understanding of the complex role of olfaction in

primate social interactions. Particularly under natural and semi-natural conditions, controlled

presentations of conspecific odors are challenging to implement [e.g. 55], as stimuli need to be

presented in a manner that is salient and realistic enough to elicit interest and closer inspection

without leading to quick habituation. Promising next steps would be multi-modal experiments and

experimental manipulation of *in vivo* odors to investigate how olfactory discrimination of social

information falls into place between the other senses, i.e. whether macaques are able to

discriminate such information based on odors alone, and/or if body odor signatures improve the

reliability of other sensory cues.

In conclusion, our results add to the limited evidence that socially relevant information may not only

be encoded in odor secretions and excretions, but also in the volatile and semi-volatile components

emanating from the body surface of primates. Based on chemical profiles alone, i.e. without knowing

what the monkeys can make of it, data suggest that Barbary macaque body odor could directly

contribute to recognizing individuals and groups, while body odor signatures may not be pronounced

or robust enough to allow differentiating kinship, rank, sex or age of conspecifics from surface body

odor alone. However, given the central role of individual recognition in forming and maintaining

social relationships, surface body odor could potentially still contribute to managing social

relationships, if monkeys associate social knowledge obtained via other sensory modalities or direct

social interactions with individual odor signatures.

Acknowledgements

We thank Roland & Mamisolo Hilgartner and the staff of the Affenberg Salem for continuous support

and Ellen Merz for permission to conduct this study. We further thank Roger Mundry for an

introduction to multi-membership models and help with the centred log-ratio transformation.

Hendrikje Westphal established the methods for genetic paternity analyses and helped with faecal

sample collection. Stefanie Bley did the allele calling and managed the paternity analyses. Antonia

Krüger and Anja Schiffner conducted DNA extractions and PCR analyses, and helped with GC-MS

measurements and TD tube cleaning. Annika Freudiger helped with the implementation of the TRACE

software. Ramona Oehme and Susan Billig in the mass spectrometry core facility at the Faculty of

Chemistry and Mineralogy at Leipzig University provided technical support in GC-MS maintenance.

This work was supported by the German Research Foundation, DFG (grant no. SCHL 2011/2-1

awarded to B. M. Weiß). The GC-MS instrument was funded by the European Fund for Regional

Structure Development, EFRE ("Europe funds Saxony", grant #100195810 to A. Widdig).

References

1.   Rendall, D., Rodman, P. S. & Emond, R. E. Vocal recognition of individuals and kin in free-
ranging rhesus monkeys. *Anim. Behav.* **51**, 1007–1015 (1996).
2.   Scheiber, I. B. R., Hohnstein, A., Kotrschal, K. & Weiß, B. M. Juvenile Greylag Geese (Anser
anser) Discriminate between Individual Siblings. *PLOS ONE* **6**, e22853 (2011).
3.   Tibbetts, E. A. & Dale, J. Individual recognition: it is good to be different. *Trends Ecol. Evol.* **22**,
529–537 (2007).
4.   Ward, A. & Webster, M. *Sociality: The Behaviour of Group-Living Animals*. (Springer, 2016).

5.   Silk, J. B. Ties that bond: the role of kinship in primate societies. in *New directions in*
anthropological kinship (ed. Stone, L.) 71–92 (Rowman & Littlefield Publishers, Inc., Boulder,
2001).

6.   Goffe, A. S., Zinner, D. & Fischer, J. Sex and friendship in a multilevel society: behavioural
patterns and associations between female and male Guinea baboons. *Behav. Ecol. Sociobiol.*
70, 323–336 (2016).

7.   Ostner, J. & Schülke, O. The evolution of social bonds in primate males. *Behaviour* **151**, 871–906
(2014).

8.   Seyfarth, R. M., Cheney, D. L. & Bergman, T. J. Primate social cognition and the origins of
language. *Trends Cogn. Sci.* **9**, 264–266 (2005).

9.   Pfefferle, D., Kazem, A. J. N., Brockhausen, R. R., Ruiz-Lambides, A. V. & Widdig, A. Monkeys
spontaneously discriminate their unfamiliar paternal kin under natural conditions using facial
cues. *Curr. Biol.* **24**, 1806–1810 (2014).

10.   Pfefferle, D., Ruiz-Lambides, A. V. & Widdig, A. Female rhesus macaques discriminate unfamiliar
paternal sisters in playback experiments – support for acoustic phenotype matching. *Proc. R.*
Soc. B **281**, 1–9 (2014).

11.   Cheney, D. L. & Seyfarth, R. M. Recognition of other individuals' social relationships by female
baboons. *Anim. Behav.* **58**, 67–75 (1999).

12.   Cheney, D. L., Seyfarth, R. M. & Palombit, R. The function and mechanisms underlying baboon
'contact' barks. *Anim. Behav.* **52**, 507–518 (1996).

13.   Moreira, L. A. A., Pessoa, D. M. A. & Sousa, M. B. C. Socio-sexual communication: A review of
the sensory modalities used by non-human primates. *Oecologia Aust.* **17**, 303–319 (2013).

14.   Yorzinski, J. L. The cognitive basis of individual recognition. *Curr. Opin. Behav. Sci.* **16**, 53–57
(2017).

15.   Wyatt, T. *Pheromones and Animal Behavior: Chemical Signals and Signatures*. (Cambridge
University Press, 2014).

16.   Stevenson, R. J. An Initial Evaluation of the Functions of Human Olfaction. *Chem. Senses* **35**, 3–
20 (2010).

17.   Bates, L. A. *et al.* African elephants have expectations about the locations of out-of-sight family
members. *Biol. Lett.* **4**, 34–36 (2008).

18.   Linklater, W. L., Mayer, K. & Swaisgood, R. R. Chemical signals of age, sex and identity in black
rhinoceros. *Anim. Behav.* **85**, 671–677 (2013).

19.   delBarco-Trillo, J., Sacha, C. R., Dubay, G. R. & Drea, C. M. Eulemur, me lemur: the evolution of
scent-signal complexity in a primate clade. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **367**, 1909–
1922 (2012).

20.   Janda, E. D., Perry, K. L., Hankinson, E., Walker, D. & Vaglio, S. Sex differences in scent-marking
in captive red-ruffed lemurs. *Am. J. Primatol.* **0**, e22951 (2019).

21.   Scordato, E. S. & Drea, C. M. Scents and sensibility: information content of olfactory signals in
the ringtailed lemur, Lemur catta. *Anim. Behav.* **73**, 301–314 (2007).

22.   Drea, C. M. & Scordato, E. S. Olfactory Communication in the Ringtailed Lemur (Lemur catta):
Form and Function of Multimodal Signals. in *Chemical signals in Vertebrates* (eds. Hurst, J. L.,
Beynon, R. J., Roberts, S. C. & Wyatt, T. D.) vol. 11 91–102 (Springer, New York, NY, 2008).

23.   Fontani, S., Kaburu, S. S. K., Marliani, G., Accorsi, P. A. & Vaglio, S. Anogenital scent-marking
signals fertility in a captive female Alaotran gentle lemur. *Front. Vet. Sci.* **9**, (2022).

24.   Kücklich, M., Weiß, B. M., Birkemeyer, C., Einspanier, A. & Widdig, A. Chemical cues of female
fertility states in a non-human primate. *Sci. Rep.* **9**, 1–12 (2019).

25.   Heymann, E. W. Scent marking strategies of new world primates. *Am. J. Primatol.* **68**, 650–661
(2006).

26.   Veilleux, C. C., Dominy, N. J. & Melin, A. D. The sensory ecology of primate food perception,
revisited. *Evol. Anthropol. Issues News Rev.* **31**, 281–301 (2022).

27.   Kawamura, S. Color vision diversity and significance in primates inferred from genetic and field
studies. *Genes Genomics* **38**, 779–791 (2016).

28. Freeman, N. J., Pasternak, G. M., Rubi, T. L., Barrett, L. & Henzi, S. P. Evidence for scent marking
in vervet monkeys? *Primates* **53**, 311–315 (2012).
29. Smith, T. D., Laitman, J. T. & Bhatnagar, K. P. The shrinking anthropoid nose, the human
vomeronasal organ, and the language of anatomical reduction. *Anat. Rec.* 2196–2204 (2014)
doi:DOI: 10.1002/ar.23035.
30. Drea, C. M. D'scent of man: A comparative survey of primate chemosignaling in relation to sex.
Horm. Behav. **68**, 117–133 (2015).
31. Tang-Martinez, Z. The mechanisms of kin discrimination and the evolution of kin recognition in
vertebrates: a critical re-evaluation. *Behav. Processes* **53**, 21–40 (2001).
32. Laska, M. & Seibt, A. Olfactory sensitivity for aliphatic alcohols in squirrel monkeys and pigtail
macaques. *J. Exp. Biol.* **205**, 1633–1643 (2002).
33. Laska, M. & Seibt, A. Olfactory sensitivity for aliphatic esters in squirrel monkeys and pigtail
macaques. *Behav. Brain Res.* **134**, 165–174 (2002).
34. Heymann, E. W. The neglected sense-olfaction in primate behavior, ecology, and evolution. *Am.
J. Primatol.* **68**, 519–524 (2006).
35. Loos, H. M. *et al.* Past, Present, and Future of Human Chemical Communication Research.
Perspect. Psychol. Sci. 17456916231188147 (2023) doi:10.1177/17456916231188147.
36. Zetzsche, M. *et al.* Combined perceptual and chemical analyses show no compelling evidence
for ovulatory cycle shifts in women's axillary odour. *Proc. R. Soc. B Biol. Sci.* **291**, 20232712
(2024).
37. Setchell, J. M. *et al.* Chemical composition of scent-gland secretions in an old world monkey
(Mandrillus sphinx): influence of sex, male status, and individual identity. *Chem. Senses* **35**,
205–220 (2010).
38. Vaglio, S. *et al.* Sternal Gland Scent-Marking Signals Sex, Age, Rank, and Group Identity in
Captive Mandrills. *Chem. Senses* **41**, 177–186 (2016).
39. Henkel, S., Lambides, A., Berger, A., Thomsen, R. & Widdig, A. Rhesus macaques (*Macaca
mulatta*) recognize group membership via olfactory cues alone. *Behav. Ecol. Sociobiol.* **69**,
2019–2034 (2015).
40. Weiß, B. M. *et al.* Chemical composition of axillary odorants reflects social and individual
attributes in rhesus macaques. *Behav. Ecol. Sociobiol.* **72**, 65 (2018).
41. Henkel, S. & Setchell, J. M. Group and kin recognition via olfactory cues in chimpanzees (Pan
troglodytes). *Proc R Soc B* **285**, 20181527 (2018).
42. Vaglio, S., Minicozzi, P., Kessler, S. E., Walker, D. & Setchell, J. M. Olfactory signals and fertility
in olive baboons. *Sci. Rep.* **11**, 8506 (2021).
43. Rigaill, L. *et al.* Chemical cues of identity and reproductive status in Japanese macaques. *Am. J.
Primatol.* **84**, e23411 (2022).
44. Kuester, J. & Paul, A. Male migration in Barbary macaques (*Macaca sylvanus*) at Affenberg
Salem. *Int. J. Primatol.* **20**, 85–106 (1999).
45. von Segesser, F., Menard, N., Gaci, B. & Martin, R. D. Genetic differentiation within and
between isolated Algerian subpopulations of Barbary macaques (Macaca sylvanus): evidence
from microsatellites. *Mol. Ecol.* **8**, 433–442 (1999).
46. Roubová, V., Konečná, M., Šmilauer, P. & Wallner, B. Whom to Groom and for What? Patterns
of Grooming in Female Barbary Macaques (*Macaca sylvanus*). *PLoS ONE* **10**, e0117298 (2015).
47. Kuester, J., Paul, A. & Arnemann, J. Kinship, familiarity and mating avoidance in Barbary
macaques (*Macaca sylvanus*). *Anim. Behav.* **48**, 1183–1194 (1994).
48. Widdig, A. Paternal kin discrimination: The evidence and likely mechanisms. *Biol. Rev.* **82**, 319–
334 (2007).
49. Paul, A. & Kuester, J. Sex ratio adjustment in a seasonally breeding primate species: evidence
from Barbary macaque Population at Affenberg Salem. *Ethology* **74**, 117–132 (1987).
50. Prud'homme, J. & Chapais, B. Aggressive interventions and matrilineal dominance relations in
semifree-ranging Barbary macaques (Macaca sylvanus). *Primates* **34**, 271–283 (1993).

51.   Berghänel, A., Ostner, J. & Schülke, O. Coalitions destabilize dyadic dominance relationships in
male Barbary macaques (Macaca sylvanus). *Behaviour* **148**, 1256–1274 (2011).

52.   Young, C., Majolo, B., Schülke, O. & Ostner, J. Male social bonds and rank predict supporter
selection in cooperative aggression in wild Barbary macaques. *Anim. Behav.* **95**, 23–32 (2014).

53.   Paul, A., Kuester, J. & Arnemann, J. The sociobiology of male-infant interactions in Barbary
macaques, *Macaca sylvanus*. *Anim. Behav.* **51**, 155–170 (1996).

54.   Fischer, J. Emergence of individual recognition in young macaques. *Anim. Behav.* **67**, 655–661
(2004).

55.   Ritter, N. Multimodal perception of social stimuli in Barbary macaques at the Affenberg Salem.
(Leipzig University, Leipzig, 2022).

56.   Schell, A., Rieck, K., Schell, K., Hammerschmidt, K. & Fischer, J. Adult but not juvenile Barbary
macaques spontaneously recognize group members from pictures. *Anim. Cogn.* **14**, 503–509
(2011).

57.   Hammerschmidt, K. & Fischer, J. Maternal discrimination of offspring vocalizations in Barbary
macaques (*Macaca sylvanus*). *Primates* **39**, 231–236 (1998).

58.   Small, M. Promiscuity in Barbary macaques (*Macaca sylvanus*). *Am. J. Primatol.* **20**, 267–282
(1990).

59.   Paul, A., Kuester, J. & Arnemann, J. DNA fingerprinting reveals that infant care by male Barbary
macaques (Macaca sylvanus) is not paternal investment. *Folia Primatol. (Basel)* **58**, 93–98
(1992).

60.   Simon, M., Widdig, A. & Weiß, B. M. Sniffing behavior of semi free-ranging Barbary macaques
(Macaca sylvanus). *Am. J. Primatol.* **86**, e23611 (2024).

61.   Burgener, N., Dehnhard, M., Hofer, H. & East, M. L. Does anal gland scent signal identity in the
spotted hyaena? *Anim. Behav.* **77**, 707–715 (2009).

62.   Majolo, B., Lehmann, J., de Bortoli Vizioli, A. & Schino, G. Fitness-related benefits of dominance
in primates. *Am. J. Phys. Anthropol.* **147**, 652–660 (2012).

63.   Mehta, P. H. & Josephs, R. A. Testosterone and cortisol jointly regulate dominance: Evidence for
a dual-hormone hypothesis. *Horm. Behav.* **58**, 898–906 (2010).

64.   Naud, A. *et al.* Relations between Spatial Distribution, Social Affiliations and Dominance
Hierarchy in a Semi-Free Mandrill Population. *Front. Psychol.* **7**, (2016).

65.   Schoof, V. A. M., Jack, K. M. & Carnegie, S. D. Rise to Power: A Case Study of Male Fecal
Androgen and Cortisol Levels before and after a Non-Aggressive Rank Change in a Group of
Wild White-Faced Capuchins (Cebus capucinus). *Folia Primatol. (Basel)* **82**, 299–307 (2011).

66.   de Turckheim, G. & Merz, E. Breeding Barbary macaques in outdoor open enclosures. in *The*
Barbary Macaque: a Case Study in Conservation (ed. Fa, J. F.) 241–261 (Plenum Press, New
York, 1984).

67.   Kücklich, M. *et al.* Different Methods for Volatile Sampling in Mammals. *PLOS ONE* **12**,
e0183440 (2017).

68.   Weiß, B. M. *et al.* A non-invasive method for sampling the body odour of mammals. *Methods*
Ecol. Evol. **9**, 420–429 (2018).

69.   Stein, S. E. An integrated method for spectrum extraction and compound identification from
gas chromatography/mass spectrometry data. *J. Am. Soc. Mass Spectrom.* **10**, 770–781 (1999).

70.   Westphal, H. Paternity assessment in Barbary macaques. (Leipzig University, Leipzig, 2020).

71.   Westphal, H., Freudiger, A., Gatter, T., Stadler, P. F. & Widdig, A. TRACE - Tool for pedigree
Relatedness Analysis and Coefficient Estimation. (2023).

72.   Newton-Fisher, N. E. Modeling Social Dominance: Elo-Ratings, Prior History, and the Intensity of
Aggression. *Int. J. Primatol.* **38**, 427–447 (2017).

73.   Berghänel, A., Ostner, J., Schröder, U. & Schülke, O. Social bonds predict future cooperation in
male Barbary macaques, Macaca sylvanus. *Anim. Behav.* **81**, 1109–1116 (2011).

74.   Neumann, C. *et al.* Assessing dominance hierarchies: Validation and advantages of progressive
evaluation with Elo-rating. *Anim. Behav.* **82**, 911–921 (2011).

75. R Core Team. R: A Language and Environment for Statistical Computing. R foundation for
statistical computing (2024).
76. Oksanen, J. *et al.* vegan: Community Ecology Package. (2022).
77. Stoffel, M. A. *et al.* Chemical fingerprints encode mother–offspring similarity, colony
membership, relatedness, and genetic quality in fur seals. *Proc. Natl. Acad. Sci.* **112**, E5005–
E5012 (2015).
78. Chung, H. & Beretvas, S. N. The impact of ignoring multiple membership data structures in
multilevel models. *Br. J. Math. Stat. Psychol.* **65**, 185–200 (2012).
79. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *J. Stat. Softw.* **80**,
1–28 (2017).
80. Jamil, T., Ozinga, W. A., Kleyer, M. & ter Braak, C. J. F. Selecting traits that explain species–
environment relationships: a generalized linear mixed model approach. *J. Veg. Sci.* **24**, 988–
1000 (2013).
81. Boogaart, K. G. van den & Tolosana-Delgado, R. *Analyzing Compositional Data with R*. (Springer,
Heidelberg, 2013).
82. Schielzeth, H. Simple means to improve the interpretability of regression coefficients. *Methods*
Ecol. Evol. **1**, 103–113 (2010).
83. Fox, J. & Weisberg, S. An {R} companion to applied regression. in (Sage, Thousand Oaks CA,
2019).
84. Vehtari, A., Gelman, A., Simpson, D., Carpenter, B. & Bürkner, P.-C. Rank-Normalization,
Folding, and Localization: An Improved Rˆ for Assessing Convergence of MCMC (with
Discussion). *Bayesian Anal.* **16**, 667–718 (2021).
85. Smith, T. Individual olfactory signatures in common marmosets (Callithrix jacchus). *Am. J.*
Primatol. **68**, 585–604 (2006).
86. Buesching, C. D., Waterhouse, J. S. & Macdonald, D. W. Gas-chromatographic analyses of the
subcaudal gland secretion of the European badger (*Meles meles*) part II: time-related variation
in the individual-specific composition. *J. Chem. Ecol.* **28**, 57–69 (2002).
87. Caspers, B., Franke, S. & Voigt, C. C. The Wing-Sac Odour of Male Greater Sac-Winged Bats
Saccopteryx bilineata (Emballonuridae) as a Composite Trait: Seasonal and Individual
Differences. in *Chemical Signals in Vertebrates 11* (eds. Hurst, J. L., Beynon, R. J., Roberts, S. C.
& Wyatt, T. D.) 151–160 (Springer New York, New York, 2008).
88. Kwak, J. *et al.* Genetically-Based Olfactory Signatures Persist Despite Dietary Variation. *PLoS*
ONE **3**, e3591 (2008).
89. Kulahci, I. G., Drea, C. M., Rubenstein, D. I. & Ghazanfar, A. A. Individual recognition through
olfactory-auditory matching in lemurs. *Proc. R. Soc. B Biol. Sci.* **281**, 20140071 (2014).
90. Palagi, E. & Dapporto, L. Beyond Odor Discrimination: Demonstrating Individual Recognition by
Scent in *Lemur Catta*. *Chem. Senses* **31**, 437–443 (2006).
91. Spence-Aizenberg, A., Kimball, B. A., Williams, L. E. & Fernandez-Duque, E. Chemical
composition of glandular secretions from a pair-living monogamous primate: Sex, age, and
gland differences in captive and wild owl monkeys (Aotus spp.). *Am. J. Primatol.* **80**, e22730
(2018).
92. Scordato, E. S., Dubay, G. & Drea, C. M. Chemical composition of scent marks in the ringtailed
lemur (*Lemur catta*): glandular differences, seasonal variation, and individual signatures. *Chem.*
Senses **32**, 493–504 (2007).
93. Penn, D. J. *et al.* Individual and gender fingerprints in human body odour. *J. R. Soc. Interface* **4**,
331–340 (2007).
94. MacDonald, E. A., Fernandez-Duque, E., Evans, S. & Hagey, L. R. Sex, age, and family differences
in the chemical composition of owl monkey (*Aotus nancymaae*) subcaudal scent secretions.
Am. J. Primatol. **70**, 12–18 (2008).
95. Theis, K. R., Schmidt, T. M. & Holekamp, K. E. Evidence for a bacterial mechanism for group-
specific social odors among hyenas. *Sci. Rep.* **2**, 1–8 (2012).

96.   Boulet, M., Charpentier, M. & Drea, C. Decoding an olfactory mechanism of kin recognition and
inbreeding avoidance in a primate. *BMC Evol. Biol.* **9**, 281 (2009).
97.   Charpentier, M. J. E., Boulet, M. & Drea, C. M. Smelling right: the scent of male lemurs
advertises genetic quality and relatedness. *Mol. Ecol.* **17**, 3225–3233 (2008).
98.   Kazem, A. J. N. & Widdig, A. Visual phenotype matching: cues to paternity are present in rhesus
macaque faces. *PLoS ONE* **8**, e55846 (2013).
99.   Kazem, A. J. N., Barth, Y., Pfefferle, D., Kulik, L. & Widdig, A. Parent–offspring facial resemblance
increases with age in rhesus macaques. *Proc R Soc B* **285**, 20181208 (2018).
100. Kean, E. F., Müller, C. T. & Chadwick, E. A. Otter scent signals age, sex, and reproductive status.
Chem. Senses **36**, 555–564 (2011).
101. Leclaire, S., Jacob, S., Greene, L. K., Dubay, G. R. & Drea, C. M. Social odours covary with
bacterial community in the anal secretions of wild meerkats. *Sci. Rep.* **7**, 3240 (2017).
102. Jänig, S., Weiß, B. M., Birkemeyer, C. & Widdig, A. Comparative chemical analysis of body odor
in great apes. *Am. J. Primatol.* **0**, e22976 (2019).
103. Marneweck, C., Jürgens, A. & Shrader, A. M. Dung odours signal sex, age, territorial and
oestrous state in white rhinos. *Proc R Soc B* **284**, 20162376 (2017).
104. Drea, C. M. *et al.* The "Secret" in Secretions: Methodological Considerations in Deciphering
Primate Olfactory Communication. *Am. J. Primatol.* **75**, 621–642 (2013).
105. Charpentier, M. J. E., Barthes, N., Proffit, M., Bessière, J.-M. & Grison, C. Critical thinking in the
chemical ecology of mammalian communication: roadmap for future studies. *Funct. Ecol.* **26**,
769–774 (2012).
106. Costello, B. de L. *et al.* A review of the volatiles from the healthy human body. *J. Breath Res.* **8**,
014001 (2014).
107. Mansur, M. A. A., Siddiqi, M. M. A., Akbor, M. A. & Saha, K. Phytochemical Screening and GC-
MS Chemical Profiling of Ethyl Acetate Extract of Seed and Stem of Anethum sowa Linn. *Dhaka*
Univ. J. Pharm. Sci. **16**, 187–194 (2018).
108. Dunbar, R. I. M. Social structure as a strategy to mitigate the costs of group living: a comparison
of gelada and guereza monkeys. *Anim. Behav.* **136**, 53–64 (2018).
109. Roberts, S. C. *et al.* Emotional expression in human odour. *Evol. Hum. Sci.* 1–35 (2022)
doi:10.1017/ehs.2022.44.
110. Crawford, J. C., Boulet, M. & Drea, C. M. Smelling wrong: hormonal contraception in lemurs
alters critical female odour cues. *Proc. R. Soc. B Biol. Sci.* **278**, 122–130 (2011).
111. Charpentier, M. J. E., Mboumba, S., Ditsoga, C. & Drea, C. M. Nasopalatine Ducts and Flehmen
Behavior in the Mandrill: Reevaluating Olfactory Communication in Old World Primates. *Am. J.*
Primatol. **75**, 703–714 (2013).
112. Maidodou, L. *et al.* Unraveling the potential of breath and sweat VOC capture devices for
human disease detection: a systematic-like review of canine olfaction and GC-MS analysis.
Front. Chem. **11**, 1282450 (2023).
113. Garner, C. E. *et al.* Volatile organic compounds from feces and their potential for diagnosis of
gastrointestinal disease. *FASEB J.* **21**, 1675–1688 (2007).
114. Bernier, U. R., Kline, D. L., Barnard, D. R., Schreck, C. E. & Yost, R. A. Analysis of Human Skin
Emanations by Gas Chromatography/Mass Spectrometry. 2. Identification of Volatile
Compounds That Are Candidate Attractants for the Yellow Fever Mosquito (*Aedes aegypti*).
Anal. Chem. **72**, 747–756 (2000).
115. Smith, T., Tomlinson, A., Mlotkiewicz, J. & Abbott, D. Female marmoset monkeys (*Callithrix*
jacchus) can be identified from the chemical composition of their scent marks. *Chem. Senses*
26, 449–458 (2001).
116. Burger, B. V. Mammalian Semiochemicals. *Top. Curr. Chem.* **240**, 231–278 (2005).
117. Gallagher, M. *et al.* Analyses of volatile organic compounds from human skin. *Br. J. Dermatol.*
159, 780–791 (2008).
981
982